



US008173117B2

(12) United States Patent
Hardiman et al.**(10) Patent No.: US 8,173,117 B2**
(45) Date of Patent: May 8, 2012**(54) MOUSE CX3C CHEMOKINE POLYPEPTIDES****(75) Inventors:** Gerard T. Hardiman, San Diego, CA (US); Devora L. Rossi, San Diego, CA (US); Kevin B Bacon, Kobe (JP); Bazan Fernando J., Palo Alto, CA (US); Thomas P. Schall, Menlo Park, CA (US); Albert Zlotnik, Palo Alto, CA (US)**(73) Assignee:** Schering Corporation, Kenilworth, NJ (US)**(*) Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.**(21) Appl. No.:** 12/833,069**(22) Filed:** Jul. 9, 2010**(65) Prior Publication Data**

US 2011/0189123 A1 Aug. 4, 2011

Related U.S. Application Data**(60)** Continuation of application No. 12/510,586, filed on Jul. 28, 2009, now Pat. No. 7,785,804, which is a continuation of application No. 12/198,626, filed on Aug. 26, 2008, now Pat. No. 7,585,502, which is a continuation of application No. 11/495,801, filed on Jul. 28, 2006, now Pat. No. 7,431,924, which is a continuation of application No. 10/397,559, filed on Mar. 25, 2003, now Pat. No. 7,115,379, which is a division of application No. 09/771,023, filed on Jan. 25, 2001, now Pat. No. 6,566,503, which is a division of application No. 09/093,482, filed on Jun. 8, 1998, now abandoned, which is a division of application No. 08/786,068, filed on Jan. 21, 1997, now abandoned, which is a continuation of application No. 08/649,006, filed on May 16, 1996, now Pat. No. 6,548,654, which is a continuation of application No. 08/590,828, filed on Jan. 24, 1996, now abandoned.**(51) Int. Cl.****C07K 14/52** (2006.01)**A61K 38/19** (2006.01)**C12N 15/64** (2006.01)**(52) U.S. Cl. 424/85.1; 530/351; 530/402; 435/69.5****(58) Field of Classification Search** None
See application file for complete search history.**(56) References Cited**

U.S. PATENT DOCUMENTS

6,013,257 A 1/2000 Pan
6,043,086 A 3/2000 Pan
6,548,654 B1 4/2003 Hardiman et al.
6,566,503 B2 5/2003 Hardiman et al.
7,115,379 B1 10/2006 Hardiman et al.
7,431,924 B2 10/2008 Hardiman et al.

FOREIGN PATENT DOCUMENTS

WO WO97/42224 11/1997

OTHER PUBLICATIONS

Mikayama et al. Proc. Natl. Acad. Sci. USAvol. 90, pp. 10056-10060, (1993).
Voet et al. Biochemistry John Wiley & Sons, Inc., pp. 126-128 and 228-234, (1990).
Kevin B. Bacon et al., Int. Arch. Allergy Immunol., 109:97-109, 1996. "Chemokines as Mediators of Allergic Inflammation".
J. Fernando Bazan et al., Nature, 385:640-644, Feb. 13, 1997. A new class of membrane-bound chemokine with a CX₃C motif.
D. Beier et al., Genbank Accession No. R75309, p. 1, Jun. 6, 1995.
Mark S. Boguski, Trends Biochem. Sci. (TIBS), 20:295-296, 1995. "The turning point in genome research".
Myriam C. Capone et al., J. Immunol., 157:969-973, 1996. "Identification Through Bioinformatics of cDNAs Encoding Human Thymic Shared Ag-1/Stem Cell Ag-2: A New Member of the Human Ly-6 Family.sup.1,2".
Angela M. Gronenborn et al., Protein Engineering, 4(3):263-269, 1991. "Modeling the Three-Dimensional Structure of the Monocyte Chemo-Attractant and Activating Protein MCAF/MCP-1 on the Basis of the Solution Structure of Interleukin-8".
L. Hillier et al., Genbank Accession No. H06104, pp. 1-2, Jun. 21, 1995.
Patricia J. Lodi et al., Science, 263:1762-1767, 1994. "High-Resolution Solution Structure of the .beta. Chemokine hMIP-1.beta. By Multidimensional NMR".
Kouji Matsushima et al., Cytokine, 1(1):2-13, 1989. "Interleukin 8 and MCAF: Novel Inflammatory Cytokines Inducible by IL 1 and TNF".
Michael D. Miller et al., Proc. Natl. Acad. Sci., 89:2950-2954, 1992. "The Human Cytokine I-309 is a Monocyte Chemoattractant".
Joost J. Oppenheim et al., Annu. Rev. Immunol., 9:617-648, 1991. "Properties of the Novel Proinflammatory Supergene 'intercrine' Cytokine Family".
Thomas J. Schall, The Cytokine Handbook, Angus W. Thompson, Ed., 2nd ed. The Academic Press Limited, Ch. 22, pp. 419-460, 1994. "The Chemokines".
Thomas J. Schall et al, Current Opinion in Immunol., 6:865-873, 1994. "Chemokines, Leukocyte Trafficking, and Inflammation".
Thomas J. Schall, Cytokine, 3(3):165-183, 1991. "Biology of the RANTES/SIS Cytokine Family".
Mark Y. Stoeckle et al., The New Biologist, 2(4):313-323, 1990. "Two Burgeoning Families of Platelet Factor 4-Related Proteins: Mediators of the Inflammatory Response". Steven R. Wiley et al., Immunity, 3: 673-682, 1995. "Identification and Characterization of a New Member of the TNF Family that Induces Apoptosis".
D. Rossi et al., FASEB Journal, 10(6):A1049, Abstract No. 290, from Joint Meeting of the American Society for Biochemistry and Molecular Biology, The American Society for Investigative Pathology and the American Association of Immunologists, New Orleans, LA, USA Jun. 1996. "Identification of a .gamma. or "C" chemokine from the chicken and a novel human and mouse chemokine".
L. Hillier et al., EMBL Database Entry HS940163, Accession No. H14940, Jul. 3, 1995, XP002030088. "The WashU-Merck EST project".
C. Auffray et al., EMBL Database Entry HSC20F041, Accession No. Z44443, Sep. 21, 1995, XP002030091. "Image: Integrated molecular analysis of the human genome and its expression".

(Continued)

Primary Examiner — Prema Mertz

(57) ABSTRACT

Nucleic acids encoding a new family of chemokines, the CX3C family, from a mammal, reagents related thereto, including specific antibodies, and purified proteins are described. Methods of using said reagents and related diagnostic kits are also provided.

8 Claims, No Drawings

OTHER PUBLICATIONS

- Fiorenza Cocchi et al., Science, 270:1811-1815, Dec. 15, 1995. "Identification of RANTES, MIP-1.alpha., and MIP-1-.beta. as the Major HIV-Suppressive Factors Produced by CD8.sup.+ T Cells". Derwent Geneseq Patent Sequence Database, Accession No. 97P-W34307, citing: WO 9742224 A1 dated Nov. 13, 1997; US 97-851160 dated May 5, 1997; US 96-643798 dated May 7, 1996; "Search record pertaining to human CX3C chemokine, performed in Jul. 1998".
- Derwent Geneseq Patent Sequence Database, Accession No. 19P-W23345, citing: WO 9727299 A1 dated Jul. 31, 1997; WO 97-US293 dated Jan. 23, 1997; US 96-649006 dated May 16, 1996; US 96-590828 dated Jan. 24, 1996; "Search record pertaining to human CX3C chemokine, performed in Jul. 1998".
- Derwent Geneseq Patent Sequence Database, Accession No. 97P-W34308, citing: WO 9742224 A1 dated Nov. 13, 1997; WO 97-US7737 dated May 6, 1997; US 97-851160 dated May 5, 1997; US 96-643798 dated May 7, 1996; "Search record pertaining to human CX3C chemokine, performed in Jul. 1998".
- Derwent Geneseq Patent Sequence Database, Accession No. 19P-W23347, citing: WO 9727299 A1 dated Jul. 31, 1997; WO 97-US293 dated Jan. 23, 1997; US 96-649006 dated May 16, 1996; US 96-590828 dated Jan. 24, 1996; "Search record pertaining to human CX3C chemokine, performed in Jul. 1998".
- Derwent Geneseq Patent Sequence Database, Accession No. 19P-W23344, citing: WO 9727299 A1 dated Jul. 31, 1997; WO 97-US293 dated Jan. 23, 1997; US 96-6496006 dated Jan. 6, 1996; US 96-590828 dated Jan. 24, 1996; "Search record pertaining to human CX3C chemokine, performed in Jul. 1998".
- Derwent Geneseq Patent Sequence Database, Accession No. 19P-W23346, citing: WO 9727299 A1 dated Jul. 31, 1997; WO 97-US293 dated Jan. 23, 1997; US 96-649006 dated May 16, 1996; US 96-590828 dated Jan. 24, 1996; "Search record pertaining to human CX3C chemokine, performed in Jul. 1998".
- Derwent Geneseq Patent Sequence Database, Accession No. 98P-W42072, citing: WO 9802459 A1 dated Jan. 22, 1998; WO 97-US12349 dated Jul. 15, 1997; US 96-683655 dated 960715; "Search record pertaining to human CX3C chemokine, performed in Jul. 1998".
- Derwent Geneseq Patent Sequence Database, Accession No. 97P-W26655, citing: WO 9725427 A1 dated Jul. 17, 1997; WO 97-US379 dated Jan. 10, 1997; US 96-586395 dated Jan. 12, 1996; "Search record pertaining to human CX3C chemokine, performed in Jul. 1998".
- Derwent Geneseq Patent Sequence Database, Accession No. 97P-W3810, citing: WO 9741230 A2 dated Nov. 6, 1997; WO 97-EP2217 dated Apr. 30, 1997; DE 96-19617312 dated Apr. 30, 1996; "Search record pertaining to human CX3C chemokine".
- James U. Bowie et al., Science, 247: 1306-1310, Mar. 16, 1990. "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions".
- Claude Daniel et al., Virology, 202:540-549, 1994. "Mapping of Linear Antigenic Sites on the S Glycoprotein of a Neurotropic Murine Coronavirus with Synthetic Peptides: A Combination of Nine Prediction Algorithms Fails to Identify Relevant Epitopes and Peptide Immunogenicity Is Drastically Influenced by the Nature of the Protein Carrier".
- C. Frommel et al., J. Mol. Evol., 21:233-257, 1985. "An Estimate on the Effect of Point Mutation and Natural Selection on the Rate of Amino Acid Replacement in Proteins".
- David G. George et al., Macromolecular Sequencing and Synthesis, David H. Schlesinger, Ed., Chapter 12, pp. 127-149, Alan R. Liss, Inc.: New York 1988. "Current Methods in Sequence Comparison and Analysis".
- Minoru Kanehisa, Nucleic Acids Research, 12(1):203-213, 1984. "Use of statistical criteria for screening potential homologies in nucleic acid sequences".
- J. Thomas Ngo et al., The Protein Folding Problem and Tertiary Structure Prediction, K. Mertz, Jr. and S. Le Grand, Eds., Chapter 14, pp. 491-495, Birkhauser: Boston 1994. "Computational Complexity, Protein Structure Prediction and the Levinthal Paradox".
- G. Opdenakker et al., GenBank, Accession Nos. X72308 & S57464, Jul. 25, 1994. Definition: "*H. sapiens* MCP-3 mRNA for monocyte chemotactic protein-3".
- Sambrook, et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Chapters 9 and 11, CSHL: Cold Spring Harbor, New York, 1989.
- Mikayama et al., Proc Natl Acad Sci U S A. Nov. 1, 1993;90(21):10056-60. "Molecular cloning and functional expression of a cDNA encoding glycosylation-inhibiting factor".
- Voet et al., Biochemistry. John Wiley & Sons, Inc. pp. 126-128 and 228-234.
- C. Auffray et al., Genbank Accession No. Z44443, Sep. 21, 1995. Definition: *H. Sapiens* partial cDNA sequence; clone c-20f04.

* cited by examiner

MOUSE CX3C CHEMOKINE POLYPEPTIDES

The present application is a continuation of U.S. patent application Ser. No. 12/510,586, filed Jul. 28, 2009; which is a continuation of U.S. patent application Ser. No. 12/198,626, filed Aug. 26, 2008; which is a continuation of U.S. patent application Ser. No. 11/495,801, filed Jul. 28, 2006; which is a continuation of U.S. patent application Ser. No. 10/397,559, filed Mar. 25, 2003; which is a division of U.S. patent application Ser. No. 09/771,023, filed Jan. 25, 2001, now U.S. Pat. No. 6,566,503; which is a division of U.S. patent application Ser. No. 09/093,482, filed Jun. 8, 1998; which is a division of U.S. patent application Ser. No. 08/786,068, filed Jan. 21, 1997; which is a continuation of U.S. patent application Ser. No. 08/649,006, filed May 16, 1996, now U.S. Pat. No. 6,548,654; which is a continuation of U.S. patent application Ser. No. 08/590,828 filed Jan. 24, 1996, each of which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The present invention contemplates compositions related to proteins which function in controlling development, differentiation, trafficking, and physiology of mammalian cells, e.g., cells of a mammalian immune system. In particular, it provides proteins which regulate or evidence development, differentiation, and function of various cell types, including hematopoietic cells.

BACKGROUND OF THE INVENTION

The circulating component of the mammalian circulatory system comprises various cell types, including red and white blood cells of the erythroid and myeloid cell lineages. See, e.g., Rapaport (1987) *Introduction to Hematology* (2d ed.) Lippincott, Philadelphia, Pa.; Jandl (1987) *Blood: Textbook of Hematology*, Little, Brown and Co., Boston, Mass.; and Paul (ed.) (1993) *Fundamental Immunology* (3d ed.) Raven Press, N.Y.

For some time, it has been known that the mammalian immune response is based on a series of complex cellular interactions, called the "immune network." Recent research has provided new insights into the inner workings of this network. While it remains clear that much of the response does, in fact, revolve around the network-like interactions of lymphocytes, macrophages, granulocytes, and other cells, immunologists now generally hold the opinion that soluble proteins, known as lymphokines, cytokines, or monokines, play a critical role in controlling these cellular interactions. Thus, there is considerable interest in the isolation, characterization, and mechanisms of action of cell modulatory factors, an understanding of which should lead to significant advancements in the diagnosis and therapy of numerous medical abnormalities, e.g., immune system and other disorders.

Lymphokines apparently mediate cellular activities in a variety of ways. They have been shown to support the proliferation, growth, and differentiation of pluripotential hematopoietic stem cells into vast numbers of progenitors comprising diverse cellular lineages making up a complex immune system. These interactions between cellular components are necessary for a healthy immune response. These different cellular lineages often respond in a different manner when lymphokines are administered in conjunction with other agents.

The chemokines are a large and diverse superfamily of proteins. The superfamily is subdivided into three branches,

based upon whether the first two cysteines in the classical chemokine motif are adjacent (termed the "C—C" branch) or spaced by an intervening residue ("C—X—C"), or a new branch which lacks two cysteines in the corresponding motif, represented by the chemokines known as lymphotactins. See, e.g., Schall and Bacon (1994) *Current Opinion in Immunology* 6:865-873; and Bacon and Schall (1996) *Int. Arch. Allergy & Immunol.* 109:97-109.

Many factors have been identified which influence the differentiation process of precursor cells, or regulate the physiology or migration properties of specific cell types. These observations indicate that other factors exist whose functions in immune function were heretofore unrecognized. These factors provide for biological activities whose spectra of effects may be distinct from known differentiation or activation factors. The absence of knowledge about the structural, biological, and physiological properties of the regulatory factors which regulate cell physiology in vivo prevents the modification of the effects of such factors. Thus, medical conditions where regulation of the development or physiology of relevant cells is inappropriate remain unmanageable.

SUMMARY OF THE INVENTION

The present invention reveals the existence of a previously unknown class of chemokine-motif containing molecules which are hereby designated the CX3C chemokines. The CX3Ckines have three amino acids which separate the cysteines in the corresponding region of the chemokine motif. Based on sequence analysis of the two CX3C protein sequences described below, it is apparent that the CX3Ckines do not belong to the C, C—C, or C—X—C chemokine families. They represent the first known members of a new heretofore unidentified class of chemokines designated CX3Ckines, or alternatively, the CX3C family of chemokines.

The present invention provides a composition of matter selected from an antibody binding site which specifically binds to a mammalian CX3C chemokine; an expression vector encoding a mammalian CX3C chemokine or fragment thereof; a substantially pure protein which is specifically recognized by the antibody binding site; and a substantially pure CX3C chemokine or peptide thereof, or a fusion protein comprising a 30 amino acid fragment of CX3C chemokine sequence.

In the antibody binding site embodiments, the antibody binding site may be: specifically immunoreactive with a mature protein selected from the group consisting of the polypeptides of SEQ ID NO: 2, 4, 6 and 8; raised against a purified or recombinantly produced human or mouse CX3C chemokine; in a monoclonal antibody, Fab, or F(ab)₂; immunoreactive with denatured antigen; or in a labeled antibody. In certain embodiments; the antibody binding site is detected in a biological sample by a method of: contacting a binding agent having an affinity for CX3C chemokine protein with the biological sample; incubating the binding agent with the biological sample to form a binding agent: CX3C chemokine protein complex; and detecting the complex. In a preferred embodiment, the biological sample is human, and the binding agent is an antibody.

A kit embodiment is provided possessing a composition, described above, with either instructional material for the use of the composition; or segregation of the composition into a container.

A nucleic acid embodiment of the invention includes an expression vector encoding a CX3C chemokine protein, wherein the protein specifically binds an antibody generated

against an immunogen selected from the mature polypeptide portions of SEQ ID NO: 2, 4, 6, and 8. The vector may: encode a CX3C chemokine polypeptide with complete sequence identity to a naturally occurring human CX3C chemokine domain; encode a CX3C chemokine protein comprising sequence selected from the polypeptides of SEQ ID NO: 2, 4, 6, and 8; or comprise sequence selected from the nucleic acids of SEQ ID NO: 1, 3, 5, or 7. In other embodiments, the vector is capable of selectively hybridizing to a nucleic acid encoding a CX3C chemokine protein, e.g., a mature protein coding segment of SEQ ID NO: 1, 3, 5, or 7. In various preferred embodiments, the isolated nucleic acid is detected in a biological sample by a method: contacting a biological sample with a nucleic acid probe capable of selectively hybridizing to the nucleic acid; incubating the nucleic acid probe with the biological sample to form a hybrid of the nucleic acid probe with complementary nucleic acid sequences present in the biological sample; and determining the extent of hybridization of the nucleic acid probe to the complementary nucleic acid sequences. In such method, preferably the nucleic acid probe is capable of hybridizing to a nucleic acid encoding a protein consisting of the polypeptides of SEQ ID NO: 2, 4, 6, or 8.

In protein embodiments, the isolated CX3C chemokine protein will preferably be of approximately 11,000 to 15,000 daltons when in unglycosylated form, and the CX3C chemokine protein specifically binds to an antibody generated against an immunogen; the polypeptides of SEQ ID NO: 2, 4, 6, or 8; and the CX3C chemokine lacks the cysteine structural motifs and sequence characteristic of a C, a CC, or a CXC chemokine. In various embodiments, the isolated CX3C chemokine protein is: selected from human CX3Ckine or mouse CX3Ckine; consists of a polypeptide comprising sequence from SEQ ID NO: 2, 4, 6, or 8; recombinantly produced, or a naturally occurring protein.

The present invention also embraces a cell transfected with the nucleic acid encoding a CX3C chemokine, e.g., where the nucleic acid has SEQ ID NO: 1, 3, 5, or 7.

The invention also provides a method of modulating physiology or development of a cell by contacting the cell with a CX3C chemokine, or an antagonist of the chemokine. In preferred embodiments, the physiology is attraction, and the cell is a peripheral blood monocyte or a T cell.

DETAILED DESCRIPTION

I. General

The present invention provides DNA sequences encoding mammalian proteins which exhibit structural properties or motifs characteristic of a cytokine or chemokine. For a review of the chemokine family, see, e.g., Lodi, et al. (1994) *Science* 263:1762-1767; Gronenborn and Clore (1991) *Protein Engineering* 4:263-269; Miller and Kranger (1992) *Proc. Nat'l Acad. Sci. USA* 89:2950-2954; Matsushima and Oppenheim (1989) *Cytokine* 1:2-13; Stoeckle and Baker (1990) *New Biol.* 2:313-323; Oppenheim, et al. (1991) *Ann. Rev. Immunol.* 9:617-648; Schall (1991) *Cytokine* 3:165-183; and *The Cytokine Handbook* Academic Press, NY. The proteins described herein are designated CX3Ckines because they were initially recognized as sharing significant structural features of chemokines, but whose structural features also exhibit sequence peculiarity, e.g., structural motifs, distinct from the other known branches of the chemokine molecules.

The best characterized embodiment of this family of proteins was discovered from a human and is designated human CX3C chemokine (GenBank Accession No. H14940). See, SEQ ID NO: 1-4 An additional CX3Ckine, represented by a mouse molecule, designated mouse CX3Ckine, is also described herein. See Table 1 and SEQ ID NO: 5-8. The descriptions below are directed, for exemplary purposes, to primate and rodent embodiments, e.g., human and mouse, but are likewise applicable to related embodiments from other, e.g., natural, sources. These sources should include various vertebrates, typically warm blooded animals, e.g., birds and mammals, particularly domestic animals, and primates.

In the human sequence (SEQ ID NO: 1-4), the signal sequence runs from about Met1 to Gly24, thus the mature polypeptide begins at about Gln25 and ends at about Val 397. A chemokine domain runs from about Gln25 to about Gly100; a stalk region, which possesses many potential glycosylation sites, runs from about Gly101 to about Gln341; a transmembrane region begins at about Ala342 and ends at about Thr361; and an intracellular domain, containing two tyrosine phosphorylation sites at residues 382 and 392, runs from about Tyr362 to Val397.

TABLE 1

Mouse CX3C chemokine nucleic acid (SEQ ID NO: 7) and amino acid (SEQ ID NO: 8) sequences.	
TGACTACTAG GAGCTGCGAC ACGGCCAGC CTCTGGCCG CCGAATTCT CCACTCCAGC	60
C ATG GCT CCC TCG CCG CTC GCG TGG CTG CTG CGC CTG GCC GCG TTC Met Ala Pro Ser Pro Leu Ala Trp Leu Leu Arg Leu Ala Ala Phe	106
1 5 10 15	
TTC CAT TTG TGT ACT CTG CTG CCG GGT CAG CAC CTC GGC ATG ACG AAA Phe His Leu Cys Thr Leu Leu Pro Gly Gln His Leu Gly Met Thr Lys	154
20 25 30	
TGC GAA ATC ATG TGC GGC AAG ATG ACC TCA CGA ATC CCA GTG GCT TTG Cys Glu Ile Met Cys Gly Lys Met Thr Ser Arg Ile Pro Val Ala Leu	202
35 40 45	
CTC ATC CGC TAT CAG CTA AAT CAG GAG TCC TGC GGC AAG CGT GCC ATT Leu Ile Arg Tyr Gln Leu Asn Gln Glu Ser Cys Gly Lys Arg Ala Ile	250
50 55 60	
GTC CTG GAG ACG ACA CAG CAC AGA CGC TTC TGT GCT GAC CCG AAG GAG Val Leu Glu Thr Thr Gln His Arg Arg Phe Cys Ala Asp Pro Lys Glu	298
65 70 75	

TABLE 1-continued

Mouse CX3C chemokine nucleic acid (SEQ ID NO: 7) and amino acid (SEQ ID NO: 8) sequences.														
AAA TGG GTC CAA GAC GCC ATG AAG CAT CTG GAT CAC CAG GCT GCT GCC	346													
Lys Trp Val Gln Asp Ala Met Lys His Leu Asp His Gln Ala Ala Ala														
80 85 90 95														
CTC ACT AAA AAT GGT GGC AAG TTT GAG AAG CCG GTG GAC AAT GTG ACA	394													
Leu Thr Lys Asn Gly Gly Lys Phe Glu Lys Arg Val Asp Asn Val Thr														
100 105 110														
CCT GGG ATC ACC TTG GCC ACT AGG GGA CTG TCC CCA TCT GCC CTG ACA	442													
Pro Gly Ile Thr Leu Ala Thr Arg Gly Leu Ser Pro Ser Ala Leu Thr														
115 120 125														
AAG CCT GAA TCC GCC ACA TTG GAA GAC CTT GCT TTG GAA CTG ACT ACT	490													
Lys Pro Glu Ser Ala Thr Leu Glu Asp Leu Ala Leu Glu Leu Thr Thr														
130 135 140														
ATT TCC CAG GAG GCC AGG GGG ACC ATG GGG ACT TCC CAA GAG CCA CCG	538													
Ile Ser Gln Glu Ala Arg Gly Thr Met Gly Thr Ser Gln Glu Pro Pro														
145 150 155														
GCA GCA GTG ACC GGA TCA TCT CTC TCA ACT TCC GAG GCA CAG GAT GCA	586													
Ala Ala Val Thr Gly Ser Ser Leu Ser Thr Ser Glu Ala Gln Asp Ala														
160 165 170 175														
GGG CTT ACG GCT AAG CCT CAG AGC ATT GGA AGT TTT GAG GCG GCT GAC	634													
Gly Leu Thr Ala Lys Pro Gln Ser Ile Gly Ser Phe Glu Ala Ala Asp														
180 185 190														
ATC TCC ACC ACC GTT TGG CCG AGT CCT GCT GTC TAC CAA TCT GGA TCT	682													
Ile Ser Thr Thr Val Trp Pro Ser Pro Ala Val Tyr Gln Ser Gly Ser														
195 200 205														
AGC TCC TGG GCT GAG GAA AAA GCT ACT GAG TCC CCC TCC ACT ACA GCC	730													
Ser Ser Trp Ala Glu Glu Lys Ala Thr Glu Ser Pro Ser Thr Thr Ala														
210 215 220														
CCA TCT CCT CAG GTG TCC ACT ACT TCA CCT TCA ACC CCA GAG GAA AAT	778													
Pro Ser Pro Gln Val Ser Thr Thr Ser Pro Ser Thr Pro Glu Glu Asn														
225 230 235														
GTT GGG TCC GAA GGC CAA CCC CCA TGG GTC CAG GGA CAG GAC CTC AGT	826													
Val Gly Ser Glu Gly Gln Pro Pro Trp Val Gln Gly Gln Asp Leu Ser														
240 245 250 255														
CCA GAG AAG TCT CTA GGG TCT GAG GAG ATA AAC CCA GTT CAT ACT GAT	874													
Pro Glu Lys Ser Leu Gly Ser Glu Glu Ile Asn Pro Val His Thr Asp														
260 265 270														
AAT TTC CAG GAG AGG GGG CCT GGC AAC ACA GTC CAC CCC TCA GTG GCT	922													
Asn Phe Gln Glu Arg Gly Pro Gly Asn Thr Val His Pro Ser Val Ala														
275 280 285														
CCC ATC TCC TCT GAA GAG ACC CCC AGC CCA GAG CTG GTG GCC TCG GGC	970													
Pro Ile Ser Ser Glu Glu Thr Pro Ser Pro Glu Leu Val Ala Ser Gly														
290 295 300														
AGC CAG GCT CCT AAG ATA GAG GAA CCC ATC CAT GCC ACT GCA GAT CCC	1018													
Ser Gln Ala Pro Lys Ile Glu Glu Pro Ile His Ala Thr Ala Asp Pro														
305 310 315														
CAG AAA CTG AGT GTG CTT ATC ACT CCT GTC CCC GAC ACC CAG GCA GCC	1066													
Gln Lys Leu Ser Val Leu Ile Thr Pro Val Pro Asp Thr Gln Ala Ala														
320 325 330 335														
ACA AGG AGG CAG GCA GTG GGG CTA CTG GCT TTC CTT GGT CTT CTT TTC	1114													
Thr Arg Arg Gln Ala Val Gly Leu Leu Ala Phe Leu Gly Leu Leu Phe														
340 345 350														
TGC CTA GGG GTG GCC ATG TTT GCT TAC CAG AGC CTT CAG GGC TGT CCC	1162													
Cys Leu Gly Val Ala Met Phe Ala Tyr Gln Ser Leu Gln Gly Cys Pro														
355 360 365														
CGC AAA ATG GCG GGG GAA ATG GTA GAA GGC CTC CGC TAC GTC CCC CGT	1210													
Arg Lys Met Ala Gly Glu Met Val Glu Gly Leu Arg Tyr Val Pro Arg														
370 375 380														
AGC TGT GGC AGT AAC TCA TAC GTC CTG GTG CCA GTG TGA GCTGCTTGCC	1259													

TABLE 1-continued

Mouse CX3C chemokine nucleic acid (SEQ ID NO: 7) and amino acid (SEQ ID NO: 8) sequences.												
Ser	Cys	Gly	Ser	Asn	Ser	Tyr	Val	Leu	Val	Pro	Val	*
385							390				395	
TGCCTGCGCTG	TGTCCAGAGT	GTGATTTCGGA	CAGCTGTCTG	GGGACCCCCC	CCCATCCTCA	1319						
TACCCACCTT	CATCCACGCT	GGGAAATGG	GAATGGAGAA	GCTGGACCTC	CAGGGGCTGT	1379						
GGGCTCCATC	CAATCCCCCT	TCCCCGAGG	GGTGGCCCCG	GAGGCCACCC	TAGACCACTA	1439						
TTCACCTATC	AGAGACAGAG	CAGGTGACCT	TCCAGCTCCT	CTATATTTGA	AAGAATCCTC	1499						
TGCTGCTGGC	TGGTTAGAGG	GGCCCTTGAC	ACCCCAACTC	CAGTGAACAA	TTATTTATTG	1559						
GATTCCCAGC	CCCTGCGACG	ACACCTGTTT	CCCGCGCGCA	CCGTGGTCCG	CCCATATCAC	1619						
AAGCAGCAGG	CCAGGCCTAT	CTGCCTGTCC	CCCTGACCTC	CTTGTGTCTC	CTGGCTTTGC	1679						
TGCAGTCGCC	AGCCCTTCTC	CTCCCCGGCC	AGCCGCGGTG	CTATCTGCCC	TATGTCTCCC	1739						
TCTATCCCCCT	GTACAGAGCG	CACCACCATC	ACCATCAACA	CCGCTGTTGT	GTCTTTTCTT	1799						
GCATGAGGTT	AAAGCTGTGT	TTTCTGGAGC	TCTCCGGGAA	GGGAGACAAG	CTTGCAGAGG	1859						
GGTTTAAAGT	GTTCTCCCC	AGACTTGGAT	GTGCTGTGAG	GGCATGCTGC	GTCTGAAGGA	1919						
AGGGTCCAGT	CCCCACTCGG	CTACCAGCAC	CACAAAGTGC	CCCACCTGTA	AAAGGAAAGA	1979						
AACGTGGTCC	AGAGCTGGCA	ATAACCTATG	GCCCTGACAT	CATCACTTTC	TCTGAGATCC	2039						
TTGTCTCCAC	CCCTGGGTGC	AACCCACCC	CTTATCAACA	TTAATAGTCA	CTGCCATTCC	2099						
ACTGGACTGA	CATTTTTGTA	CCCTGTGATT	CTGAGGGCTG	GCAAGGAGTG	GCTTGAGAGT	2159						
GCAGATCGTA	CCCTGTATGC	CCCCCCCCAA	TGGAGCTGA	GTTGGGGACT	TGCAGGAACA	2219						
GAGGCCAACT	CAGATGGCTT	CCCCTGTGTT	CTCACTAGAA	ACCCCTCCCC	CATGCACCAA	2279						
GGTGACAGTC	ACAGTCTGC	CCTGGCTAAA	GGACAAGCCA	CATAGGAAAG	ATTAGGACAA	2339						
GCCCCTCGGA	GGCAGAGGAT	CCAGGGTAAA	CCCCTGGAGT	GGCCACAAAC	CCAATTTAG	2399						
TGTAGGGACT	TGTGCATGTG	TGTACTTGCA	TAGTCAGACA	GAGGCTGCCA	GGGTCCTTTC	2459						
CTGTCTCTGA	GAGCAGTGT	CACGCCAAGG	ACTCACCTTT	GCCCCATTG	CAGGCAGGGC	2519						
CAGAACTCCC	ATAGCATTCT	CCAAGAGCCC	TGTGACATTT	TCTGGAAGGA	ACTCTGCCCT	2579						
GGGCGCAAAG	TGACTGCTGA	AGCAAGGAGC	AGCTGAGCAG	CACCCAGCG	GAGCTGAGCC	2639						
GGCAGGCCAC	GCCCCTCGGG	GGGGGCATT	TCTACCCGCC	CTGCTCTGAA	TAGCTCCAAC	2699						
TTCACCTTAG	GAGCCTCCA	GGGCGAGCT	TCACCCAGAA	GCCAGTGACT	CACTCCTTGA	2759						
TTGGTGGAAG	CTCAGTTGGC	TCCTGAGAGT	GAGGAAGCCA	ACCCTTTGTC	GACCTCCTC	2819						
CTGGGAAGCC	TGTGGGCGGC	TCTGATCATG	CTCCACAGAA	CCAGTTGTAG	GCCTGAGCCG	2879						
CAGCAGCCCG	AGTGCACTAT	ATCCTGGCTC	CTTCGGTGGG	GAACCTTTAA	GGGTTGGGAC	2939						
ACCCGTATC	GGACTTTGTT	GGTTCTCCTC	TCCCAGAGCA	GAATGTGGGC	CGTAACAATC	2999						
TGAGGAGGAC	TTTAAAGTT	GTTGATCCTT	TAGGGTTTTT	TTTCAAGCAT	CATTACCAAT	3059						
GTCTGT						3065						

The coding sequence runs from nucleotides 62-1249.

The signal sequence runs from about Met1 through Gly24.

The mature polypeptide runs from about Gln25 through Val1395.

The chemokine domain runs from about Gln25 through Gly100.

The stalk region runs from about Gly101 through Gln339.

The transmembrane domain runs from about Ala340 through Phe358.

The cytoplasmic domain runs from about Ala359 through Val1395.

The CX3Ckine proteins of this invention are defined in part by their physicochemical and biological properties. The biological properties of the human and mouse CX3Ckines described herein, e.g., human CX3Ckine and mouse CX3Ckine, are defined by their amino acid sequence, and mature size. They also should share biological properties. The human and mouse CX3Ckine molecules exhibit about 70-80% amino acid identity, depending on whether the signal or mature sequences are compared. One of skill will readily recognize that some sequence variations may be tolerated, e.g., conservative substitutions or positions remote from the helical structures, without altering significantly the biological activity of the molecule.

Table 2 shows a sequence alignment of human CX3Ckine amino acid sequence (CX3C) with the C—X—C chemokine Gro α (Gro), the C chemokine lymphotactin (LTn), and the C—C chemokine Macrophage inflammatory protein 1 β (MIP-1 β).

TABLE 2

Comparison of various chemokines	
<u>Exon 1</u>	
Gro	(SEQ ID NO: 9) MIPATRSLLCAALLLATSRLATG
LTn	(SEQ ID NO: 10) MRLLLLTFLGVCCLTPWV
MIP-1 β	(SEQ ID NO: 11) MKLCVSALSLLLVAFCAPGFS
CX3	(SEQ ID NO: 2) MAPISLSWLLRLATFCHLTVLLAG
<u>Exon 2</u>	
Gro	APIANELRCQCLQTM .GIHLKNIQSLKVLPSGPHCTQT
LTn	EGVGTVELEESSCVNLQTRLPVQKIKTYIIWEG . . . AMR
MIP	APMGSDPPTSCCFSTARKLPRNFVVDYETSSL . . . CSQP
CX3	QHGHVTKCNITC .SKMTSKIPVALLIHYQQNQAS . . . CGKR
<u>Exon 3</u>	
GRO	EVIATLKNGREACLDPEAPLVQKIVQKMLKGVPK
LTN	AVIFVTKRGLKICADPEAKWVLAIAIKTVDGRAST RKNMAETVPGTGAQRSTSTAITLTG
MIP	AVVFQTKRSKQVCADPSESWVQEYVYDLELN
CX3	AIILETRQHLRFADCPKEQWVKDAMQHLDRQAAALTRNG . . .

CX3Ckines are present in specific tissue types, e.g., neural tissues, and the interaction of the protein with a receptor will be important for mediating various aspects of cellular physiology or development. The cellular types which express message encoding CX3Ckines suggest that signals important in cell differentiation and development are mediated by them. See, e.g., Gilbert (1991) *Developmental Biology* (3d ed.) Sinauer Associates, Sunderland, Mass.; Browder, et al. (1991) *Developmental Biology* (3d ed.) Saunders, Philadelphia, Pa.; Russo, et al. (1992) *Development: The Molecular Genetic Approach* Springer-Verlag, New York, N.Y.; and Wilkins (1993) *Genetic Analysis of Animal Development* (2d ed.) Wiley-Liss, New York, N.Y. Moreover, CX3Ckine expression should serve to define certain cell subpopulations.

The CX3C chemokine producing profile of various cells is elucidated herein. Screening a cDNA library generated from brain provided a novel cytokine, designated human CX3Ckine. Human CX3Ckine exhibits distant similarity to members of the C, C—C, and C—X—C chemokine families, with another heretofore unrecognized number of amino acid residues separating the characteristic cysteines in the motif which is peculiar to and partially defines chemokines. These observations suggest that the CX3Ckines represent novel additions to the chemokine superfamily.

CX3C chemokine protein biochemistry was assessed in mammalian expression systems. Human embryonic kidney

293 cells (HEK 293) transfected with a mammalian expression construct encoding full-length CX3C chemokine were metabolically labeled with ³⁵S cysteine and methionine. CX3C chemokine was produced as a protein of Mr ~95 kDa; control transfected supernatants contained no such species. Neuraminidase and glycosidases reduced the Mr of CX3C chemokine from ~95 kDa to ~45 kDa, suggesting that the recombinant form, is glycosylated substantially. CX3C chemokine cDNA, encoding a predicted membrane-bound protein, encodes a glycoprotein which is released from cells by an undefined mechanism.

The pro-migratory activities of CX3C chemokine have been assessed in microchemotaxis assays. CX3C chemokine appears to be a potent attractant of peripheral blood monocytes and T cells. Pro-migratory activity for blood neutrophils has been difficult to demonstrate.

The CX3C chemokine gene has been mapped to human chromosome 16. Mapping studies also indicate the possibility of a pseudogene or related gene on human chromosome 14. Sequencing of genomic DNA fragments suggests CX3C chemokine gene has an intron which begins near or in the codon encoding Ile 64. Other intron/exon boundaries have yet to be mapped, but such will be easily accomplished by standard methods.

The membrane bound form of CX3Ckine possesses proadherent properties for circulating T cells and monocytes. A secreted or soluble form, consisting of the chemokine domain and the stalk region, is able to inhibit this proadhesive activity. This suggests that the membrane bound form of CX3Ckine may be a potent regulator of circulating leukocytes, and thus may be involved in various inflammatory diseases, e.g., arthritis. The soluble form may be used as a regulator of proadherence, especially in conditions of compromised immune response.

CX3C chemokine's properties as a T cell and monocyte chemoattractant, coupled with its distribution in brain and other organs, suggests that CX3C chemokine may be involved in the pathogenesis of such CNS inflammatory disorders as multiple sclerosis, and other pathologies involving neurogenic inflammation. Since CX3C chemokine distribution is not limited to the brain, however, the entire spectrum of inflammatory, infectious, and immunoregulatory states thought to involve other chemokines must also now be considered for CX3C chemokine. See, e.g., Frank, Et al. (eds.) (1995) *Samter's Immunologic Diseases* 5th ed., vols. I and II, Little, Brown, and Co., Boston, Mass.

II. Definitions

The term "binding composition" refers to molecules that bind with specificity to a CX3Ckine, e.g., in an antibody-antigen interaction. However, other compounds, e.g., receptor proteins, may also specifically associate with CX3Ckines to the exclusion of other molecules. Typically, the association will be in a natural physiologically relevant protein-protein interaction, either covalent or non-covalent, and may include members of a multiprotein complex, including carrier compounds or dimerization partners. The molecule may be a polymer, or chemical reagent. No implication as to whether a CX3Ckine is either the ligand or the receptor of a ligand-receptor interaction is necessarily represented, other than whether the interaction exhibits similar specificity, e.g., specific affinity. A functional analog may be a ligand with structural modifications, or may be a wholly unrelated molecule, e.g., which has a molecular shape which interacts with the appropriate ligand binding determinants. The ligands may serve as agonists or antagonists of the receptor, see, e.g.,

Goodman, et al. (eds.) (1990) *Goodman & Gilman's: The Pharmacological Bases of Therapeutics* (8th ed.) Pergamon Press, Tarrytown, N.Y.

The term "binding agent: CX3Ckine protein complex", as used herein, refers to a complex of a binding agent and a CX3Ckine protein that is formed by specific binding of the binding agent to the CX3Ckine protein, e.g., preferably the chemokine domain. Specific binding of the binding agent means that the binding agent has a specific binding site that recognizes a site on the CX3Ckine protein. For example, antibodies raised to a CX3Ckine protein and recognizing an epitope on the CX3Ckine protein are capable of forming a binding agent: CX3Ckine protein complex by specific binding. Typically, the formation of a binding agent: CX3Ckine protein complex allows the measurement of CX3Ckine protein in a mixture of other proteins and biologics. The term "antibody: CX3Ckine protein complex" refers to an embodiment in which the binding agent is an antibody. The antibody may be monoclonal, polyclonal, or a binding fragment of an antibody, e.g., an Fab or F(ab)₂ fragment. The antibody will preferably be a polyclonal antibody for cross-reactivity purposes.

"Homologous" nucleic acid sequences, when compared, exhibit significant similarity. The standards for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison and/or phylogenetic relationship, or based upon hybridization conditions. Hybridization conditions are described in greater detail below.

An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially separated from other biologic components which naturally accompany a native sequence, e.g., proteins and flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogs, or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule. An isolated nucleic acid will usually contain homogeneous nucleic acid molecules, but will, in some embodiments, contain nucleic acids with minor sequence heterogeneity. This heterogeneity is typically found at the polymer ends or portions not critical to a desired biological function or activity.

As used herein, the term "CX3Ckine protein" shall encompass, when used in a protein context, a protein having amino acid sequences, particularly from the chemokine motif portions, shown in SEQ ID NO: 2, 4, 6, or 8, or a significant fragment of such a protein, e.g., preferably the chemokine domain. The invention also embraces a polypeptide which exhibits similar structure to human or mouse CX3Ckine, e.g., which interacts with CX3Ckine specific binding components. These binding components, e.g., antibodies, typically bind to a CX3Ckine with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM.

The term "polypeptide" or "protein" as used herein includes a significant fragment or segment of chemokine motif portion of a CX3Ckine, and encompasses a stretch of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally at least 12 amino acids, often at least 14 amino acids, more often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more preferably at least 28 amino acids, and, in par-

ticularly preferred embodiments, at least about 30 or more amino acids, e.g., 35, 40, 45, 50, 60, 70, 80, etc.

A "recombinant" nucleic acid is defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the nucleotide sequence, typically selection or production. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants. Thus, for example, products made by transforming cells with any non-naturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a sequence recognition site. Alternatively, it is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms. Restriction enzyme recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode polypeptides similar to fragments of these antigens, and fusions of sequences from various different species variants. Mutation of protease cleavage sites may also be accomplished.

"Solubility" is reflected by sedimentation measured in Svedberg units, which are a measure of the sedimentation velocity of a molecule under particular conditions. The determination of the sedimentation velocity was classically performed in an analytical ultracentrifuge, but is typically now performed in a standard ultracentrifuge. See, Freifelder (1982) *Physical Biochemistry* (2d ed.) W.H. Freeman & Co., San Francisco, Calif.; and Cantor and Schimmel (1980) *Biophysical Chemistry* parts 1-3, W.H. Freeman & Co., San Francisco, Calif. As a crude determination, a sample containing a putatively soluble polypeptide is spun in a standard full sized ultracentrifuge at about 50K rpm for about 10 minutes, and soluble molecules will remain in the supernatant. A soluble particle or polypeptide will typically be less than about 30S, more typically less than about 15S, usually less than about 10S, more usually less than about 6S, and, in particular embodiments, preferably less than about 4S, and more preferably less than about 3S. Solubility of a polypeptide or fragment depends upon the environment and the polypeptide. Many parameters affect polypeptide solubility, including temperature, electrolyte environment, size and molecular characteristics of the polypeptide, and nature of the solvent. Typically, the temperature at which the polypeptide is used ranges from about 4° C. to about 65° C. Usually the temperature at use is greater than about 18° C. and more usually greater than about 22° C. For diagnostic purposes, the temperature will usually be about room temperature or warmer, but less than the denaturation temperature of components in the assay. For therapeutic purposes, the temperature will usually be body temperature, typically about 37° C. for humans, though under certain situations the temperature may be raised or lowered in situ or in vitro.

The size and structure of the polypeptide should generally be evaluated in a substantially stable state, and usually not in

a denatured state. The polypeptide may be associated with other polypeptides in a quaternary structure, e.g., to confer solubility, or associated with lipids or detergents in a manner which approximates natural lipid bilayer interactions.

The solvent will usually be a biologically compatible buffer, of a type used for preservation of biological activities, and will usually approximate a physiological solvent. Usually the solvent will have a neutral pH, typically between about 5 and 10, and preferably about 7.5. On some occasions, a detergent will be added, typically a mild non-denaturing one, e.g., CHS (cholesteryl hemisuccinate) or CHAPS (3-[3-cholamidopropyl]dimethyl-ammonio]-1-propane sulfonate), or a low enough concentration as to avoid significant disruption of structural or physiological properties of the protein.

"Substantially pure" in a protein context typically means that the protein is isolated from other contaminating proteins, nucleic acids, and other biologicals derived from the original source organism. Purity, or "isolation" may be assayed by standard methods, and will ordinarily be at least about 50% pure, more ordinarily at least about 60% pure, generally at least about 70% pure, more generally at least about 80% pure, often at least about 85% pure, more often at least about 90% pure, preferably at least about 95% pure, more preferably at least about 98% pure, and in most preferred embodiments, at least 99% pure. Similar concepts apply, e.g., to antibodies or nucleic acids.

"Substantial similarity" in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 50% of the nucleotides, generally at least 56%, more generally at least 59%, ordinarily at least 62%, more ordinarily at least 65%, often at least 68%, more often at least 71%, typically at least 74%, more typically at least 77%, usually at least 80%, more usually at least about 85%, preferably at least about 90%, more preferably at least about 95 to 98% or more, and in particular embodiments, as high as about 99% or more of the nucleotides. Alternatively, substantial similarity exists when the segments will hybridize under selective hybridization conditions, to a strand, or its complement, typically using a sequence derived from SEQ ID NO: 1, 3, 5 or 7. Typically, selective hybridization will occur when there is at least about 55% similarity over a stretch of at least about 30 nucleotides, preferably at least about 65% over a stretch of at least about 25 nucleotides, more preferably at least about 75%, and most preferably at least about 90% over about 20 nucleotides. See Kanehisa (1984) *Nuc. Acids Res.* 12:203-213. The length of similarity comparison, as described, may be over longer stretches, and in certain embodiments will be over a stretch of at least about 17 nucleotides, usually at least about 20 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 40 nucleotides, preferably at least about 50 nucleotides, and more preferably at least about 75 to 100 or more nucleotides, e.g., 150, 200, etc.

"Stringent conditions", in referring to homology or substantial similarity in the hybridization context, will be stringent combined conditions of salt, temperature, organic solvents, and other parameters, typically those controlled in hybridization reactions. The combination of parameters is more important than the measure of any single parameter. Stringent temperature conditions will usually include temperatures in excess of about 30° C., more usually in excess of about 37° C., typically in excess of about 45° C., more typically in excess of about 55° C., preferably in excess of about 65° C., and more preferably in excess of about 70° C. Stringent salt conditions will ordinarily be less than about 1000

mM, usually less than about 500 mM, more usually less than about 400 mM, typically less than about 300 mM, preferably less than about 200 mM, and more preferably less than about 150 mM. See, e.g., Wetmur and Davidson (1968) *J. Mol. Biol.* 31:349-370. A nucleic acid probe which binds to a target nucleic acid under stringent conditions is specific for said target nucleic acid. Such a probe is typically more than 11 nucleotides in length, and is sufficiently identical or complementary to a target nucleic acid over the region specified by the sequence of the probe to bind the target under stringent hybridization conditions.

CX3Ckines from other mammalian species can be cloned and isolated by cross-species hybridization of closely related species. See, e.g., below. Similarity may be relatively low between distantly related species, and thus hybridization of relatively closely related species is advisable. Alternatively, preparation of an antibody preparation which exhibits less species specificity may be useful in expression cloning approaches.

The phrase "specifically binds to an antibody" or "specifically immunoreactive with", when referring to a protein or peptide, refers to a binding reaction which is determinative of the presence of the protein in the presence of a heterogeneous population of proteins and other biological components. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein and do not significantly bind other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, antibodies raised to the human CX3Ckine protein immunogen with the amino acid sequence depicted in SEQ ID NO: 2, 4, 6, or 8 can be selected to obtain antibodies specifically immunoreactive with CX3Ckine proteins and not with other proteins. These antibodies recognize proteins highly similar to the homologous mouse CX3Ckine protein.

III. Nucleic Acids

Human CX3Ckine is exemplary of a larger class of structurally and functionally related proteins. These soluble chemokine proteins will serve to transmit signals between different cell types. The preferred embodiments, as disclosed, will be useful in standard procedures to isolate genes from different individuals or other species, e.g., warm blooded animals, such as birds and mammals. Cross hybridization will allow isolation of related genes encoding proteins from individuals, strains, or species. A number of different approaches are available to successfully isolate a suitable nucleic acid clone based upon the information provided herein. Southern blot hybridization studies can qualitatively determine the presence of homologous genes in human, monkey, rat, dog, cow, and rabbit genomes under specific hybridization conditions.

Complementary sequences will also be used as probes or primers. Based upon identification of the likely amino terminus, other peptides should be particularly useful, e.g., coupled with anchored vector or poly-A complementary PCR techniques or with complementary DNA of other peptides. Moreover, reverse translation using the redundancy in the genetic code may provide synthetic genes which may encode essentially identical proteins often with a conformation selection preferred for expression in a given host cell.

Techniques for nucleic acid manipulation of genes encoding CX3Ckine proteins, such as subcloning nucleic acid sequences encoding polypeptides into expression vectors, labelling probes, DNA hybridization, and the like are

described generally in Sambrook, et al. (1989) *Molecular Cloning: A Laboratory Manual* (2nd ed.) Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY, which is incorporated herein by reference. This manual is hereinafter referred to as "Sambrook, et al."

There are various methods of isolating DNA sequences encoding CX3CKine proteins. For example, DNA is isolated from a genomic or cDNA library using labeled oligonucleotide probes having sequences identical or complementary to the sequences disclosed herein. Full-length probes may be used, or oligonucleotide probes may be generated by comparison of the sequences disclosed. Such probes can be used directly in hybridization assays to isolate DNA encoding CX3CKine proteins, or primers can be designed, e.g., using flanking sequence, for use in amplification techniques such as PCR, for the isolation of DNA encoding CX3CKine proteins.

To prepare a cDNA library, mRNA is isolated from cells which express a CX3CKine protein. cDNA is prepared from the mRNA and ligated into a recombinant vector. The vector is transfected into a recombinant host for propagation, screening, and cloning. Methods for making and screening cDNA libraries are well known. See Gubler and Hoffman (1983) *Gene* 25:263-269 and Sambrook, et al.

For a genomic library, the DNA can be extracted from tissue and either mechanically sheared or enzymatically digested to yield fragments, e.g., of about 12-20 kb. The fragments are then separated by gradient centrifugation and cloned in bacteriophage lambda vectors. These vectors and phage are packaged in vitro, as described in Sambrook, et al. Recombinant phage are analyzed by plaque hybridization as described in Benton and Davis (1977) *Science* 196:180-182. Colony hybridization is carried out as generally described in e.g., Grunstein, et al. (1975) *Proc. Natl. Acad. Sci. USA* 72:3961-3965.

DNA encoding a CX3CKine protein can be identified in either cDNA or genomic libraries by its ability to hybridize with the nucleic acid probes described herein, e.g., in colony or plaque hybridization assays. The corresponding DNA regions are isolated by standard methods familiar to those of skill in the art. See, e.g., Sambrook, et al. Alternatively, sequence databases, e.g., GenBank, may be evaluated for similar or corresponding sequences.

Various methods of amplifying target sequences, such as the polymerase chain reaction, can also be used to prepare DNA encoding CX3CKine proteins. Polymerase chain reaction (PCR) technology is used to amplify such nucleic acid sequences directly from mRNA, from cDNA, and from genomic libraries or cDNA libraries. The isolated sequences encoding CX3CKine proteins may also be used as templates for PCR amplification.

Typically, in PCR techniques, oligonucleotide primers complementary to two 5' regions in two strands of the DNA region to be amplified are synthesized. The polymerase chain reaction is then carried out using the two opposite primers. See Innis, et al. (eds.) (1990) *PCR Protocols: A Guide to Methods and Applications* Academic Press, San Diego, Calif. Primers can be selected to amplify the entire regions encoding a full-length CX3CKine protein or to amplify smaller DNA segments as desired. Once such regions are PCR-amplified, they can be sequenced and oligonucleotide probes can be prepared from sequence obtained using standard techniques. These probes can then be used to isolate DNA's encoding CX3CKine proteins.

Oligonucleotides for use as probes are usually chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage and Carruthers (1983) *Tetrahedron Lett.* 22(20):1859-1862, or using an auto-

mated synthesizer, as described in Needham-VanDevanter, et al. (1984) *Nucleic Acids Res.* 12:6159-6168. Purification of oligonucleotides is performed, e.g., by native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson and Regnier (1983) *J. Chrom.* 255:137-149. The sequence of the synthetic oligonucleotide can be verified using, e.g., the chemical degradation method of Maxam, A. M. and Gilbert, W. in Grossman, L. and Moldave (eds.) (1980) *Methods in Enzymology* 65:499-560 Academic Press, New York.

An isolated nucleic acid encoding a human CX3CKine protein was identified. The nucleotide sequence and corresponding open reading frame are provided in SEQ ID NO: 1 and 2; with further sequences provided in SEQ ID NO: 3 and 4. Correspondingly, a mouse sequence was identified and its nucleotide and corresponding open reading frame are provided as SEQ ID NO: 5-8.

These CX3CKines exhibit limited similarity to portions of chemokines, particularly the chemokine domains. See, e.g., Matsushima and Oppenheim (1989) *Cytokine* 1:2-13; Oppenheim, et al. (1991) *Ann. Rev. Immunol.* 9:617-648; Schall (1991) *Cytokine* 3:165-183; and Gronenborn and Clore (1991) *Protein Engineering* 4:263-269. In particular, the human CX3CKine shows similarity to the C class of chemokines in the carboxyl-terminal portion, particularly with respect to length, and at the positions corresponding, in the numbering of mature human sequence, to the cys-ala-aspro sequence at positions 50-53; and the trp-val at positions 57-58. CX3CKines have a much longer carboxyl terminal tail than the members of the CC or CXC chemokine families, and this "stalk" region may play a role in chemokine presentation. Notably, the spacing of conserved cysteine residues in the CXC and CC families of chemokines are absent in the human CX3CKine embodiment. Other features of comparison are apparent between the CX3CKine and chemokine families. See, e.g., Lodi, et al. (1994) *Science* 263:1762-1766. In particular, β -sheet and α -helix residues can be determined using, e.g., RASMOL program, see Sayle and Milner-White (1995) *TIBS* 20:374-376; or Gronenberg, et al. (1991) *Protein Engineering* 4:263-269; and other structural features are defined in Lodi, et al. (1994) *Science* 263:1762-1767. These secondary and tertiary features assist in defining further the C, CC, and CXC structural features, along with spacing of appropriate cysteine residues.

Based upon the structural modeling and insights in the binding regions of the chemokines, it is predicted that residues in the mature human protein, lacking a signal of 24 residues, 26 (his), 28 (gin), 40 (ile), 42 (glu), 47 (arg) and 48 (leu) should be important for chemokine binding to cells. Residues at the amino terminus are probably not involved in receptor binding or specificity.

Moreover, exon boundaries are predicted to correspond to protein segments including the signal sequence through about the second residue (his) in the mature protein; from there to about three residues past the third cys (around the arg-ala); and from there to the end. The third exon appears to exhibit relatively high similarity to the other chemokines. The second exon would probably be most characteristic of the CX3C chemokines, and would be the preferred segment to use to search for homology in other variants, e.g., species or otherwise. In particular, segments expected to be preferred in producing CX3C chemokine specific antibodies will include peptides or sequence in the region from the second residue of the mature protein (his) through about the third residue after the third cysteine (arg). Fragments of at least about 8-10 residues in that region would be especially interesting peptides, e.g., starting at residue positions of the mature 1, 2, 3,

etc. Those fragments would typically end in that region, e.g., at residue 37, 36, 35, etc. Other interesting peptides of various lengths would include ones which begin or end in other positions of the protein, e.g., at residues 87, 86, etc., with lengths ranging, e.g., from about 8 to 20, 25, 30, 35, 40, etc. Corresponding fragments of other mammalian CX3CKine, e.g., mouse, will be preferred embodiments.

This invention provides isolated DNA or fragments to encode a CX3CKine protein. In addition, this invention provides isolated or recombinant DNA which encodes a protein or polypeptide which is capable of hybridizing under appropriate conditions, e.g., high stringency, with the DNA sequences described herein. Said biologically active protein or polypeptide can be an intact ligand, or fragment, and have an amino acid sequence as disclosed in SEQ ID NO: 2, 4, 6, or 8. Preferred embodiments will be full length natural sequences, from isolates, e.g., about 11,000 to 12,500 daltons in size when unglycosylated, or fragments of at least about 6,000 daltons, more preferably at least about 8,000 daltons. In glycosylated form, the protein may exceed 12,500 daltons. Further, this invention contemplates the use of isolated or recombinant DNA, or fragments thereof, which encode proteins which are homologous to a CX3CKine protein or which were isolated using cDNA encoding a CX3CKine protein as a probe. The isolated DNA can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others.

IV. Making CX3CKines

DNAs which encode a CX3CKine or fragments thereof can be obtained by chemical synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples. The redundancy of the genetic code provides a number of polynucleotide sequences which should encode the same protein.

These DNAs can be expressed in a wide variety of host cells for the synthesis of a full-length protein or fragments which can in turn, e.g., be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified molecules; and for structure/function studies. Each CX3CKine or its fragments, e.g., the chemokine domain, can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially purified to be free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The antigen, e.g., CX3CKine, or portions thereof, may be expressed as fusions with other proteins or possessing an epitope tag. Such is applicable also to antigen binding sites.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired antigen gene or its fragments, usually operably linked to appropriate genetic control elements that are recognized in a suitable host cell. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription

and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently from the host cell.

The vectors of this invention encompass DNAs which encode a CX3CKine, or a fragment thereof, typically encoding, e.g., a biologically active polypeptide, or protein. The DNA can be under the control of a viral promoter and can encode a selection marker. This invention further contemplates use of such expression vectors which are capable of expressing eukaryotic cDNA coding for a CX3CKine protein in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNA coding for the protein is inserted into the vector such that growth of the host containing the vector expresses the cDNA in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of copies of the desirable gene per cell. It is not always necessary to require that an expression vector replicate in a host cell, e.g., it is possible to effect transient expression of the protein or its fragments in various hosts using vectors that do not contain a replication origin that is recognized by the host cell. It is also possible to use vectors that cause integration of a CX3CKine gene or its fragments into the host DNA by recombination, onto integrate a promoter which controls expression of an endogenous gene.

Vectors, as used herein, contemplate plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles which enable the integration of DNA fragments into the genome of the host. Expression vectors are specialized vectors which contain genetic control elements that effect expression of operably linked genes. Plasmids are the most commonly used form of vector, but many other forms of vectors which serve an equivalent function are suitable for use herein. See, e.g., Pouwels, et al. (1985 and Supplements) *Cloning Vectors: A Laboratory Manual* Elsevier, N.Y.; and Rodriguez, et al. (eds.) (1988) *Vectors: A Survey of Molecular Cloning Vectors and Their Uses* Butterworth, Boston, Mass.

Suitable host cells include prokaryotes, lower eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., *E. coli* and *B. subtilis*. Lower eukaryotes include yeasts, e.g., *S. cerevisiae* and *Pichia*, and species of the genus *Dictyostelium*. Higher eukaryotes include established tissue culture cell lines from animal cells, both of non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide variety of vectors for many different species. As used herein, *E. coli* and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for amplifying DNA is pBR322 or its derivatives. Vectors that can be used to express CX3CKines or CX3CKine fragments include, but are not limited to, such vectors as those containing the lac promoter (pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing Lambda-, trp-, lac-, and Ipp-derived Promoters", in Rodriguez and Denhardt (eds.) *Vectors: A Survey of Molecular Cloning Vectors and Their Uses* 10:205-236 Butterworth, Boston, Mass.

Lower eukaryotes, e.g., yeasts and *Dictyostelium*, may be transformed with CX3CKine sequence containing vectors. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, *Saccharomyces cerevisiae*. It will be used generically to represent lower eukaryotes although a number of other strains and species are also avail-

able. Yeast vectors typically consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the desired protein or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YEp-series); integrating types (such as the YIp-series), or mini-chromosomes (such as the YCp-series).

Higher eukaryotic tissue culture cells are typically the preferred host cells for expression of the functionally active CX3Ckine protein. In principle, many higher eukaryotic tissue culture cell lines may be used, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred to achieve proper processing, both cotranslationally and post-translationally. Transformation or transfection and propagation of such cells is routine. Useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (e.g., if genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also may contain a selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as from adenovirus, SV40, parvovirus, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pcDNA1; pCD, see Okayama, et al. (1985) *Mol. Cell Biol.* 5:1136-1142; pMCIneo Poly-A, see Thomas, et al. (1987) *Cell* 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

It is likely that CX3Ckines need not be glycosylated to elicit biological responses. However, it will occasionally be desirable to express a CX3Ckine polypeptide in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., in unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For example, the CX3Ckine gene may be co-transformed with one or more genes encoding mammalian or other glycosylating enzymes. It is further understood that over glycosylation may be detrimental to CX3Ckine biological activity, and that one of skill may perform routine testing to optimize the degree of glycosylation which confers optimal biological activity.

A CX3Ckine, or a fragment thereof, may be engineered to be phosphatidyl inositol (PI) linked to a cell membrane, but can be removed from membranes by treatment with a phosphatidyl inositol cleaving enzyme, e.g., phosphatidyl inositol phospholipase-C. This releases the antigen in a biologically active form, and allows purification by standard procedures of protein chemistry. See, e.g., Low (1989) *Biochem. Biophys. Acta* 988:427-454; Tse, et al. (1985) *Science* 230:1003-1008; and Brunner, et al. (1991) *J. Cell Biol.* 114:1275-1283.

Now that CX3Ckines have been characterized, fragments or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) *Solid*

Phase Peptide Synthesis Pierce Chemical Co., Rockford, Ill.; Bodanszky and Bodanszky (1984) *The Practice of Peptide Synthesis* Springer-Verlag, New York, N.Y.; and Bodanszky (1984) *The Principles of Peptide Synthesis* Springer-Verlag, New York, N.Y. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process (for example, p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative-reductive process, or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both applicable to the foregoing processes.

The prepared protein and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, for example, by extraction, precipitation, electrophoresis and various forms of chromatography, and the like. The CX3Ckines of this invention can be obtained in varying degrees of purity depending upon its desired use. Purification can be accomplished by use of known protein purification techniques or by the use of the antibodies or binding partners herein described, e.g., in immunoabsorbent affinity chromatography. See, e.g., Coligan, et al. (eds.) (1995 and periodic supplements) *Current Protocols in Protein Science*, John Wiley and Sons, New York, N.Y. This immunoabsorbent affinity chromatography is carried out by first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate source cells, lysates of other cells expressing the ligand, or lysates or supernatants of cells producing the CX3Ckines as a result of recombinant DNA techniques, see below.

Multiple cell lines may be screened for one which expresses a CX3Ckine at a high level compared with other cells. Various cell lines, e.g., a mouse thymic stromal cell line TA4, is screened and selected for its favorable handling properties. Natural CX3Ckines can be isolated from natural sources, or by expression from a transformed cell using an appropriate expression vector. Purification of the expressed protein is achieved by standard procedures, or may be combined with engineered means for effective purification at high efficiency from cell lysates or supernatants. Epitope or other tags, e.g., FLAG or His₆ segments, can be used for such purification features.

V. Antibodies

Antibodies can be raised to various CX3Ckines, including individual, polymorphic, allelic, strain, or species variants, and fragments thereof, both in their naturally occurring (full-length) forms and in their recombinant forms. Additionally, antibodies can be raised to CX3Ckines in either their active or native forms or in their inactive or denatured forms. Antidiotypic antibodies may also be used.

A. Antibody Production

A number of immunogens may be used to produce antibodies specifically reactive with CX3Ckine proteins. Recombinant protein is a preferred immunogen for the production of monoclonal or polyclonal antibodies. Naturally occurring protein may also be used either in pure or impure form. Synthetic peptides, made using the human or mouse CX3Ckine protein sequences described herein, may also be used as an immunogen for the production of antibodies to CX3Ckines, e.g., the chemokine domains thereof. Recombinant protein can be expressed in eukaryotic or prokaryotic cells as described herein, and purified as described. Naturally folded or denatured material can be used, as appropriate, for

producing antibodies. Either monoclonal or polyclonal antibodies may be generated for subsequent use in immunoassays to measure the protein.

Methods of producing polyclonal antibodies are known to those of skill in the art. Typically, an immunogen, preferably a purified protein, is mixed with an adjuvant and animals are immunized with the mixture. The animal's immune response to the immunogen preparation is monitored by taking test bleeds and determining the titer of reactivity to the CX3CKine protein or fragment of interest. When appropriately high titers of antibody to the immunogen are obtained, usually after repeated immunizations, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the protein can be done if desired. See, e.g., Harlow and Lane; or Coligan.

Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Typically, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (see, Kohler and Milstein (1976) *Eur. J. Immunol.* 6:511-519, incorporated herein by reference). Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according, e.g., to the general protocol outlined by Huse, et al. (1989) *Science* 246:1275-1281.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of CX3CKines can be raised by immunization of animals with conjugates of the fragments with carrier proteins as described above. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective CX3CKines, or screened for agonistic or antagonistic activity, e.g., mediated through a receptor. These monoclonal antibodies will usually bind with at least a K_D of about 1 mM, more usually at least about 300 μ M, typically at least about 10 μ M, more typically at least about 30 μ M, preferably at least about 10 μ M, and more preferably at least about 3 μ M or better.

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) *Basic and Clinical Immunology* (4th ed.) Lange Medical Publications, Los Altos, Calif., and references cited therein; Harlow and Lane (1988) *Antibodies: A Laboratory Manual* CSH Press; Goding (1986) *Monoclonal Antibodies: Principles and Practice* (2d ed.) Academic Press, New York, N.Y.; and particularly in Kohler and Milstein (1975) *Nature* 256:495-497, which discusses one method of generating monoclonal antibodies. Summarized briefly, this method involves injecting an animal with an immunogen. The animal is then sacrificed and cells taken from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized

and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance.

Other suitable techniques involve selection of libraries of antibodies in phage or similar vectors. See, e.g., Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," *Science* 246: 1275-1281; and Ward, et al. (1989) *Nature* 341:544-546. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents, teaching the use of such labels include U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant immunoglobulins may be produced. See, Cabilly, U.S. Pat. No. 4,816,567; and Queen, et al. (1989) *Proc. Nat'l Acad. Sci. USA* 86:10029-10033.

The antibodies of this invention are useful for affinity chromatography in isolating CX3CKine protein. Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, SEPHADEX, or the like, where a cell lysate or supernatant may be passed through the column, the column washed, followed by increasing concentrations of a mild denaturant, whereby purified CX3CKine protein will be released.

The antibodies may also be used to screen expression libraries for particular expression products. Usually the antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding.

Antibodies to CX3CKines may be used for the identification of cell populations expressing CX3CKines. By assaying the expression products of cells expressing CX3CKines it is possible to diagnose disease, e.g., immune-compromised conditions.

Antibodies raised against each CX3CKine will also be useful to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological conditions related to expression of the respective antigens.

B. Immunoassays

A particular protein can be measured by a variety of immunoassay methods. For a review of immunological and immunoassay procedures in general, see Stites and Terr (eds.) (1991) *Basic and Clinical Immunology* (7th ed.). Moreover, the immunoassays of the present invention can be performed in many configurations, which are reviewed extensively in Maggio (ed.) (1980) *Enzyme Immunoassay* CRC Press, Boca Raton, Fla.; Tijan (1985) "Practice and Theory of Enzyme Immunoassays," *Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers B.V., Amsterdam; and Harlow and Lane *Antibodies, A Laboratory Manual*, supra, each of which is incorporated herein by reference. See also Chan (ed.) (1987) *Immunoassay: A Practical Guide* Academic Press, Orlando, Fla.; Price and Newman (eds.) (1991) *Principles and Practice of Immunoassays* Stockton Press, NY; and Ngo (ed.) (1988) *Non-isotopic Immunoassays* Plenum Press, NY.

Immunoassays for measurement of CX3CKine proteins can be performed by a variety of methods known to those

skilled in the art. In brief, immunoassays to measure the protein can be competitive or noncompetitive binding assays. In competitive binding assays, the sample to be analyzed competes with a labeled analyte for specific binding sites on a capture agent bound to a solid surface. Preferably the capture agent is an antibody specifically reactive with CX3CKine proteins produced as described above. The concentration of labeled analyte bound to the capture agent is inversely proportional to the amount of free analyte present in the sample.

In a competitive binding immunoassay, the CX3CKine protein present in the sample competes with labeled protein for binding to a specific binding agent, for example, an antibody specifically reactive with the CX3CKine protein. The binding agent may be bound to a solid surface to effect separation of bound labeled protein from the unbound labeled protein. Alternatively, the competitive binding assay may be conducted in liquid phase and a variety of techniques known in the art may be used to separate the bound labeled protein from the unbound labeled protein. Following separation, the amount of bound labeled protein is determined. The amount of protein present in the sample is inversely proportional to the amount of labeled protein binding.

Alternatively, a homogeneous immunoassay may be performed in which a separation step is not needed. In these immunoassays, the label on the protein is altered by the binding of the protein to its specific binding agent. This alteration in the labeled protein results in a decrease or increase in the signal emitted by label, so that measurement of the label at the end of the immunoassay allows for detection or quantitation of the protein.

CX3CKine proteins may also be determined by a variety of noncompetitive immunoassay methods. For example, a two-site, solid phase sandwich immunoassay may be used. In this type of assay, a binding agent for the protein, for example an antibody, is attached to a solid support. A second protein binding agent, which may also be an antibody, and which binds the protein at a different site, is labelled. After binding at both sites on the protein has occurred, the unbound labeled binding agent is removed and the amount of labeled binding agent bound to the solid phase is measured. The amount of labeled binding agent bound is directly proportional to the amount of protein in the sample.

Western blot analysis can be used to determine the presence of CX3CKine proteins in a sample. Electrophoresis is carried out, for example, on a tissue sample suspected of containing the protein. Following electrophoresis to separate the proteins, and transfer of the proteins to a suitable solid support, e.g., a nitrocellulose filter, the solid support is incubated with an antibody reactive with the protein. This antibody may be labeled, or alternatively may be detected by subsequent incubation with a second labeled antibody that binds the primary antibody.

The immunoassay formats described above employ labeled assay components. The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. A wide variety of labels and methods may be used. Traditionally, a radioactive label incorporating ^3H , ^{125}I , ^{35}S , ^{14}C , or ^{32}P was used. Non-radioactive labels include ligands which bind to labeled antibodies, fluorophores, chemiluminescent agents, enzymes, and antibodies which can serve as specific binding pair members for a labeled ligand. The choice of label depends on sensitivity required, ease of conjugation with the compound, stability requirements, and available instrumentation. For a review of various labelling or signal producing systems which may be used, see U.S. Pat. No. 4,391,904, which is incorporated herein by reference.

Antibodies reactive with a particular protein can also be measured by a variety of immunoassay methods. For a review of immunological and immunoassay procedures applicable to the measurement of antibodies by immunoassay techniques, see Stites and Terr (eds.) *Basic and Clinical Immunology* (7th ed.) supra; Maggio (ed.) *Enzyme Immunoassay*, supra; and Harlow and Lane *Antibodies, A Laboratory Manual*, supra.

In brief, immunoassays to measure antisera reactive with CX3CKine proteins can be competitive or noncompetitive binding assays. In competitive binding assays, the sample analyte competes with a labeled analyte for specific binding sites on a capture agent bound to a solid surface. Preferably the capture agent is a purified recombinant CX3CKine protein produced as described above. Other sources of CX3CKine proteins, including isolated or partially purified naturally occurring protein, may also be used. Noncompetitive assays include sandwich assays, in which the sample analyte is bound between two analyte-specific binding reagents. One of the binding agents is used as a capture agent and is bound to a solid surface. The second binding agent is labeled and is used to measure or detect the resultant complex by visual or instrument means. A number of combinations of capture agent and labelled binding agent can be used. A variety of different immunoassay formats, separation techniques, and labels can be also be used similar to those described above for the measurement of CX3CKine proteins.

VI. Purified CX3CKines

Human CX3CKine amino acid sequences are provided in SEQ ID NO: 2 and 4. Mouse nucleotide and amino acid sequences are provided in SEQ ID NO: 5, 6, 7, and 8.

Purified protein or defined peptides are useful for generating antibodies by standard methods, as described above. Synthetic peptides or purified protein, e.g., the chemokine domains, can be presented to an immune system to generate polyclonal and monoclonal antibodies. See, e.g., Coligan (1991) *Current Protocols in Immunology* Wiley/Greene, N.Y.; and Harlow and Lane (1989) *Antibodies: A Laboratory Manual* Cold Spring Harbor Press, NY, which are incorporated herein by reference. Alternatively, a CX3CKine receptor can be useful as a specific binding reagent, and advantage can be taken of its specificity of binding, for, e.g., purification of a CX3CKine ligand.

The specific binding composition can be used for screening an expression library made from a cell line which expresses a CX3CKine. Many methods for screening are available, e.g., standard staining of surface expressed ligand, or by panning. Screening of intracellular expression can also be performed by various staining or immunofluorescence procedures. The binding compositions could be used to affinity purify or sort out cells expressing the ligand.

The peptide segments, along with comparison to homologous genes, can also be used to produce appropriate oligonucleotides to screen a library. The genetic code can be used to select appropriate oligonucleotides useful as probes for screening. In combination with polymerase chain reaction (PCR) techniques, synthetic oligonucleotides will be useful in selecting desired clones from a library, including natural allelic or polymorphic variants.

The peptide sequences allow preparation of peptides to generate antibodies to recognize such segments, and allow preparation of oligonucleotides which encode such sequences. The sequence also allows for synthetic preparation, e.g., see Dawson, et al. (1994) *Science* 266:776-779.

Since CX3CKines appear to be soluble proteins, the gene will normally possess an N-terminal signal sequence, which is removed upon processing and secretion, and the putative cleavage site is between amino acids 24 (gly) and 25 (gln) in SEQ ID NO: 2 or 4, though it may be slightly in either direction. Analysis of the structural features in comparison with the most closely related reported sequences has revealed similarities with other cytokines, particularly the class of proteins known as chemokines. Within the chemokines are two subgroups, the CC and CXC subgroups. The CX3CKine family shares various features with each of these groups, but its combination of features is distinctive and defines a new family of related chemokines.

While further structural features result from the sequences provided in SEQ ID NO: 1 through 8, the "chemokine on a stick" feature is provided through the stalk region which possesses many sites which may provide a heavily glycosylated domain. The stalk structure may be important in CX3C chemokine presentation to other cells. In fact, it appears that the stalk region may be processed to release the soluble chemokine. This suggests the possibility of substituting the CX3C chemokine domain with other chemokines to effect efficient presentation to appropriate target cells.

In addition, the "stalk" regions are likely to affect solubility and pharmacological aspects of the protein. As such, this region will be the target of analysis to evaluate and modulate such features as pharmacokinetics. Truncation of that portion may affect half-life, clearance, and accessibility of the chemokine domains.

VII. Physical Variants

This invention also encompasses proteins or peptides having substantial amino acid sequence similarity with an amino acid sequence of a CX3CKine. Natural variants include individual, polymorphic, allelic, strain, or species variants.

Amino acid sequence similarity, or sequence identity, is determined by optimizing residue matches, if necessary, by introducing gaps as required. This changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences include natural polymorphic, allelic, and interspecies variations in each respective protein sequence. Typical homologous proteins or peptides will have from 50-100% similarity (if gaps can be introduced), to 75-100% similarity (if conservative substitutions are included) with the amino acid sequence of the CX3CKine. Similarity measures will be at least about 50%, generally at least 60%, more generally at least 65%, usually at least 70%, more usually at least 75%, preferably at least 80%, and more preferably at least 80%, and in particularly preferred embodiments, at least 85% or more. See also Needleham, et al. (1970) *J. Mol. Biol.* 48:443-453; Sankoff, et al. (1983) *Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison* Chapter One, Addison-Wesley, Reading, Mass.; and software packages from IntelliGenetics, Mountain View, Calif.; and the University of Wisconsin Genetics Computer Group, Madison, Wis.

Nucleic acids encoding mammalian CX3CKine proteins will typically hybridize to the nucleic acid sequence of SEQ ID NO: 1, 3, 5 or 7 under stringent conditions. For example, nucleic acids encoding human CX3CKine proteins will normally hybridize to the nucleic acid of SEQ ID NO: 1 under stringent hybridization conditions. Generally, stringent con-

ditions are selected to be about 10° C. lower than the thermal melting point (T_m) for the probe sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about 0.2 molar at pH 7 and the temperature is at least about 50° C. Other factors may significantly affect the stringency of hybridization, including, among others, base composition and size of the complementary strands, the presence of organic solvents such as formamide, and the extent of base mismatching. A preferred embodiment will include nucleic acids which will bind to disclosed sequences in 50% formamide and 200 mM NaCl at 42° C.

An isolated CX3CKine DNA can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and short inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode CX3CKine antigens, their derivatives, or proteins having highly similar physiological, immunogenic, or antigenic activity.

Modified sequences can be used to produce mutant antigens or to enhance expression. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant CX3CKine derivatives include predetermined or site-specific mutations of the respective protein or its fragments. "Mutant CX3CKine" encompasses a polypeptide otherwise falling within the homology definition of the human CX3CKine as set forth above, but having an amino acid sequence which differs from that of a CX3CKine as found in nature, whether by way of deletion, substitution, or insertion. In particular, "site specific mutant CX3CKine" generally includes proteins having significant similarity with a protein having a sequence of SEQ ID NO: 2, 4, 6, or 8, and as sharing various biological activities, e.g., antigenic or immunogenic, with those sequences, and in preferred embodiments contain most or all of the disclosed sequence. This applies also to polymorphic variants from different individuals. Similar concepts apply to different CX3CKine proteins, particularly those found in various warm blooded animals, e.g., mammals and birds. As stated before, it is emphasized that descriptions are generally meant to encompass other CX3CKine proteins, not limited to the human or mouse embodiments specifically discussed.

Although site specific mutation sites are predetermined, mutants need not be site specific. CX3CKine mutagenesis can be conducted by making amino acid insertions or deletions. Substitutions, deletions, insertions, or any combinations may be generated to arrive at a final construct. These include amino acid residue substitution levels from none, one, two, three, five, seven, ten, twelve, fifteen, etc. Insertions include amino- or carboxyl-terminal fusions, e.g. epitope tags. Random mutagenesis can be conducted at a target codon and the expressed mutants can then be screened for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis or polymerase chain reaction (PCR) techniques. See also, Sambrook, et al. (1989) and Ausubel, et al. (1987 and Supplements). The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these proteins, both the CX3CKine, or antigen binding sites. A heterologous fusion protein is a fusion of proteins or seg-

ments which are naturally not normally fused in the same manner. Thus, the fusion product of an immunoglobulin with a CX3Ckine polypeptide is a continuous protein molecule having sequences fused in a typical peptide linkage, typically made as a single translation product and exhibiting properties derived from each source peptide. A similar concept applies to heterologous nucleic acid sequences.

In addition, new constructs may be made from combining similar functional domains from other proteins. For example, protein-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) *Science* 243:1330-1336; and O'Dowd, et al. (1988) *J. Biol. Chem.* 263:15985-15992. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of protein-binding specificities and other functional domains.

VIII. Binding Agent: CX3Ckine Protein Complexes

A CX3Ckine protein that specifically binds to or that is specifically immunoreactive with an antibody generated against a defined immunogen, such as an immunogen consisting of the amino acid sequence of SEQ ID NO: 2, 4, 6, or 8, is typically determined in an immunoassay. The immunoassay uses a polyclonal antiserum which was raised to a protein of SEQ ID NO: 4, 6, or 8. This antiserum is selected to have low crossreactivity against other chemokines and any such crossreactivity is removed by immunoabsorption prior to use in the immunoassay.

In order to produce antisera for use in an immunoassay, the protein of SEQ ID NO: 2, 4, 6, or 8, is isolated as described herein. For example, recombinant protein may be produced in a mammalian cell line. An inbred strain of mice such as balb/c is immunized with the protein of SEQ ID NO: 2, 4, 6, or 8, using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see Harlow and Lane, supra). Alternatively, a synthetic peptide, preferably near full length, derived from the sequences disclosed herein and conjugated to a carrier protein can be used as an immunogen. Polyclonal sera are collected and titered against the immunogen protein in an immunoassay, for example, a solid phase immunoassay with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of 10^4 or greater are selected and tested for their cross reactivity against C, C—C, and CXC chemokines, using a competitive binding immunoassay such as the one described in Harlow and Lane, supra, at pages 570-573. Preferably two chemokines are used in this determination in conjunction with either human CX3Ckine or mouse CX3Ckine.

In conjunction with a CX3Ckine, the monocyte chemotactic protein-1 (MCP-1) and macrophage inflammatory protein-1 α (Mip-1 α) are used to identify antibodies which are specifically bound by a CX3Ckine. In conjunction with human CX3Ckine, the monocyte chemotactic protein-2 (MCP-2) and Mip-1 α are used to identify antibodies which are specifically bound by a CX3Ckine. These chemokines can be produced as recombinant proteins and isolated using standard molecular biology and protein chemistry techniques as described herein.

Immunoassays in the competitive binding format can be used for the crossreactivity determinations. For example, a protein of SEQ ID NO: 2, 4, 6, or 8 can be immobilized to a solid support. Proteins added to the assay compete with the binding of the antisera to the immobilized antigen. The ability of the above proteins to compete with the binding of the antisera to the immobilized protein is compared to the protein of SEQ ID NO: 2, 4, 6, or 8. The percent crossreactivity for the above proteins is calculated, using standard calculations.

Those antisera with less than 10% crossreactivity with each of the proteins listed above are selected and pooled. The cross-reacting antibodies are then removed from the pooled antisera by immunoabsorption with the above-listed proteins.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein to the immunogen protein (e.g., the CX3Ckine chemokine motif of SEQ ID NO: 2, 4, 6, or 8). In order to make this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required is less than twice the amount of the protein of SEQ ID NO: 2 that is required, then the second protein is said to specifically bind to an antibody generated to the immunogen.

It is understood that CX3Ckine proteins are a family of homologous proteins that comprise two or more genes. For a particular gene product, such as the human CX3Ckine protein, the term refers not only to the amino acid sequences disclosed herein, but also to other proteins that are polymorphic, allelic, non-allelic, or species variants. It is also understood that the term "human CX3Ckine" or "mouse CX3Ckine" includes nonnatural mutations introduced by deliberate mutation using conventional recombinant technology such as single site mutation, or by excising short sections of DNA encoding CX3Ckine proteins, or by substituting new amino acids, or adding new amino acids. Such minor alterations must substantially maintain the immunoidentity of the original molecule and/or its biological activity. Thus, these alterations include proteins that are specifically immunoreactive with a designated naturally occurring CX3Ckine protein, for example, the human CX3Ckine protein shown in SEQ ID NO: 2 or 4. The biological properties of the altered proteins can be determined by expressing the protein in an appropriate cell line and measuring, e.g., a chemotactic effect. Particular protein modifications considered minor would include conservative substitution of amino acids with similar chemical properties, as described above for the CX3Ckine family as a whole. By aligning a protein optimally with the protein of SEQ ID NO: 2, 4, 6, or 8, and by using the conventional immunoassays described herein to determine immunoidentity, or by using lymphocyte chemotaxis assays, one can determine the protein compositions of the invention.

IX. Functional Variants

The blocking of physiological response to CX3Ckines may result from the inhibition of binding of the protein to its receptor, e.g., through competitive inhibition. Thus, in vitro assays of the present invention will often use isolated protein, membranes from cells expressing a recombinant membrane associated CX3Ckine, soluble fragments comprising receptor binding segments of these proteins, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either binding segment mutations and modifications, or protein mutations and modifications, e.g., protein analogs. This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to antigen or receptor fragments compete with a test compound for binding to the protein. In this manner, the antibodies can be used to detect the presence of a polypeptide which shares one or more antigenic binding sites of the protein and can also be used to occupy binding sites on the protein that might otherwise interact with a receptor.

"Derivatives" of CX3Ckine antigens include amino acid sequence mutants, glycosylation variants, and covalent or aggregate conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in CX3Ckine amino acid side chains or at the N- or C-termini, by means which are well known in the art. These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or amino-group containing residues, e.g., lysine or arginine. Acyl groups are selected from the group of alkyl-moieties including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species. See, e.g., Coligan, et al. (eds.) (1995 and periodic supplements) *Current Protocols in Protein Science*, John Wiley and Sons, New York, N.Y. Covalent attachment to carrier proteins may be important when immunogenic moieties are haptens.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine, or other moieties, including ribosyl groups or cross-linking reagents.

A major group of derivatives are covalent conjugates of the CX3Ckine or fragments thereof with other proteins or polypeptides. These derivatives can be synthesized in recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred protein derivatization sites with cross-linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between CX3Ckines and other homologous or heterologous proteins are also provided. Many growth factors and cytokines are homodimeric entities, and a repeat construct may have various advantages, including lessened susceptibility to proteolytic degradation. Moreover, many receptors require dimerization to transduce a signal, and various dimeric proteins or domain repeats can be desirable. Heterologous polypeptides may be fusions between different surface markers, resulting in, e.g., a hybrid protein exhibiting receptor binding specificity. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a protein, e.g., a receptor-binding segment, so that the presence or location of the fused protein may be easily determined. See, e.g., Dull, et al., U.S. Pat. No. 4,859,609. Other gene fusion partners include bacterial β -galactosidase, trpE, Protein A, β -lactamase, alpha amylase, alcohol dehydrogenase, and yeast alpha mating factor. See, e.g., See, e.g., Dawson, et al. (1994) *Science* 266:776-779; and Godowski, et al. (1988) *Science* 241: 812-816. In particular, fusion proteins with portions from the related genes will be useful. Similar concepts of fusions with antigen binding sites are contemplated.

Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, or the addition or removal of other moieties, particularly those which have molecular shapes

similar to phosphate groups. In some embodiments, the modifications will be useful labeling reagents, or serve as purification targets, e.g., affinity ligands.

This invention also contemplates the use of derivatives of CX3Ckines other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives include: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption complexes, for example with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of ligands or other binding ligands. For example, a CX3Ckine antigen can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated SEPHAROSE, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of anti-CX3Ckine antibodies or its receptor. The CX3Ckines can also be labeled with a detectable group, e.g., radioiodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to another fluorescent moiety for use in diagnostic assays. Purification of CX3Ckines may be effected by immobilized antibodies or receptor.

Isolated CX3Ckine genes will allow transformation of cells lacking expression of corresponding CX3Ckines, e.g., either species types or cells which lack corresponding proteins and exhibit negative background activity. Expression of transformed genes will allow isolation of antigenically pure cell lines, with defined or single specie variants. This approach will allow for more sensitive detection and discrimination of the physiological effects of CX3Ckine receptor proteins. Subcellular fragments, e.g., cytoplasts or membrane fragments, can be isolated and used.

X. Uses

The present invention provides reagents which will find use in diagnostic applications as described elsewhere herein, e.g., in the general description for developmental abnormalities, or below in the description of kits for diagnosis.

CX3Ckine nucleotides, e.g., human or mouse CX3Ckine DNA or RNA, may be used as a component in a forensic assay. For instance, the nucleotide sequences provided may be labeled using, e.g., ^{32}P or biotin and used to probe standard restriction fragment polymorphism blots, providing a measurable character to aid in distinguishing between individuals. Such probes may be used in well-known forensic techniques such as genetic fingerprinting. In addition, nucleotide probes made from CX3Ckine sequences may be used in *in situ* assays to detect chromosomal abnormalities. For instance, rearrangements in the mouse chromosome encoding a CX3Ckine gene may be detected via well-known *in situ* techniques, using CX3Ckine probes in conjunction with other known chromosome markers.

Antibodies and other binding agents directed towards CX3Ckine proteins or nucleic acids may be used to purify the corresponding CX3Ckine molecule. As described in the Examples below, antibody purification of CX3Ckine components is both possible and practicable. Antibodies and other binding agents may also be used in a diagnostic fashion to determine whether CX3Ckine components are present in a tissue sample or cell population using well-known techniques described herein. The ability to attach a binding agent to a CX3Ckine provides a means to diagnose disorders associated with CX3Ckine misregulation. Antibodies and other

CX3CKine binding agents may also be useful as histological markers. As described in the examples below, CX3CKine expression is limited to specific tissue types. By directing a probe, such as an antibody or nucleic acid to a CX3CKine it is possible to use the probe to distinguish tissue and cell types in situ or in vitro.

This invention also provides reagents with significant therapeutic value. The CX3CKines (naturally occurring or recombinant), fragments thereof, and antibodies thereto, along with compounds identified as having binding affinity to a CX3CKine, are useful in the treatment of conditions associated with abnormal physiology or development, including abnormal proliferation, e.g., cancerous conditions, or degenerative conditions. Abnormal proliferation, regeneration, degeneration, and atrophy may be modulated by appropriate therapeutic treatment using the compositions provided herein. For example, a disease or disorder associated with abnormal expression or abnormal signaling by a CX3CKine is a target for an agonist or antagonist of the protein. The proteins likely play a role in regulation or development of neuronal or hematopoietic cells, e.g., lymphoid cells, which affect immunological responses.

Other abnormal developmental conditions are known in cell types shown to possess CX3CKine mRNA by northern blot analysis. See Berkow (ed.) *The Merck Manual of Diagnosis and Therapy*, Merck & Co., Rahway, N.J.; and Thorn, et al. *Harrison's Principles of Internal Medicine*, McGraw-Hill, NY. Developmental or functional abnormalities, e.g., of the neuronal or immune system, cause significant medical abnormalities and conditions which may be susceptible to prevention or treatment using compositions provided herein.

Certain chemokines have also been implicated in viral replication mechanisms. See, e.g., Cohen (1996) *Science* 272:809-810; Feng, et al. (1996) *Science* 272:872-877; and Cocchi, et al. (1995) *Science* 270:1811-1816. The CX3C chemokine may be useful in a similar context. Alternatively, the stalk structure may be very important in presentation of the ligand domain, and other chemokines may be advantageously substituted for the chemokine domain in this molecule. Modification in the "stalk" structure may affect many of the pharmacological properties of the CX3CKine, including half-life and biological activity.

Recombinant CX3CKine or CX3CKine antibodies can be purified and then administered to a patient, e.g., in sterile form. These reagents can be combined for therapeutic use with additional active or inert ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, e.g., immunogenic adjuvants, along with physiologically innocuous stabilizers and excipients. These combinations can be sterile filtered and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding fragments thereof, including forms which are not complement binding.

Drug screening using antibodies or receptor or fragments thereof can identify compounds having binding affinity to CX3CKines, including isolation of associated components. Subsequent biological assays can then be utilized to determine if the compound has intrinsic stimulating activity and is therefore a blocker or antagonist in that it blocks the activity of the protein. Likewise, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of a CX3CKine. This invention further contemplates the therapeutic use of antibodies to CX3CKines as antagonists. This approach should be particularly useful with other CX3CKine species variants.

The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds.) (1990) *Goodman and Gilman's: The Pharmacological Bases of Therapeutics* (8th ed.) Pergamon Press; and (1990) *Remington's Pharmaceutical Sciences* (17th ed.) Mack Publishing Co., Easton, Pa. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the *Merck Index*, Merck & Co., Rahway, N.J. Dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10 μ M concentrations, usually less than about 100 nM, preferably less than about 10 μ M (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or a slow release apparatus will often be utilized for continuous administration.

CX3CKines, fragments thereof, and antibodies to it or its fragments; antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in many conventional dosage formulations. While it is possible for the active ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations typically comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier should be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds.) (1990) *Goodman and Gilman's: The Pharmacological Bases of Therapeutics* (8th ed.) Pergamon Press; and (1990) *Remington's Pharmaceutical Sciences* (17th ed.) Mack Publishing Co., Easton, Pa.; Avis, et al. (eds.) (1993) *Pharmaceutical Dosage Forms: Parenteral Medications* Dekker, NY; Lieberman, et al. (eds.) (1990) *Pharmaceutical Dosage Forms: Tablets* Dekker, NY; and Lieberman, et al. (eds.) (1990) *Pharmaceutical Dosage Forms: Disperse Systems* Dekker, NY. The therapy of this invention may be combined with or used in association with other therapeutic agents.

Both the naturally occurring and the recombinant forms of the CX3CKines of this invention are particularly useful in kits and assay methods which are capable of screening compounds for binding activity to the proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds in a short period. See, e.g., Fodor, et al. (1991) *Science* 251:767-773, and other descriptions of chemical diversity libraries, which describe means for testing of binding affinity by a plurality of compounds. The development of suitable assays

can be greatly facilitated by the availability of large amounts of purified, soluble CX3CKine as provided by this invention.

For example, antagonists can normally be found once the protein has been structurally defined. Testing of potential protein analogs is now possible upon the development of highly automated assay methods using a purified receptor. In particular, new agonists and antagonists will be discovered by using screening techniques described herein. Of particular importance are compounds found to have a combined binding affinity for multiple CX3CKine receptors, e.g., compounds which can serve as antagonists for species variants of a CX3CKine.

This invention is particularly useful for screening compounds by using recombinant protein in a variety of drug screening techniques. The advantages of using a recombinant protein in screening for specific ligands include: (a) improved renewable source of the CX3CKine from a specific source; (b) potentially greater number of ligands per cell giving better signal to noise ratio in assays; and (c) species variant specificity (theoretically giving greater biological and disease specificity).

One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant DNA molecules expressing a CX3CKine receptor. Cells may be isolated which express a receptor in isolation from any others. Such cells, either in viable or fixed form, can be used for standard ligand/receptor binding assays. See also, Parce, et al. (1989) *Science* 246:243-247; and Owicki, et al. (1990) *Proc. Nat'l Acad. Sci. USA* 87:4007-4011, which describe sensitive methods to detect cellular responses. Competitive assays are particularly useful, where the cells (source of CX3CKine) are contacted and incubated with a labeled receptor or antibody having known binding affinity to the ligand, such as ¹²⁵I-antibody, and a test sample whose binding affinity to the binding composition is being measured. The bound and free labeled binding compositions are then separated to assess the degree of ligand binding. The amount of test compound bound is inversely proportional to the amount of labeled receptor binding to the known source. Any one of numerous techniques can be used to separate bound from free ligand to assess the degree of ligand binding. This separation step could typically involve a procedure such as adhesion to filters followed by washing, adhesion to plastic followed by washing, or centrifugation of the cell membranes. Viable cells could also be used to screen for the effects of drugs on CX3CKine mediated functions, e.g., second messenger levels, i.e., Ca⁺⁺; cell proliferation; inositol phosphate pool changes; and others. Some detection methods allow for elimination of a separation step, e.g., a proximity sensitive detection system. Calcium sensitive dyes will be useful for detecting Ca⁺⁺ levels, with a fluorimeter or a fluorescence cell sorting apparatus.

Another method utilizes membranes from transformed eukaryotic or prokaryotic host cells as the source of a CX3CKine. These cells are stably transformed with DNA vectors directing the expression of a CX3CKine, e.g., an engineered membrane bound form. Essentially, the membranes would be prepared from the cells and used in a receptor/ligand binding assay such as the competitive assay set forth above.

Still another approach is to use solubilized, unpurified or solubilized, purified CX3CKine from transformed eukaryotic or prokaryotic host cells. This allows for a "molecular" binding assay with the advantages of increased specificity, the ability to automate, and high drug test throughput.

Another technique for drug screening involves an approach which provides high throughput screening for compounds having suitable binding affinity to a CX3CKine antibody and

is described in detail in Geysen, European Patent Application 84/03564, published on Sep. 13, 1984. First, large numbers of different small peptide test compounds are synthesized on a solid substrate, e.g., plastic pins or some other appropriate surface, see Fodor, et al., supra. Then all the pins are reacted with solubilized, unpurified or solubilized, purified CX3CKine antibody, and washed. The next step involves detecting bound CX3CKine antibody.

Rational drug design may also be based upon structural studies of the molecular shapes of the CX3CKine and other effectors or analogs. See, e.g., *Methods in Enzymology* vols 202 and 203. Effectors may be other proteins which mediate other functions in response to ligand binding, or other proteins which normally interact with the receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) *Protein Crystallography* Academic Press, NY.

A purified CX3CKine can be coated directly onto plates for use in the aforementioned drug screening techniques. However, non-neutralizing antibodies to these ligands can be used as capture antibodies to immobilize the respective ligand on the solid phase.

XI. Kits

This invention also contemplates use of CX3CKine proteins, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting the presence of CX3CKine or a CX3CKine receptor. Typically the kit will have a compartment containing either a defined CX3CKine peptide or gene segment or a reagent which recognizes one or the other, e.g., receptor fragments or antibodies.

A kit for determining the binding affinity of a test compound to a CX3CKine would typically comprise a test compound; a labeled compound, e.g., a receptor or antibody having known binding affinity for the CX3CKine; a source of CX3CKine (naturally occurring or recombinant); and a means for separating bound from free labeled compound, such as a solid phase for immobilizing the CX3CKine. Once compounds are screened, those having suitable binding affinity to the CX3CKine can be evaluated in suitable biological assays, as are well known in the art, to determine whether they act as agonists or antagonists to the receptor. The availability of recombinant CX3CKine polypeptides also provide well defined standards for calibrating such assays.

A preferred kit for determining the concentration of, for example, a CX3CKine in a sample would typically comprise a labeled compound, e.g., receptor or antibody, having known binding affinity for the CX3CKine, a source of CX3CKine (naturally occurring or recombinant), and a means for separating the bound from free labeled compound, for example, a solid phase for immobilizing the CX3CKine. Compartments containing reagents, and instructions, will normally be provided.

Antibodies, including antigen binding fragments, specific for the CX3CKine or ligand fragments are useful in diagnostic applications to detect the presence of elevated levels of CX3CKine and/or its fragments. Such may allow diagnosis of the amounts of differently processed forms of the CX3CKine, e.g., successively degraded stalk structure. Such diagnostic assays can employ lysates, live cells, fixed cells, immunofluorescence, cell cultures, body fluids, and further can involve

the detection of antigens related to the ligand in serum, or the like. Diagnostic assays may be homogeneous (without a separation step between free reagent and antigen-CX3Ckine complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA), and the like. For example, unlabeled antibodies can be employed by using a second antibody which is labeled and which recognizes the antibody to a CX3Ckine or to a particular fragment thereof. Similar assays have also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) *Antibodies: A Laboratory Manual*, CSH Press, NY; Chan (ed.) (1987) *Immunoassay: A Practical Guide* Academic Press, Orlando, Fla.; Price and Newman (eds.) (1991) *Principles and Practice of Immunoassay* Stockton Press, NY; and Ngo (ed.) (1988) *Nonisotopic Immunoassay* Plenum Press, NY.

Anti-idiotypic antibodies may have similar use to diagnose presence of antibodies against a CX3Ckine, as such may be diagnostic of various abnormal states. For example, overproduction of CX3Ckine may result in production of various immunological or other medical reactions which may be diagnostic of abnormal physiological states, e.g., in cell growth, activation, or differentiation.

Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody or receptor, or labeled CX3Ckine is provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium providing appropriate concentrations of reagents for performing the assay.

Many of the aforementioned constituents of the drug screening and the diagnostic assays may be used without modification, or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, the protein, test compound, CX3Ckine, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as ¹²⁵I, enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating the bound from the free ligand, or alternatively the bound from the free test compound. The CX3Ckine can be immobilized on various matrices followed by washing. Suitable matrices include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the CX3Ckine to a matrix include, without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin-avidin. The last step in this approach involves the precipitation of ligand/receptor or ligand/antibody complex by any of several methods including those utilizing, e.g., an organic solvent such as polyethylene

glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) *Clin. Chem.* 30:1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678.

Methods for linking proteins or their fragments to the various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form, peptide bonds, the formation of thioethers by reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves use of oligonucleotide or polynucleotide sequences taken from the sequence of a CX3Ckine. These sequences can be used as probes for detecting levels of the CX3Ckine message in samples from natural sources, or patients suspected of having an abnormal condition, e.g., cancer or developmental problem. The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least about 18 nucleotides, and the polynucleotide probes may be up to several kilobases. Various labels may be employed, most commonly radionuclides, particularly ³²P. However, other techniques may also be employed, such as using biotin modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorophores, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected. The use of probes to the novel anti-sense RNA may be carried out using many conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART). This also includes amplification techniques such as polymerase chain reaction (PCR).

Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) *Progress in Growth Factor Res.* 1:89-97.

XII. Receptor Isolation

Having isolated a binding partner of a specific interaction, methods exist for isolating the counter-partner. See, Gearing, et al. (1989) *EMBO J.* 8:3667-3676. For example, means to label a CX3Ckine without interfering with the binding to its receptor can be determined. For example, an affinity label or epitope tag can be fused to either the amino- or carboxyl-terminus of the ligand. An expression library can be screened for specific binding of the CX3Ckine, e.g., by cell sorting, or other screening to detect subpopulations which express such a binding component. See, e.g., Ho, et al. (1993) *Proc. Nat'l*

Acad. Sci. USA 90:11267-11271. Alternatively, a panning method may be used. See, e.g., Seed and Aruffo (1987) *Proc. Nat'l Acad. Sci. USA* 84:3365-3369. A two-hybrid selection system may also be applied making appropriate constructs with the available BAS-1 sequences. See, e.g., Fields and Song (1989) *Nature* 340:245-246.

Protein cross-linking techniques with label can be applied to isolate binding partners of a CX3Ckine. This would allow identification of proteins which specifically interact with a CX3Ckine, e.g., in a ligand-receptor like manner. Typically, the chemokine family binds to receptors of the seven transmembrane receptor family, and the receptor for the CX3Ckine is likely to exhibit a similar structure. Thus, it is likely that the receptor will be found by expression in a system which is capable of expressing such a membrane protein in a form capable of exhibiting ligand binding capability.

The broad scope of this invention is best understood with reference to the following examples, which are not intended to limit the invention to specific embodiments.

EXAMPLES

I. General Methods

Many of the standard methods below are described or referenced, e.g., in Maniatis, et al. (1982) *Molecular Cloning, A Laboratory Manual* Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY; Sambrook, et al. (1989) *Molecular Cloning: A Laboratory Manual* (2d ed.) Vols. 1-3, CSH Press, NY; Ausubel, et al., *Biology* Greene Publishing Associates, Brooklyn, N.Y.; or Ausubel, et al. (1987 and Supplements) *Current Protocols in Molecular Biology* Wiley/Greene, NY; Innis, et al. (eds.) (1990) *PCR Protocols: A Guide to Methods and Applications* Academic Press, NY. Methods for protein purification include such methods as ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Deutscher (1990) "Guide to Protein Purification," *Methods in Enzymology* vol. 182, and other volumes in this series; and manufacturer's literature on use of protein purification products, e.g., Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, Calif. Combination with recombinant techniques allow fusion to appropriate segments (epitope tags), e.g., to a FLAG sequence or an equivalent which can be fused, e.g., via a protease-removable sequence. See, e.g., Hochuli (1989) *Chemische Industrie* 12:69-70; Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) *Genetic Engineering, Principle and Methods* 12:87-98, Plenum Press, NY; Crowe, et al. (1992) *QIAexpress: The High Level Expression & Protein Purification System* QIAGEN, Inc., Chatsworth, Calif.; and Coligan, et al. (eds.) (1995 and periodic supplements) *Current Protocols in Protein Science*, John Wiley and Sons, New York, N.Y.

Standard immunological techniques are described, e.g., in Coligan (1991) *Current Protocols in Immunology* Wiley/Greene, NY; and *Methods in Enzymology* volumes. 70, 73, 74, 84, 92, 93, 108, 116, 121, 132, 150, 162, and 163. Assays for neural cell biological activities are described, e.g., in Wouterlood (ed. 1995) *Neuroscience Protocols* modules 10, Elsevier; *Methods in Neurosciences* Academic Press; and *Neuromethods* Humana Press, Totowa, N.J. Methodology of developmental systems is described, e.g., in Meisami (ed.) *Handbook of Human Growth and Developmental Biology* CRC Press; and Chrispeels (ed.) *Molecular Techniques and Approaches in Developmental Biology* Interscience.

FACS analyses are described in Melamed, et al. (1990) *Flow Cytometry and Sorting* Wiley-Liss, Inc., New York, N.Y.; Shapiro (1988) *Practical Flow Cytometry* Liss, New York, N.Y.; and Robinson, et al. (1993) *Handbook of Flow Cytometry Methods* Wiley-Liss, New York, N.Y.

II. Isolation of Human CX3Ckine Clone

A clone encoding the human CX3Ckine is isolated from a natural source by many different possible methods. Given the sequences provided herein, PCR primers or hybridization probes are selected and/or constructed to isolate either genomic DNA segments or cDNA reverse transcripts. Appropriate cell sources include human tissues, e.g., brain libraries. Tissue distribution below also suggests source tissues. Genetic and polymorphic or allelic variants are isolated by screening a population of individuals.

PCR based detection is performed by standard methods, preferably using primers from opposite ends of the coding sequence, but flanking segments might be selected for specific purposes.

Alternatively, hybridization probes are selected. Particular AT or GC contents of probes are selected depending upon the expected homology and mismatching expected. Appropriate stringency conditions are selected to balance an appropriate positive signal to background ratio. Successive washing steps are used to collect clones of greater homology.

Further clones are isolated using an antibody based selection procedure. Standard expression cloning methods are applied including, e.g., FACS staining of membrane associated expression product. The antibodies are used to identify clones producing a recognized protein. Alternatively, antibodies are used to purify a CX3C chemokine, with protein sequencing and standard means to isolate a gene encoding that protein.

Genomic sequence based methods will also allow for identification of sequences naturally available, or otherwise, which exhibit homology to the provided sequences. Similar procedures will allow isolation of other primate genes.

III. Isolation of Rodent CX3Ckine Clone

Similar methods are used as above to isolate an appropriate mouse CX3C chemokine gene. Similar source materials as indicated above are used to isolate natural genes, including genetic, polymorphic, allelic, or strain variants. Species variants are also isolated using similar methods, e.g., from rats, moles, muskrats, copybaras, etc.

IV. Isolation of an Avian CX3Ckine Clone

An appropriate avian source is selected as above. Similar methods are utilized to isolate a species variant, though the level of similarity will typically be lower for avian CX3C chemokine as compared to a human to mouse sequence.

V. Expression; Purification; Characterization

With an appropriate clone from above, the coding sequence is inserted into an appropriate expression vector. This may be in a vector specifically selected for a prokaryote, yeast, insect, or higher vertebrate, e.g., mammalian expression system. Standard methods are applied to produce the gene product, preferably as a soluble secreted molecule, but will, in certain instances, also be made as an intracellular protein. Intracellular proteins typically require cell lysis to recover the pro-

tein, and insoluble inclusion bodies are a common starting material for further purification.

With a clone encoding a vertebrate CX3C chemokine, recombinant production means are used, although natural forms may be purified from appropriate sources. The protein product is purified by standard methods of protein purification, in certain cases, e.g., coupled with immunoaffinity methods. Immunoaffinity methods are used either as a purification step, as described above, or as a detection assay to determine the separation properties of the protein.

Preferably, the protein is secreted into the medium, and the soluble product is purified from the medium in a soluble form. Alternatively, as described above, inclusion bodies from prokaryotic expression systems are a useful source of material. Typically, the insoluble protein is solubilized from the inclusion bodies and refolded using standard methods. Purification methods are developed as described above.

In certain embodiments, the protein is made in a eukaryotic cell which glycosylates the protein normally. The purification methods may be affected thereby, as may biological activities. The intact protein can be processed to release the chemokine domain, probably due to a protease cleavage event somewhere in the glycosylated stalk region close to the chemokine/stalk boundary. While recombinant protein appears to be processed, the physiological processes which normally do such in native cells remain to be determined.

The product of the purification method described above is characterized to determine many structural features. Standard physical methods are applied, e.g., amino acid analysis and protein sequencing. The resulting protein is subjected to CD spectroscopy and other spectroscopic methods, e.g., NMR, ESR, mass spectroscopy, etc. The product is characterized to determine its molecular form and size, e.g., using gel chromatography and similar techniques. Understanding of the chromatographic properties will lead to more gentle or efficient purification methods.

CX3C chemokine protein biochemistry was assessed in mammalian expression systems. Human embryonic kidney 293 cells (HEK 293) transfected with a mammalian expression construct encoding full-length CX3C chemokine were metabolically labeled with ³⁵S cysteine and methionine. CX3C chemokine was produced as a protein of Mr ~95 kDa; control transfected supernatants contained no such species. Neuraminidase and glycosidases reduced the Mr of CX3C chemokine from ~95 kDa to ~45 kDa, suggesting that the recombinant form is glycosylated substantially. Thus CX3C chemokine cDNA, encoding a predicted membrane-bound protein, encodes a glycoprotein which is released from cells by an undefined mechanism.

Prediction of glycosylation sites may be made, e.g., as reported in Hansen, et al. (1995) *Biochem. J.* 308:801-813.

VI. Preparation of Antibodies Against Vertebrate CX3Ckine

With protein produce, as above, animals are immunized to produce antibodies. Polyclonal antiserum is raised using non-purified antigen, though the resulting serum will exhibit higher background levels. Preferably, the antigen is purified using standard protein purification techniques, including, e.g., affinity chromatography using polyclonal serum indicated above. Presence of specific antibodies is detected using defined synthetic peptide fragments. Preferred fragments include the chemokine domain.

Polyclonal serum is raised against a purified antigen, purified as indicated above, or using synthetic peptides. A series of overlapping synthetic peptides which encompass all of the

full length sequence, if presented to an animal, will produce serum recognizing most linear epitopes on the protein. Such an antiserum is used to affinity purify protein, which is, in turn, used to introduce intact full length protein into another animal to produce another antiserum preparation.

Similar techniques are used to generate induce monoclonal antibodies to either unpurified antigen, or, preferably, purified antigen.

VII. Cellular and Tissue Distribution

Distribution of the protein or gene products are determined, e.g., using immunohistochemistry with an antibody reagent, as produced above, or by screening for nucleic acids encoding the chemokine. Either hybridization or PCR methods are used to detect DNA, cDNA, or message content. Histochemistry allows determination of the specific cell types within a tissue which express higher or lower levels of message or DNA. Antibody techniques are useful to quantitate protein in a biological sample, including a liquid or tissue sample. Immunoassays are developed to quantitate protein.

Hybridization techniques were applied to the tissue types in Table 3 with positive or negative results, as indicated. The commercial tissue blots may have cellular contamination from resident cells, e.g., from blood or other cells which populate the tissue. The large and small transcripts correspond to sizes about 4 kb and less than about 2 kb, respectively.

TABLE 3

Tissue and cell distribution of human CX3Ckine gene.		
Commercial tissue library:		
cell type	large	small
spleen	-	-
thymus	+	-
prostate	+	+
testis	+	-
ovary	+	-
small intestine	+	+
colon	+	+
peripheral blood	-	-

Further analysis of tissue distribution indicates abundance of human message: heart +++; brain +++; placenta -; lung ++; liver -; muscle +; kidney -; pancreas +; spleen -; thymus +; prostate ++; testis +; ovary +; small intestine ++; colon ++; peripheral blood -; HL60 promyelocytic leukemia line -; HeLa cell S3 -; K562 chronic myelogenous leukemia line -; Molt4 lymphoblastic leukemia line -; Burkitts lymphoma RAJ1 line -; SW480 colorectal adenocarcinoma line +; A549 lung carcinoma line -; and G361 melanoma line -.

"Reverse northern" are blots from cDNA libraries with the inserts removed, and the size determinations are based upon the size of inserts in the cDNA library, and reflect the lengths found in the cDNA library inserts, which may be less than full length where the reverse transcription was not full length. As such, size determinations there are not reflective of the natural sizes. The results of these are: PBMC (peripheral blood mononuclear cells) +; PBMC (activated using T cell stimulation conditions, with anti-CD3 and PMA) -; Mot72 (resting Th0 clone) +; Mot 72 (activated with anti-CD28 and anti-CD3) -; Mot72 α (activated with anti-peptide, anergic clone) -; Mot81 (resting Th0 clone) -; Mot81 (activated with anti-CD28 and anti-CD3) -; HY06 (resting Th1 clone) -; HY06 (activated with anti-CD28 and anti-CD3) -; HY06 α

(activated with anti-peptide, anergic clone) -; HY935 (resting Th2 clone) -; HY935 (activated with anti-CD28 and anti-CD3) +; BC pool of EBV transformed lines +; resting splenocytes +; splenocytes + (activated using B cell stimulating conditions, with anti-CD40 and IL-4) -; NK cell pool -; NK pool (activated 6 h with PMA and ionomycin) +; NKA6 NK cell clone -; NKB1 NK cell clone -; NK non-cytotoxic cell clone +; and NK clone stimulated to be cytotoxic Other cells and tissues: CHO cells +; Jurkat cells (DNAX) +; Jurkat cells (another source) +; normal T cell pool +; TCT pool (transformed T cells) -; fetal kidney -; fetal lung -; fetal liver -; fetal heart -; fetal brain +; fetal gall bladder +; fetal small intestine +; fetal adipose +; fetal ovary -; fetal uterus +; adult placenta -; fetal testis +; fetal spleen +; and fetal brain +. Additional cells provided: U937 (resting monocyte cell line) +; C- (elutriated monocyte activated with LPS, IFN- γ , and anti-IL-10) +; C+ (elutriated monocytes activated with LPS, IFN- γ , and IL-10) +; M1 (elutriated monocytes activated with LPS 1 h) +; M6 (elutriated monocytes activated with LPS 6h) +; 30% DC (resting 30% CD1a+ dendritic cells, proliferated in TNF- α and GM-CSF) +; 70% DC (resting 70% CD1a+ dendritic cells, proliferated in TNF- α and GM-CSF) +; D1 (dendritic cells stimulated 1 h in PMA and ionomycin) -; D6 (dendritic cells stimulated 6 h in PMA and ionomycin) -; D5 DC (resting dendritic cells cultured 5 d in GM-CSF and IL-4) +; DC (dendritic cells cultured in GM-CSF and IL-4, LPS activated) +; DC (GM-CSF activated, like D5 cells) +; DC mix (dendritic cells stimulated with a mixture of cytokines) +; CD1a+ (99% pure CD1a+ dendritic cells, enriched from 70% DC) +; CD14+ (CD14+ fraction sorted from 70% DC, monocyte-like morphology) -; CD1Aa+ (95% CD1a+ and CD86+ sorted from 70% DC) -; TF1 (hematopoietic precursor line) +; Jurkat (T cell line) +; MRCS (lung fibroblast sarcoma cell line)+; JY (B cell line) +; U937 (pre-monocytic cell line) +.

Since the endothelium is a major site of chemokine action, a northern blot was performed to ascertain if CX3Ckine was expressed in this tissue. Human CX3Ckine was also shown to be expressed on human activated primary endothelial cells by both mRNA and protein expression. This suggests that CX3Ckine may be involved in leukocyte trafficking in various organs.

In summary, human CX3Ckine mRNA is found in monocytes, dendritic cells, T cells and B cells, e.g., found in certain immune cells.

VIII. Microchemotaxis Assays

The pro-migratory activities of CX3C chemokine have been assessed in microchemotaxis assays. See, e.g., Bacon, et al. (1988) *Br. J. Pharmacol.* 95:966-974. CX3C chemokine appears to be a potent attractant of peripheral blood monocytes and T cells. Pro-migratory activity for blood neutrophils has been difficult to demonstrate.

IX. Chromosomal Mapping

The CX3C chemokine gene has been mapped to human chromosome 16. A BIOS Laboratories (New Haven, Conn.) mouse somatic cell hybrid panel was combined with PCR. These mapping studies also indicate the possibility of a pseudogene or related gene on human chromosome 14. Sequencing of genomic DNA fragments suggests CX3C chemokine gene has an intron which begins near or in the codon encoding Ile 64. Other intron/exon boundaries have yet to be mapped. This location is distinct from the chromosomal

mapping locations of the other C, CC, or CXC chemokine families, consistent with the CX3Ckine being a separate gene family within the chemokines.

X. Biological Activities, Direct and Indirect

The 293 human embryonic kidney cell line (HEK 293) was transfected with either the membrane bound form of human CX3Ckine (293-CX3Ckine), the chemokine domain plus the "stalk" region, or a control vector without an insert. The transfected cells were subsequently cultured with either monocytes, T cells, or peripheral mononuclear (PMN) cells to assay relative adherence of these cells to CX3Ckine. Specifically, 5×10^4 cells per well of HEK 293 transfected cells were seeded in a 96 well plate. 2×10^5 monocytes, T cells, or PMNs, metabolically labeled with ^{35}S -methionine and cysteine (Amersham, Arlington Heights, Ill.), were added to each well. The plate was then incubated at 37° C. for various time points. The wells were washed 2 times RPMI supplemented with 1% FCS. Plates were then read in a Millipore Cytofluor at 485/530 nm.

In all cases, adherence to HEK 293 cells transfected with the membrane bound form of CX3Ckine was significantly enhanced when compared to the truncated CX3Ckine or mock transfected cells. Interestingly, only the membrane bound form possessed this proadhesive activity, leading to the conclusion that CX3Ckine, in its membrane bound form, may serve as a regulator of circulating leukocytes.

In another experiment, the recombinant soluble form of the chemokine domain of CX3Ckine (rCx3C) was added to HEK 293-CX3C cells and monocytes at a concentration of 1 μM per well, and assayed as described above. rCx3C was able to antagonize adhesion of monocytes to HEK 293-CX3C cells. A similar experiment was performed to investigate the effect on T cell adherence. Comparable results were obtained. Thus rCx3C may function as a negative regulator of circulating leukocytes.

A comparison of three different forms of human CX3Ckine was performed to analyze variations in chemoattractant activity that may be due to the structure of CX3Ckine. CX3C 1.7 (chemokine domain plus the entire stalk region), CX3C 0.7 (chemokine domain plus one-half stalk region), and CX3C CK (chemokine domain only) were subjected to the chemotactic assay described above, their ability to attract T cells was analyzed. CX3C 1.7 displayed a slightly better dose dependent ability to attract T cells relative to the other forms of CX3Ckine.

A robust and sensitive assay is selected as described above, e.g., on immune cells, neuronal cells, or stem cells. Chemokine is added to the assay in increasing doses to see if a dose response is detected. For example, in a proliferation assay, cells are plated out in plates. Appropriate culture medium is provided, and chemokine is added to the cells in varying amounts. Growth is monitored over a period of time which will detect either a direct effect on the cells, or an indirect effect of the chemokine.

Alternatively, an activation assay or attraction assay is used. An appropriate cell type is selected, e.g., hematopoietic cells, myeloid (macrophages, neutrophils, polymorphonuclear cells, etc.) or lymphoid (T cell, B cell, or NK cells), neural cells (neurons, neuroglia, oligodendrocytes, astrocytes, etc.), or stem cells, e.g., progenitor cells which differentiate to other cell types, e.g., gut crypt cells and undifferentiated cell types.

Other assays will be those which have been demonstrated with other chemokines. See, e.g., Schall and Bacon (1994) *Current Opinion in Immunology* 6:865-873; and Bacon and

Schall (1996) *Int. Arch. Allergy & Immunol.* 109:97-109. Effects of truncated stalk structures will be similarly evaluated.

XI. Structure Activity Relationship

Information on the criticality of particular residues is determined using standard procedures and analysis. Standard mutagenesis analysis is performed, e.g., by generating many different variants at determined positions, e.g., at the positions identified above, and evaluating biological activities of the variants. This may be performed to the extent of determining positions which modify activity, or to focus on specific positions to determine the residues which can be substituted to either retain, block, or modulate biological activity.

Alternatively, analysis of natural variants can indicate what positions tolerate natural mutations. This may result from populational analysis of variation among individuals, or across strains or species. Samples from selected individuals are analysed, e.g., by PCR analysis and sequencing. This allows evaluation of population polymorphisms. Particularly, as described above, many of the biological activities of the chemokine domain attached to different portions or extents of the stalk structure may result.

XII. Screening for Agonists/Antagonists

Agonists or antagonists are screened for ability to induce or block biological activity. The candidate compounds, e.g., sequence variants of natural CX3CKines, are assayed for their biological activities. Alternatively, compounds are screened, alone or in combinations, to determine effects on biological activity.

XIII. Isolation of a Receptor for CX3C Chemokine

Based on the proadherent properties of CX3CKine, 7 transmembrane G-protein receptor was found to be expressed by monocytes and T cells. It was also discovered that the chemokine domain is the only region of CX3CKine that can engage the receptor. Binding assays with known chemokine receptor revealed that CX3CKine does not engage chemokine receptors CCR 1 through 5, CXCR 1 and 2, or the Duffy antigen receptor. CX3CKine can, however, bind to a virally encoded chemokine receptor, CMV-US28.

Alternatively, CX3C chemokine can be used as a specific binding reagent to identify its binding partner, by taking advantage of its specificity of binding, much like an antibody would be used. A binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a substrate for panning methods. The typical chemokine receptor is a seven transmembrane receptor.

The purified protein is also be used to identify other binding partners of CX3CKine as described, e.g., in Fields and Song (1989) *Nature* 340:245-246.

The binding composition, e.g., chemokine, is used to screen an expression library made from a cell line which expresses a binding partner, i.e. receptor. Standard staining techniques are used to detect or sort intracellular or surface expressed receptor, or surface expressing transformed cells are screened by panning. Screening of intracellular expression is performed by various staining or immunofluorescence procedures. See also McMahan, et al. (1991) *EMBO J.* 10:2821-2832.

For example, on day 0, precoat 2-chamber permanox slides with 1 ml per chamber of fibronectin, 10 ng/ml in PBS, for 30 min at room temperature. Rinse once with PBS. Then plate

COS cells at 2-3x10⁵ cells per chamber in 1.5 ml of growth media. Incubate overnight at 37° C.

On day 1 for each sample, prepare 0.5 ml of a solution of 66 µg/ml DEAE-dextran, 66 µM chloroquine, and 4 µg DNA in serum free DME. For each set, a positive control is prepared, e.g., of human CX3C chemokine cDNA at 1 and 1/200 dilution, and a negative mock. Rinse cells with serum free DME. Add the DNA solution and incubate 5 hr at 37° C. Remove the medium and add 0.5 ml 10% DMSO in DME for 2.5 min. Remove and wash once with DME. Add 1.5 ml growth medium and incubate overnight.

On day 2, change the medium. On days 3 or 4, the cells are fixed and stained. Rinse the cells twice with Hank's Buffered Saline Solution (HBSS) and fix in 4% paraformaldehyde (PFA)/glucose for 5 min. Wash 3x with HBSS. The slides may be stored at -80° C. after all liquid is removed. For each chamber, 0.5 ml incubations are performed as follows. Add HBSS/saponin (0.1%) with 32 µl/ml of 1 M NaN₃ for 20 min. Cells are then washed with HBSS/saponin 1x. Add chemokine or chemokine/antibody complex to cells and incubate for 30 min. Wash cells twice with HBSS/saponin. If appropriate, add first antibody for 30 min. Add second antibody, e.g., Vector anti-mouse antibody, at 1/200 dilution, and incubate for 30 min. Prepare ELISA solution, e.g., Vector Elite ABC horseradish peroxidase solution, and preincubate for 30 min. Use, e.g., 1 drop of solution A (avidin) and 1 drop solution B (biotin) per 2.5 ml HBSS/saponin. Wash cells twice with HBSS/saponin. Add ABC HRP solution and incubate for 30 min. Wash cells twice with HBSS, second wash for 2 min, which closes cells. Then add Vector diaminobenzoic acid (DAB) for 5 to 10 min. Use 2 drops of buffer plus 4 drops DAB plus 2 drops of H₂O₂ per 5 ml of glass distilled water. Carefully remove chamber and rinse slide in water. Air dry for a few minutes, then add 1 drop of Crystal Mount and a cover slip. Bake for 5 min at 85-90° C.

Evaluate positive staining of pools and progressively subclone to isolation of single genes responsible for the binding.

Alternatively, chemokine reagents are used to affinity purify or sort out cells expressing a receptor. See, e.g., Sambrook, et al. or Ausubel, et al.

Another strategy is to screen for a membrane bound receptor by panning. The receptor cDNA is constructed as described above. The ligand can be immobilized and used to immobilize expressing cells. Immobilization may be achieved by use of appropriate antibodies which recognize, e.g., a FLAG sequence of a chemokine fusion construct, or by use of antibodies raised against the first antibodies. Recursive cycles of selection and amplification lead to enrichment of appropriate clones and eventual isolation of receptor expressing clones.

Phage expression libraries can be screened by chemokine. Appropriate label techniques, e.g., anti-FLAG antibodies, will allow specific labeling of appropriate clones.

SEQUENCES

SEQ ID NO: 1 is human CX3CKine nucleotide sequence.

SEQ ID NO: 2 is human CX3CKine amino acid sequence.

SEQ ID NO: 3 is more complete human CX3CKine nucleotide sequence.

SEQ ID NO: 4 is more complete human CX3CKine amino acid sequence.

SEQ ID NO: 5 is partial mouse CX3CKine nucleotide sequence.

SEQ ID NO: 6 is partial mouse CX3CKine amino acid sequence.

SEQ ID NO: 7 is complete mouse CX3Ckine nucleotide sequence.

SEQ ID NO: 8 is complete mouse CX3C amino acid sequence.

SEQ ID NO: 9 is Gro amino acid sequence.

SEQ ID NO: 10 is Ltn amino acid sequence.

SEQ ID NO: 11 is MIP-1 β amino acid sequence.

All references cited herein are incorporated herein by reference to the same extent as if each individual publication or

patent application was specifically and individually indicated to be incorporated by reference in its entirety for all purposes.

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1
 <211> LENGTH: 534
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (89)..(424)
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (462)..(462)
 <223> OTHER INFORMATION: n is a, c, g, or t
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (520)..(520)
 <223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 1

```

cccaagcttg gcacgagggc actgagctct gccgcctggc tctagccgcc tgccctggccc      60
cgccggggac tcttgcccac cctcagcc atg gct cgg ata tct ctg tgg tgg      112
                               Met Ala Pro Ile Ser Leu Ser Trp
                               1             5

ctg ctc cgc ttg gcc acc ttc tgc cat ctg act gtc ctg ctg gct gga      160
Leu Leu Arg Leu Ala Thr Phe Cys His Leu Thr Val Leu Leu Ala Gly
 10             15             20

cag cac cac ggt gtg acg aaa tgc aac atc acg tgc agc aag atg aca      208
Gln His His Gly Val Thr Lys Cys Asn Ile Thr Cys Ser Lys Met Thr
 25             30             35             40

tca aag ata cct gta gct ttg ctc atc cac tat caa cag aac cag gca      256
Ser Lys Ile Pro Val Ala Leu Leu Ile His Tyr Gln Gln Asn Gln Ala
 45             50             55

tca tgc ggc aaa cgc gca atc atc ttg gag acg aga cag cac agg ctg      304
Ser Cys Gly Lys Arg Ala Ile Ile Leu Glu Thr Arg Gln His Arg Leu
 60             65             70

ttc tgt gcc gac ccg aag gag caa tgg gtc aag gac gcg atg cag cat      352
Phe Cys Ala Asp Pro Lys Glu Gln Trp Val Lys Asp Ala Met Gln His
 75             80             85

ctg gac cgc cag gct gct gcc cta act ccg aaa tgg cgg cac ctt ccg      400
Leu Asp Arg Gln Ala Ala Leu Thr Pro Lys Trp Arg His Leu Pro
 90             95             100

aag aag cca gat cgg cga ggt tga agcccaggac caccctctgc cgccggggga      454
Lys Lys Pro Asp Arg Arg Gly
105             110

aatggacnag tctgttggtc cctggaaccc cgaaagccca caggcgaaaa gccagttacc      514
ctgganccga atccttcttc      534
    
```

<210> SEQ ID NO 2
 <211> LENGTH: 111
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

-continued

```

Met Ala Pro Ile Ser Leu Ser Trp Leu Leu Arg Leu Ala Thr Phe Cys
1           5           10           15
His Leu Thr Val Leu Leu Ala Gly Gln His His Gly Val Thr Lys Cys
          20           25           30
Asn Ile Thr Cys Ser Lys Met Thr Ser Lys Ile Pro Val Ala Leu Leu
          35           40           45
Ile His Tyr Gln Gln Asn Gln Ala Ser Cys Gly Lys Arg Ala Ile Ile
          50           55           60
Leu Glu Thr Arg Gln His Arg Leu Phe Cys Ala Asp Pro Lys Glu Gln
65           70           75           80
Trp Val Lys Asp Ala Met Gln His Leu Asp Arg Gln Ala Ala Ala Leu
          85           90           95
Thr Pro Lys Trp Arg His Leu Pro Lys Lys Pro Asp Arg Arg Gly
          100          105          110
    
```

```

<210> SEQ ID NO 3
<211> LENGTH: 1654
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (86)..(1279)
    
```

<400> SEQUENCE: 3

```

aagcttgcca cgagggcact gagctctgcc gectggctct agcgcctgc ctggcccccg      60
ccgggaactct tgcccacct cagcc atg gct ccg ata tct ctg tcg tgg ctg      112
          Met Ala Pro Ile Ser Leu Ser Trp Leu
          1           5
ctc cgc ttg gcc acc ttc tgc cat ctg act gtc ctg ctg gct gga cag      160
Leu Arg Leu Ala Thr Phe Cys His Leu Thr Val Leu Leu Ala Gly Gln
10           15           20           25
cac cac ggt gtg acg aaa tgc aac atc acg tgc agc aag atg aca tca      208
His His Gly Val Thr Lys Cys Asn Ile Thr Cys Ser Lys Met Thr Ser
          30           35           40
aag ata cct gta gct ttg ctc atc cac tat caa cag aac cag gca tca      256
Lys Ile Pro Val Ala Leu Leu Ile His Tyr Gln Gln Asn Gln Ala Ser
          45           50           55
tgc ggc aaa cgc gca atc atc ttg gag acg aga cag cac agg ctg ttc      304
Cys Gly Lys Arg Ala Ile Ile Leu Glu Thr Arg Gln His Arg Leu Phe
          60           65           70
tgt gcc gac ccg aag gag caa tgg gtc aag gac gcg atg cag cat ctg      352
Cys Ala Asp Pro Lys Glu Gln Trp Val Lys Asp Ala Met Gln His Leu
          75           80           85
gac cgc cag gct gct gcc cta act cga aat ggc ggc acc ttc gag aag      400
Asp Arg Gln Ala Ala Leu Thr Arg Asn Gly Gly Thr Phe Glu Lys
          90           95           100          105
cag atc ggc gag gtg aag ccc agg acc acc cct gcc gcc ggg gga atg      448
Gln Ile Gly Glu Val Lys Pro Arg Thr Thr Pro Ala Ala Gly Gly Met
          110          115          120
gac gag tct gtg gtc ctg gag ccc gaa gcc aca ggc gaa agc agt agc      496
Asp Glu Ser Val Val Leu Glu Pro Glu Ala Thr Gly Glu Ser Ser Ser
          125          130          135
ctg gag ccg act cct tct tcc cag gaa gca cag agg gcc ctg ggg acc      544
Leu Glu Pro Thr Pro Ser Ser Gln Glu Ala Gln Arg Ala Leu Gly Thr
          140          145          150
tcc cca gag ctg ccg acg ggc gtg act ggt tcc tca ggg acc agg ctc      592
Ser Pro Glu Leu Pro Thr Gly Val Thr Gly Ser Ser Gly Thr Arg Leu
          155          160          165
ccc ccg acg cca aag gct cag gat gga ggg cct gtg ggc acg gag ctt      640
    
```

-continued

Pro	Pro	Thr	Pro	Lys	Ala	Gln	Asp	Gly	Gly	Pro	Val	Gly	Thr	Glu	Leu	
170					175					180					185	
ttc	cga	gtg	cct	ccc	gtc	tcc	act	gcc	gcc	acg	tgg	cag	agt	tct	gct	688
Phe	Arg	Val	Pro	Pro	Val	Ser	Thr	Ala	Ala	Thr	Trp	Gln	Ser	Ser	Ala	
			190						195					200		
ccc	cac	caa	cct	ggg	ccc	agc	ctc	tgg	gct	gag	gca	aag	acc	tct	gag	736
Pro	His	Gln	Pro	Gly	Pro	Ser	Leu	Trp	Ala	Glu	Ala	Lys	Thr	Ser	Glu	
			205					210					215			
gcc	ccg	tcc	acc	cag	gac	ccc	tcc	acc	cag	gcc	tcc	act	gcg	tcc	tcc	784
Ala	Pro	Ser	Thr	Gln	Asp	Pro	Ser	Thr	Gln	Ala	Ser	Thr	Ala	Ser	Ser	
			220				225					230				
cca	gcc	cca	gag	gag	aat	gct	ccg	tct	gaa	ggc	cag	cgt	gtg	tgg	ggt	832
Pro	Ala	Pro	Glu	Glu	Asn	Ala	Pro	Ser	Glu	Gly	Gln	Arg	Val	Trp	Gly	
			235			240					245					
cag	gga	cag	agc	ccc	agg	cca	gag	aac	tct	ctg	gag	cgg	gag	gag	atg	880
Gln	Gly	Gln	Ser	Pro	Arg	Pro	Glu	Asn	Ser	Leu	Glu	Arg	Glu	Glu	Met	
				250		255				260					265	
ggt	ccc	gtg	cca	gcg	cac	acg	gat	gcc	ttc	cag	gac	tgg	ggg	cct	ggc	928
Gly	Pro	Val	Pro	Ala	His	Thr	Asp	Ala	Phe	Gln	Asp	Trp	Gly	Pro	Gly	
				270					275					280		
agc	atg	gcc	cac	gtc	tct	gtg	gtc	cct	gtc	tcc	tca	gaa	ggg	acc	ccc	976
Ser	Met	Ala	His	Val	Ser	Val	Val	Pro	Val	Ser	Ser	Glu	Gly	Thr	Pro	
				285				290						295		
agc	agg	gag	cca	gtg	gct	tca	ggc	agc	tgg	acc	cct	aag	gct	gag	gaa	1024
Ser	Arg	Glu	Pro	Val	Ala	Ser	Gly	Ser	Trp	Thr	Pro	Lys	Ala	Glu	Glu	
				300			305						310			
ccc	atc	cat	gcc	acc	atg	gac	ccc	cag	agg	ctg	ggc	gtc	ctt	atc	act	1072
Pro	Ile	His	Ala	Thr	Met	Asp	Pro	Gln	Arg	Leu	Gly	Val	Leu	Ile	Thr	
			315			320					325					
cct	gtc	cct	gac	gcc	cag	gct	gcc	acc	cgg	agg	cag	gcg	gtg	ggg	ctg	1120
Pro	Val	Pro	Asp	Ala	Gln	Ala	Ala	Thr	Arg	Arg	Gln	Ala	Val	Gly	Leu	
				330		335				340					345	
ctg	gcc	ttc	ctt	ggc	ctc	ctc	ttc	tgc	ctg	ggg	gtg	gcc	atg	ttc	acc	1168
Leu	Ala	Phe	Leu	Gly	Leu	Leu	Phe	Cys	Leu	Gly	Val	Ala	Met	Phe	Thr	
				350					355					360		
tac	cag	agc	ctc	cag	ggc	tgc	cct	cga	aag	atg	gca	gga	gag	atg	gcg	1216
Tyr	Gln	Ser	Leu	Gln	Gly	Cys	Pro	Arg	Lys	Met	Ala	Gly	Glu	Met	Ala	
				365					370					375		
gag	ggc	ctt	cgc	tac	atc	ccc	cgg	agc	tgt	ggt	agt	aat	tca	tat	gtc	1264
Glu	Gly	Leu	Arg	Tyr	Ile	Pro	Arg	Ser	Cys	Gly	Ser	Asn	Ser	Tyr	Val	
				380			385					390				
ctg	gtg	ccc	gtg	tga	actcctctgg	cctgtgtcta	ggtgtttgat	tcagacagct								1319
Leu	Val	Pro	Val													
				395												
gcctgggatac	cctcatcctc	atacccaccc	ccaccaagg	gcctggcctg	agctgggatg											1379
attggagggg	ggaggtggga	tcctccaggt	gcacaagctc	caagctccca	ggcattcccc											1439
aggaggccag	ccttgaccat	tctccacctt	ccagggacag	aggggggtggc	ctcccaactc											1499
accccagccc	caaaactctc	ctctgctgct	ggctggttag	aggttccctt	tgacgccatc											1559
ccagccccaa	tgaacaatta	tttattaaat	gcccagcccc	ttctgaaaaa	aaaaaaaaaa											1619
aaaaaaaaaa	aaaaaaaaaa	aattcctgcg	gccgc													1654

<210> SEQ ID NO 4
 <211> LENGTH: 397
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 4

Met Ala Pro Ile Ser Leu Ser Trp Leu Leu Arg Leu Ala Thr Phe Cys

-continued

1	5	10	15
His Leu Thr Val Leu Leu Ala Gly Gln His His Gly Val Thr Lys Cys	20	25	30
Asn Ile Thr Cys Ser Lys Met Thr Ser Lys Ile Pro Val Ala Leu Leu	35	40	45
Ile His Tyr Gln Gln Asn Gln Ala Ser Cys Gly Lys Arg Ala Ile Ile	50	55	60
Leu Glu Thr Arg Gln His Arg Leu Phe Cys Ala Asp Pro Lys Glu Gln	65	70	75
Trp Val Lys Asp Ala Met Gln His Leu Asp Arg Gln Ala Ala Ala Leu	85	90	95
Thr Arg Asn Gly Gly Thr Phe Glu Lys Gln Ile Gly Glu Val Lys Pro	100	105	110
Arg Thr Thr Pro Ala Ala Gly Gly Met Asp Glu Ser Val Val Leu Glu	115	120	125
Pro Glu Ala Thr Gly Glu Ser Ser Ser Leu Glu Pro Thr Pro Ser Ser	130	135	140
Gln Glu Ala Gln Arg Ala Leu Gly Thr Ser Pro Glu Leu Pro Thr Gly	145	150	155
Val Thr Gly Ser Ser Gly Thr Arg Leu Pro Pro Thr Pro Lys Ala Gln	165	170	175
Asp Gly Gly Pro Val Gly Thr Glu Leu Phe Arg Val Pro Pro Val Ser	180	185	190
Thr Ala Ala Thr Trp Gln Ser Ser Ala Pro His Gln Pro Gly Pro Ser	195	200	205
Leu Trp Ala Glu Ala Lys Thr Ser Glu Ala Pro Ser Thr Gln Asp Pro	210	215	220
Ser Thr Gln Ala Ser Thr Ala Ser Ser Pro Ala Pro Glu Glu Asn Ala	225	230	235
Pro Ser Glu Gly Gln Arg Val Trp Gly Gln Gly Gln Ser Pro Arg Pro	245	250	255
Glu Asn Ser Leu Glu Arg Glu Glu Met Gly Pro Val Pro Ala His Thr	260	265	270
Asp Ala Phe Gln Asp Trp Gly Pro Gly Ser Met Ala His Val Ser Val	275	280	285
Val Pro Val Ser Ser Glu Gly Thr Pro Ser Arg Glu Pro Val Ala Ser	290	295	300
Gly Ser Trp Thr Pro Lys Ala Glu Glu Pro Ile His Ala Thr Met Asp	305	310	315
Pro Gln Arg Leu Gly Val Leu Ile Thr Pro Val Pro Asp Ala Gln Ala	325	330	335
Ala Thr Arg Arg Gln Ala Val Gly Leu Leu Ala Phe Leu Gly Leu Leu	340	345	350
Phe Cys Leu Gly Val Ala Met Phe Thr Tyr Gln Ser Leu Gln Gly Cys	355	360	365
Pro Arg Lys Met Ala Gly Glu Met Ala Glu Gly Leu Arg Tyr Ile Pro	370	375	380
Arg Ser Cys Gly Ser Asn Ser Tyr Val Leu Val Pro Val	385	390	395

<210> SEQ ID NO 5
 <211> LENGTH: 209
 <212> TYPE: DNA
 <213> ORGANISM: Mus sp.
 <220> FEATURE:

-continued

```

<221> NAME/KEY: misc_feature
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (42)..(42)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (63)..(209)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (135)..(135)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (184)..(184)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (191)..(191)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (204)..(204)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 5

tnactactag gagctgcgac acggcccagc ctctggccc gncgaattcc tgcactccag      60
cc atg gct ccc tcg ccg ctc gcg tgg ctg ctg cgc ctg gcc gcg ttc      107
  Met Ala Pro Ser Pro Leu Ala Trp Leu Leu Arg Leu Ala Ala Phe
    1             5             10             15

ttc cat ttg tgt act ctg ctg ccg ggt nag cac ctc ggc atg acg aaa      155
Phe His Leu Cys Thr Leu Leu Pro Gly Xaa His Leu Gly Met Thr Lys
    20             25             30

tgc gaa atc atg tgc gac aag atg acc tna cga atn cca gtg gct tta      203
Cys Glu Ile Met Cys Asp Lys Met Thr Xaa Arg Xaa Pro Val Ala Leu
    35             40             45

ntc atc      209
Xaa Ile

```

```

<210> SEQ ID NO 6
<211> LENGTH: 49
<212> TYPE: PRT
<213> ORGANISM: Mus sp.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (25)..(25)
<223> OTHER INFORMATION: The 'Xaa' at location 25 stands for Lys, Glu,
or Gln.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (41)..(41)
<223> OTHER INFORMATION: The 'Xaa' at location 41 stands for Ser, or
Leu.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (43)..(43)
<223> OTHER INFORMATION: The 'Xaa' at location 43 stands for Ile, or
Met.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (48)..(48)
<223> OTHER INFORMATION: The 'Xaa' at location 48 stands for Ile, Val,
Leu, or Phe.

```

```

<400> SEQUENCE: 6

Met Ala Pro Ser Pro Leu Ala Trp Leu Leu Arg Leu Ala Ala Phe Phe
1             5             10             15

His Leu Cys Thr Leu Leu Pro Gly Xaa His Leu Gly Met Thr Lys Cys
20             25             30

```

-continued

Glu Ile Met Cys Asp Lys Met Thr Xaa Arg Xaa Pro Val Ala Leu Xaa
 35 40 45

Ile

<210> SEQ ID NO 7
 <211> LENGTH: 3065
 <212> TYPE: DNA
 <213> ORGANISM: Mus sp.
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (62)..(1249)

<400> SEQUENCE: 7

tgactactag gagctgcgac acggcccagc ctctggccg cogaattcct gcactccagc 60

c atg gct ccc tgc ccg ctc gcg tgg ctg ctg cgc ctg gcc gcg ttc ttc 109
 Met Ala Pro Ser Pro Leu Ala Trp Leu Leu Arg Leu Ala Ala Phe Phe
 1 5 10 15

cat ttg tgt act ctg ctg ccg ggt cag cac ctc ggc atg acg aaa tgc 157
 His Leu Cys Thr Leu Leu Pro Gly Gln His Leu Gly Met Thr Lys Cys
 20 25 30

gaa atc atg tgc ggc aag atg acc tca cga atc cca gtg gct ttg ctc 205
 Glu Ile Met Cys Gly Lys Met Thr Ser Arg Ile Pro Val Ala Leu Leu
 35 40 45

atc cgc tat cag cta aat cag gag tcc tgc ggc aag cgt gcc att gtc 253
 Ile Arg Tyr Gln Leu Asn Gln Glu Ser Cys Gly Lys Arg Ala Ile Val
 50 55 60

ctg gag acg aca cag cac aga cgc ttc tgt gct gac ccg aag gag aaa 301
 Leu Glu Thr Thr Gln His Arg Arg Phe Cys Ala Asp Pro Lys Glu Lys
 65 70 75 80

tgg gtc caa gac gcc atg aag cat ctg gat cac cag gct gct gcc ctc 349
 Trp Val Gln Asp Ala Met Lys His Leu Asp His Gln Ala Ala Ala Leu
 85 90 95

act aaa aat ggt ggc aag ttt gag aag cgg gtg gac aat gtg aca cct 397
 Thr Lys Asn Gly Gly Lys Phe Glu Lys Arg Val Asp Asn Val Thr Pro
 100 105 110

ggg atc acc ttg gcc act agg gga ctg tcc cca tct gcc ctg aca aag 445
 Gly Ile Thr Leu Ala Thr Arg Gly Leu Ser Pro Ser Ala Leu Thr Lys
 115 120 125

cct gaa tcc gcc aca ttg gaa gac ctt gct ttg gaa ctg act act att 493
 Pro Glu Ser Ala Thr Leu Glu Asp Leu Ala Leu Glu Leu Thr Thr Ile
 130 135 140

tcc cag gag gcc agg ggg acc atg ggg act tcc caa gag cca ccg gca 541
 Ser Gln Glu Ala Arg Gly Thr Met Gly Thr Ser Gln Glu Pro Pro Ala
 145 150 155 160

gca gtg acc gga tca tct ctc tca act tcc gag gca cag gat gca ggg 589
 Ala Val Thr Gly Ser Ser Leu Ser Thr Ser Glu Ala Gln Asp Ala Gly
 165 170 175

ctt acg gct aag cct cag agc att gga agt ttt gag gcg gct gac atc 637
 Leu Thr Ala Lys Pro Gln Ser Ile Gly Ser Phe Glu Ala Ala Asp Ile
 180 185 190

tcc acc acc gtt tgg ccg agt cct gct gtc tac caa tct gga tct agc 685
 Ser Thr Thr Val Trp Pro Ser Pro Ala Val Tyr Gln Ser Gly Ser Ser
 195 200 205

tcc tgg gct gag gaa aaa gct act gag tcc ccc tcc act aca gcc cca 733
 Ser Trp Ala Glu Glu Lys Ala Thr Glu Ser Pro Ser Thr Thr Ala Pro
 210 215 220

tct cct cag gtg tcc act act tca cct tca acc cca gag gaa aat gtt 781
 Ser Pro Gln Val Ser Thr Thr Ser Pro Ser Thr Pro Glu Glu Asn Val
 225 230 235 240

ggg tcc gaa ggc caa ccc cca tgg gtc cag gga cag gac ctc agt cca 829
 Gly Ser Glu Gly Gln Pro Pro Trp Val Gln Gly Gln Asp Leu Ser Pro

-continued

245		250		255		
gag aag tct cta ggg tct gag gag	ata aac cca gtt cat act gat aat					877
Glu Lys Ser Leu Gly Ser Glu Glu	Ile Asn Pro Val His Thr Asp Asn					
260	265			270		
ttc cag gag agg ggg cct ggc aac	aca gtc cac ccc tca gtg gct ccc					925
Phe Gln Glu Arg Gly Pro Gly Asn	Thr Val His Pro Ser Val Ala Pro					
275	280			285		
atc tcc tct gaa gag acc ccc agc	cca gag ctg gtg gcc tcg ggc agc					973
Ile Ser Ser Glu Glu Thr Pro Ser	Pro Glu Leu Val Ala Ser Gly Ser					
290	295			300		
cag gct cct aag ata gag gaa ccc	atc cat gcc act gca gat ccc cag					1021
Gln Ala Pro Lys Ile Glu Glu Pro	Ile His Ala Thr Ala Asp Pro Gln					
305	310			315		320
aaa ctg agt gtg ctt atc act cct	gtc ccc gac acc cag gca gcc aca					1069
Lys Leu Ser Val Leu Ile Thr Pro	Val Pro Asp Thr Gln Ala Ala Thr					
325	330			335		
agg agg cag gca gtg ggg cta ctg	gct ttc ctt ggt ctt ctt ttc tgc					1117
Arg Arg Gln Ala Val Gly Leu Leu	Ala Phe Leu Gly Leu Leu Phe Cys					
340	345			350		
cta ggg gtg gcc atg ttt gct tac	cag agc ctt cag ggc tgt ccc cgc					1165
Leu Gly Val Ala Met Phe Ala Tyr	Gln Ser Leu Gln Gly Cys Pro Arg					
355	360			365		
aaa atg gcg ggg gaa atg gta gaa	ggc ctc cgc tac gtc ccc cgt agc					1213
Lys Met Ala Gly Glu Met Val Glu	Gly Leu Arg Tyr Val Pro Arg Ser					
370	375			380		
tgt ggc agt aac tca tac gtc ctg	gtg cca gtg tga gctgcttgcc					1259
Cys Gly Ser Asn Ser Tyr Val Leu	Val Pro Val					
385	390			395		
tgctgctgctg tgtccagagt gtgattcgga	cagctgtctg gggaccccc cccatcctca					1319
taccacacct catccacgtg ggggaaatgg	gaatggagaa gctggacctc caggggctgt					1379
gggctccatc caatccccct tccccgagg	ggtggccccg gaggccacc tagaccacta					1439
ttcacttate agagacagag caggtgaact	tccagctcct ctatattga aagaatcctc					1499
tgctgctggc tggtagagg ggccttgac	acccaactc cagtgaacaa ttatttattg					1559
gattcccagc cctgcgacg acacctgttt	cccgcgcgca cgtggtccg cccatatcac					1619
aagcagcagg ccaggcctat ctgectgtcc	cctgacctc cttgtgtctc ctggctttgc					1679
tgcagtgcc agcccttctc ctccccggcc	agccgcggtg ctatctgccc tatgtctccc					1739
tctatcccct gtacagagcg caccaccate	accatcaaca ccgctgttgt gtcttttctt					1799
gcatgagggt aaagctgtgt tttctggagc	tctccgggaa gggagacaag cttgcgagag					1859
ggtttaaagt gttcctcccc agacttggat	gtgctgtgag ggcagctgc gtctgaagga					1919
agggtccagt cccactcgg ctaccagcac	cacaaagtgc cccacctgta aaaggaaaga					1979
aacgtgtgcc agagctggca ataacctatg	gcctgacat catcacttcc tctgagatcc					2039
ttgtctccac cctgggtgac aacccccacc	cttatcaaca ttaatagtca ctgccattcc					2099
actggactga catttttcta cctgtgatt	ctgagggctg gcaaggagtg gcttgagagt					2159
gcagatcgta cctgtatgc ccccccaaaa	tggaggctga gttggggact tgcaggaaca					2219
gaggccaact cagatggctt cccctgtgtt	ctcactagaa acccctcccc catgcaccaa					2279
ggtgacagtc acaggtctgc cctggctaaa	ggacaagcca cataggaaag attaggacaa					2339
gccccctgga ggcagaggat ccagggtaaa	cccctggagt ggccacaaac ccaatttcag					2399
tgtagggact tgtgcatgtg tgaacttga	tagtcagaca gaggctgcca gggctcttcc					2459
ctgtctctga gagcagtggt cacgccaagg	actcaccttt gccccattg caggcagggc					2519

-continued

```

cagaactccc atagcattct ccaagagccc tgtgacattt tctggaagga actctgcct 2579
ggcgcaaaag tgactgctga agcaaggagc agctgagcag caccocagcg gagctgagcc 2639
ggcaggccac gccctcggg ggggggcatt tctaccgcc ctgctctgaa tagctccaac 2699
ttcaccttag gagcctccca gggcgagct tcaccagaa gccagtgact cactccttga 2759
ttggtggaag ctcagttggc tcctgagagt gaggaagcca accctttgtc gaccctctc 2819
ctgggaagcc tgtggggcgc tctgatcatg ctccacagaa ccagttgtag gcctgagccg 2879
cagcagcccg agtgactat atcctggctc cttcgggtgg gaacctttaa gggttgggac 2939
accctgatc ggactttgtt gttctctccc tcccagagca gaatgtgggc cgtaacaatc 2999
tgaggaggac tttaaaagtt gttgatcctt tagggttttt tttcaagcat cattaccaat 3059
gtctgt 3065

```

```

<210> SEQ ID NO 8
<211> LENGTH: 395
<212> TYPE: PRT
<213> ORGANISM: Mus sp.

```

```

<400> SEQUENCE: 8

```

```

Met Ala Pro Ser Pro Leu Ala Trp Leu Leu Arg Leu Ala Ala Phe Phe
1          5          10          15
His Leu Cys Thr Leu Leu Pro Gly Gln His Leu Gly Met Thr Lys Cys
20          25          30
Glu Ile Met Cys Gly Lys Met Thr Ser Arg Ile Pro Val Ala Leu Leu
35          40          45
Ile Arg Tyr Gln Leu Asn Gln Glu Ser Cys Gly Lys Arg Ala Ile Val
50          55          60
Leu Glu Thr Thr Gln His Arg Arg Phe Cys Ala Asp Pro Lys Glu Lys
65          70          75          80
Trp Val Gln Asp Ala Met Lys His Leu Asp His Gln Ala Ala Ala Leu
85          90          95
Thr Lys Asn Gly Gly Lys Phe Glu Lys Arg Val Asp Asn Val Thr Pro
100         105         110
Gly Ile Thr Leu Ala Thr Arg Gly Leu Ser Pro Ser Ala Leu Thr Lys
115         120         125
Pro Glu Ser Ala Thr Leu Glu Asp Leu Ala Leu Glu Leu Thr Thr Ile
130         135         140
Ser Gln Glu Ala Arg Gly Thr Met Gly Thr Ser Gln Glu Pro Pro Ala
145         150         155         160
Ala Val Thr Gly Ser Ser Leu Ser Thr Ser Glu Ala Gln Asp Ala Gly
165         170         175
Leu Thr Ala Lys Pro Gln Ser Ile Gly Ser Phe Glu Ala Ala Asp Ile
180         185         190
Ser Thr Thr Val Trp Pro Ser Pro Ala Val Tyr Gln Ser Gly Ser Ser
195         200         205
Ser Trp Ala Glu Glu Lys Ala Thr Glu Ser Pro Ser Thr Thr Ala Pro
210         215         220
Ser Pro Gln Val Ser Thr Thr Ser Pro Ser Thr Pro Glu Glu Asn Val
225         230         235         240
Gly Ser Glu Gly Gln Pro Pro Trp Val Gln Gly Gln Asp Leu Ser Pro
245         250         255
Glu Lys Ser Leu Gly Ser Glu Glu Ile Asn Pro Val His Thr Asp Asn
260         265         270
Phe Gln Glu Arg Gly Pro Gly Asn Thr Val His Pro Ser Val Ala Pro

```


-continued

100	105	110
Leu Thr Gly		
115		
<210> SEQ ID NO 11 <211> LENGTH: 92 <212> TYPE: PRT <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Mip-1 beta amino acid sequence <400> SEQUENCE: 11		
Met Lys Leu Cys Val Ser Ala Leu Ser Leu Leu Leu Val Ala Ala		
1	5	10 15
Phe Cys Ala Pro Gly Phe Ser Ala Pro Met Gly Ser Asp Pro Pro Thr		
20	25	30
Ser Cys Cys Phe Ser Tyr Thr Ala Arg Lys Leu Pro Arg Asn Phe Val		
35	40	45
Val Asp Tyr Tyr Glu Thr Ser Ser Leu Cys Ser Gln Pro Ala Val Val		
50	55	60
Phe Gln Thr Lys Arg Ser Lys Gln Val Cys Ala Asp Pro Ser Glu Ser		
65	70	75 80
Trp Val Gln Glu Tyr Val Tyr Asp Leu Glu Leu Asn		
85	90	

We claim:

1. An isolated polypeptide wherein said polypeptide a) chemoattracts peripheral blood monocytes or T-cells, b) consists of an amino acid sequence comprising at least 95% amino acid sequence identity to the amino acid sequence set forth in SEQ ID NO 8; and c) comprises a chemokine domain identical to residues 25 to 100 of the amino acid sequence set forth in SEQ ID NO: 8.
2. The polypeptide of claim 1, which comprises the amino acid sequence set forth in SEQ ID NO: 8.
3. A composition comprising the polypeptide of claim 1 and a carrier.
4. The composition of claim 3 wherein the carrier is water, saline, or buffer.
5. The polypeptide fused to a heterologous protein.

30

6. The polypeptide of claim 5 wherein the heterologous protein is a FLAG peptide.

35

7. A method for producing an isolated polypeptide wherein said polypeptide a) chemoattracts peripheral blood monocytes or T-cells, b) consists of an amino acid sequence comprising at least 95% amino acid sequence identity to the amino acid sequence set forth in SEQ ID NO: 8, and c) comprises a chemokine domain identical to residues 25 to 100 of the amino acid sequence set forth in SEQ ID NO: 8; comprising introducing a vector comprising a polynucleotide encoding said polypeptide into an isolated host cell and culturing said host cell.

40

8. The method of claim 7 wherein the polypeptide comprises the amino acid sequence set forth in SEQ ID NO: 8.

* * * * *

专利名称(译)	小鼠CX3C趋化因子多肽		
公开(公告)号	US8173117	公开(公告)日	2012-05-08
申请号	US12/833069	申请日	2010-07-09
[标]申请(专利权)人(译)	先灵葆雅有限公司		
申请(专利权)人(译)	Schering Corporation的		
当前申请(专利权)人(译)	Schering Corporation的		
[标]发明人	HARDIMAN GERARD T ROSSI DEVORA L BACON KEVIN B J BAZAN FERNANDO SCHALL THOMAS P ZLOTNIK ALBERT		
发明人	HARDIMAN, GERARD T. ROSSI, DEVORA L. BACON, KEVIN B J., BAZAN FERNANDO SCHALL, THOMAS P. ZLOTNIK, ALBERT		
IPC分类号	C07K14/52 A61K38/19 C12N15/64 C12N15/09 A61K38/00 A61K39/395 A61P25/00 A61P29/00 A61P37/00 A61P43/00 C07K14/47 C07K16/24 C07K19/00 C12N5/10 C12N15/19 G01N33/53		
CPC分类号	C07K14/521 A61K38/00 C07K2319/00 A61P25/00 A61P29/00 A61P37/00 A61P43/00 Y10S435/81		
其他公开文献	US20110189123A1		
外部链接	Espacenet USPTO		

摘要(译)

描述了编码来自哺乳动物的新趋化因子家族CX3C家族的核酸，与其相关的试剂，包括特异性抗体和纯化的蛋白质。还提供了使用所述试剂和相关诊断试剂盒的方法。