



US009052322B2

**(12) United States Patent  
Yu****(10) Patent No.: US 9,052,322 B2  
(45) Date of Patent: \*Jun. 9, 2015**

- (54) **METHOD FOR IDENTIFYING LINEAGE-RELATED ANTIBODIES**
- (71) Applicant: **Epitomics, Inc.**, Cambridge (GB)
- (72) Inventor: **Guo-Liang Yu**, Hillsborough, CA (US)
- (73) Assignee: **EPITOMICS, INC.**, Burlingame, CA (US)
- (\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- This patent is subject to a terminal disclaimer.

5,639,641 A	6/1997	Pedersen et al.
5,693,761 A	12/1997	Queen et al.
5,693,762 A	12/1997	Queen et al.
5,876,961 A	3/1999	Crowe et al.
5,962,255 A	10/1999	Griffiths et al.
6,180,370 B1	1/2001	Queen et al.
6,329,551 B1	12/2001	Nakagome et al.
6,331,415 B1	12/2001	Cabilly et al.
6,342,587 B1	1/2002	Barbas, III et al.
6,372,214 B1	4/2002	Prusiner et al.
6,576,467 B1	6/2003	Stemmer
6,596,492 B2	7/2003	Avery et al.
7,871,611 B2	1/2011	Calzone et al.
8,293,483 B2	10/2012	Yu
2001/0036647 A1	11/2001	Choudary et al.
2002/0160373 A1	10/2002	Avery et al.
2002/0177170 A1	11/2002	Luo et al.
2003/0198638 A1	10/2003	Watkins
2004/0010376 A1	1/2004	Luo et al.
2004/0086979 A1	5/2004	Zhang et al.
2005/0033031 A1	2/2005	Couto
2006/0099204 A1	5/2006	Couto et al.
2006/0233812 A1	10/2006	Burnie et al.

- (21) Appl. No.: **14/177,003**
- (22) Filed: **Feb. 10, 2014**

- (65) **Prior Publication Data**  
US 2014/0155291 A1 Jun. 5, 2014

- Related U.S. Application Data**
- (63) Continuation of application No. 13/748,507, filed on Jan. 23, 2013, which is a continuation of application No. 13/552,517, filed on Jul. 18, 2012, now Pat. No. 8,617,830, which is a continuation of application No. 12/878,925, filed on Sep. 9, 2010, now Pat. No. 8,293,483.

- (60) Provisional application No. 61/241,714, filed on Sep. 11, 2009.

- (51) **Int. Cl.**  
**G01N 33/53** (2006.01)  
**C07K 16/00** (2006.01)  
**G01N 33/577** (2006.01)  
**C07K 16/28** (2006.01)  
**G01N 33/68** (2006.01)  
**G01N 33/566** (2006.01)

- (52) **U.S. Cl.**  
CPC ..... **G01N 33/577** (2013.01); **C07K 16/00** (2013.01); **C07K 16/2863** (2013.01); **C07K 2317/10** (2013.01); **C07K 2317/56** (2013.01); **C07K 2317/76** (2013.01); **G01N 33/6854** (2013.01); **G01N 33/566** (2013.01)

- (58) **Field of Classification Search**  
None  
See application file for complete search history.

- (56) **References Cited**  
U.S. PATENT DOCUMENTS

4,727,037 A	2/1988	Ring
4,816,397 A	3/1989	Boss et al.
4,816,567 A	3/1989	Cabilly et al.
4,859,595 A	8/1989	Strosberg et al.
4,977,081 A	12/1990	Raybould et al.
5,225,539 A	7/1993	Winter
5,472,868 A	12/1995	McCormack et al.
5,530,101 A	6/1996	Queen et al.
5,585,089 A	12/1996	Queen et al.
5,610,034 A	3/1997	Nyssonen et al.

(Continued)

## FOREIGN PATENT DOCUMENTS

WO	WO0029584	5/2000
WO	WO0148485	7/2001
WO	WO2006055778	5/2006
WO	WO2007056441	5/2007

## OTHER PUBLICATIONS

Bendig, Mary M., "Humanization of Rodent Monoclonal Antibodies by CDR Grafting", *Methods: A Companion to Methods in Enzymology*, vol. 8, 1995, pp. 83-93.

Bjorling, et al Human neutralizing human immunodeficiency virus type 2-specific Fab molecules generated by phage display, *J Gen Virol*, 80 ( Pt 8):1987-93, 1999.

Carmen, et al., "Concepts in antibody phage display", *Brief Funct Genomic Proteomic.*, 2002, 1(2):189-203.

(Continued)

*Primary Examiner* — Michail Belyavskiy

(74) *Attorney, Agent, or Firm* — James S. Keddie; Bozicevic, Field & Francis, LLP

- (57) **ABSTRACT**

In certain embodiments, the method may comprise: a) obtaining the antibody sequences from a population of B cells; b) grouping the antibody sequences to provide a plurality of groups of lineage-related antibodies; c) testing a single antibody from each of the groups in a bioassay and, after the first antibody has been identified, d) testing further antibodies that are in the same group as the first antibody in a second bioassay. In another embodiment, the method may comprise: a) testing a plurality of antibodies obtained from a first portion of an antibody producing organ of an animal; b) obtaining the sequence of a first identified antibody; c) obtaining from a second portion of said antibody producing organ the sequences of further antibodies that are related by lineage to said first antibody; and, c) testing the further antibodies in a second bioassay.

26 Claims, 13 Drawing Sheets

(56)

## References Cited

## U.S. PATENT DOCUMENTS

2007/0037217	A1	2/2007	Luo et al.
2007/0269868	A1	11/2007	Carvalho Jensen et al.
2008/0075712	A1	3/2008	Hattori et al.
2008/0207459	A1	8/2008	Karrer et al.
2008/0227660	A1	9/2008	Kastrup et al.
2009/0054254	A1	2/2009	Throsby et al.
2009/0081190	A1	3/2009	Stassar et al.
2009/0175846	A1	7/2009	Mi et al.
2010/0292083	A1	11/2010	Kolkman

## OTHER PUBLICATIONS

De Pascalis et al. Grafting of "abbreviated" complementarity-determining regions containing specificity-determining residues essential for ligand contact to engineer a less immunogenic humanized monoclonal antibody. *The Journal of Immunology*. 2002, vol. 169, pp. 3076-3084.

Delagrave et al. Effects of humanization by variable domain resurfacing on the antiviral activity of a single-chain antibody against respiratory syncytial virus. *Protein Engineering*. 1999, vol. 12, No. 4, pp. 357-362.

Green. Antibody engineering via genetic engineering of the mouse: Xenomouse strains are a vehicle for the facile generation of therapeutic monoclonal antibodies. *Journal of Immunological Methods*. 1999, vol. 231, pp. 11-23.

Griffiths, et al., "Isolation of high affinity human antibodies directly from large synthetic repertoires", *The EMBO Journal*. 1994, vol. 13, No. 14, pp. 3245-3260.

Kala, et al., "Phage displayed antibodies to heat stable alkaline phosphatase: framework region as a determinant of specificity", *J Biochem.*, 2002,132(4):535-41.

Kang, et al., Antibody redesign by chain shuffling from random combinatorial immunoglobulin libraries, *Proceedings of the National Academy of Sciences*, vol. 88, pp. 11120-11123, 1991.

Mehr et al. Analysis of mutational lineage trees from sites of primary and secondary Ig gene diversification in rabbits and chickens. *The Journal of Immunology*. 2004, vol. 172, pp. 4790-4796.

Morea et al. Antibody modeling: implications for engineering and design. *Methods*. 2000, vol. 20, pp. 267-279.

Paul, *Fundamental Immunology*, 3rd Edition, p. 292-295, 1993.

Popkov et al. Rabbit immune repertoires as sources for therapeutic monoclonal antibodies: the impact of kappa allotype-correlated variation in cysteine content on antibody libraries selected by phage display. *Journal of Molecular Biology*. 2003, vol. 325, pp. 325-335.

Rader, et al., A phage display approach for rapid antibody humanization: Designed combinatorial V gene libraries, *Proc Natl Acad Sci U S A*. Jul. 21, 1998; 95(15): 8910-8915.

Rader, et al., "The Rabbit Antibody Repertoire as a Novel Source for the Generation of Therapeutic Human Antibodies", *J Biol Chem.*, 2000, 275(18):13668-76.

Roguska et al. A comparison of two murine monoclonal antibodies humanized by CDR-grafting and variable domain resurfacing. *Protein Engineering*. 1996, vol. 9, pp. 895-904.

Roguska et al. Humanization of murine monoclonal antibodies through variable domain resurfacing. *PNAS*. 1994, vol. 91, pp. 969-973.

Rothe, et al., "In vitro display technologies reveal novel biopharmaceutics", *FASEB J.*, 2006, 20(10):1599-1610.

Sblattero, Exploiting recombination in single bacteria to make large phage antibody libraries, *Nature biotechnology*, vol. 18, pp. 75-80, 2000.

Smith, et al., "Antibody phage display technologies with special reference to angiogenesis", *FASEB J.*, 2005, 19 (3):331-341.

Steinberger et al. Generation and characterization of a recombinant human CCR5-specific antibody. *The Journal of Biological Chemistry*. 2000, vol. 275, No. 46, pp. 36073-36078.

Sun et al. Antibody repertoire development in fetal and neonatal piglets. I. Four VH genes account for 80 percent of VH usage during 84 days of fetal life. *Journal of Immunology*. 1998, vol. 161, pp. 5070-5078.

Telenius, et al., "Degenerate oligonucleotide-primed PCR: general amplification of target DNA by a single degenerate primer.", *Genomics* (1992), vol. 13, Issue: 3, pp. 718-725.

Vaswani et al. Humanized antibodies as potential therapeutic drugs. *Annals of Allergy, Asthma & Immunology*. 1998, vol. 81, pp. 105-119.

Yang, et al., "Evolutional selection of a combinatorial phage library displaying randomly-rearranged various single domains of immunoglobulin (Ig)-binding proteins (IBPs) with four kinds of Ig molecules", *BMC Microbiol.*, 2008, 8:137.

Yu et al. A humanized anti-VEGF rabbit monoclonal antibody inhibits angiogenesis and blocks tumor growth in xenograft models. *PLoS One*. 2010, vol. 5, pp. e9072.

Acosta, et al., Specific monoclonal antibody against human trypsin, *Hybrid Hybridomics*, 2002, 21:307-10.

Babcock, et al., A novel strategy for generating monoclonal antibodies from single, isolated lymphocytes producing antibodies of defined specificities, *Proc Natl Acad Sci USA*, 1996, 93:7843-8.

Becker, et al., Somatic diversification of immunoglobulin heavy chain VDJ genes: evidence for somatic gene conversion in rabbits, *Cell*, 1990, 63:987-97.

Bos, et al., Humoral immune response to 2,4-dinitrophenyl—key-hole limpet hemocyanin in antigen-free, germ-free and conventional BALB/c mice, *Eur J Immunol*, 1994, 24:59-65.

Calame, Plasma cells: finding new light at the end of B cell development, *Nat Immunol*, 2001, 2:1103-8.

Coronella, et al., Amplification of IgG VH and VL (Fab) from single human plasma cells and B cells, *Nucleic Acids Res.*, 2000, 28:1-7.

De Wildt, et al., A new method for the analysis and production of monoclonal antibody fragments originating from single human B cells, *J Immunol Methods*, 1997, 207:61-7.

Durocher, et al., High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human 293-EBNA1 cells, *Nucleic Acids Res*, 2002, 30:E9.

Embleton, et al., In-cell PCR from mRNA: amplifying and linking the rearranged immunoglobulin heavy and light chain V-genes within single cells, *Nucleic Acids Res.*, 1992, 20:3831-7.

Friedmann, et al., Neonatal VH, D, and JH gene usage in rabbit B lineage cells, 1994, *J Immunology*, 152:632-641.

Huse, et al., Generation of a large combinatorial library of the immunoglobulin repertoire in phage lambda, 1992, *Biotechnology*, 24:517-23.

Knight, et al., Molecular basis of the allelic inheritance of rabbit immunoglobulin VH allotypes: Implications for the generation of antibody diversity, *Cell*, 1990, 60:963-970.

Kurome, et al., Expression of recombinant mouse/human chimeric antibody specific to human GMP-140/P-selectin, *J Biochem*. 1994, 115:608-14.

Lagerkvist, et al., Single, antigen-specific B cells used to generate Fab fragments using CD40-mediated amplification or direct PCR cloning, *Bio Techniques*, 1995, 18:862-869.

Magori-Cohen, et al., Mutation parameters from DNA sequence data using graph theoretic measures on lineage trees, *Bioinformatics*, 2006, 22:e332-e340.

Marks, et al., By-passing immunization: human antibodies from V gene libraries displayed on phage, *J Mol Biol*, 1991, 222: 581-597.

Merz, et al., Generating a phage display antibody library against an identified neuron, *J Neurosci Methods*, 1995, 62:213-9.

Ochsenbein, et al., Protective long-term antibody memory by antigen-driven and T help-dependent differentiation of long-lived memory B cells to short-lived plasma cells independent of secondary lymphoid organs, *Proc. Natl. Acad. Sci.*, 2000, 97:13263-13268.

Orlandi, et al., Cloning immunoglobulin variable domains for expression by the polymerase chain reaction, *Proc. Natl. Acad. Sci. USA*, 1989, 86:3833-3837.

Owens, et al., The genetic engineering of monoclonal antibodies, *J Immunol Methods*. 1994, 168:149-65.

(56)

**References Cited**

## OTHER PUBLICATIONS

Rudikoff, et al., Single amino acid substitution altering antigen-binding specificity, Proc Natl Acad Sci US, 1982, 79:1979-1983.

Sastry, et al., Cloning of the immunological repertoire in *Escherichia coli* for generation of monoclonal catalytic antibodies: construction of a heavy chain variable region-specific cDNA library, Proc. Natl. Acad. Sci., 1989, 86:5728-5732.

Scheid, et al., Broad diversity of neutralizing antibodies isolated from memory B cells in HIV-infected individuals, Nature, 2009, 458:636-640.

Scheid, et al., Broad diversity of neutralizing antibodies isolated from memory B cells in HIV-infected individuals, Nature, 2009, 458:636-640 (supplemental material), <http://www.nature.com/nature/journal/v458/n7238/supinfo/nature07930.html>.

Sehgal, et al., Analyses of single B cells by polymerase chain reaction reveal rearranged VH with germline sequences in spleens of immunized adult rabbits: implications for B cell repertoire maintenance and renewal, J Immunol., 1998, 161:5347-5356.

Slifka, et al., Long-lived plasma cells: a mechanism for maintaining persistent antibody production, Curr Opin Immunol. 1998, 10:252-8.

Spieker-Polet, et al., Rabbit monoclonal antibodies: generating a fusion partner to produce rabbit-rabbit hybridomas, Proc Natl Acad Sci USA, 1995, 92:9348-52.

Takahashi, et al., The direct cloning of the immunoglobulin VH genes from primary cultured B cells specific for a short peptide, J Biotechnol, 1996, 49:201-10.

Wrammert, et al., Rapid cloning of high-affinity human monoclonal antibodies against influenza virus, Nature, 2008, 453:1-6.

Wrammert, et al., Rapid cloning of high-affinity human monoclonal antibodies against influenza virus, Nature, 2008, 453:1-19 (supplemental material).




FIG. 1A









FIG. 1B





CARDMSYAY AYDIW  
CARSGYASS YNLIW  
CARSDSYGC AYDIW  
CARVDSTC YDIW  
OSGYIINI W  
CARSGAGISGIT YFNIN  
CARGCEGYS DNDIN  
CARGMSLD IW  
CVRDSTGISA LFWW  
CARCGLASHE WFINW  
OSGANIENEE FNAIN  
CARSDSHDYD YFNIN  
CARSGSGSHEIF FNINW  
CARSPGGIED AFDW  
CARGWMSIN IW  
CARPLDSYGY AYDIW  
CARYGASVI YFNIN  
CARFHLVIVLV FLDLW  
CARBATHVAVG WLDLW  
CARLGLVIV ININ

Figure 3



>Clone 29--VK SEQ ID NOS: 131 and 132  
atggacacagggcccccaactcagctgtctgggctctctGCTGCTGCTGCTGCCACGGNScc  
M D T R A F T G L L G L L L W L P G A  
atatgtgaccctgtgtgaccccagactccatccctccctgtctgtgagctgtgggagggcaaa  
I C B P V L T Q T P S S V S A A V G G T  
gtccatcaattgcccagtcagtcaggggttggaggaacagctacttatcctgggttt  
V T I N C Q S S Q R V W K N S Y L S W F  
cagggaaaccaggggcccctcccagggcctgatctattatccactctgcccactct  
Q Q K P G Q P P K R L I Y Y T S I P S  
gggtcccctcggggttcaaggcagtggtgttggacacacagctcactctcaccatcagc  
G V P S R F K G S G T Q F T L T I S  
gacctggagtgacatgcttccacttactactgtCTAGGGAGTTATAGTCAATGATAA  
D I E C D D A A I Y Y C I G S Y S D I  
TATCTTtcggggagggaccggaggtggtgtaaaa  
Y S F C G G T E V V V K

**PCR primers:** Reverse for Clone 29--VK AGAATATATATCACTACTATACTCCCTAG (SEQ ID NO: 133)  
Forward for Clone 29--VK CCTCCTCTGGCTCCCGGTTG (SEQ ID NO: 134)

1 2 3 4 5 6 7 8 9 10  
1234567890123456789012345678901234567890123456789012345678901234567  
29F-48 DPVITQIPSSVAAVGGTTTINCQSSQRVWKN5.....YLSWFOQXPGQPPKRIYYTSTTIDSGVDSRPFKGGSGTGTPTTISDIHCDADATYYLGGSYSDDI.....YSPGGGTRFVVVK (SEQ ID NO: 135)  
8L 16 ..... (SEQ ID NO: 135)  
9L-24 ..... (SEQ ID NO: 135)  
16-63 ..... (SEQ ID NO: 135)  
32-20 .....G-Y---K.....V---A.....-GWQ---N..... (SEQ ID NO: 136)

Figure 4B













## METHOD FOR IDENTIFYING LINEAGE-RELATED ANTIBODIES

### CROSS REFERENCE TO RELATED APPLICATION

This patent application is a continuation of U.S. application Ser. No. 13/748,507, filed on Jan. 23, 2013, granted U.S. Pat. No. 8,969,013, which is a continuation of U.S. application Ser. No. 13/552,517, filed Jul. 18, 2012, granted U.S. Pat. No. 8,617,830, which is a continuation of U.S. application Ser. No. 12/878,925, filed on Sep. 9, 2010, granted U.S. Pat. No. 8,293,483, which claims the priority benefit of U.S. provisional application Ser. No. 61/241,714, filed on Sep. 11, 2009, all of which are incorporated by reference herein in their entirety.

### INTRODUCTION

Antibodies are proteins that bind a specific antigen. Generally, antibodies are specific for their targets, have the ability to mediate immune effector mechanisms, and have a long half-life in serum. Such properties make antibodies powerful therapeutics. Monoclonal antibodies are used therapeutically for the treatment of a variety of conditions including cancer, inflammation, and other diseases. There are currently over two dozen therapeutic antibody products on the market and hundreds in development.

There is a constant need for new antibodies and methods for making the same.

### SUMMARY

In certain embodiments, the method may comprise: a) obtaining the antibody heavy chain sequences and the antibody light chain sequences from a population of B cells of an animal, wherein the population of B cells is enriched for B cells that produce antibodies that specifically bind to a target antigen; b) grouping the heavy and light chain sequences on the basis of sequence similarity to provide a plurality of groups of antibodies that are related by lineage; c) testing a single antibody from each of the groups in a first bioassay to identify a first antibody that has a biological activity; and, after the first antibody has been identified, d) testing further antibodies that are in the same group as the first antibody in a second bioassay, thereby identifying a second antibody that has the biological activity.

In other embodiment, the method may comprise: a) testing a plurality of antibodies obtained from a first portion of an antibody producing organ of an animal in a first bioassay to identify a first antibody that has a biological activity; b) obtaining the sequence of the first antibody; c) obtaining from a second portion of said antibody producing organ the heavy and light chain amino acid sequences of further antibodies that are related by lineage to said first antibody by PCR, using probes are designed using the sequence of the first antibody; and, c) testing a plurality of the further antibodies in a second bioassay to identify a second antibody that has said biological activity.

In certain embodiments, the method provides a means by which significant portion of the entire antibody repertoire of an animal can be screened to identify an antibody with desirable properties. In certain embodiments the method involves first identifying a single antibody with desirable properties, and then screening other antibodies in same lineage group (i.e., a clonally related group of antibodies) as the identified antibody, to identify other antibodies that may have even

more desirable properties relative to the identified antibody. As such, the method provides an efficient way to screen for and identify new, biologically active antibodies. After identification, the second antibody may be tested in further assays, and, if it is suitable for use as a therapy, may be humanized, for example.

### BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A and 1B schematically illustrate two embodiments.

FIG. 2 shows the amino acid sequences of selected KDR-binding antibodies. Page 1 of FIG. 2 shows amino acid sequences of the heavy chains. Page 2 of FIG. 2 shows amino acid sequences of the corresponding light chains. The amino acid sequences shown in FIG. 2 are of antibodies that specifically bind to KDR and block VEGF activity. From top to bottom, FIG. 2 (page 1 of 2) SEQ ID NOS: 1-47 and FIG. 2 (page 2 of 2) SEQ ID NOS: 48-94.

FIG. 3 shows the amino acid sequence of 20 exemplary VH3 regions of unrelated rabbit antibodies. From top to bottom SEQ ID NOS: 95-114.

FIGS. 4A-4H show exemplary methods by which related antibodies can be amplified.

### DEFINITIONS

Before the present subject invention is described further, it is to be understood that this invention is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

It must be noted that as used herein and in the appended claims, the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "an antibody" includes a plurality of such antibodies and reference to "a framework region" includes reference to one or more framework regions and equivalents thereof known to those skilled in the art, and so forth.

The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

The term "nucleic acid" encompasses DNA, RNA, single stranded or double stranded and chemical modifications thereof. The terms "nucleic acid" and "polynucleotide" are used interchangeably herein.

The term "expression", as used herein, refers to the process by which a polypeptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation.

The term "expression cassette" refers to a nucleic acid construct capable of directing the expression of a gene/coding sequence of interest, which is operably linked to a promoter of the expression cassette. Such cassettes can be a linear nucleic acid or can be present in a "vector", "vector construct", "expression vector", or "gene transfer vector", in order to transfer the expression cassette into target cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

The term "operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, a given signal peptide that is operably linked to a polypeptide directs the secretion of the polypeptide from a cell. In the case of a promoter, a promoter that is operably linked to a coding sequence will direct the expression of the coding sequence. The promoter or other control elements need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. For example, intervening untranslated yet transcribed sequences can be present between the promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

The term "plurality" refers to more than 1, for example more than 2, more than about 5, more than about 10, more than about 20, more than about 50, more than about 100, more than about 200, more than about 500, more than about 1000, more than about 2000, more than about 5000, more than about 10,000, more than about 20,000, more than about 50,000, more than about 100,000, usually no more than about 200,000. A "population" contains a plurality of items.

The term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection", or "transformation", or "transduction" and includes reference to the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell wherein the nucleic acid sequence may be present in the cell transiently or may be incorporated into the genome of the cell (e.g., chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon.

The terms "antibody" and "immunoglobulin" are used interchangeably herein. These terms are well understood by those in the field, and refer to a protein consisting of one or more polypeptides that specifically binds an antigen. One form of antibody constitutes the basic structural unit of an antibody. This form is a tetramer and consists of two identical pairs of antibody chains, each pair having one light and one heavy chain. In each pair, the light and heavy chain variable regions are together responsible for binding to an antigen, and the constant regions are responsible for the antibody effector functions.

The recognized immunoglobulin polypeptides include the kappa and lambda light chains and the alpha, gamma (IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>3</sub>, IgG<sub>4</sub>), delta, epsilon and mu heavy chains or equivalents in other species. Full-length immunoglobulin "light chains" (of about 25 kDa or about 214 amino acids) comprise a variable region of about 110 amino acids at the NH<sub>2</sub>-terminus and a kappa or lambda constant region at the COOH-terminus. Full-length immunoglobulin "heavy chains" (of about 50 kDa or about 446 amino acids), similarly comprise a variable region (of about 116 amino acids) and one of the aforementioned heavy chain constant regions, e.g., gamma (of about 330 amino acids).

The terms "antibodies" and "immunoglobulin" include antibodies or immunoglobulins of any isotype, fragments of antibodies which retain specific binding to antigen, including, but not limited to, Fab, Fv, scFv, and Fd fragments, chimeric

antibodies, humanized antibodies, single-chain antibodies, and fusion proteins comprising an antigen-binding portion of an antibody and a non-antibody protein. The antibodies may be detectably labeled, e.g., with a radioisotope, an enzyme which generates a detectable product, a fluorescent protein, and the like. The antibodies may be further conjugated to other moieties, such as members of specific binding pairs, e.g., biotin (member of biotin-avidin specific binding pair), and the like. The antibodies may also be bound to a solid support, including, but not limited to, polystyrene plates or beads, and the like. Also encompassed by the term are Fab', Fv, F(ab')<sub>2</sub>, and other antibody fragments that retain specific binding to antigen, and monoclonal antibodies.

Antibodies may exist in a variety of other forms including, for example, Fv, Fab, and (Fab')<sub>2</sub>, as well as bi-functional (i.e. bi-specific) hybrid antibodies (e.g., Lanzavecchia et al., *Eur. J. Immunol.* 17, 105 (1987)) and in single chains (e.g., Huston et al., *Proc. Natl. Acad. Sci. U.S.A.*, 85, 5879-5883 (1988) and Bird et al., *Science*, 242, 423-426 (1988), which are incorporated herein by reference). (See, generally, Hood et al., "Immunology", Benjamin, N.Y., 2nd ed., 1984, and Hunkapiller and Hood, *Nature*, 323, 15-16, 1986).

An immunoglobulin light or heavy chain variable region consists of a framework region (FR) interrupted by three hypervariable regions, also called "complementarity determining regions" or "CDRs". The extent of the framework region and CDRs have been precisely defined (see, "Sequences of Proteins of Immunological Interest," E. Kabat et al., U.S. Department of Health and Human Services, 1991). The sequences of the framework regions of different light or heavy chains are relatively conserved within a species. The framework region of an antibody, that is the combined framework regions of the constituent light and heavy chains, serves to position and align the CDRs. The CDRs are primarily responsible for binding to an epitope of an antigen.

The term "chimeric antibodies" refer to antibodies whose light and heavy chain genes have been constructed, typically by genetic engineering, from antibody variable and constant region genes belonging to different species. For example, the variable segments of the genes from a mouse monoclonal antibody may be joined to human constant segments, such as gamma 1 and gamma 3. An example of a therapeutic chimeric antibody is a hybrid protein composed of the variable or antigen-binding domain from a rabbit antibody and the constant or effector domain from a human antibody, although other mammalian species may be used.

The term "humanized antibody" or "humanized immunoglobulin" refers to a non-human (e.g., mouse or rabbit) antibody containing one or more amino acids (in a framework region, a constant region or a CDR, for example) that have been substituted with a correspondingly positioned amino acid from a human antibody. In general, humanized antibodies produce a reduced immune response in a human host, as compared to a non-humanized version of the same antibody.

The terms "polypeptide" and "protein", used interchangeably herein, refer to a polymeric form of amino acids of any length, which can include coded and non-coded amino acids, chemically or biochemically modified or derivatized amino acids, and polypeptides having modified peptide backbones.

The term "natural" antibody refers to an antibody in which the heavy and light chains of the antibody have been made and paired by the immune system of a multi-cellular organism. Spleen, lymph nodes, bone marrow and serum are examples of tissues that produce natural antibodies. For example, the antibodies produced by the antibody producing cells isolated from a first animal immunized with an antigen are natural antibodies.

The term “non-naturally paired”, with respect to VH and VL chains of an engineered antibody, refers to a VH and VL pair that is not found in a natural antibody. Thus, a non-naturally paired antibody is a combination of VH and VL chain of two different natural antibodies. The VH and VL chains of a non-naturally paired antibody are not mutated relative to the VH and VL chains of the two different antibodies which provided the VH and VL chains. For example, the “non-naturally paired” IgH and IgL chains of the engineered antibody may contain the IgH variable chain from a first antibody producing cell obtained from an animal and the IgL variable chain of second antibody producing cell obtained from the same animal, where the amino acid sequence of the antibody produced by the first cell is different from the amino acid sequence of the antibody produced by the second cell. In this example, the IgH and IgL chains may be from the same lineage group. An antibody containing “non-naturally paired” IgH and IgL chains may or not be made by phage display. As such, antibodies may or may not contain viral (e.g., bacteriophage M13)-derived sequences.

The term “lineage-related antibodies” and “antibodies that related by lineage” as well as grammatically-equivalent variants thereof, are antibodies that are produced by cells that share a common B cell ancestor. Related antibodies produced by related antibody producing cells bind to the same epitope of an antigen and are typically very similar in sequence, particularly in their L3 and H3 CDRs. Both the H3 and L3 CDRs of lineage-related antibodies have an identical length and a near identical sequence (i.e., differ by up to 5, i.e., 0, 1, 2, 3, 4 or 5 residues). In certain cases, the B cell ancestor contains a genome having a rearranged light chain VJC region and a rearranged heavy chain VDJC region, and produces an antibody that has not yet undergone affinity maturation. “Naïve” or “virgin” B cells present in spleen tissue, are exemplary B cell common ancestors. Related antibodies are related via a common antibody ancestor, e.g., the antibody produced in the naïve B cell ancestor. The term “related antibodies” is not intended to describe a group of antibodies that are not produced by cells that arise from the same ancestor B-cell. A “lineage group” contains a group of antibodies that are related to one another by lineage.

The terms “treating” or “treatment” of a condition or disease refer to providing a clinical benefit to a subject, and include: (1) preventing at least one symptom of the conditions, i.e., causing a clinical symptom to not significantly develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease, (2) inhibiting the disease, i.e., arresting or reducing the development of the disease or its symptoms, or (3) relieving the disease, i.e., causing regression of the disease or its clinical symptoms.

#### DETAILED DESCRIPTION OF EXEMPLARY EMBODIMENTS

One embodiment of the subject method is illustrated in FIG. 1A. With reference to FIG. 1A, this embodiment of the method may involve immunizing an antibody-producing animal with a selected antigen, and enriching from a larger population of antibody-producing cells that bind to the antigen. In FIG. 1, six different antibody producing cells A-F that produce antibodies that bind to a target antigen are enriched from a larger population of antibody producing cells. However, in many embodiments, there may be several hundred or several thousand enriched cells. Each of these cells produces a natural antibody that contains a naturally paired IgH and IgL chain. The amino acid sequences of the heavy and light chains

of the antibodies produced by the enriched cells are obtained by sequencing the nucleic acids encoding the IgH and IgL chains of the antibodies, and the sequences are analyzed and put into lineage groups which, as discussed above, are groups of antibodies that are produced by cells that share a common B cell ancestor. Such antibodies generally have very similar sequences, and have H3 CDRs of identical length and near identical sequence as well as L3 CDRs of identical length and a near identical sequence. In the embodiment shown in FIG. 1A, the six antibody producing cells produce antibodies (AbA to AbF) that are in two lineage groups (i.e., lineage groups 1 and 2, where AbA and AbB are in lineage group 1 and AbC, AbD, AbE and AbF are in lineage group 2). After the antibodies have been placed into lineage groups, a single antibody (or, in certain cases, multiple antibodies (for example 2-10) from each lineage group) from at least one of the lineage group, e.g., AbA from lineage group 1 and AbC from lineage group 2, is selected for testing in a bioassay, where a bioassay identifies an antibody with a biological activity (e.g., a blocking or neutralizing activity). Once an antibody having a biological activity has been identified, e.g., AbC, other antibodies from the same lineage group as the identified antibody are tested to identify a second antibody that has the same biological activity as the first antibody. In the example shown in FIG. 1, antibodies D, E and F, which belong to the same lineage group as antibody C, were tested.

Many warm-blooded animals, in particular mammals such as humans, rabbits, mice, rats, sheep, cows, pigs and ayes such as chickens and turkeys, may be used as a source of antibody-produced cells. However, in certain embodiments a rabbit or mice is used because of their ease in handling, well-defined genetic traits, and the fact that they may be readily sacrificed. Procedures for immunizing animals are well known in the art, and are described in Harlow et al., (*Antibodies: A Laboratory Manual*, First Edition (1988) Cold Spring Harbor, N.Y.).

Suitable antigens include extracellularly-exposed fragments of Her2, GD2, EGFR, CEA, CD52, CD20, Lym-1, CD6, complement activating receptor (CAR), EGP40, VEGF, tumor-associated glycoprotein TAG-72 AFP (alpha-fetoprotein), BlyS (TNF and APOL—related ligand), CA125 (carcinoma antigen 125), CEA (carcinoembryonic antigen), CD2 (T-cell surface antigen), CD3 (heteromultimer associated with the TCR), CD4, CD11a (integrin alpha-L), CD14 (monocyte differentiation antigen), CD20, CD22 (B-cell receptor), CD23 (low affinity IgE receptor), CD25 (IL-2 receptor alpha chain), CD30 (cytokine receptor), CD33 (myeloid cell surface antigen), CD40 (tumor necrosis factor receptor), CD44v6 (mediates adhesion of leukocytes), CD52 (CAMPATH-1), CD80 (costimulator for CD28 and CTLA-4), complement component C5, CTLA, EGFR, eotaxin (cytokine A11), HER2/neu, HLA-DR, HLA-DR10, HLA Class II, IgE, GPIIb/IIIa (integrin), Integrin  $\alpha$ V $\beta$ 3, Integrins  $\alpha$ 4 $\beta$ 1 and  $\alpha$ 4 $\beta$ 7, Integrin  $\beta$ 2, IFN-gamma, IL-1 $\beta$ , IL-4, IL-5, IL-6R (IL6 receptor), IL-12, IL-15, KDR (VEGFR-2), lewisy, mesothelin, MUC1, MUC18, NCAM (neural cell adhesion molecule), oncofetal fibronectin, PDGF $\beta$ R (Beta platelet-derived growth factor receptor), PMSA, renal carcinoma antigen G250, RSV, E-Selectin, TGF $\beta$ 1, TGF $\beta$ 2, TNF $\alpha$ , TRAIL-R1, VAP-1 (vascular adhesion protein 1) or TNF $\alpha$ , or the like. In many embodiments, a peptide having the amino acid sequence corresponding to a portion of an extracellular domain of one of the above-listed proteins is employed as an antigen.

Antibody-producing cells may also be obtained from a subject which has generated the cells during the course of a selected disease or condition. For instance, antibody-produc-

ing cells from a human with a disease of unknown cause, such as rheumatoid arthritis, may be obtained and used in an effort to identify antibodies which have an effect on the disease process or which may lead to identification of an etiological agent or body component that is involved in the cause of the disease. Similarly, antibody-producing cells may be obtained from subjects with disease due to known etiological agents such as malaria or AIDS. These antibody-producing cells may be derived from the blood, lymph nodes or bone marrow, as well as from other diseased or normal tissues. Antibody-producing cells may also be prepared from blood collected with an anticoagulant such as heparin or EDTA. The antibody-producing cells may be further separated from erythrocytes and polymorphs using standard procedures such as centrifugation with Ficoll-Hypaque (Pharmacia, Uppsala, Sweden). Antibody-producing cells may also be prepared from solid tissues such as lymph nodes or tumors by dissociation with enzymes such as collagenase and trypsin in the presence of EDTA.

In exemplary embodiments, an affinity purification method is utilized to isolate antibody producing cells that produce antibodies that bind to an antigen. The antigen with which the animal was immunized may be immobilized on a solid phase and used to selectively retain antibody producing cells that express an antibody on their surface that binds to the antigen, while other cells are washed away. The retained cells may then be eluted by a variety of methods, such as by using an excess of the antigen, chaotropic agents, changing the pH, salt concentration, etc. Any of the well known methods for immobilizing or coupling antigen to a solid phase may be used. For example, when the antigen is a cancer cell, appropriately treated microtiter plate that will bind to cells may be used, such as microtiter plates for cell culture. In the instances where the antigen is a protein, the protein may be covalently attached to a solid phase, for example, sepharose beads, by well known techniques, etc. Alternatively, a labeled antigen may be used to specifically label cells that express an antibody that binds to the antigen and the labeled cells may then be isolated by cell sorting (e.g., by FACS). In certain cases, methods for antibody purification may be adapted to isolate antibody producing cells. Such methods are well known and are described in, for example, *J Immunol Methods*. 2003 November; 282(1-2):45-52; *J Chromatogr A*. 2007 Aug. 10; 1160(1-2):44-55; *J Biochem Biophys Methods*. 2002 May 31; 51(3):217-31. Cells may also be isolated using magnetic beads or by any other affinity solid phase capture method, protocols for which are known. In some embodiments, antigen-specific antibody producing cells may be obtained from blood by flow cytometry using the methods described in Wrammert (*Nature* 2008 453: 667-672), Scheid (*Nature* 2009 458: 636-640), Tiller (*J. Immunol. Methods* 2008 329 112-124) or Scheid (*Proc. Natl. Acad. Sci.* 2008 105: 9727-9732), for example, which are incorporated by reference for disclosure of those methods. Exemplary antibody-producing cell enrichment methods include performing flow cytometry (FACS) of cell populations obtained from a spleen, bone marrow, lymph node or other lymph organs, e.g., through incubating the cells with labeled anti-rabbit IgG and sorting the labeled cells using a FACSVantage SE cell sorter (Becton-Dickinson, San Jose, Calif.). In some embodiments, single or nearly single antibody-producing cells are deposited in microtiter plates. If the FACS system is employed, sorted cells may be deposited after enrichment directly into a microtiter plate.

Enrichment may decrease the size of the cell population by at least 50%, e.g., at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 99% and in certain cases, the

plurality of enriched antibody producing cells may be substantially pure, i.e., substantially free of other cells that do not produce an antibody that binds to the antigen, where the term "substantially pure" refers to an isolated population of antibody producing cells, in which cells that express antibodies that specifically bind to the antigen make up at least 5%, 10%, 20%, 30%, at least 40%, at least 50%, at least 60%, at least 70% or more of the total population of cells. The enriched population of antibody producing cells may be employed as a mixture of cells, or alternatively, they may be used as single cells, e.g., by dilution and deposition into individual wells of a microtiter plate.

The enriched population of antibody producing cells may comprise at least 5, at least 10, at least 30, at least 60, at least 100, at least 300, at least 500, at least 1000, at least 5,000, at least 10,000 or at least 100,000, or more antibody producing cells.

The isolated antibody-producing cells may be optionally cultured (i.e. grown in media that supports at least one, at least 5 or at least 10 or more cell divisions of the cell) by methods known to one of skill in the art after they have been deposited (see e.g. WO 01/55216).

In certain embodiments, the antibodies produced by the enriched cells are not well characterized. As such, although the antibody-producing cells are isolated based on the production of antibodies that specifically bind to the antigen, the epitope(s) to which these antibodies bind is unknown, and it is not known if the antibodies have any biological activity (e.g., a neutralizing or blocking activity). Additionally, the nucleic acid sequence or the amino acid sequence of the variable regions of IgH and IgL chains of these antibodies are not known.

Sequences encoding heavy and light chains may be amplified from the cDNA using techniques well known in the art, such as Polymerase Chain Reaction (PCR). See Mullis, U.S. Pat. No. 4,683,195; Mullis et al., U.S. Pat. No. 4,683,195; *Polymerase Chain Reaction: Current Communication in Molecular Biology*, Cold Springs Harbor Press, Cold Spring Harbor, N.Y., 1989. Briefly, cDNA segments encoding the variable domain of the antibody are exponentially amplified by performing sequential reactions with a DNA polymerase. The reaction is primed by a 5' primer and a 3' DNA primer. In some embodiments, the 3' antisense primer corresponding to a DNA sequence in the constant (or joining) region of the immunoglobulin chain and the 5' primer (or panel of related primers) corresponding to a DNA sequence in the variable region of the immunoglobulin chain. This combination of oligonucleotide primers has been used in the PCR amplification of murine immunoglobulin cDNAs of unknown sequence (see Sastry et al., *Proc Natl. Acad. Sci.* 86:5728-5732, 1989 and Orlandi et al., *Proc. Natl. Acad. Sci.* 86:3833-3837, 1989). Alternatively, an "anchored polymerase chain reaction" may be performed (see Loh et al., *Science* 243:217-220, 1989). In this procedure, the first strand cDNA is primed with a 3' DNA primer as above, and a poly(dG tail) is then added to the 3' end of the strand with terminal deoxynucleotidyl transferase. The product is then amplified by PCR using the specific 3' DNA primer and another oligonucleotide consisting of a poly(dC) tail attached to a sequence with convenient restriction sites. In many embodiments, however, the entire polynucleotide encoding a heavy or light chain is amplified using primers spanning the start codons and stop codons of both of the immunoglobulin cDNAs, however, depending on the amplification products desired, suitable primers may be used. Exemplary primers for use with rabbit antibody-producing cells are as follows: heavy chain, 5' end (CACCATGGAGACTGGGCTGCGCTGGCTTCTCCTG-

GTCGCTGTG; SEQ ID NO:177); heavy chain, 3' end (CTC-CCGCTCTCCGGGTAAATGAGCGCTGTGCCGGCGA; SEQ ID NO:178); light chain kappa, 5' end (CAGGCAGGACCCAGCATGGACACGAGGGCCCCCACT; SEQ ID NO:179); and L kappa, 3' end (TCAATAGGGGTGACTGT-TAGAGCGAGACGCCTGC; SEQ ID NO:180). Suitable restriction sites and other tails may be engineered into the amplification oligonucleotides to facilitate cloning and further processing of the amplification products. Amplification procedures using nested primers may also be used, where such nested primers are well known to one of skill in the art. Exemplary methods for amplifying antibody-encoding nucleic acid is also described in Wrammert (Nature 2008 453: 667-672) and Scheid (Nature 2009 458: 636-640), for example. In this embodiment, the enriched cells may be combined before sequencing (in which case the initial amplification product will contain a mixture of a plurality of different products that can be discriminated by cloning the products or using single molecule sequencing technologies), or the cells may be kept separate from one another (in which case the initial amplification product amplified from a single cell may contain a single species that can be sequenced).

In certain embodiments, at least 1,000 heavy chain sequences and at least 1,000 light chain sequences are obtained.

Once the amino acid sequence of heavy and light chains of the antibodies has been obtained, antibodies can be grouped on the basis of sequence similarity to provide a plurality of groups of antibodies that are related by lineage. Methods for performing clonal analysis of antibody sequences are well known and are described in a number of publications including Magori-Cohen (Bioinformatics 2006 22: e332-40), Manske (Clin. Immunol. 2006 120:106-20), Kleinstein (J. Immunol. 2003 171: 4639-49), Clement (Mol. Ecol. 2000 9: 1657-1659), Mehr (J. Immunol. 2004 172 4790-6), Wrammert (Nature 2008 453: 667-672), Scheid (Nature 2009 458: 636-640), which are incorporated by reference herein for disclosure of those methods. The antibodies placed into lineage groups should all be from a single animal, i.e., an individual mouse or rabbit.

In some embodiments, the amino acid positions of an antibody are numbered using a suitable numbering system, such as that provided by Chothia (JMol Biol 1998; 278: 457-79) or Kabat (1991, Sequences of Proteins of Immunological Interest, DHHS, Washington, DC). CDR and/or framework residues may be identified using these methods. The numbered sequences may be aligned by eye, or by employing an alignment program such as one of the CLUSTAL suite of programs (Thompson et al Nucleic Acids Research, 22:4673-4680). The variable regions of antibodies within a related group of antibodies have amino acid sequences that are very similar. For example, the VH or VL domains of antibodies within a related group of antibodies may have amino acid sequences that are at least about 80% identical (e.g., at least 85% identical, at least 90% identical, at least 95% or at least 98% or at least 99% identical), ignoring any gaps or insertions made to facilitate alignment of the sequences. Antibodies within a related group of antibodies have a VL domains that are similar to each other, as well as VH domains that are similar to each other. In other words, in certain embodiments the VH or VL domains of two different related antibodies usually contain up to about ten (i.e., one, two, three, four or five or more) amino acid differences. An amino acid difference may be present at any position of the variable domain, including in any CDR or in any framework region. Certain related antibodies have H3 CDRs that are almost identical, as well as L3 CDRs that are almost identical. In these embodiments, any two antibodies

that are related will have L3 and H3 CDRs that are each identical in length and have near identical sequences (i.e., that contain 0, 1, 2, 3, 4 or 5 amino acid changes). In other words the L3 CDRs of the two antibodies are identical in length and near identical in sequence and the H3 CDRs of the two antibodies are identical in length and near identical in sequence. Two exemplary sets of related antibodies are shown in FIG. 2, and the sequences of 20 exemplary VH3 regions of unrelated rabbit antibodies are shown for comparison in FIG. 3.

In certain embodiments, the heavy chain sequences may or may not be grouped independently of the light chain sequences. If the heavy and light chain sequences are grouped independently of one another, the heavy and light chain groups may be matched up by analysis of lineage trees.

Depending how many sequences are obtained, in certain embodiments the enriched antibodies may be grouped into at least 5 groups, at least 10 groups, at least 20 groups, at least 50 groups, or at least 100 groups or more, e.g., up to 200 or 500 groups or more. Depending how many sequences are obtained, each group may contain from 2 to several hundred or more antibodies.

Once the antibodies have been grouped, a single antibody from each of at least some of the groups (e.g., at least 20%, at least 50 or at least 80% of the groups) is tested in a first bioassay to identify a first antibody that has a biological activity. The bioassay may determine whether the antibody has a biological effect, e.g., an ability to inhibit an interaction between a receptor and an a ligand by either binding to the receptor and blocking binding of the ligand, or by binding to the ligand and neutralizing it, or by promoting or inhibiting a cellular phenotype, e.g., cell growth, cell proliferation, cell migration, cell viability (e.g., apoptosis), cell differentiation, cell adherence, cell shape changes (e.g., tubular cell formation), complement dependant cytotoxicity CDC, antibody-dependent cell-mediated cytotoxicity ADCC, receptor activation, gene expression changes, changes in post-translational modification (e.g., phosphorylation), changes in protein targeting (e.g., NFκB localization etc.), etc., or inhibition of receptor multimerization (e.g., dimer or trimerization) or receptor-ligand interactions, etc. Such bioassays are well known in the art. The term "bioassay" is intended to exclude assays in which only the ability of an antibody to bind to a target is read. Bioassays useful in this method are numerous, and include but are not limited to cellular assays in which a cellular phenotype is measured, e.g., gene expression assays; and in vivo assays that involve a particular animal (which, in certain embodiments may be an animal model for a condition related to the target). In certain cases, the assay may be a vascularization assay.

In this embodiment, the antibodies tested in the bioassay may contain naturally paired heavy and light chain variable domains, or non-naturally paired heavy and light chains (i.e., heavy and light chain variable domains from different antibodies of the same lineage group). Since the antibodies are from the same lineage group, it is expected that such antibodies will be functional.

After a first antibody that has a biological activity has been identified, further antibodies that are in the same lineage group as the first antibody are tested in a second bioassay, thereby identifying a second antibody that has the same biological activity as the first antibody. In certain cases at least 10%, at least 20%, at least 50%, or at least 80% of the antibodies in the same lineage group are tested. The first bioassay may be the same as or different to the second bioassay. In certain embodiments, a plurality of antibodies is tested, and the antibody with the best properties is chosen for future use.

In particular embodiments, the further antibodies may contain naturally paired heavy and light chain variable domains, or non-naturally paired heavy and light chain variable domains (i.e., heavy and light chain variable domains from different antibodies of the same lineage group). Since the antibodies are from the same lineage group, it is expected that such antibodies will be functional. In particular embodiments, the pairing of the heavy and light chains may be systematic (e.g., every heavy chain is tested in combination with every light chain) or random (e.g., every heavy chain is tested with randomly selected light chains), for example.

Exemplary VEGF bioassays include assays using isolated protein in a cell free systems, *in vitro* using cultured cells or *in vivo* assays. Exemplary VEGF assays include, but are not limited to a receptor tyrosine kinase inhibition assay (see, e.g., Cancer Research Jun. 15, 2006; 66:6025-6032), an *in vitro* HUVEC proliferation assay (FASEB Journal 2006; 20:2027-2035), an *in vivo* solid tumor disease assay (U.S. Pat. No. 6,811,779) and an *in vivo* angiogenesis assay (FASEB Journal 2006; 20:2027-2035). These assays are well known in the art. The descriptions of these assays are hereby incorporated by reference.

Exemplary TNF- $\alpha$  bioassays include *in vitro* assays using cell free systems or using cultured cells or *in vivo* assays. As such, TNF- $\alpha$  assays include *in vitro* human whole blood assay and cell mediated cytotoxicity assay (U.S. Pat. No. 6,090,382), *in vitro* tumor human killing assay (see, e.g., published U.S. patent application 20040185047), *in vivo* tumor regression assay (U.S. patent application 20040002589). Additional TNF- $\alpha$  assays are described in a variety of publications, including 20040151722, 20050037008, 20040185047, 20040138427, 20030187231, 20030199679, and Balazovich (Blood 1996 88: 690-696).

A subject antibody inhibits at least one activity of its target in the range of about 20% to 100%, e.g., by at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, usually up to about 70%, up to about 80%, up to about 90% or more. In certain assays, a subject antibody may inhibit its target with an  $IC_{50}$  of  $1 \times 10^{-7}$  M or less (e.g.,  $1 \times 10^{-7}$  M or less,  $1 \times 10^{-8}$  M or less,  $1 \times 10^{-9}$  M or less, usually to  $1 \times 10^{-12}$  M or  $1 \times 10^{-13}$  M). In assays in which a mouse is employed, a subject antibody may have an  $ED_{50}$  of less than 1  $\mu$ g/mouse (e.g., 10 ng/mouse to about 1  $\mu$ g/mouse). In certain embodiments, a subject antibody may be contacted with a cell in the presence of a ligand, and a ligand response phenotype of the cell is monitored.

In certain embodiments, particularly if the antigen elicits a strong response in the animal, the method may be practiced in the absence of any antigen-based enrichment of antibody producing cells prior to the first bioassay. In these embodiments, the method may involve: a) obtaining the antibody heavy chain sequences and the antibody light chain sequences from a population of B cells of an animal, wherein said population of B cells is not enriched for B cells that produce antibodies that specifically bind to a target antigen, b) grouping the heavy and light chain sequences on the basis of sequence similarity to provide a plurality of groups of antibodies that are related by lineage; c) testing a single antibody from each of the groups in a first bioassay to identify a first antibody that has a biological activity; and, after the first antibody has been identified; and d) testing further antibodies that are in the same group as the first antibody in a second bioassay, thereby identifying a second antibody that has the biological activity.

Another embodiment of is illustrated in FIG. 1B. With reference to FIG. 1B, this embodiment of the subject method may involve immunizing an antibody-producing animal with

a selected antigen, and testing a plurality of antibodies produced by a first portion of an antibody producing organ of the animal (e.g., a first portion of the spleen, a first portion of the lymph nodes, a first portion of bone marrow, or a first portion of the peripheral blood mononuclear cell (PBMC) population in the bloodstream of the animal, etc.) in a bioassay to identify a first antibody that has a biological activity. In this embodiment, the first and second portions of an antibody-producing organ need not be spatially separated in the organ. Rather, since a first portion or an organ can be made by, for example, making a single cell suspension of the organ and then removing part of the suspension, the first and second portions of an organ may be interspersed with one another in the organ. In the example shown in FIG. 1B, antibody A is identified as having a biological activity. The nucleotide sequence encoding the IgH and IgL chain of the antibody is obtained. Based on these sequences, PCR primers that are specific for the heavy and light chains of antibodies that are in the same lineage group as the identified antibody are designed, and used to obtain from the second portion of the antibody producing organ the sequences of further antibodies that are in the same lineage group as the identified antibody. The further antibodies are tested, and a second antibody from the same lineage group and also having the same biological activity as the first antibody is identified.

Many exemplary aspects of this alternative method, e.g., which antigens and bioassays can be employed in the method, etc., are discussed above. In certain embodiments, a lead antibody obtained from a first portion of an antibody-producing organ is identified using a bioassay. In this embodiment, the antibodies obtained from the first portion of the organ are screened using a hybridoma-based method or by a method that does not require production of hybridomas, e.g., by phage display or by the method described in US20040067496 and other references, for example, to identify a biologically active antibody. In one embodiment, a portion of the splenocytes of a spleen of a single animal is fused with a fusion partner to produce hybridomas that are then screened to identify a biologically active antibody. In another embodiment, heavy and light chain sequences are directly amplified from PBMCs, and recombinant antibodies are expressed in a different cell (e.g., as described in US20040067496) prior to screening. In another embodiment, a phage display library is constructed from the RNA made from a portion of the spleen of an animal, and the phage display library is screened. The first, biologically active antibody is identified, and the nucleic acid encoding that antibody is sequenced.

In certain embodiments, polynucleotides encoding the variable heavy and variable light domains of lineage-related antibodies may be amplified from the same animal as the first antibody by "CDR-anchored PCR", i.e., using pairs of primers that each contains a primer that is complementary to a CDR-encoding region of the parent antibody cDNA. In these embodiments, the method may include: a) obtaining the nucleotide sequences of: i. a heavy chain-encoding nucleic acid that encodes the variable heavy chain of a first antibody of an immunized animal; and ii. a variable light chain-encoding nucleic acid that encodes the light chain of the first antibody; b) obtaining the amino acid sequence of the variable domains of the heavy and light chains of further antibodies from the immunized animal, using: i. a first primer pair that includes a first primer that is complementary to a CDR-encoding region of the heavy chain-encoding nucleic acid; and ii. a second primer pair that includes a second primer that is complementary to a CDR-encoding region of the light chain-encoding nucleic acid. After the amino acid sequences of the variable domains of the further antibodies have been

determined by translation of the obtained nucleotide sequences, the amino acid may be analyzed using the above methods to confirm that they are related by lineage to the first antibody (e.g., analyzed to determine whether the amino acid sequences of the heavy and light chains are at least 80% identical to those of the parent antibody and whether the heavy and light chain CDR3 regions are of identical length of near identical sequence etc. as discussed above).

As would be readily apparent, a variety of techniques are available for amplifying sequences that encode further antibodies from an animal after the nucleotide sequence encoding a first antibody has been obtained from that animal. For example, sequences encoding the heavy and light chains of the second antibody may be amplified using inverse PCR (e.g., using two primers that face away from each other) or by anchored PCR using a specific (where a specific primer may be complementary to a different sequence of the first antibody, e.g., a different CDR sequence) or "universal" primer (where a universal primer is complementary to a sequence that is present in a plurality of different antibody-encoding polynucleotides), where one of the primers is complementary to first CDR-encoding region using cDNA as a template. In certain cases, a universal primer may be complementary to a sequence that is in at least 10% (e.g., at least 20% at least 40% at least 50% or at least 80%) of all heavy or light chain encoding cDNAs obtainable from the animal (e.g., complementary to nucleic acid encoding a conserved sequence that is present in the constant region or secretion signal of the antibodies). In other embodiments, the universal primer may be complementary to flanking sequences in the vector into which cDNA from the animal is cloned or to linkers ligated onto the cDNA, for example.

In one embodiment, two amplification reactions are performed using cDNA as a template, where the first reaction amplifies the heavy chain variable domain-encoding nucleic acid for the second antibody and the second reaction amplifies the light chain variable domain-encoding nucleic acid for the second antibody. In this embodiment: a) the first reaction uses: i. a CDR-specific primer that is complementary to a CDR-encoding region (i.e., the CDR1, CDR2 or CDR3 region) of the heavy chain-encoding nucleic acid of the first antibody and ii. a universal second primer that is complementary to a non-variable domain-encoding region of the antibody heavy chain cDNA, e.g., to a sequence that encodes the constant domain or secretion signal of the heavy chain of the first antibody, as illustrated in the examples section of this disclosure; and b) the second reaction uses i. a CDR-specific primer that is complementary to a CDR-encoding region (i.e., the CDR1, CDR2 or CDR3 region) of the light chain-encoding nucleic acid of the first antibody and ii. a universal second primer that is complementary to a non-variable domain-encoding region of the antibody light chain cDNA, e.g., to a sequence that encodes the constant domain or secretion signal of the light chain of the first antibody, as illustrated in the examples section of this disclosure.

Several strategies for cloning antibody sequences by PCR are known and may be readily adapted for use in the instant method (e.g., by using a CDR-specific primer in addition to a disclosed primer). Such strategies include those described by: LeBoeuf (*Cloning and sequencing of immunoglobulin variable-region genes using degenerate oligodeoxyribonucleotides and polymerase chain reaction*. Gene. 1989 82:371-7), Dattamajumdar (*Rapid cloning of any rearranged mouse immunoglobulin variable genes* Immunogenetics. 1996 43:141-51), Kettleborough (*Optimization of primers for cloning libraries of mouse immunoglobulin genes using the polymerase chain reaction* Eur. J. Immunol. 1993 23:206-11),

Babcock (A novel strategy for generating monoclonal antibodies from single, isolated lymphocytes producing antibodies of defined specificities Proc. Natl. Acad. Sci. 1996 93: 7843-7848) and Williams (*Structural diversity in domains of the immunoglobulin superfamily*. Cold Spring Harb. Symp. Quant. Biol. 1989 54:637-47) as well as many others. In certain cases, the second primer may be a mixture of different primers or degenerate primers, for example.

The heavy chain CDR-specific primer may be complementary to the sequence that encodes the CDR1, CDR2 or CDR3 region of the heavy chain of the first antibody and, likewise, the light chain CDR-specific primer may be complementary to the sequence that encodes the CDR1, CDR2 or CDR3 region of the light chain of the first antibody. In certain embodiments, a particular CDR-specific primer may be chosen because the CDR sequence to which it binds may be known to be less variable than other CDR sequences.

Such CDR-anchored amplification method described in U.S. patent application Ser. No. 61/151,052, filed Feb. 9, 2009, which is incorporated by reference in its entirety for disclosure of those methods.

The above-described CDR-anchored method is effective because most sequence diversity between the variable domains in different families of antibodies that are related by lineage is in the CDR regions (i.e., the CDRs are quite variable between different families of antibodies), whereas the sequence of the CDR regions is relatively constant within the antibodies of a single family of antibodies that are related by lineage. Because the method uses primers that are complementary to sequence that are highly variable between different families of related antibodies, only related antibodies should be successfully amplified by the method.

In this embodiment, an amplification reaction may be performed using cDNA made from a second portion of the antibody-producing organ. For example, the amplification reaction may be done using nucleic acid obtained from single cells (or cultures of the same) or nucleic acid obtained from pooled cells (e.g., pools of different antibody-producing cells that each contain cDNA). Pools may contain cDNA from at least 10, at least 100 or at least 1,000 different antibody cells, for example. In embodiments in which hybridomas are used, the identity of the hybridomas that contributed to each pool may be tracked in order to identify a hybridoma producing a second antibody if the sequence encoding the second antibody is successfully amplified. Amplification products of the expected size may be sequenced directly or cloned and sequenced using known methods.

Depending on the antigen and number of antibody-producing cells in the second portion of the antibody-producing organ, the heavy and light chain variable sequences for at least 5, at least 10, at least 20, at least 50 or at least 100 or more, e.g., up to 200, up to 500, 1,000, 5,000 or 10,000 or more sequences may be obtained.

The further antibodies are tested in a second bioassay to identify a second antibody that has the same biological activity as the first antibody. As noted above, the first and second bioassays may be the same or different. In certain cases at least 30% (e.g., at least 70%, at least 80%, or at least 90%) of the lineage-related antibodies are tested in the bioassay. In this embodiment, the further antibodies may contain naturally paired heavy and light chain variable domains, or non-naturally paired heavy and light chains (i.e., heavy and light chain variable domains from different antibodies of the same lineage group). Since the antibodies are from the same lineage group, it is expected that such antibodies will be functional. In particular embodiments, the pairing of the heavy and light chains may be systematic (e.g., every heavy chain is

tested in combination with every light chain) or random (e.g., every heavy chain is tested with randomly selected light chains), for example.

An antibody produced by the instant methods finds use in diagnostics, in antibody imaging, and in treating diseases treatable by monoclonal antibody-based therapy. In particular, an antibody humanized by the instant methods may be used for passive immunization or the removal of unwanted cells or antigens, such as by complement mediated lysis or antibody mediated cytotoxicity (ADCC), all without substantial immune reactions (e.g., anaphylactic shock) associated with many prior antibodies. For example, the antibodies of the present invention may be used as a treatment for a disease where the surface of an unwanted cell specifically expresses a protein recognized the antibody (e.g. HER2, or any other cancer-specific marker) or the antibodies may be used to neutralize an undesirable toxin, irritant or pathogen. Humanized antibodies are particularly useful for the treatment of many types of cancer, for example colon cancer, lung cancer, breast cancer prostate cancer, etc., where the cancers are associated with expression of a particular cellular marker. Since most, if not all, disease-related cells and pathogens have molecular markers that are potential targets for antibodies, many diseases are potential indications for humanized antibodies. These include autoimmune diseases where a particular type of immune cells attack self-antigens, such as insulin-dependent diabetes mellitus, systemic lupus erythematosus, pernicious anemia, allergy and rheumatoid arthritis; transplantation related immune activation, such as graft rejection and graft-vs-host disease; other immune system diseases such as septic shock; infectious diseases, such as viral infection or bacteria infection; cardiovascular diseases such as thrombosis and neurological diseases such as Alzheimer's disease.

An antibody of particular interest is one that modulates, i.e., reduces or increases a symptom of the animal model disease or condition by at least about 10%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 80%, at least about 90%, or more, when compared to a control in the absence of the antibody. In general, a monoclonal antibody of interest will cause a subject animal to be more similar to an equivalent animal that is not suffering from the disease or condition. Monoclonal antibodies that have therapeutic value that have been identified using the methods and compositions of the invention are termed "therapeutic" antibodies.

## EXAMPLES

The following examples are provided in order to demonstrate and further illustrate certain embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

### Example 1

#### Method of Producing a Library of Engineered-Antibody Producing Cells

##### Isolation of Antibody Producing Cells

Rabbits are immunized with an antigen using a standard immunization protocol. At about 10 days after the second booster immunization, antibody titers are determined using ELISA. Two booster immunizations are usually sufficient for obtaining high antibody titers. As soon as a high titer (detect-

able signal at 1:100000 dilution) is observed, the rabbit is sacrificed and bone marrow cells are collected from the femur and/or other large bones. Spleen cells and peripheral blood mononuclear cells (PBMCs) are also collected and frozen in 10% DMSO/90% FBS for analysis at a later time. Very large numbers of bone marrow cells (>2 billion) are obtained from a single rabbit. After washing, clearing of debris, and red-cell lysis, the antibody producing cells, which bind to the antigen with which the rabbit was immunized, are purified using FACS. Briefly, the antigen is conjugated to a fluorescent dye and the labeled antigen is incubated with the cells obtained above. The cells are briefly rinsed to wash off any antigen non-specifically attached to the cell. After rinsing, fluorescent cells are separated from unlabeled cells using FACS. These fluorescent cells express antibodies on their surface that specifically binds to the antigen with which the animal was immunized.

#### RT-PCR to Obtain IgH and IgL Chain cDNA

##### Primer Design:

In rabbit, the 5' coding sequences of rabbit immunoglobulin heavy chain are primarily derived from only one gene. Antibody diversity is created by gene conversion and somatic mutation, but this does not affect the 5' end of the antibody cDNA. Thus, most rabbit IgG H chains have very similar or identical signal peptide sequences, and the same is true for L chains. On the 3' side, primers hybridizing to the constant domains, which also have identical sequences in most rabbit antibodies (rabbit constant domains are not divided into subclasses). As a result, only one pair of primers each is required for amplifying the vast majority of rabbit IgG H and L sequences. Typical priming sites are shown below, although any primer sites are used so long as the a variable domain-encoding polynucleotide is amplified. Typical primers for use with rabbit antibody-producing cells are as follows: heavy chain, 5' end (CACCATGGAGACTGGGCTGCGCTGGCT-TCTCCTGGTCGCTGTG (SEQ ID NO: 181)); heavy chain, 3' end (CTCCCCTCTCCGGTAAATGAGCGCT-GTGCCGCGA (SEQ ID NO: 182)); light chain kappa, 5'end (CAGGCAGGACCCAGCATGGACAC-GAGGGCCCCACT (SEQ ID NO: 183)); and L kappa, 3'end (TCAATAGGGGTGACTGTTAGAGC-GAGACGCCTGC (SEQ ID NO: 184)).

Note that the 3' H chain primer spans the 3' end of the coding region, the stop codon, and the beginning of the 3' UTR. Thus, this primer is specific for the secreted form of IgG, and does not recognize the transmembrane form, which does not contain this sequence due to alternative splicing. Therefore, the method is unlikely to recover IgG from memory B cells, which express predominantly the transmembrane form.

##### RT-PCR Conditions:

Cell lysis is done heating in a buffer containing RNase inhibitors, followed by DNA degradation and reverse transcription performed at high temperature (60° C.) using a thermostable reverse transcriptase. Reverse transcription is primed by primers specific for the 3' region of the IgG mRNAs. A single-step RT-PCR protocol is used, utilizing a thermostable enzyme that has both reverse transcriptase and DNA polymerase activities (MasterAmp™ RT-PCR Kit for High Sensitivity, Epicentre Technologies, Madison, Wis.). PCR products are analyzed by agarose gel electrophoresis. If required, a second round of PCR is performed with nested primers. In some PCR applications, this step is required to produce sufficient amounts of specific product.

##### Co-Amplification of H and L Chain cDNAs:

Different combinations of primers are tried, to accomplish efficient PCR amplification of H and L chain cDNAs in the

same reaction. A 'head start' approach is often used, where PCR cycling is started with H chain primers alone; after a number of cycles (5 to 10) the L chain primers are added to the mix. Using these methods, similar yields of H and L chain are produced. Alternatively, a nested PCR approach is used for the H chain, by performing an initial round of PCR with primers amplifying the full-length cDNA, and a second round with primers amplifying only the vH-CH1-hinge portion of the H chain. This method should yield a product similar in size to the L chain cDNA. Expression of this product yields the F(ab')<sub>2</sub> fragment of IgG, which is divalent and fully active for antigen-binding.

IgG heavy and light chain PCR products are joined with CMV promoter and BGH3'pA (bovine growth hormone polyadenylation/transcription termination) sequences.

Method a) Overlap Extension PCR.

CMV Promoter Segment:

To prepare the CMV promoter fragment, the expression vector pcDNA-3 (which contains the CMV promoter and BGH3'pA segments) is used as a template, and the following PCR setup:

Primer 1 (5' AATTCACATTGATTATTGAG 3'; SEQ ID NO: 185) corresponding to the 5' end of the CMV promoter;

Primer 2 (5' CAGCGCAGCCCAGTCTCCATCCCGT-AAGCAGTGGGTTCTC 3'; SEQ ID NO: 186) corresponding to the 3' end of the CMV promoter, and containing a 5' extension (underlined) complementary to the 5' end of the rabbit Ig H signal peptide sequence is performed.

PCR amplification with these primers produces a linear DNA fragment consisting of the CMV promoter (610 nt) and a 20 nt extension on the 3' end, which is complementary to the 5' end of the IgG vH coding region. As mentioned above, most rabbit IgGs contain 5' vH (signal peptide) regions with nearly identical sequences. Therefore, only one primer pair is needed to amplify the majority of rabbit IgG cDNAs.

BGH3' pA Segment.

A similar approach is used to prepare the BGH3'pA segment. Again, the pcDNA3 expression vector is used as a template, and the following primers are used:

Primer 3 (5' CCGGGTAAATGAGCGCTGTGGTT-TAAACCCGCTGATCAGC 3'; SEQ ID NO: 187), corresponding to 5' end of the BGH3'pA domain extended by a 20 nt sequence complementary to the 3' end of the IgG heavy chain coding region, and including 11 nt of the 3' untranslated domain.

Primer 4 (5' AAGCCATAGAGCCGACCGCA 3'; SEQ ID NO: 188) corresponding to the 3' end of the BGH polyadenylation domain.

PCR amplification results in a 250 nt fragment containing the BGH3'pA sequence and a 20 nt extension that overlaps with the 3' end of the IgG heavy chain sequence.

Overlap Extension PCR:

The IgG heavy chain PCR product are mixed with the CMV promoter and BGH3'pA segments. The mixture is subjected to 10 cycles of PCR. The overlapping segments anneal, followed by extension of the overlapping 3' ends. At the end of the 10 cycles, the outside primers (primers 1 and 4) are added to the mixture, and another 30 cycles of PCR are performed. The product is a 2100 nt fragment consisting of the CMV promoter, the IgG H coding sequence, and the BGH terminator.

IgG Light Chain:

The process are carried out in an analogous manner to produce 1500 nt fragments consisting of CMV promoter, kappa light chain coding sequence, and BGH terminator. A separate set of primers for lambda light chains can also be used to amplify and clone lambda light chains.

A low concentration of primers in the initial PCR reaction may be used. In some embodiments, primers are be designed such that amplification of the heavy chain results in a nucleotide encoding a form of the IgG H chain that is truncated at the 3' end of the hinge domain. This fragment would be similar in size to the v kappa light chain. Co-expression of these fragments results in the secretion of F(ab')<sub>2</sub> fragments of IgG.

Method b) Topoisomerase I Coupling.

This method is used as an alternative to overlap extension PCR. The overall experimental strategy is as described above. Commercially available topoisomerase-modified CMV promoter and BGH3'pA segments will be used (Invitrogen, San Diego, Calif.). The CMV promoter element (610 nt) is provided in a modified form with the topoisomerase recognition site (CCCTT) at its 3' end, and a six base pair single-stranded overhang at the 3' end (GCCTTG) which is used for directional coupling with the PCR product. The topoisomerase I enzyme is bound to the recognition site CCCTT. In order to be joined to the Topo-modified CMV promoter, the PCR product needs to contain the sequence CGGAACAAGGG (SEQ ID NO: 189) at its 5'end. This sequence is cleaved by topoisomerase, resulting in a 6-base single-strand overhang that is complementary to the single-strand overhang of the CMV promoter element. These overhangs anneal and the fragments are covalently joined by the enzyme.

In order to link the IgG cDNA fragment to the CMV promoter, the 5' primer used in the last round of IgG amplification are extended at its 5' end with the sequence CGGAACAAGGG (SEQ ID NO: 190).

The linkage of the 3' end of the IgG fragment with the BGH3'pA element is performed in an analogous manner, except that a different single-stranded overhang (GACTCA) is being used. This provides for directionality and selective joining of the 5' end with the CMV promoter and the 3' with the BGH terminator.

The joining reaction is carried out by mixing the 5' CMV element, IgG PCR product, and 3'BGH element at a 1:2:1 ratio, and adding the 10x reaction buffer. The reaction proceeds rapidly and is usually complete within 10 min at room temperature. Following the reaction, a secondary PCR reaction is carried out, using primers corresponding to the 5' end of the CMV promoter and the 3' end of the BGH terminator (primers 1 and 4, see above). This results in the formation of the 2.1 kb IgG H expression cassette, or the 1.5 kb IgG L expression cassette. Conditions for co-production of H and L IgG expression cassettes in the same reaction are also envisioned.

The IgG H expression cassettes are cloned into a vector carrying a hygromycin resistance marker to generate an IgG H expression cassette library. The IgG L expression cassettes are cloned into a vector carrying a G418 resistance marker to generate an IgG L expression cassette library.

Equimolar amounts of the IgG H and IgG L expression cassette libraries are mixed and transfected into CHO cells. The transfected CHO cells are plated into 96-well or 384-well microtiter plates such that each well contains approximately one cell. Cells are maintained in media containing both hygromycin and G418. Cells that survive the double selection contain at least one expression cassette pair.

These cells are cultured and the antibodies produced by these cells are tested for binding to the antigen with which the rabbit was immunized.

## Example 2

### Related Antibodies

Antibodies were obtained from rabbit hybridoma cells producing anti-KDR antibodies that block the interaction of

VEGF with its receptor (KDR). The hybridoma cells were generated by fusing immunized rabbit splenocytes with the rabbit hybridoma fusion partner 240E-W2.

New Zealand white rabbits were immunized with a fusion protein containing the rabbit Fc region and the extracellular domain of KDR. Each rabbit received a primary immunization by subcutaneous injection of 0.4 mg of the purified protein with complete Freund's or TiterMax adjuvant. The animals were then boosted by subcutaneous injection of 0.2 mg of the protein with incomplete Freund's or TiterMax once every three weeks. The final boost (0.4 mg protein in saline) was given intravenously 4 days before splenectomy.

Cell fusions were performed following the conventional protocol of Spieker-Polet using PEG. The ratio of splenocytes to the fusion partner was 2:1. The fused cells were plated in 96-well plates and HAT was added after 48 hrs to select for hybridomas. Direct ELISA was performed to identify antibodies that block binding of VEGF to a KDR fusion protein coated onto a microtiter plate. In this assay, the Fc-KDR ECD fusion protein was coated onto a 96-well ELISA plate and goat anti-rabbit IgG FEB conjugated to alkaline phosphatase was used to detect antibody binding to KDR. Antibodies identified in this assay were then screened for blocking VEGF interaction with KDR in a ligand-receptor assay. The blocking antibodies were identified by their inhibition of binding of VEGF in solution to KDR coated on plates.

cDNAs coding the heavy and light chains of the antibodies were cloned and sequenced. The polypeptides encoded by the cDNAs were aligned and this alignment is shown in FIG. 2. FIG. 2 shows that two groups of related anti-KDR rabbit monoclonal Abs were obtained. Antibodies 69, 6, 71, 43, 81, 4, 30, 54, 57, 50, 68, 56, 83, 36, 77, 95, 14, 42, 27 belong to one group. Antibodies 2, 17, 3, 6, 9 belong to a different group.

FIG. 3 is a multiple sequence alignment of the H3 region of ten rabbit antibody sequences extracted from the Kabat database to illustrate the expected variation in unrelated antibodies.

### Example 3

#### CDR-Anchored Amplification of Polynucleotides Encoding Related Antibodies

Several examples illustrating a method by which the amino acid sequences of related rabbit antibodies may be obtained by PCR are set forth in FIGS. 4A-4H. In the examples shown in FIGS. 4A-4D, reverse primers that are complementary to the CDR3 regions of the light chain of antibodies 31 (FIG. 4A), 29 (FIG. 4b), 27 (FIG. 4c) and 20 (FIG. 4d) were

designed and can be used along with a universal forward primer (SEQ ID NO: 118) that binds to a site that is present in all rabbit antibody heavy chain sequences to amplify coding sequences for related antibodies. In the example shown in FIG. 4A, the primers designed against sequences that encode antibody 31 are expected to amplify light chain variable domain sequences for antibodies 11, 12, 2, 25, 22, 27, 3, 1, 19, 24, 23, 18, 13, 10 and 21, which are all from the same animal as antibody 31 and are related to antibody 31 by lineage. In the example shown in FIG. 4B, the primers designed against sequences that encode antibody 29 are expected to amplify light chain variable domain sequences for antibodies 8, 9, 16 and 32, which are all from the same animal as antibody 29 and are related to antibody 29 by lineage. In the example shown in FIG. 4C, the primers designed against sequences that encode antibody 27 are expected to amplify light chain variable domain sequences for other antibodies which are all from the same animal as antibody 27 and are related to antibody 27 by lineage. In the example shown in FIG. 4D, the primers designed against sequences that encode antibody 20 are expected to amplify light chain variable domain sequences for other antibodies which are all from the same animal as antibody 20 and are related to antibody 20 by lineage.

In the examples shown in FIGS. 4E-4H, reverse primers that are complementary to the CDR3 regions of the heavy chain of antibodies 31 (FIG. 4E), 29 (FIG. 4F), 27 (FIG. 4G) and 21 (FIG. 4H) were designed and can be used along with a universal forward primer (SEQ ID NO: 148) that binds to a site that is present in all rabbit antibody heavy chain sequences to amplify coding sequences for related antibodies. In the example shown in FIG. 4E, the primers designed against sequences that encode antibody 31 are expected to amplify heavy chain variable domain sequences for antibodies 2, 17, 22, 25, 12, 1, 24, 19, 25, 11, 31, 3, 10, 13, 21, 18 and 23, which are all from the same animal as antibody 31 and are related to antibody 31 by lineage. In the example shown in FIG. 4F, the primers designed against sequences that encode antibody 29 are expected to amplify heavy chain variable domain sequences for antibodies 8, 9, 16 and 32, which are all from the same animal as antibody 29 and are related to antibody 29 by lineage. In the example shown in FIG. 4G, the primers designed against sequences that encode antibody 27 are expected to amplify heavy chain variable domain sequences for other antibodies which are all from the same animal as antibody 27 and are related to antibody 27 by lineage. In the example shown in FIG. 4H, the primers designed against sequences that encode antibody 20 are expected to amplify heavy chain variable domain sequences for other antibodies which are all from the same animal as antibody 20 and are related to antibody 20 by lineage.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 190

<210> SEQ ID NO 1

<211> LENGTH: 119

<212> TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 1

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Ser Tyr Ala  
20 25 30

-continued

---

```

Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly
   35                               40                               45

Ile Ile Arg Gly Ser Gly Ser Ile Tyr Tyr Ala Asn Trp Ala Lys Gly
   50                               55                               60

Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Arg Phe Ala
  65                               70                               75                               80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu
                               85                               90                               95

Trp Pro Gly Ser Val Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly
  100                               105                               110

Thr Leu Val Thr Val Ser Ser
  115

```

```

<210> SEQ ID NO 2
<211> LENGTH: 119
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

```

```

<400> SEQUENCE: 2

```

```

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro
  1                               5                               10                               15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Asn Ser Tyr Ala
  20                               25                               30

Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly
  35                               40                               45

Leu Ile Arg Arg Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly
  50                               55                               60

Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Phe Thr
  65                               70                               75                               80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu
                               85                               90                               95

Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly
  100                               105                               110

Thr Leu Val Thr Val Ser Ser
  115

```

```

<210> SEQ ID NO 3
<211> LENGTH: 119
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

```

```

<400> SEQUENCE: 3

```

```

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro
  1                               5                               10                               15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Asn Asn Tyr Ala
  20                               25                               30

Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly
  35                               40                               45

Leu Ile Arg Ser Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly
  50                               55                               60

Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Phe Thr
  65                               70                               75                               80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu
                               85                               90                               95

Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly
  100                               105                               110

```

-continued

Thr Leu Val Thr Val Ser Ala  
115

<210> SEQ ID NO 4  
<211> LENGTH: 119  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 4

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15  
Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Tyr Ala  
20 25 30  
Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
35 40 45  
Leu Ile Arg Ser Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly  
50 55 60  
Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Phe Thr  
65 70 75 80  
Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
85 90 95  
Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
100 105 110  
Thr Leu Val Thr Val Ser Ser  
115

<210> SEQ ID NO 5  
<211> LENGTH: 119  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 5

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15  
Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Thr Tyr Ala  
20 25 30  
Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
35 40 45  
Leu Ile Arg Ser Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly  
50 55 60  
Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Phe Thr  
65 70 75 80  
Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
85 90 95  
Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
100 105 110  
Thr Leu Val Thr Val Ser Ser  
115

<210> SEQ ID NO 6  
<211> LENGTH: 119  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 6

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15  
Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Tyr Ala

-continued

---

	20	25	30
Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly	35	40	45
Leu Ile Arg Ser Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly	50	55	60
Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Phe Thr	65	70	75
Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu	85	90	95
Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly	100	105	110
Ser Leu Val Thr Val Ser Ser	115		

<210> SEQ ID NO 7  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 7

	5	10	15
Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro			
Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Asn Asn Tyr Ala	20	25	30
Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly	35	40	45
Leu Ile Arg Ser Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly	50	55	60
Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Phe Thr	65	70	75
Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu	85	90	95
Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly	100	105	110
Thr Leu Val Thr Val Ser Ser	115		

<210> SEQ ID NO 8  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 8

	5	10	15
Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro			
Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Asn Asn Tyr Ala	20	25	30
Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly	35	40	45
Leu Ile Arg Ser Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly	50	55	60
Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Phe Thr	65	70	75
Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu	85	90	95
Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly			

-continued

---

100 105 110

Thr Leu Val Thr Val Ser Ser  
115

<210> SEQ ID NO 9  
<211> LENGTH: 119  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 9

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15

Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Asn Asn Tyr Ala  
20 25 30

Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
35 40 45

Leu Ile Arg Ser Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly  
50 55 60

Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Phe Thr  
65 70 75 80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
85 90 95

Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
100 105 110

Thr Leu Val Thr Val Ser Ser  
115

<210> SEQ ID NO 10  
<211> LENGTH: 119  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 10

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15

Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Asn Asn Tyr Ala  
20 25 30

Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
35 40 45

Leu Ile Arg Ser Ser Gly Ala Ala Tyr Asp Ala Pro Trp Ala Lys Gly  
50 55 60

Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Phe Thr  
65 70 75 80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
85 90 95

Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
100 105 110

Thr Leu Val Thr Val Ser Ser  
115

<210> SEQ ID NO 11  
<211> LENGTH: 119  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 11

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15

-continued

---

Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Asn Asn Tyr Ala  
                   20                  25                  30

Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
           35                  40                  45

Leu Ile Arg Ser Gly Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly  
       50                  55                  60

Arg Phe Thr Ile Ser Lys Val Ser Thr Thr Val Asp Leu Lys Phe Thr  
   65                  70                  75                  80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
                   85                  90                  95

Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
           100                  105                  110

Thr Leu Val Thr Val Ser Ser  
       115

<210> SEQ ID NO 12  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 12

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
   1                  5                  10                  15

Leu Thr Leu Thr Cys Ala Val Ser Gly Ile Asp Leu Ser Ser Tyr Ala  
           20                  25                  30

Met Ser Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
       35                  40                  45

Leu Ile Arg Ser Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly  
       50                  55                  60

Arg Phe Thr Ile Ser Lys Ala Ser Thr Thr Val Asp Leu Lys Met Thr  
   65                  70                  75                  80

Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
                   85                  90                  95

Trp Pro Gly Asp Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
           100                  105                  110

Thr Leu Val Thr Val Ser Ser  
       115

<210> SEQ ID NO 13  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 13

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
   1                  5                  10                  15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Ala Tyr Ala  
           20                  25                  30

Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
       35                  40                  45

Leu Ile Arg Ser Ser Gly Ala Thr Tyr Tyr Ala Ser Trp Ala Lys Gly  
       50                  55                  60

Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
   65                  70                  75                  80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
                   85                  90                  95

-continued

Trp Pro Gly Glu Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
 100 105 110

Thr Leu Val Thr Val Ser Ser  
 115

<210> SEQ ID NO 14  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 14

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Asn Asn Tyr Ala  
 20 25 30

Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45

Ile Ile Gly Ser Ser Gly Ser Ile Phe Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60

Arg Ile Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Phe Thr  
 65 70 75 80

Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
 85 90 95

Trp Ala Gly Ser Val Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
 100 105 110

Thr Leu Val Thr Val Ser Ser  
 115

<210> SEQ ID NO 15  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 15

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15

Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Tyr Ala  
 20 25 30

Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45

Ile Ile Ser Ser Ser Gly Asn Thr Tyr Phe Ala Ser Trp Ala Lys Gly  
 50 55 60

Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Phe Thr  
 65 70 75 80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
 85 90 95

Trp Ala Gly Ser Val Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
 100 105 110

Thr Leu Val Thr Val Ser Ser  
 115

<210> SEQ ID NO 16  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 16

-continued

---

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Tyr Ala  
 20 25 30  
 Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45  
 Ile Ile Arg Pro Gly Gly Asn Thr Tyr Ser Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Phe Thr  
 65 70 75 80  
 Ser Pro Thr Pro Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
 85 90 95  
 Trp Ala Gly Asp Val Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
 100 105 110  
 Thr Leu Val Thr Val Ser Ser  
 115

<210> SEQ ID NO 17  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 17

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asp Tyr Ala  
 20 25 30  
 Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45  
 Met Val Arg Asp Thr Gly Val Thr Phe Tyr Ala Gly Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Phe Thr  
 65 70 75 80  
 Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
 85 90 95  
 Trp Ala Gly Tyr Val Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
 100 105 110  
 Thr Leu Val Thr Val Ser Ser  
 115

<210> SEQ ID NO 18  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 18

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ser Leu Ser Asp Tyr Ala  
 20 25 30  
 Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45  
 Ile Ile Ser Asn Ser Gly Ile Thr Phe Tyr Ala Gly Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Phe Thr  
 65 70 75 80

-continued

---

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Ala Leu  
85 90 95

Trp Ala Gly Tyr Ile Ala Tyr Val Tyr His Asn Ile Trp Gly Pro Gly  
100 105 110

Thr Leu Val Thr Val Ser Ser  
115

<210> SEQ ID NO 19  
<211> LENGTH: 119  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 19

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15

Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ala Leu Asn Asp Phe Ala  
20 25 30

Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
35 40 45

Met Ile Ala Ser Ser Gly Asn Thr Phe Tyr Ala Ser Trp Ala Lys Gly  
50 55 60

Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Phe Thr  
65 70 75 80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Val Leu  
85 90 95

Trp Pro Gly Tyr Ile Ala Tyr Ala Tyr His Asn Ile Trp Gly Pro Gly  
100 105 110

Thr Leu Val Thr Val Ser Ser  
115

<210> SEQ ID NO 20  
<211> LENGTH: 116  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 20

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Ser Tyr Ala  
20 25 30

Met Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu His Ile Gly  
35 40 45

Ile Ile Thr Ala Ser Gly Gly Ile Tyr Tyr Ala Ser Trp Ala Lys Gly  
50 55 60

Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Arg Ile Pro  
65 70 75 80

Ser Pro Thr Thr Glu Asp Thr Gly Thr Tyr Phe Cys Ala Arg Thr Glu  
85 90 95

Asn Ser Tyr Phe Leu Tyr Phe Thr Ile Trp Gly Pro Gly Thr Leu Val  
100 105 110

Thr Val Ser Ser  
115

<210> SEQ ID NO 21  
<211> LENGTH: 119  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

-continued

&lt;400&gt; SEQUENCE: 21

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Ser Tyr Ala  
 20 25 30  
 Met Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45  
 Phe Ile Ser Ser Ser Gly Ser Thr Tyr Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Ile Thr  
 65 70 75 80  
 Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Asn Asn Tyr  
 85 90 95  
 Asp Asp Tyr Gly Asp Phe Leu His Tyr Phe Asn Ile Trp Gly Pro Gly  
 100 105 110  
 Thr Leu Val Thr Val Ser Ser  
 115

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 110

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 22

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Ser Asn Ala  
 20 25 30  
 Met Ala Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45  
 Ile Ile Asp Gly Asp Val Ser Pro Tyr Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Ser Thr Thr Val Asp Leu Lys Met  
 65 70 75 80  
 Thr Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Pro  
 85 90 95  
 Phe Asn Ile Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105 110

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 116

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 23

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Lys Asn Ala  
 20 25 30  
 Ile Ser Trp Val Arg Gln Val Pro Gly Glu Gly Leu Glu Trp Ile Gly  
 35 40 45  
 Ile Ile Tyr Gly Asp Gly Asn Arg Asp Tyr Ala Asn Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Val Thr  
 65 70 75 80  
 Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Thr

-continued

---

	85	90	95
Thr Ile Trp Ser Asp Tyr Leu Asp Ile Trp Gly Pro Gly Thr Leu Val	100	105	110
Thr Ile Ser Ser	115		

<210> SEQ ID NO 24  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus  
 <400> SEQUENCE: 24

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro	5	10	15
Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Asn Ala	20	25	30
Met Ser Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Gly	35	40	45
Thr Ile Ser Ser Ser Gly Ser Thr Tyr Tyr Ala Thr Trp Val Lys Gly	50	55	60
Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Met Thr	65	70	75
Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Asp	85	90	95
Asp Asp Val Ser Asp Tyr Phe Tyr Tyr Phe Pro Ile Trp Gly Pro Gly	100	105	110
Thr Leu Val Thr Val Ser Ser	115		

<210> SEQ ID NO 25  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus  
 <400> SEQUENCE: 25

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro	5	10	15
Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Asn Ala	20	25	30
Met Ser Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Gly	35	40	45
Thr Ile Ser Ser Ser Gly Ser Thr Tyr Tyr Ala Thr Trp Val Lys Gly	50	55	60
Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Met Thr	65	70	75
Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Asp	85	90	95
Asp Asp Val Ser Asp Tyr Phe Tyr Tyr Phe Pro Ile Trp Gly Pro Gly	100	105	110
Thr Leu Val Thr Val Ser Ser	115		

<210> SEQ ID NO 26  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus  
 <400> SEQUENCE: 26

-continued

---

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Asn Ala  
 20 25 30  
 Met Ser Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Gly  
 35 40 45  
 Thr Ile Ser Ser Ser Gly Ser Thr Tyr Tyr Ala Thr Trp Val Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
 65 70 75 80  
 Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Asp  
 85 90 95  
 Asp Asp Val Ser Asp Tyr Phe Tyr Tyr Phe Pro Ile Trp Gly Pro Gly  
 100 105 110  
 Thr Leu Val Thr Val Ser Ser  
 115

<210> SEQ ID NO 27  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 27

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Asn Ala  
 20 25 30  
 Met Ser Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Gly  
 35 40 45  
 Thr Ile Ser Ser Ser Gly Ser Thr Tyr Tyr Ala Thr Trp Val Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
 65 70 75 80  
 Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Asp  
 85 90 95  
 Asp Asp Val Ser Asp Tyr Phe Tyr Tyr Phe Pro Ile Trp Gly Pro Gly  
 100 105 110  
 Thr Leu Val Thr Val Ser Ser  
 115

<210> SEQ ID NO 28  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 28

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Asn Ala  
 20 25 30  
 Met Ser Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Gly  
 35 40 45  
 Thr Ile Ser Ser Ser Gly Ser Thr Tyr Tyr Ala Thr Trp Val Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
 65 70 75 80

-continued

---

Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Asp  
                   85                                  90                                  95

Asp Asp Val Ser Asp Tyr Phe Tyr Tyr Phe Pro Ile Trp Gly Pro Gly  
                   100                                  105                                  110

Thr Leu Val Thr Val Ser Ser  
                   115

<210> SEQ ID NO 29  
 <211> LENGTH: 117  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 29

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Gly Ser  
 1                  5                                  10                                  15

Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ser Leu Thr Ser Tyr Trp  
                   20                                  25                                  30

Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
                   35                                  40                                  45

Ala Val Ser Asn Ser Gly Thr Thr Tyr Tyr Ala Ser Trp Ala Lys Gly  
                   50                                  55                                  60

Arg Phe Thr Ile Ser Arg Thr Ser Thr Thr Val Asp Leu Arg Met Thr  
 65                                  70                                  75                                  80

Arg Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Asp Ala  
                   85                                  90                                  95

Gly Asp Asn Tyr Phe Thr Trp Leu Asp Leu Trp Gly Gln Gly Thr Leu  
                   100                                  105                                  110

Val Thr Val Ser Ser  
                   115

<210> SEQ ID NO 30  
 <211> LENGTH: 121  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 30

Glu Glu Gln Leu Lys Glu Ser Gly Gly Asp Leu Val Lys Pro Glu Gly  
 1                  5                                  10                                  15

Ser Leu Thr Leu Thr Cys Lys Ala Ser Gly Phe Asp Leu Ser Ser Glu  
                   20                                  25                                  30

Phe Tyr Ile Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp  
                   35                                  40                                  45

Ile Gly Cys Ile Ala Thr Val Ser Ser Arg Arg Leu Tyr Ala Ser Trp  
                   50                                  55                                  60

Val Asn Gly Arg Phe Thr Ile Ser Lys Thr Ser Ser Thr Thr Val Thr  
 65                                  70                                  75                                  80

Leu Gln Met Pro Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys  
                   85                                  90                                  95

Ala Arg Asp Asp Ser Ala Arg Asn Trp Phe Tyr Phe Tyr Leu Trp Gly  
                   100                                  105                                  110

Pro Gly Thr Leu Val Thr Val Ser Ser  
                   115                                  120

<210> SEQ ID NO 31  
 <211> LENGTH: 118  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

-continued

&lt;400&gt; SEQUENCE: 31

Gln Ser Leu Glu Glu Ser Gly Gly Asp Leu Val Lys Pro Gly Thr Ser  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ser Phe Ser Ala Arg His  
 20 25 30  
 Phe Met Tyr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile  
 35 40 45  
 Gly Cys Ile Asp Ile Gly Ser Gly Ser Thr Tyr Tyr Thr Ser Trp Ala  
 50 55 60  
 Lys Asp Arg Phe Thr Ile Ser Lys Pro Ser Ser Thr Thr Val Thr Leu  
 65 70 75 80  
 Gln Met Thr Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys Ala  
 85 90 95  
 Arg Ser Ser Gly Tyr Pro Tyr Tyr Phe Thr Leu Trp Gly Pro Gly Thr  
 100 105 110  
 Leu Val Thr Val Ser Ser  
 115

&lt;210&gt; SEQ ID NO 32

&lt;211&gt; LENGTH: 122

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 32

Gln Gln Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Ala  
 1 5 10 15  
 Ser Leu Thr Leu Thr Cys Lys Ala Ser Gly Phe Pro Val Thr His Trp  
 20 25 30  
 Trp Met Cys Trp Val Arg Gln Ala Pro Arg Lys Gly Leu Glu Leu Ile  
 35 40 45  
 Ala Cys Ala Tyr Thr Gly Asp Leu Thr Thr Tyr His Ala Ser Trp Ala  
 50 55 60  
 Ile Gly Arg Phe Thr Ile Ser Thr Ser Ser Ser Thr Met Val Thr Leu  
 65 70 75 80  
 Gln Met Thr Ser Leu Thr Val Ala Asp Thr Ala Thr Tyr Phe Cys Ala  
 85 90 95  
 Thr Trp Gly Ala His Asn Gln Gly Tyr Trp Asp Gly Phe Asp Pro Trp  
 100 105 110  
 Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 115 120

&lt;210&gt; SEQ ID NO 33

&lt;211&gt; LENGTH: 122

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 33

Gln Glu Gln Leu Val Glu Ser Gly Gly Asp Leu Val Gln Pro Glu Gly  
 1 5 10 15  
 Ser Leu Thr Leu Thr Cys Lys Ala Ser Gly Leu Asp Phe Ser Ser Ser  
 20 25 30  
 Tyr Trp Ile Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu  
 35 40 45  
 Ile Ala Cys Ile Tyr Thr Asp Ser Gly Gly Ile Trp Tyr Thr Ser Trp  
 50 55 60



-continued

---

```

<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 36

Gln Ser Leu Glu Glu Ser Gly Gly Asp Leu Val Lys Pro Gly Ala Ser
 1             5             10             15
Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ser Phe Ser Ser Asn Tyr
          20             25             30
Tyr Met Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Pro Glu Trp Ile
          35             40             45
Ala Cys Ile Tyr Gly Gly Ser Ile Gly Asp Pro Ser Tyr Ala Ser Trp
          50             55             60
Ala Lys Gly Arg Phe Thr Ile Ser Lys Ala Ser Ser Thr Thr Val Thr
 65             70             75             80
Leu Gln Met Thr Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys
          85             90             95
Ala Arg Glu Glu Val Gly Val Ser Ala Pro Ser Arg Gly Trp Gly Leu
          100            105            110
Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser
          115            120

```

```

<210> SEQ ID NO 37
<211> LENGTH: 121
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 37

Gln Ser Leu Glu Glu Ser Gly Gly Asp Leu Val Lys Pro Gly Ala Ser
 1             5             10             15
Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ser Phe Ser Ser Gly Tyr
          20             25             30
Tyr Met Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
          35             40             45
Ala Cys Ile Gly Val Ser Thr Gln Gly Ala Tyr Tyr Ala Ser Trp Thr
          50             55             60
Glu Gly Arg Phe Thr Ile Ser Lys Thr Ser Ser Thr Thr Val Thr Leu
 65             70             75             80
Gln Met Thr Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys Ala
          85             90             95
Arg Thr Ala Gly Ala Pro Ala Asp Ser Leu Tyr Phe Thr Leu Trp Gly
          100            105            110
Pro Gly Thr Leu Leu Thr Val Ser Ser
          115            120

```

```

<210> SEQ ID NO 38
<211> LENGTH: 119
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 38

Gln Ser Ser Glu Glu Ser Gly Gly Asp Leu Val Lys Pro Gly Ala Ser
 1             5             10             15
Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ser Leu Ser Ser Gly Tyr
          20             25             30
Asp Met Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
          35             40             45
Gly Cys Ile Lys Thr Gly Ala Thr Asn Glu Tyr Tyr Ala Ser Trp Ala

```

-continued

---

50                    55                    60  
 Lys Gly Arg Phe Thr Ile Ser Arg Thr Ser Ser Thr Thr Val Thr Leu  
 65                    70                    75                    80  
 Arg Met Thr Ser Leu Thr Ala Ala Asp Thr Ala Thr Tyr Phe Cys Ala  
                   85                    90                    95  
 Arg Glu Asp Thr Asn Asn Trp Gly Ser Leu Asn Leu Trp Gly Pro Gly  
                   100                    105                    110  
 Thr Leu Val Thr Val Ser Ser  
                   115

<210> SEQ ID NO 39  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 39

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1                    5                    10                    15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Gly Ser Phe Ala  
                   20                    25                    30  
 Val Gly Trp Val Arg Gln Pro Pro Gly Glu Gly Leu Glu Trp Ile Gly  
                   35                    40                    45  
 Leu Ile Asn Ala Asp Glu Ala Arg Tyr Tyr Ala Ser Trp Ala Lys Gly  
                   50                    55                    60  
 Arg Phe Thr Ile Ser Arg Thr Ser Thr Thr Val Asp Leu Arg Ile Thr  
 65                    70                    75                    80  
 Ser Pro Thr Ile Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Ala Pro  
                   85                    90                    95  
 Asp Asn Phe Phe Tyr Tyr Phe Ser Met Trp Gly Pro Gly Thr Leu Val  
                   100                    105                    110  
 Thr Val Ser Ser  
                   115

<210> SEQ ID NO 40  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 40

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1                    5                    10                    15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Gly Ser Phe Ala  
                   20                    25                    30  
 Val Gly Trp Val Arg Gln Pro Pro Gly Glu Gly Leu Glu Trp Ile Gly  
                   35                    40                    45  
 Leu Ile Asn Ala Asp Glu Ala Arg Tyr Tyr Ala Ser Trp Ala Lys Gly  
                   50                    55                    60  
 Arg Phe Thr Ile Ser Arg Thr Ser Thr Thr Val Asp Leu Arg Ile Thr  
 65                    70                    75                    80  
 Ser Pro Thr Ile Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Ala Pro  
                   85                    90                    95  
 Asp Asn Phe Phe Tyr Tyr Phe Ser Met Trp Gly Pro Gly Thr Leu Val  
                   100                    105                    110  
 Thr Val Ser Ser  
                   115

-continued

---

<210> SEQ ID NO 41  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*  
  
 <400> SEQUENCE: 41  
  
 Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
  
 Leu Thr Leu Thr Cys Ser Val Ser Gly Phe Ser Leu Asn Thr Tyr Ala  
 20 25 30  
  
 Val Ile Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Ile Gly  
 35 40 45  
  
 Val Val Ala Gly Gly Gly Tyr Ile Tyr Phe Thr Thr Trp Ala Lys Gly  
 50 55 60  
  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
 65 70 75 80  
  
 Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Asp  
 85 90 95  
  
 Val Asp Asp Asn Ala Asp Tyr Ser Arg Leu Asp Ile Trp Gly Pro Gly  
 100 105 110  
  
 Thr Leu Val Thr Val Ser Leu  
 115

<210> SEQ ID NO 42  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*  
  
 <400> SEQUENCE: 42  
  
 Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Asn Asn Tyr Ala  
 20 25 30  
  
 Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45  
  
 Leu Ile Tyr Val Ser Gly Ile Thr Ser Tyr Ala Ser Trp Val Asn Gly  
 50 55 60  
  
 Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
 65 70 75 80  
  
 Ser Leu Thr Val Ser Asp Thr Ala Thr Tyr Phe Cys Ala Arg Asp Leu  
 85 90 95  
  
 Tyr Gly Gly Asp His Tyr Tyr Ile Ile Trp Gly Pro Gly Thr Leu Val  
 100 105 110  
  
 Thr Val Ser Ser  
 115

<210> SEQ ID NO 43  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*  
  
 <400> SEQUENCE: 43  
  
 Gln Ser Met Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Val Asp Leu Ser Ile Tyr Ala  
 20 25 30  
  
 Met Ser Trp Val Arg Leu Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45

-continued

---

Val Ile Leu Ser Ser Gly Arg Ser Val Tyr Thr Ser Trp Ala Lys Gly  
 50 55 60

Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
 65 70 75 80

Ser Pro Thr Ile Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Tyr  
 85 90 95

Asp Asp Tyr Val Ala Leu Phe Asn Met Trp Gly Pro Gly Thr Leu Val  
 100 105 110

Thr Val Ser Ser  
 115

<210> SEQ ID NO 44  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 44

Gln Ser Met Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15

Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Asp Leu Ser Ile Tyr Ala  
 20 25 30

Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45

Val Ile Leu Ser Ser Gly Arg Ser Val Tyr Ala Gly Trp Ala Lys Gly  
 50 55 60

Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Ile Thr  
 65 70 75 80

Ser Pro Thr Ile Glu Asp Thr Ala Ala Tyr Phe Cys Ala Arg Gly Tyr  
 85 90 95

Asp Asp Tyr Val Ala Leu Phe Asn Met Trp Gly Pro Gly Thr Leu Val  
 100 105 110

Thr Val Ser Ser  
 115

<210> SEQ ID NO 45  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 45

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Gly Ser  
 1 5 10 15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Ser Tyr Ala  
 20 25 30

Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly  
 35 40 45

Ile Ile Thr Ser Arg Ala Ile Thr Tyr Tyr Ala Thr Trp Ala Lys Gly  
 50 55 60

Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Glu Leu Lys Ile Thr  
 65 70 75 80

Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Tyr  
 85 90 95

Asp Asp Tyr Val Ala Leu Phe Asn Met Trp Gly Pro Gly Thr Leu Val  
 100 105 110

Thr Val Ser Ser  
 115

-continued

<210> SEQ ID NO 46  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 46

```

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro
 1           5           10           15
Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Ser Tyr Ala
 20           25           30
Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly
 35           40           45
Ile Ile Ser Ser Thr Gly Asn Thr Cys Tyr Ala Asn Trp Ala Lys Gly
 50           55           60
Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Lys Ile Thr
 65           70           75           80
Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Tyr
 85           90           95
Asp Asp Tyr Val Ala Leu Phe Asn Met Trp Gly Pro Gly Thr Leu Val
 100          105          110
Thr Val Ser Ser
 115
  
```

<210> SEQ ID NO 47  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 47

```

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro
 1           5           10           15
Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Ser Tyr Ala
 20           25           30
Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Gly
 35           40           45
Ile Ile Phe Ser Ser Gly Asn Ile Val Tyr Ala Arg Arg Ala Lys Gly
 50           55           60
Arg Phe Thr Ile Ser Lys Thr Ser Thr Thr Val Asp Leu Gln Ile Thr
 65           70           75           80
Ser Pro Thr Ile Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Tyr
 85           90           95
Asp Asp Tyr Val Ala Leu Phe Asn Met Trp Gly Pro Gly Thr Leu Val
 100          105          110
Thr Val Ser Ser
 115
  
```

<210> SEQ ID NO 48  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 48

```

Asp Pro Val Leu Thr Gln Thr Pro Ala Ser Val Ser Glu Pro Val Gly
 1           5           10           15
Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Glu Asp Leu Gly Ser Asn
 20           25           30
  
```

-continued

---

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
                   35                                  40                                  45

Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
           50                                  55                                  60

Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Gly Val Gln Cys  
   65                                  70                                  75                                  80

Asp Asp Ala Ala Thr Tyr Tyr Cys Gln Asp Thr Tyr Tyr Gly Asn Thr  
                   85                                  90                                  95

Tyr Leu Gly Ala Phe Gly Gly Gly Thr Glu Val Val Val Arg  
                   100                                  105                                  110

<210> SEQ ID NO 49  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus  
 <400> SEQUENCE: 49

Asp Leu Val Met Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly  
   1                  5                                  10                                  15

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Gly Asn  
                   20                                  25                                  30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
                   35                                  40                                  45

Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
           50                                  55                                  60

Ser Gly Phe Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
   65                                  70                                  75                                  80

Asp Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Thr  
                   85                                  90                                  95

Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys  
                   100                                  105                                  110

<210> SEQ ID NO 50  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus  
 <400> SEQUENCE: 50

Asp Leu Val Met Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly  
   1                  5                                  10                                  15

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Gly Asn  
                   20                                  25                                  30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Gln Leu Leu Ile  
                   35                                  40                                  45

Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
           50                                  55                                  60

Ser Gly Phe Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
   65                                  70                                  75                                  80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Thr  
                   85                                  90                                  95

Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys  
                   100                                  105                                  110

<210> SEQ ID NO 51  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

-continued

&lt;400&gt; SEQUENCE: 51

Asp Leu Val Met Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly  
 1 5 10 15  
 Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Gly Asn  
 20 25 30  
 Leu Ala Trp Tyr Leu Gln Lys Pro Gly Gln Pro Pro Gln Leu Leu Ile  
 35 40 45  
 Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60  
 Ser Gly Phe Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80  
 Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Ser  
 85 90 95  
 Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

&lt;210&gt; SEQ ID NO 52

&lt;211&gt; LENGTH: 110

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 52

Asp Leu Val Met Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly  
 1 5 10 15  
 Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Gly Asn  
 20 25 30  
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Gln Leu Leu Ile  
 35 40 45  
 Tyr Ser Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60  
 Ser Gly Phe Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80  
 Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Ser  
 85 90 95  
 Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

&lt;210&gt; SEQ ID NO 53

&lt;211&gt; LENGTH: 110

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 53

Asp Leu Val Met Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly  
 1 5 10 15  
 Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Gly Asn  
 20 25 30  
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45  
 Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60  
 Ser Gly Phe Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80  
 Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Ser  
 85 90 95  
 Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys

-continued

---

100 105 110

<210> SEQ ID NO 54  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 54

Asp Leu Val Met Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly  
 1 5 10 15

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Gly Asn  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45

Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60

Ser Gly Phe Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Ser  
 85 90 95

Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 55  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 55

Asp Leu Val Met Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly  
 1 5 10 15

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Asp Asn  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45

Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60

Ser Gly Phe Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Ser  
 85 90 95

Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 56  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 56

Asp Leu Val Met Thr Gln Thr Pro Ser Pro Val Ser Ala Pro Val Gly  
 1 5 10 15

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Gly Asn  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45

Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60



-continued

---

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Gly Asn  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45

Tyr Ser Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60

Ser Gly Ser Gly Thr Glu Tyr Ser Leu Ser Ile Ser Asp Leu Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Ser  
 85 90 95

Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 60  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 60

Asp Val Val Met Thr Gln Thr Pro Ala Ser Val Ser Glu Pro Val Gly  
 1 5 10 15

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Ser Asn  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45

Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60

Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Asn  
 85 90 95

Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 61  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 61

Asp Val Val Met Thr Gln Thr Pro Ala Ser Val Ser Glu Pro Val Gly  
 1 5 10 15

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Ala Ser Asn  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45

Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60

Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Phe Asn  
 85 90 95

Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 62  
 <211> LENGTH: 110

-continued

---

```

<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 62

Asp Val Val Met Thr Gln Thr Pro Ala Ser Val Ser Glu Pro Val Gly
 1          5          10          15
Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Ile Gly Gly Asn
          20          25          30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
          35          40          45
Tyr Ser Ala Ser Ser Leu Thr Ser Gly Val Pro Ser Arg Phe Lys Gly
          50          55          60
Ser Glu Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys
65          70          75          80
Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Ser Tyr Tyr Gly Phe Asn
          85          90          95
Tyr Val Gly Pro Phe Gly Gly Gly Thr Glu Val Val Val Lys
          100          105          110

```

```

<210> SEQ ID NO 63
<211> LENGTH: 110
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

```

```

<400> SEQUENCE: 63

Gly Pro Val Met Thr Gln Thr Pro Ala Ser Val Ser Glu Pro Val Gly
 1          5          10          15
Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Glu Asp Ile Gly Ser Asn
          20          25          30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Asn Leu Leu Val
          35          40          45
Tyr Ser Ala Ser Thr Leu Thr Ser Gly Val Pro Ser Arg Phe Ser Gly
          50          55          60
Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys
65          70          75          80
Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Asp Thr Tyr Tyr Gly Asn Thr
          85          90          95
Tyr Leu Gly Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys
          100          105          110

```

```

<210> SEQ ID NO 64
<211> LENGTH: 110
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

```

```

<400> SEQUENCE: 64

Gly Pro Val Met Thr Gln Thr Pro Ala Ser Val Ser Glu Pro Val Gly
 1          5          10          15
Gly Thr Val Thr Met Lys Cys Gln Ala Ser Glu Asp Ile Gly Ser Asn
          20          25          30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Asn Leu Leu Val
          35          40          45
Phe Ser Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Ala Gly
          50          55          60
Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys
65          70          75          80
Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Asp Ser Tyr Tyr Gly Asn Ser

```

-continued

---

	85	90	95
Tyr Leu Gly Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys	100	105	110

<210> SEQ ID NO 65  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 65

Asp Pro Val Met Thr Gln Thr Pro Ala Ser Val Ser Glu Pro Val Gly			
1	5	10	15
Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Glu Leu Gly Ser Asn	20	25	30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile	35	40	45
Tyr Ser Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys Gly	50	55	60
Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys	65	70	75
Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Phe Gly Ser Ser	85	90	95

	100	105	110
--	-----	-----	-----

<210> SEQ ID NO 66  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 66

Asp Pro Val Leu Thr Gln Thr Pro Ala Ser Val Ser Ala Val Val Gly			
1	5	10	15
Gly Thr Val Thr Ile Asn Cys Gln Thr Ser Glu Asp Ile Ala Ser Asn	20	25	30
Leu Ala Trp Tyr Gln Gln Lys Ser Gly Gln Arg Pro Lys Leu Leu Ile	35	40	45
Tyr Glu Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Lys Gly	50	55	60
Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys	65	70	75
Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Tyr Thr Tyr Tyr Gly Ser Ser	85	90	95

	100	105	110
--	-----	-----	-----

<210> SEQ ID NO 67  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 67

Ala Val Val Leu Thr Gln Thr Ala Ser Pro Val Ser Gly Ala Val Gly			
1	5	10	15
Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Gln Asn Ile Tyr Asn Asn	20	25	30
Leu Ala Trp Tyr Arg Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile	35	40	45

-continued

---

Tyr Ala Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60

Ser Arg Ser Gly Thr Gln Phe Thr Leu Ser Ile Ser Asp Leu Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Tyr Phe Asp Ser Ser  
 85 90 95

Ser Thr Asp Ala Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 68  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 68

Ala Val Val Leu Thr Gln Thr Ala Ser Pro Val Ser Gly Ala Val Gly  
 1 5 10 15

Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Gln Asn Ile Tyr Asn Asn  
 20 25 30

Leu Ala Trp Tyr Arg Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45

Tyr Ala Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60

Ser Arg Ser Gly Thr Gln Phe Thr Leu Ser Ile Ser Asp Leu Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Tyr Phe Asp Ser Ser  
 85 90 95

Ser Thr Asp Ala Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 69  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 69

Ala Val Val Leu Thr Gln Thr Ala Ser Pro Val Ser Gly Ala Val Gly  
 1 5 10 15

Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Gln Asn Ile Tyr Asn Asn  
 20 25 30

Leu Ala Trp Tyr Arg Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45

Tyr Ala Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60

Ser Arg Ser Gly Thr Gln Phe Thr Leu Ser Ile Ser Asp Leu Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Tyr Phe Asp Ser Ser  
 85 90 95

Ser Thr Asp Ala Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 70  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 70

-continued

---

Ala Val Val Leu Thr Gln Thr Ala Ser Pro Val Ser Gly Ala Val Gly  
 1 5 10 15  
 Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Gln Asn Ile Tyr Asn Asn  
 20 25 30  
 Leu Ala Trp Tyr Arg Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45  
 Tyr Ala Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60  
 Ser Arg Ser Gly Thr Gln Phe Thr Leu Ser Ile Ser Asp Leu Glu Cys  
 65 70 75 80  
 Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Tyr Phe Asp Ser Ser  
 85 90 95  
 Ser Thr Asp Ala Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 71  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 71

Ala Val Val Leu Thr Gln Thr Ala Ser Pro Val Ser Gly Ala Val Gly  
 1 5 10 15  
 Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Gln Asn Ile Tyr Asn Asn  
 20 25 30  
 Leu Ala Trp Tyr Arg Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45  
 Tyr Ala Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60  
 Ser Arg Ser Gly Thr Gln Phe Thr Leu Ser Ile Ser Asp Leu Glu Cys  
 65 70 75 80  
 Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Tyr Phe Asp Ser Ser  
 85 90 95  
 Ser Thr Asp Ala Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 72  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 72

Ala Val Val Leu Thr Gln Thr Ala Ser Pro Val Ser Gly Ala Val Gly  
 1 5 10 15  
 Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Gln His Ile Tyr Ser Asn  
 20 25 30  
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Arg Pro Lys Leu Leu Ile  
 35 40 45  
 Tyr Ala Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60  
 Ser Arg Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80  
 Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Tyr Tyr Ser Ser Asp  
 85 90 95  
 Ser Thr Asp Asn Thr Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

-continued

---

```

<210> SEQ ID NO 73
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 73
Ala Val Val Leu Thr Gln Thr Pro Ser Ser Val Ser Ala Ala Val Gly
 1             5             10             15
Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Asn Ile Tyr Ser Asn
          20             25             30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
          35             40             45
Tyr Gly Ala Ser Ala Leu Pro Ser Gly Val Pro Ser Arg Phe Ser Gly
 50             55             60
Ser Arg Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Asp Leu Glu Cys
 65             70             75             80
Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Phe Tyr Ser Ser Ser
          85             90             95
Asn Asp Asp Asn Pro Phe Gly Gly Gly Thr Glu Val Ala Val Lys
 100            105            110

```

```

<210> SEQ ID NO 74
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 74
Ala Val Val Leu Thr Gln Thr Thr Ser Ser Val Ser Ala Asp Val Gly
 1             5             10             15
Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Glu Asn Ile Tyr Ser Leu
          20             25             30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
          35             40             45
Tyr Ser Ala Ser Asp Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly
 50             55             60
Ser Gly Phe Gly Thr Glu Phe Thr Leu Thr Ile Ser Asp Leu Glu Cys
 65             70             75             80
Ala Asp Ala Ala Thr Tyr His Cys Gln Ser Tyr Tyr Tyr Ser Gly Ser
          85             90             95
Ser Ala Asp Thr Gly Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys
 100            105            110

```

```

<210> SEQ ID NO 75
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 75
Ala Val Val Leu Thr Gln Thr Ala Ser Pro Val Ser Gly Ala Val Gly
 1             5             10             15
Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Gln Asn Ile Tyr Ser Asn
          20             25             30
Leu Ala Trp Tyr Gln His Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
          35             40             45
Tyr Lys Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys Gly
 50             55             60
Ser Gly Ser Gly Thr Glu Tyr Ile Leu Thr Ile Ser Asp Leu Glu Cys

```

-continued

---

```

65              70              75              80
Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Thr Asn Tyr Tyr Ser Ile Asn
      85              90              95
Gly Gly Glu Val Thr Phe Gly Gly Gly Thr Glu Val Val Val Lys
      100              105              110

```

```

<210> SEQ ID NO 76
<211> LENGTH: 110
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 76

```

```

Asp Val Val Met Thr Gln Thr Pro Ala Ser Val Ser Glu Pro Val Gly
 1              5              10              15
Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Gln Ser Ile Arg Leu Ser
      20              25              30
Asp Leu Ala Trp Tyr Gln Gln Lys Pro Gly His Pro Pro Lys Leu Leu
      35              40              45
Ile Tyr Lys Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Lys
      50              55              60
Gly Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Gly Val Gln
65              70              75              80
Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Ile Asp Tyr Asp Asn
      85              90              95
Tyr Val Phe Phe Phe Gly Gly Gly Thr Glu Val Val Val Lys
      100              105              110

```

```

<210> SEQ ID NO 77
<211> LENGTH: 110
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 77

```

```

Ala Leu Val Met Thr Gln Thr Pro Ala Ser Val Ser Ala Ala Val Gly
 1              5              10              15
Gly Thr Val Thr Ile Ser Cys Gln Ala Ser Gln Ser Ile His Ser Trp
      20              25              30
Val Ser Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
      35              40              45
Tyr Tyr Ala Ala Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys Gly
      50              55              60
Ser Gly Ser Gly Lys Gln Phe Thr Leu Thr Ile Ser Asp Leu Glu Cys
65              70              75              80
Asp Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Asp Phe Gly Gly Ser Asp
      85              90              95
Val Asp Asn Thr Phe Gly Gly Gly Thr Glu Val Val Val Ala
      100              105              110

```

```

<210> SEQ ID NO 78
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 78

```

```

Gln Val Leu Thr Gln Ser Pro Ser Pro Val Ser Ala Ala Val Gly Gly
 1              5              10              15
Thr Val Thr Ile Asn Cys Gln Ala Ser Gln Ser Val Tyr Asn Gly Asp
      20              25              30

```

-continued

Trp Leu Gly Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Ser Ser Leu Ile Asn  
 50 55 60  
 Cys Asn Gly Ser Gly Thr Gln Trp Thr Leu Thr Ile Ser Gly Val Gln  
 65 70 75 80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Gln Gly Glu Phe Ser Cys Ser  
 85 90 95  
 Ser Ala Asp Cys Val Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 79  
 <211> LENGTH: 113  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 79

Ala Asp Ile Val Leu Thr Gln Thr Pro Ala Ser Val Ser Ala Thr Val  
 1 5 10 15  
 Gly Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Gln Asn Ile Tyr Ser  
 20 25 30  
 Asn Leu Ala Trp Tyr Gln Gln Arg Pro Gly Gln Pro Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Ala Ala Ser Asn Leu Ala Ser Gly Val Ser Ser Arg Phe Ser  
 50 55 60  
 Gly Ser Arg Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Asp Leu Glu  
 65 70 75 80  
 Cys Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Tyr Tyr Ser Ser  
 85 90 95  
 Asp Asp Asn Tyr Leu Asn Thr Phe Gly Gly Gly Thr Glu Val Val Val  
 100 105 110

Lys

<210> SEQ ID NO 80  
 <211> LENGTH: 113  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 80

Ala Asp Ile Val Leu Thr Gln Thr Pro Ala Ser Val Ser Ala Thr Val  
 1 5 10 15  
 Gly Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Gln Asn Ile Tyr Ser  
 20 25 30  
 Asn Leu Ala Trp Tyr Gln Gln Arg Pro Gly Gln Pro Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Ala Ala Ser Asn Leu Ala Ser Gly Val Ser Ser Arg Phe Ser  
 50 55 60  
 Gly Ser Arg Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Asp Leu Glu  
 65 70 75 80  
 Cys Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Tyr Tyr Ser Ser  
 85 90 95  
 Asp Asp Asn Tyr Leu Asn Thr Phe Gly Gly Gly Thr Glu Val Val Val  
 100 105 110

Lys

-continued

---

<210> SEQ ID NO 81  
 <211> LENGTH: 107  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 81

Gln Val Leu Thr Gln Thr Pro Ala Ser Val Ser Ala Thr Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Lys Cys Gln Ala Ser Gln Asn Ile Tyr Ser Asn Leu  
 20 25 30  
 Ala Trp Tyr Gln Gln Thr Pro Gly Gln Pro Pro Lys Leu Leu Ile Tyr  
 35 40 45  
 Gly Ala Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Lys Gly Ser  
 50 55 60  
 Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Asp Leu Glu Cys Ala  
 65 70 75 80  
 Asp Ala Gly Thr Tyr Tyr Cys Gln Asp Tyr Tyr Gly Thr Ser Arg Tyr  
 85 90 95  
 Ile Phe Gly Gly Gly Thr Glu Val Val Val Gly  
 100 105

<210> SEQ ID NO 82  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 82

Ala Asp Ile Val Met Thr Gln Thr Pro Ala Ser Val Ser Glu Pro Val  
 1 5 10 15  
 Gly Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Gln Ser Ile Tyr Asn  
 20 25 30  
 Tyr Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Ala Ala Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
 50 55 60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Asp Leu Glu  
 65 70 75 80  
 Cys Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Asn Tyr Gly Thr Ser  
 85 90 95  
 Ser Ala Ser Tyr Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 83  
 <211> LENGTH: 107  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 83

Asp Val Val Met Thr Gln Thr Pro Phe Ser Val Ser Ala Ala Val Gly  
 1 5 10 15  
 Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Ser Ile Tyr Ile Asn  
 20 25 30  
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45  
 Tyr Asp Ala Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60  
 Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80

-continued

Ala Asp Ala Thr Thr Tyr Tyr Cys Gln Asn Tyr Tyr Gly Ser Ser Tyr  
85 90 95

Asp Phe Gly Gly Gly Thr Glu Val Val Val Lys  
100 105

<210> SEQ ID NO 84  
<211> LENGTH: 111  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 84

Asp Val Val Met Thr Gln Thr Pro Ala Ser Val Glu Ala Ala Val Gly  
1 5 10 15

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Asn Ile Tyr Ser Gly  
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
35 40 45

Tyr Asp Ala Ser Asn Leu Glu Thr Gly Val Pro Ser Arg Phe Lys Gly  
50 55 60

Ser Gly Ser Gly Ala Gln Phe Thr Leu Thr Ile Ser Gly Val Gln Cys  
65 70 75 80

Asp Asp Ala Ala Thr Tyr Tyr Cys Gln Cys Ser Glu Tyr Asp Ser Ser  
85 90 95

Tyr Val Pro Asn Val Phe Gly Gly Gly Thr Glu Val Val Val Lys  
100 105 110

<210> SEQ ID NO 85  
<211> LENGTH: 107  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 85

Gln Val Leu Thr Gln Thr Pro Ala Ser Val Ser Ala Ala Val Gly Gly  
1 5 10 15

Thr Val Thr Ile Lys Cys Gln Ala Ser Glu Ser Val Ser Ser Arg Leu  
20 25 30

Ala Trp Tyr Gln Gln Lys Ala Gly Gln Pro Pro Lys Leu Leu Ile Tyr  
35 40 45

Ser Ala Ser Thr Leu Ala Ser Gly Val Ser Ser Arg Phe Lys Gly Ser  
50 55 60

Gly Ser Gly Thr Glu Phe Ser Leu Thr Ile Ser Asp Leu Glu Cys Ala  
65 70 75 80

Asp Ala Ala Thr Tyr Tyr Cys Gln Ser Tyr Tyr Arg Ile Asp Arg Asn  
85 90 95

Leu Phe Gly Glu Gly Thr Glu Val Val Val Lys  
100 105

<210> SEQ ID NO 86  
<211> LENGTH: 110  
<212> TYPE: PRT  
<213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 86

Ala Val Val Leu Thr Gln Thr Pro Ala Ser Val Ser Ala Ala Val Gly  
1 5 10 15

Gly Thr Val Thr Ile Asn Cys Gln Ala Ser Gln Ser Ile Asn Ser Gly  
20 25 30

-continued

---

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile  
 35 40 45

Tyr Gln Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys Gly  
 50 55 60

Ser Gly Ser Gly Thr Glu Tyr Thr Leu Thr Ile Ser Asp Leu Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Thr Tyr Asp Thr Ser Asp  
 85 90 95

Leu Asp Asn Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 87  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*  
 <400> SEQUENCE: 87

Ala Tyr Asp Met Thr Gln Thr Pro Ala Ser Val Glu Val Ala Val Gly  
 1 5 10 15

Gly Thr Val Thr Ile Lys Cys Gln Ala Ser Glu Asp Ile Glu Ser Tyr  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Arg Pro Leu Leu Leu Ile  
 35 40 45

Tyr Ala Ala Ser Thr Leu Ala Ser Gly Ala Ser Ser Arg Phe Lys Gly  
 50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Gly Val Glu Cys  
 65 70 75 80

Ala Asp Ala Ala Thr Tyr Tyr Cys His Gln Ala Tyr Thr Thr Thr Gly  
 85 90 95

Leu Asp Ala Gly Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 88  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*  
 <400> SEQUENCE: 88

Ile Val Met Thr Gln Thr Pro Ser Ser Lys Ser Val Pro Val Gly Asp  
 1 5 10 15

Thr Val Thr Ile Asn Cys Gln Ala Ser Glu Ser Val Tyr Asn Gly Asp  
 20 25 30

Arg Leu Ala Trp Phe Gln Gln Lys Pro Gly Gln Pro Pro Lys Arg Leu  
 35 40 45

Ile Tyr Leu Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
 50 55 60

Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Asp Val Val  
 65 70 75 80

Cys Ala Asp Ala Ala Thr Tyr Tyr Cys Ala Gly Tyr Arg Ser Ala Ser  
 85 90 95

Thr Asp Gly Asn Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 89  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

-continued

&lt;400&gt; SEQUENCE: 89

Ala Val Leu Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Ser Cys Gln Ser Ser Lys Ser Val Tyr Ala Asp Asn  
 20 25 30  
 Cys Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Arg Pro Arg Leu Leu  
 35 40 45  
 Ile Tyr Glu Ala Ser Lys Leu Pro Ser Gly Val Pro Ser Arg Phe Lys  
 50 55 60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Asp Val Gln  
 65 70 75 80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Ala Gly Asp Tyr Ser Pro Thr  
 85 90 95  
 Ser Asp Asn Ser Phe Ser Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

&lt;210&gt; SEQ ID NO 90

&lt;211&gt; LENGTH: 110

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 90

Ala Val Leu Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Lys Cys Gln Ser Ser Gln Ser Val Tyr Asn His Asn  
 20 25 30  
 Leu Leu Ser Trp Tyr Gln His Lys Pro Gly Gln Pro Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
 50 55 60  
 Gly Asn Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Gln  
 65 70 75 80  
 Cys Ala Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Gly Tyr Asp Asp Asp  
 85 90 95  
 Ala Asp Thr Ala Phe Gly Gly Gly Thr Glu Val Val Val Arg  
 100 105 110

&lt;210&gt; SEQ ID NO 91

&lt;211&gt; LENGTH: 110

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 91

Ala Val Leu Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Lys Cys Gln Ser Ser Gln Ser Val Tyr Asn Asn Asn  
 20 25 30  
 Leu Leu Ser Trp Tyr Gln His Lys Pro Gly Gln Pro Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
 50 55 60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Gln  
 65 70 75 80  
 Cys Ala Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Gly Tyr Asp Asp Asp  
 85 90 95  
 Ala Asp Thr Ala Phe Gly Gly Gly Thr Glu Val Val Val Arg

-continued

---

100                      105                      110  
 <210> SEQ ID NO 92  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*  
 <400> SEQUENCE: 92  
 Ala Val Leu Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly Gly  
 1                      5                      10                      15  
 Thr Val Thr Ile Lys Cys Gln Ser Ser Gln Ser Val Tyr Asn Asn Ala  
                     20                      25                      30  
 Leu Leu Ser Trp Tyr Gln Gln Lys Ser Gly Gln Pro Pro Lys Leu Leu  
                     35                      40                      45  
 Ile Tyr Asp Ala Ser Lys Leu Ala Ser Gly Val Pro Asp Arg Phe Ser  
                     50                      55                      60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Gln  
 65                      70                      75                      80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Gly Tyr Asp Asp Asp  
                     85                      90                      95  
 Ala Asp Thr Thr Phe Gly Gly Gly Thr Glu Val Val Val Glu  
                     100                      105                      110

<210> SEQ ID NO 93  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*  
 <400> SEQUENCE: 93  
 Ala Val Leu Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly Gly  
 1                      5                      10                      15  
 Thr Val Thr Ile Lys Cys Gln Ser Ser Gln Ser Val Tyr Asn Asn Asn  
                     20                      25                      30  
 Ala Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu  
                     35                      40                      45  
 Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Asp Arg Phe Ser  
                     50                      55                      60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Gln  
 65                      70                      75                      80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Gly Tyr Asp Asp Asp  
                     85                      90                      95  
 Ala Asp Thr Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
                     100                      105                      110

<210> SEQ ID NO 94  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*  
 <400> SEQUENCE: 94  
 Ala Val Val Thr Gln Thr Pro Ser Pro Val Ser Ala Thr Val Gly Gly  
 1                      5                      10                      15  
 Thr Val Thr Ile Ser Cys Gln Ser Ser Glu Ser Val Tyr Asn Asp Val  
                     20                      25                      30  
 Cys Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Arg Pro Lys Leu Leu  
                     35                      40                      45  
 Ile Tyr Asp Ala Phe Thr Leu Ala Ser Gly Val Ser Ser Arg Phe Lys  
                     50                      55                      60



-continued

---

<211> LENGTH: 17  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 100

Cys Ala Arg Gly Gly Ala Gly Ile Ser Gly Tyr Thr Tyr Phe Asn Ile  
 1 5 10 15

Trp

<210> SEQ ID NO 101  
 <211> LENGTH: 14  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 101

Cys Ala Arg Gly Cys Pro Gly Tyr Gly Asp Asn Asp Ile Trp  
 1 5 10

<210> SEQ ID NO 102  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 102

Cys Ala Arg Gly Tyr Trp Ser Leu Asp Ile Trp  
 1 5 10

<210> SEQ ID NO 103  
 <211> LENGTH: 15  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 103

Cys Val Arg Asp Ser Thr Gly Ile Ser Ala Leu Phe Asn Val Trp  
 1 5 10 15

<210> SEQ ID NO 104  
 <211> LENGTH: 16  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 104

Cys Ala Arg Arg Gly Ala Thr Ala Ser His Arg Trp Phe Thr Ile Trp  
 1 5 10 15

<210> SEQ ID NO 105  
 <211> LENGTH: 16  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 105

Cys Gly Ser Gly Ala Asn Ile Glu Asn Glu Phe Phe Asn Ala Ile Trp  
 1 5 10 15

-continued

---

<210> SEQ ID NO 106  
 <211> LENGTH: 16  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 106

Cys Ala Arg Gly Asp Arg Ser His Asp Tyr Asp Tyr Phe Lys Ile Trp  
 1 5 10 15

<210> SEQ ID NO 107  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 107

Cys Ala Arg Ser Gln Asp Ser Gly Ser His Asp Asp Phe Pro Phe Asn  
 1 5 10 15

Ile Trp

<210> SEQ ID NO 108  
 <211> LENGTH: 15  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 108

Cys Ala Arg Ser Pro Gly Gly Ile Gly Asp Ala Phe Asp Pro Trp  
 1 5 10 15

<210> SEQ ID NO 109  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 109

Cys Ala Arg Gly Trp Val Gly Leu Asn Ile Trp  
 1 5 10

<210> SEQ ID NO 110  
 <211> LENGTH: 15  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 110

Cys Ala Arg Arg Ala Asp Ser Tyr Gly Tyr Ala Tyr Asp Ile Trp  
 1 5 10 15

<210> SEQ ID NO 111  
 <211> LENGTH: 14  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 111

Cys Ala Arg Tyr Gly Ala Ser Val Thr Tyr Phe Asn Ile Trp  
 1 5 10

-continued

<210> SEQ ID NO 112  
 <211> LENGTH: 17  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 112

Cys Ala Arg Phe Arg Ile Leu Val Ile Val Leu Val Pro Leu Asp Leu  
 1 5 10 15

Trp

<210> SEQ ID NO 113  
 <211> LENGTH: 17  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 113

Cys Ala Arg Gly Ala Thr Met Thr Met Val Arg Gly Trp Leu Asp Leu  
 1 5 10 15

Trp

<210> SEQ ID NO 114  
 <211> LENGTH: 13  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: VH3 region

<400> SEQUENCE: 114

Cys Ala Arg Leu Gly Leu Val Val Val Ile Asn Ile Trp  
 1 5 10

<210> SEQ ID NO 115  
 <211> LENGTH: 405  
 <212> TYPE: DNA  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 115

atggacacga gggccccac tcagctgctg gggctcctgc tgctctggct cccaggtgcc 60  
 acatttgtc aactgctgac ccagactgca tcgccctgt ctacagctgt gggaggcaca 120  
 gtcaccatca agtgccagtc cagtcagagt gtttttaaga ggaagtctt atcctggtat 180  
 cagcagaaac cagggcaggc tcccaaactc ctgatctacg atgcatccac tctggcatct 240  
 ggggtcccat cacggttcag tggcagtgga tctgggacac agttcactct caccatcagc 300  
 ggcggtcagt gtgacgatgc tgccaactac tactgtctag gcagttttga ttgtactagt 360  
 gctgattgtc atgttttcgg cggagggacc gaggtggtgg tcaaa 405

<210> SEQ ID NO 116  
 <211> LENGTH: 135  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 116

Met Asp Thr Arg Ala Pro Thr Gln Leu Leu Gly Leu Leu Leu Leu Trp  
 1 5 10 15

Leu Pro Gly Ala Thr Phe Ala Gln Leu Leu Thr Gln Thr Ala Ser Pro  
 20 25 30



-continued

&lt;400&gt; SEQUENCE: 120

Gln Val Leu Thr Gln Thr Ala Ser Pro Val Ser Ala Ala Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Lys Cys Gln Ser Ser Gln Ser Val Tyr Lys Arg Lys  
 20 25 30  
 Ser Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Ser  
 50 55 60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Gln  
 65 70 75 80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Phe Asp Cys Thr  
 85 90 95  
 Ser Ala Asp Cys His Val Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

&lt;210&gt; SEQ ID NO 121

&lt;211&gt; LENGTH: 112

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 121

Gln Val Leu Thr Gln Thr Ala Ser Pro Val Ser Ala Ala Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Asn Cys Gln Ser Ser Gln Ser Val Tyr Lys Arg Lys  
 20 25 30  
 Ser Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
 50 55 60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Gln  
 65 70 75 80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Phe Asp Cys Thr  
 85 90 95  
 Arg Ala Asp Cys His Val Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

&lt;210&gt; SEQ ID NO 122

&lt;211&gt; LENGTH: 112

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 122

Gln Val Leu Thr Gln Thr Ala Ser Pro Val Ser Ala Ala Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Lys Cys Gln Ser Ser Gln Ser Val Tyr Lys Arg Lys  
 20 25 30  
 Ser Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
 50 55 60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Gln  
 65 70 75 80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Phe Asp Cys Thr  
 85 90 95

-continued

---

Arg Ala Asp Cys His Val Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 123  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 123

Gln Val Leu Thr Gln Thr Ala Ser Pro Val Ser Ala Ala Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Lys Cys Gln Ser Ser Gln Ser Val Tyr Lys Arg Lys  
 20 25 30  
 Ser Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
 50 55 60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Gln  
 65 70 75 80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Phe Asp Cys Thr  
 85 90 95  
 Ser Ala Asp Cys His Val Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 124  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 124

Gln Val Leu Thr Gln Thr Ala Ser Pro Val Ser Ala Ala Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Lys Cys Gln Ser Ser Gln Ser Val Tyr Lys Ser Lys  
 20 25 30  
 His Cys Ser Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
 50 55 60  
 Gly Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Gly Val Gln  
 65 70 75 80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Tyr Asp Cys Ser  
 85 90 95  
 Thr Ala Asp Cys His Val Phe Gly Gly Gly Thr Gly Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 125  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 125

Gln Val Leu Thr Gln Thr Pro Ser Ser Val Ser Ala Ala Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Lys Cys Gln Ser Ser Gln Ser Val Tyr Lys Ser Lys  
 20 25 30  
 His Leu Ser Trp Phe Gln Gln Lys Pro Gly Gln Ala Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Asp Thr Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Ser

-continued

50	55	60																	
Gly	Ser	Gly	Ser	Gly	Thr	Gln	Phe	Thr	Leu	Thr	Ile	Ser	Gly	Val	Gln				
65					70					75					80				
Cys	Asp	Asp	Ala	Ala	Ala	Tyr	Tyr	Cys	Leu	Gly	Ser	Tyr	Asp	Cys	Leu				
			85						90					95					
Ser	Ala	Asp	Cys	His	Val	Phe	Gly	Gly	Gly	Thr	Glu	Val	Val	Val	Lys				
			100					105						110					

<210> SEQ ID NO 126  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 126

Gln	Val	Leu	Thr	Gln	Thr	Pro	Ser	Ser	Val	Ser	Ala	Ala	Val	Gly	Gly				
1				5					10					15					
Thr	Val	Thr	Ile	Lys	Cys	Gln	Ser	Ser	Gln	Ser	Val	Tyr	Lys	Ser	Lys				
			20					25					30						
His	Leu	Ser	Trp	Phe	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Lys	Leu	Leu				
		35					40					45							
Ile	Tyr	Asp	Thr	Ser	Thr	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Lys				
	50					55					60								
Gly	Ser	Gly	Ser	Gly	Thr	Gln	Phe	Thr	Leu	Thr	Ile	Ser	Gly	Val	Gln				
65					70					75					80				
Cys	Asp	Asp	Ala	Ala	Ala	Tyr	Tyr	Cys	Leu	Gly	Ser	Tyr	Asp	Cys	Leu				
			85						90					95					
Ser	Ala	Asp	Cys	His	Val	Phe	Gly	Gly	Gly	Thr	Glu	Val	Val	Val	Lys				
			100					105						110					

<210> SEQ ID NO 127  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 127

Gln	Val	Leu	Thr	Gln	Thr	Pro	Ser	Pro	Val	Ser	Ala	Ala	Val	Gly	Gly				
1				5					10					15					
Thr	Val	Thr	Ile	Asn	Cys	Gln	Ala	Ser	Gln	Ser	Val	Tyr	Lys	Ser	Lys				
			20					25					30						
His	Leu	Ser	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Pro	Pro	Lys	Leu	Leu				
		35					40					45							
Ile	Tyr	Asp	Ala	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Lys				
	50					55					60								
Gly	Ser	Gly	Ser	Gly	Thr	His	Phe	Thr	Leu	Thr	Ile	Ser	Gly	Val	Gln				
65					70					75					80				
Cys	Asp	Asp	Ala	Ala	Thr	Tyr	Tyr	Cys	Leu	Gly	Ser	Tyr	Asp	Cys	Ser				
			85						90					95					
Arg	Ala	Asp	Cys	His	Val	Phe	Gly	Gly	Gly	Thr	Glu	Val	Val	Val	Lys				
			100					105						110					

<210> SEQ ID NO 128  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 128

Gln	Val	Leu	Thr	Gln	Thr	Pro	Ser	Pro	Val	Ser	Ala	Ala	Val	Gly	Gly				
1				5					10					15					

-continued

---

Thr Val Thr Ile Asn Cys Arg Ala Ser Gln Thr Val Tyr Lys Ser Lys  
                   20                  25                  30

His Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu  
           35                  40                  45

Ile Tyr Asp Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
       50                  55                  60

Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Glu  
   65                  70                  75                  80

Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Tyr Asp Cys Ser  
           85                  90                  95

Ser Thr Asp Cys His Val Phe Gly Gly Gly Thr Glu Val Val Val Lys  
       100                  105                  110

<210> SEQ ID NO 129  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 129

Gln Val Leu Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly Gly  
   1                  5                  10                  15

Thr Val Thr Ile Asn Cys Gln Ala Ser Gln Arg Val Tyr Lys Asn Lys  
                   20                  25                  30

His Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Arg Pro Lys Leu Leu  
       35                  40                  45

Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Lys  
       50                  55                  60

Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Leu Ser Gly Val Gln  
   65                  70                  75                  80

Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Tyr Asp Cys Thr  
           85                  90                  95

Ile Thr Asp Cys His Val Phe Gly Gly Gly Thr Glu Val Val Val Lys  
       100                  105                  110

<210> SEQ ID NO 130  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 130

Gln Val Leu Thr Gln Thr Pro Ser Pro Val Ser Ala Ala Val Gly Gly  
   1                  5                  10                  15

Thr Val Thr Ile Asn Cys Gln Ala Ser Gln Ser Val Tyr Lys Arg Lys  
                   20                  25                  30

Tyr Leu Ser Trp Tyr Gln Lys Lys Pro Gly Gln Ala Pro Lys Leu Leu  
       35                  40                  45

Ile Tyr Asp Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Ser  
       50                  55                  60

Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val Gln  
   65                  70                  75                  80

Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Tyr Asp Cys Ser  
           85                  90                  95

Arg Thr Asp Cys His Val Phe Ala Gly Gly Thr Glu Leu Val Val Lys  
       100                  105                  110

<210> SEQ ID NO 131

-continued

---

<211> LENGTH: 396  
 <212> TYPE: DNA  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 131

```

atggacacga gggccccac tcagctgctg gggctcctgc tgctctggct cccaggtgcc      60
atatgtgacc ctgtgctgac ccagactcca tcctccgtgt ctgcagctgt gggaggcaca      120
gtcaccatca attgccagtc cagtcagagg gtttgaaga acagctactt atcctggttt      180
cagcagaaac cagggcagcc tccaagcgc ctgatatatt atacatccac tctgccatct      240
ggggtccecat cgcggttcaa aggcagtgga tctgggacac agttcactct caccatcagc      300
gacctggagt gtgacgatgc tgccacttac tactgtctag ggagttatag tgatgatata      360
tattctttcg gcggagggac cgaggtggtg gtcaaa                                396
  
```

<210> SEQ ID NO 132  
 <211> LENGTH: 132  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 132

```

Met Asp Thr Arg Ala Pro Thr Gln Leu Leu Gly Leu Leu Leu Leu Trp
  1          5          10          15
Leu Pro Gly Ala Ile Cys Asp Pro Val Leu Thr Gln Thr Pro Ser Ser
          20          25          30
Val Ser Ala Ala Val Gly Gly Thr Val Thr Ile Asn Cys Gln Ser Ser
          35          40          45
Gln Arg Val Trp Lys Asn Ser Tyr Leu Ser Trp Phe Gln Gln Lys Pro
          50          55          60
Gly Gln Pro Pro Lys Arg Leu Ile Tyr Tyr Thr Ser Thr Leu Pro Ser
          65          70          75          80
Gly Val Pro Ser Arg Phe Lys Gly Ser Gly Ser Gly Thr Gln Phe Thr
          85          90          95
Leu Thr Ile Ser Asp Leu Glu Cys Asp Asp Ala Ala Thr Tyr Tyr Cys
          100          105          110
Leu Gly Ser Tyr Ser Asp Asp Ile Tyr Ser Phe Gly Gly Gly Thr Glu
          115          120          125
Val Val Val Lys
          130
  
```

<210> SEQ ID NO 133  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 133

```

agaatatata tcatcactat aactccctag                                30
  
```

<210> SEQ ID NO 134  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 134

```

gctgctctgg ctcccaggtg                                20
  
```

-continued

<210> SEQ ID NO 135  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 135

Asp Pro Val Leu Thr Gln Thr Pro Ser Ser Val Ser Ala Ala Val Gly  
 1 5 10 15  
 Gly Thr Val Thr Ile Asn Cys Gln Ser Ser Gln Arg Val Trp Lys Asn  
 20 25 30  
 Ser Tyr Leu Ser Trp Phe Gln Gln Lys Pro Gly Gln Pro Pro Lys Arg  
 35 40 45  
 Leu Ile Tyr Tyr Thr Ser Thr Leu Pro Ser Gly Val Pro Ser Arg Phe  
 50 55 60  
 Lys Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Asp Leu  
 65 70 75 80  
 Glu Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Tyr Ser Asp  
 85 90 95  
 Asp Ile Tyr Ser Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 136  
 <211> LENGTH: 110  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 136

Asp Pro Val Leu Thr Gln Thr Pro Ser Ser Val Ser Ala Ala Val Gly  
 1 5 10 15  
 Gly Thr Val Thr Ile Asn Cys Gln Ser Ser Gln Ser Val Tyr Lys Asn  
 20 25 30  
 Lys Tyr Leu Ser Trp Phe Gln Gln Lys Pro Gly Gln Pro Pro Lys Arg  
 35 40 45  
 Leu Ile Tyr Tyr Val Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe  
 50 55 60  
 Lys Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Gly Val  
 65 70 75 80  
 Gln Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Leu Gly Ser Tyr Ser Asn  
 85 90 95  
 Asp Ile Tyr Ser Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 137  
 <211> LENGTH: 402  
 <212> TYPE: DNA  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 137

atggacacga gggcccccac tcagctgctg gggtcctctg tgctctggct cccaggtgcc 60  
 acatttgctg aagtgtgac ccagactgca tgcacctgt ctgcacctgt gggaggcaca 120  
 gtcaccatca attgccagtc cagtcagagt gtttataata acaacgaatt atcttggtat 180  
 cagcagaaac caggacagcc tcccaagctc ctgatctatg ctgcatccat tttggcatct 240  
 ggggtcccat tgcggttcaa aggcagtgga tctgggacac agttcactct caccatcagc 300  
 gacctggagt gtgacgatgc tgccacttac tactgtcaag gcagttatta tagtgggtgt 360  
 tggtaaatg ctttcggcgg agggaccgag gtggtggtca aa 402

-continued

<210> SEQ ID NO 138  
 <211> LENGTH: 134  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*  
  
 <400> SEQUENCE: 138  
 Met Asp Thr Arg Ala Pro Thr Gln Leu Leu Gly Leu Leu Leu Leu Trp  
 1 5 10 15  
 Leu Pro Gly Ala Thr Phe Ala Gln Val Leu Thr Gln Thr Ala Ser Pro  
 20 25 30  
 Val Ser Ala Pro Val Gly Gly Thr Val Thr Ile Asn Cys Gln Ser Ser  
 35 40 45  
 Gln Ser Val Tyr Asn Asn Asn Glu Leu Ser Trp Tyr Gln Gln Lys Pro  
 50 55 60  
 Gly Gln Pro Pro Lys Leu Leu Ile Tyr Ala Ala Ser Ile Leu Ala Ser  
 65 70 75 80  
 Gly Val Pro Leu Arg Phe Lys Gly Ser Gly Ser Gly Thr Gln Phe Thr  
 85 90 95  
 Leu Thr Ile Ser Asp Leu Glu Cys Asp Asp Ala Ala Thr Tyr Tyr Cys  
 100 105 110  
 Gln Gly Ser Tyr Tyr Ser Gly Gly Trp Tyr Asn Ala Phe Gly Gly Gly  
 115 120 125  
 Thr Glu Val Val Val Lys  
 130

<210> SEQ ID NO 139  
 <211> LENGTH: 36  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 139

agcattgtac caaccaccac tataataact gccttg

36

<210> SEQ ID NO 140  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 140

Gln Val Leu Thr Gln Thr Ala Ser Pro Val Ser Ala Pro Val Gly Gly  
 1 5 10 15  
 Thr Val Thr Ile Asn Cys Gln Ser Ser Gln Ser Val Tyr Asn Asn Asn  
 20 25 30  
 Glu Leu Ser Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu  
 35 40 45  
 Ile Tyr Ala Ala Ser Ile Leu Ala Ser Gly Val Pro Leu Arg Phe Lys  
 50 55 60  
 Gly Ser Gly Ser Gly Thr Gln Phe Thr Leu Thr Ile Ser Asp Leu Glu  
 65 70 75 80  
 Cys Asp Asp Ala Ala Thr Tyr Tyr Cys Gln Gly Ser Tyr Tyr Ser Gly  
 85 90 95  
 Gly Trp Tyr Asn Ala Phe Gly Gly Gly Thr Glu Val Val Val Lys  
 100 105 110

<210> SEQ ID NO 141

-continued

---

<211> LENGTH: 396  
 <212> TYPE: DNA  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 141

```

atggacacga gggcccccac tcagctgctg gggctcctgc tgctctggct cccaggtgcc      60
acatttgctc aagtgtgac ccagactcca ccctccgtgt ctgcagctgt gggaggcaca      120
gtcaccatca gttgccagtc cagtcagagc gtttataata ataactgggtt aggctggtat      180
cagcagaaat cagggcagcc tccaagctc ctgatctatt atgcatccac tctggcatct      240
ggggtctcat cgcggttcaa aggcagtgga tctgggacac agttcactct caccatcagc      300
gacctggagt gtgacgatgc tgccacttat tattgtgcag gcggttatag tgatatgatg      360
aatgctttcg gcggagggac tgaggtggtg gttaaa                                396
  
```

<210> SEQ ID NO 142  
 <211> LENGTH: 132  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 142

```

Met Asp Thr Arg Ala Pro Thr Gln Leu Leu Gly Leu Leu Leu Leu Trp
 1           5           10           15
Leu Pro Gly Ala Thr Phe Ala Gln Val Leu Thr Gln Thr Pro Pro Ser
           20           25           30
Val Ser Ala Ala Val Gly Gly Thr Val Thr Ile Ser Cys Gln Ser Ser
           35           40           45
Gln Ser Val Tyr Asn Asn Asn Trp Leu Gly Trp Tyr Gln Gln Lys Ser
           50           55           60
Gly Gln Pro Pro Lys Leu Leu Ile Tyr Tyr Ala Ser Thr Leu Ala Ser
65           70           75           80
Gly Val Ser Ser Arg Phe Lys Gly Ser Gly Ser Gly Thr Gln Phe Thr
           85           90           95
Leu Thr Ile Ser Asp Leu Glu Cys Asp Asp Ala Ala Thr Tyr Tyr Cys
           100          105          110
Ala Gly Gly Tyr Ser Asp Met Met Asn Ala Phe Gly Gly Gly Thr Glu
           115          120          125
Val Val Val Lys
           130
  
```

<210> SEQ ID NO 143  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 143

```

gcagggcggtt atagtgatat gatgaatgct                                30
  
```

<210> SEQ ID NO 144  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 144

```

Gln Val Leu Thr Gln Thr Pro Pro Ser Val Ser Ala Ala Val Gly Gly
 1           5           10           15
Thr Val Thr Ile Ser Cys Gln Ser Ser Gln Ser Val Tyr Asn Asn Asn
  
```



-continued

&lt;400&gt; SEQUENCE: 147

tgaggagacg gtgaccaggg tgccctgggcc ccacaaactc cattg 45

&lt;210&gt; SEQ ID NO 148

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Primer

&lt;400&gt; SEQUENCE: 148

ctgcgctggc ttctctggt c 21

&lt;210&gt; SEQ ID NO 149

&lt;211&gt; LENGTH: 109

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Oryctolagus cuniculus

&lt;400&gt; SEQUENCE: 149

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Thr Tyr Ala  
20 25 30Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Ala  
35 40 45Ala Phe Gly Thr Ser Gly Thr Thr Asn Tyr Ala Thr Trp Ala Lys Gly  
50 55 60Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Met Asp Leu Arg Ile Thr  
65 70 75 80Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
85 90 95Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
100 105

&lt;210&gt; SEQ ID NO 150

&lt;211&gt; LENGTH: 109

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Oryctolagus cuniculus

&lt;400&gt; SEQUENCE: 150

Gln Ser Val Glu Glu Phe Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
1 5 10 15Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Thr Val Ala  
20 25 30Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Ala  
35 40 45Ala Phe Gly Thr Ser Gly Thr Thr Asn Tyr Ala Thr Trp Ala Lys Gly  
50 55 60Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Val Asp Leu Arg Ile Thr  
65 70 75 80Arg Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
85 90 95Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
100 105

&lt;210&gt; SEQ ID NO 151

&lt;211&gt; LENGTH: 109

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Oryctolagus cuniculus

-continued

&lt;400&gt; SEQUENCE: 151

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Thr Val Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Ala  
 35 40 45  
 Ala Phe Gly Thr Ser Gly Thr Thr Asn Tyr Ala Thr Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Val Asp Leu Arg Ile Thr  
 65 70 75 80  
 Arg Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
 85 90 95  
 Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105

&lt;210&gt; SEQ ID NO 152

&lt;211&gt; LENGTH: 109

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 152

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ser Leu Ser Thr Val Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Gln Trp Ile Ala  
 35 40 45  
 Ala Phe Gly Thr Arg Gly Thr Thr Asn Tyr Ala Thr Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Val Asp Leu Arg Ile Thr  
 65 70 75 80  
 Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
 85 90 95  
 Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105

&lt;210&gt; SEQ ID NO 153

&lt;211&gt; LENGTH: 109

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Oryctolagus cuniculus*

&lt;400&gt; SEQUENCE: 153

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Val Ser Leu Arg Gly Tyr Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Lys Trp Ile Ala  
 35 40 45  
 Ala Phe Gly Thr Ser Gly Thr Thr Asn Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Val Asp Leu Lys Ile Thr  
 65 70 75 80  
 Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
 85 90 95

-continued

---

Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105

<210> SEQ ID NO 154  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 154

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Ile Ser Gly Val Ser Leu Arg Gly Tyr Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Lys Trp Ile Ala  
 35 40 45  
 Ala Phe Gly Thr Ser Gly Thr Thr Asn Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Val Asp Leu Lys Ile Thr  
 65 70 75 80  
 Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
 85 90 95  
 Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105

<210> SEQ ID NO 155  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 155

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Arg Ser Tyr Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Ala  
 35 40 45  
 Ala Phe Gly Thr Ser Gly Thr Thr Asn Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Val Tyr Leu Lys Ile Thr  
 65 70 75 80  
 Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
 85 90 95  
 Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105

<210> SEQ ID NO 156  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 156

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Arg Ser Tyr Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Ala  
 35 40 45  
 Ala Phe Gly Thr Ser Gly Thr Thr Asn Tyr Ala Ser Trp Ala Lys Gly

-continued

---

50	55	60	
Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Val Tyr Leu Arg Ile Thr			
65	70	75	80
Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp			
	85	90	95
Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser			
	100	105	

<210> SEQ ID NO 157  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 157

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro			
1	5	10	15
Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Tyr Ala			
	20	25	30
Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Ala			
	35	40	45
Ala Phe Gly Thr Ser Gly Ser Thr Asn Tyr Ala Ser Trp Ala Lys Gly			
	50	55	60
Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Met His Leu Lys Ile Thr			
65	70	75	80
Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp			
	85	90	95
Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser			
	100	105	

<210> SEQ ID NO 158  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 158

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro			
1	5	10	15
Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Asn Tyr Ala			
	20	25	30
Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Glu Trp Ile Ala			
	35	40	45
Ala Phe Gly Thr Ser Gly Ser Ala Ser Tyr Ala Ser Trp Ala Lys Gly			
	50	55	60
Arg Phe Thr Ile Ser Lys Thr Ser Asn Thr Val Asp Leu Lys Ile Thr			
65	70	75	80
Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp			
	85	90	95
Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser			
	100	105	

<210> SEQ ID NO 159  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 159

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro			
1	5	10	15

-continued

---

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Lys Tyr Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Glu Gly Leu Lys Trp Ile Ala  
 35 40 45  
 Ala Leu Gly Ala Ser Gly Thr Thr Asn Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Arg Thr Ser Asn Thr Val Asp Leu Lys Ile Thr  
 65 70 75 80  
 Ser Pro Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
 85 90 95  
 Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105

<210> SEQ ID NO 160  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 160

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Tyr Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ala  
 35 40 45  
 Ala Phe Gly Thr Ser Gly Thr Thr Asn Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Lys Thr Ser Asn Thr Val Asp Leu Lys Ile Thr  
 65 70 75 80  
 Ser Pro Thr Pro Gly Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
 85 90 95  
 Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Ile Ser Ser  
 100 105

<210> SEQ ID NO 161  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 161

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Ser Tyr Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ala  
 35 40 45  
 Ala Phe Gly Thr Ser Gly Thr Arg Asn Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Arg Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
 65 70 75 80  
 Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
 85 90 95  
 Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105

<210> SEQ ID NO 162

-continued

---

<211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 162

Gln Ser Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr Pro  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Gly Ser Tyr Ala  
 20 25 30  
 Met Ile Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ala  
 35 40 45  
 Ala Phe Gly Thr Ser Gly Thr Arg Asn Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Arg Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
 65 70 75 80  
 Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gln Trp  
 85 90 95  
 Ser Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105

<210> SEQ ID NO 163  
 <211> LENGTH: 399  
 <212> TYPE: DNA  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 163

atggagactg ggtcgcgctg gcttctcctg gtcgctgtgc tcaaagggtg ccagtgtcag 60  
 tcgctggagg agtccggggg tcgectggta acgcctggag gatecctgac actcaectgc 120  
 acagtctctg gaatcgacct cagtacctat ccaatgggct gggtcgcca ggtccaggg 180  
 aaggggctgg aatacatcgg aatcgttttt cctagtcttg gctcatatta cgcgagctgg 240  
 gcaaaaggcc gattcaccat ctccaaaacc tcgtcaacca cggtgatct gcgcatgacc 300  
 agtctgacaa ccgaggacac ggccacctat ttctgtgcca gaggggtaac taatagttgg 360  
 gatccctggg gccccaggcac cctggtcacc gtctcctca 399

<210> SEQ ID NO 164  
 <211> LENGTH: 133  
 <212> TYPE: PRT  
 <213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 164

Met Glu Thr Gly Leu Arg Trp Leu Leu Leu Val Ala Val Leu Lys Gly  
 1 5 10 15  
 Val Gln Cys Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro  
 20 25 30  
 Gly Gly Ser Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser  
 35 40 45  
 Thr Tyr Pro Met Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu  
 50 55 60  
 Tyr Ile Gly Ile Val Phe Pro Ser Leu Gly Ser Tyr Tyr Ala Ser Trp  
 65 70 75 80  
 Ala Lys Gly Arg Phe Thr Ile Ser Lys Thr Ser Ser Thr Thr Val Asp  
 85 90 95  
 Leu Arg Met Thr Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys  
 100 105 110  
 Ala Arg Gly Val Thr Asn Ser Trp Asp Pro Trp Gly Pro Gly Thr Leu

-continued

---

115                      120                      125

Val Thr Val Ser Ser  
130

<210> SEQ ID NO 165  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 165

cccagggatc ccaactatta gttacc                      26

<210> SEQ ID NO 166  
<211> LENGTH: 114  
<212> TYPE: PRT  
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 166

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Ser Pro Gly Gly Ser  
1                      5                      10                      15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Thr Tyr Pro  
                    20                      25                      30

Met Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Ile Gly  
                    35                      40                      45

Ile Val Phe Pro Ser Leu Gly Ser Tyr Tyr Ala Ser Trp Ala Lys Gly  
50                      55                      60

Arg Phe Thr Ile Ser Lys Thr Ser Ser Thr Thr Val Asp Leu Arg Met  
65                      70                      75                      80

Thr Ser Leu Thr Ala Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly  
                    85                      90                      95

Val Thr Asn Ser Trp Asp Pro Trp Gly Pro Gly Thr Val Val Thr Val  
                    100                      105                      110

Ser Ser

<210> SEQ ID NO 167  
<211> LENGTH: 114  
<212> TYPE: PRT  
<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 167

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Gly Ser  
1                      5                      10                      15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Thr Tyr Pro  
20                      25                      30

Met Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Ile Gly  
35                      40                      45

Ile Val Phe Pro Ser Leu Gly Ser Tyr Tyr Ala Ser Trp Ala Lys Gly  
50                      55                      60

Arg Phe Thr Ile Ser Lys Thr Ser Ser Thr Thr Val Asp Leu Arg Met  
65                      70                      75                      80

Thr Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly  
85                      90                      95

Val Thr Asn Ser Trp Asp Pro Trp Gly Pro Gly Thr Leu Val Thr Val  
100                      105                      110

Ser Ser

-continued

<210> SEQ ID NO 168  
 <211> LENGTH: 114  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 168

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Gly Ser  
 1 5 10 15  
 Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Thr Tyr Pro  
 20 25 30  
 Met Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Ile Gly  
 35 40 45  
 Ile Val Phe Pro Asn Ile Gly Ser Tyr Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60  
 Arg Phe Thr Ile Ser Ser Thr Ser Ser Thr Thr Val Asp Leu Arg Met  
 65 70 75 80  
 Thr Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly  
 85 90 95  
 Val Thr Asn Ser Trp Asp Pro Trp Gly Pro Gly Thr Leu Val Thr Val  
 100 105 110  
 Ser Ser

<210> SEQ ID NO 169  
 <211> LENGTH: 393  
 <212> TYPE: DNA  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 169

atggagactg ggctgcgctg gcttctcctg gtcgctgtgc tcaaagggtg ccagtgtcag 60  
 tcgctggagg agtccggggg tcgctggta acgcctggag gatccctgac actcacctgc 120  
 acagtctctg gaatcgacct cagtagctat ggaatgggct gggtccgcca ggctccaggg 180  
 aagggtctgg aatacatcgc aatcattagt tatggtggta gagcatacta cgcgagctgg 240  
 gcgaaaggcc gattcaccat ctccagaact tcgaccacgg tggatctgaa aatgaccagt 300  
 ctgacaaccg aggacacggc cacctatttc tgtgccagag gatttagcgc ctttaacttg 360  
 tggggcccag gcaccctggt caccgtctcc tca 393

<210> SEQ ID NO 170  
 <211> LENGTH: 131  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 170

Met Glu Thr Gly Leu Arg Trp Leu Leu Leu Val Ala Val Leu Lys Gly  
 1 5 10 15  
 Val Gln Cys Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro  
 20 25 30  
 Gly Gly Ser Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser  
 35 40 45  
 Ser Tyr Gly Met Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu  
 50 55 60  
 Tyr Ile Ala Ile Ile Ser Tyr Gly Gly Arg Ala Tyr Tyr Ala Ser Trp  
 65 70 75 80  
 Ala Lys Gly Arg Phe Thr Ile Ser Arg Thr Ser Thr Thr Val Asp Leu  
 85 90 95

-continued

Lys Met Thr Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala  
 100 105 110

Arg Gly Phe Ser Ala Phe Asn Leu Trp Gly Pro Gly Thr Leu Val Thr  
 115 120 125

Val Ser Ser  
 130

<210> SEQ ID NO 171  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 171

gccccacaag ttaaaggcgc taaatc

26

<210> SEQ ID NO 172  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 172

Gln Ser Leu Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Gly Ser  
 1 5 10 15

Leu Thr Leu Thr Cys Thr Val Ser Gly Ile Asp Leu Ser Ser Tyr Gly  
 20 25 30

Met Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Ile Ala  
 35 40 45

Ile Ile Ser Tyr Gly Gly Arg Ala Tyr Tyr Ala Ser Trp Ala Lys Gly  
 50 55 60

Arg Phe Thr Ile Ser Arg Thr Ser Thr Thr Val Asp Leu Lys Met Thr  
 65 70 75 80

Ser Leu Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg Gly Phe  
 85 90 95

Ser Ala Phe Asn Leu Trp Gly Pro Gly Thr Leu Val Thr Val Ser Ser  
 100 105 110

<210> SEQ ID NO 173  
 <211> LENGTH: 399  
 <212> TYPE: DNA  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 173

atggagactg ggctgcgctg gcttctcctg gtcgctgtgc tcaaaggtgt ccagtgtcag 60

tcggtggtgg aggagtccgg gggtcgcctg gtcacgcctg ggacaccct gacactcacc 120

tgcacagcct ctggattctc cctcagtagg tttgcaatga ggtgggtccg ccaggctcca 180

gggaaggggc tggaatacat cggagccatc gagactgatg gtaggacata ctacgcgagg 240

tgggcgaaag gccgattcac cattccaag acctcgaccg cggtgcatct gaagttcacc 300

agtccgacaa ccgaggacac gggcacgtat ttctgtacca gagggctggt tacaatttct 360

actttgtggg gcccaggcac cctggtcacc gtctctca 399

<210> SEQ ID NO 174  
 <211> LENGTH: 133  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 174

-continued

Met Glu Thr Gly Leu Arg Trp Leu Leu Leu Val Ala Val Leu Lys Gly  
 1 5 10 15  
 Val Gln Cys Gln Ser Val Val Glu Glu Ser Gly Gly Arg Leu Val Thr  
 20 25 30  
 Pro Gly Thr Pro Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ser Leu  
 35 40 45  
 Ser Arg Phe Ala Met Arg Trp Val Arg Gln Ala Pro Gly Lys Gly Leu  
 50 55 60  
 Glu Tyr Ile Gly Ala Ile Glu Thr Asp Gly Arg Thr Tyr Tyr Ala Arg  
 65 70 75 80  
 Trp Ala Lys Gly Arg Phe Thr Ile Ser Lys Thr Ser Thr Ala Val His  
 85 90 95  
 Leu Lys Phe Thr Ser Pro Thr Thr Glu Asp Thr Gly Thr Tyr Phe Cys  
 100 105 110  
 Thr Arg Gly Leu Val Thr Ile Ser Thr Leu Trp Gly Pro Gly Thr Leu  
 115 120 125  
 Val Thr Val Ser Ser  
 130

<210> SEQ ID NO 175  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 175

gccccacaaa gtagaattg taaccagc

28

<210> SEQ ID NO 176  
 <211> LENGTH: 114  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryctolagus cuniculus*

<400> SEQUENCE: 176

Gln Ser Val Val Glu Glu Ser Gly Gly Arg Leu Val Thr Pro Gly Thr  
 1 5 10 15  
 Pro Leu Thr Leu Thr Cys Thr Ala Ser Gly Phe Ser Leu Ser Arg Phe  
 20 25 30  
 Ala Met Arg Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Tyr Ile  
 35 40 45  
 Gly Ala Ile Glu Thr Asp Gly Arg Thr Tyr Tyr Ala Arg Trp Ala Lys  
 50 55 60  
 Gly Arg Phe Thr Ile Ser Lys Thr Ser Thr Ala Val His Leu Lys Phe  
 65 70 75 80  
 Thr Ser Pro Thr Thr Glu Asp Thr Gly Thr Tyr Phe Cys Thr Arg Gly  
 85 90 95  
 Leu Val Thr Ile Ser Thr Leu Trp Gly Pro Gly Thr Leu Val Thr Val  
 100 105 110  
 Ser Ser

<210> SEQ ID NO 177  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer

-continued

---

<400> SEQUENCE: 177

caccatggag actgggctgc gctggcttct cctggtcgct gtg 43

<210> SEQ ID NO 178  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 178

ctcccgtct cgggtaaat gagegctgtg ccggcga 37

<210> SEQ ID NO 179  
<211> LENGTH: 36  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 179

caggcaggac ccagcatgga cagcaggcc cccact 36

<210> SEQ ID NO 180  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 180

tcaatagggg tgactgtag agcgagcgc ctgc 34

<210> SEQ ID NO 181  
<211> LENGTH: 43  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 181

caccatggag actgggctgc gctggcttct cctggtcgct gtg 43

<210> SEQ ID NO 182  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 182

ctcccgtct cgggtaaat gagegctgtg ccggcga 37

<210> SEQ ID NO 183  
<211> LENGTH: 36  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 183

caggcaggac ccagcatgga cagcaggcc cccact 36

-continued

---

<210> SEQ ID NO 184  
 <211> LENGTH: 34  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer  
  
 <400> SEQUENCE: 184  
  
 tcaatagggg tgactgtag agcgagacgc ctgc 34

<210> SEQ ID NO 185  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer  
  
 <400> SEQUENCE: 185  
  
 aattcacatt gattattgag 20

<210> SEQ ID NO 186  
 <211> LENGTH: 40  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer  
  
 <400> SEQUENCE: 186  
  
 cagcgagcc cagtctccat cccgtaagca gtgggttctc 40

<210> SEQ ID NO 187  
 <211> LENGTH: 40  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer  
  
 <400> SEQUENCE: 187  
  
 ccgggtaaat gagcgctgtg gtttaaaccg gctgatcagc 40

<210> SEQ ID NO 188  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer  
  
 <400> SEQUENCE: 188  
  
 aagccataga gccgaccgca 20

<210> SEQ ID NO 189  
 <211> LENGTH: 11  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 189  
  
 cggaacaagg g 11

<210> SEQ ID NO 190  
 <211> LENGTH: 11  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence

-continued

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic oligonucleotide

&lt;400&gt; SEQUENCE: 190

cggaacaagg g

11

What is claimed is:

1. A method for identifying an antibody that binds to an antigen, comprising:

a) obtaining: (i) antibody heavy chain variable domain sequences and (ii) antibody light chain variable domain sequences by sequencing heavy and light chain-encoding cDNAs made from a population of antibody-producing B cells of an animal that has been immunized by the antigen;

b) grouping the heavy chain variable domain sequences of step a) into a plurality of heavy chain groups and, independently, grouping the light chain variable domain sequences of step a) into a plurality of light chain groups, wherein:

(i) heavy chain sequences that comprise CDR3 regions that are of the same length and contain up to five amino acid substitutions relative to one another are grouped together; and

(ii) light chain sequences that comprise CDR3 regions that are of the same length and contain up to five amino acid substitutions relative to one another are grouped together;

c) selecting a heavy chain group and a light chain group from the groups produced by step b);

d) pairing a heavy chain variable domain sequence and a light chain variable domain sequence from the groups selected in c); and

e) testing an antibody comprising the heavy and light chain sequences paired in d) for binding to the antigen.

2. The method of claim 1, wherein the heavy and light chain variable domain sequences paired in step d) are not naturally paired together in the animal of step a).

3. The method of claim 1, wherein:

(i) heavy chain sequences that comprise CDR3 regions that are of the same length and contain 0, 1 or 2 amino acid substitutions relative to one another are grouped together; and

(ii) light chain sequences that comprise CDR3 regions that are of the same length and contain 0, 1 or 2 amino acid substitutions relative to one another are grouped together.

4. The method of claim 1, wherein the grouping step (b) further comprises comparing the entire lengths of the heavy and light chain variable domain sequences to one another.

5. The method of claim 1, wherein the obtaining step a) is done by sequencing cDNAs encoding the heavy and light chain variable domain sequences from single B cells, or cultures of the same.

6. The method of claim 1, wherein the obtaining step a) is done by sequencing cDNAs encoding the heavy and light chain variable domain sequences made from a plurality of pools of B cells.

7. The method of claim 1, wherein the heavy and light chain groups of selecting step c) are selected by analysis of lineage trees.

8. The method of claim 1, wherein, prior to step a), the antibody-producing B cells are enriched by binding to the antigen.

9. The method of claim 1, wherein, prior to step a), the antibody-producing B cells are not enriched by binding to the antigen.

10. The method of claim 1, wherein the selected heavy chain and light chain groups selected in step c) both contain at least 10 members.

11. The method of claim 1, wherein the testing step e) comprises testing the antibody of e) in a blocking assay and/or a neutralization assay.

12. The method of claim 1, wherein the testing step e) comprises testing the antibody of e) in an ELISA.

13. The method of claim 1, wherein the obtaining step a) comprises obtaining at least 1,000 different antibody heavy chain variable domain sequences and at least 1,000 different antibody light chain variable domain sequences.

14. The method of claim 1, wherein the antibody-producing B cells are cells obtained from spleen, lymph node or peripheral blood.

15. The method of claim 1, wherein the animal is a rabbit.

16. A method for identifying a monoclonal antibody comprising:

a) obtaining a population of antibody-producing B cells of an animal that has been immunized by an antigen;

b) making pools of the antibody-producing B cells obtained in step a) to produce a plurality of pools;

c) sequencing antibody heavy chain and antibody light chain-encoding cDNAs from each of the pools of step b);

d) grouping the heavy chain variable domain sequences of step c) into a plurality of heavy chain groups and, independently, grouping the light chain variable domain sequences of step c) into a plurality of light chain groups, wherein:

(i) heavy chain sequences that comprise CDR3 regions that are of the same length and contain up to five amino acid substitutions relative to one another are grouped together; and

(ii) light chain sequences that comprise CDR3 regions that are of the same length and contain up to five amino acid substitutions relative to one another are grouped together;

e) selecting a heavy chain group and a light chain group from the groups produced by step d);

f) pairing a heavy chain variable domain sequence and a light chain variable domain sequence from the groups selected in step e); and

g) testing an antibody comprising the heavy and light chain sequences paired in step f) for binding to the antigen.

17. The method of claim 16, wherein the pools comprise at least 1,000 different antibody-producing B cells.

18. The method of claim 16, wherein the sequencing step c) provides at least 10,000 different heavy chain variable domain sequences and at least 10,000 different light chain variable domain sequences.

19. The method of claim 16, wherein, in step d):

(i) heavy chain sequences that are grouped together comprise CDR3 regions that are of the same length and contain 0, 1, 2 or 3 amino acid substitutions relative to one another; and

(ii) light chain sequences that are grouped together comprise CDR3 regions that are of the same length and contain 0, 1, 2 or 3 amino acid substitutions relative to one another.

20. The method of claim 16, wherein the grouping step d) 5 further comprises comparing the entire lengths of the heavy and light chain variable domain sequences to one another.

21. The method of claim 16, wherein the heavy and light chain groups selected in step e) are selected by analysis of lineage trees. 10

22. The method of claim 16, wherein, prior to step a), the B cells are enriched by binding to the antigen.

23. The method of claim 16, wherein, prior to step a), the B cells are not enriched by binding to the antigen.

24. The method of claim 16, wherein the grouping of step 15 a) results in at least 10 groups of heavy chain sequences and at least 10 groups of light chain sequences.

25. The method of claim 16, wherein the B cells are cells obtained from spleen, lymph node or peripheral blood.

26. The method of claim 16, wherein the animal is a rabbit. 20

\* \* \* \* \*

专利名称(译)	鉴定谱系相关抗体的方法		
公开(公告)号	<a href="#">US9052322</a>	公开(公告)日	2015-06-09
申请号	US14/177003	申请日	2014-02-10
[标]申请(专利权)人(译)	Epitomics公司INC ABCAM		
申请(专利权)人(译)	Epitomics公司, INC. ( C / O ABCAM PLC )		
当前申请(专利权)人(译)	Epitomics公司, INC.		
[标]发明人	YU GUO LIANG		
发明人	YU, GUO-LIANG		
IPC分类号	G01N33/53 C07K16/28 G01N33/577 C07K16/00 G01N33/68 G01N33/566		
CPC分类号	G01N33/577 C07K16/00 C07K16/2863 G01N33/6854 G01N33/566 C07K2317/10 C07K2317/56 C07K2317/76 C12Q1/6876		
代理人(译)	凯迪, JAMES S.		
优先权	61/241714 2009-09-11 US		
其他公开文献	US20140155291A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

在某些实施方案中，该方法可包括：a) 从B细胞群获得抗体序列；b) 对抗体序列进行分组以提供多组谱系相关抗体；c) 在生物测定中测试来自每个组的单个抗体，并且在已经鉴定第一抗体后，d) 在第二生物测定中测试与第一抗体在同一组中的其他抗体。在另一个实施方案中，该方法可以包括：a) 测试从动物的抗体产生器官的第一部分获得的多种抗体；b) 获得第一个鉴定的抗体的序列；c) 从所述抗体生成器官的第二部分获得通过谱系与所述第一抗体相关的其他抗体的序列；c) 在第二次生物测定中测试其他抗体。

