



US 20100279433A1

(19) **United States**

(12) **Patent Application Publication**
Holtzman et al.

(10) **Pub. No.: US 2010/0279433 A1**
(43) **Pub. Date: Nov. 4, 2010**

(54) **ASSAY METHOD FOR ALZHEIMER'S DISEASE**

(76) Inventors: **David M. Holtzman**, St. Louis, MO (US); **Ronald DeMattos**, Noblesville, IN (US); **Kelly R. Bales**, Coatsville, IN (US); **David J. Cummins**, Indianapolis, IN (US); **Steven M. Paul**, Carmel, IN (US)

Correspondence Address:
MCANDREWS HELD & MALLOY, LTD
500 WEST MADISON STREET, SUITE 3400
CHICAGO, IL 60661

(21) Appl. No.: **12/834,271**

(22) Filed: **Jul. 12, 2010**

Related U.S. Application Data

(63) Continuation of application No. 10/486,908, filed on Feb. 17, 2004, now Pat. No. 7,771,722, filed as application No. PCT/US02/26321 on Aug. 16, 2002.

(60) Provisional application No. 60/313,221, filed on Aug. 17, 2001, provisional application No. 60/334,987, filed on Oct. 23, 2001.

Publication Classification


(51) **Int. Cl.**
G01N 33/53 (2006.01)

(52) **U.S. Cl.** **436/501**

(57) **ABSTRACT**


A diagnostic test for preclinical and clinical Alzheimer's disease is based on plasma levels of $A\beta_{40}$, $A\beta_{42}$, their ratio, or their rate of entry following administration of antibodies that sequester $A\beta$. Alterations of any of these parameters from control values identifies preclinical or clinical Alzheimer's disease.

A




Parameter	PB 40
Number of XY Pairs	42
Pearson r	0.02586
95% confidence interval	-0.2804 to 0.3273
P value (two-tailed)	0.8709
P value summary	ns
Is the correlation significant? (alpha=0.05)	No
R squared	0.0006685

B



Parameter	PB 42
Number of XY Pairs	47
Pearson r	-0.07367
95% confidence interval	-0.3536 to 0.2180
P value (two-tailed)	0.6217
P value summary	ns
Is the correlation significant? (alpha=0.05)	No
R squared	0.005456


C



Parameter	PB ratio
Number of XY Pairs	40
Pearson r	0.1213
95% confidence interval	-0.1978 to 0.4171
P value (two-tailed)	0.4560
P value summary	ns
Is the correlation significant? (alpha=0.05)	No
R squared	0.01471


Figure 1

A



Parameter	1-Hour 40
Number of XY Pairs	52
Pearson r	0.6567
95% confidence interval	0.4676 to 0.7884
P value (two-tailed)	P<0.0001
P value summary	***
Is the correlation significant? (alpha=0.05)	Yes
R squared	0.4313


B



Parameter	1-Hour Ratio
Number of XY Pairs	52
Pearson r	0.7565
95% confidence interval	0.6093 to 0.8533
P value (two-tailed)	P<0.0001
P value summary	***
Is the correlation significant? (alpha=0.05)	Yes
R squared	0.5723


Figure 2

A




Parameter	24-Hour 40
Number of XY Pairs	52
Pearson r	0.6628
95% confidence interval	0.4759 to 0.7924
P value (two-tailed)	P<0.0001
P value summary	***
Is the correlation significant? (alpha=0.05)	Yes
R squared	0.4393

B



Parameter	24-Hour 42
Number of XY Pairs	52
Pearson r	0.4039
95% confidence interval	0.1471 to 0.6096
P value (two-tailed)	0.0030
P value summary	**
Is the correlation significant? (alpha=0.05)	Yes
R squared	0.1631

C




Parameter	24-Hour Ratio
Number of XY Pairs	52
Pearson r	0.7987
95% confidence interval	0.6724 to 0.8799
P value (two-tailed)	P<0.0001
P value summary	***
Is the correlation significant? (alpha=0.05)	Yes
R squared	0.6380

Figure 3

A


$A\beta_{40}$ Entry Rate Into Plasma



Parameter	40 rate slope
Number of XY Pairs	52
Pearson r	0.6360
95% confidence interval	0.4394 to 0.7745
P value (two-tailed)	P<0.0001
P value summary	***
Is the correlation significant? (alpha=0.05)	Yes
R squared	0.4046

B


$A\beta_{42}$ Entry Rate Into Plasma



Parameter	42 rate slope
Number of XY Pairs	52
Pearson r	0.4062
95% confidence interval	0.1499 to 0.6114
P value (two-tailed)	0.0028
P value summary	**
Is the correlation significant? (alpha=0.05)	Yes
R squared	0.1650

C

Ratio of the $A\beta_{40}/A\beta_{42}$
Entry Rate into Plasma



Parameter	Ratio Rate
Number of XY Pairs	52
Pearson r	0.6551
95% confidence interval	0.4653 to 0.7873
P value (two-tailed)	P<0.0001
P value summary	***
Is the correlation significant? (alpha=0.05)	Yes
R squared	0.4291

Figure 4





Figure 5

<u>Plasma Aβ Correlation's with Alzheimer-Like Pathology in Hippocampus</u>							
Plasma A β correlation with A β load and fibrillar amyloid							
	<u>Pre-Bleed</u>	<u>5-Min</u>	<u>1-Hour</u>	<u>3-Hour</u>	<u>6-Hour</u>	<u>24-Hour</u>	<u>AUC</u>
<u>Plasma Aβ40:</u>							
A β Load:	Pearson r	-0.0158	0.5527	0.5904	0.4310	0.5533	0.5932
	P value	0.9209	<0.0001	<0.0001	0.0014	<0.0001	<0.0001
Amyloid Load:	Pearson r	0.1535	0.7420	0.6257	0.7053	0.6684	0.7432
	P value	0.3378	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
<u>Plasma Aβ42:</u>							
A β Load:	Pearson r	-0.0614	0.2223	-0.0036	0.1309	0.4551	0.3391
	P value	0.6817	0.1207	0.9798	0.3549	0.0008	0.0139
Amyloid Load:	Pearson r	0.0443	0.4790	0.2321	0.3996	0.4476	0.6062
	P value	0.7698	0.0005	0.1013	0.0037	0.0011	<0.0001
<u>Aβ40/42 Ratio:</u>							
A β Load:	Pearson r	0.0369	0.5223	0.6888	0.4215	0.1754	0.7190
	P value	0.8236	<0.0001	<0.0001	0.0019	0.2183	<0.0001
Amyloid Load:	Pearson r	0.1293	0.4825	0.5047	0.4364	0.2843	0.6029
	P value	0.4393	0.0004	0.0002	0.0014	0.0454	<0.0001

Figure 6

ASSAY METHOD FOR ALZHEIMER'S DISEASE

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the priority of U.S. provisional applications 60/334,987, filed Oct. 23rd, 2001, 60/313,221, filed Aug. 17th, 2001, and 60/313,224, filed Aug. 17th, 2001, the contents of which are incorporated herein by reference.

TECHNICAL FIELD

[0002] The invention relates to an assay which permits diagnosis of preclinical and clinical Alzheimer's disease. The test relies on assessing the levels of amyloid beta (A β) peptide in plasma following administration of certain anti-A β antibodies to a subject.

BACKGROUND ART

[0003] A number of symptomologies which result in cognitive deficits, stroke, brain hemorrhage, and general mental debilitation appear to be associated with neuritic and cerebrovascular plaques in the brain containing the amyloid beta peptide (A β). Among these conditions are both preclinical and clinical Alzheimer's disease, Down's syndrome, and preclinical and clinical cerebral amyloid angiopathy (CAA). The amyloid plaques are formed from amyloid beta peptides. These peptides circulate in the blood and in the cerebrospinal fluid (CSF). The A β peptide in circulating form is composed of 39-43 amino acids (mostly 40 or 42 amino acids) resulting from the cleavage of a common precursor protein, amyloid precursor protein, often designated APP.

[0004] Evidence suggests that A β can be transported back and forth between brain and the blood (Ghersi-Egea, J-F., et al., *J. Neurochem.* (1996) 67:880-883; Zlokovic, B. V., et al., *Biochem. Biophys. Res. Comm.* (1993) 67:1034-1040; Shibata, M., et al., *J. Clin. Invest.* (2000) 106:1489-1499. Further A β in plaques is in an equilibrium with soluble A β in the brain and blood (Kawarabayashi, T., et al., *J. Neurosci.* (2001) 21:372-381), DeMattos et al., *Proc. Natl. Acad. Sci USA* (2001) 98:8850-8855.

[0005] As described in PCT application US00/35681 and U.S. Ser. No. 09/153,130 both incorporated herein by reference, total circulating levels of A β peptide in CSF are similar in normal individuals and individuals predisposed to exhibit the symptoms of Alzheimer's. However, A β ₄₂ levels are lower on average in individuals with Alzheimer's disease (Nitsch, R. M., et al., *Ann. Neurol.* (1995) 37:512-518). It is known that A β ₄₂ is more prone to aggregate than is A β ₄₀, and when this happens, adverse consequences such as A β deposition in amyloid plaques, conversion of A β to toxic forms, nerve cell damage, and behavioral impairment such as dementia ensue (Golde, T. E., et al., *Biochem. Biophys. Acta.* (2000) 1502: 172-187).

[0006] PCT application PCT/US01/06191 entitled "Humanized Antibodies That Sequester A β Peptide" filed 26 Feb. 2001 and incorporated herein by reference describes antibodies which do not appreciably cross the blood-brain barrier and which sequester A β peptides circulating in biological fluids. These antibodies are described as useful for preventive and therapeutic treatment of conditions associated with the formation of A β -containing diffuse, neuritic, and cerebrovascular plaques in the brain. The application

describes administering the antibodies and then measuring circulating levels of A β peptide in blood in order to assess the progress of therapy. There is no clear suggestion, however, that the levels of A β peptide following administration of the antibodies are diagnostic of the condition itself. The present invention resides in the surprising result that enhanced levels of both A β ₄₀ and A β ₄₂ as well as the A β ₄₀/A β ₄₂ ratio correlate with the levels of A β peptide deposition in the brain when the antibodies are administered to an individual. Thus, measurement of these components in the blood after administration of the antibody provides a simple straightforward diagnostic test for both clinical and preclinical Alzheimer's disease and related neurological disorders.

[0007] There are additional relevant publications concerning the behavior of A β peptide antibodies. For example, PCT publication WO99/27944 published 10 Jun. 1999 describes methods to induce an immune response in order to reduce amyloid deposits. Publication No. WO99/60024 published 25 Nov. 1999, describes methods for amyloid removal using anti-amyloid antibodies. Additional PCT publications, including WO00/72880, WO00/72876 and WO00/77178 all describe various activities of anti-A β peptide antibodies. Antibodies directed to the N-terminus of this peptide are said to reduce plaques in a transgenic murine model; immunization with the amyloid itself is described as are antibodies designed to catalyze hydrolysis of the peptide.

[0008] It has been shown that one pathway for A β metabolism is via transport from CNS to the plasma (Zlokovic, B. V., et al., *Proc. Natl. Acad. Sci (USA)* (1996) 93:4229-4234; Ghersi-Egea, J-F., et al., *J. Neurochem.* (1996) 67:880-883). Additionally, it has been shown that A β in plasma can cross the blood-brain-barrier and enter the brain (Zlokovic, B. V., et al., *Biochem. Biophys. Res. Comm.* (1993) 67:1034-1040). It has also been shown that administration of certain polyclonal and monoclonal A β antibodies decreases A β deposition in amyloid plaques in the App^{V717F} transgenic mouse model of Alzheimer's disease (Bard, F., et al., *Nature Med.* (2000) 6:916-919). This was said to be due to certain anti-A β antibodies crossing the blood-brain-barrier and stimulating phagocytosis of amyloid plaques by microglial cells. In Bard's experiments, assays of brain slices ex vivo showed that the presence of added A β antibody, along with exogenously added microglia, induced phagocytosis of A β , resulting in removal of A β deposits.

[0009] The levels of both soluble A β ₄₀ and A β ₄₂ in CSF and blood can readily be detected using standardized assays using antibodies directed against epitopes along the A β chain. Such assays have been reported, for example, in U.S. Pat. Nos. 5,766,846; 5,837,672; and 5,593,846. These patents describe the production of murine monoclonal antibodies to the central domain of the A β peptide, and these were reported to have epitopes around and including positions 16 and 17. Antibodies directed against the N-terminal region were described as well. Several monoclonal antibodies were asserted to immunoreact with positions 13-28 of the A β peptide; these did not bind to a peptide representing positions 17-28, thus, according to the cited patents, establishing that it is this region, including positions 16-17 (the γ -secretase site) that was the target of these antibodies. Among antibodies known to bind between amino acids 13 and 28 of A β are mouse antibodies 266 (m266), 4G8, and 1C2.

DISCLOSURE OF THE INVENTION

[0010] It has now been found that antibodies which are useful for performing assays for A β peptide, and which are

useful in treatment of conditions associated with amyloid plaques in the brain can elicit a response which results in a marked increase in the level of A β peptide in the blood and this level can be used as a diagnostic marker for clinical and preclinical Alzheimer's disease. These antibodies, which may or may not be humanized, sequester A β peptide from its bound, circulating form in blood and alter clearance of soluble and bound forms of A β in central nervous system and plasma. These antibodies, and fragments thereof, specifically bind to an epitope between amino acids 13 and 28 of the A β molecule. The CDR of these antibodies can be derived from mouse monoclonal antibody 266 (SEQ ID NO:1 through SEQ ID NO:6). Useful antibodies include antibodies and fragments thereof, wherein the variable regions have sequences comprising the CDR from mouse antibody 266 and specific human framework sequences (SEQ ID NO:7 through SEQ ID NO:10), wherein the antibodies retain approximately the binding properties of the mouse antibody and have in vitro and in vivo properties functionally equivalent to the mouse antibody 266. Especially useful are humanized antibodies and fragments thereof, wherein the light chain is SEQ ID NO:11 and the heavy chain is SEQ ID NO:12.

[0011] Thus, in one aspect, the invention is directed to a method to diagnose Alzheimer's disease in a subject at both a clinical and preclinical stage which method comprises administering to said subject an amount of an antibody that sequesters A β peptide from its bound, circulating form in blood, and alters clearance of soluble and bound forms of A β in the central nervous system in plasma, or which specifically binds an epitope contained within positions 13-28 of A β , preferably an antibody having an immunoreactivity equivalent to mouse antibody 266 effective to alter the levels of circulating A β peptides in the blood of said subject when said subject is in a clinical or preclinical stage of Alzheimer's disease followed by measuring the level of A β_{40} , A β_{42} , or the ratio of A β_{40} /A β_{42} in the blood of said subject, wherein an enhanced concentration of A β_{40} , A β_{42} and/or A β_{40} /A β_{42} ratio in said subject identifies said subject as in a preclinical or clinical stage of Alzheimer's disease or cerebral amyloid angiopathy. In other aspects, the invention is directed to kits containing the appropriate materials for conducting the diagnostic method.

BRIEF DESCRIPTION OF THE DRAWINGS

[0012] FIGS. 1A, B and C are graphs showing the levels of A β_{40} (FIG. 1A), A β_{42} (FIG. 1B), and A β_{40} /A β_{42} ratio (FIG. 1C) in plasma of transgenic mice prior to administration of the antibody m266, and the lack of correlation with brain A β deposits.

[0013] FIGS. 2A and B are graphs showing plasma A β_{40} (FIG. 2A) and plasma A β_{40} /A β_{42} ratio (FIG. 2B) in transgenic mice one hour after injection of antibody m266, and the significant correlation with brain A β deposits.

[0014] FIGS. 3A, B and C are graphs showing the significant correlations of the two A β peptides (FIGS. 3A and 3B) and their ratio (FIG. 3C) with A β peptide deposition in the brain 24 hours after injection with monoclonal antibody m266.

[0015] FIGS. 4A, B and C are graphs showing the significant correlations of entry rates into the circulation of the two A β peptides (FIGS. 4A and 4B) and their ratio (FIG. 4C) and A β peptide deposition in transgenic mice.

[0016] FIGS. 5A and B are graphs showing an alternative graphical representation of A β_{40} levels in the plasma 24 hours

(FIG. 5A) and 1 hour (FIG. 5B) after m266 injection correlated with the percentage hippocampus covered by A β deposits.

[0017] FIG. 6 is a table showing Pearson correlation coefficients (Pearson r) and significance (P value) determined between plasma A β values (pre and post injection of m266) and hippocampal A β or amyloid load.

MODES OF CARRYING OUT THE INVENTION

[0018] The A β peptides that circulate in human biological fluids represent a carboxy terminal region of a precursor protein encoded on chromosome 21. It has been reported from the results of in vitro experiments that the A β peptide has poor solubility in physiological solutions, since it contains a stretch of hydrophobic amino acids which are a part of the region that anchors its longer precursor to the lipid membranes of cells. It is thus not surprising that circulating A β peptide is normally complexed with other moieties that prevent it from aggregating. This has resulted in difficulties in detecting circulating A β peptide in biological fluids.

[0019] The above-mentioned patent documents (U.S. Pat. Nos. 5,766,846; 5,837,672 and 5,593,846) describe the preparation of antibodies, including a monoclonal antibody, designated clone 266 (m266), which was raised against, and has been shown to bind specifically to, a peptide comprising amino acids 13-28 of the A β peptide. Applicants have found that after administering m266 to APP^{P717F} mice, a mouse model of Alzheimer's disease, they can measure levels of A β peptides in the circulation that are diagnostic of the levels of amyloid plaques in the brain. Thus, these antibodies are useful not only in conducting assays for circulating A β peptides per se, but also for eliciting circulating blood levels which are diagnostic of the amount of amyloid plaque in the brain, and thus useful in identifying individuals in clinical and preclinical stages of Alzheimer's disease. One such antibody, m266, binds to the mid-region of A β peptide.

[0020] By "monoclonal antibody that bonds to the mid-region of A β peptide" is meant a monoclonal antibody (Mab or Mabs) that binds an amino acid sequence representing an epitope contained between positions 13-28 of A β . The entire region need not be targeted. As long as the antibody binds at least an epitope within this region (especially, e.g., including the α -secretase site 16-17 or the site at which antibody 266 binds), such antibodies are effective in the method of the invention.

[0021] By "antibody" is meant a monoclonal antibody per se, or an immunologically effective fragment thereof, such as an F_{ab}, F_{ab'}, or F_{(ab')₂} fragment thereof. In some contexts, herein, fragments will be mentioned specifically for emphasis; nevertheless, it will be understood that regardless of whether fragments are specified, the term "antibody" includes such fragments as well as single-chain forms. As long as the protein retains the ability specifically to bind its intended target, and in this case, to sequester A β peptide from its carrier proteins in blood, it is included within the term "antibody." Also included within the definition "antibody" for example, are single chain forms, generally designated F_v, regions, of antibodies with this specificity. Preferably, but not necessarily, the antibodies useful in the invention are produced recombinantly, as manipulation of the typically murine or other non-human antibodies with the appropriate specificity is required in order to convert them to humanized form. Antibodies may or may not be glycosylated, though glycosylated antibodies are preferred. Antibodies are properly cross-linked via disulfide bonds, as is well-known.

1 5 10 15
Arg Ser Ser Gln Ser Leu Ile Tyr Ser Asp Gly Asn Ala Tyr Leu His

amino acid sequence, in which the framework originated from human germline VH segments DP53 and J segment JH4, with several amino acid substitutions to the consensus amino acids in the same human subgroup to reduce potential immunogenicity:

(SEQ ID NO: 8)

1	5	10	15
Xaa Val Gln Leu Val Glu Xaa Gly Gly Gly Leu Val Gln Pro Gly			
	20	25	30
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser			
	35	40	45
Arg Tyr Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu			
	50	55	60
Xaa Leu Val Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr			
	65	70	75
Pro Asp Xaa Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Xaa			
	80	85	90
Xaa Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Xaa Asp			
	95	100	105
Thr Ala Val Tyr Tyr Cys Ala Ser Gly Asp Tyr Trp Gly Gln Gly			
	110		
Thr Xaa Val Thr Val Ser Ser			

wherein:

Xaa at position 1 is Glu or Gln;

Xaa at position 7 is Ser or Leu;

Xaa at position 46 is Glu, Val, Asp, or Ser;

Xaa at position 63 is Thr or Ser;

Xaa at position 75 is Ala, Ser, Val, or Thr;

Xaa at position 76 is Lys or Arg;

Xaa at position 89 is Glu or Asp; and

Xaa at position 107 is Leu or Thr.

[0033] A particularly preferred light chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline Vk segments DPK18 and J segment Jk1, with several amino acid substitutions to the consensus amino acids in the same human V subgroup to reduce potential immunogenicity:

(SEQ ID NO: 9)

1	5	10	15
Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu			
	20	25	30
Gly Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Ile			
	35	40	45
Tyr Ser Asp Gly Asn Ala Tyr Leu His Trp Phe Leu Gln Lys Pro			
	50	55	60
Gly Gln Ser Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe			
	65	70	75
Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp			
	80	85	90
Phe Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Val Gly Val			
	95	100	105
Tyr Tyr Cys Ser Gln Ser Thr His Val Pro Trp Thr Phe Gly Gln			
	110		
Gly Thr Lys Val Glu Ile Lys Arg.			

[0034] A particularly preferred heavy chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline VH segments DP53 and J segment JH4:

(SEQ ID NO: 10)

1	5	10	15
Glu Val Gln Leu	Val Glu Ser Gly Gly	Gly Leu Val Gln	Pro Gly
20	25	30	
Gly Ser Leu Arg	Leu Ser Cys Ala Ala	Ser Gly Phe Thr	Phe Ser
35	40	45	
Arg Tyr Ser Met	Ser Trp Val Arg Gln	Ala Pro Gly Lys	Gly Leu
50	55	60	
Glu Leu Val Ala	Gln Ile Asn Ser Val	Gly Asn Ser Thr	Tyr Tyr
65	70	75	
Pro Asp Thr Val	Lys Gly Arg Phe Thr	Ile Ser Arg Asp	Asn Ala
80	85	90	
Lys Asn Thr Leu	Tyr Ser Leu Gln Met	Asn Ser Leu Arg	Ala Glu Asp
95	100	105	
Thr Ala Val Tyr	Tyr Cys Ala Ser Gly	Asp Tyr Trp Gly	Gln Gly
110			
Thr Leu Val Thr	Val Ser Ser.		

[0035] A preferred light chain for a humanized antibody of the present invention has the amino acid sequence:

(SEQ ID NO: 11)

1	5	10	15
Asp Val Val Met	Thr Gln Ser Pro Leu	Ser Leu Pro Val	Thr Leu
20	25	30	
Gly Gln Pro Ala	Ser Ile Ser Cys Arg	Ser Ser Gln Ser	Leu Ile
35	40	45	
Tyr Ser Asp Gly	Asn Ala Tyr Leu His	Trp Phe Leu Gln	Lys Pro
50	55	60	
Gly Gln Ser Pro	Arg Leu Leu Ile Tyr	Lys Val Ser Asn	Arg Phe
65	70	75	
Ser Gly Val Pro	Asp Arg Phe Ser Gly	Ser Gly Ser Gly	Thr Asp
80	85	90	
Phe Thr Leu Lys	Ile Ser Arg Val Glu	Ala Glu Asp Val	Gly Val
95	100	105	
Tyr Tyr Cys Ser	Gln Ser Thr His Val	Pro Trp Thr Phe	Gly Gln
110	115	120	
Gly Thr Lys Val	Glu Ile Lys Arg Thr	Val Ala Ala Pro	Ser Val
125	130	135	
Phe Ile Phe Pro	Pro Ser Asp Glu Gln	Leu Lys Ser Gly	Thr Ala
140	145	150	
Ser Val Val Cys	Leu Leu Asn Asn Phe	Tyr Pro Arg Glu	Ala Lys
155	160	165	
Val Gln Trp Lys	Val Asp Asn Ala Leu	Gln Ser Gly Asn	Ser Gln
170	175	180	
Glu Ser Val Thr	Glu Gln Asp Ser Lys	Asp Ser Thr Tyr	Ser Leu
185	190	195	

-continued

Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys
 200 205 210
 Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val
 215
 Thr Lys Ser Phe Asn Arg Gly Glu Cys.

[0036] A preferred heavy chain for a humanized antibody
 of the present invention has the amino acid sequence:

(SEQ ID NO: 12)

1 5 10 15
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 20 25 30
 Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
 35 40 45
 Arg Tyr Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Leu Val Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr
 65 70 75
 Pro Asp Thr Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala
 80 85 90
 Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 95 100 105
 Thr Ala Val Tyr Tyr Cys Ala Ser Gly Asp Tyr Trp Gly Gln Gly
 110 115 120
 Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 125 130 135
 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala
 140 145 150
 Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
 155 160 165
 Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe
 170 175 180
 Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
 185 190 195
 Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys
 200 205 210
 Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val
 215 220 225
 Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
 230 235 240
 Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 245 250 255
 Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr
 260 265 270
 Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe
 275 280 285
 Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys
 290 295 300
 Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val
 305 310 315

-continued

```

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
      320              325              330
Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr
      335              340              345
Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
      350              355              360
Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu
      365              370              375
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
      380              385              390
Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
      395              400              405
Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu
      410              415              420
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
      425              430              435
Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
      440
Leu Ser Leu Ser Pro Gly Lys.

```

[0037] Other sequences are possible for the light and heavy chains for the humanized antibodies of the present invention and for humanized 266. The immunoglobulins can have two pairs of light chain/heavy chain complexes, at least one chain comprising one or more mouse complementarity determining regions functionally joined to human framework region segments.

[0038] Starting at position 56 of the heavy chain variable region, both m266 and humanized 266 contain the sequence Asn-Ser-Thr. This sequence is an example of the Asn-X-Ser/Thr signal for N-linked glycosylation, wherein the Asn is the site of attachment of N-linked glycosyl chains. Both m266 and humanized 266 are extensively glycosylated at this site. Quite unpredictably and advantageously, the affinity of humanized 266 that is deglycosylated in the heavy chain CDR2 for Aβ peptide is markedly higher than that of humanized 266. The heavy chain CDR2 of deglycosylated humanized 266 has the following amino acid sequences:

heavy chain CDR2:

```

      1              5              10              15
Gln Ile Asn Ser Val Gly Xaa Xaa Xaa Tyr Tyr Pro Asp Thr Val Lys Gly
      (SEQ ID NO: 13)

```

[0039] wherein:

[0040] Xaa at position 7 is any amino acid, provided that if Xaa at position 8 is neither Asp nor Pro and Xaa at position 9 is Ser or Thr, then Xaa at position 7 is not Asn;

[0041] Xaa at position 8 is any amino acid, provided that if Xaa at position 7 is Asn and Xaa at position 9 is Ser or Thr, then Xaa at position 8 is Asp or Pro; and

[0042] Xaa at position 9 is any amino acid, provided that if Xaa at position 7 is Asn and Xaa at position 8 is neither Asp nor Pro, then Xaa at position 9 is neither Ser nor Thr;

[0043] By "any amino acid" is meant any naturally-occurring amino acid. Preferred naturally-occurring amino acids are Ala, Cys, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr.

[0044] A preferred deglycosylated humanized antibody is a humanized form of m266, wherein the deglycosylated heavy chain CDR2 is SEQ ID NO:13, wherein:

[0045] Xaa at position 7 of SEQ ID NO:13 is selected from the group consisting of Ala, Cys, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr, provided that if Xaa at position 8 is neither Asp nor Pro and Xaa at position 9 is Ser or Thr, then Xaa at position 7 is not Asn;

[0046] Xaa at position 8 of SEQ ID NO:13 is selected from the group consisting of Ala, Cys, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr, provided that if Xaa at position 7 is Asn and Xaa at position 9 is Ser or Thr, then Xaa at position 8 is Asp or Pro; and

[0047] Xaa at position 9 of SEQ ID NO:13 is selected from the group consisting of Ala, Cys, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr,

provided that if Xaa at position 7 is Asn and Xaa at position 8 is neither Asp nor Pro, then Xaa at position 9 is neither Ser nor Thr.

[0048] A preferred heavy chain variable region of a deglycosylated humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline VH segment DP53 and J segment JH4, with several amino acid substitutions to the consensus amino acids in the same human subgroup to reduce potential immunogenicity and wherein the N-glycosylation site in heavy chain CDR2 is modified so that it cannot be N-glycosylated:

(SEQ ID NO: 14)

1	5	10	15
Xaa Val Gln Leu Val Glu Xaa Gly Gly Leu Val Gln Pro Gly			
20	25	30	
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser			
35	40	45	
Arg Tyr Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu			
50	55	60	
Xaa Leu Val Ala Gln Ile Asn Ser Val Gly Xaa Xaa Xaa Tyr Tyr			
65	70	75	
Pro Asp Xaa Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Xaa			
80	85	90	
Xaa Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Xaa Asp			
95	100	105	
Thr Ala Val Tyr Tyr Cys Ala Ser Gly Asp Tyr Trp Gly Gln Gly			
110			
Thr Xaa Val Thr Val Ser Ser			

[0049] wherein:

[0050] Xaa at position 1 is Glu or Gln;

[0051] Xaa at position 7 is Ser or Leu;

[0052] Xaa at position 46 is Glu, Val, Asp, or Ser;

[0053] Xaa at position 56 is any amino acid, provided that if Xaa at position 57 is neither Asp nor Pro and Xaa at position 59 is Ser or Thr, then Xaa at position 56 is not Asn;

[0061] A particularly preferred heavy chain variable region of a deglycosylated humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline VH segment DP53 and J segment JH4 and wherein the N-glycosylation site in heavy chain CDR2 is modified so that it cannot be N-glycosylated:

(SEQ ID NO: 15)

1	5	10	15
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly			
20	25	30	
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser			
35	40	45	
Arg Tyr Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu			
50	55	60	
Glu Leu Val Ala Gln Ile Asn Ser Val Gly Xaa Xaa Xaa Tyr Tyr			
65	70	75	
Pro Asp Thr Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala			
80	85	90	
Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp			
95	100	105	
Thr Ala Val Tyr Tyr Cys Ala Ser Gly Asp Tyr Trp Gly Gln Gly			
110			
Thr Leu Val Thr Val Ser Ser.			

[0054] Xaa at position 57 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 58 is Ser or Thr, then Xaa at position 57 is Asp or Pro; and

[0055] Xaa at position 58 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 57 is neither Asp nor Pro, then Xaa at position 58 is neither Ser nor Thr

[0056] Xaa at position 63 is Thr or Ser;

[0057] Xaa at position 75 is Ala, Ser, Val, or Thr;

[0058] Xaa at position 76 is Lys or Arg;

[0059] Xaa at position 89 is Glu or Asp; and

[0060] Xaa at position 107 is Leu or Thr.

[0062] wherein:

[0063] Xaa at position 56 is any amino acid, provided that if Xaa at position 57 is neither Asp nor Pro and Xaa at position 59 is Ser or Thr, then Xaa at position 56 is not Asn;

[0064] Xaa at position 57 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 58 is Ser or Thr, then Xaa at position 57 is Asp or Pro; and

[0065] Xaa at position 58 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 57 is neither Asp nor Pro, then Xaa at position 58 is neither Ser nor Thr.

[0066] A preferred heavy chain for a deglycosylated humanized antibody of the present invention, wherein the

N-glycosylation site in heavy chain CDR2 is modified so that it cannot be N-glycosylated, has the amino acid sequence:

(SEQ ID NO: 16)

1	5	10	15
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly			
20	25	30	
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser			
35	40	45	
Arg Tyr Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu			
50	55	60	
Glu Leu Val Ala Gln Ile Asn Ser Val Gly Xaa Xaa Xaa Tyr Tyr			
65	70	75	
Pro Asp Thr Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala			
80	85	90	
Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp			
95	100	105	
Thr Ala Val Tyr Tyr Cys Ala Ser Gly Asp Tyr Trp Gly Gln Gly			
110	115	120	
Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val			
125	130	135	
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala			
140	145	150	
Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr			
155	160	165	
Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe			
170	175	180	
Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val			
185	190	195	
Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys			
200	205	210	
Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val			
215	220	225	
Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro			
230	235	240	
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro			
245	250	255	
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr			
260	265	270	
Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe			
275	280	285	
Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys			
290	295	300	
Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val			
305	310	315	
Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys			
320	325	330	
Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr			
335	340	345	
Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr			
350	355	360	
Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu			

-continued

365	370	375
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu		
380	385	390
Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro		
395	400	405
Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu		
410	415	420
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys		
425	430	435
Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser		
440		
Leu Ser Leu Ser Pro Gly Lys		

[0067] wherein:

[0068] Xaa at position 56 is any amino acid, provided that if Xaa at position 57 is neither Asp nor Pro and Xaa at position 59 is Ser or Thr, then Xaa at position 56 is not Asn;

[0069] Xaa at position 57 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 58 is Ser or Thr, then Xaa at position 57 is Asp or Pro; and

[0070] Xaa at position 58 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 57 is neither Asp nor Pro, then Xaa at position 58 is neither Ser nor Thr.

[0071] Preferred deglycosylated 266 antibodies having the heavy variable region according to SEQ ID NO:14, SEQ ID NO:15, and SEQ ID NO:16 are those wherein:

[0072] Xaa at position 56 is selected from the group consisting of Ala, Gly, His, Asn, Gln, Ser, and Thr, provided that if Xaa at position 58 is Ser or Thr, then Xaa at position 56 is not Asn;

[0073] Xaa at position 57 is selected from the group consisting of Ala, Gly, His, Asn, Gln, Ser, and Thr; and

[0074] Xaa at position 58 is selected from the group consisting of Ala, Gly, His, Asn, Gln, Ser, and Thr, provided that if Xaa at position 56 is Asn, then Xaa at position 58 is neither Ser nor Thr.

[0075] Preferred sequences for CDR2 (positions 56, 57, and 58) of the heavy chain SEQ ID NO:14, SEQ ID NO:15, and SEQ ID NO:16 include those in which only a single amino acid is changed, those in which only two amino acids are changed, or all three are changed. It is preferred to replace Asn at position 56. It is preferred to replace Thr at position 58 with an amino acid other than Ser. It is preferred to not destroy the N-glycosylation site in the CDR2 of the 266 heavy chain by replacing Ser at position 57 with Pro or Asp. Conservative substitutions at one, two, or all three positions are preferred. The most preferred species are those in which Asn at position 56 is replaced with Ser or Thr. Particularly preferred antibodies are those in which Ser or Thr is at position 56, Ser is at position 57, and Thr is at position 58 of SEQ ID NO:14, SEQ ID NO:15, or SEQ ID NO:16.

[0076] Especially preferred deglycosylated species are antibodies comprising a light chain of SEQ ID NO:11 and a heavy chain of SEQ ID NO:16, wherein in SEQ ID NO:16, Xaa at position 56 is Ser, Xaa at position 57 is Ser, and Xaa at position 58 is Thr ("N56S"), or wherein in SEQ ID NO:16, Xaa at position 56 is Thr, Xaa at position 57 is Ser, and Xaa at position 58 is Thr ("N56T").

[0077] Production of the antibodies useful in the invention typically involves recombinant techniques, as is described in PCT/US01/06191 cited above and incorporated herein by reference.

[0078] The antibodies (including immunologically reactive fragments) are administered to a subject to be evaluated for conditions associated with A β deposits such as clinical or preclinical Alzheimer's disease, or clinical or preclinical amyloid angiopathy, using standard administration techniques, preferably peripherally (i.e. not by administration into the central nervous system) by intravenous, intraperitoneal, subcutaneous, pulmonary, transdermal, intramuscular, intranasal, buccal, sublingual, or suppository administration.

[0079] The compositions for administration are designed to be appropriate for the selected mode of administration, and pharmaceutically acceptable excipients such as dispersing agents, buffers, surfactants, preservatives, solubilizing agents, isotonicity agents, stabilizing agents and the like are used as appropriate. Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton Pa., latest edition, incorporated herein by reference, provides a compendium of formulation techniques as are generally known to practitioners. It may be particularly useful to alter the solubility characteristics of the antibodies of the invention, making them more lipophilic, for example, by encapsulating them in liposomes or by blocking polar groups.

[0080] Peripheral systemic delivery by intravenous or intraperitoneal or subcutaneous injection is preferred. Suitable vehicles for such injections are straightforward. In addition, however, administration may also be effected through the mucosal membranes by means of nasal aerosols or suppositories. Suitable formulations for such modes of administration are well known and typically include surfactants that facilitate cross-membrane transfer. Such surfactants are often derived from steroids or are cationic lipids, such as N-[1-(2,3-dioleoyl)propyl]-N,N,N-trimethyl ammonium chloride (DOTMA) or various compounds such as cholesterol hemisuccinate, phosphatidyl glycerols and the like.

[0081] The concentration of the humanized antibody in formulations from as low as about 0.1% to as much as 15 or 20% by weight and will be selected primarily based on fluid volumes, viscosities, and so forth, in accordance with the particular mode of administration selected. Thus, a typical composition for injection could be made up to contain 1 mL sterile buffered water of phosphate buffered saline and 1-1000 mg, preferably 10-100 mg, of the humanized antibody of the present invention. The formulation could be sterile

filtered after making the formulation, or otherwise made microbiologically acceptable. A typical composition for intravenous infusion could have volumes between 1-250 mL of fluid, such as sterile Ringer's solution, and 1-100 mg per mL, or more in antibody concentration. Therapeutic agents of the invention can be frozen or lyophilized for storage and reconstituted in a suitable sterile carrier prior to use. Lyophilization and reconstitution can lead to varying degrees of antibody activity loss (e.g. with conventional immune globulins, IgM antibodies tend to have greater activity loss than IgG antibodies). Dosages may have to be adjusted to compensate. The pH of the formulation will be selected to balance antibody stability (chemical and physical) and comfort to the patient when administered. Generally, pH between 4 and 8 is tolerated.

[0082] Although the foregoing methods appear the most convenient and most appropriate for administration of proteins such as humanized antibodies, by suitable adaptation, other techniques for administration, such as transdermal administration and oral administration may be employed provided proper formulation is designed.

[0083] In addition, it may be desirable to employ controlled release formulations using biodegradable films and matrices, or osmotic mini-pumps, or delivery systems based on dextran beads, alginate, or collagen.

[0084] In summary, formulations are available for administering the antibodies of the invention and are well-known in the art and may be chosen from a variety of options.

[0085] Typical dosage levels can be optimized using standard clinical techniques and will be dependent on the mode of administration.

[0086] After administration of the antibody to the subject, blood samples are withdrawn at periodic intervals over the succeeding minutes, hours, or days. Suitable time periods may be as short as a few minutes, 10 minutes, 30 minutes, or 1 hour, several hours, or days may be allowed to elapse before withdrawal of the blood sample. Measurement after less than 3 hours is preferred. If desired, the plasma fraction can be obtained for ease of analysis. Standard analytic techniques for analysis of the $A\beta_{40}$, $A\beta_{42}$ and the ratio thereof are used. These techniques are described, for example, in U.S. Pat. No. 5,766,846. Any suitable technique for analysis, however, can be employed, such as chromatographic separation, Western blotting, ELISA assays, homogenous assays and the like.

[0087] The concentration of the $A\beta_{40}$, $A\beta_{42}$, or their ratio is then compared to these values in a control. Typical controls include individuals known to be free of conditions associated with the amyloid plaques, such as teenagers or very young adults and in addition, age-matched cognitively normal controls are obtained by averaging values from the general population. While some elderly age-matched cognitively normal controls have pre-clinical AD, most do not. Thus, the average values from such a population will be useful and critical to obtain. Design of standard controls is a process that is well known to the ordinary practitioner. Individuals who have elevated levels of the stated peptides or of the ratio of $A\beta_{40}$ to $A\beta_{42}$ as compared to the control values are then identified as having a high likelihood of clinical or preclinical conditions associated with the formation of amyloid plaques.

[0088] It may be desirable to package the components for carrying out the assay of the invention into convenient kits. Such kits will include containers such as bottles or vials which contain samples of the antibody to be administered as well as the appropriate reagents for carrying out the assay on

the withdrawn blood sample. The kit will also contain instructions for conducting the assay and, optionally, charts of control values.

[0089] The following examples are intended to illustrate but not to limit the invention.

[0090] The examples hereinbelow employ, among others, a murine monoclonal antibody designated "266" which was originally prepared by immunization with a peptide comprised of residues 13-28 of human $A\beta$ peptide. The antibody was confirmed to immunoreact with this peptide, but had previously been reported to not react with the peptide containing only residues 17-28 of human $A\beta$ peptide, or at any other epitopes within the $A\beta$ peptide. The preparation of this antibody is described in U.S. Pat. No. 5,766,846, incorporated herein by reference. As the examples here describe experiments conducted in murine systems, the use of murine monoclonal antibodies is satisfactory. However, in the treatment methods of the invention intended for human use, humanized forms of the antibodies with the immunospecificity corresponding to that of antibody 266 are preferred.

Example 1

Correlation of Circulating Peptide Levels with Plaques

[0091] A murine model for Alzheimer's disease, APP V717F transgenic mice, was used in this assay. These mice are described by Games, D., et al., *Nature* (1995) 373:523-527; Bales, K. R., et al., *Nature Genet.* (1997) 17:263-264; and by Holtzman, D. M., et al., *Proc. Natl. Acad. Sci. U.S.A.* (2000) 97:2892-2897. In this model, a mutant form of the human APP gene is expressed and results in an early onset form of familial Alzheimer's disease. Although the brains of these mice appear normal initially, $A\beta$ deposition in the form of diffuse and neuritic plaques occurs at 6-15 months, although mice homozygous for the transgene show variability in that at 9-14 months of age, some mice develop $A\beta$ deposits while others do not.

[0092] 53 homozygous mice at 12 months were used in this study.

[0093] Plasma levels of $A\beta_{40}$, $A\beta_{42}$, and $A\beta_{40}/A\beta_{42}$ ratios were measured by ELISA in the plasma of these mice prior to administration of 500 μ g of m266 and at various time intervals up to 24 hours after administering this antibody. After 24 hours, the mice were sacrificed, and the amount of $A\beta$ deposition in the brain was assessed in the hippocampus and cortex as described by DeMattos, et al. *Proc. Nat'l. Acad. Sci. USA* (2001) 98:8850-8855, and evaluated as a percentage of brain covered by $A\beta$ deposits.

[0094] As shown in FIGS. 1A, B and C, if the percentage $A\beta$ coverage due to deposition in the hippocampus is plotted on the x-axis against the levels of the peptides and their ratio in plasma on the y-axis prior to administration of the antibody, no correlation is found. Regardless of whether the percent $A\beta$ deposition was essentially zero (0) or over 75%, the average level of $A\beta_{40}$ was approximately 250 (pg/ml) and of $A\beta_{42}$ approximately 400 (pg/ml). The ratio of $A\beta_{40}$ to $A\beta_{42}$ was thus approximately 0.5-0.6.

[0095] As shown in FIGS. 2 A and B, however, the plasma level of $A\beta_{40}$ strongly correlated with the percentage of $A\beta$ deposition in hippocampus one hour after m266 injection, as did the ratio of $A\beta_{40}$ to $A\beta_{42}$.

[0096] FIGS. 3 A, B and C show similar results obtained 24 hours post injection. The levels obtained of $A\beta_{40}$ and the

$A\beta_{40}/A\beta_{42}$ ratio strongly correlated with the % $A\beta$ deposition in hippocampus. The $A\beta_{42}$ levels also correlated with % $A\beta$ deposition but not as well as $A\beta_{40}$ levels.

[0097] FIGS. 4 A, B and C show analogous results with respect to entry rate of the two $A\beta$ peptides into the plasma and the calculated values for the entry rate as a function of the ratio of these peptides. The best correlations with $A\beta$ deposition were rate of $A\beta_{40}$ entry and the ratio of $A\beta_{40}/A\beta_{42}$.

[0098] FIGS. 5 A and B show an alternate presentation of the data for plasma levels of $A\beta_{40}$ 24 hours and 1 hour after m266 injection. When the mice were grouped according to low, medium, or high $A\beta$ coverage in the hippocampus, the animals with low $A\beta$ deposition could be completely distinguished from those with high deposition as a function of the level of plasma $A\beta_{40}$.

Example 2

[0099] In a study similar to that set forth in Example 1, a cohort of 49 homozygous APP V717F mice were used. Before and after injection of 500 μ g IV of m266, plasma samples were obtained at 5 minutes, 1 hour, 3 hours, 6 hours and 24 hours and levels of $A\beta_{40}$ and $A\beta_{42}$ were assessed as described in Example 1. The mice were sacrificed after 24 hours and 1 hemisphere was assessed for the percentage of the area of the hippocampus or cingulate cortex occupied by $A\beta$

peptide (using quantitative $A\beta$ immunofluorescence staining) and the area occupied by amyloid (by thioflavine-S (amyloid) staining). The regions from the other hemisphere were assessed for $A\beta$ peptide by ELISA.

[0100] The Pearson correlation coefficient (Pearson r) and significance (P value) were determined between plasma $A\beta$ values (pre and post injection of m266) and hippocampal $A\beta$ or amyloid load using GraphPad Prism software (version 3.00 for Windows, San Diego, USA). $A\beta$ load is defined as the percentage area of the hippocampus covered by $A\beta$ -immunoreactive deposits. Amyloid load is defined as the percentage area of the hippocampus covered by thioflavine-S positive deposits. Correlations were also determined between the plasma $A\beta$ accumulation over 24 hours (area under curve, AUC) and hippocampal $A\beta$ load or amyloid load.

[0101] FIG. 6 shown the results obtained. Briefly, it was found that the base line levels (prior to injection) of $A\beta_{40}$, $A\beta_{42}$ and the calculated $A\beta_{40}/A\beta_{42}$ ratio prior to injection with m266 did not correlate with percentage $A\beta$ or amyloid deposition. However, following administration of m266, there were significant correlations between plasma $A\beta_{40}$, $A\beta_{42}$, and $A\beta_{40}/A\beta_{42}$ ratio with both $A\beta$ and amyloid burden in the hippocampus and cingulate cortex.

[0102] Statistical analysis of the results permits accurate prediction of hippocampal $A\beta$ load in these mice based on plasma $A\beta_{40}$ levels 24 hours following m266 injection.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Mus sp.
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(16)
 <223> OTHER INFORMATION: Light Chain CDR1

<400> SEQUENCE: 1

Arg Ser Ser Gln Ser Leu Ile Tyr Ser Asp Gly Asn Ala Tyr Leu His
 1 5 10 15

<210> SEQ ID NO 2
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: mus sp.
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(7)
 <223> OTHER INFORMATION: Light Chain CDR2

<400> SEQUENCE: 2

Lys Val Ser Asn Arg Phe Ser
 1 5

<210> SEQ ID NO 3
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: mus sp.
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(9)
 <223> OTHER INFORMATION: Light Chain CDR3

-continued

<400> SEQUENCE: 3

Ser Gln Ser Thr His Val Pro Trp Thr
1 5

<210> SEQ ID NO 4
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: mus sp.
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(5)
<223> OTHER INFORMATION: Heavy Chain CDR1

<400> SEQUENCE: 4

Arg Tyr Ser Met Ser
1 5

<210> SEQ ID NO 5
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: mus sp.
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Heavy Chain CDR2

<400> SEQUENCE: 5

Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Thr Val Lys
1 5 10 15

Gly

<210> SEQ ID NO 6
<211> LENGTH: 3
<212> TYPE: PRT
<213> ORGANISM: mus sp.
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(3)
<223> OTHER INFORMATION: Heavy Chain CDR3

<400> SEQUENCE: 6

Gly Asp Tyr
1

<210> SEQ ID NO 7
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(113)
<223> OTHER INFORMATION: Humanized Antibody Light Chain Variable Region
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa at position 2 is Val or Ile
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa at position 7 is Ser or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Xaa at position 14 is Thr or Ser

-continued

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: Xaa at position 15 is Leu or Pro
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (30)..(30)
<223> OTHER INFORMATION: Xaa at position 30 is Ile or Val
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(50)
<223> OTHER INFORMATION: Xaa at position 50 is Arg, Gln, or Lys
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (88)..(88)
<223> OTHER INFORMATION: Xaa at position 88 is Val or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (105)..(105)
<223> OTHER INFORMATION: Xaa at position 105 is Gln or Gly
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (108)..(108)
<223> OTHER INFORMATION: Xaa at position 108 is Lys or Arg
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (109)..(109)
<223> OTHER INFORMATION: Xaa at position 109 is Val or Leu

<400> SEQUENCE: 7

Asp Xaa Val Met Thr Gln Xaa Pro Leu Ser Leu Pro Val Xaa Xaa Gly
1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Xaa Tyr Ser
20 25 30

Asp Gly Asn Ala Tyr Leu His Trp Phe Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Xaa Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Xaa Gly Val Tyr Tyr Cys Ser Gln Ser
85 90 95

Thr His Val Pro Trp Thr Phe Gly Xaa Gly Thr Xaa Xaa Glu Ile Lys
100 105 110

Arg

<210> SEQ ID NO 8
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Humanized antibody
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(112)
<223> OTHER INFORMATION: Humanized Antibody Heavy Chain Variable Region
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa at position 1 is Glu or Gln
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa at position 7 is Ser or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (46)..(46)

-continued

```

<223> OTHER INFORMATION: Xaa at position 46 is Glu, Val, Asp, or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (63)..(63)
<223> OTHER INFORMATION: Xaa at position 63 is Thr or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (75)..(75)
<223> OTHER INFORMATION: Xaa at position 75 is Ala, Ser, Val, or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (76)..(76)
<223> OTHER INFORMATION: Xaa at position 76 is Lys or Arg
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (89)..(89)
<223> OTHER INFORMATION: Xaa at position 89 is Glu or Asp
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (107)..(107)
<223> OTHER INFORMATION: Xaa at position 107 is Leu or Thr

<400> SEQUENCE: 8

Xaa Val Gln Leu Val Glu Xaa Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr
20          25          30

Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Xaa Leu Val
35          40          45

Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Xaa Val
50          55          60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Xaa Xaa Asn Thr Leu Tyr
65          70          75          80

Leu Gln Met Asn Ser Leu Arg Ala Xaa Asp Thr Ala Val Tyr Tyr Cys
85          90          95

Ala Ser Gly Asp Tyr Trp Gly Gln Gly Thr Xaa Val Thr Val Ser Ser
100         105         110

<210> SEQ ID NO 9
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Humanized antibody
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(113)
<223> OTHER INFORMATION: Humanized Antibody Light Chain Variable Region

<400> SEQUENCE: 9

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly
1          5          10          15

Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Ile Tyr Ser
20          25          30

Asp Gly Asn Ala Tyr Leu His Trp Phe Leu Gln Lys Pro Gly Gln Ser
35          40          45

Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50          55          60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65          70          75          80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser
85          90          95

```

-continued

Thr His Val Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105 110

Arg

<210> SEQ ID NO 10
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Humanized Antibody
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(112)
 <223> OTHER INFORMATION: Humanized Antibody Heavy Chain Variable Region
 <400> SEQUENCE: 10

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr
 20 25 30
 Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val
 35 40 45
 Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Thr Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ser Gly Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 100 105 110

<210> SEQ ID NO 11
 <211> LENGTH: 219
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Humanized antibody
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(219)
 <223> OTHER INFORMATION: Humanized Antibody Light Chain
 <400> SEQUENCE: 11

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly
 1 5 10 15
 Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Ile Tyr Ser
 20 25 30
 Asp Gly Asn Ala Tyr Leu His Trp Phe Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser
 85 90 95
 Thr His Val Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105 110

-continued

```

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
   115                               120               125

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
   130                               135               140

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
   145                               150               155               160

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
   165                               170               175

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
   180                               185               190

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
   195                               200               205

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
   210                               215

```

```

<210> SEQ ID NO 12
<211> LENGTH: 442
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Humanized antibody
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(442)
<223> OTHER INFORMATION: Humanized Antibody Heavy Chain

```

```

<400> SEQUENCE: 12

```

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1      5      10      15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr
20     25     30

Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val
35     40     45

Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Thr Val
50     55     60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
65     70     75     80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85     90     95

Ala Ser Gly Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
100    105    110

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
115    120    125

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
130    135    140

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
145    150    155    160

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
165    170    175

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
180    185    190

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
195    200    205

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
210    215    220

```

-continued

```

Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
225                230                235                240

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
                245                250                255

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
                260                265                270

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
                275                280                285

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
290                295                300

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
305                310                315                320

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
                325                330                335

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu
                340                345                350

Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
                355                360                365

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
                370                375                380

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
385                390                395                400

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
                405                410                415

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
                420                425                430

Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
                435                440

```

```

<210> SEQ ID NO 13
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: mus variant
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Heavy Chain CDR2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa at position 7 is any amino acid, provided
that Xaa at position 8 is neither Asp nor Pro and Xaa at position
9 is Ser or Thr, then Xaa at position 7 is not Asn
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa at position 8 is any amino acid, provided
that Xaa at position 7 is Asn and Xaa at position 9 is Ser or
Thr, then Xaa at position 8 is Asp or Pro
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa at position 9 is any amino acid, provided
that Xaa at position 7 is Asn and Xaa at position 8 is neither
Asp nor Pro, then Xaa at position 9 is neither Ser nor Thr

<400> SEQUENCE: 13

```

```

Gln Ile Asn Ser Val Gly Xaa Xaa Xaa Tyr Tyr Pro Asp Thr Val Lys
1           5           10           15

```

-continued

<210> SEQ ID NO 14
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(112)
<223> OTHER INFORMATION: Deglycosylated Humanized Antibody Heavy Chain Variable Region
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa at position 1 is Glu or Gln
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa at position 7 is Ser or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (46)..(46)
<223> OTHER INFORMATION: Xaa at position 46 is Glu, Val, Asp, or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (56)..(56)
<223> OTHER INFORMATION: Xaa at position 56 is any amino acid, provided that if Xaa at position 57 is neither Asp nor Pro and Xaa at position 58 is Ser or Thr, then Xaa at position 56 is not Asn
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (57)..(57)
<223> OTHER INFORMATION: Xaa at position 57 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 58 is Ser or Thr, then Xaa at position 57 is Asp or Pro
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (58)..(58)
<223> OTHER INFORMATION: Xaa at position 58 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 57 is neither Asp nor Pro, then Xaa at position 58 is neither Ser nor Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (63)..(63)
<223> OTHER INFORMATION: Xaa at position 63 is Thr or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (75)..(75)
<223> OTHER INFORMATION: Xaa at position 75 is Ala, Ser, Val, or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (76)..(76)
<223> OTHER INFORMATION: Xaa at position 76 is Lys or Arg
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (89)..(89)
<223> OTHER INFORMATION: Xaa at position 89 is Glu or Asp
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (107)..(107)
<223> OTHER INFORMATION: Xaa at position 107 is Leu or Thr

<400> SEQUENCE: 14

Xaa Val Gln Leu Val Glu Xaa Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr
20 25 30

Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Xaa Leu Val
35 40 45

Ala Gln Ile Asn Ser Val Gly Xaa Xaa Xaa Tyr Tyr Pro Asp Xaa Val

-continued

50	55	60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Xaa Xaa Asn Thr Leu Tyr		
65	70	75 80
Leu Gln Met Asn Ser Leu Arg Ala Xaa Asp Thr Ala Val Tyr Tyr Cys		
	85	90 95
Ala Ser Gly Asp Tyr Trp Gly Gln Gly Thr Xaa Val Thr Val Ser Ser		
	100	105 110

<210> SEQ ID NO 15
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(112)
 <223> OTHER INFORMATION: Deglycosylated Humanized Antibody Heavy Chain Variable Region
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (56)..(56)
 <223> OTHER INFORMATION: Xaa at position 56 is any amino acid, provided that if Xaa at position 57 is neither Asp nor Pro and Xaa at position 58 is Ser or Thr, then Xaa at position 56 is not Asn
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (57)..(57)
 <223> OTHER INFORMATION: Xaa at position 57 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 58 is Ser or Thr, then Xaa at position 57 is Asp or Pro
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (58)..(58)
 <223> OTHER INFORMATION: Xaa at position 58 is any amino acid, provided that if Xaa at position 56 is Asn and Xaa at position 57 is neither Asp nor Pro, then Xaa at position 58 is neither Ser nor Thr
 <400> SEQUENCE: 15

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr
20 25 30
Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val
35 40 45
Ala Gln Ile Asn Ser Val Gly Xaa Xaa Xaa Tyr Tyr Pro Asp Thr Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ser Gly Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
100 105 110

<210> SEQ ID NO 16
 <211> LENGTH: 442
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Humanized antibody
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(442)

-continued

```

<223> OTHER INFORMATION: Humanized Antibody Heavy Chain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (56)..(56)
<223> OTHER INFORMATION: Xaa at position 56 is any amino acid, provided
      that Xaa at position 57 is neither Asp nor Pro and Xaa at
      position 58 is Ser or Thr, then Xaa at position 56 is not Asn
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (57)..(57)
<223> OTHER INFORMATION: Xaa at position 57 is any amino acid, provided
      that Xaa at position 56 is Asn and Xaa at position 58 is Ser or
      Thr, then Xaa at position 57 is Asp or Pro
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (58)..(58)
<223> OTHER INFORMATION: Xaa at position 58 is any amino acid, provided
      that Xaa at position 56 is Asn and Xaa at position 57 is neither
      Asp nor Pro, then Xaa at position 58 is neither Ser nor Thr

<400> SEQUENCE: 16

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr
20          25          30

Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val
35          40          45

Ala Gln Ile Asn Ser Val Gly Xaa Xaa Xaa Tyr Tyr Pro Asp Thr Val
50          55          60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
65          70          75          80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85          90          95

Ala Ser Gly Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
100         105         110

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
115         120         125

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
130         135         140

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
145         150         155         160

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
165         170         175

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
180         185         190

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
195         200         205

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
210         215         220

Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
225         230         235         240

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
245         250         255

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
260         265         270

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
275         280         285

```

-continued

```

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 290                295                300

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 305                310                315                320

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
                325                330                335

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu
                340                345                350

Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 355                360                365

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 370                375                380

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 385                390                395                400

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
                405                410                415

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
                420                425                430

Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 435                440

```

```

<210> SEQ ID NO 17
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Mus sp.
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Light chain CDR1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Light Chain CDR1

```

```

<400> SEQUENCE: 17

```

```

Arg Ser Ser Gln Ser Leu Val Tyr Ser Asp Gly Asn Ala Tyr Leu His
 1          5          10          15

```

```

<210> SEQ ID NO 18
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Mus sp.
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(17)
<223> OTHER INFORMATION: Heavy chain CDR2

```

```

<400> SEQUENCE: 18

```

```

Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Ser Val Lys
 1          5          10          15

```

```

Gly

```

1. A method to diagnose preclinical or clinical Alzheimer's disease in a subject, which method comprises administering to said subject an amount of an antibody which specifically binds an epitope contained within positions 13-28 of A β or an antibody that sequesters A β peptide from its bound, circulating form in the blood and alters clearance of soluble and bound forms of A β in the central nervous system in plasma;

effective to alter the levels of circulating A β peptides in the blood of said subject when said subject is in a preclinical or clinical stage of Alzheimer's disease, followed by measuring the level of, A β_{40} , A β_{42} , or the ratio of A β_{40} /A β_{42} in the blood of said subject at a time interval after said administering; and comparing the level of A β_{40} , A β_{42} , or the ratio of A β_{40} /A β_{42} in said subject with a control value of said levels, wherein

differing levels of $A\beta_{40}$, $A\beta_{42}$ or $A\beta_{40}/A\beta_{42}$ ratio in said subject as compared to control levels or ratio identifies said subject as in a preclinical or clinical stage of Alzheimer's disease.

2. The method of claim 1, wherein said time interval is less than 1 week.

3. The method of claim 1, wherein said time interval is less than or equal to 24 hours.

4. The method of claim 3, wherein the time interval is less than or equal to 3 hours.

5. The method of claim 1, wherein said administering is by injection of said antibodies.

6. The method of claim 1, wherein the subject is human and the antibody is a humanized antibody or a fragment thereof.

7. The method of claim 6, wherein the humanized antibody or fragment thereof comprises a light chain of the sequence given by SEQ ID NO:11 and a heavy chain of the sequence given by SEQ ID NO:12.

8. The method of claim 6, wherein the humanized antibody or fragment thereof comprises a light chain of the sequence given by SEQ ID NO:11 and a heavy chain of the sequence given by SEQ ID NO:16.

9. The method of claim 6, wherein the humanized antibody or fragment thereof comprises a light chain comprising a

variable region of the sequence given by SEQ ID NO:7 and a heavy chain comprising a variable region of the sequence given by SEQ ID NO:14.

10. The method of claim 1, wherein said antibody is a fragment.

11. The method of claim 1, wherein the antibody specifically binds to an epitope of $A\beta$ to which antibody 266 specifically binds.

12. The method of claim 1, wherein the antibody is a single-chain antibody.

13. A kit for the diagnosis of clinical or preclinical Alzheimer's disease in a subject which comprises a container containing an antibody which specifically binds an epitope contained within positions 13-28 of $A\beta$ or an antibody that sequesters $A\beta$ peptide from its bound, circulating form in the blood and alters clearance of soluble and bound forms of $A\beta$ in the central nervous system and in plasma and instructions for administering the antibody.

14. The kit of claim 13, which further contains a reagent for assessing the level of $A\beta_{40}$ and/or $A\beta_{42}$ in the blood.

15. The kit of claim 13, which further contains a description of control values for $A\beta_{40}$, $A\beta_{42}$, and/or $A\beta_{40}/A\beta_{42}$ ratios in blood of normal subjects.


* * * * *

专利名称(译)	阿尔茨海默病的测定方法		
公开(公告)号	US20100279433A1	公开(公告)日	2010-11-04
申请号	US12/834271	申请日	2010-07-12
[标]申请(专利权)人(译)	霍尔茨曼DAVID中号 DEMATTOS RONALD BALES KELLY - [R 康明斯大卫· PAUL史蒂芬中号		
申请(专利权)人(译)	霍尔茨曼DAVID中号 DEMATTOS RONALD BALES KELLY - [R 康明斯大卫· PAUL史蒂芬中号		
当前申请(专利权)人(译)	礼来公司 华盛顿大学		
[标]发明人	HOLTZMAN DAVID M DEMATTOS RONALD BALES KELLY R CUMMINS DAVID J PAUL STEVEN M		
发明人	HOLTZMAN, DAVID M. DEMATTOS, RONALD BALES, KELLY R. CUMMINS, DAVID J. PAUL, STEVEN M.		
IPC分类号	G01N33/53 G01N33/68		
CPC分类号	A61K2039/505 C07K16/18 C07K2317/24 G01N2800/2821 C07K2317/56 C07K2317/92 G01N33/6896 C07K2317/41		
优先权	PCT/US2002/026321 2002-08-16 WO 60/313221 2001-08-17 US 60/334987 2001-10-23 US 60/313224 2001-08-17 US		
其他公开文献	US8444977		
外部链接	Espacenet USPTO		


摘要(译)

临床前和临床阿尔茨海默氏病的诊断测试基于A β 40，A β 42的血浆水平，它们的比例或在施用螯合A β 的抗体后其进入速率。来自对照值的任何这些参数的改变确定了临床前或临床阿尔茨海默氏病。

A



B



C




Figure 1