



(19) **United States**

(12) **Patent Application Publication**
MACDONALD et al.

(10) **Pub. No.: US 2011/0154512 A1**

(43) **Pub. Date: Jun. 23, 2011**

(54) **HUMANIZED FC GAMMA R MICE**

Publication Classification

(75) Inventors: **Lynn MACDONALD**, White Plains, NY (US); **Naxin Tu**, Pleasantville, NY (US); **Cagan Gurer**, Valhalla, NY (US); **Sean Stevens**, San Francisco, CA (US); **Andrew J. Murphy**, Croton-on-Hudson, NY (US)

(51) **Int. Cl.**
A01K 67/027 (2006.01)
G01N 33/00 (2006.01)
C12Q 1/18 (2006.01)
G01N 33/53 (2006.01)

(52) **U.S. Cl.** **800/3; 800/18; 435/32; 435/7.1**

(57) **ABSTRACT**

(73) Assignee: **Regeneron Pharmaceuticals, Inc.**, Tarrytown, NY (US)


Genetically modified non-human animals and methods and compositions for making and using them are provided, wherein the genetic modification comprises a deletion of the endogenous low affinity FcγR locus, and wherein the mouse is capable of expressing a functional FcRγ-chain. Genetically modified mice are described, including mice that express low affinity human FcγR genes from the endogenous FcγR locus, and wherein the mice comprise a functional FcRγ-chain. Genetically modified mice that express up to five low affinity human FcγR genes on accessory cells of the host immune system are provided.

(21) Appl. No.: **12/971,080**

(22) Filed: **Dec. 17, 2010**

Related U.S. Application Data

(60) Provisional application No. 61/288,562, filed on Dec. 21, 2009.






Figure 1




Figure 2




Figure 3A





Figure 3B





Figure 3C





Figure 3D




Figure 4




Figure 5A




Figure 5B




Figure 6




Figure 7

HUMANIZED FC GAMMA R MICE

[0001] This application claims the benefit of the filing date under 35 USC §119(e), and is a nonprovisional, of U.S. Provisional Patent Application Ser. No. 61/288,562, filed 21 Dec. 2009, which provisional application is hereby incorporated by reference.

FIELD OF INVENTION

[0002] The field of invention is genetically modified non-human animals that lack endogenous murine FcγR genes, including genetically modified animals that comprise a replacement of endogenous FcγR genes with human FcγR genes, and including mice that are capable of expressing at least two, three, four, or five functional human low affinity FcγR genes, and including genetically modified mice comprising immune cells that do not express endogenous low affinity FcγR genes.

BACKGROUND

[0003] Fc receptors (FcRs) are proteins found on the surface of cells of the immune system that carry out a variety of functions of the immune system in mammals. FcRs exist in a variety of types, on a variety of cells, and mediate a variety of immune functions such as, for example, binding to antibodies that are attached to infected cells or invading pathogens, stimulating phagocytic or cytotoxic cells to destroy microbes, or infected cells by antibody-mediated phagocytosis or antibody-dependent cell-mediated cytotoxicity (ADCC).

[0004] ADCC is a process whereby effector cells of the immune system lyse a target cell bound by antibodies. This process depends on prior exposure to a foreign antigen or cell, resulting in an antibody response. ADCC can be mediated through effector cells such as, for example, natural killer (NK) cells, by binding of FcR expressed on the surface of the effector cell to the Fc portion of the antibody which itself is bound to the foreign antigen or cell. Because of the central role that FcRs play in the immune response, useful non-human animals that co-express multiple human FcRs are needed, including non-human animals that co-express multiple human low affinity FcRs. There exists a need for non-human animal models of human FcR function and human processes of ADCC for the study and elucidation of human disease therapies, in particular anti-tumor therapies and therapies for treating autoimmune diseases, and pharmaceutical drug development, in particular in the development, design, and testing of human antibody pharmaceuticals.

BRIEF DESCRIPTION OF THE FIGURES

[0005] FIG. 1 is a schematic depiction of a wild type low affinity FcγR locus in a mouse, showing mouse FcγRIIB, FcγRIV and FcγRIII genes and a mouse FcγR targeting vector used for a targeted deletion of these genes, which includes a neomycin cassette flanked by site-specific recombination sites.

[0006] FIG. 2 shows histograms of splenocytes gated for B cells (anti-CD19), NK cells (anti-NKp46) and macrophages (anti-F4/80) including expression of endogenous mFcγRII and mFcγRIII genes for wild type and low affinity FcγR α-chain gene-deficient mice (mFcγR KO).

[0007] FIGS. 3A-3D show in vivo depletion of B cells with a human anti-human CD20 antibody with mouse Fc (Ab 168)

or human Fc (Ab 735) in humanized CD20 mice (hCD20) and humanized CD20 mice bred to FcγR knockout mice (hCD20/FcγR KO) in several lymphocyte compartments: bone marrow (FIG. 3A), blood (FIG. 3B), lymph node (FIG. 3C) and spleen (FIG. 3D). For each graph, the y-axis shows the percent of gated B cells (B220⁺/IgM⁺ or B220⁺/CD19⁺) and the x-axis shows the antibody dose for each animal group: 10 mg/kg Control antibody (C), 2 mg/kg human anti-human CD20 antibody (2 Ab) and 10 mg/kg human anti-human CD20 antibody (10 Ab).

[0008] FIG. 4 is a schematic depiction of a neomycin-targeted deletion of the low-affinity mouse FcγR locus and a second targeting vector for inserting two human low affinity FcγR genes (hFcγRIIIA and hFcγRIIA) into the deleted mouse locus, which includes a hygromycin cassette flanked by site-specific recombination sites. For expression of hFcγRIIA on platelets, an extended promoter region operably linked to the hFcγRIIA gene of the Human FcγRIIIA-IIA Targeting Vector is employed; to prevent expression of hFcγRIIA on platelets, the promoter region is omitted or substantially omitted.

[0009] FIG. 5A shows histograms of splenocytes gated for NK cells (anti-NKp46) and macrophages (anti-F4/80) including expression of human FcγRIIIA for wild type and human FcγRIIIA-IIA homozygote mice (Human FcγRIIIA/FcγRIIA HO).

[0010] FIG. 5B shows histograms of splenocytes gated for neutrophils (anti-Ly6G) and macrophages (anti-F4/80) including expression of human FcγRIIA for wild type and human FcγRIIIA-IIA homozygote mice (Human FcγRIIIA/FcγRIIA HO).

[0011] FIG. 6 is a schematic depiction of a hygromycin-targeted deletion of the low affinity mouse FcγR locus including an insertion of two low affinity human FcγR genes (hFcγRIIIA and hFcγRIIA) and a third targeting vector for inserting three additional low affinity human FcγR genes (hFcγRIIB, hFcγRIIIB and hFcγRIIC) and a neomycin cassette flanked by site-specific recombination sites.

[0012] FIG. 7 shows histograms of splenocytes gated for B cells (anti-CD19) and neutrophils (anti-Ly6G) including expression of human FcγRIIB and human FcγRIIIB for wild type and human FcγRIIIA-IIIB-IIA-IIIB-IIIC homozygote mice (Human FcγRIIIA/FcγRIIIB/FcγRIIA/FcγRIIB/FcγRIIC HO).

SUMMARY

[0013] Genetically modified cells, non-human embryos, non-human animals and methods and compositions for making and using them are provided. In various aspects, the non-human animals comprise a human FcγR receptor, a deletion of an endogenous low affinity FcγR receptor, and/or a replacement of an endogenous FcγR receptor with a human FcγR receptor at an endogenous mouse low affinity FcγR locus.

[0014] In one aspect, genetically modified cells, non-human embryos, and non-human animals are provided that comprise a functional FcR γ-chain, wherein the cells, embryos, and animals comprise a further modification comprising a replacement of the low affinity endogenous non-human FcγR gene sequences (e.g., FcγRIIB, FcγRIV and FcγRIII) with one or more low affinity human FcγR gene sequences (e.g., selected from FcγRIIA, FcγRIIB, FcγRIIC, FcγRIIIA, FcγRIIIB, and a combination thereof).

[0015] In one embodiment, the cells, non-human embryos, and non-human animals are murine. In one embodiment, the functional FcR γ -chain is a mouse FcR γ -chain. In one embodiment, the mouse FcR γ -chain is an FcR γ -chain endogenous to the mouse, the cell, or the embryo.

[0016] In one embodiment, the cells, embryos, and animals are mice, and the mice express a functional α -chain of a human low affinity Fc γ R receptor and a functional endogenous mouse γ -chain.

[0017] In one aspect, a genetically modified mouse is provided, wherein the mouse does not express an endogenous α -chain selected from an Fc γ RIIB α -chain, an Fc γ RIV α -chain, an Fc γ RIII α -chain, and a combination thereof; wherein the mouse expresses a functional endogenous mouse γ -chain.

[0018] In a specific embodiment, the mouse does not express a functional Fc γ RIIB α -chain, does not express a functional Fc γ RIV α -chain, and does not express a functional Fc γ RIII α -chain.

[0019] In one embodiment, the mouse genome comprises a deletion of an endogenous Fc γ RIIB α -chain, a deletion of an endogenous Fc γ RIV α -chain, and a deletion of an endogenous Fc γ RIII α -chain.

[0020] In one embodiment, the mouse comprises a deletion of an endogenous Fc γ RIIB α -chain, a deletion of an endogenous Fc γ RIV α -chain, and a deletion of an endogenous Fc γ RIII α -chain, and further comprises a reduced ability to make an immune response to an antigen as compared with a wild type mouse's ability with respect to the same antigen. In one embodiment, the reduced immune response includes a decreased antibody-dependent cell-mediated cytotoxicity (ADCC). In one embodiment, the reduced immune response includes a reduced ability in a cell killing assay to achieve antibody-dependent NK cell killing. In specific embodiments, the reduction in ADCC or antibody-dependent NK cell killing is at least 50%, in one embodiment at least 75%, in one embodiment at least 90%.

[0021] In one embodiment, the mouse comprises a deletion of an endogenous Fc γ RIIB α -chain, a deletion of an endogenous Fc γ RIV α -chain, and a deletion of an endogenous Fc γ RIII α -chain, and further comprises an increased humoral antibody response upon immunization with an antigen as compared to a wild type mouse, e.g., a mouse of the same or similar strain that does not comprise the deletion. In one embodiment, the increased humoral antibody response is 2-fold as compared to a wild type mouse. In one embodiment, the increased humoral antibody response is 3-fold as compared to a wild type mouse. In one embodiment, the increased humoral antibody response is 5-fold as compared to a wild type mouse. In one embodiment, the increased humoral antibody response is 7-fold as compared to a wild type mouse. In one embodiment, the increased humoral antibody response is 10-fold as compared to a wild type mouse. In a specific embodiment, humoral antibody response is measured by micrograms of antibody that specifically binds an antigen (with which the mouse has been immunized) per microgram of serum protein from the mouse. In one embodiment, the increased humoral antibody response is with respect to an antigen to which a wild type mouse exhibits tolerance, or to an antigen which in a wild type mouse exhibits a poor or minimal humoral immune response. In a specific embodiment, the antigen is a mouse antigen. In a specific embodiment, the antigen is a human antigen that exhibits an identity with a mouse protein of at least about 95%, 96%, 97%, 98%, or 99%.

[0022] In one aspect, a genetically modified mouse is provided, comprising a replacement of a low affinity mouse Fc γ R α -chain gene with a low affinity human Fc γ R α -chain gene, wherein the replacement is at the endogenous mouse Fc γ R α -chain gene locus. In one embodiment, the low affinity mouse Fc γ R α -chain gene is selected from an Fc γ RIIB, Fc γ RIV and an Fc γ RIII α -chain gene. In a specific embodiment, a genetically modified mouse is provided, wherein the mouse expresses an endogenous FcR γ -chain, and wherein the low affinity human Fc γ R α -chain gene is Fc γ RIIA α -chain. In another specific embodiment, the genetically modified mouse expresses an endogenous FcR γ -chain and a functional human Fc γ RIIA α -chain on NK cells. In a specific embodiment, the functionality of Fc γ RIIA α -chain on NK cells is reflected by human antibody-mediated NK killing (e.g., ADCC mediated by a human antibody).

[0023] In one aspect, a genetically modified cell, non-human embryo, or non-human animal is provided, wherein the genetic modification comprises a replacement of at least one endogenous low affinity Fc γ R α -chain gene with a human Fc γ R α -chain gene, and the cell, embryo, or animal expresses a functional FcR γ -chain. In one embodiment, the functional FcR γ -chain is an endogenous FcR γ -chain. In one embodiment, the low affinity human Fc γ R α -chain gene is selected from an Fc γ RIIA α -chain gene, an Fc γ RIIA α -chain gene, and a combination thereof. In a specific embodiment, the human Fc γ RIIA gene comprises a polymorphism, wherein the polymorphism is selected from a 131His low responder polymorphism and a 131Arg high responder polymorphism. In a specific embodiment, the Fc γ RIIA polymorphism is the 131H is low responder polymorphism. In one embodiment, the Fc γ RIIA gene is a specific allelic variant, wherein the allelic variant is selected from a 158Val variant and a 158Phe variant. In a specific embodiment, the Fc γ RIIA allelic variant is the 158Val variant.

[0024] In one embodiment the low affinity human Fc γ R gene is selected from an Fc γ RIIB, Fc γ RIIC, an Fc γ RIIB gene, and a combination thereof. In a specific embodiment, the human Fc γ RIIB gene comprises an amino acid substitution, wherein the substitution is selected from an 232Ile or a 232Thr substitution. In another specific embodiment, amino acid substitution is a 232Ile substitution. In a specific embodiment, the Fc γ RIIB gene is a specific allelic variant, wherein the allelic variant is selected from a NA1 variant and a NA2 variant. In another specific embodiment, the Fc γ RIIB allelic variant is a NA2 variant.

[0025] In one embodiment the low-affinity human Fc γ R α -chain gene is selected from a Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIA, Fc γ RIIB α -chain gene, and a combination thereof.

[0026] In one embodiment, the low affinity mouse Fc γ RIV α -chain gene and the Fc γ RIII α -chain gene are replaced with at least one low affinity human Fc γ R α -chain gene. In one embodiment, the low affinity mouse Fc γ RIV α -chain gene and the Fc γ RIIB α -chain gene are replaced with at least one low affinity human Fc γ R α -chain gene. In one embodiment, the low affinity mouse Fc γ RIIB α -chain gene and the Fc γ RIII α -chain gene are replaced with at least one low affinity human Fc γ R α -chain gene. In a specific embodiment, the at least one low affinity human Fc γ R α -chain gene is selected from an Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIA, Fc γ RIIB α -chain gene, and a combination thereof. In another specific embodiment, the at least one low affinity human Fc γ R α -chain gene is selected from an Fc γ RIIA α -chain gene, an Fc γ RIIA

α -chain gene, and a combination thereof. In another specific embodiment, the at least one low affinity human Fc γ R α -chain gene is selected from an Fc γ RIIB, Fc γ RIIC, Fc γ RIIIB α -chain gene, and a combination thereof. In another specific embodiment, the low affinity mouse Fc γ R genes are replaced with a human Fc γ RIIA α -chain gene and a human Fc γ RIIA α -chain gene. In another specific embodiment, the low affinity human Fc γ RIIA and Fc γ RIIA α -chain genes comprise variants, wherein the Fc γ RIIA α -chain gene comprises a 131H is variant and the Fc γ RIIA α -chain gene comprises a 158Val variant. In another specific embodiment, the low affinity mouse Fc γ R α -chain genes are replaced with the following low affinity human Fc γ R α -chain genes: Fc γ RIIB, Fc γ RIIC and Fc γ RIIIB. In another specific embodiment, the low affinity human Fc γ RIIB α -chain gene and Fc γ RIIIB α -chain gene comprise variants, wherein the Fc γ RIIB α -chain gene comprises a 232Ile variant and the Fc γ RIIIB α -chain gene comprises an NA2 variant.

[0027] In one embodiment, the genetic modifications comprise a replacement of syntenic genomic sequences of mouse and human chromosome 1. In a specific embodiment, the genetic modifications comprise a replacement of a genomic fragment comprising endogenous low affinity mouse Fc γ R genes with a genomic fragment comprising low affinity human Fc γ R genes. In another specific embodiment, the mouse genome from chromosome 1:172,889,983 to chromosome1:172,989,911 is replaced with a human genomic fragment comprising human chromosome 1:161,474,729 to chromosome1:161,620,458.

[0028] In one aspect, a genetically modified cell, non-human embryo, or non-human animal is provided, wherein the genetic modification comprises a knockout of one or more endogenous low affinity receptor α -chain genes, and the presence of an episome comprising one or more human Fc γ R α -chain genes. In a specific embodiment, the cell, embryo, or animal expresses a functional FcR γ -chain. In a specific embodiment, the episome is a mini chromosome. In one embodiment, the functional FcR γ -chain is endogenous to the cell, embryo, or animal.

[0029] In one aspect, a genetically modified mouse is provided, comprising a replacement of a low affinity mouse Fc γ R α -chain gene with a low affinity human Fc γ R α -chain gene, and the mouse comprises a mouse FcR γ -chain gene, and the mouse expresses a functional human low affinity Fc γ R receptor. In one embodiment, the functional low affinity Fc γ R receptor is expressed on a cell type in which the low affinity Fc γ R receptor is expressed in humans. In a specific embodiment, the functional human low affinity Fc γ R receptor is Fc γ RIIA and the Fc γ RIIA is expressed on NK cells.

[0030] In one embodiment, the mouse comprises a deletion of two mouse Fc γ R α -chain genes. In another embodiment, the mouse comprises a deletion of three mouse Fc γ R α -chain genes.

[0031] In one embodiment, the mouse comprises a replacement of three mouse Fc γ R α -chain genes with at least one human Fc γ R α -chain gene. In another embodiment, the mouse comprises a replacement of two mouse Fc γ R α -chain genes with at least one human Fc γ R α -chain gene. In a specific embodiment, the mouse comprises a replacement of three mouse Fc γ R α -chain genes with at least two human Fc γ R α -chain genes. In another specific embodiment, the three mouse Fc γ R α -chain genes are replaced with three human Fc γ R α -chain genes. In another specific embodiment, the mouse comprises a replacement of two mouse Fc γ R

α -chain genes with at least two human Fc γ R α -chain genes. In yet another specific embodiment, the two mouse Fc γ R α -chain genes are replaced with at least three human Fc γ R α -chain genes.

[0032] In one embodiment, the low affinity mouse Fc γ R α -chain gene is selected from an Fc γ RIIB, Fc γ RIV, Fc γ RIII α -chain gene, and a combination thereof.

[0033] In one embodiment, the low affinity human Fc γ R α -chain gene is selected from an Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIA, Fc γ RIIIB α -chain gene, and a combination thereof. In one embodiment, the low affinity human Fc γ R α -chain gene is selected from an Fc γ RIIA, an Fc γ RIIA α -chain gene, and a combination thereof. In one embodiment, the low affinity human Fc γ R α -chain gene is selected from an Fc γ RIIB, Fc γ RIIC, an Fc γ RIIIB α -chain gene, and a combination thereof.

[0034] In one embodiment, the low affinity mouse Fc γ RIV α -chain gene and the Fc γ RIII α -chain gene are replaced with at least one human Fc γ R α -chain gene. In one embodiment, the low-affinity mouse Fc γ RIV α -chain gene and the Fc γ RIIB α -chain gene are replaced with at least one human Fc γ R α -chain gene. In one embodiment, the low affinity mouse Fc γ RIIB α -chain gene and the Fc γ RIIIB α -chain gene are replaced with at least one human Fc γ R α -chain gene. In a specific embodiment, the at least one human Fc γ R α -chain gene is selected from an Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIA, Fc γ RIIIB α -chain gene, and a combination thereof. In another specific embodiment, the at least one human Fc γ R α -chain gene is selected from an Fc γ RIIA, an Fc γ RIIA α -chain gene, and a combination thereof. In another specific embodiment, the at least one human Fc γ R α -chain gene is selected from an Fc γ RIIB, Fc γ RIIC, Fc γ RIIIB α -chain gene, and a combination thereof. In another specific embodiment, the mouse α -chain genes are replaced with the following human Fc γ R α -chain genes: Fc γ RIIA and Fc γ RIIA. In yet another specific embodiment, the mouse α -chain genes are replaced with the following human Fc γ R α -chain genes: Fc γ RIIB, Fc γ RIIC and Fc γ RIIIB.

[0035] In one aspect, a genetically modified mouse is provided, comprising a low affinity human Fc γ R α -chain and a mouse FcR γ -chain subunit, wherein the mouse expresses the human Fc γ R α -chain on a cell selected from a neutrophil, an eosinophil, a basophil, a monocyte, a macrophage, a platelet, a Langerhans cell, a dendritic cell, an NK cell, a mast cell, a B cell, a T cell, and a combination thereof. In one embodiment, the mouse expresses a human Fc γ RIIA α -chain on a cell selected from a neutrophil, a macrophage, an eosinophil, a platelet, a dendritic cell, a Langerhans cell, and a combination thereof. In one embodiment, the mouse is capable of phagocytosis, ADCC and cellular activation initiated or mediated through the expressed human Fc γ RIIA α -chain. In one embodiment the mouse expresses a human Fc γ RIIA α -chain on a cell selected from a macrophage, an NK cell, a monocyte, a mast cell, an eosinophil, a dendritic cell, a Langerhans cell, at least one T cell type, and a combination thereof. In one embodiment, the mouse is capable of ADCC mediated through the human Fc γ RIIA α -chain expressed on NK cells. In a specific embodiment, the mouse exhibits hFc γ RIIA-mediated ADCC in response to an antibody comprising a human Fc.

[0036] In one embodiment, the mouse expresses both a human Fc γ RIIA α -chain and a human Fc γ RIIA α -chain. In one embodiment, the human Fc γ RIIA α -chain is expressed

on platelets and the human Fc γ RIIIA α -chain is expressed on NK cells. In one embodiment, the mouse is capable of ADCC mediated by an antibody comprising a human Fc, wherein the mediation is through either the human Fc γ RIIA α -chain or through the human Fc γ RIIIA α -chain expressed on the surface of accessory cells. In one embodiment, the human Fc γ RIIA α -chain is not expressed on platelets. In a specific embodiment wherein the human Fc γ RIIA α -chain is not expressed on platelets, the mouse lacks or substantially lacks a human promoter sequence that operably linked to the human Fc γ RIIA α -chain in a human genome.

[0037] In one embodiment, the mouse expresses a human Fc γ RIIB α -chain on a cell selected from a B cell, a mast cell, a basophil, a macrophage, an eosinophil, a neutrophil, a dendritic cell, a Langerhans cell, and a combination thereof. In a specific embodiment, the mouse expresses a human Fc γ RIIB α -chain on a B cell and a mast cell. In another specific embodiment, the mouse is capable of endocytosis of immune complexes mediated through the expressed human Fc γ RIIB α -chain. In one embodiment, the mouse expresses a human Fc γ RIIC α -chain on a cell selected from a neutrophil, a macrophage, an eosinophil, a platelet, a dendritic cell, a Langerhans cell, and a combination thereof. In a specific embodiment, the mouse is capable of phagocytosis, ADCC and cellular activation initiated through the expressed human Fc γ RIIC α -chain.

[0038] In one embodiment, the mouse expresses a human Fc γ RIIIB α -chain on neutrophils and eosinophils. In a specific embodiment, the mouse is capable of cellular activation, phagocytosis, ADCC and degranulation, wherein the activation, phagocytosis, ADCC, and degranulation are mediated through the expressed human Fc γ RIIIB α -chain.

[0039] In one aspect, a mouse is provided that comprises a deletion of the endogenous Fc γ RIIB, Fc γ RIV and Fc γ RIII genes and insertion of human Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIIA, and Fc γ RIIIB genes, and wherein the mouse comprises a functional mouse FcR γ -chain gene.

[0040] In one embodiment, the mouse comprises a deletion of the α -chains encoded by endogenous Fc γ RIIB, Fc γ RIV and Fc γ RIII genes and insertion of the α -chains encoded by human Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIIA, and Fc γ RIIIB genes.

[0041] In one embodiment, the insertion of the human Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIIA, and Fc γ RIIIB α -chain genes is at a random location within the mouse genome.

[0042] In one embodiment, the insertion of the human Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIIA, and Fc γ RIIIB α -chain genes is at the endogenous mouse low affinity Fc γ R α -chain locus.

[0043] In one embodiment, the mouse expresses human Fc γ RIIIA on NK cells and macrophages. In a specific embodiment, all or substantially all NK cells from a splenocyte sample of the mouse express human Fc γ RIIIA. In a specific embodiment, all or substantially all macrophages from a splenocyte sample of the mouse express human Fc γ RIIIA.

[0044] In one embodiment, the mouse expresses a human Fc γ R selected from human Fc γ RIIA, human Fc γ RIIIA, and a combination thereof, on a cell type selected from neutrophils, macrophages, and a combination thereof. In a specific embodiment, the mouse expresses human Fc γ RIIA and human FORMA on all or substantially all neutrophils and macrophages of a splenocyte sample from the mouse.

[0045] In one embodiment, the mouse expresses human Fc γ RIIB and human Fc γ RIIIB on B cells and neutrophils of B cells from a B cell-gated splenocyte sample from the mouse. In a specific embodiment, the mouse expresses Fc γ RIIIB and Fc γ RIIB on all or substantially all B cells and neutrophils from a B cell-gated splenocyte sample from the mouse.

[0046] In one embodiment, the mouse further comprises a humanized CD20 gene. In one embodiment, the mouse that further comprises the humanized CD20 gene following treatment with an anti-CD20 binding protein that comprises an Fc exhibits depletion (in vivo) of B cells. In one embodiment, the depletion is in a compartment selected from bone marrow, blood, lymph node, spleen, and a combination thereof. In one embodiment, the Fc is a human Fc. In one embodiment, the Fc is a mouse Fc. In one embodiment, the anti-CD20 binding protein is an anti-CD20 antibody.

[0047] In one aspect, a cell is provided comprising a genetic modification as described herein. In one embodiment, the cell is selected from an embryonic stem (ES) cell, a pluripotent cell, an induced pluripotent cell, and a totipotent cell. In one embodiment, the cell is selected from a mouse cell and a rat cell. In a specific embodiment, the cell is an ES cell. In a more specific embodiment, the cell is a mouse ES cell.

[0048] In one aspect, a non-human embryo is provided, comprising a genetic modification as described herein. In one embodiment, the non-human embryo is selected from a mouse embryo and a rat embryo.

[0049] In one aspect, a method is provided for determining efficacy of a therapeutic.

[0050] In one embodiment, the therapeutic is an antibody (e.g., mono-, bi-, tri-, multispecific) comprising a human Fc. In one embodiment, the therapeutic is a human antibody. In one embodiment, the efficacy is efficacy of therapeutic-mediated cell killing (e.g., ADCC). In a specific embodiment, the human therapeutic is a fusion protein comprising an Fc of a human immunoglobulin heavy chain. In one embodiment, the therapeutic is administered to a mouse as described herein and a level of therapeutic-dependent ADCC is measured. In one embodiment, the mouse is used to assess the ADCC activity of a therapeutic by administering the therapeutic to the mouse and then detecting (e.g., in vitro from a sample (e.g., blood) taken from the animal) binding of the therapeutic to a human low affinity Fc γ R on an Fc γ R-expressing cell. In a specific embodiment, accessory cells of the mouse are isolated from the mouse and tested for the ability, in the presence and absence of the therapeutic, to mediate therapeutic-dependent ADCC.

[0051] In one aspect, a method is provided for determining whether a low affinity Fc γ R is associated with a human disease or disorder, comprising a step of determining a trait associated with the human disease or disorder in a mouse according to the invention. In one embodiment, the trait is a phenotype associated with the absence or loss of a function of one or more low affinity Fc γ Rs. In a specific embodiment, the disease or disorder is an autoimmune disease or disorder. In a specific embodiment, the autoimmune disease or disorder is selected from Rheumatoid Arthritis (RA), Systemic Lupus Erythematosus (SLE), type I diabetes, Guillain-Barré syndrome, sclerosis, multiple sclerosis, Goodpasture's syndrome, Wegener's Granulomatosis and experimental autoimmune encephalomyelitis (EAE). In a specific embodiment, the mouse comprises a polymorphism in a low affinity Fc γ R, and the trait is selected from an enhanced ability to mediate ADCC in comparison to the majority of the human popula-

tion that does not bear the polymorphism, and a reduced ability to mediate ADCC in comparison to the majority of the human population that does not bear the polymorphism.

[0052] In one aspect, a method for making an anti-human FcR α -chain antibody in a mouse is provided, comprising exposing a mouse according to the invention to a human FcR as described herein. In one embodiment, an antibody that recognizes the human FcR is isolated from the mouse. In another embodiment, a nucleic acid sequence that encodes all or part of a variable region of an antibody that recognizes the human FcR is identified and cloned.

[0053] In one aspect, a method for determining ability of anti-human FcR antibodies to target molecules to FcR-expressing cells for phagocytosis of the target molecule is provided, comprising exposing a mouse as described herein to an agent comprising an anti-human FcR antibody, and measuring phagocytosis of the target molecule.

[0054] In one aspect, a method is provided for making an antibody, in a mouse, to an antigen that is poorly immunogenic in a mouse that is wild type with respect to one or more Fc γ Rs, comprising exposing a mouse as described herein that lacks a mouse low affinity FcR but expresses an Fc γ R γ -chain to the antigen that is poorly immunogenic in the mouse that is wild type with respect to one or more Fc γ Rs, and identifying an antibody that recognizes the poorly antigenic antigen. In one embodiment, the method comprises isolating the antibody from the mouse. In another embodiment, a nucleic acid sequence that encodes all or part of a variable region of the antibody is identified and cloned.

[0055] In one aspect, a method for making a mouse capable of making antibodies comprising human variable regions is provided, comprising a step of breeding a first mouse as described herein with a second mouse that comprises (a) one or more human immunoglobulin variable region gene segments and one or more human constant region genes; or, (b) one or more human immunoglobulin variable region gene segments operably linked to a mouse constant region gene, wherein the human gene segments replace variable region gene segments at the mouse variable region gene segment locus.

[0056] In one embodiment, the second mouse (a) comprises a transgene that comprises one or more human immunoglobulin light chain variable region gene segments and a human light chain constant gene, and a transgene that comprises one or more human immunoglobulin heavy chain variable region gene segments and one or more human heavy chain constant genes. In one embodiment, the transgene that comprises one or more human immunoglobulin heavy chain variable region gene segments comprises two or more heavy chain constant genes and is capable of class switching. In a specific embodiment, the mouse comprises an inactivated endogenous light chain locus and/or an inactivated endogenous heavy chain locus. In a specific embodiment, the mouse comprises a deletion of an endogenous light chain locus and/or a deletion of an endogenous heavy chain locus.

[0057] In one embodiment, the second mouse (b) comprises human heavy and human light variable region gene segments, at the heavy and light mouse loci, respectively.

[0058] In one aspect, a method is provided for selecting an anti-tumor antibody, comprising a step of determining the ability of an antibody to mediate ADCC, wherein the ability of the antibody to mediate ADCC is tested by determining ADCC mediated by a cell of a mouse as described herein, and the antibody is selected if it mediates ADCC employing a cell

of a genetically modified mouse as described herein. In a specific embodiment, binding of the antibody to the cell of the genetically modified mouse is determined, and the anti-tumor antibody is selected for its ability to bind a human FOR on the cell. In a specific embodiment, the human FOR is a low affinity Fc γ R.

[0059] In one embodiment, the anti-tumor antibody is identified by its enhanced ability to mediate ADCC through a cell of the mouse as compared to ability of the anti-tumor antibody to mediate ADCC through a cell of a wild type mouse. In a specific embodiment, the anti-tumor antibody is identified by its ability to mediate ADCC through NK cells. In a specific embodiment, the NK cells express human Fc γ RIIIA.

[0060] In one embodiment, a method is provided for selecting an anti-tumor agent, comprising a step of administering an agent comprising a human Fc or a modified human Fc to a first non-human animal wherein the first non-human animal is genetically modified in accordance with the invention and comprises a human tumor; a step of administering the agent to a second non-human animal comprising the tumor; and determining the ability of the first non-human animal and the second non-human animal to retard growth of the human tumor following administration of the agent, wherein the agent is selected as an anti-tumor agent if it exhibits an enhanced ability to retard growth of the human tumor in the first non-human animal but not in the second non-human animal.

[0061] In one embodiment, the first non-human animal is modified to comprise a deletion of an endogenous FcR α -subunit, and is modified to comprise a human FcR α -subunit selected from the group consisting of an Fc γ RIIA α -subunit, an Fc γ RIIB α -subunit, an Fc γ RIIC α -subunit, an Fc γ RIIIA α -subunit, an Fc γ RIIIB α -subunit, and a combination thereof. In one embodiment, the second animal is a wild type animal. In one embodiment, the first non-human animal expresses an endogenous FcR γ -chain.

[0062] In one embodiment, the first non-human animal expresses a functional endogenous Fc γ RI.

[0063] In one aspect, a method is provided for making a mouse that lacks a low affinity mouse Fc γ R, expresses a functional FcR γ -chain, and comprises genes encoding α -chains of the human Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIIA, and Fc γ RIIIB, comprising a step of replacing the low affinity mouse Fc γ R α -chains with human Fc γ R α -chains, at the mouse Fc γ R α -chain locus.

[0064] In one embodiment, a first step comprises deleting the α -chains of the endogenous Fc γ RIIB, Fc γ RIV and Fc γ RIII genes and inserting the α -chains of the human Fc γ RIIA and Fc γ RIIIA genes; a second step comprises inserting the α -chains of the human Fc γ RIIB, Fc γ RIIC and Fc γ RIIIB genes into the mouse genome that results from the first step; wherein the mouse comprises a functional mouse FcR γ -chain gene. In a specific embodiment, the α -chains of the human Fc γ RIIB, Fc γ RIIC and Fc γ RIIIB genes of the second step are inserted 5' relative to the α -chains of the human Fc γ RIIA and Fc γ RIIIA genes of the first step.

[0065] In one aspect, a method for determining cell killing by a human therapeutic in a non-primate is provided, comprising a step of exposing a cell, non-human embryo, or non-human animal to a human therapeutic that comprises a human Fc, wherein the cell, embryo, or animal comprises a functional FcR γ -chain and comprises a replacement of one or more endogenous low affinity Fc γ R α -chain genes with one or more human Fc γ R α -chains, and determining the ability of

the human therapeutic to mediate cell killing through a low affinity human Fc γ R of the cell, embryo, or animal.

[0066] In one embodiment, the non-primate is a mouse. In a specific embodiment, endogenous mouse Fc γ R α -chain genes Fc γ RIIB, Fc γ RIV and Fc γ RIII are replaced with human Fc γ R α -chain genes Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIIA, and Fc γ RIIIB.

[0067] In one embodiment, the cell is selected from a B cell, a mast cell, a basophil, a macrophage, an eosinophil, a neutrophil, a dendritic cell, a Langerhans cell, and a combination thereof. In a specific embodiment, the cell is an NK cell and NK cell-mediated ADCC by a human or a humanized antibody is determined. In a specific embodiment, the low affinity human Fc γ R is a human Fc γ RIIIA.

[0068] In one aspect, a method for determining therapeutic-dependent thrombosis is provided, comprising exposing a first non-human animal that expresses a human Fc γ RIIA on a platelet to a therapeutic; exposing a second non-human animal that does not express the human Fc γ RIIA on a platelet to said therapeutic; measuring in the first non-human animal and in the second non-human animal an amount of therapeutic-dependent thrombosis; and, determining a difference in therapeutic-dependent thrombosis.

[0069] In one embodiment, the non-human animal is selected from a mouse and a rat.

[0070] In one embodiment, the determined difference in therapeutic-dependent thrombosis is employed to identify a risk associated with administering the therapeutic to a human. In one embodiment, the determined difference results in a change of administration of the therapeutic to a human patient in need thereof.

DETAILED DESCRIPTION

[0071] The invention is not limited to particular methods, and experimental conditions described, as such methods and conditions may vary. The terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the claims.

[0072] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, particular methods and materials are now described. All publications mentioned herein are incorporated herein by reference in their entirety.

[0073] The phrase “targeting construct” includes a polynucleotide molecule that comprises a targeting region. A targeting region comprises a sequence that is substantially homologous to a sequence in a target cell, tissue or animal and provides for integration of the targeting construct into a position within the genome of the cell, tissue or animal. In a specific embodiment, the targeting construct further comprises a nucleic acid sequence or gene of particular interest, a selectable marker, control and or regulatory sequences, and other nucleic acid sequences that allow for recombination mediated through the exogenous addition of proteins that aid in or facilitate recombination involving such sequences. In another specific embodiment, the targeting construct further comprises a gene of interest, wherein the gene of interest is a heterologous gene that encodes a protein that has a similar function as a protein encoded by the endogenous sequence.

[0074] The term “replacement” includes wherein a DNA sequence is placed into a genome of a cell in such a way as to replace a sequence within a genome, at the locus of the genomic sequence, with a heterologous sequence (e.g., a human sequence in a mouse), unless otherwise indicated. The DNA sequence so placed may include one or more regulatory sequences that are part of source DNA used to obtain the sequence so placed (e.g., promoters, enhancers, 5'- or 3'-untranslated regions, etc.). For example, in various embodiments, the replacement is a substitution of an endogenous sequence for a heterologous sequence that results in the production of a gene product from the DNA sequence so placed (comprising the heterologous sequence), but not expression of the endogenous sequence; the replacement is of an endogenous genomic sequence with a DNA sequence that encodes a protein that has a similar function as a protein encoded by the endogenous genomic sequence (e.g., the endogenous genomic sequence encodes a low affinity mouse Fc γ R receptor, and the DNA fragment encodes one or more human low affinity Fc γ R receptors, such as, e.g., a human Fc γ RIIC and/or an Fc γ RIIIB).

[0075] The term “Fc γ R” includes a receptor for an Fc, e.g., an Fc portion of an IgG immunoglobulin. The Fc γ R genes include an α -chain that is expressed on the surface of the cell and serves as a ligand-binding domain, and associates with either a homodimer of the FcR γ -chain or a heterodimer of the FcR γ -chain and the δ -chain. There are several different Fc γ R genes and they can be categorized into low affinity and high affinity types according to preferential binding to IgG in immune complexes. Low affinity Fc γ R genes in humans include Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIIA and Fc γ RIIIB and within most of these genes naturally occurring genetic differences, or polymorphisms, have been described in human subjects with autoimmune diseases. Persons of skill upon reading this disclosure will recognize that one or more endogenous low affinity Fc γ R genes in a genome (or all) can be replaced by one or more heterologous low affinity Fc γ R genes (e.g., variants or polymorphisms such as allelic forms, genes from another species, chimeric forms, etc.).

[0076] The phrase “allelic variants” includes variations of a normal sequence of a gene resulting in a series of different forms of the same gene. The different forms may comprise differences of up to, e.g., 20 amino acids in the sequence of a protein from a gene. For example, alleles can be understood to be alternative DNA sequences at the same physical gene locus, which may or may not result in different traits (e.g., heritable phenotypic characteristics) such as susceptibility to certain diseases or conditions that do not result in other alleles for the same gene or result in varying degrees in the other alleles.

[0077] An “accessory cell” includes an immune cell that is involved in the effector functions of the immune response. Exemplary immune cells include a cell of lymphoid or myeloid origin, e.g., lymphocytes, natural killer (NK) cells, monocytes, macrophages, neutrophils, eosinophils, basophils, platelets, Langerhans cells, dendritic cells, mast cells etc. Accessory cells carry out specific functions of the immune system through receptors, e.g., FcRs, expressed on their surfaces. In a specific embodiment, an accessory cell is capable of triggering ADCC mediated through an FcR, e.g., a low affinity Fc γ R, expressed on the cell surface. For example, macrophages expressing FcRs are involved in phagocytosis and destruction of antibody-coated bacteria. Accessory cells might also be capable of releasing an agent that mediates

other immune processes. For example, mast cells can be activated by antibody bound to FcRs to release granules, e.g., inflammatory molecules (e.g., cytokines) at a site of infection. In various other embodiments, the expression of FcRs on accessory cells can be regulated by other factors (e.g., cytokines). For example, Fc γ RI and Fc γ RIII expression can be induced by stimulation with interferon- γ (IFN- γ).

Mouse and Human FcRs

[0078] The receptors for the Fc (i.e., constant) regions of immunoglobulins (FcRs) play an important role in the regulation of the immune response. FcRs are present on accessory cells of the host's immune system to effectively dispose of foreign antigens bound by an antibody. FcRs also play important roles in balancing both activating and inhibitory responses of the accessory cells of the immune system. FcRs are involved in phagocytosis by macrophages, degranulation of mast cells, uptake of antibody-antigen complexes and modulation of the immune response, as well as other immune system processes.

[0079] In mice and humans, distinct FcRs are differentially expressed on the surface of different accessory cells that are each specific for the immunoglobulin isotypes present in the expressed antibody repertoire. For example, immunoglobulin G (IgG) antibodies mediate effector functions through IgG receptors (Fc γ Rs). Fc γ Rs have been classified into three groups: high affinity activating Fc γ RI (CD64), low affinity inhibitory Fc γ RII (CD32) and low affinity activating Fc γ RIII (CD16). Although each group is present in both mice and humans, the number of isoforms and subsets of immune cells on which they are present are different. For example, Fc γ RIIA and Fc γ RIIIB are expressed on accessory cells in humans but are reportedly absent from mice. Further, affinities of the different IgG isotypes (e.g., IgG1) for each Fc γ R is different in mice and humans.

[0080] Activation or inhibition of cell signaling through Fc γ Rs and the effector functions associated with antibody binding to Fc γ Rs are believed to be mediated by specific sequence motifs of intracellular domains of Fc γ Rs, or of the subunits of co-receptors. Activating receptors are most commonly associated with the common γ -chain (FcR γ -chain) which contains an immunoreceptor tyrosine-based activation motif (ITAM). ITAMs contain a specific sequence of about 9-12 amino acids that include tyrosine residues that are phosphorylated in response to antibody binding to an FcR. Phosphorylation leads to a signal transduction cascade. Mice that lack a gene encoding an FcR γ -chain (FcR γ -chain KO) have been reported (e.g., see Takai et al. (1994) FcR γ Chain Depletion Results in Pleiotropic Effector Cell Defects, *Cell* 76:519-529; van Vugt et al. (1996) FcR γ -Chain Is Essential for Both Surface Expression and Function of Human Fc γ RI (CD64) In Vivo, *Blood* 87(9):3593-3599; and Park et al. (1998) Resistance of Fc Receptor-deficient Mice to Fatal Glomerulonephritis, *J. Clin. Invest.* 102(6):1229-1238). The FcR γ -chain is reportedly essential for proper surface expression and function (e.g., signal transduction, phagocytosis, etc.) of most of the FcRs; FcR γ -chain KO mice lack Fc γ RI according to some reports. However, other reports reveal that FcR γ -chain KO mice indeed express Fc γ RI on the surface of certain accessory cells, and the Fc γ RI expressed reportedly appears functional in that it binds IgG in mice in the absence of expressed FcR γ -chain (Barnes et al. (2002) Fc γ RI-Deficient Mice Show Multiple Alterations to Inflammatory and Immune Responses, *Immunity* 16:379-389).

[0081] In contrast, Fc γ RIIB is an inhibitory receptor that contains an immunoreceptor tyrosine-based inhibitory motif (ITIM) in its cytoplasmic domain. Like ITAMs, ITIMs are sequence motifs that include phosphorylatable tyrosine residues. However, downstream events following phosphorylation of an ITM lead to inhibition, not activation, of immune cell functions. Mice deficient in Fc γ RIIB reportedly exhibit an increased antibody response in comparison to wild type mice (Takai et al. (1996) Augmented humoral and anaphylactic responses in Fc γ RII-deficient mice, *Nature* 379:346-349), an observation that supports the role of Fc γ RIIB as a downregulator of the B cell antibody response.

[0082] In humans, Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIIA and Fc γ RIIIB are considered the classical low affinity Fc γ R genes and are located together on the same chromosome (Su et al. (2002) Genomic organization of classical human low-affinity Fc γ receptor genes, *Genes and Immunity* 3 (Suppl 1):S51-S56). These genes exhibit several polymorphisms associated with distinct phenotypes, e.g., an alteration of ligand binding and function of the receptor. Some polymorphisms are associated with autoimmune diseases, e.g., systemic lupus erythematosus (SLE), rheumatoid arthritis (RA), and multiple sclerosis (MS). Transgenic mice for different human Fc γ Rs (hFc γ Rs) have been developed and used as disease models, generating high affinity antibodies, testing therapeutic antibodies for ability to elicit specific cellular responses, screening compounds that ameliorate aberrant immune responses, etc. (e.g., see Heijnen et al. (1996) A Human Fc γ RI/CD64 Transgenic Model for In Vivo Analysis of (Bispecific) Antibody Therapeutics, *J. Hematother.* 4:351-356; Heijnen and van de Winkel (1996) Antigen Targeting to Myeloid-specific Human Fc γ RI/CD64 Triggers Enhanced Antibody Responses in Transgenic, *J. Clin. Invest.* 97(2):331-338; U.S. Pat. Nos. 6,111,166, 6,676,927, 7,351,875, 7,402,728, and 7,416,726).

[0083] Despite the significant roles of the FcRs in providing the bridge between antibodies and accessory cells of the immune system, no model system currently exists in which all the low affinity hFc γ Rs are expressed. A mouse in which all the low-affinity hFc γ Rs are co-expressed—including mice that lack endogenous mouse Fc γ Rs—in various embodiments could be used to accurately reflect effects of a human antibody therapeutic, including ADCC-mediated effects. Such a mouse would serve as a vital tool in the engineering, analysis and evaluation of therapeutic antibodies for treatment of human diseases such as, e.g., RA, type 1 diabetes, SLE, and autoimmunity, by providing an animal model capable of achieving a more accurate assessment of immunological processes in humans, particularly in the context of testing human antibody therapeutics. The mouse will also be a valuable source of cells bearing the low affinity receptors, which cells can be used in *in vitro* assays for assessing therapeutic-dependent cell killing for therapeutics that bind the low affinity receptors, and thus for identifying useful human therapeutics.

Endogenous Low Affinity Fc γ R Gene Deficient Mice

[0084] Genetically modified non-human animals are provided that do not express endogenous low affinity mouse Fc γ R genes, but that express an endogenous mouse FcR γ -chain. In various embodiments, the FcR γ -chain is expressed in a distribution (i.e., in cell types) and at a level in the mouse that is the same or substantially the same as in a wild type mouse. Endogenous low affinity Fc γ R genes can be

expressed either on the surface of immune cells or in a soluble manner in the periphery of the animals. Genetic modifications for making a non-human animal that does not express endogenous low affinity mouse Fc γ R genes are conveniently described by using the mouse as an illustration. A genetically modified mouse according to the invention can be made in a variety of ways, particular embodiments of which are discussed herein.

[0085] A schematic illustration (not to scale) of low affinity mouse Fc γ R gene locus is provided in FIG. 1 (top) to show Fc γ R gene arrangement at the endogenous locus. As illustrated, low affinity mouse Fc γ R genes Fc γ RIIB, Fc γ RIV and Fc γ RIII are present together in close proximity on one chromosome. Each of these genes comprise the α -chain or ligand binding domain responsible for the binding the Fc portion of an antibody molecule.

[0086] A genetically modified mouse lacking a nucleotide sequence encoding an α -chain of the endogenous low affinity Fc γ R genes can be made by any method known in the art. For example, a targeting vector can be made that deletes the low affinity mouse Fc γ R α -chain genes with selectable marker gene. FIG. 1 illustrates a mouse genome (bottom) targeted by a targeting construct having a 5' homology arm containing sequence upstream of the endogenous low affinity Fc γ R α -chain locus, followed by a drug selection cassette (e.g. a neomycin resistance gene flanked by loxP sequences), and a 3' homology arm containing sequence downstream of the endogenous low affinity Fc γ R α -chain locus. Upon homologous recombination at the locus, the endogenous low affinity Fc γ R α -chain locus is replaced by a drug selection cassette (bottom of FIG. 1). The endogenous low affinity Fc γ R α -chain gene locus is thereby deleted resulting in a cell or non-human animal that does not express endogenous low-affinity mouse Fc γ R α -chain genes. The drug selection cassette may optionally be removed by the subsequent addition of a recombinase (e.g., by Cre treatment).

[0087] Genetically modifying a mouse to render an endogenous low-affinity mouse Fc γ R α -chain gene or genes non-functional, in various embodiments, results in a mouse that exhibits defects in immune responses, making the mouse useful for evaluating cooperative, as well as individual, roles of the endogenous low-affinity mouse Fc γ R genes in normal and disordered immune function, IgG-mediated processes, and autoimmune disease. In various embodiments, modifying the α -chains of the endogenous low-affinity mouse Fc γ R genes, but not the FcR γ -chain, avoids a potential reduction of other endogenous FcR genes (e.g., high affinity Fc γ RI) that require the FcR γ -chain for surface expression and function, thus maintaining various other immunological functions and processes mediated through γ -chain-dependent processes.

[0088] According to some reports, FcR γ -chain deficient mice lack surface expression of Fc γ RIII and Fc γ RI. However, Fc γ RI has reportedly been detected on the cell surface in FcR γ -chain deficient mice and is reportedly at least partially functional. In contrast, mice according to the present invention contain unmodified endogenous FcR γ -chain, which preserves natural cell surface expression patterns and cellular functions of other FcR genes that require FcR γ -chain.

[0089] In various embodiments, mice of the present invention present an advantage over other Fc γ R gene-deficient mice in that the genetic modifications that they bear result in the maintenance of other genes necessary for other immunological functions not entirely devoted to low affinity Fc γ R genes. For example, with a functional FcR γ -chain, other

γ -chain-dependent proteins (e.g., Fc γ RI) will be able to associate with the FcR γ -chain and participate in effector cell functions in the immune response. In various genetically modified mice in accordance with the invention, it is believed that maintaining such functions (due to the presence of a functional FcR γ -chain) while deleting endogenous low affinity Fc γ R genes (one or more α -subunits) enables a more precise elucidation of the roles of FcRs in autoimmunity.

Low Affinity Fc γ R Humanized Mice

[0090] Genetically modified non-human animals are provided that express low-affinity human Fc γ R genes. Low affinity human Fc γ R genes can be expressed either on the surface of accessory cells of the animal's immune system or in a soluble manner in the periphery of the animals.

[0091] The genetic modification, in various embodiments, comprises a deletion of a functional α -chain of one or more low-affinity mouse Fc γ R genes, and in some embodiments a further modification comprising a replacement with two or more, with three or more, with four or more, or with five low-affinity human Fc γ R α -subunit genes, wherein the non-human animal expresses a functional mouse FcR γ -chain gene. Genetically modified non-human embryos, cells, and targeting constructs for making the non-human animals, non-human embryos, and cells are also provided.

[0092] Compositions and methods for making a mouse that expresses a human Fc γ R gene, including specific polymorphic forms or allelic variants (e.g., single amino acid differences), are provided, including compositions and method for making a mouse that expresses such genes from a human promoter and a human regulatory sequence. The methods include selectively rendering an endogenous low affinity mouse Fc γ R gene nonfunctional (e.g., by a deletion of its α -chain), and employing an α -chain of a low affinity human Fc γ R gene at the endogenous low affinity mouse Fc γ R gene locus to express a low affinity human Fc γ R α -subunit gene in a mouse. The deletion of the low affinity mouse Fc γ R gene is made by deletion of one or more α -chain genes, but not an FcR γ -chain gene. The approach selectively renders one or more endogenous low affinity Fc γ R α -chain genes nonfunctional while retaining a functional endogenous FcR γ -chain.

[0093] The endogenous Fc γ R α -chain replacement approach employs a relatively minimal disruption in natural Fc γ R-mediated signal transduction in the animal, in various embodiments, because the genomic sequence of the Fc γ R α -chains are replaced in a single fragment and therefore retain normal functionality by including necessary regulatory sequences. Thus, in such embodiments, the Fc γ R α -chain modification does not affect other endogenous FcRs dependent upon functional FcR γ -chain molecules. Further, in various embodiments, the modification does not affect the assembly of a functional receptor complex involving an Fc γ R α -chain and the endogenous FcR γ -chain, which is believed to be required for proper expression of some Fc γ R α -chains on the cell surface and for downstream signaling resulting from an activated receptor. Because the FcR γ -chain is not deleted, in various embodiments animals containing a replacement of endogenous Fc γ R α -chain genes with human Fc γ R α -chain genes should be able to process normal effector functions from antibodies through binding of the Fc portion of IgG immunoglobulins to the human Fc γ R α -chains present on the surface of accessory cells.

[0094] A schematic illustration (not to scale) of a deleted endogenous low affinity mouse Fc γ R gene is provided in FIG.

4 (top). As illustrated, low affinity human FcγR genes FcγRIIA and FcγRIIA are inserted into the deleted endogenous low affinity mouse FcγR gene locus by a targeting construct (Human FcγRIIA-IIA Targeting Vector) with a genomic fragment containing the human low affinity human FcγRIIA and FcγRIIA genes. Each of these genes comprise the α-chain or ligand-binding domain of the human FcγR genes responsible for the binding the Fc portion of an antibody molecule.

[0095] A genetically modified mouse that expresses low affinity human FcγR genes at the endogenous low affinity mouse FcγR locus can be made by any method known in the art. For example, a targeting vector can be made that introduces low affinity human FcγR genes (e.g., FcγRIIA and FcγRIIA) with a selectable marker gene. FIG. 4 illustrates a mouse genome comprising a deletion of the endogenous low affinity FcγR locus (top). As illustrated, the targeting construct contains a 5' homology arm containing sequence upstream of the endogenous low affinity mouse FcγR locus, followed by a drug selection cassette (e.g., a hygromycin resistance gene flanked on both sides by loxP sequences), a genomic fragment containing a human FcγRIIA gene, human HSP76 gene and human FcγRIIA gene, and a 3' homology arm containing sequence downstream of the endogenous low affinity mouse FcγR locus. Upon homologous recombination at the deleted locus, the drug selection cassette is replaced by the sequence contained in the targeting vector (bottom of FIG. 4). The endogenous low affinity FcγR gene locus is thus replaced with low affinity human FcγR genes resulting in a cell or animal that expresses low-affinity human FcγR genes. The drug selection cassette may optionally be removed by the subsequent addition of a recombinase (e.g., by Cre treatment).

[0096] For expression of hFcγRIIA on platelets, the targeting construct Human hFcγRIIA-IIA Targeting Vector comprises an extended sequence that includes, e.g., all or substantially all of the human promoter region operably linked to the hFcγRIIA gene in a human genome. For preventing expression of hFcγRIIA on platelets, the targeting construct lacks all or substantially all of the human promoter region operably linked to the hFcγRIIA gene in a human.

[0097] Further modifications to the chimeric locus (bottom of FIG. 4) can be achieved using similar techniques as described for replacement with two human FcγR genes. The modification to replace the endogenous low affinity FcγR gene locus with two human FcγR genes can further provide a starting point for incorporation of other low affinity human FcγR genes. For example, a schematic illustration (not to scale) of an endogenous low affinity FcγR locus replaced with two human low affinity FcγR genes is provided in FIG. 6 (top). As illustrated, low affinity human FcγR genes FcγRIIB, FcγRIIC and FcγRIIB are inserted into the modified endogenous low affinity mouse FcγR gene locus by another targeting construct (Human FcγRIIB-IIIB-IIC Targeting Vector) with a genomic fragment containing the low affinity human FcγRIIB, FcγRIIC and FcγRIIB genes. Each of these genes comprise the α-chain or ligand-binding domain of the human FcγR genes responsible for the binding the Fc portion of an antibody molecule.

[0098] A genetically modified mouse that expresses five low affinity human FcγR genes at the endogenous low affinity mouse FcγR locus can be made by any method known in the art. For example, a targeting vector can be made that introduces low affinity human FcγR genes (e.g., FcγRIIB, FcγRIIC

and FcγRIIB) with a selectable marker gene. FIG. 6 illustrates a mouse genome comprising a replacement of the endogenous low affinity FcγR locus with two low affinity human FcγR genes (top). As illustrated, the targeting construct contains a 5' homology arm containing sequence upstream of the endogenous low affinity mouse FcγR locus, followed by a drug selection cassette (e.g., a neomycin resistance gene flanked on both sides by loxP sequences), a genomic fragment containing a human FcγRIIB gene, a human FcγRIIB, a human HSP77 gene, a human FcγRIIC gene, followed by a 3' homology arm containing sequence upstream of the low affinity human FcγRIIA gene present at the endogenous locus. Upon homologous recombination at the modified locus, a human FcγRIIB, FcγRIIB and FcγRIIC gene are inserted 5' to the human FcγRIIA and FcγRIIA genes previously present at the endogenous low affinity FcγR gene locus by the sequence contained in the targeting vector (bottom of FIG. 6). The modified endogenous low affinity FcγR gene locus is thus further modified to incorporate three additional low affinity human FcγR genes resulting in a cell or animal that expresses five low-affinity human FcγR genes. The drug selection cassette may optionally be removed by the subsequent addition of a recombinase (e.g., by Cre treatment). FIG. 6 (bottom) shows the structure of the resulting locus, which will express five low affinity human FcγR genes that can be detected on the surface of accessory cells of the animal's immune system and independently associate, as appropriate, with an endogenous Fcγ-chain.

Experimental Models of FcγR Deficient Mice and FcγR Humanized Mice

[0099] Genetically modified non-human animals that do not express endogenous low affinity mouse FcγR genes are useful, e.g., to elucidate the various functions of the individual low affinity FcγR genes in the immune response, to measure the efficacy of a human therapeutic antibody via cell-mediated immunity (e.g., ADCC), to determine an FcγR's role in immune diseases or disorder, to serve as models of immune diseases or disorders, to generate antibodies against one or more FcγR proteins, and to serve as breeding mates to generate other genetically modified mice of interest.

[0100] In one embodiment, a mouse according to the invention can be used to determine a cytotoxic effect lost (in comparison to a wild type mouse) by a mouse that does not express low affinity FcγR genes by administering an agent to such a mouse, where the agent is known to trigger an FcγR-dependent cytotoxic effect in wild type mice. In one embodiment, a mouse of the present invention is implanted with tumor cells and, after a subsequent period of time, injected with an antibody specific for an antigen expressed on the surface of the tumor cells. The isotype of the antibody is known prior to injection and the animals are analyzed for impairment of FcγR-dependent ADCC by comparison to ADCC observed in wild type animals.

[0101] In another aspect, mice deficient in endogenous low affinity receptors could be combined (e.g., by breeding) with other immune deficient mice to develop in vivo models of autoimmune disease. For example, Severe Combined Immunodeficiency (SCID) mice are routinely used in the art as model organisms for studying the immune system. SCID mice have an impaired ability to make T or B lymphocytes, or activate some components of the complement system, and cannot efficiently fight infections, reject tumors, and reject transplants. Low affinity FcγR α-subunit gene-deficient mice

of the present invention may be bred to SCID mice to ascertain cell depletion in a host animal in response to administration of an antibody therapeutic (e.g., an anti-tumor antibody), which would determine the roles of ADCC and complement-dependent cytotoxicity (CDC) in tumor cell depletion in vivo.

[0102] In another aspect, genetically modified non-human animals comprising a replacement of the endogenous low affinity Fc γ R genes with low-affinity human Fc γ R genes are provided. Such animals are useful for studying the pharmacokinetics of fully human antibodies and hFc γ R-mediated ADCC. In addition, human Fc γ R genes have been shown to exhibit polymorphisms or allelic variants associated with disease (e.g., SLE, RA, Wegener's granulomatosis, Guillain-Barré syndrome and Multiple Sclerosis). Thus, genetically modified non-human animals that comprise a replacement of the endogenous low affinity Fc γ R genes with specific allelic or polymorphic forms of human Fc γ R genes can be used to study human autoimmune diseases, and traits associated with the polymorphisms, in the animal. In a specific embodiment, the allelic forms of human Fc γ R genes are associated with enhanced efficacy for human IgG.

[0103] In another specific embodiment, the effect of a human low affinity Fc γ R polymorphism on the efficacy of a human antibody therapeutic is determined. In a specific embodiment, an anti-tumor antibody is administered to a first humanized mouse comprising a first polymorphism of a human Fc γ R and also to a second humanized mouse comprising a second polymorphism of a human Fc γ R, wherein the first and the second mice each comprise a human tumor cell; and the anti-tumor activity of the anti-tumor antibody is assessed in the first mouse and in the second mouse. In a specific embodiment, a treatment option is selected by a physician with respect to treating a human having the first or the second polymorphism and having a tumor corresponding to the human tumor cell, based on the assessment of efficacy of the anti-tumor antibody in the first mouse and in the second mouse.

[0104] Suitable polymorphisms of human Fc γ R genes include all those known in the art. For the human Fc γ RIIA gene, polymorphisms include, e.g., the high responder and low responder phenotype reported by the ability of T cells to proliferate in response to IgG. The high responder polymorphism is characterized by an arginine residue at position 131 (131Arg) while the low responder is characterized by a histidine residue at position 131 (131His). In a specific embodiment, the human Fc γ RIIA sequence comprises the 131His polymorphism. A representative protein sequence of the human Fc γ RIIA α -chain is shown in SEQ ID NO:32.

[0105] Single-nucleotide substitutions of the human Fc γ RIIB gene result in mis-sense substitutions in the ligand-binding domain (α -chain) and putatively affect the binding ability of an Fc portion of an IgG to bind to the α -chain of Fc γ RIIB on the cell surface. For example, substitution of a threonine residue for an isoleucine at position 232 (Ile232Thr) within the transmembrane domain of the Fc γ RIIB gene in mice has been shown to impair the signaling ability of the receptor. In a specific embodiment, the human Fc γ RIIB gene comprises the isoleucine variant (232Ile). A representative protein sequence of the human Fc γ RIIB α -chain is shown in SEQ ID NO:33.

[0106] Allelic variants of the human Fc γ RIIIA gene are proposed to be involved in susceptibility to SLE and RA. This allelic variant includes a phenylalanine substitution for valine at position 158 (Val158Phe). The valine allelic variant

(158Val) is characterized to have a higher affinity for IgG1 and IgG3 than the phenylalanine allelic variant (158Phe). The 158Phe allelic variant has been proposed to lead to a reduced clearance of immune complexes. In a specific embodiment, the human Fc γ RIIIA gene comprises the 158Val allelic variant. A representative protein sequence of the human Fc γ RIIIA α -chain is shown in SEQ ID NO:35.

[0107] Allelic variants of the human Fc γ RIIIB gene include the neutrophil antigen 1 (NA1) and neutrophil antigen 2 (NA2) alleles. These allelic variants have been proposed to be involved in blood-transfusion reactions, alloimmune neutropenia, SLE and Wegener's granulomatosis. The NA2 allelic variant is characterized by a diminished ability to mediate phagocytosis. In a specific embodiment, the human Fc γ RIIIB gene comprises the NA2 allelic variant. A representative protein sequence of the human Fc γ RIIIB α -chain is shown in SEQ ID NO:36.

[0108] In one aspect, the genetically modified non-human animals are useful for optimizing Fc γ R-mediated functions triggered by the Fc portion of therapeutic antibodies. The Fc regions of antibodies can be modified by any method known in the art. For example, amino acid residues within the Fc portion (e.g., CH2 and CH3 domains) can be modified to selectively enhance the binding affinity to human Fc γ RIIIA. Thus, the resulting antibody should have enhanced Fc γ RIIIA-dependent ADCC. In a specific embodiment, an animal expressing human Fc γ RIIIA of the present invention is used to evaluate the enhanced ADCC ability of a modified human antibody by administering a modified human antibody to the animal, detecting (e.g., in vitro) antibody binding to Fc γ RIIIA-expressing cells and comparing the ADCC activity observed to the ADCC activity observed from that determined in a wild type animal.

EXAMPLES

Example 1

Generation of Low Affinity Fc γ R Gene Deficient Mice

[0109] A targeting construct for introducing a deletion of the endogenous low affinity mouse Fc γ R locus (described below) was constructed (FIG. 1).

[0110] The targeting construct was made using VELOCIGENE® technology (see, e.g., U.S. Pat. No. 6,586,251 and Valenzuela et al. (2003) High-throughput engineering of the mouse genome coupled with high-resolution expression analysis, *Nature Biotech.* 21(6):652-659) to modify the Bacterial Artificial Chromosome (BAC) RP23-395f6 (Invitrogen). RP23-395f6 BAC DNA was modified to delete the endogenous low affinity Fc γ RIIB, Fc γ RIV and FORM genes comprising the α -chain of each of the Fc γ Rs.

[0111] Briefly, upstream and downstream homology arms were made employing primers mFcR 5-up-1 (5'-ACCAGGATAT GACCTGTAGA G; SEQ ID NO:1) and mFcR 3-up-1a (GTCCATGGGT AAGTAGAAAC A; SEQ ID NO:2), and mFcR 5-DN (ATGCGAGCTC ATGCATCTATG TCGGGT-GCGG AGAAAGAGGT AATGCATTCT TGCCCAATAC TTAC; SEQ ID NO:3) and mFcR 3-DN (ACTCATGGAG CCTCAACAGG A; SEQ ID NO:4), respectively. These homology arms were used to make a cassette that deleted the α -chains of the endogenous low affinity Fc γ RIIB, Fc γ RIV and Fc γ RIII genes. The targeting construct included a loxod neomycin resistance gene comprising homology arms comprising sequence homologous to a 5' and a 3' region with

respect to the endogenous locus. Genes and/or sequences upstream of the endogenous FcγRIIB gene and downstream of the endogenous FcγRIII gene (see FIG. 1) were unmodified by the targeting construct.

[0112] The targeted deletion was confirmed by polymerase chain reaction (PCR) using primers outside the deleted region and within the targeting construct. The upstream region of the deleted locus was confirmed by PCR using primers to mFcR-up-detect (ATCCTGAGTA TACTATGACA AGA; SEQ ID NO:5) and PGK-up-detect (ACTAGTGAGA CGTGC-TACTT C; SEQ ID NO:6), whereas the downstream region of the deleted locus was confirmed using primers pA-DN-detect (CTCCCACTCA TGATCTATAG A; SEQ ID NO:7) and mFcR-DN-detect (TGGAGCCTCA ACAGGACTCC A; SEQ ID NO:8). The nucleotide sequence across the upstream deletion point included the following, which indicates endogenous mouse sequence downstream of the FcγRIIB gene (contained within the parentheses below) linked contiguously to cassette sequence present at the deletion point: (GTC-CATGGGT AAGTAGAAAC A)TTCGCTACC TTAGGAC-CGT TA (SEQ ID NO:9). The nucleotide sequence across the downstream deletion point included the following, which indicates cassette sequence contiguous with endogenous mouse sequence upstream of the FcγRIII gene (contained within the parentheses below): CGGGTGCGGA GAAA-GAGGTA AT(GCATTCTT GCCCAATACT TA) (SEQ ID NO:10).

[0113] Mice deficient in FcγRIIB, FcγRIII and FcγRIV were generated through electroporation of a targeted BAC DNA (described above) into mouse ES cells. Positive ES cell clones are confirmed by Taqman™ screening and karyotyping. Positive ES cell clones were then used to implant female mice to give rise to a litter of pups deficient in low affinity FcγR genes.

Example 2

Characterization of Low Affinity FcγR Gene Deficient Mice

[0114] Spleens were harvested from FcγR deficient and wild type mice and perfused with 10 mL Collagenase-D in sterile disposable bags. Each bag containing a single spleen was then placed into a Stomacher® (Seward) and homogenized at a medium setting for 30 seconds. Homogenized spleens were transferred to 10 cm petri dishes and incubated for 25 minutes at 37° C. Cells were separated with a pipette using a 1:50 dilution of 0.5 M EDTA, followed by another incubation for five minutes at 37° C. Cells were then pelleted with a centrifuge (1000 rpm for 10 minutes) and red blood cells were lysed in 4 mL ACK buffer (Invitrogen) for three minutes. Splenocytes were diluted with RPMI-1640 (Sigma) and centrifuged again. Pelleted cells were resuspended in 10 mL RPMI-1640 and filtered with a 0.2 μm cell strainer.

[0115] Flow Cytometry. Lymphocyte cell populations were identified by FACs on the BD LSR II System (BD Bioscience) with the following fluorescence conjugated cell surface markers: anti-CD19 (B cells), anti-CD3 (T cells), anti-NKp46 (NK cells) and anti-F4/80 (macrophages). Lymphocytes were gated for specific cell lineages and analyzed for expression of endogenous FcγRIII and FcγRIIB with a rat anti-mouse FcγRIII/II antibody (clone 2.4G2, BD Biosciences). Clone 2.4G2 recognizes a common polymorphic epitope on the extracellular domains of murine FcγRIII and FcγRII. The results show that there was no detectable murine

low affinity FORM or FcγRII on B-cells, NK cells and macrophages in mFcγR KO mice (FIG. 2).

[0116] ADCC Assay. Splenocytes isolated from FcγR gene deficient and wild type mice were analyzed for their ability to perform ADCC in a cell-killing assay. Cell populations were isolated and separated using MACS® Technology (Miltenyi Biotec). Briefly, T-cells were depleted from splenocytes using magnetically labeled anti-mouse CD3 beads. The T-cell depleted splenocytes were then enriched for NK cells using magnetically labeled anti-mouse CD49B beads. Separately, Raji cells (expressing human CD20) were coated with varying concentrations (ranging from 0.1 to 10 μg/mL) of mouse anti-human CD20 antibody (Clone B1; Beckman Coulter) for 30 minutes at 4° C. The antibody-coated Raji cells were incubated with the enriched NK cells at ratios (NK:Raji) of 100:1 and 50:1 for four hours at 37° C. Cell death was measured using the CytoTox-Glo™ Cytotoxicity Assay (Promega). Luminescence signal is derived from lysed cells and proportional to the number of dead cells. Luminescence from controls (no anti-CD20 antibody) was determined for background dead cell count for each ratio and subtracted from measurements for wild type and KO mice. Average cell death was calculated and percent decrease in cell killing (% ADCC) was determined by comparison to wild type. Results are shown in Table 1.

TABLE 1

	mFcγR KO	% ADCC		
		10 μg/mL B1 Antibody	1 μg/mL B1 Antibody	0.1 μg/mL B1 Antibody
NK cell:Raji cell	100:1	42	53	35
	50:1	15	0	0

Example 3

In Vivo Depletion of B cells in Low Affinity FcγR Gene Deficient Mice

[0117] The effect of human or murine Fc isotypes on B cell depletion through the ADCC pathway was determined for various B cell compartments in low affinity FcγR gene deficient mice engineered to express human CD20 using a human anti-human CD20 antibody. Mice expressing human CD20 were separately engineered using techniques known in the art. Mice that express human CD20 on B cells and deficient in low affinity FcγR genes (described in Example 1) were made by standard breeding techniques of the two engineered strains.

[0118] Separate groups of mice that expressed human CD20 and had a full complement of endogenous low affinity FcγR genes were each administered one of the following: (1) 10 mg/kg control antibody (N=4; human antibody not specific for human CD20 having a mouse IgG2a); (2) 2 mg/kg Ab 168 (N=3; human anti-hCD20 antibody with a mouse IgG2a; heavy and light chain variable region sequences found in SEQ ID NOs: 339 and 347, respectively, of US Patent Publication No. 2009/0035322); (3) 10 mg/kg Ab 168; (4) 2 mg/kg Ab 735 (N=3; Ab 168 with human IgG1); (5) 10 mg/kg Ab 735. In a similar set of experiments, groups of mice that expressed human CD20 and had a deletion of the endogenous low affinity FcγR genes were administered the control and human anti-hCD20 antibodies (described above).

[0119] Mice in each group were administered the antibodies by intra-peritoneal injections. Seven days post-injection, animals were euthanized and the remaining B cell contents of bone marrow (B220⁺/IgM⁺), peripheral blood (B220⁺/CD19⁺), lymph node (B220⁺/CD19⁺) and spleen (B220⁺/CD19⁺) were identified by multi-color FACS performed on a LSR-II flow cytometer and analyzed using Flow-Jo software (as described above). The results of the B cell depletion experiments are shown in FIGS. 3A-3D.

[0120] As shown in FIGS. 3A-3D, Ab 735 depleted B cells with a lower efficiency than Ab 168 in mice containing a complete complement of low affinity FcγR genes. Further, for both antibodies (mouse and human Fc), B cell depletion was significantly reduced in mice lacking a complete complement of low affinity FcγR genes. This Example shows that the ability to deplete B cells through the ADCC pathway requires low affinity FcγRs and demonstrate that measuring ADCC efficiency for antibodies containing human constant regions in mice is more suitable by the use of genetically engineered mice containing a full complement of human low affinity FcγR genes.

Example 4

Generation of FcγRIIIA/FcγRIIA Humanized Mice

[0121] A targeting construct for introducing two low affinity human FcγR genes into a deleted endogenous low affinity mouse FcγR locus (described below) was constructed (FIG. 4).

[0122] A targeting construct comprising human FcγRIIA and FcγRIIIA genes was made using similar methods (see Example 1) through modification of BAC RP23-395f6 and CTD-2514j12 (Invitrogen). BAC DNA of both BACs was modified to introduce a deletion of the α-chains of the low affinity human FcγRIIA and FcγRIIIA genes into the deleted endogenous low affinity FcγR locus.

[0123] In a similar fashion, upstream and downstream homology arms were made employing primers h14 (GC-CAGCCACA AAGGAGATAA TC; SEQ ID NO:11) and h15 (GCAACATTTA GGACAACTCG GG; SEQ ID NO:12), and h4 (GATTTCTTAA CCACCTACCC C; SEQ ID NO:13) and h5 (TCTTTCCAA TGGCAGTTG; SEQ ID NO:14), respectively. These homology arms were used to make a cassette that introduced the α-chains of low affinity human FcγRIIA and FcγRIIIA genes into the endogenous mouse low affinity FcγR locus. The targeting construct included a 5' homology arm including sequence 5' to the deleted endogenous low affinity FcγR locus, a FRT⁺ hygromycin resistance gene, followed by a human genomic fragment from BAC CTD-2514j12 comprising low affinity human FcγRIIA and FcγRIIIA α-chain genes, and a 3' homology arm comprising mouse sequence 3' to the deleted endogenous low affinity FcγR locus (middle of FIG. 4). For a mouse that expresses FcγRIIA on mouse platelets, a targeting construct was made in a similar manner (using the same BACs) except that the construct comprises an extended promoter sequence operably linked to the human FcγRIIA gene in the human genome, e.g., up to about 18 kb or more, using a hygromycin cassette that is flanked on both sides by lox2372 sites, wherein the junction of the promoter region and the first lox2372 site is ATCGGGGATA GAGATGTTT (CC)GC-GATCGC GGTACCGGGC (SEQ ID NO:37 human/lox2372 junction in parentheses) and wherein the junction of the second lox2372 site and mouse sequence is TTATACGAAG

TTATACCGG(T G)CATTCTTGC CCAATACTTA (SEQ ID NO:38 lox2372/mouse junction in parentheses). Suitable primers were used to genotype the humanization comprising the promoter region.

[0124] Targeted insertion of the human FcγRIIA and FcγRIIIA α-chain genes was confirmed by PCR (as described above). The upstream region of the partially humanized locus was confirmed by PCR using primers h16 (CCCAGGTAAG TCGTGATGAA ACAG; SEQ ID NO:15) and pA-DN-detect (CTCCCACTCA TGATCTATAG A; SEQ ID NO:16), whereas the downstream region of the partially humanized locus was confirmed using primers mFcR DN-detect-9 (TG-GAGCCTCA ACAGGACTCC A; SEQ ID NO:17) and h6 (CACACATCTC CTGGTGACTT G; SEQ ID NO:18). The nucleotide sequence across the downstream junction included the following, which indicates a novel insertion point of endogenous human sequence upstream of the hFcγRIIA gene (contained within the parentheses below) contiguous with endogenous mouse sequence 3' of the deleted low affinity FcγR locus: (CAACTGCCAT TGGAAAAGA)C TCGAGTGCCA TTTCATTACC TC (SEQ ID NO:19). The upstream junction includes two novel sequences. One point of the upstream junction includes the following, which indicates nucleotide sequence of the hygromycin cassette contiguous with human genomic sequence (contained within the parentheses below) that comprises the upstream region of the inserted hFcγRIIIA gene: TAAACCCGCG GTG-GAGCTC(G CCAGCCACAA AGGAGATAAT CA) (SEQ ID NO:20). The second point of the upstream junction includes the following, which indicates a nucleotide sequence of an endogenous mouse sequence (contained within the parentheses below) from the upstream region of the deleted low affinity FcγR locus contiguous with a nucleotide sequence within the hygromycin cassette: (CCATGGGTAA GTAGAAAC)TC TAGACCCCGG GGCTCGATAA CT (SEQ ID NO:21).

[0125] Mice containing two low affinity human FcγR genes (hFcγRIIA, lacking extended promoter region, and hFcγRIIIA) in place of the endogenous low affinity mouse FcγR locus were generated through electroporation of the targeted BAC DNA (described above) into mouse ES cells. Positive ES cell clones were confirmed by TaqmanTM screening and karyotyping. Positive ES cell clones were then used to implant female mice using the VELOCIMOUSE[®] method (described below) to generate a litter of pups containing a replacement of the endogenous low affinity FcγR genes with the two human low affinity FcγR genes.

[0126] Targeted ES cells described above were used as donor ES cells and introduced into an 8-cell stage mouse embryo by the VELOCIMOUSE[®] method (see, e.g., U.S. Pat. No. 7,294,754 and Poueymirou et al. (2007) F0 generation mice that are essentially fully derived from the donor gene-targeted ES cells allowing immediate phenotypic analyses Nature Biotech. 25(1):91-99. VELOCIMICE[®] (F0 mice fully derived from the donor ES cell) bearing hFcγRIIA and hFcγRIIIA were identified by genotyping using a modification of allele assay (Valenzuela et al., supra) that detected the presence of the hFcγR genes.

[0127] Mice bearing the hFcγR genes can be bred to a Cre deleter mouse strain (see, e.g., International Patent Application Publication No. WO 2009/114400) in order to remove any loxed neo cassette introduced by the targeting construct that is not removed, e.g., at the ES cell stage or in the embryo. Optionally, the neomycin cassette is retained in the mice.

[0128] Pups are genotyped and a pup heterozygous for the hFcγR genes is selected for characterizing FcγRIIA and FcγRIIA humanizations.

Example 5

Characterization of FcγRIIA/FcγRIIA Humanized Mice

[0129] Spleens were harvested from humanized FcγRIIA/FcγRIIA (heterozygotes, lacking the extended FcγRIIA promoter region) and wild type mice and prepared for FACs (as described above).

[0130] Flow Cytometry. Lymphocytes were gated for specific cell lineages and analyzed for expression of hFcγRII and hFcγRIII using a mouse anti-human FcγRII antibody (Clone FL18.26; BD Biosciences) and a mouse anti-human FcγRIII antibody (Clone 3G8; BD Biosciences), respectively. Relative expression (++, +) or no expression (-) observed for each lymphocyte subpopulation is shown in Table 2.

TABLE 2

Lymphocyte Lineage	hFcγRIII	hFcγRII
B cells	-	-
NK cells	++	-
Macrophages	+	+
Neutrophils	-	+

[0131] In a similar experiment, spleens were harvested from humanized FcγRIIA/FcγRIIA (homozygotes, lacking the extended FcγRIIA promoter region) and wild type mice and prepared for FACs (as described above). Results are shown in FIGS. 5A and 5B. Percent of separate lymphocyte cell populations expressing human FcγRIIA or both in FcγRIIA/FcγRIIA homozygote mice is shown in Table 3.

TABLE 3

Lymphocyte Lineage	hFcγRIII	hFcγRII	hFcγRII/hFcγRIII
NK cells	97	—	—
Macrophages	26	14	39
Neutrophils	—	94	—

[0132] As shown in this Example, genetically modified mice (both heterozygote and homozygote genotypes) generated in accordance with Example 3 expressed human FcγRIIA on NK cells and macrophages; and human FcγRIIA on neutrophils and macrophages, but not platelets. Human FcγRIIA was highly expressed on NK cells. The expression pattern of human FcγR genes shown in this Example is consistent with the expression patterns of these genes in human accessory cells.

Example 6

Generation of Low Affinity FcγR Humanized Mice

[0133] A targeting construct for introducing three additional low affinity human FcγR genes into a partially humanized endogenous low affinity FcγR locus (described below) was constructed (FIG. 6).

[0134] A targeting construct comprising human FcγRIIB, FcγRIIB and FcγRIIC genes was made using similar methods (see Example 1) through modification of BAC RP-23 395f6 and RP-11 697e5 (Invitrogen). BAC DNA of both

BACs was modified to introduce the α-chains of the low affinity human FcγRIIB, FcγRIIB and FcγRIIC genes into the partially humanized endogenous low affinity FcγR locus containing two human low affinity FcγR genes.

[0135] In a similar fashion, upstream and downstream homology arms were made employing primers mFcR up-1 (ACCAGGATAT GACCTGTAGA G; SEQ ID NO:22) and mFcR2b NheI-2 (GTTTCTACTT ACCCATGGAC; SEQ ID NO:23), and h10 (AAATACACAC TGCCACAGAC AG; SEQ ID NO:24) and h11 (CCTCTTTTGT GAGTTTCCTG TG; SEQ ID NO:25), respectively. These homology arms were used to make a cassette that introduced DNA sequences encoding the α-chains of low affinity human FcγRIIB, FcγRIIB and FcγRIIC. The targeting construct included a 5' homology arm including mouse sequence 5' to the deleted endogenous low affinity FcγR locus, a loxed neomycin resistance gene, followed by a human genomic fragment from BAC RP-11 697e5 comprising low affinity human FcγRIIB, FcγRIIB and FcγRIIC α-chain genes, and a 3' homology arm comprising human sequence 5' to the low affinity human FcγRIIA α-chain gene (middle of FIG. 6).

[0136] Targeted insertion of three additional low affinity human FcγR genes was confirmed by PCR (as described above). The upstream region of the fully humanized locus was confirmed by PCR using primers mFcR up-detect-3 (GAGTACTA TGACAAGAGC ATC; SEQ ID NO:26) and PGK up-detect (ACTAGTGAGA CGTGCTACTT C; SEQ ID NO:27), whereas the downstream region of the fully humanized locus was confirmed using primers neo detect (CTCCCACTCA TGATCTATAG A; SEQ ID NO:28) and h12 (CTTTTATGG TCCCAATC AG; SEQ ID NO:29). The nucleotide sequence across the downstream junction included the same human genomic sequence upstream of the hFcγRIIA α-chain gene (see Example 3; SEQ ID NO:19). The nucleotide sequence across the upstream junction included the following, which indicates two novel junctions of mouse and cassette sequences and cassette and human genomic sequences at the insertion point. The junction of genomic mouse sequence (contained within the parentheses below) and the upstream region of the neo cassette sequence is: (GTCCATGGGT AAGTAGAAAC A)TTCGCTACC TTAGGACCGT TA (SEQ ID NO:30). The second novel junction includes the joining of the 3' end of neo cassette (contained within the parentheses below) and a human genomic sequence downstream of the hFcγRIIB α-chain gene: (GCTTATCGAT ACCGTCGAC)A AATACACACT GCCACAGACA GG; SEQ ID NO:31). These junctions are shown in FIG. 6 (middle) within the targeting construct. The resulting modified genome of the fully humanized low affinity FcγR locus is shown in FIG. 6 (bottom).

[0137] Mice containing five low affinity human FcγR genes in place of the endogenous low affinity mouse FcγR locus were generated through electroporation of the targeted BAC DNA (described above) into mouse ES cells. Positive ES cell clones were confirmed by Taqman™ screening and karyotyping. Positive ES cell clones were then used to implant female mice (as described above) to give rise to a litter of pups containing a replacement of the endogenous low affinity FcγR genes for five human low affinity FcγR genes.

Example 7

Characterization of Low Affinity FcγR Humanized Mice

[0138] Spleens were harvested from fully humanized FcγR (heterozygotes) and wild type mice and prepared for FACs (as described above).

[0139] Flow Cytometry. Lymphocytes were gated for specific cell lineages and analyzed for expression of human Fc γ RIIA and Fc γ RIIIA using a mouse anti-human Fc γ RII antibody (Clone FL18.26; BD Biosciences) and a mouse anti-human Fc γ RIII antibody (Clone 3G8; BD Biosciences), respectively. Relative expression (++, +) or no expression (–) observed for each lymphocyte subpopulation is shown in Table 4.

TABLE 4

Lymphocyte Lineage	hFc γ RIII	hFc γ RII
B cells	–	+
NK cells	+	+
Macrophages	+	+
Neutrophils	+	+

[0140] In a similar experiment, spleens were harvested from fully humanized Fc γ R (homozygotes) and wild type mice and prepared for FACs (as described above). Results are shown in FIG. 7. Percent of separate lymphocyte cell populations expressing human Fc γ RIIIA, human Fc γ RIIIB, human Fc γ RIIA, human Fc γ RIIB, human Fc γ RIIC or a combination thereof in fully humanized Fc γ R homozygote mice is shown in Table 5.

TABLE 5

Lymphocyte Lineage	hFc γ RIII	hFc γ RII	hFc γ RII/hFc γ RIII
B cells		100	
NK cells	30	—	—
Macrophages	<1	55	26
Neutrophils	—		100

[0141] As shown in this Example, genetically modified mice (both heterozygote and homozygote genotypes) generated in accordance with Example 5 expressed human Fc γ RIIIA on NK cells and macrophages, human Fc γ RIIIB on neutrophils, human Fc γ RIIA on neutrophils and macroph-

ages, human Fc γ RIIB on B cells, and human Fc γ RIIC on NK cells. The expression pattern of human Fc γ R genes shown in this Example is consistent with the expression patterns of these genes in human accessory cells.

Example 8

ADCC in Humanized Fc γ R Mice

[0142] Splenocytes isolated from Fc γ R gene deficient (i.e. knockout), Fc γ RIIIA/Fc γ RIIA (homozygotes), Fc γ RIIIA/Fc γ RIIIB/Fc γ RIIA/Fc γ RIIB/Fc γ RIIC (homozygotes) and wild type mice were analyzed for their ability to perform ADCC in a cell-killing assay (as described above in Example 2).

[0143] Briefly, cell populations were isolated and separated using MACS[®] Technology (Miltenyi Biotec). Briefly, T and B cell depleted splenocytes were cultured for two weeks in the presence of mouse IL-2 (500 U/mL). The resulting expanded NK cells were used as effector cells in the ADCC assays at a ratio of 50:1 (NK:Raji). Raji cells were coated with 10 μ g/mL of Ab 168 or Ab 735 (as described above in Example 3). Results are shown in Table 6.

TABLE 6

NK Cell Genotype	% ADCC	
	10 μ g/mL Ab 168	10 μ g/mL Ab 735
Wild Type	89	72
Mouse Fc γ R KO	13	14
Human Fc γ RIIIA-IIA HO	78	85
Human Fc γ RIIIA-IIIB-IIA-IIIB-IIIC HO	81	59

[0144] The present invention is not to be limited in scope by the specific embodiments describe herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and the accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 38

<210> SEQ ID NO 1
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 1

accaggatat gacctgtaga g

21

<210> SEQ ID NO 2
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 2

-continued

gtccatgggt aagtagaaac a 21

<210> SEQ ID NO 3
<211> LENGTH: 65
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 3

atgcgagctc atgcatctat gtcgggtgcg gagaaagagg taatgcattc ttgcccaata 60
cttac 65

<210> SEQ ID NO 4
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 4

actcatggag cctcaacagg a 21

<210> SEQ ID NO 5
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 5

atcctgagta tactatgaca aga 23

<210> SEQ ID NO 6
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 6

actagtgaga cgtgctactt c 21

<210> SEQ ID NO 7
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 7

ctcccactca tgatctatag a 21

<210> SEQ ID NO 8
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 8

tggagcctca acaggactcc a 21

-continued

<210> SEQ ID NO 9
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 9

gtccatgggt aagtagaaac attcgtacc ttaggaccgt ta 42

<210> SEQ ID NO 10
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 10

cggtgcgga gaaagaggta atgcattctt gcccaatact ta 42

<210> SEQ ID NO 11
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 11

gccagccaca aaggagataa tc 22

<210> SEQ ID NO 12
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 12

gcaacattta ggacaactcg gg 22

<210> SEQ ID NO 13
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 13

gatttcctaa ccacctacc c 21

<210> SEQ ID NO 14
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 14

tcttttccaa tggcagttg 19

<210> SEQ ID NO 15
<211> LENGTH: 24
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 15

cccaggtaag tcgtgatgaa acag 24

<210> SEQ ID NO 16
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 16

ctcccactca tgatctatag a 21

<210> SEQ ID NO 17
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 17

tggagcctca acaggactcc a 21

<210> SEQ ID NO 18
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 18

cacacatctc ctggtgactt g 21

<210> SEQ ID NO 19
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 19

caactgccat tggaaaagac tcgagtgcca tttcattacc tc 42

<210> SEQ ID NO 20
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 20

taaaccgcg gtggagctcg ccagccacaa aggagataat ca 42

<210> SEQ ID NO 21
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

-continued

<400> SEQUENCE: 21
 ccatgggtaa gtagaaactc tagacccccg ggctcgataa ct 42

<210> SEQ ID NO 22
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 22
 accaggatat gacctgtaga g 21

<210> SEQ ID NO 23
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 23
 gtttctactt acccatggac 20

<210> SEQ ID NO 24
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 24
 aaatacacac tgccacagac ag 22

<210> SEQ ID NO 25
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 25
 cctettttgt gagtttcctg tg 22

<210> SEQ ID NO 26
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 26
 gagtatacta tgacaagagc atc 23

<210> SEQ ID NO 27
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 27
 actagtgaga cgtgctactt c 21

-continued

<210> SEQ ID NO 28
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

 <400> SEQUENCE: 28

 ctccactca tgatctatag a 21

<210> SEQ ID NO 29
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

 <400> SEQUENCE: 29

 ctttttatgg tcccacaatc ag 22

<210> SEQ ID NO 30
 <211> LENGTH: 42
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

 <400> SEQUENCE: 30

 gtccatgggt aagtagaaac attcgtacc ttaggaccgt ta 42

<210> SEQ ID NO 31
 <211> LENGTH: 42
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

 <400> SEQUENCE: 31

 gcttatcgat accgtcgaca aatacacact gccacagaca gg 42

<210> SEQ ID NO 32
 <211> LENGTH: 317
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

 <400> SEQUENCE: 32

 Met Thr Met Glu Thr Gln Met Ser Gln Asn Val Cys Pro Arg Asn Leu
 1 5 10 15

 Trp Leu Leu Gln Pro Leu Thr Val Leu Leu Leu Ala Ser Ala Asp
 20 25 30

 Ser Gln Ala Ala Ala Pro Pro Lys Ala Val Leu Lys Leu Glu Pro Pro
 35 40 45

 Trp Ile Asn Val Leu Gln Glu Asp Ser Val Thr Leu Thr Cys Gln Gly
 50 55 60

 Ala Arg Ser Pro Glu Ser Asp Ser Ile Gln Trp Phe His Asn Gly Asn
 65 70 75 80

 Leu Ile Pro Thr His Thr Gln Pro Ser Tyr Arg Phe Lys Ala Asn Asn
 85 90 95

-continued

Asn Asp Ser Gly Glu Tyr Thr Cys Gln Thr Gly Gln Thr Ser Leu Ser
 100 105 110
 Asp Pro Val His Leu Thr Val Leu Ser Glu Trp Leu Val Leu Gln Thr
 115 120 125
 Pro His Leu Glu Phe Gln Glu Gly Glu Thr Ile Met Leu Arg Cys His
 130 135 140
 Ser Trp Lys Asp Lys Pro Leu Val Lys Val Thr Phe Phe Gln Asn Gly
 145 150 155 160
 Lys Ser Gln Lys Phe Ser His Leu Asp Pro Thr Phe Ser Ile Pro Gln
 165 170 175
 Ala Asn His Ser His Ser Gly Asp Tyr His Cys Thr Gly Asn Ile Gly
 180 185 190
 Tyr Thr Leu Phe Ser Ser Lys Pro Val Thr Ile Thr Val Gln Val Pro
 195 200 205
 Ser Met Gly Ser Ser Ser Pro Met Gly Ile Ile Val Ala Val Val Ile
 210 215 220
 Ala Thr Ala Val Ala Ala Ile Val Ala Ala Val Val Ala Leu Ile Tyr
 225 230 235 240
 Cys Arg Lys Lys Arg Ile Ser Ala Asn Ser Thr Asp Pro Val Lys Ala
 245 250 255
 Ala Gln Phe Glu Pro Pro Gly Arg Gln Met Ile Ala Ile Arg Lys Arg
 260 265 270
 Gln Leu Glu Glu Thr Asn Asn Asp Tyr Glu Thr Ala Asp Gly Gly Tyr
 275 280 285
 Met Thr Leu Asn Pro Arg Ala Pro Thr Asp Asp Asp Lys Asn Ile Tyr
 290 295 300
 Leu Thr Leu Pro Pro Asn Asp His Val Asn Ser Asn Asn
 305 310 315

<210> SEQ ID NO 33
 <211> LENGTH: 303
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 33

Met Gly Ile Leu Ser Phe Leu Pro Val Leu Ala Thr Glu Ser Asp Trp
 1 5 10 15
 Ala Asp Cys Lys Ser Pro Gln Pro Trp Gly His Met Leu Leu Trp Thr
 20 25 30
 Ala Val Leu Phe Leu Ala Ala Pro Pro Lys Ala Val Leu Lys Leu Glu
 35 40 45
 Pro Gln Trp Ile Asn Val Leu Gln Glu Asp Ser Val Thr Leu Thr Cys
 50 55 60
 Arg Gly Thr His Ser Pro Glu Ser Asp Ser Ile Gln Trp Phe His Asn
 65 70 75 80
 Gly Asn Leu Ile Pro Thr His Thr Gln Pro Ser Tyr Arg Phe Lys Ala
 85 90 95
 Asn Asn Asn Asp Ser Gly Glu Tyr Thr Cys Gln Thr Gly Gln Thr Ser
 100 105 110
 Leu Ser Asp Pro Val His Leu Thr Val Leu Ser Glu Trp Leu Val Leu
 115 120 125

-continued

Gln Thr Pro His Leu Glu Phe Gln Glu Gly Glu Thr Ile Val Leu Arg
130 135 140

Cys His Ser Trp Lys Asp Lys Pro Leu Val Lys Val Thr Phe Phe Gln
145 150 155 160

Asn Gly Lys Ser Lys Lys Phe Ser Arg Ser Asp Pro Asn Phe Ser Ile
165 170 175

Pro Gln Ala Asn His Ser His Ser Gly Asp Tyr His Cys Thr Gly Asn
180 185 190

Ile Gly Tyr Thr Leu Tyr Ser Ser Lys Pro Val Thr Ile Thr Val Gln
195 200 205

Ala Pro Ser Ser Ser Pro Met Gly Ile Ile Val Ala Val Val Thr Gly
210 215 220

Ile Ala Val Ala Ala Ile Val Ala Ala Val Val Ala Leu Ile Tyr Cys
225 230 235 240

Arg Lys Lys Arg Ile Ser Ala Leu Pro Gly Tyr Pro Glu Cys Arg Glu
245 250 255

Met Gly Glu Thr Leu Pro Glu Lys Pro Ala Asn Pro Thr Asn Pro Asp
260 265 270

Glu Ala Asp Lys Val Gly Ala Glu Asn Thr Ile Thr Tyr Ser Leu Leu
275 280 285

Met His Pro Asp Ala Leu Glu Glu Pro Asp Asp Gln Asn Arg Ile
290 295 300

<210> SEQ ID NO 34
 <211> LENGTH: 323
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 34

Met Gly Ile Leu Ser Phe Leu Pro Val Leu Ala Thr Glu Ser Asp Trp
1 5 10 15

Ala Asp Cys Lys Ser Pro Gln Pro Trp Gly His Met Leu Leu Trp Thr
20 25 30

Ala Val Leu Phe Leu Ala Pro Val Ala Gly Thr Pro Ala Ala Pro Pro
35 40 45

Lys Ala Val Leu Lys Leu Glu Pro Gln Trp Ile Asn Val Leu Gln Glu
50 55 60

Asp Ser Val Thr Leu Thr Cys Arg Gly Thr His Ser Pro Glu Ser Asp
65 70 75 80

Ser Ile Pro Trp Phe His Asn Gly Asn Leu Ile Pro Thr His Thr Gln
85 90 95

Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn Asp Ser Gly Glu Tyr Thr
100 105 110

Cys Gln Thr Gly Gln Thr Ser Leu Ser Asp Pro Val His Leu Thr Val
115 120 125

Leu Ser Glu Trp Leu Val Leu Gln Thr Pro His Leu Glu Phe Gln Glu
130 135 140

Gly Glu Thr Ile Val Leu Arg Cys His Ser Trp Lys Asp Lys Pro Leu
145 150 155 160

Val Lys Val Thr Phe Phe Gln Asn Gly Lys Ser Lys Lys Phe Ser Arg
165 170 175

-continued

Ser Asp Pro Asn Phe Ser Ile Pro Gln Ala Asn His Ser His Ser Gly
 180 185 190

Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr Thr Leu Tyr Ser Ser Lys
 195 200 205

Pro Val Thr Ile Thr Val Gln Ala Pro Ser Ser Ser Pro Met Gly Ile
 210 215 220

Ile Val Ala Val Val Thr Gly Ile Ala Val Ala Ala Ile Val Ala Ala
 225 230 235 240

Val Val Ala Leu Ile Tyr Cys Arg Lys Lys Arg Ile Ser Ala Asn Ser
 245 250 255

Thr Asp Pro Val Lys Ala Ala Gln Phe Glu Pro Pro Gly Arg Gln Met
 260 265 270

Ile Ala Ile Arg Lys Arg Gln Pro Glu Glu Thr Asn Asn Asp Tyr Glu
 275 280 285

Thr Ala Asp Gly Gly Tyr Met Thr Leu Asn Pro Arg Ala Pro Thr Asp
 290 295 300

Asp Asp Lys Asn Ile Tyr Leu Thr Leu Pro Pro Asn Asp His Val Asn
 305 310 315 320

Ser Asn Asn

<210> SEQ ID NO 35
 <211> LENGTH: 290
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 35

Met Gly Gly Gly Ala Gly Glu Arg Leu Phe Thr Ser Ser Cys Leu Val
 1 5 10 15

Gly Leu Val Pro Leu Gly Leu Arg Ile Ser Leu Val Thr Cys Pro Leu
 20 25 30

Gln Cys Gly Ile Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu
 35 40 45

Leu Val Ser Ala Gly Met Arg Thr Glu Asp Leu Pro Lys Ala Val Val
 50 55 60

Phe Leu Glu Pro Gln Trp Tyr Arg Val Leu Glu Lys Asp Ser Val Thr
 65 70 75 80

Leu Lys Cys Gln Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Gln Trp
 85 90 95

Phe His Asn Glu Ser Leu Ile Ser Ser Gln Ala Ser Ser Tyr Phe Ile
 100 105 110

Asp Ala Ala Thr Val Asp Asp Ser Gly Glu Tyr Arg Cys Gln Thr Asn
 115 120 125

Leu Ser Thr Leu Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp
 130 135 140

Leu Leu Leu Gln Ala Pro Arg Trp Val Phe Lys Glu Glu Asp Pro Ile
 145 150 155 160

His Leu Arg Cys His Ser Trp Lys Asn Thr Ala Leu His Lys Val Thr
 165 170 175

Tyr Leu Gln Asn Gly Lys Gly Arg Lys Tyr Phe His His Asn Ser Asp
 180 185 190

Phe Tyr Ile Pro Lys Ala Thr Leu Lys Asp Ser Gly Ser Tyr Phe Cys

-continued

```

<210> SEQ ID NO 37
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 37

atcggggata gagatgtttg cgcgatcgc ggtaccgggc                40

<210> SEQ ID NO 38
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 38

ttatacgaag ttataccggt gcattcttgc ccaatactta                40

```

We claim:

1. A mouse comprising a genetic modification, wherein the genetic modification comprises a replacement of endogenous low affinity Fc γ R genes with at least two low affinity human Fc γ R genes placed at the endogenous mouse low affinity Fc γ R gene locus, and wherein the mouse comprises a functional FcR γ -chain.

2. The mouse of claim 1, wherein the at least two low affinity human Fc γ R genes are selected from Fc γ RIIA, Fc γ RIIB, Fc γ RIIC, Fc γ RIIA and Fc γ RIIB.

3. The mouse of claim 1, wherein the at least two low affinity human Fc γ R genes are Fc γ RIIA and Fc γ RIIA.

4. The mouse of claim 3, wherein the human Fc γ RIIA is expressed on NK cells of the mouse.

5. The mouse of claim 1, wherein the at least two low affinity human Fc γ R genes are Fc γ RIIB, Fc γ RIIC, and Fc γ RIIB.

6. The mouse of claim 3, wherein the Fc γ RIIA and Fc γ RIIA genes comprise allelic variants or polymorphisms.

7. The mouse of claim 5, wherein the mouse comprises human low affinity Fc γ R allelic variants, and the allelic variants are selected from the group consisting of (a) Fc γ RIIA allelic variant 131H, (b) Fc γ RIIA allelic variant 158V, (c) Fc γ RIIB allelic variant 232I, (d) Fc γ RIIB allelic variant characterized by a neutrophil antigen 2 (NA2) allele, and, (e) a combination thereof.

8. A genetically modified mouse, wherein the genetic modification comprises a replacement of endogenous mouse genes encoding Fc γ RIIB, Fc γ RIII, and Fc γ RIV α -subunits with a gene encoding a human Fc γ RIIA α -subunit, wherein the mouse expresses a human Fc γ RIIA protein associated with an endogenous mouse FcR γ -subunit, wherein the human Fc γ RIIA protein is expressed on the surface of a mouse NK cell.

9. The genetically modified mouse of claim 8, wherein mouse NK cell is a circulating NK cell in the blood of the mouse, and the Fc γ RIIA protein expressed on the surface of the mouse NK cell binds an immunoadhesin or an antibody that comprises a human Fc or a modified human Fc, wherein

the immunoadhesin or antibody specifically binds a target cell in the mouse, and binding of the immunoadhesin or the antibody to the Fc γ RIIA of the NK cell mediates NK cell killing of the target cell.

10. The genetically modified mouse of claim 9, wherein the target cell is infected with a human pathogen, and the immunoadhesin or antibody specifically binds an epitope of the human pathogen.

11. The genetically modified mouse of claim 9, wherein the target cell is a tumor cell, and the immunoadhesin or antibody specifically binds an epitope of the tumor cell.

12. A method for measuring therapeutic-dependent cell killing, comprising:

- (a) administering to a genetically modified mouse a therapeutic agent that specifically binds a target cell that is not an NK cell in the mouse, wherein the therapeutic agent comprises a human Fc;
- (b) measuring a level of NK-mediated target cell killing in the mouse or in a tissue sample of the mouse; and,
- (c) determining the amount of target cell killing that is mediated by the therapeutic agent, thereby measuring therapeutic-dependent cell killing;

wherein the genetic modification comprises a deletion of a gene encoding the α -subunit of mouse Fc γ RIIB, a deletion of a gene encoding the α -subunit of mouse Fc γ RIII, and a deletion of a gene encoding the α -subunit of mouse Fc γ RIV, and wherein the mouse comprises a gene encoding an α -subunit of a human Fc γ RIIA at the endogenous mouse α -subunit locus and wherein the mouse expresses a functional FcR γ -chain subunit.

13. The method of claim 12, wherein the mouse further comprises at the endogenous mouse α -subunit locus a gene encoding a human Fc γ R α -subunit selected from an Fc γ RIIA α -subunit, Fc γ RIIB α -subunit, an Fc γ RIIC α -subunit, an Fc γ RIIB α -subunit, and a combination thereof.

14. The method of claim 12, wherein the target cell is a human pathogen, and the therapeutic specifically binds an epitope of the human pathogen.

15. The method of claim **12**, wherein the target cell is a mouse cell infected by a human pathogen, and the therapeutic specifically binds an epitope of the human pathogen.

16. The method of claim **12**, wherein the target cell is a human cell infected by a human pathogen, and the therapeutic specifically binds an epitope of the human pathogen.

17. The method of claim **12**, wherein the target cell is a human tumor cell, and the therapeutic specifically binds an epitope of the human tumor cell.

18. The method of claim **12**, wherein the tissue sample is a blood sample.

19. The method of claim **12**, wherein the therapeutic is an antibody.

20. The method of claim **17**, wherein the antibody is a human antibody.

* * * * *

专利名称(译)	人源化FcγR小鼠		
公开(公告)号	US20110154512A1	公开(公告)日	2011-06-23
申请号	US12/971080	申请日	2010-12-17
[标]申请(专利权)人(译)	再生元医药公司		
申请(专利权)人(译)	REGENERON制药公司.		
当前申请(专利权)人(译)	REGENERON制药公司.		
[标]发明人	MACDONALD LYNN TU NAXIN GURER CAGAN STEVENS SEAN MURPHY ANDREW J		
发明人	MACDONALD, LYNN TU, NAXIN GURER, CAGAN STEVENS, SEAN MURPHY, ANDREW J.		
IPC分类号	A01K67/027 G01N33/00 C12Q1/18 G01N33/53		
CPC分类号	A01K67/0276 A01K67/0278 A01K2207/15 A01K2217/072 A01K2217/075 G01N33/5014 A01K2227/105 A01K2267/0387 C07K14/70535 A61K49/0008 A01K2217/15 A01K2267/03 A61K38/17 A61K38/18 C07K14/435 C07K14/475 C07K14/705 C07K16/00 C07K16/18 C07K16/22 C07K16/24 C07K16/28 C07K16/30		
优先权	61/288562 2009-12-21 US		
其他公开文献	US8658154		
外部链接	Espacenet USPTO		

摘要(译)

提供了遗传修饰的非人动物以及用于制备和使用它们的方法和组合物，其中遗传修饰包括内源低亲和力FcγR基因座的缺失，并且其中所述小鼠能够表达功能性FcRγ链。描述了遗传修饰的小鼠，包括从内源FcγR基因座表达低亲和力人FcγR基因的小鼠，并且其中小鼠包含功能性FcRγ链。提供了在宿主免疫系统的辅助细胞上表达多达五种低亲和力人FcγR基因的遗传修饰小鼠。

