



US 20140303021A1

(19) **United States**

(12) **Patent Application Publication**
LaBrie et al.

(10) **Pub. No.: US 2014/0303021 A1**
(43) **Pub. Date: Oct. 9, 2014**

(54) **METHODS AND DEVICES FOR DETECTING OBSTRUCTIVE UROPATHY AND ASSOCIATED DISORDERS**

(60) Provisional application No. 61/327,389, filed on Apr. 23, 2010, provisional application No. 61/232,091, filed on Aug. 7, 2009.

(71) Applicant: **Rules-Based Medicine, Inc.**, Austin, TX (US)

Publication Classification

(72) Inventors: **Samuel T. LaBrie**, Austin, TX (US);
James P. Mapes, Austin, TX (US);
Michael D. Spain, Austin, TX (US);
Ralph L. McDade, Austin, TX (US);
Dominic P. Eisinger, Lake Placid, NY (US);
Karri L. Ballard, Austin, TX (US)

(51) **Int. Cl.**
G01N 33/566 (2006.01)
G01N 33/53 (2006.01)
(52) **U.S. Cl.**
CPC **G01N 33/566** (2013.01); **G01N 33/5302** (2013.01); **G01N 2800/34** (2013.01)
USPC **506/9**; 435/7.92; 436/501; 422/69; 435/287.2

(21) Appl. No.: **14/243,659**

(57) **ABSTRACT**

(22) Filed: **Apr. 2, 2014**

Methods and devices for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder in a mammal are described. In particular, methods and devices for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder using measured concentrations of a combination of three or more analytes in a test sample taken from the mammal are described.

Related U.S. Application Data

(63) Continuation of application No. 12/852,236, filed on Aug. 6, 2010, now Pat. No. 8,735,080.





FIG. 1

FIG. 2A








FIG. 2D



ABNORMAL vs. NORMAL kidney plasma



relative importance by random forest

FIG. 3




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 6A




FIG. 6B






FIG. 6C



relative importance by random forest

FIG. 7A



relative importance by random forest

FIG. 7B




FIG. 7C




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 12A




FIG. 12B




FIG. 12C




FIG. 13A




FIG. 13B




FIG. 13C

METHODS AND DEVICES FOR DETECTING OBSTRUCTIVE UROPATHY AND ASSOCIATED DISORDERS

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the priority of U.S. provisional application Ser. No. 61/327,389, filed Apr. 23, 2010, and U.S. provisional application Ser. No. 61/232,091, filed Aug. 7, 2009, each of which is hereby incorporated by reference in its entirety and is related to U.S. patent application Ser. Nos. [Not Yet Assigned], entitled Computer Methods and Devices for Detecting Kidney Damage, Methods and Devices for Detecting Glomerulonephritis and Associated Disorders, Methods and Devices for Detecting Kidney Damage, Devices for Detecting Renal Disorders, Methods and Devices for Detecting Kidney Transplant Rejection, Methods and Devices for Detecting Diabetic Nephropathy and Associated Disorders, Attorney Docket Nos. 060075-, filed on the same date as this application, the entire contents of which are incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The invention encompasses methods and devices for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder in a mammal. In particular, the present invention provides methods and devices for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder using measured concentrations of a combination of three or more analytes in a test sample taken from the mammal.

BACKGROUND OF THE INVENTION

[0003] The urinary system, in particular the kidneys, perform several critical functions such as maintaining electrolyte balance and eliminating toxins from the bloodstream. In the human body, the pair of kidneys together process roughly 20% of the total cardiac output, amounting to about 1 L/min in a 70-kg adult male. Because compounds in circulation are concentrated in the kidney up to 1000-fold relative to the plasma concentration, the kidney is especially vulnerable to injury due to exposure to toxic compounds.

[0004] An obstruction causing the flow of urine to back up into the kidneys can cause permanent damage to the kidneys, and may result in renal failure. Existing diagnostic tests such as BUN and serum creatine tests typically detect only advanced stages of kidney damage. Other diagnostic tests such as kidney tissue biopsies or CAT scans have the advantage of enhanced sensitivity to earlier stages of kidney damage, but these tests are also generally costly, slow, and/or invasive.

[0005] A need exists in the art for a fast, simple, reliable, and sensitive method of detecting obstructive uropathy or an associated disorder. In a clinical setting, the early detection of kidney damage would help medical practitioners to diagnose and treat kidney damage more quickly and effectively.

SUMMARY OF THE INVENTION

[0006] The present invention provides methods and devices for diagnosing, monitoring, or determining a renal disorder in a mammal. In particular, the present invention provides methods and devices for diagnosing, monitoring, or determining a

renal disorder using measured concentrations of a combination of three or more analytes in a test sample taken from the mammal.

[0007] One aspect of the invention encompasses a method for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder in a mammal. The method typically comprises providing a test sample comprising a sample of bodily fluid taken from the mammal. Then, the method comprises determining a combination of sample concentrations for three or more sample analytes in the test sample, wherein the sample analytes are selected from the group consisting of alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GST-alpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF. The combination of sample concentrations may be compared to a data set comprising at least one entry, wherein each entry of the data set comprises a list comprising three or more minimum diagnostic concentrations indicative of obstructive uropathy or an associated disorder. Each minimum diagnostic concentration comprises a maximum of a range of analyte concentrations for a healthy mammal. Next, the method comprises determining a matching entry of the dataset in which all minimum diagnostic concentrations are less than the corresponding sample concentrations and identifying an indicated disorder comprising the particular disorder of the matching entry.

[0008] Another aspect of the invention encompasses a method for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder in a mammal. The method generally comprises providing a test sample comprising a sample of bodily fluid taken from the mammal. Then the method comprises determining the concentrations of three or more sample analytes in a panel of biomarkers in the test sample, wherein the sample analytes are selected from the group consisting of alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GST-alpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF. Diagnostic analytes are identified in the test sample, wherein the diagnostic analytes are the sample analytes whose concentrations are statistically different from concentrations found in a control group of humans who do not suffer from obstructive uropathy or an associated disorder. The combination of diagnostic analytes is compared to a dataset comprising at least one entry, wherein each entry of the dataset comprises a combination of three or more diagnostic analytes reflective of obstructive uropathy or an associated disorder. The particular disorder having the combination of diagnostic analytes that essentially match the combination of sample analytes is then identified.

[0009] Yet another aspect of the invention encompasses a computer readable media encoded with an application comprising modules executable by a processor and configured to diagnose, monitor, or determine obstructive uropathy or an associated disorder in a mammal. The application usually comprises an analyte input module to receive three or more sample analyte concentrations selected from the group consisting of alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GST-alpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF. The media also comprises a comparison module to compare each sample analyte concentration to an entry of an obstructive uropathy or an associated disorder database, wherein each entry comprises a list of minimum diagnostic concentrations reflective of obstructive uropathy

or an associated disorder. The media further comprises an analysis module to determine a most likely disorder by combining the particular disorders identified by the comparison module for all of the sample analyte concentrations.

[0010] Still another aspect of the invention encompasses a system for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder in a mammal. The system typically comprises, in part, a database to store a plurality of obstructive uropathy or an associated disorder database entries. The system also comprises a processing device comprising a disorder diagnosis application comprising modules executable by the processing device. The disorder diagnosis application comprises an analyte input module to receive three or more sample analyte concentrations selected from the group consisting of alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GST-alpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF; a comparison module to compare each sample analyte concentration to an entry of the obstructive uropathy or an associated disorder database, wherein each entry comprises a list of minimum diagnostic concentrations reflective of obstructive uropathy or an associated disorder; and an analysis module to determine a most likely disorder by combining the particular disorders identified by the comparison module for all of the sample analyte concentrations.

[0011] An additional aspect of the invention encompasses a method for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder in a mammal. The method usually comprises providing an analyte concentration measurement device comprising three or more detection antibodies. Each detection antibody comprises an antibody coupled to an indicator, wherein the antigenic determinants of the antibodies are sample analytes associated with obstructive uropathy or an associated disorder. The sample analytes are generally selected from the group consisting of alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GST-alpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF. The method next comprises providing a test sample comprising three or more sample analytes and a bodily fluid taken from the mammal. The test sample is contacted with the detection antibodies and the detection antibodies are allowed to bind to the sample analytes. The concentrations of the sample analytes are determined by detecting the indicators of the detection antibodies bound to the sample analytes in the test sample. The concentrations of each sample analyte correspond to a corresponding minimum diagnostic concentration reflective of obstructive uropathy or an associated disorder.

[0012] Other aspects and iterations of the invention are described in more detail below.

DESCRIPTION OF FIGURES

[0013] FIG. 1 shows the four different disease groups from which samples were analyzed, and a plot of two different estimations on eGFR outlining the distribution within each group.

[0014] FIG. 2 is a number of scatter plots of results on selected proteins in urine and plasma. The various groups are indicated as follows—control: blue, AA: red, DN: green, GN: yellow, OU: orange. (A) A1M in plasma, (B) cystatin C in plasma, (C) B2M in urine, (D) cystatin C in urine.

[0015] FIG. 3 depicts the multivariate analysis of the disease groups and their respective matched controls using plasma results. Relative importance shown using the random forest model.

[0016] FIG. 4 depicts three graphs showing the mean AUROC and its standard deviation (A) for plasma samples, and mean error rates (B) and mean AUROC (C) from urine samples for each classification method used to distinguish disease samples vs. normal samples. Disease encompasses analgesic abuse (AA), glomerulonephritis (GN), obstructive uropathy (OU), and diabetic nephropathy (DN). Normal=NL.

[0017] FIG. 5 depicts three graphs showing the average importance of analytes and clinical variables from 100 bootstrap runs measured by random forest (A and B) or boosting (C) to distinguish disease (AA+GN+ON+DN) samples vs. normal samples from plasma (A) and urine (B and C).

[0018] FIG. 6 depicts three graphs showing the mean AUROC and its standard deviation (A) for plasma samples, and mean error rates (B) and mean AUROC (C) from urine samples for each classification method used to distinguish obstructive uropathy samples vs. normal samples. Abbreviations as in FIG. 4.

[0019] FIG. 7 depicts three graphs showing the average importance of analytes and clinical variables from 100 bootstrap runs measured by random forest (A and B) or boosting (C) to distinguish obstructive uropathy samples vs. normal samples from plasma (A) and urine (B and C).

[0020] FIG. 8 depicts three graphs showing the mean AUROC and its standard deviation (A) for plasma samples, and mean error rates (B) and mean AUROC (C) from urine samples for each classification method used to distinguish analgesic abuse samples vs. obstructive uropathy samples. Abbreviations as in FIG. 4.

[0021] FIG. 9 depicts three graphs showing the average importance of analytes and clinical variables from 100 bootstrap runs measured by random forest (A and B) or boosting (C) to distinguish analgesic abuse samples vs. obstructive uropathy samples from plasma (A) and urine (B and C).

[0022] FIG. 10 depicts three graphs showing the mean AUROC and its standard deviation (A) for plasma samples, and mean error rates (B) and mean AUROC (C) from urine samples for each classification method used to distinguish obstructive uropathy samples vs. glomerulonephritis samples. Abbreviations as in FIG. 4.

[0023] FIG. 11 depicts three graphs showing the average importance of analytes and clinical variables from 100 bootstrap runs measured by random forest (A and B) or boosting (C) to distinguish obstructive uropathy samples vs. glomerulonephritis samples from plasma (A) and urine (B and C).

[0024] FIG. 12 depicts three graphs showing the mean AUROC and its standard deviation (A) for plasma samples, and mean error rates (B) and mean AUROC (C) from urine samples for each classification method used to distinguish diabetic nephropathy samples vs. obstructive uropathy samples. Abbreviations as in FIG. 4.

[0025] FIG. 13 depicts three graphs showing the average importance of analytes and clinical variables from 100 bootstrap runs measured by random forest (A and B) or boosting (C) to distinguish diabetic nephropathy samples vs. obstructive uropathy samples from plasma (A) and urine (B and C).

DETAILED DESCRIPTION OF THE INVENTION

[0026] It has been discovered that a multiplexed panel of three, six, or preferably 16, biomarkers may be used to detect

obstructive uropathy and associated disorders. As used herein, the term "obstructive uropathy" refers to a structural or functional hindrance of normal urine flow. The term may encompass chronic unilateral obstructive uropathy, chronic bilateral obstructive uropathy, acute unilateral obstructive uropathy, or acute bilateral obstructive uropathy. Additionally, the present invention encompasses biomarkers that may be used to detect a disorder associated with obstructive uropathy. As used herein, the phrase "a disorder associated with obstructive uropathy" refers to a disorder that stems from a structural or functional hindrance of normal urine flow. For instance, non-limiting examples of associated disorders may include hydronephrosis and obstructive nephropathy.

[0027] The biomarkers included in a multiplexed panel of the invention are analytes known in the art that may be detected in the urine, serum, plasma and other bodily fluids of mammals. As such, the analytes of the multiplexed panel may be readily extracted from the mammal in a test sample of bodily fluid. The concentrations of the analytes within the test sample may be measured using known analytical techniques such as a multiplexed antibody-based immunological assay. The combination of concentrations of the analytes in the test sample may be compared to empirically determined combinations of minimum diagnostic concentrations and combinations of diagnostic concentration ranges associated with healthy kidney function or obstructive uropathy or an associated disorder to determine whether obstructive uropathy is indicated in the mammal.

[0028] One embodiment of the present invention provides a method for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder in a mammal that includes determining the presence or concentration of a combination of three or more sample analytes in a test sample containing the bodily fluid of the mammal. The measured concentrations of the combination of sample analytes is compared to the entries of a dataset in which each entry contains the minimum diagnostic concentrations of a combination of three or more analytes reflective of obstructive uropathy or an associated disorder. Other embodiments provide computer-readable media encoded with applications containing executable modules, systems that include databases and processing devices containing executable modules configured to diagnose, monitor, or determine a renal disorder in a mammal. Still other embodiments provide antibody-based devices for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder in a mammal.

[0029] The analytes used as biomarkers in the multiplexed assay, methods of diagnosing, monitoring, or determining a renal disorder using measurements of the analytes, systems and applications used to analyze the multiplexed assay measurements, and antibody-based devices used to measure the analytes are described in detail below.

I. Analytes in Multiplexed Assay

[0030] One embodiment of the invention measures the concentrations of three, six, or preferably sixteen, biomarker analytes within a test sample taken from a mammal and compares the measured analyte concentrations to minimum diagnostic concentrations to diagnose, monitor, or determine obstructive uropathy or an associated disorder in a mammal. In this aspect, the biomarker analytes are known in the art to occur in the urine, plasma, serum and other bodily fluids of mammals. The biomarker analytes are proteins that have known and documented associations with early renal damage

in humans. As defined herein, the biomarker analytes include but are not limited to alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GST-alpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF. A description of each biomarker analyte is given below.

(a) Alpha-1 Microglobulin (A1M)

[0031] Alpha-1 microglobulin (A1M, Swiss-Prot Accession Number P02760) is a 26 kDa glycoprotein synthesized by the liver and reabsorbed in the proximal tubules. Elevated levels of A1M in human urine are indicative of glomerulotubular dysfunction. A1M is a member of the lipocalin superfamily and is found in all tissues. Alpha-1-microglobulin exists in blood in both a free form and complexed with immunoglobulin A (IgA) and heme. Half of plasma A1M exists in a free form, and the remainder exists in complexes with other molecules including prothrombin, albumin, immunoglobulin A and heme. Nearly all of the free A1M in human urine is reabsorbed by the megalin receptor in proximal tubular cells, where it is then catabolized. Small amounts of A1M are excreted in the urine of healthy humans. Increased A1M concentrations in human urine may be an early indicator of renal damage, primarily in the proximal tubule.

(b) Beta-2 Microglobulin (B2M)

[0032] Beta-2 microglobulin (B2M, Swiss-Prot Accession Number P61769) is a protein found on the surfaces of all nucleated cells and is shed into the blood, particularly by tumor cells and lymphocytes. Due to its small size, B2M passes through the glomerular membrane, but normally less than 1% is excreted due to reabsorption of B2M in the proximal tubules of the kidney. Therefore, high plasma levels of B2M occur as a result of renal failure, inflammation, and neoplasms, especially those associated with B-lymphocytes.

(c) Calbindin

[0033] Calbindin (Calbindin D-28K, Swiss-Prot Accession Number P05937) is a Ca-binding protein belonging to the troponin C superfamily. It is expressed in the kidney, pancreatic islets, and brain. Calbindin is found predominantly in subpopulations of central and peripheral nervous system neurons, in certain epithelial cells involved in Ca²⁺ transport such as distal tubular cells and cortical collecting tubules of the kidney, and in enteric neuroendocrine cells.

(d) Clusterin

[0034] Clusterin (Swiss-Prot Accession Number P10909) is a highly conserved protein that has been identified independently by many different laboratories and named SGP2, S35-S45, apolipoprotein J, SP-40, 40, ADHC-9, gp80, GPIII, and testosterone-repressed prostate message (TRPM-2). An increase in clusterin levels has been consistently detected in apoptotic heart, brain, lung, liver, kidney, pancreas, and retinal tissue both in vivo and in vitro, establishing clusterin as a ubiquitous marker of apoptotic cell loss. However, clusterin protein has also been implicated in physiological processes that do not involve apoptosis, including the control of complement-mediated cell lysis, transport of beta-amyloid precursor protein, shuttling of aberrant beta-amyloid across the blood-brain barrier, lipid scavenging, membrane remodeling, cell aggregation, and protection from immune detection and tumor necrosis factor induced cell death.

(e) Connective Tissue Growth Factor (CTGF)

[0035] Connective tissue growth factor (CTGF, Swiss-Prot Accession Number P29279) is a 349-amino acid cysteine-rich polypeptide belonging to the CCN family. In vitro studies have shown that CTGF is mainly involved in extracellular matrix synthesis and fibrosis. Up-regulation of CTGF mRNA and increased CTGF levels have been observed in various diseases, including diabetic nephropathy and cardiomyopathy, fibrotic skin disorders, systemic sclerosis, biliary atresia, liver fibrosis and idiopathic pulmonary fibrosis, and nondiabetic acute and progressive glomerular and tubulointerstitial lesions of the kidney. A recent cross-sectional study found that urinary CTGF may act as a progression promoter in diabetic nephropathy.

(f) Creatinine

[0036] Creatinine is a metabolite of creatine phosphate in muscle tissue, and is typically produced at a relatively constant rate by the body. Creatinine is chiefly filtered out of the blood by the kidneys, though a small amount is actively secreted by the kidneys into the urine. Creatinine levels in blood and urine may be used to estimate the creatinine clearance, which is representative of the overall glomerular filtration rate (GFR), a standard measure of renal function. Variations in creatinine concentrations in the blood and urine, as well as variations in the ratio of urea to creatinine concentration in the blood, are common diagnostic measurements used to assess renal function.

(g) Cystatin C (Cyst C)

[0037] Cystatin C (Cyst C, Swiss-Prot Accession Number P01034) is a 13 kDa protein that is a potent inhibitor of the C1 family of cysteine proteases. It is the most abundant extracellular inhibitor of cysteine proteases in testis, epididymis, prostate, seminal vesicles and many other tissues. Cystatin C, which is normally expressed in vascular wall smooth muscle cells, is severely reduced in both atherosclerotic and aneurysmal aortic lesions.

(h) Glutathione S-Transferase alpha (GST-alpha)

[0038] Glutathione S-transferase alpha (GST-alpha, Swiss-Prot Accession Number P08263) belongs to a family of enzymes that utilize glutathione in reactions contributing to the transformation of a wide range of compounds, including carcinogens, therapeutic drugs, and products of oxidative stress. These enzymes play a key role in the detoxification of such substances.

(i) Kidney Injury Molecule-1 (KIM-1)

[0039] Kidney injury molecule-1 (KIM-1, Swiss-Prot Accession Number Q96D42) is an immunoglobulin superfamily cell-surface protein highly upregulated on the surface of injured kidney epithelial cells. It is also known as TIM-1 (T-cell immunoglobulin mucin domain-1), as it is expressed at low levels by subpopulations of activated T-cells and hepatitis A virus cellular receptor-1 (HAVCR-1). KIM-1 is increased in expression more than any other protein in the injured kidney and is localized predominantly to the apical membrane of the surviving proximal epithelial cells.

(j) Microalbumin

[0040] Albumin is the most abundant plasma protein in humans and other mammals. Albumin is essential for main-

taining the osmotic pressure needed for proper distribution of body fluids between intravascular compartments and body tissues. Healthy, normal kidneys typically filter out albumin from the urine. The presence of albumin in the urine may indicate damage to the kidneys. Albumin in the urine may also occur in patients with long-standing diabetes, especially type 1 diabetes. The amount of albumin eliminated in the urine has been used to differentially diagnose various renal disorders. For example, nephrotic syndrome usually results in the excretion of about 3.0 to 3.5 grams of albumin in human urine every 24 hours. Microalbuminuria, in which less than 300 mg of albumin is eliminated in the urine every 24 hours, may indicate the early stages of diabetic nephropathy.

(k) Neutrophil Gelatinase-Associated Lipocalin (NGAL)

[0041] Neutrophil gelatinase-associated lipocalin (NGAL, Swiss-Prot Accession Number P80188) forms a disulfide bond-linked heterodimer with MMP-9. It mediates an innate immune response to bacterial infection by sequestering iron. Lipocalins interact with many different molecules such as cell surface receptors and proteases, and play a role in a variety of processes such as the progression of cancer and allergic reactions.

(l) Osteopontin (OPN)

[0042] Osteopontin (OPN, Swiss-Prot Accession Number P10451) is a cytokine involved in enhancing production of interferon-gamma and IL-12, and inhibiting the production of IL-10. OPN is essential in the pathway that leads to type I immunity, OPN appears to form an integral part of the mineralized matrix. OPN is synthesized within the kidney and has been detected in human urine at levels that may effectively inhibit calcium oxalate crystallization. Decreased concentrations of OPN have been documented in urine from patients with renal stone disease compared with normal individuals.

(m) Tamm-Horsfall Protein (THP)

[0043] Tamm-Horsfall protein (THP, Swiss-Prot Accession Number P07911), also known as uromodulin, is the most abundant protein present in the urine of healthy subjects and has been shown to decrease in individuals with kidney stones. THP is secreted by the thick ascending limb of the loop of Henley. THP is a monomeric glycoprotein of ~85 kDa with ~30% carbohydrate moiety that is heavily glycosylated. THP may act as a constitutive inhibitor of calcium crystallization in renal fluids.

(n) Tissue Inhibitor of Metalloproteinase-1 (TIMP-1)

[0044] Tissue inhibitor of metalloproteinase-1 (TIMP-1, Swiss-Prot Accession Number P01033) is a major regulator of extracellular matrix synthesis and degradation. A certain balance of MMPs and TIMPs is essential for tumor growth and health. Fibrosis results from an imbalance of fibrogenesis and fibrolysis, highlighting the importance of the role of the inhibition of matrix degradation role in renal disease.

(o) Trefoil Factor 3 (TFF3)

[0045] Trefoil factor 3 (TFF3, Swiss-Prot Accession Number Q07654), also known as intestinal trefoil factor, belongs to a small family of mucin-associated peptides that include TFF1, TFF2, and TFF3. TFF3 exists in a 60-amino acid monomeric form and a 118-amino acid dimeric form. Under

normal conditions TFF3 is expressed by goblet cells of the intestine and the colon. TFF3 expression has also been observed in the human respiratory tract, in human goblet cells and in the human salivary gland. In addition, TFF3 has been detected in the human hypothalamus.

(p) Vascular Endothelial Growth Factor (VEGF)

[0046] Vascular endothelial growth factor (VEGF, Swiss-Prot Accession Number P15692) is an important factor in the pathophysiology of neuronal and other tumors, most likely functioning as a potent promoter of angiogenesis. VEGF may also be involved in regulating blood-brain-barrier functions under normal and pathological conditions. VEGF secreted from the stromal cells may be responsible for the endothelial cell proliferation observed in capillary hemangioblastomas, which are typically composed of abundant microvasculature and primitive angiogenic elements represented by stromal cells.

II. Combinations of Analytes Measured by Multiplexed Assay

[0047] The method for diagnosing, monitoring, or determining a renal disorder involves determining the presence or concentrations of a combination of sample analytes in a test sample. The combinations of sample analytes, as defined herein, are any group of three or more analytes selected from the biomarker analytes, including but not limited to alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GST-alpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF. In one embodiment, the combination of analytes may be selected to provide a group of analytes associated with obstructive uropathy or an associated disorder.

[0048] In one embodiment, the combination of sample analytes may be any three of the biomarker analytes. In other embodiments, the combination of sample analytes may be any four, any five, any six, any seven, any eight, any nine, any ten, any eleven, any twelve, any thirteen, any fourteen, any fifteen, or all sixteen of the sixteen biomarker analytes. In some embodiments, the combination of sample analytes comprises alpha-1 microglobulin, beta-2 microglobulin, cystatin C, KIM-1, THP, and TIMP-1. In another embodiment, the combination of sample analytes may comprise a combination listed in Table A.

TABLE A

alpha-1 microglobulin	beta-2 microglobulin	calbindin
alpha-1 microglobulin	beta-2 microglobulin	clusterin
alpha-1 microglobulin	beta-2 microglobulin	CTGF
alpha-1 microglobulin	beta-2 microglobulin	creatinine
alpha-1 microglobulin	beta-2 microglobulin	cystatin C
alpha-1 microglobulin	beta-2 microglobulin	GST-alpha
alpha-1 microglobulin	beta-2 microglobulin	KIM-1
alpha-1 microglobulin	beta-2 microglobulin	microalbumin
alpha-1 microglobulin	beta-2 microglobulin	NGAL
alpha-1 microglobulin	beta-2 microglobulin	osteopontin
alpha-1 microglobulin	beta-2 microglobulin	THP
alpha-1 microglobulin	beta-2 microglobulin	TIMP-1
alpha-1 microglobulin	beta-2 microglobulin	TFF-3
alpha-1 microglobulin	beta-2 microglobulin	VEGF
alpha-1 microglobulin	beta-2 microglobulin	clusterin
alpha-1 microglobulin	calbindin	CTGF
alpha-1 microglobulin	calbindin	creatinine
alpha-1 microglobulin	calbindin	cystatin C
alpha-1 microglobulin	calbindin	GST-alpha
alpha-1 microglobulin	calbindin	KIM-1

TABLE A-continued

alpha-1 microglobulin	calbindin	microalbumin
alpha-1 microglobulin	calbindin	NGAL
alpha-1 microglobulin	calbindin	osteopontin
alpha-1 microglobulin	calbindin	THP
alpha-1 microglobulin	calbindin	TIMP-1
alpha-1 microglobulin	calbindin	TFF-3
alpha-1 microglobulin	calbindin	VEGF
alpha-1 microglobulin	clusterin	CTGF
alpha-1 microglobulin	clusterin	creatinine
alpha-1 microglobulin	clusterin	cystatin C
alpha-1 microglobulin	clusterin	GST-alpha
alpha-1 microglobulin	clusterin	KIM-1
alpha-1 microglobulin	clusterin	microalbumin
alpha-1 microglobulin	clusterin	NGAL
alpha-1 microglobulin	clusterin	osteopontin
alpha-1 microglobulin	clusterin	THP
alpha-1 microglobulin	clusterin	TIMP-1
alpha-1 microglobulin	clusterin	TFF-3
alpha-1 microglobulin	clusterin	VEGF
alpha-1 microglobulin	clusterin	CTGF
alpha-1 microglobulin	CTGF	creatinine
alpha-1 microglobulin	CTGF	cystatin C
alpha-1 microglobulin	CTGF	GST-alpha
alpha-1 microglobulin	CTGF	KIM-1
alpha-1 microglobulin	CTGF	microalbumin
alpha-1 microglobulin	CTGF	NGAL
alpha-1 microglobulin	CTGF	osteopontin
alpha-1 microglobulin	CTGF	THP
alpha-1 microglobulin	CTGF	TIMP-1
alpha-1 microglobulin	CTGF	TFF-3
alpha-1 microglobulin	CTGF	VEGF
alpha-1 microglobulin	creatinine	cystatin C
alpha-1 microglobulin	creatinine	GST-alpha
alpha-1 microglobulin	creatinine	KIM-1
alpha-1 microglobulin	creatinine	microalbumin
alpha-1 microglobulin	creatinine	NGAL
alpha-1 microglobulin	creatinine	osteopontin
alpha-1 microglobulin	creatinine	THP
alpha-1 microglobulin	creatinine	TIMP-1
alpha-1 microglobulin	creatinine	TFF-3
alpha-1 microglobulin	creatinine	VEGF
alpha-1 microglobulin	cystatin C	GST-alpha
alpha-1 microglobulin	cystatin C	KIM-1
alpha-1 microglobulin	cystatin C	microalbumin
alpha-1 microglobulin	cystatin C	NGAL
alpha-1 microglobulin	cystatin C	osteopontin
alpha-1 microglobulin	cystatin C	THP
alpha-1 microglobulin	cystatin C	TIMP-1
alpha-1 microglobulin	cystatin C	TFF-3
alpha-1 microglobulin	cystatin C	VEGF
alpha-1 microglobulin	GST-alpha	KIM-1
alpha-1 microglobulin	GST-alpha	microalbumin
alpha-1 microglobulin	GST-alpha	NGAL
alpha-1 microglobulin	GST-alpha	osteopontin
alpha-1 microglobulin	GST-alpha	THP
alpha-1 microglobulin	GST-alpha	TIMP-1
alpha-1 microglobulin	GST-alpha	TFF-3
alpha-1 microglobulin	GST-alpha	VEGF
alpha-1 microglobulin	KIM-1	microalbumin
alpha-1 microglobulin	KIM-1	NGAL
alpha-1 microglobulin	KIM-1	osteopontin
alpha-1 microglobulin	KIM-1	THP
alpha-1 microglobulin	KIM-1	TIMP-1
alpha-1 microglobulin	KIM-1	TFF-3
alpha-1 microglobulin	KIM-1	VEGF
alpha-1 microglobulin	microalbumin	NGAL
alpha-1 microglobulin	microalbumin	osteopontin
alpha-1 microglobulin	microalbumin	THP
alpha-1 microglobulin	microalbumin	TIMP-1
alpha-1 microglobulin	microalbumin	TFF-3
alpha-1 microglobulin	microalbumin	VEGF
alpha-1 microglobulin	NGAL	osteopontin
alpha-1 microglobulin	NGAL	THP
alpha-1 microglobulin	NGAL	TIMP-1
alpha-1 microglobulin	NGAL	TFF-3
alpha-1 microglobulin	NGAL	VEGF
alpha-1 microglobulin	osteopontin	THP
alpha-1 microglobulin	osteopontin	TIMP-1

TABLE A-continued

alpha-1 microglobulin	osteopontin	TFF-3
alpha-1 microglobulin	osteopontin	VEGF
alpha-1 microglobulin	THP	TIMP-1
alpha-1 microglobulin	THP	TFF-3
alpha-1 microglobulin	THP	VEGF
alpha-1 microglobulin	TIMP-1	TFF-3
alpha-1 microglobulin	TIMP-1	VEGF
alpha-1 microglobulin	TFF-3	VEGF
beta-2 microglobulin	calbindin	clusterin
beta-2 microglobulin	calbindin	CTGF
beta-2 microglobulin	calbindin	creatinine
beta-2 microglobulin	calbindin	cystatin C
beta-2 microglobulin	calbindin	GST-alpha
beta-2 microglobulin	calbindin	KIM-1
beta-2 microglobulin	calbindin	microalbumin
beta-2 microglobulin	calbindin	NGAL
beta-2 microglobulin	calbindin	osteopontin
beta-2 microglobulin	calbindin	THP
beta-2 microglobulin	calbindin	TIMP-1
beta-2 microglobulin	calbindin	TFF-3
beta-2 microglobulin	calbindin	VEGF
beta-2 microglobulin	clusterin	CTGF
beta-2 microglobulin	clusterin	creatinine
beta-2 microglobulin	clusterin	cystatin C
beta-2 microglobulin	clusterin	GST-alpha
beta-2 microglobulin	clusterin	KIM-1
beta-2 microglobulin	clusterin	microalbumin
beta-2 microglobulin	clusterin	NGAL
beta-2 microglobulin	clusterin	osteopontin
beta-2 microglobulin	clusterin	THP
beta-2 microglobulin	clusterin	TIMP-1
beta-2 microglobulin	clusterin	TFF-3
beta-2 microglobulin	clusterin	VEGF
beta-2 microglobulin	CTGF	creatinine
beta-2 microglobulin	CTGF	cystatin C
beta-2 microglobulin	CTGF	GST-alpha
beta-2 microglobulin	CTGF	KIM-1
beta-2 microglobulin	CTGF	microalbumin
beta-2 microglobulin	CTGF	NGAL
beta-2 microglobulin	CTGF	osteopontin
beta-2 microglobulin	CTGF	THP
beta-2 microglobulin	CTGF	TIMP-1
beta-2 microglobulin	CTGF	TFF-3
beta-2 microglobulin	CTGF	VEGF
beta-2 microglobulin	creatinine	cystatin C
beta-2 microglobulin	creatinine	GST-alpha
beta-2 microglobulin	creatinine	KIM-1
beta-2 microglobulin	creatinine	microalbumin
beta-2 microglobulin	creatinine	NGAL
beta-2 microglobulin	creatinine	osteopontin
beta-2 microglobulin	creatinine	THP
beta-2 microglobulin	creatinine	TIMP-1
beta-2 microglobulin	creatinine	TFF-3
beta-2 microglobulin	creatinine	VEGF
beta-2 microglobulin	cystatin C	GST-alpha
beta-2 microglobulin	cystatin C	KIM-1
beta-2 microglobulin	cystatin C	microalbumin
beta-2 microglobulin	cystatin C	NGAL
beta-2 microglobulin	cystatin C	osteopontin
beta-2 microglobulin	cystatin C	THP
beta-2 microglobulin	cystatin C	TIMP-1
beta-2 microglobulin	cystatin C	TFF-3
beta-2 microglobulin	cystatin C	VEGF
beta-2 microglobulin	GST-alpha	KIM-1
beta-2 microglobulin	GST-alpha	microalbumin
beta-2 microglobulin	GST-alpha	NGAL
beta-2 microglobulin	GST-alpha	osteopontin
beta-2 microglobulin	GST-alpha	THP
beta-2 microglobulin	GST-alpha	TIMP-1
beta-2 microglobulin	GST-alpha	TFF-3
beta-2 microglobulin	GST-alpha	VEGF
beta-2 microglobulin	KIM-1	microalbumin
beta-2 microglobulin	KIM-1	NGAL
beta-2 microglobulin	KIM-1	osteopontin
beta-2 microglobulin	KIM-1	THP
beta-2 microglobulin	KIM-1	TIMP-1
beta-2 microglobulin	KIM-1	TFF-3

TABLE A-continued

beta-2 microglobulin	KIM-1	VEGF
beta-2 microglobulin	microalbumin	NGAL
beta-2 microglobulin	microalbumin	osteopontin
beta-2 microglobulin	microalbumin	THP
beta-2 microglobulin	microalbumin	TIMP-1
beta-2 microglobulin	microalbumin	TFF-3
beta-2 microglobulin	microalbumin	VEGF
beta-2 microglobulin	NGAL	osteopontin
beta-2 microglobulin	NGAL	THP
beta-2 microglobulin	NGAL	TIMP-1
beta-2 microglobulin	NGAL	TFF-3
beta-2 microglobulin	NGAL	VEGF
beta-2 microglobulin	osteopontin	THP
beta-2 microglobulin	osteopontin	TIMP-1
beta-2 microglobulin	osteopontin	TFF-3
beta-2 microglobulin	osteopontin	VEGF
beta-2 microglobulin	THP	TIMP-1
beta-2 microglobulin	THP	TFF-3
beta-2 microglobulin	THP	VEGF
beta-2 microglobulin	TIMP-1	TFF-3
beta-2 microglobulin	TIMP-2	VEGF
beta-2 microglobulin	TFF-3	VEGF
calbindin	clusterin	CTGF
calbindin	clusterin	creatinine
calbindin	clusterin	cystatin C
calbindin	clusterin	GST-alpha
calbindin	clusterin	KIM-1
calbindin	clusterin	microalbumin
calbindin	clusterin	NGAL
calbindin	clusterin	osteopontin
calbindin	clusterin	THP
calbindin	clusterin	TIMP-1
calbindin	clusterin	TFF-3
calbindin	clusterin	VEGF
calbindin	clusterin	VEGF
calbindin	CTGF	creatinine
calbindin	CTGF	cystatin C
calbindin	CTGF	GST-alpha
calbindin	CTGF	KIM-1
calbindin	CTGF	microalbumin
calbindin	CTGF	NGAL
calbindin	CTGF	osteopontin
calbindin	CTGF	THP
calbindin	CTGF	TIMP-1
calbindin	CTGF	TFF-3
calbindin	CTGF	VEGF
calbindin	creatinine	cystatin C
calbindin	creatinine	GST-alpha
calbindin	creatinine	KIM-1
calbindin	creatinine	microalbumin
calbindin	creatinine	NGAL
calbindin	creatinine	osteopontin
calbindin	creatinine	THP
calbindin	creatinine	TIMP-1
calbindin	creatinine	TFF-3
calbindin	creatinine	VEGF
calbindin	cystatin C	GST-alpha
calbindin	cystatin C	KIM-1
calbindin	cystatin C	microalbumin
calbindin	cystatin C	NGAL
calbindin	cystatin C	osteopontin
calbindin	cystatin C	THP
calbindin	cystatin C	TIMP-1
calbindin	cystatin C	TFF-3
calbindin	cystatin C	VEGF
calbindin	GST-alpha	KIM-1
calbindin	GST-alpha	microalbumin
calbindin	GST-alpha	NGAL
calbindin	GST-alpha	osteopontin
calbindin	GST-alpha	THP
calbindin	GST-alpha	TIMP-1
calbindin	GST-alpha	TFF-3
calbindin	GST-alpha	VEGF
calbindin	GST-alpha	VEGF
calbindin	KIM-1	microalbumin
calbindin	KIM-1	NGAL
calbindin	KIM-1	osteopontin
calbindin	KIM-1	THP
calbindin	KIM-1	TIMP-1
calbindin	KIM-1	TFF-3

TABLE A-continued

calbindin	KIM-1	TFF-3
calbindin	KIM-1	VEGF
calbindin	microalbumin	NGAL
calbindin	microalbumin	osteopontin
calbindin	microalbumin	THP
calbindin	microalbumin	TIMP-1
calbindin	microalbumin	TFF-3
calbindin	microalbumin	VEGF
calbindin	NGAL	osteopontin
calbindin	NGAL	THP
calbindin	NGAL	TIMP-1
calbindin	NGAL	TFF-3
calbindin	NGAL	VEGF
calbindin	osteopontin	THP
calbindin	osteopontin	TIMP-1
calbindin	osteopontin	TFF-3
calbindin	osteopontin	VEGF
calbindin	THP	TIMP-1
calbindin	THP	TFF-3
calbindin	THP	VEGF
calbindin	TIMP-1	TFF-3
calbindin	TIMP-1	VEGF
calbindin	TFF-3	VEGF
clusterin	CTGF	creatinine
clusterin	CTGF	cystatin C
clusterin	CTGF	GST-alpha
clusterin	CTGF	KIM-1
clusterin	CTGF	microalbumin
clusterin	CTGF	NGAL
clusterin	CTGF	osteopontin
clusterin	CTGF	THP
clusterin	CTGF	TIMP-1
clusterin	CTGF	TFF-3
clusterin	CTGF	VEGF
clusterin	creatinine	cystatin C
clusterin	creatinine	GST-alpha
clusterin	creatinine	KIM-1
clusterin	creatinine	microalbumin
clusterin	creatinine	NGAL
clusterin	creatinine	osteopontin
clusterin	creatinine	THP
clusterin	creatinine	TIMP-1
clusterin	creatinine	TFF-3
clusterin	creatinine	VEGF
clusterin	cystatin C	GST-alpha
clusterin	cystatin C	KIM-1
clusterin	cystatin C	microalbumin
clusterin	cystatin C	NGAL
clusterin	cystatin C	osteopontin
clusterin	cystatin C	THP
clusterin	cystatin C	TIMP-1
clusterin	cystatin C	TFF-3
clusterin	cystatin C	VEGF
clusterin	GST-alpha	KIM-1
clusterin	GST-alpha	microalbumin
clusterin	GST-alpha	NGAL
clusterin	GST-alpha	osteopontin
clusterin	GST-alpha	THP
clusterin	GST-alpha	TIMP-1
clusterin	GST-alpha	TFF-3
clusterin	GST-alpha	VEGF
clusterin	KIM-1	microalbumin
clusterin	KIM-1	NGAL
clusterin	KIM-1	osteopontin
clusterin	KIM-1	THP
clusterin	KIM-1	TIMP-1
clusterin	KIM-1	TFF-3
clusterin	KIM-1	VEGF
clusterin	microalbumin	NGAL
clusterin	microalbumin	osteopontin
clusterin	microalbumin	THP
clusterin	microalbumin	TIMP-1
clusterin	microalbumin	TFF-3
clusterin	microalbumin	VEGF
clusterin	NGAL	osteopontin
clusterin	NGAL	THP
clusterin	NGAL	TIMP-1

TABLE A-continued

clusterin	NGAL	TFF-3
clusterin	NGAL	VEGF
clusterin	osteopontin	THP
clusterin	osteopontin	TIMP-1
clusterin	osteopontin	TFF-3
clusterin	osteopontin	VEGF
clusterin	THP	TIMP-1
clusterin	THP	TFF-3
clusterin	THP	VEGF
clusterin	TIMP-1	TFF-3
clusterin	TIMP-1	VEGF
clusterin	TFF-3	VEGF
CTGF	creatinine	cystatin C
CTGF	creatinine	GST-alpha
CTGF	creatinine	KIM-1
CTGF	creatinine	microalbumin
CTGF	creatinine	NGAL
CTGF	creatinine	osteopontin
CTGF	creatinine	THP
CTGF	creatinine	TIMP-1
CTGF	creatinine	TFF-3
CTGF	creatinine	VEGF
CTGF	cystatin C	GST-alpha
CTGF	cystatin C	KIM-1
CTGF	cystatin C	microalbumin
CTGF	cystatin C	NGAL
CTGF	cystatin C	osteopontin
CTGF	cystatin C	THP
CTGF	cystatin C	TIMP-1
CTGF	cystatin C	TFF-3
CTGF	cystatin C	VEGF
CTGF	GST-alpha	KIM-1
CTGF	GST-alpha	microalbumin
CTGF	GST-alpha	NGAL
CTGF	GST-alpha	osteopontin
CTGF	GST-alpha	THP
CTGF	GST-alpha	TIMP-1
CTGF	GST-alpha	TFF-3
CTGF	GST-alpha	VEGF
CTGF	KIM-1	microalbumin
CTGF	KIM-1	NGAL
CTGF	KIM-1	osteopontin
CTGF	KIM-1	THP
CTGF	KIM-1	TIMP-1
CTGF	KIM-1	TFF-3
CTGF	KIM-1	VEGF
CTGF	microalbumin	NGAL
CTGF	microalbumin	osteopontin
CTGF	microalbumin	THP
CTGF	microalbumin	TIMP-1
CTGF	microalbumin	TFF-3
CTGF	microalbumin	VEGF
CTGF	NGAL	osteopontin
CTGF	NGAL	THP
CTGF	NGAL	TIMP-1
CTGF	NGAL	TFF-3
CTGF	NGAL	VEGF
CTGF	osteopontin	THP
CTGF	osteopontin	TIMP-1
CTGF	osteopontin	TFF-3
CTGF	osteopontin	VEGF
CTGF	THP	TIMP-1
CTGF	THP	TFF-3
CTGF	THP	VEGF
CTGF	TIMP-1	TFF-3
CTGF	TIMP-1	VEGF
CTGF	TFF-3	VEGF
creatinine	cystatin C	GST-alpha
creatinine	cystatin C	KIM-1
creatinine	cystatin C	microalbumin
creatinine	cystatin C	NGAL
creatinine	cystatin C	osteopontin
creatinine	cystatin C	THP
creatinine	cystatin C	TIMP-1
creatinine	cystatin C	TFF-3
creatinine	cystatin C	VEGF
creatinine	GST-alpha	KIM-1
creatinine	GST-alpha	microalbumin

TABLE A-continued

creatinine	GST-alpha	microalbumin
creatinine	GST-alpha	NGAL
creatinine	GST-alpha	osteopontin
creatinine	GST-alpha	THP
creatinine	GST-alpha	TIMP-1
creatinine	GST-alpha	TFF-3
creatinine	GST-alpha	VEGF
creatinine	KIM-1	microalbumin
creatinine	KIM-1	NGAL
creatinine	KIM-1	osteopontin
creatinine	KIM-1	THP
creatinine	KIM-1	TIMP-1
creatinine	KIM-1	TFF-3
creatinine	KIM-1	VEGF
creatinine	microalbumin	NGAL
creatinine	microalbumin	osteopontin
creatinine	microalbumin	THP
creatinine	microalbumin	TIMP-1
creatinine	microalbumin	TFF-3
creatinine	microalbumin	VEGF
creatinine	NGAL	osteopontin
creatinine	NGAL	THP
creatinine	NGAL	TIMP-1
creatinine	NGAL	TFF-3
creatinine	NGAL	VEGF
creatinine	osteopontin	THP
creatinine	osteopontin	TIMP-1
creatinine	osteopontin	TFF-3
creatinine	osteopontin	VEGF
creatinine	THP	TIMP-1
creatinine	THP	TFF-3
creatinine	THP	VEGF
creatinine	TIMP-1	TFF-3
creatinine	TIMP-1	VEGF
creatinine	TFF-3	VEGF
cystatin C	GST-alpha	KIM-1
cystatin C	GST-alpha	microalbumin
cystatin C	GST-alpha	NGAL
cystatin C	GST-alpha	osteopontin
cystatin C	GST-alpha	THP
cystatin C	GST-alpha	TIMP-1
cystatin C	GST-alpha	TFF-3
cystatin C	GST-alpha	VEGF
cystatin C	KIM-1	microalbumin
cystatin C	KIM-1	NGAL
cystatin C	KIM-1	osteopontin
cystatin C	KIM-1	THP
cystatin C	KIM-1	TIMP-1
cystatin C	KIM-1	TFF-3
cystatin C	KIM-1	VEGF
cystatin C	microalbumin	NGAL
cystatin C	microalbumin	osteopontin
cystatin C	microalbumin	THP
cystatin C	microalbumin	TIMP-1
cystatin C	microalbumin	TFF-3
cystatin C	microalbumin	VEGF
cystatin C	NGAL	osteopontin
cystatin C	NGAL	THP
cystatin C	NGAL	TIMP-1
cystatin C	NGAL	TFF-3
cystatin C	NGAL	VEGF
cystatin C	NGAL	VEGF
cystatin C	osteopontin	THP
cystatin C	osteopontin	TIMP-1
cystatin C	osteopontin	TFF-3
cystatin C	osteopontin	VEGF
cystatin C	osteopontin	VEGF
cystatin C	THP	TIMP-1
cystatin C	THP	TFF-3
cystatin C	THP	VEGF
cystatin C	TIMP-1	TFF-3
cystatin C	TIMP-1	VEGF
cystatin C	TFF-3	VEGF
GST-alpha	KIM-1	microalbumin
GST-alpha	KIM-1	NGAL
GST-alpha	KIM-1	osteopontin
GST-alpha	KIM-1	THP
GST-alpha	KIM-1	TIMP-1
GST-alpha	KIM-1	TFF-3

TABLE A-continued

GST-alpha	KIM-1	VEGF
GST-alpha	microalbumin	NGAL
GST-alpha	microalbumin	osteopontin
GST-alpha	microalbumin	THP
GST-alpha	microalbumin	TIMP-1
GST-alpha	microalbumin	TFF-3
GST-alpha	microalbumin	VEGF
GST-alpha	NGAL	osteopontin
GST-alpha	NGAL	THP
GST-alpha	NGAL	TIMP-1
GST-alpha	NGAL	TFF-3
GST-alpha	NGAL	VEGF
GST-alpha	osteopontin	THP
GST-alpha	osteopontin	TIMP-1
GST-alpha	osteopontin	TFF-3
GST-alpha	osteopontin	VEGF
GST-alpha	THP	TIMP-1
GST-alpha	THP	TFF-3
GST-alpha	THP	VEGF
GST-alpha	TIMP-1	TFF-3
GST-alpha	TIMP-1	VEGF
GST-alpha	TFF-3	VEGF
KIM-1	microalbumin	NGAL
KIM-1	microalbumin	osteopontin
KIM-1	microalbumin	THP
KIM-1	microalbumin	TIMP-1
KIM-1	microalbumin	TFF-3
KIM-1	microalbumin	VEGF
KIM-1	NGAL	osteopontin
KIM-1	NGAL	THP
KIM-1	NGAL	TIMP-1
KIM-1	NGAL	TFF-3
KIM-1	NGAL	VEGF
KIM-1	osteopontin	THP
KIM-1	osteopontin	TIMP-1
KIM-1	osteopontin	TFF-3
KIM-1	osteopontin	VEGF
KIM-1	THP	TIMP-1
KIM-1	THP	TFF-3
KIM-1	THP	VEGF
KIM-1	TIMP-1	TFF-3
KIM-1	TIMP-1	VEGF
KIM-1	TFF-3	VEGF
microalbumin	NGAL	osteopontin
microalbumin	NGAL	THP
microalbumin	NGAL	TIMP-1
microalbumin	NGAL	TFF-3
microalbumin	NGAL	VEGF
microalbumin	osteopontin	THP
microalbumin	osteopontin	TIMP-1
microalbumin	osteopontin	TFF-3
microalbumin	osteopontin	VEGF
microalbumin	THP	TIMP-1
microalbumin	THP	TFF-3
microalbumin	THP	VEGF
microalbumin	TIMP-1	TFF-3
microalbumin	TIMP-1	VEGF
microalbumin	TFF-3	VEGF
NGAL	osteopontin	THP
NGAL	osteopontin	TIMP-1
NGAL	osteopontin	TFF-3
NGAL	osteopontin	VEGF
NGAL	THP	TIMP-1
NGAL	THP	TFF-3
NGAL	THP	VEGF
NGAL	TIMP-1	TFF-3
NGAL	TIMP-1	VEGF
NGAL	TFF-3	VEGF
osteopontin	THP	TIMP-1
osteopontin	THP	TFF-3
osteopontin	THP	VEGF
osteopontin	TIMP-1	TFF-3
osteopontin	TIMP-1	VEGF
osteopontin	TFF-3	VEGF
THP	TIMP-1	TFF-3
THP	TIMP-1	VEGF
THP	TFF-3	VEGF

TABLE A-continued

THP	TIMP-1	VEGF
THP	TFF-3	VEGF
TIMP-1	TFF-3	VEGF

[0049] In one exemplary embodiment, the combination of sample analytes may include creatinine, KIM-1, and THP. In another exemplary embodiment, the combination of sample analytes may include microalbumin, creatinine, and KIM-1. In yet another exemplary embodiment, the combination of sample analytes may include creatinine, THP, and A1M. In still another exemplary embodiment, the combination of sample analytes may include microalbumin, TIMP-1, and osteopontin. In an alternative exemplary embodiment, the combination of sample analytes may include creatinine, THP, alpha 1 microglobulin, clusterin, NGAL, and osteopontin.

III. Test Sample

[0050] The method for diagnosing, monitoring, or determining a renal disorder involves determining the presence of sample analytes in a test sample. A test sample, as defined herein, is an amount of bodily fluid taken from a mammal. Non-limiting examples of bodily fluids include urine, blood, plasma, serum, saliva, semen, perspiration, tears, mucus, and tissue lysates. In an exemplary embodiment, the bodily fluid contained in the test sample is urine, plasma, or serum.

(a) Mammals

[0051] A mammal, as defined herein, is any organism that is a member of the class Mammalia. Non-limiting examples of mammals appropriate for the various embodiments may include humans, apes, monkeys, rats, mice, dogs, cats, pigs, and livestock including cattle and oxen. In an exemplary embodiment, the mammal is a human.

(b) Devices and Methods of Taking Bodily Fluids from Mammals

[0052] The bodily fluids of the test sample may be taken from the mammal using any known device or method so long as the analytes to be measured by the multiplexed assay are not rendered undetectable by the multiplexed assay. Non-limiting examples of devices or methods suitable for taking bodily fluid from a mammal include urine sample cups, urethral catheters, swabs, hypodermic needles, thin needle biopsies, hollow needle biopsies, punch biopsies, metabolic cages, and aspiration.

[0053] In order to adjust the expected concentrations of the sample analytes in the test sample to fall within the dynamic range of the multiplexed assay, the test sample may be diluted to reduce the concentration of the sample analytes prior to analysis. The degree of dilution may depend on a variety of factors including but not limited to the type of multiplexed assay used to measure the analytes, the reagents utilized in the multiplexed assay, and the type of bodily fluid contained in the test sample. In one embodiment, the test sample is diluted by adding a volume of diluent ranging from about 1/2 of the original test sample volume to about 50,000 times the original test sample volume.

[0054] In one exemplary embodiment, if the test sample is human urine and the multiplexed assay is an antibody-based capture-sandwich assay, the test sample is diluted by adding a volume of diluent that is about 100 times the original test sample volume prior to analysis. In another exemplary embodiment, if the test sample is human serum and the mul-

tiplexed assay is an antibody-based capture-sandwich assay, the test sample is diluted by adding a volume of diluent that is about 5 times the original test sample volume prior to analysis. In yet another exemplary embodiment, if the test sample is human plasma and the multiplexed assay is an antibody-based capture-sandwich assay, the test sample is diluted by adding a volume of diluent that is about 2,000 times the original test sample volume prior to analysis.

[0055] The diluent may be any fluid that does not interfere with the function of the multiplexed assay used to measure the concentration of the analytes in the test sample. Non-limiting examples of suitable diluents include deionized water, distilled water, saline solution, Ringer's solution, phosphate buffered saline solution, TRIS-buffered saline solution, standard saline citrate, and HEPES-buffered saline.

IV. Multiplexed Assay Device

[0056] In one embodiment, the concentration of a combination of sample analytes is measured using a multiplexed assay device capable of measuring the concentrations of three or more, preferably sixteen, of the biomarker analytes. A multiplexed assay device, as defined herein, is an assay capable of simultaneously determining the concentration of three or more different sample analytes using a single device and/or method. Any known method of measuring the concentration of the biomarker analytes may be used for the multiplexed assay device. Non-limiting examples of measurement methods suitable for the multiplexed assay device may include electrophoresis, mass spectrometry, protein microarrays, surface plasmon resonance and immunoassays including but not limited to western blot, immunohistochemical staining, enzyme-linked immunosorbent assay (ELISA) methods, and particle-based capture-sandwich immunoassays.

(a) Multiplexed Immunoassay Device

[0057] In one embodiment, the concentrations of the analytes in the test sample are measured using a multiplexed immunoassay device that utilizes capture antibodies marked with indicators to determine the concentration of the sample analytes.

(i) Capture Antibodies

[0058] In the same embodiment, the multiplexed immunoassay device includes three or more capture antibodies. Capture antibodies, as defined herein, are antibodies in which the antigenic determinant is one of the biomarker analytes. Each of the at least three capture antibodies has a unique antigenic determinant that is one of the biomarker analytes. When contacted with the test sample, the capture antibodies form antigen-antibody complexes in which the analytes serve as antigens.

[0059] The term "antibody," as used herein, encompasses a monoclonal ab, an antibody fragment, a chimeric antibody, and a single-chain antibody.

[0060] In some embodiments, the capture antibodies may be attached to a substrate in order to immobilize any analytes captured by the capture antibodies. Non-limiting examples of suitable substrates include paper, cellulose, glass, or plastic strips, beads, or surfaces, such as the inner surface of the well of a microtitration tray. Suitable beads may include polystyrene or latex microspheres.

(ii) Indicators

[0061] In one embodiment of the multiplexed immunoassay device, an indicator is attached to each of the three or more capture antibodies. The indicator, as defined herein, is any compound that registers a measurable change to indicate the presence of one of the sample analytes when bound to one of the capture antibodies. Non-limiting examples of indicators include visual indicators and electrochemical indicators.

[0062] Visual indicators, as defined herein, are compounds that register a change by reflecting a limited subset of the wavelengths of light illuminating the indicator, by fluorescing light after being illuminated, or by emitting light via chemiluminescence. The change registered by visual indicators may be in the visible light spectrum, in the infrared spectrum, or in the ultraviolet spectrum. Non-limiting examples of visual indicators suitable for the multiplexed immunoassay device include nanoparticulate gold, organic particles such as polyurethane or latex microspheres loaded with dye compounds, carbon black, fluorophores, phycoerythrin, radioactive isotopes, nanoparticles, quantum dots, and enzymes such as horseradish peroxidase or alkaline phosphatase that react with a chemical substrate to form a colored or chemiluminescent product.

[0063] Electrochemical indicators, as defined herein, are compounds that register a change by altering an electrical property. The changes registered by electrochemical indicators may be an alteration in conductivity, resistance, capacitance, current conducted in response to an applied voltage, or voltage required to achieve a desired current. Non-limiting examples of electrochemical indicators include redox species such as ascorbate (vitamin C), vitamin E, glutathione, polyphenols, catechols, quercetin, phytoestrogens, penicillin, carbazole, murrans, phenols, carbonyls, benzoates, and trace metal ions such as nickel, copper, cadmium, iron and mercury.

[0064] In this same embodiment, the test sample containing a combination of three or more sample analytes is contacted with the capture antibodies and allowed to form antigen-antibody complexes in which the sample analytes serve as the antigens. After removing any uncomplexed capture antibodies, the concentrations of the three or more analytes are determined by measuring the change registered by the indicators attached to the capture antibodies.

[0065] In one exemplary embodiment, the indicators are polyurethane or latex microspheres loaded with dye compounds and phycoerythrin.

(b) Multiplexed Sandwich Immunoassay Device

[0066] In another embodiment, the multiplexed immunoassay device has a sandwich assay format. In this embodiment, the multiplexed sandwich immunoassay device includes three or more capture antibodies as previously described. However, in this embodiment, each of the capture antibodies is attached to a capture agent that includes an antigenic moiety. The antigenic moiety serves as the antigenic determinant of a detection antibody, also included in the multiplexed immunoassay device of this embodiment. In addition, an indicator is attached to the detection antibody.

[0067] In this same embodiment, the test sample is contacted with the capture antibodies and allowed to form antigen-antibody complexes in which the sample analytes serve as antigens. The detection antibodies are then contacted with the test sample and allowed to form antigen-antibody com-

plexes in which the capture agent serves as the antigen for the detection antibody. After removing any uncomplexed detection antibodies the concentration of the analytes are determined by measuring the changes registered by the indicators attached to the detection antibodies.

(c) Multiplexing Approaches

[0068] In the various embodiments of the multiplexed immunoassay devices, the concentrations of each of the sample analytes may be determined using any approach known in the art. In one embodiment, a single indicator compound is attached to each of the three or more antibodies. In addition, each of the capture antibodies having one of the sample analytes as an antigenic determinant is physically separated into a distinct region so that the concentration of each of the sample analytes may be determined by measuring the changes registered by the indicators in each physically separate region corresponding to each of the sample analytes.

[0069] In another embodiment, each antibody having one of the sample analytes as an antigenic determinant is marked with a unique indicator. In this manner, a unique indicator is attached to each antibody having a single sample analyte as its antigenic determinant. In this embodiment, all antibodies may occupy the same physical space. The concentration of each sample analyte is determined by measuring the change registered by the unique indicator attached to the antibody having the sample analyte as an antigenic determinant.

(d) Microsphere-Based Capture-Sandwich Immunoassay Device

[0070] In an exemplary embodiment, the multiplexed immunoassay device is a microsphere-based capture-sandwich immunoassay device. In this embodiment, the device includes a mixture of three or more capture-antibody microspheres, in which each capture-antibody microsphere corresponds to one of the biomarker analytes. Each capture-antibody microsphere includes a plurality of capture antibodies attached to the outer surface of the microsphere. In this same embodiment, the antigenic determinant of all of the capture antibodies attached to one microsphere is the same biomarker analyte.

[0071] In this embodiment of the device, the microsphere is a small polystyrene or latex sphere that is loaded with an indicator that is a dye compound. The microsphere may be between about 3 μm and about 5 μm in diameter. Each capture-antibody microsphere corresponding to one of the biomarker analytes is loaded with the same indicator. In this manner, each capture-antibody microsphere corresponding to a biomarker analyte is uniquely color-coded.

[0072] In this same exemplary embodiment, the multiplexed immunoassay device further includes three or more biotinylated detection antibodies in which the antigenic determinant of each biotinylated detection antibody is one of the biomarker analytes. The device further includes a plurality of streptavidin proteins complexed with a reporter compound. A reporter compound, as defined herein, is an indicator selected to register a change that is distinguishable from the indicators used to mark the capture-antibody microspheres.

[0073] The concentrations of the sample analytes may be determined by contacting the test sample with a mixture of capture-antigen microspheres corresponding to each sample analyte to be measured. The sample analytes are allowed to

form antigen-antibody complexes in which a sample analyte serves as an antigen and a capture antibody attached to the microsphere serves as an antibody. In this manner, the sample analytes are immobilized onto the capture-antigen microspheres. The biotinylated detection antibodies are then added to the test sample and allowed to form antigen-antibody complexes in which the analyte serves as the antigen and the biotinylated detection antibody serves as the antibody. The streptavidin-reporter complex is then added to the test sample and allowed to bind to the biotin moieties of the biotinylated detection antibodies. The antigen-capture microspheres may then be rinsed and filtered.

[0074] In this embodiment, the concentration of each analyte is determined by first measuring the change registered by the indicator compound embedded in the capture-antigen microsphere in order to identify the particular analyte. For each microsphere corresponding to one of the biomarker analytes, the quantity of analyte immobilized on the microsphere is determined by measuring the change registered by the reporter compound attached to the microsphere.

[0075] For example, the indicator embedded in the microspheres associated with one sample analyte may register an emission of orange light, and the reporter may register an emission of green light. In this example, a detector device may measure the intensity of orange light and green light separately. The measured intensity of the green light would determine the concentration of the analyte captured on the microsphere, and the intensity of the orange light would determine the specific analyte captured on the microsphere.

[0076] Any sensor device may be used to detect the changes registered by the indicators embedded in the microspheres and the changes registered by the reporter compound, so long as the sensor device is sufficiently sensitive to the changes registered by both indicator and reporter compound. Non-limiting examples of suitable sensor devices include spectro-

photometers, photosensors, colorimeters, cyclic coulometry devices, and flow cytometers. In an exemplary embodiment, the sensor device is a flow cytometer.

V. Method for Diagnosing, Monitoring, or Determining a Renal Disorder

[0077] In one embodiment, a method is provided for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder that includes providing a test sample, determining the concentration of a combination of three or more sample analytes, comparing the measured concentrations of the combination of sample analytes to the entries of a dataset, and identifying obstructive uropathy or an associated disorder based on the comparison between the concentrations of the sample analytes and the minimum diagnostic concentrations contained within each entry of the dataset.

(a) Diagnostic Dataset

[0078] In an embodiment, the concentrations of the sample analytes are compared to the entries of a dataset. In this embodiment, each entry of the dataset includes a combination of three or more minimum diagnostic concentrations indicative of a particular renal disorder. A minimum diagnostic concentration, as defined herein, is the concentration of an analyte that defines the limit between the concentration range corresponding to normal, healthy renal function and the concentration reflective of a particular renal disorder. In one embodiment, each minimum diagnostic concentration is the maximum concentration of the range of analyte concentrations for a healthy, normal individual. The minimum diagnostic concentration of an analyte depends on a number of factors including but not limited to the particular analyte and the type of bodily fluid contained in the test sample. As an illustrative example, Table 1 lists the expected normal ranges of the biomarker analytes in human plasma, serum, and urine.

TABLE 1

Normal Concentration Ranges In Human Plasma, Serum, and Urine Samples							
Analyte	Units	Plasma		Sera		Urine	
		low	high	low	high	low	high
Calbindin	ng/ml	—	<5.0	—	<2.6	4.2	233
Clusterin	μg/ml	86	134	37	152	—	<0.089
CTGF	ng/ml	2.8	7.5	—	<8.2	—	<0.90
GST-alpha	ng/ml	6.7	62	1.2	52	—	<26
KIM-1	ng/ml	0.053	0.57	—	<0.35	0.023	0.67
VEGF	pg/ml	222	855	219	1630	69	517
B2M	μg/ml	0.68	2.2	1.00	2.6	—	<0.17
Cyst C	ng/ml	608	1170	476	1250	3.9	79
NGAL	ng/ml	89	375	102	822	2.9	81
OPN	ng/ml	4.1	25	0.49	12	291	6130
TIMP-1	ng/ml	50	131	100	246	—	<3.9
A1M	μg/ml	6.2	16	5.7	17	—	<4.2
THP	μg/ml	0.0084	0.052	0.0079	0.053	0.39	2.6
TFF3	μg/ml	0.040	0.49	0.021	0.17	—	<21
Creatinine	mg/dL	—	—	—	—	13	212
Microalbumin	μg/ml	—	—	—	—	—	>16

[0079] In one embodiment, the high values shown for each of the biomarker analytes in Table 1 for the analytic concentrations in human plasma, sera and urine are the minimum diagnostics values for the analytes in human plasma, sera, and urine, respectively. In one exemplary embodiment, the minimum diagnostic concentration in human plasma of alpha-1 microglobulin is about 16 $\mu\text{g/ml}$, beta-2 microglobulin is about 2.2 $\mu\text{g/ml}$, calbindin is greater than about 5 ng/ml , clusterin is about 134 $\mu\text{g/ml}$, CTGF is about 16 ng/ml , cystatin C is about 1170 ng/ml , GST-alpha is about 62 ng/ml . KIM-1 is about 0.57 ng/ml , NGAL is about 375 ng/ml , osteopontin is about 25 ng/ml , THP is about 0.052 $\mu\text{g/ml}$, TIMP-1 is about 131 ng/ml , TFF-3 is about 0.49 $\mu\text{g/ml}$, and VEGF is about 855 pg/ml .

[0080] In another exemplary embodiment, the minimum diagnostic concentration in human sera of alpha-1 microglobulin is about 17 $\mu\text{g/ml}$, beta-2 microglobulin is about 2.6 $\mu\text{g/ml}$, calbindin is greater than about 2.6 ng/ml , clusterin is about 152 $\mu\text{g/ml}$, CTGF is greater than about 8.2 ng/ml , cystatin C is about 1250 ng/ml , GST-alpha is about 52 ng/ml , KIM-1 is greater than about 0.35 ng/ml , NGAL is about 822 ng/ml , osteopontin is about 12 ng/ml , THP is about 0.053 $\mu\text{g/ml}$, TIMP-1 is about 246 ng/ml , TFF-3 is about 0.17 $\mu\text{g/ml}$, and VEGF is about 1630 pg/ml .

[0081] In yet another exemplary embodiment, the minimum diagnostic concentration in human urine of alpha-1 microglobulin is about 233 $\mu\text{g/ml}$, beta-2 microglobulin is greater than about 0.17 $\mu\text{g/ml}$, calbindin is about 233 ng/ml , clusterin is greater than about 0.089 $\mu\text{g/ml}$, CTGF is greater than about 0.90 ng/ml , cystatin C is about 1170 ng/ml , GST-alpha is greater than about 26 ng/ml , KIM-1 is about 0.67 ng/ml , NGAL is about 81 ng/ml , osteopontin is about 6130 ng/ml , THP is about 2.6 $\mu\text{g/ml}$, TIMP-1 is greater than about 3.9 ng/ml , TFF-3 is greater than about 21 $\mu\text{g/ml}$, and VEGF is about 517 pg/ml .

[0082] In one embodiment, the minimum diagnostic concentrations represent the maximum level of analyte concentrations falling within an expected normal range. Obstructive uropathy or an associated disorder may be indicated if the concentration of an analyte is higher than the minimum diagnostic concentration for the analyte.

[0083] If diminished concentrations of a particular analyte are known to be associated with obstructive uropathy or an associated disorder, the minimum diagnostic concentration may not be an appropriate diagnostic criterion for identifying obstructive uropathy or an associated disorder indicated by the sample analyte concentrations. In these cases, a maximum diagnostic concentration may define the limit between the expected normal concentration range for the analyte and a sample concentration reflective of obstructive uropathy or an associated disorder. In those cases in which a maximum diagnostic concentration is the appropriate diagnostic criterion, sample concentrations that fall below a maximum diagnostic concentration may indicate obstructive uropathy or an associated disorder.

[0084] A critical feature of the method of the multiplexed analyte panel is that a combination of sample analyte concentrations may be used to diagnose obstructive uropathy or an associated disorder. In addition to comparing subsets of the biomarker analyte concentrations to diagnostic criteria, the analytes may be algebraically combined and compared to corresponding diagnostic criteria. In one embodiment, two or more sample analyte concentrations may be added and/or subtracted to determine a combined analyte concentration. In

another embodiment, two or more sample analyte concentrations may be multiplied and/or divided to determine a combined analyte concentration. To identify obstructive uropathy or an associated disorder, the combined analyte concentration may be compared to a diagnostic criterion in which the corresponding minimum or maximum diagnostic concentrations are combined using the same algebraic operations used to determine the combined analyte concentration.

[0085] In yet another embodiment, the analyte concentration measured from a test sample containing one type of body fluid may be algebraically combined with an analyte concentration measured from a second test sample containing a second type of body fluid to determine a combined analyte concentration. For example, the ratio of urine calbindin to plasma calbindin may be determined and compared to a corresponding minimum diagnostic urine:plasma calbindin ratio to identify a particular renal disorder.

[0086] A variety of methods known in the art may be used to define the diagnostic criteria used to identify obstructive uropathy or an associated disorder. In one embodiment, any sample concentration falling outside the expected normal range indicates obstructive uropathy or an associated disorder. In another embodiment, the multiplexed analyte panel may be used to evaluate the analyte concentrations in test samples taken from a population of patients having obstructive uropathy or an associated disorder and compared to the normal expected analyte concentration ranges. In this same embodiment, any sample analyte concentrations that are significantly higher or lower than the expected normal concentration range may be used to define a minimum or maximum diagnostic concentration, respectively. A number of studies comparing the biomarker concentration ranges of a population of patients having a renal disorder to the corresponding analyte concentrations from a population of normal healthy subjects are described in the examples section below.

[0087] In an exemplary embodiment, an analyte value in a test sample higher than the minimum diagnostic value for the top 3 analytes of the particular sample type (e.g. plasma, urine, etc.), wherein the top 3 are determined by the random forest classification method may result in a diagnosis of obstructive uropathy.

VI. Automated Method for Diagnosing, Monitoring, or Determining a Renal Disorder

[0088] In one embodiment, a system for diagnosing, monitoring, or determining obstructive uropathy or an associated disorder in a mammal is provided that includes a database to store a plurality of renal disorder database entries, and a processing device that includes the modules of a renal disorder determining application. In this embodiment, the modules are executable by the processing device, and include an analyte input module, a comparison module, and an analysis module.

[0089] The analyte input module receives three or more sample analyte concentrations that include the biomarker analytes. In one embodiment, the sample analyte concentrations are entered as input by a user of the application. In another embodiment, the sample analyte concentrations are transmitted directly to the analyte input module by the sensor device used to measure the sample analyte concentration via

a data cable, infrared signal, wireless connection or other methods of data transmission known in the art.

[0090] The comparison module compares each sample analyte concentration to an entry of a renal disorder database. Each entry of the renal disorder database includes a list of minimum diagnostic concentrations reflective of a particular renal disorder. The entries of the renal disorder database may further contain additional minimum diagnostic concentrations to further define diagnostic criteria including but not limited to minimum diagnostic concentrations for additional types of bodily fluids, additional types of mammals, and severities of a particular disorder.

[0091] The analysis module determines a most likely renal disorder by combining the particular renal disorders identified by the comparison module for all of the sample analyte concentrations. In one embodiment, the most likely renal disorder is the particular renal disorder from the database entry having the most minimum diagnostic concentrations that are less than the corresponding sample analyte concentrations. In another embodiment, the most likely renal disorder is the particular renal disorder from the database entry having minimum diagnostic concentrations that are all less than the corresponding sample analyte concentrations. In yet other embodiments, the analysis module combines the sample analyte concentrations algebraically to calculate a combined sample analyte concentration that is compared to a combined minimum diagnostic concentration calculated from the corresponding minimum diagnostic criteria using the same algebraic operations. Other combinations of sample analyte concentrations from within the same test sample, or combinations of sample analyte concentrations from two or more different test samples containing two or more different bodily fluids may be used to determine a particular renal disorder in still other embodiments.

[0092] The system includes one or more processors and volatile and/or nonvolatile memory and can be embodied by or in one or more distributed or integrated components or systems. The system may include computer readable media (CRM) on which one or more algorithms, software, modules, data, and/or firmware is loaded and/or operates and/or which operates on the one or more processors to implement the systems and methods identified herein. The computer readable media may include volatile media, nonvolatile media, removable media, non-removable media, and/or other media or mediums that can be accessed by a general purpose or special purpose computing device. For example, computer readable media may include computer storage media and communication media, including but not limited to computer readable media. Computer storage media further may include volatile, nonvolatile, removable, and/or non-removable media implemented in a method or technology for storage of information, such as computer readable instructions, data structures, program modules, and/or other data. Communication media may, for example, embody computer readable instructions, data structures, program modules, algorithms, and/or other data, including but not limited to as or in a modulated data signal. The communication media may be embodied in a carrier wave or other transport mechanism and may include an information delivery method. The communication media may include wired and wireless connections and technologies and may be used to transmit and/or receive wired or wireless communications. Combinations and/or

sub-combinations of the above and systems, components, modules, and methods and processes described herein may be made.

[0093] The following examples are included to demonstrate preferred embodiments of the invention.

EXAMPLES

[0094] The following examples illustrate various iterations of the invention.

Example 1

Least Detectable Dose and Lower Limit of Quantitation of Assay for Analytes Associated with Renal Disorders

[0095] To assess the least detectable doses (LDD) and lower limits of quantitation (LLOQ) of a variety of analytes associated with renal disorders, the following experiment was conducted. The analytes measured were alpha-1 microglobulin (A1M), beta-2 microglobulin (B2M), calbindin, clusterin, CTGF, cystatin C, GST-alpha, KIM-1, NGAL, osteopontin (OPN), THP, TIMP-1, TFF-3, and VEGF.

[0096] The concentrations of the analytes were measured using a capture-sandwich assay using antigen-specific antibodies. For each analyte, a range of standard sample dilutions ranging over about four orders of magnitude of analyte concentration were measured using the assay in order to obtain data used to construct a standard dose response curve. The dynamic range for each of the analytes, defined herein as the range of analyte concentrations measured to determine its dose response curve, is presented below.

[0097] To perform the assay, 5 μL of a diluted mixture of capture-antibody microspheres were mixed with 5 μL of blocker and 10 μL of pre-diluted standard sample in each of the wells of a hard-bottom microtiter plate. After incubating the hard-bottom plate for 1 hour, 10 μL of biotinylated detection antibody was added to each well, and then the hard-bottom plate was incubated for an additional hour. 10 μL of diluted streptavidin-phycoerythrin was added to each well and then the hard-bottom plate was incubated for another 60 minutes.

[0098] A filter-membrane microtiter plate was pre-wetted by adding 100 μL wash buffer, and then aspirated using a vacuum manifold device. The contents of the wells of the hard-bottom plate were then transferred to the corresponding wells of the filter-membrane plate. All wells of the hard-bottom plate were vacuum-aspirated and the contents were washed twice with 100 μL of wash buffer. After the second wash, 100 μL of wash buffer was added to each well, and then the washed microspheres were resuspended with thorough mixing. The plate was then analyzed using a Luminex 100 Analyzer (Luminex Corporation, Austin, Tex., USA). Dose response curves were constructed for each analyte by curve-fitting the median fluorescence intensity (MFI) measured from the assays of diluted standard samples containing a range of analyte concentrations.

[0099] The least detectable dose (LDD) was determined by adding three standard deviations to the average of the MFI signal measured for 20 replicate samples of blank standard solution (i.e. standard solution containing no analyte). The MFI signal was converted to an LDD concentration using the dose response curve and multiplied by a dilution factor of 2.

[0100] The lower limit of quantification (LLOQ), defined herein as the point at which the coefficient of variation (CV) for the analyte measured in the standard samples was 30%, was determined by the analysis of the measurements of increasingly diluted standard samples. For each analyte, the standard solution was diluted by 2 fold for 8 dilutions. At each stage of dilution, samples were assayed in triplicate, and the CV of the analyte concentration at each dilution was calculated and plotted as a function of analyte concentration. The LLOQ was interpolated from this plot and multiplied by a dilution factor of 2.

[0101] The LDD and LLOQ results for each analyte are summarized in Table 2:

TABLE 2

LDD, LLOQ, and Dynamic Range of Analyte Assay					
Analyte	Units	LDD	LLOQ	Dynamic Range	
				minimum	maximum
Calbindin	ng/mL	1.1	3.1	0.516	2580
Clusterin	ng/mL	2.4	2.3	0.676	3378
CTGF	ng/mL	1.3	3.8	0.0794	400
GST-alpha	ng/mL	1.4	3.6	0.24	1,200
KIM-1	ng/mL	0.016	0.028	0.00478	24
VEGF	pg/mL	4.4	20	8.76	44,000
β-2M	μg/mL	0.012	0.018	0.0030	15
Cystatin C	ng/mL	2.8	3.7	0.60	3,000
NGAL	ng/mL	4.1	7.8	1.2	6,000
Osteopontin	ng/mL	29	52	3.9	19,500
TIMP-1	ng/mL	0.71	1.1	0.073	365
A-1M	μg/mL	0.059	0.29	0.042	210
THP	μg/mL	0.46	0.30	0.16	800
TFF-3	μg/mL	0.06	0.097	0.060	300

[0102] The results of this experiment characterized the least detectible dose and the lower limit of quantification for fourteen analytes associated with various renal disorders using a capture-sandwich assay.

Example 2

Precision of Assay for Analytes Associated with Renal Disorders

[0103] To assess the precision of an assay used to measure the concentration of analytes associated with renal disorders, the following experiment was conducted. The analytes measured were alpha-1 microglobulin (A1M), beta-2 microglobulin (B2M), calbindin, clusterin, CTGF, cystatin C, GST-alpha, KIM-1, NGAL, osteopontin (OPN), THP, TIMP-1, TFF-3, and VEGF. For each analyte, three concentration levels of standard solution were measured in triplicate during three runs using the methods described in Example 1. The percent errors for each run at each concentration are presented in Table 3 for all of the analytes tested:

TABLE 3

Precision of Analyte Assay					
Analyte	Average concentration (ng/mL)	Run 1 Error (%)	Run 2 Error (%)	Run 2 Error (%)	Interrun Error (%)
	36	5	3	2	7
	281	1	6	0	3

TABLE 3-continued

Precision of Analyte Assay					
Analyte	Average concentration (ng/mL)	Run 1 Error (%)	Run 2 Error (%)	Run 2 Error (%)	Interrun Error (%)
	39	5	1	6	8
	229	1	3	0	2
CTGF	1.2	10	17	4	14
	2.5	19	19	14	14
	18	7	5	13	9
GST-alpha	3.9	14	7	5	10
	16	13	7	10	11
	42	1	16	6	8
KIM-1	0.035	2	0	5	13
	0.32	4	5	2	8
	2.9	0	5	7	4
VEGF	65	10	1	6	14
	534	9	2	12	7
	5,397	1	13	14	9
β-2M	0.040	6	1	8	5
	0.43	2	2	0	10
	6.7	6	5	11	6
Cystatin C	10.5	4	1	7	13
	49	0	0	3	9
	424	2	6	2	5
NGAL	18.1	11	3	6	13
	147	0	0	6	5
	1,070	5	1	2	5
Osteopontin	44	1	10	2	11
	523	9	9	9	7
	8,930	4	10	1	10
TIMP-1	2.2	13	6	3	13
	26	1	1	4	14
	130	1	3	1	4
A-1M	1.7	11	7	7	14
	19	4	1	8	9
	45	3	5	2	4
THP	9.4	3	10	11	11
	15	3	7	8	6
	37	4	5	0	5
TFF-3	0.3	13	3	11	12
	4.2	5	8	5	7
	1.2	3	7	0	13

[0104] The results of this experiment characterized the precision of a capture-sandwich assay for fourteen analytes associated with various renal disorders over a wide range of analyte concentrations. The precision of the assay varied between about 1% and about 15% error within a given run, and between about 5% and about 15% error between different runs. The percent errors summarized in Table 2 provide information concerning random error to be expected in an assay measurement caused by variations in technicians, measuring instruments, and times of measurement.

Example 3

Linearity of Assay for Analytes Associated with Renal Disorders

[0105] To assess the linearity of an assay used to measure the concentration of analytes associated with renal disorders, the following experiment was conducted. The analytes measured were alpha-1 microglobulin (A1M), beta-2 microglobulin (B2M), calbindin, clusterin, CTGF, cystatin C, GST-alpha, KIM-1, NGAL, osteopontin (OPN), THP, TIMP-1, TFF-3, and VEGF. For each analyte, three concentration levels of standard solution were measured in triplicate during three runs using the methods described in Example 1. Linearity of the assay used to measure each analyte was determined

by measuring the concentrations of standard samples that were serially-diluted throughout the assay range. The % recovery was calculated as observed vs. expected concentration based on the dose-response curve. The results of the linearity analysis are summarized in Table 4.

TABLE 4

Linearity of Analyte Assay				
Analyte	Dilution	Expected concentration	Observed concentration	Recovery (%)
Calbindin (ng/mL)	1:2	61	61	100
	1:4	30	32	106
	1:8	15	17	110
Clusterin (ng/mL)	1:2	41	41	100
	1:4	21	24	116
	1:8	10	11	111
CTGF (ng/mL)	1:2	1.7	1.7	100
	1:4	0.84	1.0	124
	1:8	0.42	0.51	122
GST-alpha (ng/mL)	1:2	25	25	100
	1:4	12	14	115
	1:8	6.2	8.0	129
KIM-1 (ng/mL)	1:2	0.87	0.87	100
	1:4	0.41	0.41	101
	1:8	0.21	0.19	93
VEGF (pg/mL)	1:2	2,525	2,525	100
	1:4	1,263	1,340	106
	1:8	631	686	109
β-2M (μg/mL)	1:100	0.63	0.63	100
	1:200	0.31	0.34	106
	1:400	0.16	0.17	107
Cystatin C (ng/mL)	1:100	249	249	100
	1:200	125	122	102
	1:400	62	56	110
NGAL (ng/mL)	1:100	1,435	1,435	100
	1:200	718	775	108
	1:400	359	369	103
Osteopontin (ng/mL)	1:100	6,415	6,415	100
	1:200	3,208	3,275	102
	1:400	1,604	1,525	95
TIMP-1 (ng/mL)	1:100	35	35	100
	1:200	18	18	100
	1:400	8.8	8.8	100
A-1M (μg/mL)	1:2000	37	37	100
	1:4000	18	18	99
	1:8000	9.1	9.2	99
THP (μg/mL)	1:2000	28	28	100
	1:4000	14	14	96
	1:8000	6.7	7.1	94
TFF-3 (μg/mL)	1:2000	8.8	8.8	100
	1:4000	3.8	4.4	86
	1:8000	1.9	2.2	86

[0106] The results of this experiment demonstrated reasonably linear responses of the sandwich-capture assay to variations in the concentrations of the analytes in the tested samples.

Example 4

Spike Recovery of Analytes Associated with Renal Disorders

[0107] To assess the recovery of analytes spiked into urine, serum, and plasma samples by an assay used to measure the concentration of analytes associated with renal disorders, the following experiment was conducted. The analytes measured were alpha-1 microglobulin (A1M), beta-2 microglobulin (B2M), calbindin, clusterin, CTGF, cystatin C, GST-alpha, KIM-1, NGAL, osteopontin (OPN), THP, TIMP-1, TFF-3, and VEGF. For each analyte, three concentration levels of

standard solution were spiked into known urine, serum, and plasma samples. Prior to analysis, all urine samples were diluted 1:2000 (sample:diluent), all plasma samples were diluted 1:5 (sample:diluent), and all serum samples were diluted 1:2000 (sample:diluent).

[0108] The concentrations of the analytes in the samples were measured using the methods described in Example 1. The average % recovery was calculated as the proportion of the measurement of analyte spiked into the urine, serum, or plasma sample (observed) to the measurement of analyte spiked into the standard solution (expected). The results of the spike recovery analysis are summarized in Table 5.

TABLE 5

Spike Recovery of Analyte Assay in Urine, Serum, and Plasma Samples				
Analyte	Spike Concentration	Recovery in Urine Sample (%)	Recovery in Serum Sample (%)	Recovery in Plasma Sample (%)
Calbindin (ng/mL)	66	76	82	83
	35	91	77	71
	18	80	82	73
	average	82	80	76
Clusterin (ng/mL)	80	72	73	75
	37	70	66	72
	20	90	73	70
	average	77	70	72
CTGF (ng/mL)	8.4	91	80	79
	4.6	114	69	78
	2.4	76	80	69
	average	94	77	75
GST-alpha (ng/mL)	27	75	84	80
	15	90	75	81
	7.1	82	84	72
	average	83	81	78
KIM-1 (ng/mL)	0.63	87	80	83
	.029	119	74	80
	0.14	117	80	78
	average	107	78	80
VEGF (pg/mL)	584	88	84	82
	287	101	77	86
	123	107	84	77
	average	99	82	82
β-2M (μg/mL)	0.97	117	98	98
	0.50	124	119	119
	0.24	104	107	107
	average	115	108	105
Cystatin C (ng/mL)	183	138	80	103
	90	136	97	103
	40	120	97	118
	average	131	91	108
NGAL (ng/mL)	426	120	105	111
	213	124	114	112
	103	90	99	113
	average	111	106	112
Osteopontin (ng/mL)	1,245	204	124	68
	636	153	112	69
	302	66	103	67
	average	108	113	68
TIMP-1 (ng/mL)	25	98	97	113
	12	114	89	103
	5.7	94	99	113
	average	102	95	110
A-1M (μg/mL)	0.0028	100	101	79
	0.0012	125	80	81
	0.00060	118	101	82
	Average	114	94	81
THP (μg/mL)	0.0096	126	108	90
	0.0047	131	93	91
	0.0026	112	114	83
	average	123	105	88

TABLE 5-continued

Spike Recovery of Analyte Assay in Urine, Serum, and Plasma Samples				
Analyte	Spike Concentration	Recovery in Urine Sample (%)	Recovery in Serum Sample (%)	Recovery in Plasma Sample (%)
TFF-3 ($\mu\text{g/mL}$)	0.0038	105	114	97
	0.0019	109	104	95
	0.0010	102	118	93
	average	105	112	95

[0109] The results of this experiment demonstrated that the sandwich-type assay is reasonably sensitive to the presence of all analytes measured, whether the analytes were measured in standard samples, urine samples, plasma samples, or serum samples.

Example 5

Matrix Interferences of Analytes Associated with Renal Disorders

[0110] To assess the matrix interference of hemoglobin, bilirubin, and triglycerides spiked into standard samples, the following experiment was conducted. The analytes measured were alpha-1 microglobulin (A1M), beta-2 microglobulin (B2M), calbindin, clusterin, CTGF, cystatin C, GST-alpha, KIM-1, NGAL, osteopontin (OPN), THP, TIMP-1, TFF-3, and VEGF. For each analyte, three concentration levels of standard solution were spiked into known urine, serum, and plasma samples. Matrix interference was assessed by spiking hemoglobin, bilirubin, and triglyceride into standard analyte samples and measuring analyte concentrations using the methods described in Example 1. A % recovery was determined by calculating the ratio of the analyte concentration measured from the spiked sample (observed) divided by the analyte concentration measured from the standard sample (expected). The results of the matrix interference analysis are summarized in Table 6.

TABLE 6

Matrix Interference of Hemoglobin, Bilirubin, and Triglyceride on the Measurement of Analytes			
Analyte	Matrix Compound Spiked into Sample	Maximum Spike Concentration	Overall Recovery (%)
Calbindin (mg/mL)	Hemoglobin	500	110
	Bilirubin	20	98
	Triglyceride	500	117
Clusterin (mg/mL)	Hemoglobin	500	125
	Bilirubin	20	110
	Triglyceride	500	85
CTGF (mg/mL)	Hemoglobin	500	91
	Bilirubin	20	88
	Triglyceride	500	84
GST-alpha (mg/mL)	Hemoglobin	500	100
	Bilirubin	20	96
	Triglyceride	500	96
KIM-1 (mg/mL)	Hemoglobin	500	108
	Bilirubin	20	117
	Triglyceride	500	84
VEGF (mg/mL)	Hemoglobin	500	112
	Bilirubin	20	85
	Triglyceride	500	114

TABLE 6-continued

Matrix Interference of Hemoglobin, Bilirubin, and Triglyceride on the Measurement of Analytes			
Analyte	Matrix Compound Spiked into Sample	Maximum Spike Concentration	Overall Recovery (%)
β -2M ($\mu\text{g/mL}$)	Hemoglobin	500	84
	Bilirubin	20	75
	Triglyceride	500	104
Cystatin C (ng/mL)	Hemoglobin	500	91
	Bilirubin	20	102
	Triglyceride	500	124
NGAL (ng/mL)	Hemoglobin	500	99
	Bilirubin	20	92
	Triglyceride	500	106
Osteopontin (ng/mL)	Hemoglobin	500	83
	Bilirubin	20	86
	Triglyceride	500	106
TIMP-1 (ng/mL)	Hemoglobin	500	87
	Bilirubin	20	86
	Triglyceride	500	93
A-1M ($\mu\text{g/mL}$)	Hemoglobin	500	103
	Bilirubin	20	110
	Triglyceride	500	112
THP ($\mu\text{g/mL}$)	Hemoglobin	500	108
	Bilirubin	20	101
	Triglyceride	500	121
TFF-3 ($\mu\text{g/mL}$)	Hemoglobin	500	101
	Bilirubin	20	101
	Triglyceride	500	110

[0111] The results of this experiment demonstrated that hemoglobin, bilirubin, and triglycerides, three common compounds found in urine, plasma, and serum samples, did not significantly degrade the ability of the sandwich-capture assay to detect any of the analytes tested.

Example 6

Sample Stability of Analytes Associated with Renal Disorders

[0112] To assess the ability of analytes spiked into urine, serum, and plasma samples to tolerate freeze-thaw cycles, the following experiment was conducted. The analytes measured were alpha-1 microglobulin (A1M), beta-2 microglobulin (B2M), calbindin, clusterin, CTGF, cystatin C, GST-alpha, KIM-1, NGAL, osteopontin (OPN), THP, TIMP-1, TFF-3, and VEGF. Each analyte was spiked into known urine, serum, and plasma samples at a known analyte concentration. The concentrations of the analytes in the samples were measured using the methods described in Example 1 after the initial addition of the analyte, and after one, two and three cycles of freezing and thawing. In addition, analyte concentrations in urine, serum and plasma samples were measured immediately after the addition of the analyte to the samples as well as after storage at room temperature for two hours and four hours, and after storage at 4° C. for 2 hours, four hours, and 24 hours.

[0113] The results of the freeze-thaw stability analysis are summarized in Table 7. The % recovery of each analyte was calculated as a percentage of the analyte measured in the sample prior to any freeze-thaw cycles.

TABLE 7

Freeze-Thaw Stability of the Analytes in Urine, Serum, and Plasma							
Analyte	Period and Temp	Urine Sample		Serum Sample		Plasma Sample	
		Concentration	Recovery (%)	Concentration	Recovery (%)	Concentration	Recovery (%)
Calbindin (ng/mL)	Control	212	100	31	100	43	100
	1X	221	104	30	96	41	94
	2X	203	96	30	99	39	92
	3X	234	110	30	97	40	93
Clusterin (ng/mL)	0	315	100	232	100	187	100
	1X	329	104	227	98	177	95
	2X	341	108	240	103	175	94
	3X	379	120	248	107	183	98
CTGF (ng/mL)	0	6.7	100	1.5	100	1.2	100
	1X	7.5	112	1.3	82	1.2	94
	2X	6.8	101	1.4	90	1.2	100
	3X	7.7	115	1.2	73	1.3	107
GST-alpha (ng/mL)	0	12	100	23	100	11	100
	1X	13	104	24	105	11	101
	2X	14	116	21	92	11	97
	3X	14	111	23	100	12	108
KIM-1 (ng/mL)	0	1.7	100	0.24	100	0.24	100
	1X	1.7	99	0.24	102	0.22	91
	2X	1.7	99	0.22	94	0.19	78
	3X	1.8	107	0.23	97	0.22	93
VEGF (pg/mL)	0	1,530	100	1,245	100	674	100
	1X	1,575	103	1,205	97	652	97
	2X	1,570	103	1,140	92	612	91
	3X	1,700	111	1,185	95	670	99
β-2 M (μg/mL)	0	0.0070	100	1.2	100	15	100
	1X	0.0073	104	1.1	93	14	109
	2X	0.0076	108	1.2	103	15	104
	3X	0.0076	108	1.1	97	13	116
Cystatin C (ng/mL)	0	1,240	100	1,330	100	519	100
	1X	1,280	103	1,470	111	584	113
	2X	1,410	114	1,370	103	730	141
	3X	1,420	115	1,380	104	589	113
NGAL (ng/mL)	0	45	100	245	100	84	100
	1X	46	102	179	114	94	112
	2X	47	104	276	113	91	108
	3X	47	104	278	113	91	109
Osteopontin (ng/mL)	0	38	100	1.7	100	5.0	100
	1X	42	110	1.8	102	5.5	110
	2X	42	108	1.5	87	5.5	109
	3X	42	110	1.3	77	5.4	107
TIMP-1 (ng/mL)	0	266	100	220	100	70	100
	1X	265	100	220	10	75	108
	2X	255	96	215	98	77	110
	3X	295	111	228	104	76	109
A-1 M (μg/mL)	0	14	100	26	100	4.5	100
	1X	13	92	25	96	4.2	94
	2X	15	107	25	96	4.3	97
	3X	16	116	23	88	4.0	90
THP (μg/mL)	0	4.6	100	31	100	9.2	100
	1X	4.4	96	31	98	8.8	95
	2X	5.0	110	31	100	9.2	100
	3X	5.2	114	27	85	9.1	99
TFF-3 (μg/mL)	0	4.6	100	24	100	22	100
	1X	4.4	96	23	98	22	103
	2X	5.0	110	24	103	22	101
	3X	5.2	114	19	82	22	102

[0114] The results of the short-term stability assessment are summarized in Table 8. The % recovery of each analyte was calculated as a percentage of the analyte measured in the sample prior to any short-term storage.

TABLE 8

Short-Term Stability of Analytes in Urine, Serum, and Plasma							
Analyte	Storage Time/ Temp	Urine Sample		Serum Sample		Plasma Sample	
		Sample Conc.	Recovery (%)	Sample Conc.	Recovery (%)	Sample Conc.	Recovery (%)
Cal- bindin (ng/mL)	Control	226	100	33	100	7	100
	2 hr/room temp	242	107	30	90	6.3	90
	2 hr. @ 4° C.	228	101	29	89	6.5	93
	4 hr @ room temp	240	106	28	84	5.6	79
	4 hr. @ 4° C.	202	89	29	86	5.5	79
	24 hr. @ 4° C.	199	88	26	78	7.1	101
Clus- terin (ng/mL)	Control	185	100	224	100	171	100
	2 hr @ room temp	173	94	237	106	180	105
	2 hr. @ 4° C.	146	79	225	100	171	100
	4 hr @ room temp	166	89	214	96	160	94
	4 hr. @ 4° C.	157	85	198	88	143	84
	24 hr. @ 4° C.	185	100	207	92	162	94
CTGF (ng/mL)	Control	1.9	100	8.8	100	1.2	100
	2 hr @ room temp	1.9	99	6.7	76	1	83
	2 hr. @ 4° C.	1.8	96	8.1	92	1.1	89
	4 hr @ room temp	2.1	113	5.6	64	1	84
	4 hr. @ 4° C.	1.7	91	6.4	74	0.9	78
	24 hr. @ 4° C.	2.2	116	5.9	68	1.1	89
GST- alpha (ng/mL)	Control	14	100	21	100	11	100
	2 hr @ room temp	11	75	23	107	11	103
	2 hr. @ 4° C.	13	93	22	104	9.4	90
	4 hr @ room temp	11	79	21	100	11	109
	4 hr. @ 4° C.	12	89	21	98	11	100
	24 hr. @ 4° C.	13	90	22	103	14	129
KIM-1 (ng/mL)	Control	1.5	100	0.23	100	0.24	100
	2 hr @ room temp	1.2	78	0.2	86	0.22	90
	2 hr. @ 4° C.	1.6	106	0.23	98	0.21	85
	4 hr @ room temp	1.3	84	0.19	82	0.2	81
	4 hr. @ 4° C.	1.4	90	0.22	93	0.19	80
	24 hr. @ 4° C.	1.1	76	0.18	76	0.23	94
VEGF (pg/mL)	Control	851	100	1215	100	670	100
	2 hr @ room temp	793	93	1055	87	622	93
	2 hr. @ 4° C.	700	82	1065	88	629	94
	4 hr @ room temp	704	83	1007	83	566	84
	4 hr. @ 4° C.	618	73	1135	93	544	81

TABLE 8-continued

Short-Term Stability of Analytes in Urine, Serum, and Plasma							
Analyte	Storage Time/ Temp	Urine Sample		Serum Sample		Plasma Sample	
		Sample Conc.	Recovery (%)	Sample Conc.	Recovery (%)	Sample Conc.	Recovery (%)
β -2 M (μ g/mL)	24 hr. @ 4° C.	653	77	1130	93	589	88
	Control	0.064	100	2.6	100	1.2	100
	2 hr @ room temp	0.062	97	2.4	92	1.1	93
	2 hr. @ 4° C.	0.058	91	2.2	85	1.2	94
	4 hr @ room temp	0.064	101	2.2	83	1.2	94
	4 hr. @ 4° C.	0.057	90	2.2	85	1.2	98
	24 hr. @ 4° C.	0.06	94	2.5	97	1.3	103
	Control	52	100	819	100	476	100
Cys- tatin C (ng/mL)	2 hr @ room temp	50	96	837	102	466	98
	2 hr. @ 4° C.	44	84	884	108	547	115
	4 hr @ room temp	49	93	829	101	498	105
	4 hr. @ 4° C.	46	88	883	108	513	108
	24 hr. @ 4° C.	51	97	767	94	471	99
	Control	857	100	302	100	93	100
	2 hr @ room temp	888	104	287	95	96	104
	2 hr. @ 4° C.	923	108	275	91	92	100
NGAL (ng/mL)	4 hr @ room temp	861	101	269	89	88	95
	4 hr. @ 4° C.	842	98	283	94	94	101
	24 hr. @ 4° C.	960	112	245	81	88	95
	Control	2243	100	6.4	100	5.2	100
	2 hr @ room temp	2240	100	6.8	107	5.9	114
	2 hr. @ 4° C.	2140	95	6.4	101	6.2	120
	4 hr @ room temp	2227	99	6.9	108	5.8	111
	4 hr. @ 4° C.	2120	95	7.7	120	5.2	101
Osteo- pontin (ng/mL)	24 hr. @ 4° C.	2253	100	6.5	101	6	116
	Control	17	100	349	100	72	100
	2 hr @ room temp	17	98	311	89	70	98
	2 hr. @ 4° C.	16	94	311	89	68	95
	4 hr @ room temp	17	97	306	88	68	95
	4 hr. @ 4° C.	16	93	329	94	74	103
	24 hr. @ 4° C.	18	105	349	100	72	100
	Control	17	100	349	100	72	100
TIMP-1 (ng/mL)	2 hr @ room temp	17	98	311	89	70	98
	2 hr. @ 4° C.	16	94	311	89	68	95
	4 hr @ room temp	17	97	306	88	68	95
	4 hr. @ 4° C.	16	93	329	94	74	103
	24 hr. @ 4° C.	18	105	349	100	72	100
	Control	3.6	100	2.2	100	1	100
	2 hr @ room temp	3.5	95	2	92	1	105
	2 hr. @ 4° C.	3.4	92	2.1	97	0.99	99
A-1 M (μ g/mL)	4 hr @ room temp	3.2	88	2.2	101	0.99	96
	4 hr. @ 4° C.	3	82	2.2	99	0.97	98
	24 hr. @ 4° C.	3	83	2.2	100	1	101

TABLE 8-continued

Short-Term Stability of Analytes in Urine, Serum, and Plasma							
Analyte	Storage Time/ Temp	Urine Sample		Serum Sample		Plasma Sample	
		Sample Conc.	Recovery (%)	Sample Conc.	Recovery (%)	Sample Conc.	Recovery (%)
THP ($\mu\text{g/mL}$)	Control	1.2	100	34	100	2.1	100
	2 hr @ room temp	1.2	99	34	99	2	99
	2 hr @ 4° C.	1.1	90	34	100	2	98
	4 hr @ room temp	1.1	88	27	80	2	99
	4 hr @ 4° C.	0.95	79	33	97	2	95
	24 hr @ 4° C.	0.91	76	33	98	2.4	116
	TFF-3 ($\mu\text{g/mL}$)	Control	1230	100	188	100	2240
2 hr @ room temp	1215	99	179	95	2200	98	
2 hr @ 4° C.	1200	98	195	104	2263	101	
4 hr @ room temp	1160	94	224	119	2097	94	
4 hr @ 4° C.	1020	83	199	106	2317	103	
24 hr @ 4° C.	1030	84	229	122	1940	87	

[0115] The results of this experiment demonstrated that the analytes associated with renal disorders tested were suitably stable over several freeze/thaw cycles, and up to 24 hrs of storage at a temperature of 4° C.

Example 8

Analysis of Kidney Biomarkers in Plasma and Urine from Patients with Renal Injury

[0116] A screen for potential protein biomarkers in relation to kidney toxicity/damage was performed using a panel of biomarkers, in a set of urine and plasma samples from patients with documented renal damage. The investigated patient groups included diabetic nephropathy (DN), obstructive uropathy (OU), analgesic abuse (AA) and glomerulonephritis (GN) along with age, gender and BMI matched control groups. Multiplexed immunoassays were applied in order to quantify the following protein analytes: Alpha-1 Microglobulin (α1M), KIM-1, Microalbumin, Beta-2-Microglobulin (β2M), Calbindin, Clusterin, CystatinC, TrefoilFactor-3 (TFF-3), CTGF, GST-alpha, VEGF, Calbindin, Osteopontin, Tamm-HorsfallProtein (THP), TIMP-1 and NGAL.

[0117] Li-Heparin plasma and mid-stream spot urine samples were collected from four different patient groups. Samples were also collected from age, gender and BMI matched control subjects. 20 subjects were included in each

group resulting in a total number of 160 urine and plasma samples. All samples were stored at -80°C . before use. Glomerular filtration rate for all samples was estimated using two different estimations (Modification of Diet in Renal Disease or MDRD, and the Chronic Kidney Disease Epidemiology Collaboration or CKD-EPI) to outline the eGFR (estimated glomerular filtration rate) distribution within each patient group (FIG. 1). Protein analytes were quantified in human plasma and urine using multiplexed immunoassays in the Luminex xMAP™ platform. The microsphere-based multiplex immunoassays consist of antigen-specific antibodies and optimized reagents in a capture-sandwich format. Output data was given as g/ml calculated from internal standard curves. Because urine creatinine (uCr) correlates with renal filtration rate, data analysis was performed without correction for uCr. Univariate and multivariate data analysis was performed comparing all case vs. control samples as well as cases vs. control samples for the various disease groups.

[0118] The majority of the measured proteins showed a correlation to eGFR. Measured variables were correlated to eGFR using Pearson's correlations coefficient, and samples from healthy controls and all disease groups were included in the analysis. 11 and 7 proteins displayed P-values below 0.05 for plasma and urine (Table 9) respectively.

TABLE 9

Correlation analysis of eGFR and variables for all case samples					
URINE			PLASMA		
Variable	Pearson's r	P-Value	Variable	Pearson's r	P-Value
Alpha-1-Microglobulin	-0.08	0.3	Alpha-1-Microglobulin	-0.33	< 0.001

TABLE 9-continued

Correlation analysis of eGFR and variables for all case samples					
URINE			PLASMA		
Variable	Pearson's r	P-Value	Variable	Pearson's r	P-Value
Beta-2-Microglobulin	-0.23	0.003	Beta-2-Microglobulin	-0.39	< 0.001
Calbindin	-0.16	0.04	Calbindin	-0.18	< 0.02
Clusterin	-0.07	0.4	Clusterin	-0.51	< 0.001
CTGF	-0.08	0.3	CTGF	-0.05	0.5
Creatinine	-0.32	< 0.001	Cystatin-C	-0.42	< 0.001
Cystatin-C	-0.24	0.002	GST-alpha	-0.12	0.1
GST-alpha	-0.11	0.2	KIM-1	-0.17	0.03
KIM-1	-0.08	0.3	NGAL	-0.28	<0.001
Microalbumin_UR	-0.17	0.03	Osteopontin	-0.33	< 0.001
NGAL	-0.15	0.07	THP	-0.31	< 0.001
Osteopontin	-0.19	0.02	TIMP-1	-0.28	<0.001
THP	-0.05	0.6	TFF3	-0.38	< 0.001
TIMP-1	-0.19	0.01	VEGF	-0.14	0.08
TFF2	-0.09	0.3			
VEGF	-0.07	0.4			

P values <0.0001 are shown in bold italics

P values <0.005 are shown in bold

P values <0.05 are shown in italics

[0119] For the various disease groups, univariate statistical analysis revealed that in a direct comparison (T-test) between cases and controls, a number of proteins were differentially expressed in both urine and plasma (Table 10 and FIG. 2). In particular, clusterin showed a marked differential pattern in plasma.

TABLE 10

Differentially regulated proteins by univariate statistical analysis				
Group	Matrix	Protein	p-value	
AA	Urine	Calbindin	0.016	
AA	Urine	NGAL	0.04	
AA	Urine	Osteopontin	0.005	
AA	Urine	Creatinine	0.001	
AA	Plasma	Calbindin	0.05	
AA	Plasma	Clusterin	0.003	
AA	Plasma	KIM-1	0.03	
AA	Plasma	THP	0.001	
AA	Plasma	TIMP-1	0.02	
DN	Urine	Creatinine	0.04	
DN	Plasma	Clusterin	0.006	
DN	Plasma	KIM-1	0.01	
GN	Urine	Creatinine	0.004	
GN	Urine	Microalbumin	0.0003	
GN	Urine	NGAL	0.05	
GN	Urine	Osteopontin	0.05	
GN	Urine	TFF3	0.03	
GN	Plasma	Alpha 1 Microglobulin	0.002	
GN	Plasma	Beta 2 Microglobulin	0.03	
GN	Plasma	Clusterin	0.00	
GN	Plasma	Cystatin C	0.01	
GN	Plasma	KIM-1	0.003	
GN	Plasma	NGAL	0.03	
GN	Plasma	THP	0.001	
GN	Plasma	TIMP-1	0.003	
GN	Plasma	TFF3	0.01	
GN	Plasma	VEGF	0.02	
OU	Urine	Clusterin	0.02	
OU	Urine	Microalbumin	0.007	
OU	Plasma	Clusterin	0.00	

[0120] Application of multivariate analysis yielded statistical models that predicted disease from control samples (plasma results are shown in FIG. 3)

[0121] In conclusion, these results form a valuable base for further studies on these biomarkers in urine and plasma both regarding baseline levels in normal populations and regarding the differential expression of the analytes in various disease groups. Using this panel of analytes, error rates from ada-boosting and/or random forest were low enough (<10%) to allow a prediction model to differentiate between control and disease patient samples. Several of the analytes showed a greater correlation to eGFR in plasma than in urine.

Example 9

Statistical Analysis of Kidney Biomarkers in Plasma and Urine from Patients with Renal Injury

[0122] Urine and plasma samples were taken from 80 normal control group subjects and 20 subjects from each of four disorders: analgesic abuse, diabetic nephropathy, glomerulonephritis, and obstructive uropathy. The samples were analyzed for the quantity and presence of 16 different proteins (alpha-1 microglobulin (α 1M), beta-2 microglobulin (β 2M), calbindin, clusterin, CTGF, creatinine, cystatin C, GST-alpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF) as described in Example 1 above. The goal was to determine the analytes that distinguish between a normal sample and a diseased sample, a normal sample and an obstructive uropathy (OU) sample, and finally, an obstructive uropathy sample from the other disease samples (diabetic nephropathy (DN), analgesic abuse (AA), and glomerulonephritis (GN)).

[0123] From the above protein analysis data, bootstrap analysis was used to estimate the future performance of several classification algorithms. For each bootstrap run, training data and testing data was randomly generated. Then, the following algorithms were applied on the training data to generate models and then apply the models to the testing data to make predictions: automated Matthew's classification

algorithm, classification and regression tree (CART), conditional inference tree, bagging, random forest, boosting, logistic regression, SVM, and Lasso. The accuracy rate and ROC areas were recorded for each method on the prediction of the testing data. The above was repeated 100 times. The mean and the standard deviation of the accuracy rates and of the ROC areas were calculated.

[0124] The mean error rates and AUROC were calculated from urine and AUROC was calculated from plasma for 100 runs of the above method for each of the following comparisons: disease (AA+GN+OU+DN) vs. normal (FIG. 4, Table 11), OU vs. normal (FIG. 6, Table 13), OU vs. AA (FIG. 8, Table 15), OU vs. GN (FIG. 10, Table 17), and OU vs. DN (FIG. 12, Table 19).

[0125] The average relative importance of 16 different analytes (alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GST-alpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF) and 4 different clinical variables (weight, BMI, age, and gender) from 100 runs were analyzed with two different statistical methods—random forest (plasma and urine samples) and boosting (urine samples)—for each of the following comparisons: disease (AA+GN+OU+DN) vs. normal (FIG. 5, Table 12), OU vs. normal (FIG. 7, Table 14), OU vs. AA (FIG. 9, Table 16), OU vs. GN (FIG. 11, Table 18), and OU vs. DN (FIG. 13, Table 20).

TABLE 11

disease v. NL		
method	Mean AUROC	Standard deviation of AUROC
random forest	0.931	0.039
bagging	0.919	0.045
svm	0.915	0.032
boosting	0.911	0.06
lasso	0.897	0.044
logistic regression	0.891	0.041
ctree	0.847	0.046
cart	0.842	0.032
matt	0.83	0.023

TABLE 12

disease v. NL	
analyte	relative importance
Creatinine	11.606
Kidney_Injury_M	8.486
Tamm_Horsfall_P	8.191
Total_Protein	6.928
Osteopontin	6.798
Neutrophil_Gela	6.784
Tissue_Inhibito	6.765
Vascular_Endoth	6.716
Trefoil_Factor_	6.703
Cystatin_C	6.482
Alpha_1_Microgl	6.418
Beta_2_Microglo	6.228
Glutathione_S_T	6.053
clusterin	5.842

TABLE 13

OU v. NL		
method	Mean AUROC	Standard deviation of AUROC
cart	1	0
bagging	1	0
boosting	1	0
random forest	1	0.003
lasso	0.997	0.012
ctree	0.991	0.048
svm	0.917	0.079
logistic regression	0.911	0.099
matt	0.853	0.107

TABLE 14

OU v. NL	
analyte	
Creatinine	19.055
Tamm_Horsfall_P	7.682
Alpha_1_Microgl	7.664
clusterin	6.863
Neutrophil_Gela	6.825
Osteopontin	6.517
Kidney_Injury_M	6.156
Total_Protein	6.021
Vascular_Endoth	5.971
Tissue_Inhibito	5.941
Trefoil_Factor_	5.779
Cystatin_C	5.683
Glutathione_S_T	5.097
Beta_2_Microglo	4.746

TABLE 15

OU v. AA		
method	Mean AUROC	Standard deviation of AUROC
random forest	0.814	0.11
bagging	0.792	0.115
svm	0.788	0.112
lasso	0.786	0.118
boosting	0.757	0.117
matt	0.687	0.111
logistic regression	0.683	0.116
cart	0.665	0.097
ctree	0.659	0.118

TABLE 16

OU v. AA	
analyte	Relative importance
Total_Protein	11.502
Tissue_Inhibito	9.736

TABLE 16-continued

OU v. AA	
analyte	Relative importance
Cystatin_C	9.161
Alpha_1_Microgl	8.637
Trefoil_Factor_	7.329
Osteopontin	7.326
Beta_2_Microglo	6.978
Neutrophil_Gela	6.577
Glutathione_S_T	6.100
Tamm_Horsfall_P	6.066
Kidney_Injury_M	6.038
Vascular_Endoth	5.946
clusterin	4.751
Creatinine	3.854

TABLE 17

OU v. GN		
method	Mean AUROC	Standard deviation of AUROC
random	0.946	0.061
forest		
boosting	0.941	0.071
bagging	0.933	0.078
svm	0.932	0.07
lasso	0.888	0.092
cart	0.879	0.091
ctree	0.879	0.104
matt	0.872	0.099
logistic	0.869	0.134
regression		

TABLE 18

OU v. GN	
analyte	Relative importance
Total_Protein	14.391
Alpha_1_Microgl	11.033
Kidney_Injury_M	8.453
Tissue_Inhibito	7.934
Creatinine	7.714
clusterin	7.353
Osteopontin	7.123
Vascular_Endoth	6.321
Neutrophil_Gela	6.244
Cystatin_C	5.632
Beta_2_Microglo	4.962
Tamm_Horsfall_P	4.829
Trefoil_Factor_	4.594
Glutathione_S_T	3.418

TABLE 19

DN v. OU		
method	mean_AUROC	std_AUROC
lasso	0.993	0.019
random	0.986	0.027
forest		

TABLE 19-continued

DN v. OU		
method	mean_AUROC	std_AUROC
boosting	0.986	0.027
bagging	0.977	0.04
cart	0.962	0.045
ctree	0.954	0.05
svm	0.95	0.059
logistic	0.868	0.122
regression		
matt	0.862	0.111

TABLE 20

DN v. OU	
analyte	Relative importance
Creatinine	18.278
Alpha_1_Microgl	9.808
clusterin	9.002
Beta_2_Microglo	8.140
Cystatin_C	7.101
Osteopontin	6.775
Glutathione_S_T	5.731
Neutrophil_Gela	5.720
Trefoil_Factor_	5.290
Kidney_Injury_M	5.031
Total_Protein	5.030
Vascular_Endoth	4.868
Tissue_Inhibito	4.615
Tamm_Horsfall_P	4.611

[0126] It should be appreciated by those of skill in the art that the techniques disclosed in the examples above represent techniques discovered by the inventors to function well in the practice of the invention. Those of skill in the art should, however, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention, therefore all matter set forth or shown in the accompanying drawings is to be interpreted as illustrative and not in a limiting sense.

1-23. (canceled)

24. A method for diagnosing, monitoring, or determining obstructive uropathy in a mammal, the method comprising:

- a. providing a test sample comprising a sample of bodily fluid taken from the mammal;
- b. determining sample concentrations of three or more sample analytes in a panel of biomarkers in the test sample, wherein the sample analytes are selected from the group consisting of alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GSTalpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF;
- c. comparing the concentrations of the sample analytes with concentrations of the sample analytes in a panel of biomarkers from a control sample;
- d. identifying the particular disorder based on a statistically significant difference in the concentrations of the sample analytes compared the concentrations of the sample analytes from the control sample; and
- e. treating the mammal identified as having the particular disorder.

25. The method of claim 24, wherein the mammal is selected from the group consisting of humans, apes, monkeys, rats, mice, dogs, cats, pigs, and livestock including cattle and oxen.

26. The method of claim 24, wherein the bodily fluid is selected from the group consisting of urine, blood, plasma, serum, saliva, semen, and tissue lysates.

27. The method of claim 24, wherein a combination of sample concentrations for six or more sample analytes in the test sample is determined.

28. The method of claim 24, wherein a combination of sample concentrations for sixteen sample analytes in the test sample is determined.

29. A method for diagnosing, monitoring, or determining obstructive uropathy in a mammal, the method comprising:

- a. providing a test sample comprising a sample of bodily fluid taken from the mammal;
- b. determining sample concentrations of three or more sample analytes in a panel of biomarkers in the test sample, wherein the sample analytes are selected from the group consisting of alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GSTalpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF;
- c. comparing the concentrations of the sample analytes with concentrations of the sample analytes in a panel of biomarkers from a control sample; and
- d. identifying the particular disorder based on a statistically significant difference in the concentrations of the sample analytes compared the concentrations of the sample analytes from the control sample.

30. The method of claim 29, wherein the mammal is selected from the group consisting of humans, apes, monkeys, rats, mice, dogs, cats, pigs, and livestock including cattle and oxen.

31. The method of claim 29, wherein the bodily fluid is selected from the group consisting of urine, blood, plasma, serum, saliva, semen, and tissue lysates.

32. The method of claim 29, wherein a combination of sample concentrations for six or more sample analytes in the test sample is determined.

33. The method of claim 29, wherein a combination of sample concentrations for sixteen sample analytes in the test sample is determined.

34. A system for diagnosing, monitoring, or determining obstructive uropathy in a mammal, the method comprising:

- a. providing a test sample comprising a sample of bodily fluid taken from the mammal;
- b. determining sample concentrations for three or more sample analytes in the test sample, wherein the sample analytes are alpha-1 microglobulin, beta-2 microglobulin, calbindin, clusterin, CTGF, creatinine, cystatin C, GSTalpha, KIM-1, microalbumin, NGAL, osteopontin, THP, TIMP-1, TFF-3, and VEGF;
- c. inputting the sample concentrations into an analyte input module;
- d. comparing the sample concentrations to a data set comprising at least one entry stored in a comparison module,

wherein each entry of the data set comprises a list comprising three or more minimum diagnostic concentrations indicative of obstructive uropathy or an associated disorder, wherein each minimum diagnostic concentration comprises a maximum of a range of analyte concentrations for a healthy mammal;

- e. determining a matching entry of the dataset in which all minimum diagnostic concentrations are less than the corresponding sample concentrations; and,
- f. identifying an indicated disorder comprising the particular disorder of the matching entry.

35. The method of claim 34, wherein the mammal is selected from the group consisting of humans, apes, monkeys, rats, mice, dogs, cats, pigs, and livestock including cattle and oxen.

36. The method of claim 34, wherein the minimum diagnostic concentration in human plasma of alpha-1 microglobulin is about 16 $\mu\text{g/ml}$, beta-2 microglobulin is about 2.2 $\mu\text{g/ml}$, calbindin is greater than about 5 ng/ml , clusterin is about 134 $\mu\text{g/ml}$, CTGF is about 16 ng/ml , cystatin C is about 1170 ng/ml , GST-alpha is about 62 ng/ml , KIM-1 is about 0.57 ng/ml , NGAL is about 375 ng/ml , osteopontin is about 25 ng/ml , THP is about 0.052 $\mu\text{g/ml}$, TIMP-1 is about 131 ng/ml , TFF-3 is about 0.49 $\mu\text{g/ml}$, and VEGF is about 855 pg/ml .

37. The method of claim 34, wherein the minimum diagnostic concentration in human sera of alpha-1 microglobulin is about 17 $\mu\text{g/ml}$, beta-2 microglobulin is about 2.6 $\mu\text{g/ml}$, calbindin is greater than about 2.6 ng/ml , clusterin is about 152 $\mu\text{g/ml}$, CTGF is greater than about 8.2 ng/ml , cystatin C is about 1250 ng/ml , GST-alpha is about 52 ng/ml , KIM-1 is greater than about 0.35 ng/ml , NGAL is about 822 ng/ml , osteopontin is about 12 ng/ml , THP is about 0.053 $\mu\text{g/ml}$, TIMP-1 is about 246 ng/ml , TFF-3 is about 0.17 $\mu\text{g/ml}$, and VEGF is about 1630 pg/ml .

38. The method of claim 34, wherein the minimum diagnostic concentration in human urine of alpha-1 microglobulin is about 233 $\mu\text{g/ml}$, beta-2 microglobulin is greater than about 0.17 $\mu\text{g/ml}$, calbindin is about 233 ng/ml , clusterin is greater than about 0.089 $\mu\text{g/ml}$, CTGF is greater than about 0.90 ng/ml , cystatin C is about 1170 ng/ml , GST-alpha is greater than about 26 ng/ml , KIM-1 is about 0.67 ng/ml , NGAL is about 81 ng/ml , osteopontin is about 6130 ng/ml , THP is about 2.6 $\mu\text{g/ml}$, TIMP-1 is greater than about 3.9 ng/ml , TFF-3 is greater than about 21 $\mu\text{g/ml}$, and VEGF is about 517 pg/ml .

39. The method of claim 34, wherein the bodily fluid is selected from the group consisting of urine, blood, plasma, serum, saliva, semen, and tissue lysates.

40. The method of claim 34, wherein a combination of sample concentrations for six or more sample analytes in the test sample is determined.

41. The method of claim 34, wherein a combination of sample concentrations for sixteen sample analytes in the test sample is determined.

* * * * *

专利名称(译)	用于检测阻塞性尿路病和相关病症的方法和装置		
公开(公告)号	US20140303021A1	公开(公告)日	2014-10-09
申请号	US14/243659	申请日	2014-04-02
申请(专利权)人(译)	规则为基础的医学, INC.		
当前申请(专利权)人(译)	规则为基础的医学, INC.		
[标]发明人	LABRIE SAMUEL T MAPES JAMES P SPAIN MICHAEL D MCDADE RALPH L EISINGER DOMINIC P BALLARD KARRI L		
发明人	LABRIE, SAMUEL T. MAPES, JAMES P. SPAIN, MICHAEL D. MCDADE, RALPH L. EISINGER, DOMINIC P. BALLARD, KARRI L.		
IPC分类号	G01N33/566 G01N33/53		
CPC分类号	G01N33/6893 G01N33/566 G01N33/5302 G01N2800/34 G01N2333/47 G01N2333/4703 G01N2333/4706 G01N2333/4725 G01N2333/4727 G01N2333/475 G01N2333/52 G01N2333/70503 G01N2333/70539 G01N2333/765 G01N2333/775 G01N2333/8139 G01N2333/8146 G01N2333/82 G01N2333/91177 G01N2800/347 G01N2800/52 G01N2800/56 G01N2800/60 Y10T436/147777		
优先权	61/232091 2009-08-07 US 61/327389 2010-04-23 US		
外部链接	Espacenet USPTO		

摘要(译)

描述了用于诊断, 监测或确定哺乳动物中的阻塞性尿路病或相关病症的方法和装置。具体地, 描述了使用从哺乳动物取得的测试样品中测量浓度的三种或更多种分析物的组合来诊断, 监测或确定阻塞性尿路病或相关病症的方法和装置。




FIG. 1