



US 20070077251A1

(19) **United States**(12) **Patent Application Publication**
MacPhee et al.(10) **Pub. No.: US 2007/0077251 A1**(43) **Pub. Date: Apr. 5, 2007**(54) **LIPOPROTEIN ASSOCIATED
PHOSPHOLIPASE A2, INHIBITORS
THEREOF AND USE OF THE SAME IN
DIAGNOSIS AND THERAPY**

tinuation of application No. 08/387,858, filed on Feb. 24, 1995, now Pat. No. 5,981,252, filed as 371 of international application No. PCT/GB94/01374, filed on Jun. 24, 1994.

(76) Inventors: **Colin H. MacPhee**, Berwyn, PA (US);
David Graham Tew, Saffron Walden
(GB)(30) **Foreign Application Priority Data**

Jan. 11, 1994 (GB) 9400413.2
Jun. 25, 1993 (GB) 9313144.9

Correspondence Address:
LICATA & TYRRELL P.C.
66 E. MAIN STREET
MARLTON, NJ 08053 (US)

Publication Classification(21) Appl. No.: **11/603,741**

(51) **Int. Cl.**
A61K 39/395 (2006.01)
C12Q 1/34 (2006.01)
C07H 21/04 (2006.01)
C12P 21/06 (2006.01)
C12N 9/18 (2006.01)


(22) Filed: **Nov. 22, 2006****Related U.S. Application Data**

(52) **U.S. Cl.** **424/146.1**; 435/197; 435/69.1;
435/320.1; 435/325; 536/23.2;
530/388.26; 435/18

(60) Division of application No. 10/973,467, filed on Oct. 26, 2004, which is a continuation of application No. 10/173,233, filed on Jun. 14, 2002, now Pat. No. 7,052,862, which is a continuation of application No. 09/569,899, filed on May 12, 2000, which is a continuation of application No. 09/294,384, filed on Apr. 20, 1999, now Pat. No. 6,177,257, which is a con-

(57) **ABSTRACT**

The enzyme Lp-PLA₂ in purified form, an isolated nucleic acid molecule encoding Lp-PLA₂, the use of an inhibitor of the enzyme Lp-PLA₂ in therapy and a method of screening compounds to identify those compounds.



Compound 4 inhibits copper ($5\mu\text{M}$) - stimulated LDL ($150\mu\text{g/ml}$) oxidation.

FIGURE 1



Lanes 2, 4 and 6 contain adjacent fractions of purified native Lp-PLA2. Lanes 1, 3 and 5 are fractions 2, 4 and 6 respectively after N-deglycosylation.


FIGURE 2

**LIPOPROTEIN ASSOCIATED PHOSPHOLIPASE
A₂, INHIBITORS THEREOF AND USE OF THE
SAME IN DIAGNOSIS AND THERAPY**

[0001] The present invention relates to the use of inhibitors of an enzyme in the therapy, in particular in the treatment of atherosclerosis. The present invention also relates to the isolation and purification of the enzyme, to isolated nucleic acids encoding the enzyme, to recombinant host cells transformed with DNA encoding the enzyme, to the use of the enzyme in diagnosing a patient's susceptibility to atherosclerosis, and to the use of the enzyme in identifying compounds which are potentially useful for the treatment of atherosclerosis.

[0002] Lipoprotein Associated Phospholipase A₂ (Lp-PLA₂), also previously known in the art as Platelet Activating Factor Acetyl Hydrolase (PAF acetyl hydrolase). During the conversion of LDL to its oxidised form, Lp-PLA₂ is responsible for hydrolysing the sn-2 ester of oxidatively modified phosphatidylcholine to give lyso-phosphatidylcholine and an oxidatively modified fatty acid. Both of these products of Lp-PLA₂ action are potent chemoattractants for circulating monocytes. As such, this enzyme is thought to be responsible for the accumulation of cells loaded with cholesterol ester in the arteries, causing the characteristic 'fatty streak' associated with the early stages of atherosclerosis. Inhibition of the Lp-PLA₂ enzyme would therefore be expected to stop the build up of this fatty streak (by inhibition of the formation of lysophosphatidylcholine), and so be useful in the treatment of atherosclerosis. In addition, it is proposed that Lp-PLA₂ plays a direct role in LDL oxidation. This is due to the poly unsaturated fatty acid-derived lipid peroxide products of Lp-PLA₂ action contributing to and enhancing the overall oxidative process. In keeping with this idea, Lp-PLA₂ inhibitors inhibit LDL oxidation. Lp-PLA₂ inhibitors may therefore have a general application in any disorder that involves lipid peroxidation in conjunction with the enzyme activity, for example in addition to conditions such as atherosclerosis and diabetes other conditions such as rheumatoid arthritis, stroke, myocardial infarction, reperfusion injury and acute and chronic inflammation.

[0003] The present invention therefore provides in a first aspect an inhibitor of the enzyme lipoprotein associated Lp-PLA₂ for use in therapy, in particular in the treatment of atherosclerosis. Suitable compounds able to inhibit the Lp-PLA₂ enzyme are known in the art and include for example, the following compounds of structure (I):



(I)

in which R is C₁₋₆alkylCONR²;

[0004] R² is hydrogen or C₁₋₆alkyl;

[0005] X is oxygen, sulphur or —O(CO)—;

[0006] R¹ is C₈₋₂₀alkyl;

[0007] Z is N(R³)₂, [⊕]N(R³)₃, SR³, [⊖](R³)₂, in which each group R³ is the same or different and is C₁₋₆ alkyl, OR², C₁₋₄alkanoyl, imidazolyl or N-methylimidazolyl

[0008] Suitably R² is hydrogen or C₁₋₆ alkyl; preferably R² is hydrogen.

[0009] Suitably X is oxygen, sulphur or —O(CO)—; preferably X is oxygen

[0010] Suitably R¹ is C₈₋₂₀alkyl; preferably R¹ is C₁₆₋₁₈ alkyl

[0011] Suitably Z is N(R³)₂, [⊕]N(R³)₃, [⊕]S(R³)₂, in which each group R³ is the same or different and is C₁₋₆ alkyl, OR², C₁₋₄alkanoyl, imidazolyl or N-methylimidazolyl; preferably Z is SR³ in which R³ is methyl or OR² in which R² is hydrogen

[0012] The compounds of structure (I) can be prepared by processes known to those skilled in the art, for example as described in J Chem Soc Chem Comm.,1993, 70-72; J Org Chem, 1983, 48, 1197 and Chem Phys Lipids, 1984,35,29-37 or procedures analogous thereto.

[0013] When used in therapy, the compounds of structure (I) are formulated in accordance with standard pharmaceutical practice.

[0014] The compounds of structure (I) and their pharmaceutically acceptable salts which are active when given orally can be formulated as liquids, for example syrups, suspensions or emulsions, tablets, capsules and, lozenges.

[0015] A liquid formulation will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable liquid carrier(s) for example, ethanol, glycerine, non-aqueous solvent, for example polyethylene glycol oils, or water with a suspending agent, preservative, flavouring or colouring agent.

[0016] A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

[0017] A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

[0018] Typical parenteral compositions consist of a solution or suspension of the compound or pharmaceutically acceptable salt in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

[0019] A typical suppository formulation comprises a compound of formula (I) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent such as polymeric

glycols, gelatins or cocoa butter or other low-melting vegetable or synthetic waxes or fats. Preferably the composition is in unit dose form such as a tablet or capsule.

[0020] Each dosage unit for oral administration contains preferably from 1 to 250 mg (and for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of the formula (I) or a pharmaceutically acceptable salt thereof calculated as the free base.

[0021] The daily dosage regimen for an adult patient may be, for example, an oral dose of between 1 mg and 500 mg, preferably between 1 mg and 250 mg, or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 25 mg, of the compound of the formula (I) or a pharmaceutically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy.

[0022] The enzyme, lipoprotein associated Lp-PLA₂ has not hitherto been available in isolated purified form. The present invention therefore provides in a further aspect, the enzyme lipoprotein associated Lp-PLA₂ in purified form. By purified form is meant at least 80%, more preferably 90%, still more preferably 95% and most preferably 99% pure with respect to other protein contaminants.

[0023] The enzyme Lp-PLA₂ may be characterised by one or more partial peptide sequences selected from SEQ ID NOs:1, 2, 3, 4, 10 and 11 or by the partial peptide sequence comprising residues 271 to 441 or consisting of residues 1 to 441 of SEQ ID NO:9. The enzyme Lp-PLA₂ may further or alternatively characterised by its molecular weight found to be 45 kDa, at least 45 kDa, 45-47 kDa, 46-47 kDa or 45-50 kDa.

[0024] The invention also provides fragments of the enzyme having Lp-PLA₂ activity.

[0025] The enzyme can be isolated and purified using the methods hereafter described. Once isolated, the protein sequence of the enzyme can be obtained using standard techniques. In identifying said sequence, a number of protein fragments have been identified, each of which comprises part of the whole sequence of the enzyme. These sequences are themselves novel and form a further aspect of the invention.

[0026] This invention also provides isolated nucleic acid molecules encoding the enzyme, including mRNAs, DNAs, cDNAs as well as antisense analogs thereof and biologically active and diagnostically or therapeutically useful fragments thereof.

[0027] In particular, the invention provides an isolated nucleic acid molecule consisting of bases 1 to 1361 or 38 to 1361 or comprising the sequence corresponding to bases 848 to 1361 of SEQ ID NO: 9.

[0028] This invention also provides recombinant vectors, such as cloning and expression plasmids useful as reagents in the recombinant production of the enzyme, as well as recombinant prokaryotic and/or eukaryotic host cells comprising the novel nucleic acid sequence.

[0029] This invention also provides nucleic acid probes comprising nucleic acid molecules of sufficient length to specifically hybridize to the novel nucleic acid sequences.

[0030] This invention also provides an antisense oligonucleotide having a sequence capable of binding with mRNAs encoding the enzyme so as to prevent the translation of said mRNA.

[0031] This invention also provides transgenic non-human animals comprising a nucleic acid molecule encoding the enzyme. Also provided are methods for use of said transgenic animals as models for mutation and SAR (structure/activity relationship) evaluation as well as in drug screens.

[0032] This invention further provides a method of screening compounds to identify those compounds which inhibit the enzyme comprising contacting isolated enzyme with a test compound and measuring the rate of turnover of an enzyme substrate as compared with the rate of turnover in the absence of test compound.

[0033] "Recombinant" polypeptides refer to polypeptides produced by recombinant DNA techniques; i.e., produced from cells transformed by an exogenous DNA construct encoding the desired polypeptide. "Synthetic" polypeptides are those prepared by chemical synthesis.

[0034] A "replicon" is any genetic element (e.g., plasmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*; i.e., capable of replication under its own control.

[0035] A "vector" is a replicon, such as a plasmid, phage, or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

[0036] A "double-stranded DNA molecule" refers to the polymeric form of deoxyribonucleotides (bases adenine, guanine, thymine, or cytosine) in a double-stranded helix, both relaxed and supercoiled. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the sense strand of DNA.

[0037] A DNA "coding sequence of" or a "nucleotide sequence encoding" a particular protein, is a DNA sequence which is transcribed and translated into a polypeptide when placed under the control of appropriate regulatory sequences.

[0038] A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. Within the promoter sequence will be found a transcription initiation site (conveniently defined by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes.

[0039] DNA "control sequences" refers collectively to promoter sequences, ribosome binding sites, polyadenylation signals, transcription termination sequences, upstream regulatory domains, enhancers, and the like, which collectively provide for the expression (i.e., the transcription and translation) of a coding sequence in a host cell.

[0040] A control sequence “directs the expression” of a coding sequence in a cell when RNA polymerase will bind the promoter sequence and transcribe the coding sequence into mRNA, which is then translated into the polypeptide encoded by the coding sequence.

[0041] A “host cell” is a cell which has been transformed or transferred, or is capable of transformation or transfection by an exogenous DNA sequence.

[0042] A cell has been “transformed” by exogenous DNA when such exogenous DNA has been introduced inside the cell membrane. Exogenous DNA may or may not be integrated (covalently linked) into chromosomal DNA making up the genome of the cell. In prokaryotes and yeasts, for example, the exogenous DNA may be maintained on an episomal element, such as a plasmid. With respect to eukaryotic cells, a stably transformed or transfected cell is one in which the exogenous DNA has become integrated into the chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cell containing the exogenous DNA.

[0043] A “clone” is a population of cells derived from a single cell or common ancestor by mitosis. A “cell line” is a clone of a primary cell that is capable of stable growth in vitro for many generations.

[0044] Two DNA or polypeptide sequences are “substantially homologous” or “substantially the same” when at least about 85% (preferably at least about 90%, and most preferably at least about 95%) of the nucleotides or amino acids match over a defined length of the molecule and includes allelic variations. As used herein, substantially homologous also refers to sequences showing identity to the specified DNA or polypeptide sequence. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., “Current Protocols in Mol. Biol.” Vol. I & II, Wiley Interscience. Ausbel et al. (ed.) (1992). Protein sequences that are substantially the same can be identified by proteolytic digestion, gel electrophoresis and microsequencing.

[0045] The term “functionally equivalent” intends that the amino acid sequence of the subject protein is one that will exhibit enzymatic activity of the same kind as that of Lp-PLA₂.

[0046] A “heterologous” region of a DNA construct is an identifiable segment of DNA within or attached to another DNA molecule that is not found in association with the other molecule in nature.

[0047] This invention provides an isolated nucleic acid molecule encoding the enzyme Lp-PLA₂. One means for isolating the coding nucleic acid is to probe a human genomic or cDNA library with a natural or artificially designed probe using art recognized procedures (See for example: “Current Protocols in Molecular Biology”, Ausubel, F. M. et al. (eds.) Greene Publishing Assoc and John Wiley Interscience, New York, 1989, 1992); for example using the protein fragment information disclosed herein. The enzyme of this invention may be made by recombinant genetic engineering techniques. The isolated

nucleic acids particularly the DNAs can be introduced into expression vectors by operatively linking the DNA to the necessary expression control regions (e.g. regulatory regions) required for gene expression. The vectors can be introduced into the appropriate host cells such as prokaryotic (e.g., bacterial), or eukaryotic (e.g. yeast, insect or mammalian) cells by methods well known in the art (Ausubel et al., supra). The coding sequences for the desired proteins having been prepared or isolated, can be cloned into a suitable vector or replicon. Numerous cloning vectors are known to those of skill in the art, and the selection of an appropriate cloning vector is a matter of choice. Examples of recombinant DNA vectors for cloning and host cells which they can transform include the bacteriophage λ (*E. coli*), pBR322 (*E. coli*), pACYC177 (*E. coli*), pKT230 (gram-negative bacteria), pGV1106 (gram-negative bacteria), pLAFR1 (gram-negative bacteria), pME290 (non-*E. coli* gram-negative bacteria), pHV14 (*E. coli* and *Bacillus subtilis*), pBD9 (*Bacillus*), pIJ61 (*Streptomyces*), pUC6 (*Streptomyces*), YIp5 (*Saccharomyces*), a baculovirus insect cell system, YCp19 (*Saccharomyces*). See, generally, “DNA Cloning”: Vols. I & II, Glover et al. ed. IRL Press Oxford (1985) (1987) and; T. Maniatis al. (“Molecular Cloning” Cold Spring Harbor Laboratory (1982).

[0048] The gene can be placed under the control of a promoter, ribosome binding site (for bacterial expression) and, optionally, an operator (collectively referred to herein as “control” elements), so that the DNA sequence encoding the desired protein is transcribed into RNA in the host cell transformed by a vector containing this expression construction. The coding sequence may or may not contain a signal peptide or leader sequence. The protein sequences of the present invention can be expressed using, for example, the *E. coli* tac promoter or the protein A gene (spa) promoter and signal sequence. Leader sequences can be removed by the bacterial host in post-translational processing See. e.g., U.S. Pat. Nos. 4,431,739; 4,425,437; 4,338,397.

[0049] In addition to control sequences, it may be desirable to add regulatory sequences which allow for regulation of the expression of the protein sequences relative to the growth of the host cell. Regulatory sequences are known to those of skill in the art, and examples include those which cause the expression of a gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Other types of regulatory elements may also be present in the vector, for example, enhancer sequences.

[0050] An expression vector is constructed so that the particular coding sequence is located in the vector with the appropriate regulatory sequences, the positioning and orientation of the coding sequence with respect to the control sequences being such that the coding sequence is transcribed under the “control” of the control sequences (i.e., RNA polymerase which binds to the DNA molecule at the control sequences transcribes the coding sequence). Modification of the coding sequences may be desirable to achieve this end. For example, in some cases it may be necessary to modify the sequence so that it may be attached to the control sequences with the appropriate orientation; i.e., to maintain the reading frame. The control sequences and other regulatory sequences may be ligated to the coding sequence prior to insertion into a vector, such as the cloning vectors described above. Alternatively, the coding sequence can be

cloned directly into an expression vector which already contains the control sequences and an appropriate restriction site. Modification of the coding sequences may also be performed to alter codon usage to suit the chosen host cell, for enhanced expression.

[0051] In some cases, it may be desirable to add sequences which cause the secretion of the polypeptide from the host organism, with subsequent cleavage of the secretory signal. It may also be desirable to produce mutants or analogs of the enzyme of interest. Mutants or analogs may be prepared by the deletion of a portion of the sequence encoding the protein, by insertion of a sequence, and/or by substitution of one or more nucleotides within the sequence. Techniques for modifying nucleotide sequences, such as site-directed mutagenesis, are well known to those skilled in the art. See, e.g., T. Maniatis et al., *supra*; *DNA Cloning*, Vols. I and II, *supra*; *Nucleic Acid Hybridization*, *supra*.

[0052] A number of prokaryotic expression vectors are known in the art. See, e.g., U.S. Pat. Nos. 4,578,355; 4,440,859; 4,436,815; 4,431,740; 4,431,739; 4,428,941; 4,425,437; 4,418,149; 4,411,994; 4,366,246; 4,342,832; U.K. Patent Applications GB 2,121,054; GB 2,008,123; GB 2,007,675; and European Patent Application 103,395. Yeast expression vectors are also known in the art. See, e.g., U.S. Pat. Nos. 4,446,235; 4,443,539; 4,430,428; see also European Patent Applications 103,409; 100,561; 96,491. pSV2neo (as described in *J. Mol. Appl. Genet* 1:327-341) which uses the SV40 late promoter to drive expression in mammalian cells or pCDNA1neo, a vector derived from pCDNA1 (*Mol. Cell Biol.* 7:4125-29) which uses the CMV promoter to drive expression. Both these latter two vectors can be employed for transient or stable (using G418 resistance) expression in mammalian cells. Insect cell expression systems, e.g., *Drosophila* are also useful, see for example, PCT applications US 89105155 and US 91/06838 as well as EP application 88/304093.3.

[0053] Depending on the expression system and host selected, the enzyme of the present invention may be produced by growing host cells transformed by an expression vector described above under conditions whereby the protein of interest is expressed. The protein is then isolated from the host cells and purified. If the expression system secretes the protein into growth media, the protein can be purified directly from the media. If the protein is not secreted, it is isolated from cell lysates or recovered from the cell membrane fraction. Where the protein is localized to the cell surface, whole cells or isolated membranes can be used as an assayable source of the desired gene product. Protein expressed bacterial hosts such as *E. coli* may require isolation from inclusion bodies and refolding. The selection of the appropriate growth conditions and recovery methods are within the skill of the art.

[0054] The identification of this novel target for the treatment of atherosclerosis, also leads to a novel diagnostic method to diagnose a patient's susceptibility to developing atherosclerotic disease.

[0055] The present invention therefore provides in a still farther aspect a diagnostic method comprising isolating a sample of blood from the patient, and assaying said sample for Lp-PLA₂ activity. Patients that are susceptible to atherosclerotic disease are expected to have elevated levels of the Lp-PLA₂ enzyme, and hence the levels of Lp-PLA₂ provides an indication of the patient's susceptibility to atherosclerotic disease. Moreover, Lp-PLA₂ is found located predominantly on dense subfraction(s) of LDL which are

known to be very atherogenic. Plasma Lp-PLA₂ levels could therefore provide a ready measure of these very atherogenic small dense LDL particles.


[0056] It is expected that the presence of the enzyme in the blood sample of the patient can be assayed by analysis of enzyme activity (i.e. by an assay set up against the purified enzyme as standard); or alternatively by assaying protein content of the sample by using polyclonal or monoclonal antibodies prepared against the purified enzyme. Monoclonal (and polyclonal) antibodies raised against the purified enzyme or fragments thereof are themselves novel and form a further aspect of the invention.

DATA AND EXAMPLES

[0057] 1. Screen for Lp-PLA₂ inhibition.

[0058] Enzyme activity was determined by measuring the rate of turnover of the artificial substrate (A) at 37° C. in 50 mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulphonic acid) buffer containing 150 mM NaCl, pH 7.4.

(A)



Assays were performed in 96 well titre plates.

[0059] Lp-PLA₂ was pre-incubated at 37° C. with vehicle or test compound for 10 min in a total volume of 180 μ l. The reaction was then initiated by the addition of 20 μ l 10 $\times$ substrate (A) to give a final substrate concentration of 20 μ M. The reaction was followed at 405 nm for 20 minutes using a plate reader with automatic mixing. The rate of reaction was measured as the rate of change of absorbance.

[0060] Results:

Compound No	XR ¹	R	Z	IC ₅₀ (μ M)
1	O(CH ₂) ₁₅ CH ₃	CH ₃ CONH	N ⁺ (CH ₃) ₃	0.8
2	O(CH ₂) ₁₇ CH ₃	CH ₃ CONH	N ⁺ (CH ₃) ₃	3.5
3	O(CH ₂) ₁₇ CH ₃	CH ₃ CONH	S ⁺ (CH ₃) ₂	1.0
4	O(CH ₂) ₁₇ CH ₃	CH ₃ CONH	SCH ₃	0.08
5	O(CH ₂) ₁₇ CH ₃	CH ₃ CONH	OH	0.45
6	O(CH ₂) ₁₇ CH ₃	CH ₃ CONH	OAc	0.2
7	O(CH ₂) ₁₇ CH ₃	CH ₃ CONH		0.5
8	O(CH ₂) ₁₇ CH ₃	CH ₃ CONH		0.55
9	O(CH ₂) ₁₇ CH ₃	CF ₃ CONH	N ⁺ (CH ₃) ₃	2.5

[0061] 2. Copper Stimulated LDL Oxidation:

[0062] Copper stimulated oxidation of LDL is routinely measured by following the increase in conjugated diene

formation by monitoring the change in absorption at 234 nm. This assay can be used to study inhibitors of oxidative modification of LDL. FIG. 1 demonstrates that Lp-PLA₂ inhibitors are effective inhibitors of LDL oxidation through a prolongation of the lag phase, using compound 4 as an example.

[0063] 3. Inhibition of Cu²⁺ stimulated lyso-phosphatidylcholine (lyso-PtdCho) formation.

[0064] A 1 ml aliquot of human LDL (0.25 mg protein/ml) was incubated for 15 min at 37° C. with compound or vehicle. 5 μM Cu²⁺ was then added to allow oxidation/lyso-PtdCho formation to occur. The incubation was terminated by the addition of 3.75 ml chloroform/methanol/c HCl (200:400:5,v/v/v). Following the addition of 1.25 ml chloroform and 1.25 ml 0.1M HCl, the mixture was vortexed and centrifuged. The lower phase was carefully removed and the upper phase re-extracted with an equal volume of synthetic lower phase. The extracts were pooled and dried under nitrogen.

[0065] Phospholipids were reconstituted into 50 μl chloroform/methanol (2:1 v/v). 10 μl aliquots were spotted on to pre-run silica gel HPTLC plates and then developed in chloroform/methanol 25-30% methylamine (60:20:5 v/v/v). Plates were subsequently sprayed with the fluorescent indicator, 2-p-toluidinylnaphthalene-6-sulphonic acid (1 mM in 50 mM Tris/HCl, pH 7.4) to identify phospholipid components. Fluorescence was measured at 222 nm using a CAMAG TLC scanner. Lipoprotein lyso/PtdCho content was quantified using a standard curve (0.05-0.5 μg) prepared in parallel.

[0066] Compound 4 dose dependently inhibits LDL lyso-PtdCho accumulation stimulated by copper ions with an IC₅₀ value of ~30 μM.

[0067] 4. Purification of lipoprotein associated Lp-PLA₂

[0068] Low density lipoprotein (LDL) was obtained by plasma apheresis. The LDL was dialysed against 0.5 M NaCl 50 mM MES (4-morpholine ethane sulphonic acid), pH=6.0 overnight at 4° C. Solid CHAPS (3-[(3-cholamidopropyl)dimethylamino]-1-propane sulphonate) was added to 10 mM and the LDL stirred for 30 minutes to effect solubilisation. The solubilised LDL was pumped onto a pre-equilibrated Blue Sepharose 6FF (Pharmacia) column (26x20 cm). The column was then washed with 50 mM MES, 10 mM CHAPS, 0.5 M NaCl, pH=6.0 followed by 50 mM Tris, 10 mM CHAPS, 0.5 M NaCl, pH=8.0 until the absorbance (280 nm) of the eluate reached zero. Lp-PLA₂ was eluted using 50 mM Tris, 10 mM CHAPS, 15 M NaCl, pH=8.0. The Lp-PLA₂ fraction was then concentrated and dialysed overnight against 50 mM Tris, 10 mM CHAPS, pH=8.0.

[0069] The dialysed Lp-PLA₂ was submitted to anion exchange chromatography on a mono Q column (Pharmacia) using 50 mM Tris, 10 mM CHAPS, pH=8.0 with a NaCl gradient from zero to 0.3 M. The Lp-PLA₂ fractions obtained from the mono Q column were applied directly to a Hi Trap Blue cartridge (Pharmacia). The cartridge was washed with 50 mM Tris, 10 mM CHAPS, 0.5 M NaCl pH=8.0 until the absorbance of the eluate (280 nm) was zero. Lp-PLA₂ was then eluted using 50 mM Tris, 10 mM CHAPS, 1.5 M NaCl, pH=8.0. This gave Lp-PLA₂ which is

greater than 95% pure as shown in FIG. 2 This also demonstrates that the native enzyme is extensively glycosylated.

[0070] 5. Enzyme Sequence

[0071] The purity of the final enzyme preparation was verified by five criteria 1) SDS-polyacrylamide gel electrophoresis gave one band for both native and de-glycosylated forms. 2) Reverse phase high pressure liquid chromatography (RP-HPLC) gave a single peak, 3) The intact preparation gave no results by protein sequencing, implying that the protein was N-terminally blocked and free of any contaminants with open N-terminals, 4) By laser desorption mass spectrometry only one broad peak was observed with de-glycosylated protein, and 5) none of the sections of extended peptide data from sequencing gave any database matches indicative of contaminating proteins. Three cleavage strategies were used to obtain internal sequence information; trypsin (after de-glycosylation), cyanogen bromide (methionine cleavage) and BNPS-Skatol (tryptophan cleavage). The resulting peptides were separated by RP-HPLC, collected and sequenced. The accumulated sequence data allowed several extended stretches of primary structure of the Lp-PLA₂ enzyme to be verified. These are shown below as Peptides 1, 2, 3 and 4 (SEQ ID Nos 1 to 4). When searched against the National Centre for Biotechnological information (NCBI) non-redundant peptide sequence databases no high similarity matches were obtained. Estimation of the molecular weight of pure, de-glycosylated protein by laser desorption mass spectrometry gives values in the region of 45-47 kDa (separately 45 kDa and 46-47 kDa), indicating that the sequences constitute approximately 15 to 20% of the protein.

[0072] 6. Gene Sequence

[0073] Three expressed sequence tags (ESTs) from human cDNA libraries have been found to have extensive alignments with the Peptide Sequences 1 to 3. These EST's are shown below as Nucleotide Sequences 1 to 3 (SEQ ID Nos: 5 to 7) Nucleotide Sequence 1 is a 420 base sequence derived from a human foetal spleen library. Nucleotide Sequence 2 is a 379 base sequence derived from a 12-week human embryo library. Nucleotide Sequence 3 is a 279 base sequence derived from a T-cell lymphoma library. The identities at both the nucleic acid and amino acid level justified an overlapping alignment of the cDNA of all three ESTs, Nucleotide Sequences 3 (bases 1-278), 1 (bases 1-389) (in reverse orientation) and 2 (bases 1-304) with the Peptide Sequences 1, 2 and 3 (partially). Beyond these limits, the poor resolution of the raw sequence data precludes accurate base calling.

[0074] There are two remaining unassigned peptide sections from Peptides 3 and 4, both of which are expected to be present in the complete protein, -Q-Y-I-N-P-A-V-, and W-L-M-G-N-I-L-R-L-L-F-G-S-M-T-T-P-A-N-.

[0075] 7. Isolation of full-length Lp-PLA₂ cDNA

[0076] The full DNA sequence was determined for the clone (HLTA145) from which the Lymphoma EST (SEQ ID No:7) was derived, giving a total of 572 bases; SEQ ID No:8. There is one base difference between this sequence and the EST (between bases 1 to 256 of the EST); at position 27 of HLTA145 there is an A compared with a T in the EST. This would cause a coding change; L in HLTA145 compared

with F in the EST. Clone HLTA145 was used as a radiolabelled probe to screen the Lymphoma cDNA library in order to isolate the full-length Lp-PLA₂ clone. The library was prepared in the bacteriophage λ vector, Unizap XR (Stratagene).

Preparation of the Filters for Screening

[0077] The library was plated out at a density of 20,000 plaques per 150 mm petri dish onto *E. coli* XL-1 Blue host cells (ie. 200,000 plaques on 10 dishes). An overnight of XL-1 Blue was prepared in 100 mls LB/0.2% w/v Maltose/10 mM MgSO₄. The cells were pelleted, resuspended in 50 mls 10 mM MgSO₄ and stored on ice. 180 μ l of the library bacteriophage stock (23,400 pfu's) were added to 7 mls XL-1 Blue cells, mixed and divided into 10 aliquots of 615 μ l. The 10 tubes were incubated at 37° C. for 30 minutes. 7 mls of molten (@45° C.) top agarose (0.7% w/v agarose in LB) were added, mixed well and poured onto 150 mm LE agar plates (1.5% w/v agar in LB). The plates were inverted and incubated at 37° C. for approximately 7.5 hours. The plates were held at 4° C. until needed.

[0078] The plaques were transferred to 132 mm Hybond-N nylon filters (Amersham International) by laying the filters on the plates for 2 minutes (4 minutes for the duplicate). The DNA's on the filters were denatured for 2 minutes (0.5M NaCl, 1.5M NaOH), neutralised for 4 minutes (1.5M NaCl, 1.0M Tris pH7.4) and the filters placed on 2 $\times$ SSC for 1 minute. The filters were then dried and the DNA cross-linked to the filter using a Stratalinker UV 2400 (Stratagene) at 120,000 μ Joules/cm².

[0079] The filters were pre-hybridised in 1 mM EDTA, 0.5M NaHPO₄, 7% SDS (Church, G M. and Gilbert, W. (1984) PNAS USA 81 p 1991-1995) in a Techn HB2 hybridisation oven at 55° C. for 3 hours. Each bottle contained 2 filters and 25 mls prehybridization solution.

Preparation of the Radiolabelled Probe

[0080] The probe cDNA (from HLTA 145) was excised from pBluescript II SK+/- as an approximately 600 bp EcoRI-XhoI fragment and approximately 100 ng of gel purified fragment were labelled using 12MBq ³²P dATP and 1.2MBq ³²P dCTP by PCR labelling using Taq DNA polymerase (Boehringer Mannheim) and primers designed to prime at the 5' and 3' ends of the EST sequence. The labelling reaction was carried out in a total volume of 200 μ l and included unlabelled dNTP's at the following concentrations:

dATP	20 μ M
dCTP	20 μ M
dGTP	200 μ M
dTTP	200 μ M

The PCR reaction was carried out over 35 cycles of:

[0081] 94° C. for 30 s

[0082] 60° C. for 30 s

[0083] 72° C. for 30 s

Screening

[0084] The radiolabelled probe was denatured at 98° C. for 5 minutes and divided into 10 aliquots of 20 μ l. One aliquot was added per hybridisation bottle. Hybridisation was carried out over 16 hours at 55° C. The filters were washed at 60° C. (2 $\times$ 10 minutes) with 0.1% w/v SDS, 0.1 $\times$ SSC (50 mls per wash per bottle). The filters were autoradiographed and the films (Fuji Medical X-Ray Film) developed after 5 days exposure.

[0085] Duplicate positives were progressed to a secondary screen. The plaques were cored out into 1 ml SM (100 mM NaCl, 10 mM MgSO₄, 1 M Tris, pH7.4), titrated and plated onto 90 mm petri dishes at between 20 and 200 pfu's per dish. The secondary screen was carried out as described for the primary screen except the filters were washed at 65° C. The autoradiographs were developed after 16 hours exposure.

DNA Sequencing

[0086] The duplicated positive clones from the secondary screen were excised from the λ Unizap XR bacteriophage vector into the Bluescript phagemid (according to the Stratagene manual) for characterisation. One of the clones, carrying an insert of approximately 1.5 kb, was sequenced on both strands (using the USB Sequenase 2.0 DNA sequencing kit) by primer walking (SEQ ID No:9). The cDNA has an open reading frame with the potential to code for a polypeptide of 441 amino acids.

[0087] The 3' region of the full-length cDNA aligns with the HLTA145 sequence with the exception of 3 mismatches (see below). The predicted polypeptide sequence of the lymphoma Lp-PLA₂ is shown as SEQ ID No:9.

[0088] Inspection of the full length cDNA (SEQ ID No: 9) reveals probable errors in Peptide 3. One of these errors is in the assignment of continuity between V-M which is incompatible with the perfect sequence match with the cDNA after this position. It seems likely that a short peptide, containing the sequence -Q-Y-I-N-P-, co-purified with the longer cyanogen bromide partial cleavage peptide and, by being present in greater quantity, was assigned as the major sequence and contiguous with the subsequent amino acids. The remaining section of Peptide 3 and the whole of Peptide 4 can be identified in the predicted full length enzyme sequence (SEQ ID No:9). It thus seems likely that Peptide 3 is in fact two separate Peptides 5 (SEQ ID No:10) and 6 (SEQ ID No:11). The second probable error has occurred in the transcription from the raw data for Peptide 3 which on checking was consistent with Peptide 5 having the sequence -Q-Y-I-N-P-V-A, rather than Q-Y-I-N-P-A-V-.

[0089] The 3 base differences are as follows:

[0090] 1) T at 859 is A in HLTA145; aminoacid change F in full-length, L in HLTA145. (Note that the original EST is identical with the full-length cDNA at position 859).

[0091] 2) C at 1173 is T in HLTA145; aminoacid change A in full-length, V in HLTA145.

[0092] 3) T at 1203 is C in HLTA145; aminoacid change L in full-length, S in HLTA145.

[0093] The peptide data and the Foetal Spleen EST sequence (SEQ ID No:5) support the full-length cDNA

sequence for differences (2) and (3) although the Human Embryo EST (SEQ ID No:6) is identical to the Lymphoma EST (SEQ ID No:7) at position 1173. The Human Embryo EST (SEQ ID No:6) has a further difference (4) corresponding to position 1245 in the full-length Lymphoma sequence (SEQ ID No:9)(comparison between bases 2 to 304 of the Human Embryo EST and the full-length Lymphoma cDNA).

[0094] 4) A at 1245 is T in the Embryo EST (SEQ ID No:6)(amino acid change D to V in the Embryo EST). Peptide data covering this region supports the Lymphoma DNA sequence (SEQ ID No:9).

The Lp-PLA₂ DNA sequence from 848 to 1361 of SEQ ID No:9 (amino acid residues 271 to 441 of SEQ ID No:9) is the region for which all major data sets agree substantially, ie. the peptide data, the Foetal spleen, full-length Lymphoma and it includes the known active site and is therefore believed to be a significant characterising region for the Lp-PLA₂ enzyme

[0095] The predicted MW for the full reading frame is 50090. This is in excess of that determined for the deglycosylated, purified protein but post-translational events could account for this discrepancy. The most likely of these are the removal of an N-terminal signal peptide and/or limited proteolytic degradation of the protein C-terminal. The latter could occur in-vivo, during purification, or under the conditions of de-glycosylation.

Diagnostic Method

[0096] A sample of blood is taken from a patient, the plasma/serum sample prepared and passed through a dextran sulphate column pre-equilibrated with 0.9% (w/v) NaCl solution. Following washes with the same salt solution Lp-PLA₂ is eluted with a 4.1% (w/v) NaCl solution. Heparin agarose columns can also be used with the wash and elution solutions containing 0.7% and 2.9% NaCl, respectively. Enzyme present in the sample is determined by assaying for either

[0097] (a) enzyme activity:

[0098] The substrate (A) (see structure in 1) is used to assay Lp-PLA₂ activity by monitoring the absorbance change at 400 nm. Purified enzyme is pre-incubated at 37° C. and substrate (50 µM) is added after 5 minutes. The absorbance change at 400 nm is monitored for 20 minutes. This substrate has previously been reported as a substrate for classical calcium-dependent PLA₂s. (Washburn, W. N. and Dennis, E. A., J. Amer Chem. Soc., 1990, 112, 2040-2041); or

[0099] (b) protein content

[0100] Total protein content (i.e. enzyme content) can be determined using polyclonal antiserum raised against purified human Lp-PLA₂. The antiserum recognises both native and glycosylated enzyme as measured by immunoprecipitation of activity and Western Blot analysis.

[0101] Polyclonal antiserum was prepared as follows. Immunisation of rabbits involved mixing 0.5 ml of purified human Lp-PLA₂ (=100 µg) with an equal volume of Freund's complete adjuvant. The final emulsion was given subcutaneously in 4x0.25 ml injections. Boosting using a Freund's incomplete adjuvant antigen mixture (4x0.25 ml subcut; dosage=50 µg) took place 4 weeks later. Adequate titre was evident at between 6-8 weeks from initial injection.

IN THE FIGURES

[0102] FIG. 1 is a graph of absorbance at 234 nm against time (min) in a study of inhibition of copper (5 µM)-stimulated LDL (150 µg/ml) oxidation by compound 4 vs control vehicle.

[0103] FIG. 2 is an analysis the purified Lp-PLA₂ material of Example 4 following separation by polyacrylamide gel electrophoresis. Lanes 2, 4 and 6 contain adjacent fractions of purified native Lp-PLA₂. Lanes 1, 3 and 5 are fractions 2, 4 and 6 respectively after N-deglycosylation.

SEQUENCE DATA

[0104]

```
SEQ. ID. No: 1 - Peptide 1
-M-L-K-L-K-G-D-I-D-S-N-A-A-I-D-L-S-N-K-A-S-L-A-F-L-Q-K-H-L-G-L-H-K-D-F-D-Q-

SEQ. ID. No: 2 - Peptide 2
-W-M-F-P-L-G-D-E-V-Y-S-R-I-P-Q-P-L-F-F-I-N-S-E-Y-F-Q-Y-P-A-N-

SEQ. ID. No: 3 - Peptide 3
-Q-Y-I-N-P-A-V-M-I-T-I-R-G-S-V-H-Q-N-F-A-D-F-T-F-A-T-G-

SEQ. ID. No: 4 - Peptide 4
-W-L-M-G-N-I-L-R-L-L-F-G-S-M-T-T-P-A-N

SEQ. ID. No: 5 - Nucleotide Sequence 1
1 AAAAAACCTA TTTTAATCCT AATTGTATT CTCTATTCCT GAAGAGTTCT
51 GTAACATGAT GTGTTGATTG GTTGTGTAA TGTGGTCCC TGAATAAGA
101 TTCTCATCAT CTCCTTCAAT CAAGCAGTCC CACTGATCAA AATCTTTATG
151 AAGTCCTAAA TGCTTTTGTA AGAATGCTAA TGAAGCTTGT TGCTAAGAT
201 CAATAGCTGC ATTTGAATCT ATGTCTCCCT TTAATTTGAG CATGTGTCCA
251 ATTATTTTGC CAGTNGCAAA AGTGAAGTCA GCAAATCTCT GGTGGACTGA
301 ACCCCTGATT GTAATCATCT TTCTTTCTTT ATCAGGTGAG TAGCATTTTT
```

-continued

351 TCATTTTAT GATATTAGCA GGATATTGGA AATATTCAGN GTTGNTAAAA

401 AGNGGNGGCT GAGGGATTCT

SEQ. ID. No: 6 - Nucleotide Sequence 2

1 TGCTAATATC ATAAAAATGA AAAAATGCTA CTCACCTGAT AAAGAAAGAA

51 AGATGATTAC AATCAGGGGT TCAGTCCACC AGANTTTTGC TGACTTCACT

101 TTGCAACTG GCAAAATAAT TGGACACATG CTCAAATTAA AGGGAGACAT

151 AGATTCAAAT GTAGCTATTG ATCTTAGCAA CAAAGCTTCA TTAGCATTCT

201 TACAAAAGCA TTTAGGACTT CATAAAGATT TTGTTCACTG GGACTGCTTG

251 ATTGAAGGAG ATGATGAGAA TCTTATCCA GGGACCAACA TTAACACAAC

301 CAATTCAACA CATCATGTTT ACAGAACTTC TTCCAGGGAA TAGGAGGAAA

351 TACAATTGGG GTTTAAAATA GGTTTTTTTT

SEQ. ID. No: 7 - Nucleotide Sequence 3

1 GAAGAATGCA TTAGATTTAA AGTTTGATAT GGAACAACCTG AAGGACTCTA

51 TTGATAGGGA AAAAATAGCA GTAATTGGAC ATTCTTTTGG TGGAGCAACG

101 GTTATTCAGA CTCTTAGTGA AGATCAGAGA TTCAGATGTG GTATTGCCCT

151 GGATGCATGG ATGTTTCCAC TGGGTGATGA AGTATATTCC AGAATTCCTC

201 AGCCCTCTT TTTTATCAAC TCTGAATATT TCCAATATCC TGCTAATATC

251 ATAAAANTGG AAAAATGCTA CTCACCTGG

Seq. ID. No: 8 - DNA sequence of HLTA145

10 20 30 40 50
AAAATAGCAG TAATTGGACA TTCTTTAGGT GGAGCAACGG TTATTGAGC

60 70 80 90 100
TCTTAGTGAA GATCAGAGAT TCAGATGTGG TATTGCCCTG GATGCATGGA

110 120 130 140 150
TGTTTCCACT GGGTGATGAA GTATATTCCA GAATTCCTCA GCCCTCTTT

160 170 180 190 200
TTTATCAACT CTGAATATTT CCAATATCCT GCTAATATCA TAAAAATGAA

210 220 230 240 250
AAAATGCTAC TCACCTGATA AAGAAAGAAA GATGATTACA ATCAGGGGTT

260 270 280 290 300
CAGTCCACCA GAATTTTGCT GACTTCACCT TTGCAACTGG CAAAAAATT

310 320 330 340 350
GGACACATGC TCAAATTAAA GGGAGACATA GATTCAAATG TAGCTATTGA

360 370 380 390 400
TCTTAGCAAC AAAGCTTCAT CAGCATTCTT ACAAAGCAT TTAGGACTTC

410 420 430 440 450
ATAAAGATTT TGATCAGTGG GACTGCTTGA TTGAAGGAGA TGATGAGAAT

460 470 480 490 500
CTTATTCCAG GGACCAACAT TAACACAACC AATCAACACA TCATGTTACA

510 520 530 540 550
GAACTCTTCA GGAATAGAGA AATACAATTA GGATTAATAAT AGGTTTTTTA

560 570
AAAAAAAAAA AAAAAAACT CG

SEQ. ID. No:9 - cDNA Sequence of Lymphoma Lp-PLA2

10 20 30 40 50
TGAGAGACTAAGCTGAACTGCTGCTCAGCTCCCAAGATGGTGCCACCCA
M V P P K

60 70 80 90 100

-continued

AATTGCATGTGCTTTTCTGCCTCTGCGGCTGCCTGGCTGTGGTTTATCCT
L H V L F C L C G C L A V V Y P

110 120 130 140 150
TTTGACTGGCAATACATAAATCCTGTTGCCATATGAAATCATCAGCATG
F D W Q Y I N P V A H M K S S A W

160 170 180 190 200
GGTCAACAAAATACAAGTACTGATGGCTGCTGCAAGCTTTGGCCAACTA
V N K I Q V L M A A A S F G Q T K

210 220 230 240 250
AAATCCCCGGGGAAATGGGCCTTATTCGTTGGTTGTACAGACTTAATG
I P R G N G P Y S V G C T D L M

260 270 280 290 300
TTTGATCACACTAATAAGGCACCTTCTTGCGTTTATATTATCCATCCCA
F D H T N K G T F L R L Y Y P S Q

310 320 330 340 350
AGATAATGATCGCCTTGACACCTTTGGATCCCAAATAAAGAATATTTT
D N D P L D T L W I P N K E Y F W

360 370 380 390 400
GGGTCTTTAGCAAATTTCTTGGAACACACTGGCTTATGGGCAACATTTTG
G L S K F L G T H W L M G N I L

410 420 430 440 450
AGGTACTCTTTGGTTCAATGACAACTCCTGCAAACTGGAATTCCTCT
R L L F G S M T T P A N W N S P L

460 470 480 490 500
GAGGCCTGGTGAAAAATATCCACTTGTGTTTTTCTCATGGTCTTGGGG
R P G E K Y P L V V F S H G L G A

510 520 530 540 550
CATTCAGGACACTTTATCTGCTATTGGCATTGACCTGGCATCTCATGGG
F R T L Y S A I G I D L A S H G

560 570 580 590 600
TTTATAGTTGCTGCTGTAGAACACAGAGATAGATCTGCATCTGCAACTTA
F I V A A V E H R D R S A S A T Y

610 620 630 640 650
CTATTTCAAGGACCAATCTGCTGCAGAAATAGGGGACAAGTCTTGGCTCT
Y F K D Q S A A E I G D K S W L Y

660 670 680 690 700
ACCTTAGAACCTGAAACAAGAGGAGAGACATATACGAAATGAGCAG
L R T L K Q E E E T H I R N E Q

710 720 730 740 750
GTACGGCAAAGAGCAAAGAAATGTTCCCAAGCTCTCAGTCTGATTCTTGA
V R Q R A K E C S Q A L S L I L D

760 770 780 790 800
CATTGATCATGGAAGCCAGTGAAGAATGCATTAGATTTAAAGTTTGATA
I D H G K P V K N A L D L K F D M

810 820 830 840 850
TGGAACAACCTGAAGGACTCTATTGATAGGGAAAAATAGCAGTAATTGGA
E Q L K D S I D R E K I A V I G

860 870 880 890 900
CATTCCTTTTGGTGGAGCAACGGTTATTCAGACTCTTAGTGAAGATCAGAG
H S F G G A T V I Q T L S E D Q R

910 920 930 940 950
ATTCAGATGTGGTATTGCCCTGGATGCATGGATGTTCCACTGGGTGATG
F R C G I A L D A W M F P L G D E

960 970 980 990 1000
AAGTATATTCCAGAATTCCTCAGCCCTCTTTTATCAACTCTGAATAT
V Y S R I P Q P L F F I N S E Y

1010 1020 1030 1040 1050
TTCCAATATCTGCTAATATCATAAAAATGAAAAATGCTACTCACCTGA

-continued

F Q Y P A N I I K M K K C Y S P D

1060 1070 1080 1090 1100
TAAAGAAAGAAAGATGATTACAATCAGGGGTTTCAGTCCACCAGAATTTTG
K E R K M I T I R G S V H Q N F A

1110 1120 1130 1140 1150
CTGACTTCACTTTTGCAACTGGCAAAATAATTGGACACATGCTCAAATTA
D F T F A T G K I I G H M L K L

1160 1170 1180 1190 1200
AAGGGAGACATAGATTCAAATGCAGCTATTGATCTTAGCAACAAAGCTTC
K G D I D S N A A I D L S N K A S

1210 1220 1230 1240 1250
ATTAGCATTCTTACAAAAGCATTTAGGACTTCATAAAGATTTTGATCAGT
L A F L Q K H L G L H K D P D Q W

1260 1270 1280 1290 1300
GGGACTGCTTGATTGAAGGAGATGATGAGAATCTTATTCAGGGACCAAC
D C L I H G D D E N L I P G T N

1310 1320 1330 1340 1350
ATTAACAACAACCAATCAACACATCATGTTACAGAACTCTTCAGGAATAGA
I N T T N Q H I M L Q N S S G I E

1360
GAAATACAATT
K Y N

SEQ. ID. No: 10 - Peptide 5
-Q-Y-I-N-P-V-A-

SEQ. ID. No: 11 - Peptide 6
-M-I-T-I-R-G-S-V-H-Q-N-F-A-D-F-T-F-A-T-G

[0105]

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 11

(2) INFORMATION FOR SEQ ID NO: 1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 37 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Met Leu Lys Leu Lys Gly Asp Ile Asp Ser Asn Ala Ala Ile Asp Le
1 5 10 15

Ser Asn Lys Ala Ser Leu Ala Phe Leu Gln Lys His Leu Gly Leu Hi
20 25 30

-continued

Lys Asp Phe Asp Gln
35

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Trp Met Phe Pro Leu Gly Asp Glu Val Tyr Ser Arg Ile Pro Gln Pr
1 5 10 15

Leu Phe Phe Ile Asn Ser Glu Tyr Phe Gln Tyr Pro Ala Asn
20 25 30

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Gln Tyr Ile Asn Pro Ala Val Met Ile Thr Ile Arg Gly Ser Val Hi
1 5 10 15

Gln Asn Phe Ala Asp Phe Thr Phe Ala Thr Gly
20 25

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Trp Leu Met Gly Asn Ile Leu Arg Leu Leu Phe Gly Ser Met Thr Th
1 5 10 15

Pro Ala Asn

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 420 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

```
AAAAAACCTA TTTAATCCT AATTGTATTT CTCTATTCCT GAAGAGTCT GTAACATGAT    60
GTGTTGATTG GTTGTTTAA TGTGGTCCC TGAATAAGA TTCTCATCAT CTCCTTCAAT    120
CAAGCAGTCC CACTGATCAA AATCTTTATG AAGTCCTAAA TGCTTTTGTGA AGAATGCTAA    180
TGAAGCTTTG TTGCTAAGAT CAATAGCTGC ATTTGAATCT ATGTCTCCCT TTAATTTGAG    240
CATGTGTCCA ATTATTTTGC CAGTNGCAA AGTGAAGTCA GCAAATCTCT GGTGGACTGA    300
ACCCCTGATT GTAATCATCT TTCTTTCTTT ATCAGGTGAG TAGCATTTTT TCATTTTAT    360
GATATTAGCA GGATATTGGA AATATTCAGN GTTGNTAAAA AGNGGNGGCT GAGGGATTCT    420
```

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 379 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

```
TGCTAATATC ATAAAAATGA AAAAATGCTA CTCACCTGAT AAAGAAAGAA AGATGATTAC    60
AATCAGGGGT TCAGTCCACC AGANTTTTGC TGAATCACT TTTGCAACTG GCAAATAAT    120
TGGACACATG CTCAAATTAA AGGAGACAT AGATTCAAAT GTAGCTATTG ATCTTAGCAA    180
CAAAGCTTCA TTAGCATTCT TACAAAAGCA TTAGGACTT CATAAGATT TTGTTCACTG    240
GGACTGCTTG ATTGAAGGAG ATGATGAGAA TCTATTCCA GGGACCAACA TTAACACAAC    300
CAATTCAACA CATCATGTTT ACAGAACTTC TTCCAGGGAA TAGGAGGAAA TACAATTGGG    360
GTTTAAATAA GGTTTTTTT    379
```

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 279 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

```
GAAGAATGCA TTAGATTTAA AGTTTGATAT GGAACAACAG AAGGACTCTA TTGATAGGGA    60
AAAAATAGCA GTAATTGGAC ATTCTTTTGG TGGAGCAACG GTTATTGAGA CTCTTAGTGA    120
AGATCAGAGA TTCAGATGTG GTATTGCCCT GGATGCATGG ATGTTTCCAC TGGGTGATGA    180
```

-continued

```

AGTATATTCC AGAATTCCTC AGCCCCCTCT TTTTATCAAC TCTGAATATT TCCAATATCC 240
TGCTAATATC ATAAAAANTGG AAAAATGCTA CTCACCTGG 279

```

(2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 572 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

```

AAAATAGCAG TAATTGGACA TTCTTTAGGT GGAGCAACGG TTATTCAGAC TCTTAGTGAA 60
GATCAGAGAT TCAGATGTGG TATTGCCCTG GATGCATGGA TGTTCCTACT GGGTGATGAA 120
GTATATTCCA GAATTCCTCA GCCCCTCTTT TTTATCAACT CTGAATATTT CCAATATCCT 180
GCTAATATCA TAAAAATGAA AAAATGCTAC TCACCTGATA AAGAAAGAAA GATGATTACA 240
ATCAGGGGTT CAGTCCACCA GAATTTTGCT GACTTCACCT TTGCAACTGG CAAAATAATT 300
GGACACATGC TCAAAATAAA GGGAGACATA GATTCAAATG TAGCTATTGA TCTTAGCAAC 360
AAAGCTTCAT CAGCATTCTT ACAAAGCAT TTAGGACTTC ATAAAGATTT TGATCAGTGG 420
GACTGCTTGA TTGAAGGAGA TGATGAGAAT CTTATTCCAG GGACCAACAT TAACACAACC 480
AATCAACACA TCATGTTACA GAACTCTTCA GGAATAGAGA AATACAATTA GGATTAAAT 540
AGGTTTTTTT AAAAAAAAAA AAAAAAACT CG 572

```

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1361 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 38..1360

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

```

TGAGAGACTA AGCTGAACT GCTGCTCAGC TCCCAAG ATG GTG CCA CCC AAA TTG 55
                               Met Val Pro Pro Lys Leu
                               1           5
CAT GTG CTT TTC TGC CTC TGC GGC TGC CTG GCT GTG GTT TAT CCT TTT 103
His Val Leu Phe Cys Leu Cys Gly Cys Leu Ala Val Val Tyr Pro Phe
    10           15           20
GAC TGG CAA TAC ATA AAT CCT GTT GCC CAT ATG AAA TCA TCA GCA TGG 151
Asp Trp Gln Tyr Ile Asn Pro Val Ala His Met Lys Ser Ser Ala Trp
    25           30           35
GTC AAC AAA ATA CAA GTA CTG ATG GCT GCT GCA AGC TTT GGC CAA ACT 199

```


-continued

Val	Asn	Lys	Ile	Gln	Val	Leu	Met	Ala	Ala	Ala	Ser	Phe	Gly	Gln	Thr	
40						45					50					
AAA	ATC	CCC	CGG	GGA	AAT	GGG	CCT	TAT	TCC	GTT	GGT	TGT	ACA	GAC	TTA	247
Lys	Ile	Pro	Arg	Gly	Asn	Gly	Pro	Tyr	Ser	Val	Gly	Cys	Thr	Asp	Leu	
55					60					65					70	
ATG	TTT	GAT	CAC	ACT	AAT	AAG	GGC	ACC	TTC	TTG	CGT	TTA	TAT	TAT	CCA	295
Met	Phe	Asp	His	Thr	Asn	Lys	Gly	Thr	Phe	Leu	Arg	Leu	Tyr	Tyr	Pro	
				75					80					85		
TCC	CAA	GAT	AAT	GAT	CGC	CTT	GAC	ACC	CTT	TGG	ATC	CCA	AAT	AAA	GAA	343
Ser	Gln	Asp	Asn	Asp	Arg	Leu	Asp	Thr	Leu	Trp	Ile	Pro	Asn	Lys	Glu	
				90				95					100			
TAT	TTT	TGG	GGT	CTT	AGC	AAA	TTT	CTT	GGA	ACA	CAC	TGG	CTT	ATG	GGC	391
Tyr	Phe	Trp	Gly	Leu	Ser	Lys	Phe	Leu	Gly	Thr	His	Trp	Leu	Met	Gly	
		105					110					115				
AAC	ATT	TTG	AGG	TTA	CTC	TTT	GGT	TCA	ATG	ACA	ACT	CCT	GCA	AAC	TGG	439
Asn	Ile	Leu	Arg	Leu	Leu	Phe	Gly	Ser	Met	Thr	Thr	Pro	Ala	Asn	Trp	
		120					125					130				
AAT	TCC	CCT	CTG	AGG	CCT	GGT	GAA	AAA	TAT	CCA	CTT	GTT	GTT	TTT	TCT	487
Asn	Ser	Pro	Leu	Arg	Pro	Gly	Glu	Lys	Tyr	Pro	Leu	Val	Val	Phe	Ser	
135					140					145					150	
CAT	GGT	CTT	GGG	GCA	TTC	AGG	ACA	CTT	TAT	TCT	GCT	ATT	GGC	ATT	GAC	535
His	Gly	Leu	Gly	Ala	Phe	Arg	Thr	Leu	Tyr	Ser	Ala	Ile	Gly	Ile	Asp	
				155					160					165		
CTG	GCA	TCT	CAT	GGG	TTT	ATA	GTT	GCT	GCT	GTA	GAA	CAC	AGA	GAT	AGA	583
Leu	Ala	Ser	His	Gly	Phe	Ile	Val	Ala	Ala	Val	Glu	His	Arg	Asp	Arg	
			170					175					180			
TCT	GCA	TCT	GCA	ACT	TAC	TAT	TTC	AAG	GAC	CAA	TCT	GCT	GCA	GAA	ATA	631
Ser	Ala	Ser	Ala	Thr	Tyr	Tyr	Phe	Lys	Asp	Gln	Ser	Ala	Ala	Glu	Ile	
			185				190					195				
GGG	GAC	AAG	TCT	TGG	CTC	TAC	CTT	AGA	ACC	CTG	AAA	CAA	GAG	GAG	GAG	679
Gly	Asp	Lys	Ser	Trp	Leu	Tyr	Leu	Arg	Thr	Leu	Lys	Gln	Glu	Glu	Glu	
		200					205				210					
ACA	CAT	ATA	CGA	AAT	GAG	CAG	GTA	CGG	CAA	AGA	GCA	AAA	GAA	TGT	TCC	727
Thr	His	Ile	Arg	Asn	Glu	Gln	Val	Arg	Gln	Arg	Ala	Lys	Glu	Cys	Ser	
215					220					225					230	
CAA	GCT	CTC	AGT	CTG	ATT	CTT	GAC	ATT	GAT	CAT	GGA	AAG	CCA	GTG	AAG	775
Gln	Ala	Leu	Ser	Leu	Ile	Leu	Asp	Ile	Asp	His	Gly	Lys	Pro	Val	Lys	
				235					240					245		
AAT	GCA	TTA	GAT	TTA	AAG	TTT	GAT	ATG	GAA	CAA	CTG	AAG	GAC	TCT	ATT	823
Asn	Ala	Leu	Asp	Leu	Lys	Phe	Asp	Met	Glu	Gln	Leu	Lys	Asp	Ser	Ile	
				250				255					260			
GAT	AGG	GAA	AAA	ATA	GCA	GTA	ATT	GGA	CAT	TCT	TTT	GGT	GGA	GCA	ACG	871
Asp	Arg	Glu	Lys	Ile	Ala	Val	Ile	Gly	His	Ser	Phe	Gly	Gly	Ala	Thr	
		265					270					275				
GTT	ATT	CAG	ACT	CTT	AGT	GAA	GAT	CAG	AGA	TTC	AGA	TGT	GGT	ATT	GCC	919
Val	Ile	Gln	Thr	Leu	Ser	Glu	Asp	Gln	Arg	Phe	Arg	Cys	Gly	Ile	Ala	
		280					285				290					
CTG	GAT	GCA	TGG	ATG	TTT	CCA	CTG	GGT	GAT	GAA	GTA	TAT	TCC	AGA	ATT	967
Leu	Asp	Ala	Trp	Met	Phe	Pro	Leu	Gly	Asp	Glu	Val	Tyr	Ser	Arg	Ile	
295					300					305					310	
CCT	CAG	CCC	CTC	TTT	TTT	ATC	AAC	TCT	GAA	TAT	TTC	CAA	TAT	CCT	GCT	1015
Pro	Gln	Pro	Leu	Phe	Phe	Ile	Asn	Ser	Glu	Tyr	Phe	Gln	Tyr	Pro	Ala	
				315					320					325		
AAT	ATC	ATA	AAA	ATG	AAA	AAA	TGC	TAC	TCA	CCT	GAT	AAA	GAA	AGA	AAG	1063
Asn	Ile	Ile	Lys	Met	Lys	Lys	Cys	Tyr	Ser	Pro	Asp	Lys	Glu	Arg	Lys	
				330					335				340			
ATG	ATT	ACA	ATC	AGG	GGT	TCA	GTC	CAC	CAG	AAT	TTT	GCT	GAC	TTC	ACT	1111

-continued

Met Ile Thr Ile Arg Gly Ser Val His Gln Asn Phe Ala Asp Phe Thr	
345 350 355	
TTT GCA ACT GGC AAA ATA ATT GGA CAC ATG CTC AAA TTA AAG GGA GAC	1159
Phe Ala Thr Gly Lys Ile Ile Gly His Met Leu Lys Leu Lys Gly Asp	
360 365 370	
ATA GAT TCA AAT GCA GCT ATT GAT CTT AGC AAC AAA GCT TCA TTA GCA	1207
Ile Asp Ser Asn Ala Ala Ile Asp Leu Ser Asn Lys Ala Ser Leu Ala	
375 380 385 390	
TTC TTA CAA AAG CAT TTA GGA CTT CAT AAA GAT TTT GAT CAG TGG GAC	1255
Phe Leu Gln Lys His Leu Gly Leu His Lys Asp Phe Asp Gln Trp Asp	
395 400 405	
TGC TTG ATT GAA GGA GAT GAT GAG AAT CTT ATT CCA GGG ACC AAC ATT	1303
Cys Leu Ile Glu Gly Asp Asp Glu Asn Leu Ile Pro Gly Thr Asn Ile	
410 415 420	
AAC ACA ACC AAT CAA CAC ATC ATG TTA CAG AAC TCT TCA GGA ATA GAG	1351
Asn Thr Thr Asn Gln His Ile Met Leu Gln Asn Ser Ser Gly Ile Glu	
425 430 435	
AAA TAC AAT T	1361
Lys Tyr Asn	
440	

(2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Gln Tyr Ile Asn Pro Val Ala
 1 5

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Met Ile Thr Ile Arg Gly Ser Val His Gln Asn Phe Ala Asp Phe Th
 1 5 10 15
 Phe Ala Thr Gly
 20

1-18. (canceled)

19. A method of diagnosing a patient's susceptibility to atherosclerotic disease which comprises analysing a sample from the patient for the presence of the enzyme lipoprotein

phospholipase A2 (Lp-PLA₂) using an antibody raised against the enzyme, wherein elevated levels of Lp-PLA₂ in said sample are indicative of the patient being susceptible to atherosclerotic disease.

20-24. (canceled)

25. The method according to claim 19 wherein the atherosclerotic disease is selected from the group comprising atherosclerosis, stroke and myocardial infarction.

26. The method according to claim 19 wherein the atherosclerotic disease is atherosclerosis.

27. The method according to claim 19 wherein the atherosclerotic disease is stroke.

28. The method according to claim 19 wherein the atherosclerotic disease is myocardial infarction.

29. The method according to claim 19 wherein the sample is selected from the group comprising blood, serum and plasma.

30. The method according to claim 19 wherein the sample is blood.

31. The method according to claim 19 wherein the sample is serum.

32. The method according to claim 19 wherein the sample is plasma.

33. The method according to claim 19 wherein the sample is assayed in a 96 well titre plate.

34. The method according to claim 19 wherein the antibody recognizes native or glycosylated Lp-PLA₂.

35. The method according to claim 19 wherein the antibody is a polyclonal antibody.


36. The method according to claim 19 wherein the antibody is a monoclonal antibody.

* * * * *

专利名称(译)	脂蛋白相关磷脂酶A2，其抑制剂及其在诊断和治疗中的用途		
公开(公告)号	US20070077251A1	公开(公告)日	2007-04-05
申请号	US11/603741	申请日	2006-11-22
[标]申请(专利权)人(译)	MACPHEE COLIN ^ h TEW DAVID 摹		
申请(专利权)人(译)	MACPHEE COLIN ^ h TEW DAVID 摹		
当前申请(专利权)人(译)	MACPHEE COLIN ^ h TEW DAVID 摹		
[标]发明人	MACPHEE COLIN H TEW DAVID GRAHAM		
发明人	MACPHEE, COLIN H. TEW, DAVID GRAHAM		
IPC分类号	A61K39/395 C12Q1/34 C07H21/04 C12P21/06 C12N9/18 G01N33/53 A61K31/739 A61K38/00 A61K38/55 A61P9/10 C07K14/47 C07K16/40 C12N1/21 C12N9/16 C12N9/20 C12N15/00 C12N15/09 C12N15/55 C12N15/74 C12P21/08 C12Q1/44 G01N33/573		
CPC分类号	A61K38/005 C12N9/18 C12N9/20 C12Q1/44 G01N33/6893 G01N2800/323 G01N2800/324 Y10S435/81 A61P9/10		
优先权	1994000413 1994-01-11 GB 1993013144 1993-06-25 GB 08/387858 1995-02-24 US PCT/GB1994/001374 1994-06-24 WO		
其他公开文献	US7416853		
外部链接	Espacenet USPTO		

摘要(译)

纯化形式的酶Lp-PLA2，编码Lp-PLA2的分离的核酸分子，酶Lp-PLA2的抑制剂在治疗中的用途以及筛选化合物以鉴定那些化合物的方法。



(I)