



US 20110117116A1

(19) **United States**(12) **Patent Application Publication**
THOMAS et al.(10) **Pub. No.: US 2011/0117116 A1**(43) **Pub. Date: May 19, 2011**(54) **NUCLEIC ACIDS ENCODING A HOUSE DUST
MITE ALLERGEN, DER P III, AND USES
THEREFOR****Publication Classification**(51) **Int. Cl.***A61K 39/35* (2006.01)*C12P 21/02* (2006.01)*G01N 33/566* (2006.01)*A61P 37/08* (2006.01)*A61P 37/04* (2006.01)(75) Inventors: **Wayne R. THOMAS**, Nedlands
(AU); **Kaw-Yan Chua**, Taipei
(TW); **Bruce L. Rogers**, Belmont,
MA (US); **Mei-chang Kuo**,
Winchester, MA (US)(73) Assignee: **MERCK PATENT GmbH**,
Darmstadt (DE)(21) Appl. No.: **12/688,744**(22) Filed: **Jan. 15, 2010**(52) **U.S. Cl. 424/185.1; 435/69.3; 436/501**(57) **ABSTRACT****Related U.S. Application Data**(60) Division of application No. 10/846,341, filed on May
14, 2004, now Pat. No. 7,666,429, which is a continu-
ation of application No. 08/462,515, filed on Jun. 5,
1995, now Pat. No. 6,752,991, which is a division of
application No. 08/163,919, filed on Dec. 8, 1993, now
Pat. No. 6,180,771.

Isolated nucleic acids encoding an allergen of *Dermatophagoides pteronyssinus*, Der p III, are disclosed. A cDNA encoding a peptide having a Der p III activity and a predicted molecular weight of about 24,985 daltons is also described. The nucleic acids can be used as probes to detect the presence of Der p III nucleic acid in a sample or for the recombinant production of peptides having an activity of Der p III. Peptides having an activity of Der p III can be used in compositions suitable for pharmaceutical administration or methods of diagnosing sensitivity to house dust mites.

CT 2
TTTTTTTTTTTTTGA AACATTGAAAAACATTGAAGCAACTGTGTGTACAACCAGAAAG 62
ATGATCATCTATAATATTTTAATTGTTTTATTATTGGCCATTAATACATTGGCTAATCCA 122
M I I Y N I L I V L L L A I N T L A N P
-29 -20 -10

ATTCTACCAGCATCACCAAATGCAACTATTGTTGGTGGTGAAAAAGCATTAGCTGGTGAA 182
I L P A S P N A T I V G G E K A L A G E
1 10

TGTCCATATCAGATTTTCATTACAATCAAGTAGTCATTTTGTGGTGGTACTATTCTTGAT 242
C P Y Q I S L Q S S S H F C G G T I L D
20 30

GAATATTGGATTTTAAACAGCTGCACATTGTGTTGCCGGACAAACAGCAAGTAACTTTCA 302
E Y W I L T A A H C V A G Q T A S K L S
40 50

ATTCGTTACAATAGTTTAAACATTTCATTAGGTGGTGAAAAAATTTCTGTTGCTAAAATT 362
I R Y N S L K H S L G G E K I S V A K I
60 70

TTTGACATGAAAAATATGATAGTTATCAAATTGATAATGATATTGCATTGATTAAGCTT 422
F A H E K Y D S Y Q I D N D I A L I K L
80 90

AAATCACCTATGAAATTAAATCAGAAAAATGCCAAAGCTGTTGGATTACCAGCAAAAGGA 482
K S P M K L N Q K N A K A V G L P A K G
100 110

TCGGATGTAAAAGTTGGTGATCAAGTTCGTGTTTCTGGTTGGGGTTATCTTGAAGAAGGA 542
S D V K V G D Q V R V S G W G Y L E E G
120 130

AGTTATTCATTACCATCTGAATTAAGACGTGTTGATATTGCTGTTGTATCACGTAAAGAA 602
S Y S L P S E L R R V D I A V V S R K E
140 150

TGTAATGAATTATATTCAAAGCTAATGCTGAAGTTACTGATAATATGATTTGTGGTGGT 662
C N E L Y S K A N A E V T D N M I C G G
160 170

GATGTTGCAAATGGTGGTAAAGATTCTTGTCAAGGTGATTCTGGTGGACCGGTTGTTGAT 722
D V A N G G K D S C Q G D S G G P V V D
180 190

GTTAAAAATAATCAAGTTGTTGGTATTGTTTCATGGGGTTATGTTGTGCACGTAAAGGT 782
V K N N Q V V G I V S W G Y G C A R K G
200 210

TATCCAGGTGTTTATACACGTGTTGGTAATTTTATCGATTGGATTGAATCAAAACGTTCA 842
Y P G V Y T R V G N F I D W I E S K R S
220 230

CAGTGATTGATAAAAAACAATTTTCGACATATAAAAAATTGGCAAATGATGCCATTATATGTT 902
Q *

ATCATTATGTTTCTGGTGATTCTTTTCTACTTTTATCATTTTATTTTATATTTCGAGCTG 962
AAAACAAACAATCATGATTGTATAGGATTTGTTGTTGTTTTCATTTTAAAAAATCCA 1022
AAATAAATAAATAAATAATCATTTCACAAAAAAAAG 1059

Fig. 1

DerpIII	MIYNILIVL	LLAINTLANP	ILPASPNATI	VGGEKALAGE	CPYQISLQS-	49
CRAYFISH	-----	-----	-----I	VGGTDAVLGE	FPYQLSFQET	21
Consensus			I	VGG..A..GE	.PYQ.S.Q..	50
DerpIII	----SSHFCG	GTILDEYWIL	TAAHCVAG--	-QTASKLSIR	YNSLKHSL--	90
CRAYFISH	FLGFSFHFCG	ASIYNENYAI	TAGHCVYGDD	YENPSGLQIV	AGELDMSVNE	71
Consensus	S.HFCG	..I..E....	TA.HCV.G..	S.L.I.	...L..S...	100
DerpIII	GGEK-ISVAK	IFAHEKYDSY	QIDNDIALIK	LKSPMKLNQK	NAKAVGLPAK	139
CRAYFISH	GSEQTITVSK	IILHENFDYD	LLDNDISLLK	LSGSLTFN-N	NVAPIALPAQ	120
Consensus	G.E..I.V.K	I..HE..D..	..DNDI.L.K	L.....N..	N.....LPA.	150
DerpIII	GSDVKVGDQV	RVSGWGYLEE	GSYSLPSELR	RVDIAVVSRK	ECNELYSKAN	189
CRAYFISH	G-HTATGN-V	IVTGWGTTSE	GG-NTPDVLQ	KVTVPPLVSDA	ECRDDY--GA	165
Consensus	G.....G..V	.V.GWG...E	G....P..L.	.V....VS..	EC...Y....	200
DerpIII	AEVTDNMICG	GDVANGGKDS	CQGDSGGPVV	--DVKNQV	GIVSWGYGCA	237
CRAYFISH	DEIFDSMICA	G-VPEGGKDS	CQGDSGGPLA	ASDTGSTYLA	GIVSWGYGCA	214
Consensus	.E..D.MIC.	G.V..GGKDS	CQGDSGGP..	..D.....	GIVSWGYGCA	250
DerpIII	RKGYPGVYTR	VGNFIDWIES	KRSQ			261
CRAYFISH	RPGYPGVYTE	VSYHVDWIKA	N-AV			237
Consensus	R.GYPGVYT.	V....DWI..				274

Fig. 2




Fig. 3

NUCLEIC ACIDS ENCODING A HOUSE DUST MITE ALLERGEN, DER P III, AND USES THEREFOR

BACKGROUND OF THE INVENTION

[0001] Approximately 10% of the population become hypersensitized (allergic) upon exposure to antigens from a variety of environmental sources. Those antigens that induce immediate and/or delayed types of hypersensitivity are known as allergens (King, T. P., (1976) *Adv. Immunol.*, 22:77-105). Allergens can include products of grasses, trees, weeds, animal dander, insects, food, drugs, and chemicals. Genetic predisposition of an individual is believed to play a role in the development of immediate allergic responses (Young, R. P. et al., (1990) *Clin. Sci.*, 72:19) such as atopy and anaphylaxis, whose symptoms include hay fever, asthma, and hives.

[0002] The antibodies involved in atopic allergy belong primarily to the IgE class of immunoglobulins. IgE binds to basophils, mast cells and dendritic cells via a specific, high-affinity receptor FcεRI (Kinet, J. P., (1990) *Curr. Opin. Immunol.*, 1499-505). Upon combination of an allergen acting as a ligand with its cognate receptor IgE, FcεRI bound to the IgE may be cross-linked on the cell surface, resulting in physiological manifestations of the IgE-allergen interaction. These physiological effects include the release of, among other substances, histamine, serotonin, heparin, chemotactic factor(s) for eosinophilic leukocytes and/or leukotrienes C4, D4, and E4, which cause prolonged constriction of bronchial smooth muscle cells (Hood, L. E. et al., *Immunology* (2nd ed.), The Benjamin/Cummings Publishing Co., Inc. (1984)). Hence, the ultimate consequence of the interaction of an allergen with IgE is allergic symptoms triggered by the release of the aforementioned mediators. Such symptoms may be systemic or local in nature, depending on the route of entry of the antigen and the pattern of deposition of IgE on mast cells or basophils. Local manifestations generally occur on epithelial surfaces at the site of entry of the allergen. Systemic effects can induce anaphylaxis (anaphylactic shock) which results from IgE-basophil response to circulating (intravascular) antigen.

[0003] Studies with purified allergens have shown that about 80% of patients allergic to the mite *Dermatophagoides pteronyssinus* produce IgE reactive to Der p I and Der p II (Chapman M. D. et al., *J. Immunol.* (1980) 125:587-92; Lind P., *J. Allergy Clin. Immunol.* (1985) 76:753-61; Van der Zee J. S. et al., *Allergy Clin. Immunol.* (1988) 81:884-95). For about half the patients these specificities constitute 50% of the IgE antimitte antibody. The allergen Der p III, recently identified as trypsin, (Stewart G. A. et al., *Immunology* (1992) 75:29-35) reacts with a similar or higher frequency (Stewart G. A. et al., supra; Ford S. A. et al., *Clin. Exp. Allergy* (1989) 20:27-31). However, in the only quantitative study reported to date, the investigators determined that the level of IgE binding to Der p III was considerably less than for Der p I. Electrophoretic techniques (Ford S. A. et al., supra; Bengtsson A. et al., *Int. Arch. Allergy Appl. Immunol.* (1986) 80:383-90; Lind P. et al., *Scand. J. Immunol.* (1983) 17:263-73; Tovey E. R. et al., *J. Allergy Clin. Immunol.* (1987) 79:93-102) have shown that most sera contain IgE which recognize other allergens.

[0004] The significance of the IgE reactivity to Der p III remains uncertain. The reactivity of this group of allergens has been reported to be as low as 16% using a fluid phase assay (Heymann et al., *J. Allergy Clin. Immunol.* (1989) 83:1055-1067) and as high as 100% using RAST assay (Stewart et al., *Immunology* (1992) 75: 29-35). Several others have

reported IgE reactivity between 60-83% (Tovey et al., *J. Allergy Clin. Immunol.* (1987) 79: 93-102; Thomas et al., *Exp. Appl. Acarol.* (1992) 16:153-164; Yasueda et al., *Clin. Exp. Allergy* (1993) 23:384-390). The discrepancies in the frequency of IgE reactivity to the group III allergens may be attributable to either the differences in the purity of the allergen preparation studied or the differences in sensitivity of the assay techniques used. In order to determine the importance of particular specificities in the allergic reactions, there is a need for quantities of pure allergen, which would enable quantitative IgE binding tests and studies of the frequency and lymphokine profile of T cell responses to the allergen.

[0005] Many patients with sensitivity to house dust mite allergens are treated currently by administration of small, gradually increasing doses of house dust mite extracts. Use of these extracts has multiple drawbacks, including potential anaphylaxis during treatment and the necessity of continuing therapy, often for a period of several years to build up sufficient tolerance and significant diminution of clinical symptoms. The ability to substitute compositions of house dust mite allergen, Der p III, would overcome several of these drawbacks. Thus, a source of pure allergen that could be provided in quantity for use as a diagnostic or therapeutic reagent and therapeutic methods that would overcome the drawbacks associated with house dust mite extracts are highly desirable.

SUMMARY OF INVENTION

[0006] This invention provides isolated nucleic acids encoding peptides having at least one biological activity of Der p III, an allergen of the species *Dermatophagoides pteronyssinus* (house dust mite). A preferred nucleic acid is a cDNA having a nucleotide sequence shown in FIG. 1 (SEQ ID NO:1). The invention also pertains to peptides encoded by all or a portion of such cDNA (SEQ ID NO:1) and having at least one biological activity of Der p III. Also contemplated are isolated nucleic acids which hybridize under high stringency conditions (e.g., equivalent to 20-27° C. below T_m and 1M NaCl) to a nucleic acid having a nucleotide sequence shown in FIG. 1 (SEQ ID NO:1) or which encodes a peptide comprising all or a portion of an amino acid sequence of FIG. 1 (SEQ ID NO:2). Nucleic acids which encode peptides having an activity of Der p III and having at least 50% homology with a sequence shown in FIG. 1 (SEQ ID NO:2) are also featured. Peptides having a Der p III activity produced by recombinant expression of a nucleic acid of the invention, and peptides having a Der p III activity prepared by chemical synthesis are also featured by this invention. Preferred peptides have the ability to induce a T cell response which can include T cell stimulation (measured by, for example, T cell proliferation or cytokine secretion) or T cell non-responsiveness (i.e., contact with the peptide or a complex of the peptide with an MHC molecule of an antigen presenting cell induces the T cell to become unresponsive to stimulatory signals or incapable of proliferation). Other preferred peptides either apart from or in addition to the ability to induce a T cell response, have the ability to bind specific IgE of house dust mite-allergic subjects. Such peptides are useful in diagnosing sensitivity to house dust mites in a subject. Still other peptides, either apart from or in addition to the ability to induce a T cell response, have a significantly reduced or negligible ability to bind house dust mite-allergic IgE. Such peptides are particularly useful as therapeutic agents.

[0007] Other preferred peptides comprise an amino acid sequence shown in FIG. 1 (SEQ ID NO:2). In one embodiment, peptides having a Der p III activity and comprising a portion of the amino acid sequence of FIG. 1 (SEQ ID NO:2) are at least about 8-30 amino acids in length, preferably about 10-20 amino acids in length, and most preferably about 10-16 amino acids in length.

[0008] Another aspect of the invention features antibodies specifically reactive with peptides having a Der p III activity. Peptides having an activity of Der p III can be used in compositions suitable for pharmaceutical administration. For example, such compositions can be used in a manner similar to house dust mite extracts to treat or prevent allergic reactions to house dust mites in a subject. Nucleic acids of the invention and peptides having an activity of Der p III can also be used for diagnosing sensitivity in a subject to house dust mites.

BRIEF DESCRIPTION OF THE DRAWINGS

[0009] FIG. 1 shows the complete nucleotide (SEQ ID NO:1) and deduced amino acid (SEQ ID NO:2) sequences of the Der p III clone.

[0010] FIG. 2 shows the amino acid sequences of Der p III (SEQ ID NO:2) and a trypsin protein from crayfish (SEQ ID NO:3).

[0011] FIG. 3 shows the results of SDS-PAGE analysis of various concentrations of recombinant Der p III (lane 1, 4.3 µg; lane 2, 8.7 µg; lane 3, 13.8 µg; lane 4, 17.4 µg; markers are indicated by M).

DETAILED DESCRIPTION OF THE INVENTION

[0012] This invention pertains to isolated nucleic acids encoding peptides having at least one biological activity of Der p III, a Group III allergen of the species *Dermatophagoides pteronyssinus*. Preferably, the nucleic acid is a cDNA comprising a nucleotide sequence shown in FIG. 1 (SEQ ID NO:1).

[0013] The cDNA shown in FIG. 1 (SEQ ID NO:1) encodes a Der p III peptide which includes a predicted 29 amino acid residue pre-pro region encoded by nucleotides 63 through 149. This leader sequence is not found in the mature Der p III which is encoded by nucleotides 150 through 845. The deduced amino acid sequence based on this cDNA is also shown in FIG. 1 (SEQ ID NO:2). The cDNA encodes a 232 residue mature peptide having a predicted molecular weight of 24,985 daltons including seven cysteine residues. A polyadenylation signal sequence 179 nucleotides after the last base is present in the cDNA (See FIG. 1). A culture of *E. coli* transfected with an expression vector containing the cDNA encoding Der p III was deposited under the Budapest Treaty with the American Type Culture Collection on Oct. 15, 1993 and assigned accession number 69472.

[0014] Accordingly, one aspect of this invention pertains to isolated nucleic acids comprising a nucleotide sequence encoding Der p III, fragments thereof encoding peptides having at least one biological activity of Der p III and/or equivalents of such nucleic acids. The term nucleic acid as used herein is intended to include such fragments and equivalents. The term equivalent is intended to include nucleotide sequences encoding functionally equivalent Der p III peptides having an activity of Der p III. As defined herein, a peptide having an activity of Der p III has at least one biological activity of the Der p III allergen. Equivalent nucleotide

sequences include sequences that differ by one or more nucleotide substitutions, additions or deletions, such as allelic variants, and include sequences that differ from the nucleotide sequence encoding Der p III shown in FIG. 1 (SEQ ID NO:1) due to the degeneracy of the genetic code. Equivalents includes nucleotide sequences that hybridize under stringent conditions (i.e., equivalent to about 20-27° C. below melting temperature (T_m) and about 1M NaCl) to the nucleotide sequence of Der p III shown in FIG. 1 (SEQ ID NO:1).

[0015] Peptides referred to herein as having an activity of Der p III or having a Der p III activity are defined herein as peptides that have an amino acid sequence corresponding to all or a portion of the amino acid sequence of Der p III shown in FIG. 1 (SEQ ID NO:2). For example, a peptide having an activity of Der p III may have the ability to induce a response in Der p III restricted T cells such as stimulation (e.g., T cell proliferation or cytokine secretion) or induce T cell non-responsiveness. Alternatively, or additionally, a peptide having an activity of Der p III may have the ability to bind (to be recognized by) immunoglobulin E antibodies of house dust mite-allergic subjects. Peptides which bind IgE are useful in methods of detecting allergic sensitivity to Der p III in a subject. Peptides that do not bind IgE, or bind IgE to a lesser extent than a purified, native Der p III protein binds IgE are particularly useful as therapeutic agents.

[0016] In one embodiment, the nucleic acid is a cDNA encoding a peptide having an activity of Der p III. Preferably, the nucleic acid is a cDNA molecule comprising at least a portion of the nucleotide sequence encoding Der p III shown in FIG. 1 (SEQ ID NO:1). A preferred portion of the cDNA molecule of FIG. 1 (SEQ ID NO:1) includes the coding region of the molecule.

[0017] In another embodiment, the nucleic acid of the invention encodes a peptide having an activity of Der p III and comprising an amino acid sequence shown in FIG. 1 (SEQ ID NO:2). Preferred nucleic acids encode a peptide having a Der p III activity and having at least about 50% homology, more preferably at least about 60% homology and most preferably at least about 70% homology with the sequence shown in FIG. 1 (SEQ ID NO:2). Nucleic acids which encode peptides having a Der p III activity and having at least about 90%, more preferably at least about 95%, and most preferably at least about 98-99% homology with a sequence set forth in FIG. 1 (SEQ ID NO:2) are also within the scope of the invention. Homology refers to sequence similarity between two peptides having an activity of Der p III or between two nucleic acid molecules. Homology can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequence is occupied by the same base or amino acid, then the molecules are homologous at that position. A degree of homology between sequences is a function of the number of matching or homologous positions shared by the sequences.

[0018] Another aspect of the invention provides a nucleic acid that hybridizes under high or low stringency conditions to a nucleic acid encoding a peptide having all or a portion of an amino acid sequence shown in FIG. 1 (SEQ ID NO:2). Appropriate stringency conditions which promote DNA hybridization, for example, 6.0× sodium chloride/sodium citrate (SSC) at about 45° C., followed by a wash of 2.0×SSC at 50° are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of

about 2.0×SSC at 50° C. to a high stringency of about 0.2×SSC at 50° C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22° C., to high stringency conditions at about 65° C.

[0019] Isolated nucleic acids encoding a peptide having an activity of Der p III, as described herein, and having a sequence that differs from the nucleotide sequence shown in FIG. 1 (SEQ ID NO:1) due to degeneracy in the genetic code are also within the scope of the invention. Such nucleic acids differ in sequence from the sequence of FIG. 1 (SEQ ID NO:1) encode functionally equivalent peptides (i.e., peptides having an activity of Der p III), but due to degeneracy in the genetic code. For example, a number of amino acids are designated by more than one triplet. Codons that specify the same amino acid, or synonyms (for example, CAU and CAC are synonyms for histidine) may result in “silent” mutations that do not affect the amino acid sequence of the Der p III protein. However, it is expected that DNA sequence polymorphisms that do lead to changes in the amino acid sequence of Der p III will exist within the house dust mite population. One skilled in the art will appreciate that these variations in one or more nucleotides (up to about 3-4% of the nucleotides) of the nucleic acids encoding peptides having an activity of Der p III may exist among individual house dust mites due to natural allelic variation. Any and all such nucleotide variations and resulting amino acid polymorphisms are within the scope of this invention. Furthermore, there may be one or more isoforms or related, cross-reacting family members of Der p III. Such isoforms or family members are defined as proteins related in function and amino acid sequence to Der p III, but are encoded by genes at different loci.

[0020] Fragments of a nucleic acid encoding Der p III are also within the scope of the invention. As used herein, a fragment of a nucleic acid encoding Der p III refers to a nucleotide sequence having fewer nucleotides than the nucleotide sequence encoding the entire amino acid sequence of Der p III protein and which encodes a peptide having an activity of Der p III (i.e., a peptide having at least one biological activity of the Der p III allergen) as defined herein.

[0021] Preferred nucleic acid fragments encode peptides of at least 10 amino acid residues in length, preferably about 10-20 amino acid residues in length, and more preferably about 10-16 amino acid residues in length. Nucleic acid fragments that encode peptides having a Der p III activity of at least about 30 amino acid residues in length, at least about 40 amino acid residues in length, at least about 60 amino acid residues in length, at least about 80 amino acid residues in length, at least about 100 amino acid residues in length, and at least about 200 amino acid residues in length or more are also within the scope of this invention.

[0022] Nucleic acid fragments within the scope of the invention include those capable of hybridizing under high or low stringency with nucleic acids from other animal species for use in screening protocols to detect Der p III or allergens that are cross-reactive with Der p III. Generally, the nucleic acid encoding a peptide having an activity of Der p III will be selected from the bases encoding the mature protein, however, in some instances, it may be desirable to select all or part of a peptide from the leader sequence portion of the nucleic acid of the invention. Nucleic acids within the scope of the invention may also contain linker sequences, modified restriction endonuclease sites and other sequences useful for

molecular cloning, expression or purification of recombinant peptides having an activity of Der p III.

[0023] A nucleic acid encoding a peptide having an activity of Der p III may be obtained from mRNA present in house dust mites of the species *Dermatophagoides pteronyssinus*. It should also be possible to obtain nucleic acids encoding Der p III from *Dermatophagoides pteronyssinus* genomic DNA. For example, the gene encoding Der p III can be cloned from either a cDNA or a genomic library in accordance with protocols herein described. A cDNA encoding Der p III can be obtained by isolating total mRNA from *Dermatophagoides pteronyssinus*. Double stranded cDNAs can then be prepared from the total mRNA. Subsequently, the cDNAs can be inserted into a suitable plasmid or bacteriophage vector using any one of a number of known techniques. Genes encoding Der p III can also be cloned using established polymerase chain reaction techniques in accordance with the nucleotide sequence information provided by the invention. The nucleic acids of the invention can be DNA or RNA. A preferred nucleic acid is a cDNA encoding Der p III having the sequence depicted in FIG. 1 (SEQ ID NO:1).

[0024] This invention also provides expression vectors containing a nucleic acid encoding a peptide having an activity of Der p III operably linked to at least one regulatory sequence. Operably linked is intended to mean that the nucleotide sequence is linked to a regulatory sequence in a manner which allows expression of the nucleotide sequence. Regulatory sequences are art-recognized and are selected to direct expression of the peptide having an activity of Der p III by a transfected host cell. Accordingly, the term regulatory sequence includes promoters, enhancers and other expression control elements. Such regulatory sequences are described in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990). It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transformed and/or the type of protein desired to be expressed. In one embodiment, the expression vector includes a DNA encoding a peptide having an activity of Der p III. Such expression vectors can be used to transfect cells to thereby produce proteins or peptides, including fusion proteins or peptides encoded by nucleic acids as described herein.

[0025] This invention further pertains to a host cell transfected to express a peptide having an activity of Der p III. The host cell may be any procaryotic or eucaryotic cell. For example, a peptide having an activity of Der p III may be expressed in bacterial cells such as *E. coli*, insect cells (baculovirus), yeast, or mammalian cells such as Chinese hamster ovary cell (CHO). Other suitable host cells are referred to in Goeddel (1990), *supra*, or known to those skilled in the art.

[0026] Expression in eucaryotic cells such as mammalian, yeast, or insect cells can lead to partial or complete glycosylation and/or formation of relevant inter- or intra-chain disulfide bonds of a recombinant peptide product. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, et al., (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz et al., (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, Calif.), Baculovirus vectors available for expression of proteins in cultured insect cells (SF 9 cells) include the pAc series (Smith et al., (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow, V. A., and Summers, M. D., (1989) *Virology* 170:31-39). Generally, COS

cells (Gluzman, Y., (1981) *Cell* 23:175-182) are used in conjunction with such vectors as pCDM 8 (Aruffo, A. and Seed, B., (1987) *Proc. Natl. Acad. Sci. USA* 84:8573-8577) for transient amplification/expression in mammalian cells, while CHO (dhfr⁻ Chinese Hamster Ovary) cells are used with vectors such as pMT2PC (Kaufman et al., (1987), *EMBO J.* 6:187-195) for stable amplification/expression in mammalian cells. Vector DNA can be introduced into mammalian cells via conventional techniques such as calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, or electroporation. Suitable methods for transforming host cells can be found in Sambrook et al., (*Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory textbooks.

[0027] Expression in procaryotes is most often carried out in *E. coli* with either fusion or non-fusion inducible expression vectors. Fusion vectors usually add a number of NH₂ terminal amino acids to the expressed target gene. These NH₂ terminal amino acids often are referred to as a reporter group. Such reporter groups usually serve two purposes: 1) to increase the solubility of the target recombinant protein; and 2) to aid in the purification of the target recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the reporter group and the target recombinant protein to enable separation of the target recombinant protein from the reporter group subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Amrad Corp., Melbourne, Australia), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) which fuse glutathione S-transferase, maltose E binding protein, or protein A, respectively, to the target recombinant protein. A preferred reporter group is poly(His), which renders the recombinant fusion protein easily purifiable by metal chelate chromatography.

[0028] Inducible non-fusion expression vectors include pTrc (Amann et al., (1988) *Gene* 69:301-315) and pET11d (Studier et al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 60-89). While target gene expression relies on host RNA polymerase transcription from the hybrid trp-lac fusion promoter in pTrc, expression of target genes inserted into pET11d relies on transcription from the T7 gn10-lac 0 fusion promoter mediated by coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7 gn1 under the transcriptional control of the lacUV 5 promoter.

[0029] One strategy to maximize recombinant Der p III expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant to protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 119-128). Another strategy would be to alter the nucleic acid encoding Der p III to be inserted into an expression vector so that the individual codons for each amino acid would be those preferentially utilized in highly expressed *E. coli* proteins (Wada (1992) *Nuc. Acids Res.* 20:2111-2118). Such alteration of nucleic acids of the invention can be carried out by standard DNA synthesis techniques.

[0030] The nucleic acids of the invention can also be chemically synthesized using standard techniques. Various methods of chemically synthesizing polydeoxynucleotides are known, including solid-phase synthesis which, like peptide synthesis, has been fully automated in commercially available DNA synthesizers (See, e.g., Itakura et al., U.S. Pat. No. 4,598,049; Caruthers et al. U.S. Pat. No. 4,458,066; and Itakura U.S. Pat. Nos. 4,401,796 and 4,373,071, incorporated by reference herein).

[0031] The present invention further pertains to methods of producing peptides that have an activity of Der p III. For example, a host cell transfected with a nucleic acid vector directing expression of a nucleotide sequence encoding a peptide having an activity of Der p III can be cultured under appropriate conditions to allow expression of the peptide to occur. The peptide may be secreted and isolated from a mixture of cells and medium containing the peptide. Alternatively, the peptide may be retained cytoplasmically and the cells harvested, lysed and the protein isolated. A cell culture includes host cells, media and other byproducts. Suitable media for cell culture are well known in the art. Peptides of the invention can be isolated from cell culture medium, host cells, or both using techniques known in the art for purifying proteins including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis, and immunoaffinity purification with antibodies specific for such peptides.

[0032] Another aspect of the invention pertains to isolated peptides having an activity of Der p III. A peptide having an activity of Der p III has at least one biological activity of the Der p III allergen. For example, a peptide having an activity of Der p III may have the ability to induce a response in Der p III specific T cells such as stimulation (T cell proliferation or cytokine secretion) or induce T cell non-responsiveness. In one embodiment, a peptide having an activity of Der p III stimulates T cells as evidenced by, for example, T cell proliferation or cytokine secretion. In another embodiment, peptides having a Der p III activity induce T cell non-responsiveness in which T cells are unresponsive to a subsequent challenge with a Der p III peptide or native Der p III protein following exposure to the peptide. In yet another embodiment, a peptide having a Der p III activity has reduced IgE binding activity compared to purified, native Der p III protein. A peptide having an activity of Der p III may differ in amino acid sequence from the Der p III sequence depicted in FIG. 1 (SEQ ID NO:2) but such differences result in a modified protein which functions in the same or similar manner as a native Der p III protein or which has the same or similar characteristics of Der p III protein. Various modifications of the Der p III protein to produce these and other functionally equivalent peptides are described in detail herein. The term peptide, as used herein, refers to peptides, proteins, and polypeptides.

[0033] A peptide can be produced by modification of the amino acid sequence of the Der p III protein shown in FIG. 1 (SEQ ID NO:2), such as by a substitution, addition, or deletion of one or more amino acid residues. Such modifications may be directed to amino acid residues not involved in a biological activity of the peptide, or the modifications may be directed to such amino acid residues in order to enhance, or eliminate a particular biological activity. Peptides of the invention are at least about 8 amino acid residues in length, preferably about 10-20 amino acid residues in length and more preferably about 10-16 amino acid residues in length.

Peptides having an activity of Der p III of at least about 30 amino acid residues in length, at least about 40 amino acid residues in length, at least about 60 amino acid residues in length, at least about 80 amino acid residues in length, at least about 100 amino acid residues in length, and at least about 200 amino acid residues or more in length are also included within the scope of this invention.

[0034] In another embodiment of the invention, peptides from a related Group III allergen of the species *Dermatophagoides farinae*, Der f III, are provided. Such peptides have been isolated from a purified, native Der f III protein and comprise the sequences: IVGGVKAKAGDSPYQISLQSSSHFXGGSILD (SEQ ID NO:15), an N-terminal sequence; MICGGDVANGGVDSQGD (SEQ ID NO:10), an internal peptide; and MTLDQTNAPVPLPTS (SEQ ID NO:12), an internal peptide.

[0035] Yet another embodiment of the invention provides a substantially pure preparation of a peptide having an activity of Der p III. Such a preparation is substantially free of proteins and peptides with which the peptide naturally occurs (i.e., other house dust mite peptides), either in a cell or when secreted by a cell.

[0036] The term isolated as used herein refers to a nucleic acid or peptide that is substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. Such peptides are also characterized as being free of all other house dust mite proteins. Accordingly, an isolated peptide having an activity of Der p III is produced recombinantly or synthetically and is substantially free of cellular material and culture medium or substantially free of chemical precursors or other chemicals and is free of all other house dust mite proteins. An isolated nucleic acid is also free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the organism from which the nucleic acid is derived.

[0037] Peptides having an activity of Der p III can be obtained, for example, by screening peptides recombinantly produced from the corresponding fragment of the nucleic acid of Der p III encoding such peptides. In addition, fragments can be chemically synthesized using techniques known in the art such as conventional Merrifield solid phase f-Moc or t-Boc chemistry. For example, the Der p III protein may be arbitrarily divided into fragments of desired length with no overlap of the fragments, or preferably divided into overlapping fragments of a desired length. The fragments can be produced (recombinantly or by chemical synthesis) and tested to identify those peptides having a Der p III activity, i.e., the ability to induce a T cell response such as T cell stimulation (T cell proliferation, cytokine secretion) or T cell non-responsiveness, and/or has reduced IgE binding activity.

[0038] In one embodiment, peptides having an activity of Der p III can be identified by the ability of the peptide to stimulate T cells or to induce T cell non-responsiveness. Peptides which stimulate T cells, as determined by, for example, T cell proliferation or cytokine secretion are defined herein as comprising at least one T cell epitope. T cell epitopes are believed to be involved in initiation and perpetuation of the immune response to the protein allergen which is responsible for the clinical symptoms of allergy. These T cell epitopes are thought to trigger early events at the level of the T helper cell by binding to an appropriate HLA molecule on the surface of an antigen presenting cell, thereby stimulating the T cell subpopulation with the relevant T cell receptor for

the epitope. These events lead to T cell proliferation, lymphokine secretion, local inflammatory reactions, recruitment of additional immune cells to the site of antigen/T cell interaction, and activation of the B cell cascade, leading to the production of antibodies. One isotype of these antibodies, IgE, is fundamentally important to the development of allergic symptoms and its production is influenced early in the cascade of events at the level of the T helper cell, by the nature of the lymphokines secreted. A T cell epitope is the basic element, or smallest unit of recognition by a T cell receptor, where the epitope comprises amino acids essential to receptor recognition. Amino acid sequences which mimic those of the T cell epitopes and which modify the allergic response to protein allergens are within the scope of this invention.

[0039] Screening peptides for those which retain a Der p III activity as described herein can be accomplished using one or more of several different assays. For example, in vitro, Der p III T cell stimulatory activity is assayed by contacting a peptide known or suspected of having a Der p III activity with an antigen presenting cell which presents appropriate MHC molecules in a T cell culture. Presentation of a peptide having a Der p III activity in association with appropriate MHC molecules to T cells in conjunction with the necessary costimulation has the effect of transmitting a signal to the T cell that induces the production of increased levels of cytokines, particularly of interleukin-2 and interleukin-4. The culture supernatant can be obtained and assayed for interleukin-2 or other known cytokines. For example, any one of several conventional assays for interleukin-2 can be employed, such as the assay described in *Proc. Natl. Acad. Sci USA*, 86:1333 (1989), the pertinent portions of which are incorporated herein by reference. A kit for an assay for the production of interferon is also available from Genzyme Corporation (Cambridge, Mass.).

[0040] Alternatively, a common assay for T cell proliferation entails measuring tritiated thymidine incorporation. The proliferation of T cells can be measured in vitro by determining the amount of ³H-labeled thymidine incorporated into the replicating DNA of cultured cells. Therefore, the rate of DNA synthesis and, in turn, the rate of cell division can be quantified.

[0041] In another embodiment, a peptide having a Der p III activity is screened for the ability to induce T cell non-responsiveness. The ability of a peptide known to stimulate T cells (as determined by one or more of the above described assays) to inhibit or completely block the activity of purified native Der p III or portion thereof and induce a state of non-responsiveness can be determined using subsequent attempts at stimulation of the T cells with antigen presenting cells that present native Der p III, or a peptide having a Der p III activity, following exposure to the peptide having a Der p III activity. If the T cells are unresponsive to the subsequent activation attempts, as determined by interleukin-2 synthesis and/or T cell proliferation, a state of non-responsiveness has been induced. See, e.g., Gimmi, et al. (1993) *Proc. Natl. Acad. Sci USA*, 90:6586-6590; and Schwartz (1990) *Science*, 248:1349-1356, for assay systems that can be used as the basis for an assay in accordance with the present invention.

[0042] In yet another embodiment, peptides having a Der p III activity are identified by IgE binding activity. For therapeutic purposes, peptides of the invention preferably do not bind IgE specific for a house dust mite allergen or bind such IgE to a substantially lesser extent than the purified, native house dust mite allergen binds such IgE. Reduced IgE bind-

ing activity refers to IgE binding activity that is less than that of purified, native Der p III protein. If a peptide having a Der p III activity is to be used as a diagnostic reagent, it is not necessary that the peptide have reduced IgE binding activity compared to the native Der p III allergen. IgE binding activity of peptides can be determined, for example, by an enzyme-linked immunosorbent assay (ELISA) using, for example, sera obtained from a subject (i.e., an allergic subject) that has been previously exposed to the native Der p III allergen. Briefly, the peptide suspected of having a Der p III activity is coated onto wells of a microtiter plate. After washing and blocking the wells, antibody solution consisting of the plasma of an allergic subject who has been exposed to a peptide suspected of having a Der p III activity is incubated in the wells. The plasma is generally depleted of IgG before incubation. A labeled secondary antibody is added to the wells and incubated. The amount of IgE binding is then quantified and compared to the amount of IgE bound by a purified, native Der p III protein. Alternatively, the IgE binding activity of a peptide can be determined by Western blot analysis. For example, a peptide suspected of having a Der p III activity is run on a polyacrylamide gel using SDS-PAGE. The peptide is then transferred to nitrocellulose and subsequently incubated with sera from an allergic subject. After incubation with a labeled secondary antibody, the amount of IgE bound is then determined and quantified.

[0043] Another assay which can be used to determine the IgE binding activity of a peptide is a competition ELISA assay. Briefly, an IgE antibody pool is generated by combining plasma from house dust mite allergic subjects that have been shown by direct ELISA to have IgE reactive with native Der p III. This pool is used in ELISA competition assays to compare IgE binding of native Der p III and a peptide suspected of having a Der p III activity. IgE binding for the native Der p III and a peptide suspected of having a Der p III activity is determined and quantified.

[0044] If a peptide having an activity of Der p III binds IgE and is to be used as a therapeutic agent, it is preferable that such binding does not result in the release of mediators (e.g., histamines) from mast cells or basophils. To determine whether a peptide which binds IgE results in the release of mediators, a histamine release assay can be performed using standard reagents and protocols obtained from Amac, Inc. (Westbrook, Me.). Briefly, a buffered solution of a peptide suspected of having a Der p III activity is combined with an equal volume of whole heparinized blood from an allergic subject. After mixing and incubation, the cells are pelleted and the supernatants are processed and analyzed using a radioimmunoassay to determine the amount of histamine released.

[0045] Peptides having an activity of Der p III which are to be used as therapeutic agents are preferably tested in mammalian models of house dust mite atopy, such as the mouse model disclosed in Tamura et al., (1986) *Microbiol. Immunol.* 30:883-896, or in U.S. Pat. No. 4,939,239, or in the primate model disclosed in Chiba et al., (1990) *Int. Arch. Allergy Immunol.* 93:83-88. Initial screening for IgE binding to a peptide having an activity of Der p III may be performed by scratch tests or intradermal skin tests on laboratory animals or human volunteers, or in in vitro systems such as RAST, RAST inhibition, ELISA assay, RIA (radioimmunoassay), or a histamine release assay, as described above.

[0046] It is possible to modify the structure of a peptide having an activity of Der p III for such purposes as increasing

solubility, enhancing therapeutic or prophylactic efficacy, or stability (e.g., shelf life ex vivo and resistance to proteolytic degradation in vivo). Such modified peptides are considered functional equivalents of peptides having an activity of Der p III as defined herein. A modified peptide can be produced in which the amino acid sequence has been altered, such as by amino acid substitution, deletion, or addition, to modify immunogenicity and/or reduce allergenicity, or to which a component has been added for the same purpose.

[0047] For example, a peptide having an activity of Der p III can be modified so that it maintains the ability to induce T cell non-responsiveness and bind MHC proteins without the ability to induce a strong proliferative response or, possibly, any proliferative response when administered in immunogenic form. In this instance, critical binding residues for T cell receptor function can be determined using known techniques (e.g., substitution of each residue and determination of the presence or absence of T cell reactivity). Those residues shown to be essential for interaction with the T cell receptor can be modified by replacing the essential amino acid with another, preferably similar amino acid residue (a conservative substitution) whose presence is shown to enhance, diminish but not eliminate, or not affect T cell reactivity. In addition, those amino acid residues which are not essential for T cell receptor interaction can be modified by being replaced by another amino acid whose incorporation may enhance, diminish but not eliminate, or not affect T cell reactivity, but whose incorporation does not eliminate binding to relevant MHC.

[0048] Additionally, a peptide having an activity of Der p III can be modified by replacing an amino acid shown to be essential for interaction with the MHC protein complex with another, preferably similar amino acid residue (conservative substitution) whose presence is shown to enhance, diminish but not eliminate or affect T cell activity. In addition, amino acid residues which are not essential for interaction with the MHC protein complex but which still bind the MHC protein complex can be modified by being replaced by another amino acid whose incorporation may enhance, not affect, or diminish but not eliminate T cell reactivity. Preferred amino acid substitutions for non-essential amino acids include, but are not limited to substitutions with alanine, glutamic acid, or a methyl amino acid.

[0049] Another example of modification of a peptide having an activity of Der p III is substitution of cysteine residues preferably with alanine, serine, threonine, leucine or glutamic acid residues to minimize dimerization via disulfide linkages. In addition, amino acid side chains of peptides of the invention can be chemically modified. Another modification is cyclization of the peptide.

[0050] In order to enhance stability and/or reactivity, a peptide having an activity of Der p III can be modified to incorporate one or more polymorphisms in the amino acid sequence of the protein allergen resulting from any natural allelic variation. Additionally, D-amino acids, non-natural amino acids, or non-amino acid analogs can be substituted or added to produce a modified protein within the scope of this invention. Furthermore, a peptide having an activity of Der p III can be modified using polyethylene glycol (PEG) according to the method of A. Schon and co-workers (Wie et al., supra) to produce a protein conjugated with PEG. In addition, PEG can be added during chemical synthesis of the protein. Other modifications of a peptide having an activity of Der p III include reduction/alkylation (Tarr, *Methods of Protein*

Microcharacterization, J. E. Silver ed., Humana Press, Clifton N.J. 155-194 (1986)); acylation (Tarr, supra); chemical coupling to an appropriate carrier (Mishell and Shiigi, eds., *Selected Methods in Cellular Immunology*, W H Freeman, San Francisco, Calif. (1980), U.S. Pat. No. 4,939,239; or mild formalin treatment (Marsh (1971), *Int. Arch. of Allergy and Appl. Immunol.* 41:199-215).

[0051] To facilitate purification and potentially increase solubility of a peptide having an activity of Der p III, it is possible to add an amino acid fusion moiety to the peptide backbone. For example, hexa-histidine can be added to the protein for purification by immobilized metal ion affinity chromatography (Hochuli, E. et al., (1988) *Bio/Technology* 6:1321-1325). In addition, to facilitate isolation of peptides free of irrelevant sequences, specific endoprotease cleavage sites can be introduced between the sequences of the fusion moiety and the peptide. In order to successfully desensitize a subject to Der p III protein or related allergen, it may be necessary to increase the solubility of the protein by adding functional groups to the protein or by omitting hydrophobic regions of the protein.

[0052] To potentially aid proper antigen processing of T cell epitopes within Der p III, canonical protease sensitive sites can be engineered between regions, each comprising at least one T cell epitope via recombinant or synthetic methods. For example, charged amino acid pairs, such as KK or RR, can be introduced between regions within a protein or fragment during recombinant construction thereof. The resulting peptide can be rendered sensitive to cleavage by cathepsin and/or other trypsin-like enzymes which would generate portions of the protein containing one or more T cell epitopes. In addition, such charged amino acid residues can result in an increase in the solubility of the peptide.

[0053] Site-directed mutagenesis of a nucleic acid encoding a peptide having an activity of Der p III can be used to modify the structure of the peptide by methods known in the art. Such methods may, among others, include polymerase chain reaction (PCR) with oligonucleotide primers bearing one or more mutations (Ho et al., (1989) *Gene* 77:51-59) or total synthesis of mutated genes (Hostomsky, Z. et al., (1989) *Biochem. Biophys. Res. Comm* 161:1056-1063). To enhance recombinant protein expression, the aforementioned methods can be applied to change the codons present in the cDNA sequence of the invention to those preferentially utilized by the host cell in which the recombinant protein is being expressed (Wada et al., supra).

[0054] Another aspect of the invention pertains to an antibody specifically reactive with a peptide having an activity of Der p III. The antibodies of this invention can be used to standardize allergen extracts or to isolate the naturally-occurring or native form of Der p III. For example, by using peptides having an activity of Der p III based on the cDNA sequence of Der p III, anti-protein/anti-peptide antisera or monoclonal antibodies can be made using standard methods. A mammal such as a mouse, a hamster or a rabbit can be immunized with an immunogenic form of the peptide (e.g., Der p III protein or an antigenic fragment which is capable of eliciting an antibody response). Techniques for conferring immunogenicity on a protein or peptide include conjugation to carriers or other techniques are well known in the art. A peptide having an activity of Der p III can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or

serum. Standard ELISA or other immunoassays can be used with the immunogen as antigen to assess the levels of antibodies.

[0055] Following immunization, anti-Der p III antisera can be obtained and, if desired, polyclonal anti-Der p III antibodies isolated from the serum. To produce monoclonal antibodies, antibody-producing cells (lymphocytes) can be harvested from an immunized animal and fused by standard somatic cell fusion procedures with immortalizing cells such as myeloma cells to yield hybridoma cells. Such techniques are well known in the art, for example the hybridoma technique originally developed by Kohler and Milstein, (*Nature* (1975) 256:495-497) as well as other techniques such as the human B cell hybridoma technique (Kozbar *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy* (1985) Alan R. Liss, Inc. pp. 77-96). Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with a peptide having an activity of Der p III and the monoclonal antibodies isolated.

[0056] The term antibody as used herein is intended to include fragments thereof which are also specifically reactive with a peptide having an activity of Der p III. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above for whole antibodies. For example, F(ab')₂ fragments can be generated by treating antibody with pepsin. The resulting F(ab')₂ fragment can be treated to reduce disulfide bridges to produce Fab' fragments. The antibody of the present invention is further intended to include bispecific and chimeric molecules having an anti-Der p III portion.

[0057] Another aspect of this invention provides T cell clones and soluble T cell receptors specifically reactive with a peptide having an activity of Der p III. Monoclonal T cell populations (i.e., T cells genetically identical to one another and expressing identical T cell receptors) can be derived from a subject sensitive to Der p III, followed by repetitive in vitro stimulation with a Der p III protein or peptide having an activity of Der p III in the presence of MHC-matched antigen-presenting cells. Single Der p III MHC responsive cells can then be cloned by limiting dilution and permanent lines expanded and maintained by periodic in vitro restimulation. Alternatively, Der p III specific T-T hybridomas can be produced by a technique similar to B cell hybridoma production. For example, a mammal, such as a mouse, is immunized with a peptide having an activity of Der p III, T cells from the mammal can be purified and fused with an autonomously growing T cell tumor line. From the resulting hybridomas, cells responding to a peptide having an activity of Der p III are selected and cloned. Procedures for propagating monoclonal T cell populations are described in *Cellular and Molecular Immunology* (Abul K. Abbas et al. ed.), W.B. Saunders Company, Philadelphia, Pa. (1991) page 139. Soluble T cell receptors specifically reactive with a peptide having an activity of Der p III can be obtained by immunoprecipitation using an antibody against the T cell receptor as described in *Immunology: A Synthesis* (Second Edition), Edward S. Golub et al., ed., Sinauer Associates, Inc., Sunderland, Mass. (1991) pages 366-369.

[0058] T cell clones specifically reactive with a peptide having an activity of Der p III can be used to isolate and molecularly clone the gene encoding the relevant T cell receptor. In addition, a soluble T cell receptor specifically reactive

with a peptide having an activity of Der p III can be used to interfere with or inhibit antigen-dependent activation of the relevant T cell subpopulation, for example, by administration to a subject sensitive to Der p III. Antibodies specifically reactive with such a T cell receptor can be produced according to the techniques described herein. Such antibodies can be used to block or interfere with the T cell interaction with peptides presented by MHC.

[0059] Exposure of allergic subjects to peptides having an activity of Der p III and which have T cell stimulating activity may cause the appropriate T cell subpopulations to become non-responsive to the respective protein allergen (e.g., fail to stimulate an immune response upon such exposure). In addition, such administration may modify the lymphokine secretion profile as compared with exposure to the naturally-occurring protein allergen or portion thereof (e.g., result in a decrease of IL-4 and/or an increase in IL-2). Furthermore, exposure to peptides having an activity of Der p III which have T cell stimulating activity may influence T cell subpopulations which normally participate in the response to the allergen such that these T cells are drawn away from the site(s) of normal exposure to the allergen (e.g., nasal mucosa, skin, and lung) towards the site(s) of therapeutic administration of the protein or fragment derived therefrom. This redistribution of T cell subpopulations may ameliorate or reduce the ability of a subject's immune system to stimulate the usual immune response at the site of normal exposure to the allergen, resulting in a diminution in allergic symptoms.

[0060] A peptide having an activity of Der p III when administered to a subject sensitive to house dust mites is capable of modifying the B cell response, T cell response, or both the B cell and the T cell response of the subject to the allergen. As used herein, modification of the allergic response of a subject to a house dust mite allergen can be defined as non-responsiveness or diminution in symptoms to the allergen, as determined by standard clinical procedures (See e.g., Varney et. al., (1990) *British Medical Journal* 302:265-269), including diminution in house dust mite induced asthmatic symptoms. As referred to herein, a diminution in symptoms includes any reduction in the allergic response of a subject to the allergen following a treatment regimen with a peptide of the invention. This diminution in symptoms may be determined subjectively (e.g., the subject feels more comfortable upon exposure to the allergen), or clinically, such as with a standard skin test.

[0061] Peptides or antibodies of the present invention can also be used for detecting and diagnosing sensitivity to Der p III. For example, this could be done by combining blood or blood products obtained from a subject to be assessed for sensitivity with a peptide having an activity of Der p III, under conditions appropriate for binding of components in the blood (e.g., antibodies, T cells, B cells) with the peptide(s) and determining the extent to which such binding occurs. Other diagnostic methods for allergic diseases which the peptides or antibodies of the present invention can be used include radio-allergosorbent test (RAST), paper radioimmunosorbent test (PRIST), enzyme linked immunosorbent assay (ELISA), radioimmunoassays (RIA), immuno-radiometric assays (IRMA), luminescence immunoassays (LIA), histamine release assays and IgE immunoblots.

[0062] In another assay, the presence in a subject of IgE specific for Der p III and the ability of T cells of the subject to respond to T cell epitopes of Der p III can be determined by administering to the subject an Immediate Type Hypersensi-

tivity test and/or a Delayed Type Hypersensitivity test (see e.g., *Immunology* (1985) Roitt, I. M., Brostoff, J., Male, D. K. (eds.), C.V. Mosby Co., Gower Medical Publishing, London, N.Y., pp. 19.2-19.18; pp. 22.1-22.10) utilizing a peptide having an activity of Der p III, or a modified form of a peptide having an activity of Der p III, each of which binds IgE specific for the allergen. The same subjects are administered a Delayed Type Hypersensitivity test prior to, simultaneously with, or subsequent to administration of the Immediate Type Hypersensitivity test. Of course, if the Immediate Type Hypersensitivity test is administered prior to the Delayed Type Hypersensitivity test, the Delayed Type Hypersensitivity test would be given to those subjects exhibiting a specific Immediate Type Hypersensitivity reaction. The Delayed Type Hypersensitivity test utilizes a peptide having an activity of Der p III which has human T cell stimulating activity and which does not bind IgE specific for the allergen in a substantial percentage of the population of subjects sensitive to the allergen (e.g., at least about 75%). Those subjects found to have both a specific Immediate type Hypersensitivity reaction and a specific Delayed Type Hypersensitivity reaction are administered an amount of a composition suitable for pharmaceutical administration. The composition comprises the peptide having an activity of Der p III as used in the Delayed Type Hypersensitivity test and a pharmaceutically acceptable carrier or diluent.

[0063] A peptide having an activity of Der p III can be used in methods of treating and preventing allergic reactions to a house dust mite allergen or a cross-reactive protein allergen. Thus, the present invention provides compositions suitable for pharmaceutical administration comprising an amount of at least one peptide having an activity of Der p III and a pharmaceutically acceptable carrier. Administration of the compositions of the present invention to a subject to be desensitized can be carried out using known procedures, at dosages and for periods of time effective to reduce sensitivity (i.e., reduce the allergic response) of the subject to house dust mite. The term subject is intended to include living organisms in which an immune response can be elicited, e.g., mammals. Examples of subjects include humans, dogs, cats, mice, rats, and transgenic species thereof. An amount of at least one peptide having an activity of Der p III necessary to achieve a therapeutic effect may vary according to factors such as the degree of sensitivity of the subject to house dust mite, the age, sex, and weight of the subject, and the ability of a peptide having an activity of Der p III to elicit an antigenic response in the subject. Dosage regimens may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

[0064] The active compound (i.e., at least one peptide having an activity of Der p III) may be administered in a convenient manner such as by injection (subcutaneous, intravenous, etc.), oral administration, inhalation, transdermal application, or rectal administration. Depending on the route of administration, the active compound may be coated in a material to protect the compound from the action of enzymes, acids and other natural conditions which may inactivate the compound.

[0065] To administer a peptide having an activity of Der p III by other than parenteral administration, it may be necessary to coat the peptide with, or co-administer the peptide with, a material to prevent its inactivation. For example, a

peptide having an activity of Der p III may be administered to a subject in an appropriate carrier, diluent or adjuvant, co-administered with enzyme inhibitors or in an appropriate carrier such as liposomes. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Adjuvant is used in its broadest sense and includes any immune stimulating compound such as to interferon. Adjuvants contemplated herein include resorcinols, non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecyl polyethylene ether. Enzyme inhibitors include pancreatic trypsin inhibitor, diisopropylfluorophosphate (DEP) and trasylol. Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes (Strejan et al., (1984) *J. Neuroimmunol.* 7:27). For purposes of inducing T cell non-responsiveness, the composition is preferably administered in non-immunogenic form, e.g., one that does not contain adjuvant.

[0066] The active compound may also be administered parenterally or intraperitoneally. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

[0067] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. In all cases, the composition must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, sodium chloride, or polyalcohols such as mannitol and sorbitol, in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate or gelatin.

[0068] Sterile injectable solutions can be prepared by incorporating active compound (i.e., at least one peptide having an activity of Der p III) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient (i.e., at least one peptide having an activity of Der p III) plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0069] When the peptide having an activity of Der p III is suitably protected, as described above, the peptide may be

orally administered, for example, with an inert diluent or an assimilable edible carrier. The peptide and other ingredients may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the active compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. The percentage of the active ingredient in the compositions and preparations may, of course, be varied and may conveniently be between about 5 to about 80% of the weight of the unit. The amount of active compound in such therapeutically useful compositions is such that a suitable dosage will be obtained.

[0070] As used herein "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions suitable for pharmaceutical administration is contemplated. Supplementary active compounds can also be incorporated into the compositions.

[0071] It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in subjects.

[0072] The invention also provides a composition comprising at least two peptides having an activity of Der p III (e.g., a physical mixture of at least two peptides), each having T cell stimulating activity. Alternatively, a peptide having at least two regions, each having T cell stimulating activity (i.e., each region comprising at least one T cell epitope) can be administered to an allergic subject. A composition of two peptides having a Der p III activity or a composition of two peptides having at least two regions, each having T cell stimulating activity can be administered to a subject in the form of a composition with a pharmaceutically acceptable carrier as hereinbefore described. An amount of one or more of such compositions can be administered simultaneously or sequentially to a subject sensitive to a house dust mite allergen to treat such sensitivity.

[0073] The cDNA (or the mRNA which served as a template during reverse transcription) encoding a peptide having an activity of Der p III can be used to identify similar nucleic acids in any variety or type of animal and, thus, to molecularly clone genes which have sufficient sequence homology to hybridize to the cDNA encoding a peptide having an activity of Der p III. Thus, the present invention includes not only peptides having an activity of Der p III, but also other proteins which may be allergens encoded by DNA which hybridizes to DNA of the present invention.

[0074] Isolated peptides that are immunologically related to Der p III, such as by antibody cross-reactivity or T cell cross-reactivity, other than those already identified, are

within the scope of the invention. Such peptides bind antibodies specific for the protein and peptides of the invention, or stimulate T cells specific for the protein and peptides of this invention.

[0075] A peptide having an activity of Der p III (i.e., Der p III produced recombinantly or by chemical synthesis) is free of all other house dust mite proteins and, thus, is useful in the standardization of allergen extracts which are key reagents for the diagnosis and treatment of house dust mite hypersensitivity. In addition, such a peptide is of a consistent, well-defined composition and biological activity for use in preparations that will be administered for therapeutic purposes (e.g., to modify the allergic response of a subject sensitive to house dust mite). Such peptides can also be used to study the mechanism of immunotherapy of *D. pteronyssinus* allergy and to design modified derivatives or analogs useful in immunotherapy.

[0076] Work by others has shown that high doses of allergen extracts generally produce the best results during immunotherapy (i.e., best symptom relief). However, many subjects are unable to tolerate large doses of such extracts due to systemic reactions elicited by the allergens and other components within these preparations. A peptide having an activity of Der p III has the advantage of being free of all other mite protein. Thus, such a peptide can be administered for therapeutic purposes with fewer anticipated side effects.

[0077] It is now also possible to design an agent or a drug capable of blocking or inhibiting the ability of a house dust mite allergen to induce an allergic reaction in sensitive subjects. Such agents could be designed, for example, in such a manner that they would bind to relevant anti-Der p III IgE molecules, thus preventing IgE-allergen binding, and subsequent mast cell/basophil degranulation. Alternatively, such agents could bind to cellular components of the immune system, resulting in suppression or desensitization of the allergic responses to house dust mite allergens. A non-restrictive example of this is the use of peptides including B or T cell epitopes of Der p III or modifications thereof, based on the cDNA protein structure of Der p III to suppress the allergic response to a house dust mite allergen. This could be carried out by defining the structures of fragments encoding B and T cell epitopes which affect B and T cell function in in vitro studies with blood components from subjects sensitive to house dust mite.

[0078] The invention is further illustrated by the following examples which should not be construed as further limiting the subject invention. The contents of all references and published patent applications cited throughout this application are hereby incorporated by reference.

[0079] The following methodology was used throughout the Examples.

EXAMPLES

Materials and Methods

[0080] *D. pteronyssinus* Cultures

[0081] Whole mites were purchased from the Commonwealth Serum Laboratories, Parkville, Australia and spent medium was a gift from the same source.

Purification of Native Der p III

[0082] Using a method adapted from Heymann et al. (Heymann et al. *J Allergy Clin Immunol* (1989) 83: 1055-1067), 15 ml of a 50-80% saturated ammonium sulfate precipitate of *D.*

pteronyssinus spent growth medium was applied to an upward flowing, 2 cm×90 cm polyacrylamide P-100 column equilibrated in PBS (Pharmacia, LKB Biotechnology, Uppsala, Sweden). The protein content of the eluted 5 ml fractions was determined by measuring optical density (A280 nm) and analysis by SDS-PAGE. Those fractions containing predominantly bands in the 30 kDa region were pooled, concentrated by polyethylene glycol 6000 (BDH Chemicals, Aust. PTY. Ltd. Kilsyth, Vic. Aust), dialyzed against PBS and passed over the column again. This was repeated twice until the analysis by SDS-PAGE determined that the only bands detectable were a doublet of approximately molecular weight 30 kDa.

Protein Sequence Analysis

[0083] The affinity purified native Der p III and Der f III proteins were subjected to HPEC using a 12% column system (Applied Biosystems, Foster City, Calif.). The affinity purified proteins or the HPEC fractions of the appropriate molecular weight were then subjected to protein sequence analysis using Applied Biosystems model 477A gas-phase sequenator with on-line phenylthiohydantoin derivative analysis (model 120). Alternatively, the affinity purified proteins were first separated by SDS-PAGE (BioRad), transferred to Problot (Applied Biosystems) and sequenced using a Beckman model LF3000. After initial sequence analysis of the proteins, o-phthalaldehyde was applied to block N-termini except those with prolines (position 13 for Der p III) to eliminate contaminating peptide sequences and extend the N-terminal sequence unambiguously. The CNBr peptides were produced by cleaving the affinity purified proteins with 2% (w/v) CNBr in 70% (w/v) formic acid overnight at room temperature. The digested peptides were subjected to HPEC for purification and then the peptide fragments were subjected to sequence analysis.

Preferred Codon Usage

[0084] The codon usage bias for the mature proteins Der f I (Dilworth et al. *Clin Exp Allergy* (1991) 21:25-32), Der f II (Trudinger et al. *Clin Exp Allergy* (1991) 21:33-37), Der p I (Chua et al. *J Exp Med* (1988) 167:175-182) and Der p II (Chua et al. *Int Arch Allergy Appl Immunol* (1990) 91:118-123) was determined. The average percent usage of each triplet codon for each amino acid was determined for each of these four *Dermatophagoides* proteins and the results tabulated in Table 1.

TABLE 1

Preferred codon usage table for *Dermatophagoides* allergens.

Amino acid		Codon	%
Ala	A	GCT	45.4
		GCC	33
		GCA	21.6
		GCG	0
Cys	C	TGT	55.2
		TGC	44.8
Asp	D	GAT	92.3
		GAC	7.7
Glu	E	GAA	100
		GAG	0
Phe	F	TTC	82.5
		TTT	17.5

TABLE 1-continued

Preferred codon usage table for Dermatophagoides allergens.			
Amino acid		Codon	%
Gly	G	GGT	65.2
		GGA	24.5
		GGC	10.3
		GGG	0
His	H	CAT	78.8
		CAC	21.2
Ile	I	ATT	50
		ATC	48.5
		ATA	1.5
Lys	K	AAA	99
		AAG	1
Leu	L	TTG	64.5
		TTA	18.5
		CTC	8
		CTT	6.5
		CTG	2.5
		CTA	0
Met	M	ATG	100
Asn	N	AAT	52
		AAC	48
Pro	P	CCA	87.2
		CCC	9.6
		CCG	3.2
		CCT	0
Gln	Q	CAA	94.4
		CAG	5.6
Arg	R	CGA	36.5
		CGT	41.6
		CGC	16.5
		AGA	5.4
		CGG	0
		AGG	0
Ser	S	TCA	42.5
		TCT	18.4
		AGT	11.2
		TCG	11.2
		AGC	10.2
		TCC	6.5
Thr	T	ACT	39.2
		ACA	35
		ACC	18.9
		ACG	6.9
Val	V	GTT	51.2
		GTC	33.8
		GTA	12.3
		GTG	2.7
Trp	W	TGG	100
Tyr	Y	TAT	62
		TAC	38
Stop		TAA	75
		TGA	25
		TAG	0

Construction of the *D. pteronyssinus* λ gt10 cDNA Library

[0085] Polyadenylated mRNA (10 μ g) was used to synthesize cDNA by the RNase H method (Gubler et al. *Gene* (1983) 25:263-269) using a kit (Amersham International, Aylesbury, U.K.). After the addition of EcoRI restriction enzyme linkers (New England Biolabs, Beverly, U.S.A.), the cDNA was ligated to alkaline-phosphatase treated lambda gt10 arms (Promega Biotec, Madison, Wis.). The recombinant phage DNA was packaged and plated out in *E. coli* JP777 to produce a library of 5×10^5 recombinants.

Kinase End-labeling of Oligonucleotides

[0086] Oligonucleotides were synthesized using an Applied Biosystem PCR mate (Applied Biosystems, Foster City, Calif.). Twenty picomole of oligonucleotide DNA was

end labeled with (γ - 32 P) ATP using T4 polynucleotide kinase (Promega Corp., Madison, Wis.) (Maniatis, T., Fritsch, E. F. and Sambrook, J. *Molecular Cloning: A Laboratory Manual*, 2nd Ed. Cold Spring Harbor Laboratory Press, 1989). Labeled oligonucleotide was purified by 15% polyacrylamide gel electrophoresis and subsequently eluted into sterile distilled water.

Isolation of Der p III cDNA Clones from the λ gt10 cDNA Library

[0087] Screening of the library was performed with two probes designed using both N-terminal and internal protein sequence data obtained as described above for the group III allergens. The first, P3forward3 (P3F3), was a 38 mer in length with the following nucleotide sequence, 5'TCA-GAAAAAGCTTTGGCTGGTGAATCACCATATCAAAT 3' (SEQ ID NO:8). The second probe, P3reverse4 (P3R4), a 41 mer with the following nucleotide sequence, 5'GAATCAA-CACCACCATTAGCAACATCACCACCGCAAATCAT3' (SEQ ID NO:9). The library was plated at 25,000 pfu per 150 mm petri dish and the phage were lifted onto nitrocellulose (Schleicher & Schuell, Dassel, Germany). Duplicate filter lifts from each plate were denatured and baked for hybridization with a different one of the two probes (Maniatis, T., Fritsch, E. F. and Sambrook, J. *Molecular Cloning: A Laboratory Manual*, 2nd Ed. Cold Spring Harbor Laboratory Press, 1989). Hybridizations were performed in hybridization mix (6x sodium chloride/sodium citrate pH 7.0 (SSC), 0.1% Denhardt's, 100 μ g/ml denatured herring sperm DNA) at 42° C. with 10^6 cpm/ml labeled probe overnight. Filters were washed three times at 42° C. for 20 min and then at 50° C. for 10 min, in 400 ml wash solution containing 6xSSC, 0.1% Triton X-100.

Isolation of DNA from λ gt10 Der p III Clones

[0088] Phage DNA from the Der p III clones was prepared using a polyethylene glycol precipitation procedure (Chua et al. *J Exp Med* (1988) 167:175-182).

Subcloning and DNA Sequencing

[0089] Purified Der p III phage DNA was digested with EcoRI restriction enzyme (Toyobo) and the released fragment ligated to EcoRI restriction enzyme digested M13mp19 sequencing vector (Messing, *Meth. Enzymol.* (1983) 101: 20-78). The recombinant DNA was transformed using *E. coli* TG-1 and sequencing performed using the dideoxynucleotide chain termination method and Sequenase (U.S. Biochemicals) (Sanger et al., *Proc Natl Acad Sci USA* (1977) 74:5463-5467). The oligonucleotide primers used for sequencing included the universal primers, a 17-mer sequencing primer (-40)5' CAGCACTGACCCTTTTG3' (SEQ ID NO:4) and a 16-mer reverse sequencing primer (-24)5' AACAGCTATGACCATG3' (SEQ ID NO: 5), the two internal primers used for the library screening (SEQ ID NOS: 8 and 9) and two internal primers P3forward5 (P3F5) 5'AAAGCTGTTGGAT-TACCA3' (SEQ ID NO:6) and P3reverse5 (P3R5) 5'TACATCCGATCCTTTTGC3' (SEQ ID NO:7) and P3F4 5'GCGGATCCATTGTTGGTGGT 3' (SEQ ID NO:18). The two internal probes were designed to correspond to nucleotide residues 456-473 (P3F5) and 491-474 (P3R5) (FIG. 1). The primers were used to sequence the isolated clone in both orientations.

DNA and Protein Sequence Analysis

[0090] Sequence analysis was performed using the MAC VECTOR software (IBI, New Haven, U.S.A.). Computation

for sequence homology with other proteins was performed at the NCBI using the BLAST network service. The version of BLAST used was BLAST 1.3.1 OMP (7 Jul. 1993).

Example 1

Sequence of the Native Group III Allergens

[0091] The protein isolation procedure produced a Der p III sample which ran as a doublet with molecular weights 30 and 28 k when analyzed by SDS-PAGE. Both bands reacted with polyclonal mouse anti-Der f III in accord with the interspecies cross-reactivity previously reported (Thomas et al., *Exp. Appl. Acarol.* (1992) 16:153-164). Der f III isolated from a 5A12 monoclonal antibody (Heymann et al. *J. Allergy Clin. Immunol.* (1989) 1055-1067) column exhibited similar characteristics. Using the o-phthalaldehyde to eliminate the contaminating protein sequences, the N-terminal sequencing of both native Der p III and Der f III corrected and extended the known Der p III sequence and extended the Der f III sequence. The Der p III N-terminal sequence was extended to IVGGEKALAGQSPYQISLQSSSHFSGGTIL (SEQ ID NO:16). The Der f III N-terminal sequence was extended to IVGGVKAKAGDSPYQISLQSSSHFXGGSILD (SEQ ID NO:15). Comparison of the sequence data published by Stewart et al. (*Immunology* (1992) 75: 29-35) and Heymann et al., (*J. Allergy Clin. Immunol* (1989) 83: 1055-1067) for the group III allergens with the data herein indicated several errors in the published sequence. In particular, a non-conservative change at residue 8 from a positively charged lysine to a non-polar hydrophobic leucine was found (Der f III, Heymann et al., *J Allergy Clin Immunol* (1989) 83: 1055-1067). Two internal peptides of Der f III were isolated by HPEC after CNBr digestion of the natural protein, MICGGDVANG-GVDSEQGD (SEQ ID NO:10) and MTLDQTNAKPV-PLPTS (SEQ ID NO:12).

Example 2

Sequence of Recombinant Der p III

[0092] The protein sequence information obtained as described in Example 1 was used in conjunction with the preferred codon usage data (Table 1) to construct two oligonucleotides probes. These oligonucleotides were designed to hybridize to nucleotide residues 159-196 and to residues 688-648 of the Der p III clone (FIG. 2). Only clones which hybridized strongly with both probes were isolated from the λ gt10 cDNA library. The resulting nucleotide sequence for the P3WS1 clone and the deduced amino acid sequence is shown in FIG. 1. The complete nucleotide sequence was 1059 bp in length. This includes a 5' non-coding region of 62 bp, a 211 bp 3' untranslated region and an open reading frame of 786 bp with a stop codon (TGA) at nucleotide residues 846-848. There is no poly A tail but there does appear to be a polyadenylation signal (AATAAA). The open reading frame encodes a protein which includes a 29 amino acid pre-pro region and starting at the N-terminal isoleucine, a mature protein of 232 amino acid residues with a calculated molecular weight of 24,985 and pI of 8.5. The methionine residue (ATG) at amino acid position -29 is the most likely translation initiation site. Selection for this initiation site was based in part on the presence of a sequence following the methionine which encodes a classical signal peptide of 18 amino acid residues with predominantly hydrophobic residues present. In addition, the immediate sequence prior to this ATG codon,

5'GAAAGATG3', conforms loosely to the Kozak consensus sequence (CCACCATG) for the eucaryotic translation initiation site with the crucial purine (most often A) at the -3 position (Kozak, *Nucleic Acid Research* (1984) 12:857-872).

[0093] The protein sequencing data of Der p III described in Example 1 differs from the amino acid sequence deduced from the cDNA of the P3WS1 clone as a result of the substitutions of Glu/Gln at residue 11 and a possible Cys/Ser substitution at residue 17. These substitutions may be due to the existence of Der p III in different isoforms. Analysis of the deduced protein sequence for Der p III P3WS1 confirmed that the position of the Der f III CNBr peptide 1 (MI(C)GGDVANGGVDS(E)QGD) (SEQ ID NO:10) was correctly predicted to be from amino acid residues 177-183 and has 88% identity with the recombinant sequence in this region, MICGGDVANGGKDSCQD (SEQ ID NO:11). Interestingly, there are two identical non-conservative changes that exist when the native Der f III peptide sequence is aligned with the P3WS1 clone sequence. There is a change from a non-polar hydrophobic valine at 178 (according to numbering in FIG. 1) in Der f III to a positively charged lysine residue in the P3WS1 clone sequences. The other change is a glutamic acid residue at 181 in the Der f III substituted for a cysteine residue in the P3WS1 clone sequence. The cysteine forms one of the disulfide bridges in both these and most other trypsin proteins. The predicted alignment of the second Der f III peptide, (M)TL DQTNA(K)PVPL(P)(T)(S) (SEQ ID NO:12), produced by the CNBr digestion was from amino acid residues 95-110 (MKLNQKNKAVGLPAK) (SEQ ID NO:13). There is 63% homology with the P3WS1 clone in this region. These differences between the Der p III and Der f III protein sequences are consistent with the 20% sequence variation found between other homologous allergens from this species such as the Der p I and Der f I; and Der p II and Der f II.

[0094] The complete sequence for the Der p III P3WS1 clone encodes a protein of 319 amino acids. Der p III, like all known trypsins, is synthesized as a pre-pro zymogen. The cleavage site of the signal peptide is postulated to be between amino acid residues -12 and -11 as the residues around this site conform to the amino acid constraints outlined by von Heijne, (von Heijne, *Eur J Biochem* (1983) 133:17-21). Von Heijne proposed that small neutral amino acids are strongly preferred at positions -1 and -3 from the cleavage site, such as the alanine and tyrosine seen in the Der p III gene. Proline residues are never found in positions +1 to -3 thus excluding any other sites. Most mammalian trypsin proteins have been reported to have pre or signal peptides of 15 or 16 amino acids. The Der p III peptide is 18 amino acids in length. While the difference in length is relatively small, it does seem that the length of the signal peptide may be attributable to the phylogenetic diversity of the species from which the trypsin originated. This is most apparent in *Streptomyces griseus* with a 32 amino acid pre peptide (Kim et al., *Biochem Biophys Res Commun* (1991) 181:707-713). It has also been reported that for mammalian trypsins the signal peptide contains two specific clusters of two and then four hydrophobic residues (MacDonald et al., *J Biol Chem* (1982) 257:9724-9732; Le Huerou et al., *Eur J Biochem* (1990) 193:767-73). There is an abundance of hydrophobic residues within this region of the Der p III gene but they are not arranged into highly conserved clusters. The same is true of the signal peptides for both *Drosophila melangastor* (Davis et al., *Nucleic Acid Research* (1988) 13:6605-6615) and *S. griseus* (Kim et al., *Biochem Biophys Res Commun* (1991) 181:707-

713). It is possible therefore that the highly conserved clusters of hydrophobic residues are not a characteristic of all trypsin proteins but rather that of mammalian trypsins.

[0095] Downstream from the signal peptide is the 11 residue pre-activation peptide from amino acid residues -11 to -1. Comparison of this pro region with other pro regions from trypsins of various species indicates the Der p III pro peptide is unusual. Both vertebrate and invertebrate trypsins have been shown to have an octapeptide and or a hexapeptide pro region with four contiguous aspartyl residues followed by a lysine residue at the -1 position (Le Huerou et al., *Eur J Biochem* (1990) 193:767-73). It is the carboxyl region of this lysine residue which is cleaved during the activation of the trypsinogen to trypsin. The Der p III lacks this polyaspartyl-lysine sequence. The trypsinogen therefore can not be activated by the usual mechanism. There are a few other trypsins which are similar to Der p III in this respect, *D. melangastor* (Davis et al., *Nucleic Acid Research* (1988) 13:6605-6615), *S. griseus* (Kim et al., *Biochem Biophys Res Commun* (1991) 181:707-713) and most interestingly, a 32 kDa anionic trypsin from the rat pancreas (Gendry and Launay, *Biochemica et Biophysica Acta* (1988) 955:243-249). This group showed that enterokinase, an enzyme which has a highly specific affinity for the polyaspartyl-lysine residues in trypsin proteins, had no effect on the trypsin that they isolated. They suggested therefore that a different mechanism was being used for the activation process. The absence of any homology of the Der p III proenzyme region with any of the known sequences may imply a unique activation mechanism.

[0096] A search of the protein database has confirmed that Der p III P3WS1 is homologous to both vertebrate and invertebrate trypsins. The Der p III allergen is not a mite chymotrypsin (Gendry and Launay, *Biochemica et Biophysica Acta* (1988) 955:243-249) as comparison of the N-terminal protein sequences showed that they differ by as much as 50%. FIG. 2 shows a comparison of the amino acid sequences of Der p III P3WS1 and crayfish trypsins (SEQ ID NOS: 2 and 3). The crayfish has 44% homology with Der p III. There are seven cysteine residues, six of these residues are known to form disulphide bridges, 54-70, 181-198 and 210-236 (FIG. 2.) (Hartley, *Phil Trans Roy Soc Lond B* (1970) 257:77-87). Most importantly, the Der p III protein contains the highly conserved residues involved in the catalytic activity and substrate specificity of the trypsin proteins. The conserved His 40 and Ser 185 (FIG. 1) represent the, charge relay system which comprise the active catalytic site of the trypsin enzyme. The positively charged aspartic acid residue at 179, represents the trypsin specificity site. Most serine proteases have a neutral amino acid at this residue (Keil, *The Enzymes*. Academic Press, New York, 1971, pages 249-279). The glycine residues at 206 and 211 in combination with the aspartic acid at 179 are responsible for accommodating the bulky positively charged side chains of lysine and arginine, the substrate residues cleaved by trypsins. Amino acid residues 179-185 and 202-206 (FIG. 1) form the S1 binding pockets which are involved in the binding of the substrate to the trypsin molecule. The areas immediately surrounding all of these important residues are the most conserved regions of the Der p III protein. All these results are consistent with the fact that among trypsin proteins from different species structural variations occur in those regions of the molecule which are not important for catalytic activity (Vithayathil et al., *Arch Biochem Biophys* (1961) 92:532-540).

[0097] Most trypsin proteins generally have a common pairing of cysteine residues to form six disulphide bridges, with two of these bridges, Cys 15-145 and Cys 117-218 being unique to this group of proteins. Der p III like both crayfish (Titani et al., *Biochemistry* (1983) 22:1459-1465) and *S. griseus* are missing these two unique links, but more importantly Der p III is unique as it contains an extra unpaired cysteine residue. This residue may be the equivalent to Cys 15 in the bovine trypsin which unlike, Cys 145, 117 and 218 was not eliminated during evolutionary divergence of the mite trypsin.

[0098] The isolation of the gene encoding the nucleotide sequence for the group III allergen, Der p III is the first primary sequence determination for this group of *Dermatophagoides allergens*. Comparison of the sequence with other vertebrate and invertebrate serine proteases in view of substrate binding experiments (Stewart et al., *Immunology* (1992) 75: 29-35; Ando et al., *Alerugi* (1992) 41:704-707), indicates that the group III allergens are trypsin proteins. Trypsin proteins are secreted as pre-pro zymogens by the pancreatic acinar cells of all vertebrate and invertebrate species. Invertebrate trypsins have been reported to have a molecular weight ranging from 20 to 30 kDa (Graf et al., *Insect Biochem* (1985) 15:611-618). The Der p III P3WS1 trypsinogen has a calculated molecular weight of 28 kDa and the corresponding mature protein a molecular weight of 24,985 kDa while the native protein purified, existed as duplicate of 28 and 30 kDa as estimated by SDS-PAGE. There have been other reports of multiple protein bands being isolated for both Der f III (Thomas et al., *Exp Appl Acarol* (1992) 16:153-164) and Der p III (Stewart et al., *Immunology* (1992) 75: 29-35). These results are not unusual as trypsin proteins are known to exist in several isoforms. Gendry and Launay (1988) isolated duplicate protein bands of an anionic trypsin-like protein from rat pancreas. They demonstrated that the higher 32 kDa band represented the inactive trypsin and the lower 30 kDa band represented the active form of the trypsin which could autocatalyze the cleavage of the 32 kDa band. It is therefore possible that the 30 kDa band represents the inactive trypsinogen and the 28 kDa band the active trypsin. The number of isozymes for invertebrate trypsins has been reported to range from 1-12 (MacDonald et al., *J Biol Chem* (1982) 257:9724-9732). Stewart et al. (1992) proposed nine major isoforms of Der p III existed, with pI's ranging from 4 to >8.

Example 3

Expression and Purification of Recombinant Der p III

[0099] A complementary DNA insert encoding Der p III (FIG. 1) was digested with EcoRI and inserted into pUC 19. To remove the non-translated 5' sequence as well as part of the putative hydrophobic leader, the cDNA was cleaved with MscI and an EcoRI linker was added. Subsequently, the truncated Der p III fragment was EcoRI digested and subcloned into the expression vector pET11d (Studier et al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 60-89). The sequence and reading frame were verified by DNA sequencing. The Der p III coding sequence encompassed residue 13 to the stop codon (see FIG. 1). The expression vector pET11d Der III was transformed into the *E. coli* host strain BL21(DE3) and selected on plates containing 150 µg/ml ampicillin (Studier et

al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 60-89). A single transformant colony was grown up in 2 ml volume of BHIB medium containing 150 µg/ml ampicillin at 37° C., for approximately 6 hr. Ten milliliters of this culture was spread onto a selection plate and grown overnight at 37° C. The bacterial lawn was recovered in 2 ml media and added to 500 ml of BHIB medium (150 µg/ml ampicillin) and grown at 37° C. to A₆₀₀=1.0. Recombinant expression was induced by the addition of IPTG to a final concentration of 1 mM. After 2 hr growth, cells were harvested, lysed and the proteins solubilized in 6 M guanidine HCl buffer containing 100 mM 2-ME as previously described for the ragweed recombinant allergens, Amb a I.1 and Amb a II (Rogers, et al. (1991) *J. Immunol.* 147:2547-2552).

[0100] The guanidine HCl lysate containing Der p III was subjected to Ni²⁺ metal-ion affinity chromatography under denaturing conditions in 8 M urea (Hochuli, E. et al., (1988) *Bio/Technology* 6:1321-1325). After elution from the Ni²⁺ chelating support, QIAGEN NTA-Agarose (Diagen GmbH, Dusseldorf, Germany), the recombinant Der p III protein preparations were subjected to SDS-PAGE analysis (FIG. 3).

[0101] Recombinant Der p III had an expected molecular weight of approximately 27 kDa as predicted for the 252

amino acids plus amino acids used as a purification tag (MGHHHHHHEF (SEQ ID NO:17)). The pET11d expression system and the Ni²⁺ methyl-ion-affinity chromatography method generated approximately 12 mg recombinant Der p III per liter of growth medium (as assessed by A₂₈₀ measurement). This expression and purification scheme yielded a protein preparation with purity exceeding 90% as assessed by SDS-PAGE visualized by Coomassie Blue staining. (FIG. 3). The observed apparent molecular weight of approximately 32 kDa is slightly higher than that predicted from the primary structure (FIG. 1, and above). In FIG. 3, increasing concentrations of Der p III preparation were examined to verify the purity (lane 1, 4.3 µg; lane 2, 8.7 µg; lane 3, 13.8 µg; lane 4, 17.4 µg; markers are indicated by M).

EQUIVALENTS

[0102] Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific embodiments described herewith. Such equivalents are considered to be within the scope of this invention and are encompassed by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 18

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1059 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 63..848

(ix) FEATURE:

- (A) NAME/KEY: mat_peptide
- (B) LOCATION: 150..848

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CTTTTTTTTT TTTTGTGAAA ACATTGAAAA ACATTGAAGC AACTGTGTGT ACAACCAGAA	60
AC ATG ATC ATC TAT AAT ATT TTA ATT GTT TTA TTA TTG GCC ATT AAT	107
Met Ile Ile Tyr Asn Ile Leu Ile Val Leu Leu Leu Ala Ile Asn	
-29 -25 -20 -15	
ACA TTG GCT AAT CCA ATT CTA CCA GCA TCA CCA AAT GCA ACT ATT GTT	155
Thr Leu Ala Asn Pro Ile Leu Pro Ala Ser Pro Asn Ala Thr Ile Val	
-10 -5 1	
GGT GGT GAA AAA GCA TTA GCT GGT GAA TGT CCA TAT CAG ATT TCA TTA	203
Gly Gly Glu Lys Ala Leu Ala Gly Glu Cys Pro Tyr Gln Ile Ser Leu	
5 10 15	
CAA TCA AGT AGT CAT TTT TGT GGT GGT ACT ATT CTT GAT GAA TAT TGG	251
Gln Ser Ser Ser His Phe Cys Gly Gly Thr Ile Leu Asp Glu Tyr Trp	
20 25 30	

-continued

ATT TTA ACA GCT GCA CAT TGT GTT GCC GGA CAA ACA GCA AGT AAA CTT	299
Ile Leu Thr Ala Ala His Cys Val Ala Gly Gln Thr Ala Ser Lys Leu	
35 40 45 50	
TCA ATT CGT TAC AAT AGT TTA AAA CAT TCA TTA GGT GGT GAA AAA ATT	347
Ser Ile Arg Tyr Asn Ser Leu Lys His Ser Leu Gly Gly Glu Lys Ile	
55 60 65	
TCT GTT GCT AAA ATT TTT GCA CAT GAA AAA TAT GAT AGT TAT CAA ATT	395
Ser Val Ala Lys Ile Phe Ala His Glu Lys Tyr Asp Ser Tyr Gln Ile	
70 75 80	
GAT AAT GAT ATT GCA TTG ATT AAG CTT AAA TCA CCT ATG AAA TTA AAT	443
Asp Asn Asp Ile Ala Leu Ile Lys Leu Lys Ser Pro Met Lys Leu Asn	
85 90 95	
CAG AAA AAT GCC AAA GCT GTT GGA TTA CCA GCA AAA GGA TCG GAT GTA	491
Gln Lys Asn Ala Lys Ala Val Gly Leu Pro Ala Lys Gly Ser Asp Val	
100 105 110	
AAA GTT GGT GAT CAA GTT CGT GTT TCT GGT TGG GGT TAT CTT GAA GAA	539
Lys Val Gly Asp Gln Val Arg Val Ser Gly Trp Gly Tyr Leu Glu Glu	
115 120 125 130	
GGA AGT TAT TCA TTA CCA TCT GAA TTA AGA CGT GTT GAT ATT GCT GTT	587
Gly Ser Tyr Ser Leu Pro Ser Glu Leu Arg Arg Val Asp Ile Ala Val	
135 140 145	
GTA TCA CGT AAA GAA TGT AAT GAA TTA TAT TCA AAA GCT AAT GCT GAA	635
Val Ser Arg Lys Glu Cys Asn Glu Leu Tyr Ser Lys Ala Asn Ala Glu	
150 155 160	
GTT ACT GAT AAT ATG ATT TGT GGT GGT GAT GTT GCA AAT GGT GGT AAA	683
Val Thr Asp Asn Met Ile Cys Gly Gly Asp Val Ala Asn Gly Gly Lys	
165 170 175	
GAT TCT TGT CAA GGT GAT TCT GGT GGA CCG GTT GTT GAT GTT AAA AAT	731
Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Val Val Asp Val Lys Asn	
180 185 190	
AAT CAA GTT GTT GGT ATT GTT TCA TGG GGT TAT GGT TGT GCA CGT AAA	779
Asn Gln Val Val Gly Ile Val Ser Trp Gly Tyr Gly Cys Ala Arg Lys	
195 200 205 210	
GGT TAT CCA GGT GTT TAT ACA CGT GTT GGT AAT TTT ATC GAT TGG ATT	827
Gly Tyr Pro Gly Val Tyr Thr Arg Val Gly Asn Phe Ile Asp Trp Ile	
215 220 225	
GAA TCA AAA CGT TCA CAG TGATTGATAA AAACAATTTC GACATATAAA	875
Glu Ser Lys Arg Ser Gln	
230	
AATTGGCAAA TGATGCCATT ATATGTTATC ATTATGTTTC TGGTGATTCT TTTTCTACTT	935
TTATCATTTTT ATTTTATATAT CGAGCTGAAA ACAAACAATC ATGATTGTAT AGGGATTTGT	995
TGTTGTTTTT CAATTAAAA AAATCCAAAA TAAATAAATA AATAATCATT TCAACAAAAA	1055
AAAG	1059

(2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 261 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Ile Ile Tyr Asn Ile Leu Ile Val Leu Leu Leu Ala Ile Asn Thr
-29 -25 -20 -15

Leu Ala Asn Pro Ile Leu Pro Ala Ser Pro Asn Ala Thr Ile Val Gly

-continued

-10	-5	1
Gly Glu Lys Ala Leu Ala Gly Glu Cys Pro Tyr Gln Ile Ser Leu Gln 5 10 15		
Ser Ser Ser His Phe Cys Gly Gly Thr Ile Leu Asp Glu Tyr Trp Ile 20 25 30 35		
Leu Thr Ala Ala His Cys Val Ala Gly Gln Thr Ala Ser Lys Leu Ser 40 45 50		
Ile Arg Tyr Asn Ser Leu Lys His Ser Leu Gly Gly Glu Lys Ile Ser 55 60 65		
Val Ala Lys Ile Phe Ala His Glu Lys Tyr Asp Ser Tyr Gln Ile Asp 70 75 80		
Asn Asp Ile Ala Leu Ile Lys Leu Lys Ser Pro Met Lys Leu Asn Gln 85 90 95		
Lys Asn Ala Lys Ala Val Gly Leu Pro Ala Lys Gly Ser Asp Val Lys 100 105 110 115		
Val Gly Asp Gln Val Arg Val Ser Gly Trp Gly Tyr Leu Glu Glu Gly 120 125 130		
Ser Tyr Ser Leu Pro Ser Glu Leu Arg Arg Val Asp Ile Ala Val Val 135 140 145		
Ser Arg Lys Glu Cys Asn Glu Leu Tyr Ser Lys Ala Asn Ala Glu Val 150 155 160		
Thr Asp Asn Met Ile Cys Gly Gly Asp Val Ala Asn Gly Gly Lys Asp 165 170 175		
Ser Cys Gln Gly Asp Ser Gly Gly Pro Val Val Asp Val Lys Asn Asn 180 185 190 195		
Gln Val Val Gly Ile Val Ser Trp Gly Tyr Gly Cys Ala Arg Lys Gly 200 205 210		
Tyr Pro Gly Val Tyr Thr Arg Val Gly Asn Phe Ile Asp Trp Ile Glu 215 220 225		
Ser Lys Arg Ser Gln 230		

(2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 237 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Ile Val Gly Gly Thr Asp Ala Val Leu Gly Glu Phe Pro Tyr Gln Leu 1 5 10 15
Ser Phe Gln Glu Thr Phe Leu Gly Phe Ser Phe His Phe Cys Gly Ala 20 25 30
Ser Ile Tyr Asn Glu Asn Tyr Ala Ile Thr Ala Gly His Cys Val Tyr 35 40 45
Gly Asp Asp Tyr Glu Asn Pro Ser Gly Leu Gln Ile Val Ala Gly Glu 50 55 60
Leu Asp Met Ser Val Asn Glu Gly Ser Glu Gln Thr Ile Thr Val Ser 65 70 75 80

-continued

Lys	Ile	Ile	Leu	His	Glu	Asn	Phe	Asp	Tyr	Asp	Leu	Leu	Asp	Asn	Asp
				85					90					95	
Ile	Ser	Leu	Leu	Lys	Leu	Ser	Gly	Ser	Leu	Thr	Phe	Asn	Asn	Asn	Val
			100					105					110		
Ala	Pro	Ile	Ala	Leu	Pro	Ala	Gln	Gly	His	Thr	Ala	Thr	Gly	Asn	Val
			115				120					125			
Ile	Val	Thr	Gly	Trp	Gly	Thr	Thr	Ser	Glu	Gly	Gly	Asn	Thr	Pro	Asp
		130				135					140				
Val	Leu	Gln	Lys	Val	Thr	Val	Pro	Leu	Val	Ser	Asp	Ala	Glu	Cys	Arg
145					150					155				160	
Asp	Asp	Tyr	Gly	Ala	Asp	Glu	Ile	Phe	Asp	Ser	Met	Ile	Cys	Ala	Gly
				165					170					175	
Val	Pro	Glu	Gly	Gly	Lys	Asp	Ser	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro
			180					185					190		
Leu	Ala	Ala	Ser	Asp	Thr	Gly	Ser	Thr	Tyr	Leu	Ala	Gly	Ile	Val	Ser
		195					200					205			
Trp	Gly	Tyr	Gly	Cys	Ala	Arg	Pro	Gly	Tyr	Pro	Gly	Val	Tyr	Thr	Glu
	210					215					220				
Val	Ser	Tyr	His	Val	Asp	Trp	Ile	Lys	Ala	Asn	Ala	Val			
225					230					235					

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CAGCACTGAC CCTTTTG

17

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

AACAGCTATG ACCATG

16

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

AAAGCTGTTG GATTACCA

18

-continued

(2) INFORMATION FOR SEQ ID NO: 7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

TACATCCGAT CCTTTTGC

18

(2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 38 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

TCAGAAAAAG CTTTGGCTGG TGAATCACCA TATCAAAT

38

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 41 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

GAATCAACAC CACCATTAGC AACATCACCA CCGCAAATCA T

41

(2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Met	Ile	Cys	Gly	Gly	Asp	Val	Ala	Asn	Gly	Gly	Val	Asp	Ser	Glu	Gln
1			5					10				15			

Gly Asp

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

-continued

Met	Ile	Cys	Gly	Gly	Asp	Val	Ala	Asn	Gly	Gly	Lys	Asp	Ser	Cys	Gln
1			5					10					15		

Asp

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Met	Thr	Leu	Asp	Gln	Thr	Asn	Ala	Lys	Pro	Val	Pro	Leu	Pro	Thr	Ser
1			5			10						15			

(2) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Met	Lys	Leu	Asn	Gln	Lys	Asn	Ala	Lys	Ala	Val	Gly	Leu	Pro	Ala	Lys
1			5			10						15			

(2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Asn	Asn	Gln	Val	Gly	Ile
1		5			

(2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: N-terminal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

Ile	Val	Gly	Gly	Val	Lys	Ala	Lys	Ala	Gly	Asp	Ser	Pro	Tyr	Gln	Ile
1			5				10					15			

Ser	Leu	Gln	Ser	Ser	Ser	His	Phe	Xaa	Gly	Gly	Ser	Ile	Leu	Asp
	20						25					30		

-continued

(2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: N-terminal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

```

Ile Val Gly Gly Glu Lys Ala Leu Ala Gly Gln Ser Pro Tyr Gln Ile
1           5           10           15

Ser Leu Gln Ser Ser Ser His Phe Ser Gly Gly Thr Ile Leu
          20           25           30

```

(2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

```

Met Gly His His His His His His Glu Phe
1           5           10

```

(2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

GCGGATCCAT TGTTGGTGGT

20

1-47. (canceled)

48. A method of treating sensitivity to a house dust mite allergen in a mammal sensitive to such allergen, comprising administering to said mammal a therapeutically effective amount of an isolated peptide of a Der p III protein allergen consisting of a portion of the amino acid sequence of SEQ ID NO: 2 at least 30 amino acid residues in length, wherein the peptide is free of all other house dust mite proteins and has at least one of the following features: is capable of stimulating T or B cells specific for a Der p III protein allergen; or includes at least one B cell epitope recognized by a B cell receptor specific for a Der p III protein allergen and wherein the peptide does not comprise the amino acid sequence -29 to 232 or 1-232 of SEQ ID NO: 2.

49. A method of detecting sensitivity to a house dust mite allergen in a mammal, comprising administering to said mammal an isolated peptide of a Der p III protein allergen and

detecting an increase in the IgE mediated immune response to said peptide in said mammal, wherein the isolated peptide of a Der p III protein allergen consists of a portion of the amino acid sequence of SEQ ID NO: 2 at least 30 amino acid residues in length, wherein the peptide is free of all other house dust mite proteins and has at least one of the following features: is capable of stimulating T or B cells specific for a Der p III protein allergen; or includes at least one B cell epitope recognized by a B cell receptor specific for a Der p III protein allergen and wherein the peptide does not comprise the amino acid sequence -29 to 232 or 1-232 of SEQ ID NO: 2.

50. A method of detecting sensitivity in a subject to a house dust mite allergen, comprising combining a blood component obtained from the subject with an isolated peptide of a Der p III protein allergen, under conditions appropriate for binding of the blood component to the peptide and determining the extent to which such binding occurs, wherein the isolated

peptide of a Der p III protein allergen consists of a portion of the amino acid sequence of SEQ ID NO: 2 at least 30 amino acid residues in length, wherein the peptide is free of all other house dust mite proteins and has at least one of the following features: is capable of stimulating T or B cells specific for a Der p III protein allergen; or includes at least one B cell epitope recognized by a B cell receptor specific for a Der p III protein allergen and wherein the peptide does not comprise the amino acid sequence-29 to 232 or 1-232 of SEQ ID NO: 2.

51. A method of producing an isolated peptide of a Der p III protein allergen, comprising expressing the peptide from a

recombinant nucleic acid encoding the peptide, wherein the isolated peptide of a Der p III protein allergen consists of a portion of the amino acid sequence of SEQ ID NO: 2 at least 30 amino acid residues in length, wherein the peptide is free of all other house dust mite proteins and has at least one of the following features: is capable of stimulating T or B cells specific for a Der p III protein allergen; or includes at least one B cell epitope recognized by a B cell receptor specific for a Der p III protein allergen and wherein the peptide does not comprise the amino acid sequence-29 to 232 or 1-232 of SEQ ID NO: 2.

* * * * *

专利名称(译)	编码屋尘螨过敏原的核酸，der p iii，并用于此		
公开(公告)号	US20110117116A1	公开(公告)日	2011-05-19
申请号	US12/688744	申请日	2010-01-15
申请(专利权)人(译)	MERCK PATENT GMBH		
当前申请(专利权)人(译)	MERCK PATENT GMBH		
[标]发明人	THOMAS WAYNE R CHUA KAW YAN ROGERS BRUCE L KUO MEI CHANG		
发明人	THOMAS, WAYNE R. CHUA, KAW-YAN ROGERS, BRUCE L. KUO, MEI-CHANG		
IPC分类号	A61K39/35 C12P21/02 G01N33/566 A61P37/08 A61P37/04 G01N33/53 A61K38/00 A61K39/395 C07H21/04 C07K14/435 C07K16/18 C12N15/09 C12P21/08		
CPC分类号	C07K14/43531 A61K38/00 A61P37/04 A61P37/08		
其他公开文献	US8287875		
外部链接	Espacenet USPTO		

摘要(译)

公开了编码屋尘螨 (Dermatophagoides pteronyssinus) 过敏原的分离的核酸，Der p III。还描述了编码具有Der p III活性和预测分子量为约24,985道尔顿的肽的cDNA。核酸可用作探针以检测样品中Der p III核酸的存在或用于重组产生具有Der p III活性的肽。具有Der p III活性的肽可用于适于药物施用的组合物或诊断室内尘螨敏感性的方法。

```

CT      2
152
122
182
242
302
362
422
482
542
602
662
722
782
842
902
962
1022
1082
1142
1202
1262
1322
1382
1442
1502
1562
1622
1682
1742
1802
1862
1922
1982
2042
2102
2162
2222
2282
2342
2402
2462
2522
2582
2642
2702
2762
2822
2882
2942
3002
3062
3122
3182
3242
3302
3362
3422
3482
3542
3602
3662
3722
3782
3842
3902
3962
4022
4082
4142
4202
4262
4322
4382
4442
4502
4562
4622
4682
4742
4802
4862
4922
4982
5042
5102
5162
5222
5282
5342
5402
5462
5522
5582
5642
5702
5762
5822
5882
5942
6002
6062
6122
6182
6242
6302
6362
6422
6482
6542
6602
6662
6722
6782
6842
6902
6962
7022
7082
7142
7202
7262
7322
7382
7442
7502
7562
7622
7682
7742
7802
7862
7922
7982
8042
8102
8162
8222
8282
8342
8402
8462
8522
8582
8642
8702
8762
8822
8882
8942
9002
9062
9122
9182
9242
9302
9362
9422
9482
9542
9602
9662
9722
9782
9842
9902
9962
10022
10082
10142
10202
10262
10322
10382
10442
10502
10562
10622
10682
10742
10802
10862
10922
10982
11042
11102
11162
11222
11282
11342
11402
11462
11522
11582
11642
11702
11762
11822
11882
11942
12002
12062
12122
12182
12242
12302
12362
12422
12482
12542
12602
12662
12722
12782
12842
12902
12962
13022
13082
13142
13202
13262
13322
13382
13442
13502
13562
13622
13682
13742
13802
13862
13922
13982
14042
14102
14162
14222
14282
14342
14402
14462
14522
14582
14642
14702
14762
14822
14882
14942
15002
15062
15122
15182
15242
15302
15362
15422
15482
15542
15602
15662
15722
15782
15842
15902
15962
16022
16082
16142
16202
16262
16322
16382
16442
16502
16562
16622
16682
16742
16802
16862
16922
16982
17042
17102
17162
17222
17282
17342
17402
17462
17522
17582
17642
17702
17762
17822
17882
17942
18002
18062
18122
18182
18242
18302
18362
18422
18482
18542
18602
18662
18722
18782
18842
18902
18962
19022
19082
19142
19202
19262
19322
19382
19442
19502
19562
19622
19682
19742
19802
19862
19922
19982
20042
20102
20162
20222
20282
20342
20402
20462
20522
20582
20642
20702
20762
20822
20882
20942
21002
21062
21122
21182
21242
21302
21362
21422
21482
21542
21602
21662
21722
21782
21842
21902
21962
22022
22082
22142
22202
22262
22322
22382
22442
22502
22562
22622
22682
22742
22802
22862
22922
22982
23042
23102
23162
23222
23282
23342
23402
23462
23522
23582
23642
23702
23762
23822
23882
23942
24002
24062
24122
24182
24242
24302
24362
24422
24482
24542
24602
24662
24722
24782
24842
24902
24962
25022
25082
25142
25202
25262
25322
25382
25442
25502
25562
25622
25682
25742
25802
25862
25922
25982
26042
26102
26162
26222
26282
26342
26402
26462
26522
26582
26642
26702
26762
26822
26882
26942
27002
27062
27122
27182
27242
27302
27362
27422
27482
27542
27602
27662
27722
27782
27842
27902
27962
28022
28082
28142
28202
28262
28322
28382
28442
28502
28562
28622
28682
28742
28802
28862
28922
28982
29042
29102
29162
29222
29282
29342
29402
29462
29522
29582
29642
29702
29762
29822
29882
29942
30002
30062
30122
30182
30242
30302
30362
30422
30482
30542
30602
30662
30722
30782
30842
30902
30962
31022
31082
31142
31202
31262
31322
31382
31442
31502
31562
31622
31682
31742
31802
31862
31922
31982
32042
32102
32162
32222
32282
32342
32402
32462
32522
32582
32642
32702
32762
32822
32882
32942
33002
33062
33122
33182
33242
33302
33362
33422
33482
33542
33602
33662
33722
33782
33842
33902
33962
34022
34082
34142
34202
34262
34322
34382
34442
34502
34562
34622
34682
34742
34802
34862
34922
34982
35042
35102
35162
35222
35282
35342
35402
35462
35522
35582
35642
35702
35762
35822
35882
35942
36002
36062
36122
36182
36242
36302
36362
36422
36482
36542
36602
36662
36722
36782
36842
36902
36962
37022
37082
37142
37202
37262
37322
37382
37442
37502
37562
37622
37682
37742
37802
37862
37922
37982
38042
38102
38162
38222
38282
38342
38402
38462
38522
38582
38642
38702
38762
38822
38882
38942
39002
39062
39122
39182
39242
39302
39362
39422
39482
39542
39602
39662
39722
39782
39842
39902
39962
40022
40082
40142
40202
40262
40322
40382
40442
40502
40562
40622
40682
40742
40802
40862
40922
40982
41042
41102
41162
41222
41282
41342
41402
41462
41522
41582
41642
41702
41762
41822
41882
41942
42002
42062
42122
42182
42242
42302
42362
42422
42482
42542
42602
42662
42722
42782
42842
42902
42962
43022
43082
43142
43202
43262
43322
43382
43442
43502
43562
43622
43682
43742
43802
43862
43922
43982
44042
44102
44162
44222
44282
44342
44402
44462
44522
44582
44642
44702
44762
44822
44882
44942
45002
45062
45122
45182
45242
45302
45362
45422
45482
45542
45602
45662
45722
45782
45842
45902
45962
46022
46082
46142
46202
46262
46322
46382
46442
46502
46562
46622
46682
46742
46802
46862
46922
46982
47042
47102
47162
47222
47282
47342
47402
47462
47522
47582
47642
47702
47762
47822
47882
47942
48002
48062
48122
48182
48242
48302
48362
48422
48482
48542
48602
48662
48722
48782
48842
48902
48962
49022
49082
49142
49202
49262
49322
49382
49442
49502
49562
49622
49682
49742
49802
49862
49922
49982
50042
50102
50162
50222
50282
50342
50402
50462
50522
50582
50642
50702
50762
50822
50882
50942
51002
51062
51122
51182
51242
51302
51362
51422
51482
51542
51602
51662
51722
51782
51842
51902
51962
52022
52082
52142
52202
52262
52322
52382
52442
52502
52562
52622
52682
52742
52802
52862
52922
52982
53042
53102
53162
53222
53282
53342
53402
53462
53522
53582
53642
53702
53762
53822
53882
53942
54002
54062
54122
54182
54242
54302
54362
54422
54482
54542
54602
54662
54722
54782
54842
54902
54962
55022
55082
55142
55202
55262
55322
55382
55442
55502
55562
55622
55682
55742
55802
55862
55922
55982
56042
56102
56162
56222
56282
56342
56402
56462
56522
56582
56642
56702
56762
56822
56882
56942
57002
57062
57122
57182
57242
57302
57362
57422
57482
57542
57602
57662
57722
57782
57842
57902
57962
58022
58082
58142
58202
58262
58322
58382
58442
58502
58562
58622
58682
58742
58802
58862
58922
58982
59042
59102
59162
59222
59282
59342
59402
59462
59522
59582
59642
59702
59762
59822
59882
59942
60002
60062
60122
60182
60242
60302
60362
60422
60482
60542
60602
60662
60722
60782
60842
60902
60962
61022
61082
61142
61202
61262
61322
61382
61442
61502
61562
61622
61682
61742
61802
61862
61922
61982
62042
62102
62162
62222
62282
62342
62402
62462
62522
62582
62642
62702
62762
62822
62882
62942
63002
63062
63122
63182
63242
63302
63362
63422
63482
63542
63602
63662
63722
63782
63842
63902
63962
64022
64082
64142
64202
64262
64322
64382
64442
64502
64562
64622
64682
64742
64802
64862
64922
64982
65042
65102
65162
65222
65282
65342
65402
65462
65522
65582
65642
65702
65762
65822
65882
65942
66002
66062
66122
66182
66242
66302
66362
66422
66482
66542
66602
66662
66722
66782
66842
66902
66962
67022
67082
67142
67202
67262
67322
67382
67442
67502
67562
67622
67682
67742
67802
67862
67922
67982
68042
68102
68162
68222
68282
68342
68402
68462
68522
68582
68642
68702
68762
68822
68882
68942
69002
69062
69122
69182
69242
69302
69362
69422
69482
69542
69602
69662
69722
69782
69842
69902
69962
70022
70082
70142
70202
70262
70322
70382
70442
70502
70562
70622
70682
70742
70802
70862
70922
70982
71042
71102
71162
71222
71282
71342
71402
71462
71522
71582
71642
71702
71762
71822
71882
71942
72002
72062
72122
72182
72242
72302
72362
72422
72482
72542
72602
72662
72722
72782
72842
72902
72962
73022
73082
73142
73202
73262
73322
73382
73442
73502
73562
73622
73682
73742
73802
73862
73922
73982
74042
74102
74162
74222
74282
74342
74402
74462
74522
74582
74642
74702
74762
74822
74882
74942
75002
75062
75122
75182
75242
75302
75362
75422
75482
75542
75602
75662
75722
75782
75842
75902
75962
76022
76082
76142
76202
76262
76322
76382
76442
76502
76562
76622
76682
76742
76802
76862
76922
76982
77042
77102
77162
77222
77282
77342
77402
77462
77522
77582
77642
77702
77762
77822
77882
77942
78002
78062
78122
78182
78242
78302
78362
78422
78482
78542
78602
78662
78722
78782
78842
78902
78962
79022
79082
79142
79202
79262
79322
79382
79442
79502
79562
79622
79682
79742
79802
79862
79922
79982
80042
80102
80162
80222
80282
80342
80402
80462
80522
80582
80642
80702
80762
80822
80882
80942
81002
81062
81122
81182
81242
81302
81362
81422
81482
81542
81602
81662
81722
81782
81842
81902
81962
82022
82082
82142
82202
82262
82322
82382
82442
82502
82562
82622
82682
82742
82802
82862
82922
82982
83042
83102
83162
83222
83282
83342
83402
83462
83522
83582
83642
83702
83762
83822
83882
83942
84002
84062
84122
84182
84242
84302
84362
84422
84482
84542
84602
84662
84722
84782
84842
84902
84962
85022
85082
85142
85202
85262
85322
85382
85442
85502
85562
85622
85682
85742
85802
85862
85922
85982
86042
86102
86162
86222
86282
86342
86402
86462
86522
86582
86642
86702
86762
86822
86882
86942
87002
87062
87122
87182
87242
87302
87362
87422
87482
87542
87602
87662
87722
87782
87842
87902
87962
88022
88082
88142
88202
88262
88322
88382
88442
88502
88562
88622
88682
88742
88802
88862
88922
88982
89042
89102
89162
89222
89282
89342
89402
89462
89522
89582
89642
89702
89762
89822
89882
89942
90002
90062
90122
90182
90242
90302
90362
90422
90482
90542
90602
90662
90722
90782
90842
90902
90962
91022
91082
91142
91202
91262
91322
91382
91442
91502
91562
91622
91682
91742
91802
91862
91922
91982
92042
92102
92162
92222
92282
92342
92402
92462
92522
92582
92642
92702
92762
92822
92882
92942
93002
93062
93122
93182
93242
93302
93362
93422
93482
93542
93602
93662
93722
93782
93842
93902
93962
94022
94082
94142
94202
94262
94322
94382
94442
94502
94562
94622
94682
94742
94802
94862
94922
94982
95042
95102
95162
95222
95282
95342
95402
95462
95522
95582
95642
95702
95762
95822
95882
95942
96002
96062
96122
96182
96242
96302
96362
96422
96482
96542
96602
96662
96722
96782
96842
96902
96962
97022
97082
97142
97202
97262
97322
97382
97442
97502
97562
97622
97682
97742
97802
97862
97922
97982
98042
98102
98162
98222
98282
98342
98402
98462
98522
98582
98642
98702
98762
98822
98882
98942
99002
99062
99122
99182
99242
99302
99362
99422
99482
99542
99602
99662
99722
99782
99842
99902
99962
100022
100082
100142
100202
100262
100322
100382
100442
100502
100562
100622
100682
100742
100802
100862
100922
100982
101042
101102
101162
101222
101282
101342
101402
101462
101522
101582
101642
101702
101762
101822
101882
101942
102002
102062
102122
102182
102242
102302
102362
102422
102482
102542
102602
102662
102722
102782
102842
102902
102962
103022
103082
103142
103202
103262
103322
103382
103442
103502
103562
103622
103682
103742
103802
103862
103922
103982
104042
104102
104162
104222
104282
104342
104402
104462
104522
104582
104642
104702
104762
104822
104882
104942
105002
105062
105122
105182
105242
105302
105362
105422
105482
105542
105602
105662
105722
105782
105842
105902
105962
106022
106082
106142
106202
106262
106322
106382
106442
106502
106562
106622
106682
106742
106802
106862
106922
106982
107042
107102
107162
107222
107282
107342
107402
107462
107522
107582
107642
107702
107762
107822
107882
107942
108002
108062
108122
108182
108242
108302
108362
108422
108482
108542
108602
108662
108722
108782
108842
108902
108962
109022
109082
109142
109202
109262
109322
109382
109442
109502
109562
109622
109682
109742
109802
109862
109922
109982
110042
110102
110162
110222
110282
110342
110402
110462
110522
110582
110642
110702
110762
110822
110882
110942
111002
111062
111122
111182
111242
111302
111362
111422
111482
111542
111602
111662
111722
111782
111842
111902
111962
112022
112082
112142
112202
112262
112322
112382
112442
112502
11256
```