



US009201044B2

(12) **United States Patent**  
**Kearney et al.**

(10) **Patent No.:** **US 9,201,044 B2**  
(45) **Date of Patent:** **Dec. 1, 2015**

(54) **COMPOSITIONS, METHODS AND KITS FOR DIAGNOSIS OF LUNG CANCER**

FOREIGN PATENT DOCUMENTS

(71) Applicant: **Integrated Diagnostics, Inc.**, Seattle, WA (US)

WO WO-2011085163 A2 7/2011  
WO 2012075042 A1 6/2012  
WO 2013096845 A2 6/2013

(72) Inventors: **Paul Edward Kearney**, Seattle, WA (US); **Kenneth Charles Fang**, San Francisco, CA (US); **Xiao-Jun Li**, Bellevue, WA (US); **Clive Hayward**, Seattle, WA (US)

OTHER PUBLICATIONS

(73) Assignee: **Integrated Diagnostics, Inc.**, Seattle, WA (US)

Swensen et al (Radiology, 2003, 226:756-761).\*  
Wei et al (Journal of Proteome Research, 2011, 10:161-174, published online Sep. 3, 2010).\*  
Bouchal et al (Journal of Proteome Research, 2009, 8:362-373).\*  
Kitada et al (World Journal of Surgical Oncology, 2011, 9:124; internet pp. 1-5).\*  
Bouchal et al (Journal of Proteome Research, 2009, 8:362-373) Supplementary Table 2.\*  
Wang et al (Cancer Biology and Therapy, 2009, 8:1083-1094).\*  
Lange et al (2008, Molecular Systems Biology 4: 222, internet pp. 1-14).\*

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 199 days.

Jung-Hyun Rho et al., "Glycoproteomic Analysis of Human Lung Adenocarcinomas Using Glycoarrays and Tandem Mass Spectrometry: Differential Expression and Glycosylation Patterns of Vimentin and Fetuin A Isoforms", Journal of Protein Chemistry, Kluwer Academic Publishers-Plenum Publishers, NE, vol. 28, No. 3-4, May 2, 2009, pp. 148-160.

(21) Appl. No.: **13/724,823**

(22) Filed: **Dec. 21, 2012**

"Evolution of Translational Omics: Lessons Learned and the Path Forward." *Committee on the Review of Omics-Based Tests for Predicting Patient Outcomes in Clinical Trials*. Micheel et al., eds. (2012):xv-338.

(65) **Prior Publication Data**

US 2013/0217057 A1 Aug. 22, 2013

Addona et al. "A Pipeline that Integrates the Discovery and Verification of Plasma Protein Biomarkers Reveals Candidate Markers for Cardiovascular Disease." *Nat. Biotechnol.* 29.7(2011):635-643.

**Related U.S. Application Data**

(60) Provisional application No. 61/578,712, filed on Dec. 21, 2011, provisional application No. 61/589,920, filed on Jan. 24, 2012, provisional application No. 61/676,859, filed on Jul. 27, 2012, provisional application No. 61/725,153, filed on Nov. 12, 2012.

Addona et al. "Multi-Site Assessment of the Precision and Reproducibility of Multiple Reaction Monitoring-Based Measurements of Proteins in Plasma." *Nat. Biotechnol.* 27.7(2009):633-641.

(51) **Int. Cl.**

**G01N 33/53** (2006.01)  
**G01N 27/62** (2006.01)  
**G01N 33/574** (2006.01)  
**G06F 19/18** (2011.01)

Albert et al. "Evaluation of the Solitary Pulmonary Nodule." *Am. Fam. Physician.* 80.8(2009):827-831.

(52) **U.S. Cl.**

CPC ..... **G01N 27/62** (2013.01); **G01N 33/57423** (2013.01); **G01N 2800/60** (2013.01); **G06F 19/18** (2013.01)

Bigbee et al. "A Multiplexed Serum Biomarker Immunoassay Panel Discriminates Clinical Lung Cancer Patients from High-Risk Individuals Found to be Cancer-Free by CT Scanning." *J. Thorac Oncol.* 7.4(2012):698-708.

(58) **Field of Classification Search**

None  
See application file for complete search history.

Brusniak et al. "Corra: Computational Framework and Tools for LC-MS Discovery and Targeted Mass Spectrometry-Based Proteomics." *BMC Bioinformatics.* 9(2008):542.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,183,188 B2 2/2007 Kronke et al.  
2006/0257857 A1 11/2006 Keene et al.  
2007/0099251 A1 5/2007 Zhang et al.  
2007/0111322 A1 5/2007 Yang  
2007/0128598 A1 6/2007 Boender  
2007/0202539 A1 8/2007 Aebersold et al.  
2007/0269895 A1 11/2007 Aebersold et al.  
2009/0317392 A1\* 12/2009 Nakamura et al. .... 424/139.1  
2010/0093108 A1 4/2010 Khattar et al.  
2010/0184034 A1\* 7/2010 Bankaitis-Davis et al. .... 435/6  
2010/0279382 A1 11/2010 Aebersold et al.  
2012/0142558 A1\* 6/2012 Li et al. .... 506/18  
2013/0230877 A1 9/2013 Kearney

Carozzi et al. "Molecular Profile in Body Fluids in Subjects Enrolled in a Randomised Trial for Lung Cancer Screening: Perspectives of Integrated Strategies for Early Diagnosis." *Lung Cancer.* 68.2(2010):216-221.

Chapman et al. "EarlyCDT®-Lung Test: Improved Clinical Utility Through Additional Autoantibody Assays." *Tumor Biol.* 33.5(2012):1319-1326.

(Continued)

*Primary Examiner* — Laura B Goddard

(74) *Attorney, Agent, or Firm* — Cooley LLP; Ivor R. Elrifi; Matthew Pavao

(57)

**ABSTRACT**

Methods are provided for identifying biomarker proteins that exhibit differential expression in subjects with a first lung condition versus healthy subjects or subjects with a second lung condition. Also provided are compositions comprising these biomarker proteins and methods of using these biomarker proteins or panels thereof to diagnose, classify, and monitor various lung conditions. The methods and compositions provided herein may be used to diagnose or classify a subject as having lung cancer or a non-cancerous condition, and to distinguish between different types of cancer (e.g., malignant versus benign, SCLC versus NSCLC).

**15 Claims, 11 Drawing Sheets**

(56)

## References Cited

## OTHER PUBLICATIONS

- Cima et al. "Cancer Genetics-Guided Discovery of Serum Biomarker Signatures for Diagnosis and Prognosis of Prostate Cancer." *PNAS*. 108.8(2011):3342-3347.
- Desiere et al. "The PeptideAtlas Project." *Nucleic Acids Res.* 34(2006):D655-D658.
- Farrah et al. "A High-Confidence Human Plasma Proteome Reference Set with Estimated Concentrations in PeptideAtlas." *Mol. Cell. Proteomics*. 10.9(2011):M110.006353.
- Gould et al. "Evaluation of Patients with Pulmonary Nodules: When is it Lung Cancer?" *Chest*. 132.S3(2007):108S-130S.
- Halliwell et al. "Oxidative Stress and Cancer: Have We Moved Forward?" *Biochem. J.* 401.1(2007):1-11.
- Hanash et al. "Emerging Molecular Biomarkers—Blood-Based Strategies to Detect and Monitor Cancer." *Nat. Rev. Clin. Oncol.* 8.3(2011):142-150.
- Hassanein et al. "Advances in Proteomic Strategies Toward the Early Detection of Lung Cancer." *Proc. Am. Thorac. Soc.* 8.2(2011):183-188.
- Hennessey et al. "Serum MicroRNA Biomarkers for Detection of Non-Small Cell Lung Cancer." *PLoS One*. 7.2(2012):e32307.
- Henschke et al. "CT Screening for Lung Cancer: Suspiciousness of Nodules According to Size on Baseline Scans." *Radiology*. 231.1(2004):164-168.
- Henschke et al. "Early Lung Cancer Action Project: Overall Design and Findings from Baseline Screenings." *Lancet*. 354.9173(1999):99-105.
- Hüttenhain et al. "Reproducible Quantification of Cancer-Associated Proteins in Body Fluids using Targeted Proteomics." *Sci. Transl. Med.* 4.142(2012):149ra194.
- Kearney et al. "Protein Identification and Peptide Expression Resolver: Harmonizing Protein Identification with Protein Expression Data." *J. Proteome Res.* 7.1(2008):234-244.
- Kitteringham et al. "Multiple Reaction Monitoring for Quantitative Biomarker Analysis in Proteomics and Metabolomics." *J. Chromatogr. B*. 877.13(2009):1229-1239.
- Lam et al. "Early CDT-Lung: An Immunobiomarker Test as an Aid to Early Detection of Lung Cancer." *Cancer Prev. Res.* 4.7(2011):1126-1134.
- Lange et al. "Selected Reaction Monitoring for Quantitative Proteomics: A Tutorial." *Mol. Sys. Biol.* 4(2008):222.
- Lehtiö et al. "Lung Cancer Proteomics, Clinical and Technological Considerations." *J. Proteomics*. 73.10(2010):1851-1863.
- Lombardi et al. Clinical Significance of a Multiple Biomarker Assay in Patients with Lung Cancer. *Chest*. 97.3(1990):639-644.
- MacMahon et al. "Guidelines for Management of Small Pulmonary Nodules Detected on CT Scans: A Statement from the Fleischner Society." *Radiology*. 237.2(2005):395-400.
- Makawita et al. "The Bottleneck in the Cancer Biomarker Pipeline and Protein Quantification through Mass Spectrometry-Based Approaches: Current Strategies for Candidate Verification." *Clin. Chem.* 56.2(2010):212-222.
- McClish. "Analyzing a Portion of the ROC Curve." *Med. Decis. Making*. 9.3(1989):190-195.
- Miller et al. "Minimizing Unintended Consequences of Detecting Lung Nodules by Computed Tomography." *Am. J. Resp. Crit. Care Med.* 178.9(2008):891-892.
- Ocak et al. "Mass Spectrometry-Based Proteomic Profiling of Lung Cancer." *Proc. Am. Thorac. Soc.* 6.2(2009):159-170.
- Omenn et al. "Overview of the HUPO Plasma Proteome Project: Results from the Pilot Phase with 35 Collaborating Laboratories and Multiple Analytical Groups, Generating a Core Dataset of 3020 Proteins and a Publicly-Available Database." *Proteomics*. 5.13(2005):3226-3245.
- Ost et al. "Decision Making in Patients with Pulmonary Nodules." *Am. J. Respir. Crit. Care Med.* 185.4(2012):363-372.
- Ostroff et al. "Unlocking Biomarker Discovery: Large Scale Application of Aptamer Proteomic Technology for Early Detection of Lung Cancer." *PLoS One*. 5.12(2010):e15003.
- Pecot et al. "Added Value of a Serum Proteomic Signature in the Diagnostic Evaluation of Lung Nodules." *Cancer Epidemiol. Biomarkers Prev.* 21.5(2012):786-792.
- Perkins et al. "Probability-Based Protein Identification by Searching Sequence Databases Using Mass Spectrometry Data." *Electrophoresis*. 20.18(1999):3551-3567.
- Picotti et al. "High-Throughput Generation of Selected Reaction-Monitoring Assays for Proteins and Proteomes." *Nat. Meth.* 7.1(2010):43-46.
- Polanski et al. "A List of Candidate Cancer Biomarkers for Targeted Proteomics." *Biomarker Insights*. 1(2007):1-48.
- Price et al. "Highly Accurate Two-Gene Classifier for Differentiating Gastrointestinal Stromal Tumors and Leiomyosarcomas." *PNAS*. 104.9(2007):3414-3419.
- Qin et al. "SRM Targeted Proteomics in Search for Biomarkers of HCV-Induced Progression of Fibrosis to Cirrhosis in HALT-C Patients." *Proteomics*. 12.8(2012):1244-1252.
- Radulovic et al. "Informatics Platform for Global Proteomic Profiling and Biomarker Discovery Using Liquid Chromatography-Tandem Mass Spectrometry." *Mol. Cell. Proteins*. 3.10(2004):984-997.
- Reiter et al. "mProphet: Automated Data Processing and Statistical Validation for Large-Scale SRM Experiments." *Nat. Meth.* 8.5(2011):430-435.
- Rho et al. "Glycoproteomic Analysis of Human Lung Adenocarcinomas Using Glycoarrays and Tandem Mass Spectrometry: Differential Expression and Glycosylation Patterns of Vimentin and Fetuin A Isoforms." *Protein J.* 28.3-4(2009):148-160.
- Rom et al. "Identification of an Autoantibody Panel to Separate Lung Cancer from Smokers and Nonsmokers." *BMC Cancer*. 10(2010):234.
- Schauer et al. "National Council on Radiation Protection and Measurements Report Shows Substantial Medical Exposure Increase." *Radiol.* 253.2(2009):293-296.
- States et al. "Challenges in Deriving High-Confidence Protein Identifications from Data Gathered by a HUPO Plasma Proteome Collaborative Study." *Nat. Biotechnol.* 24.3(2006):333-338.
- Stern et al. "Nationwide Evaluation of X-Ray Trends (NEXT) 2000-01 Survey of Patient Radiation Exposure from Computed Tomographic (CT) Examinations in the United States." *87th Scientific Assembly and Annual Meeting of the Radiological Society of North America, Chicago*, Nov. 25-30, 2001.
- Taguchi et al. "Unleashing the Power of Proteomics to Develop Blood-Based Cancer Markers." *Clin. Chem.* 59(2013):1.
- Teutsch et al. "The Evaluation of Genomic Applications in Practice and Prevention (EGAPP) Initiative: Methods of the EGAPP Working Group." *Genet. Med.* 11.1(2009):3-14.
- Walser et al. "Smoking and Lung Cancer: The Role of Inflammation." *Proc. Am. Thorac. Soc.* 5.8(2008):811-815.
- Whiteaker et al. "A Targeted Proteomics-Based Pipeline for Verification of Biomarkers in Plasma." *Nat. Biotechnol.* 29.7(2011):625-634.
- Wiener et al. "Population-Based Risk for Complications after Trans thoracic Needle Lung Biopsy of a Pulmonary Nodule: An Analysis of Discharge Records." *Ann. Int. Med.* 155.3(2011):137-144.
- Yildiz et al. "Diagnostic Accuracy of MALDI Mass Spectrometric Analysis of Unfractionated Serum in Lung Cancer." *J. Thorac. Oncol.* 2.10(2007):893-901.
- Zeng et al. "Lung Cancer Serum Biomarker Discovery Using Glycoprotein Capture and Liquid Chromatography Mass Spectrometry." *J. Proteome Res.* 9.12(2010):6440-6449.
- Tockman M. et al., "Considerations in Bringing a Cancer Biomarker to Clinical Applications", *Cancer Research*, (1992), vol. 52, p. 2711 s-2718s.
- Ozaki Y. et al., "Expression and Immunogenicity of a Tumor-Associated Antigen, 90K/Mac-2 Binding Protein, in Lung Carcinoma", *Cancer*, (2002), vol. 95, p. 1954-1962.
- Ueda K. et al., "A Comprehensive Peptidome Profiling Technology for the Identification of Early Detection Biomarkers for Lung Adenocarcinoma", *PLoS One*, (Apr. 12, 2011), vol. 6, issue 4, e18567, p. 1-12.

(56)

**References Cited**

OTHER PUBLICATIONS

Milman et al., "The serum ferritin concentration is a significant prognostic indicator of survival in primary lung cancer", *Oncology Reports*, (2002), vol. 9, No. 1, p. 193-198.

Fracchia A. et al., "A Comparative Study on Ferritin Concentration in Serum and Bilateral Bronchoalveolar Lavage Fluid of Patients with Peripheral Lung Cancer versus Control Subjects", *Oncology*, (1999), vol. 56, p. 181-188.

\* cited by examiner





FIG. 1

FIG. 2






FIG. 3





FIG. 4

FIG. 5






FIG. 6

FIG. 7




FIG. 8




FIG. 9




FIG. 10





FIG. 11



## COMPOSITIONS, METHODS AND KITS FOR DIAGNOSIS OF LUNG CANCER

### RELATED APPLICATIONS

This application claims priority and benefit of U.S. Provisional Application No. 61/578,712 filed Dec. 21, 2011, U.S. Provisional Application No. 61/589,920 filed Jan. 24, 2012, U.S. Provisional Application No. 61/676,859 filed Jul. 27, 2012 and U.S. Provisional Application No. 61/725,153 filed Nov. 12, 2012, the contents of each of which are incorporated herein by reference in their entireties.

### INCORPORATION-BY-REFERENCE OF SEQUENCE LISTING

The contents of the text file named "44549-505001US\_ST25.txt", which was created on Feb. 25, 2013 and is 6 KB in size, are hereby incorporated by reference in their entireties.

### BACKGROUND

Lung conditions and particularly lung cancer present significant diagnostic challenges. In many asymptomatic patients, radiological screens such as computed tomography (CT) scanning are a first step in the diagnostic paradigm. Pulmonary nodules (PNs) or indeterminate nodules are located in the lung and are often discovered during screening of both high risk patients or incidentally. The number of PNs identified is expected to rise due to increased numbers of patients with access to health care, the rapid adoption of screening techniques and an aging population. It is estimated that over 3 million PNs are identified annually in the US. Although the majority of PNs are benign, some are malignant leading to additional interventions. For patients considered low risk for malignant nodules, current medical practice dictates scans every three to six months for at least two years to monitor for lung cancer. The time period between identification of a PN and diagnosis is a time of medical surveillance or "watchful waiting" and may induce stress on the patient and lead to significant risk and expense due to repeated imaging studies. If a biopsy is performed on a patient who is found to have a benign nodule, the costs and potential for harm to the patient increase unnecessarily. Major surgery is indicated in order to excise a specimen for tissue biopsy and diagnosis. All of these procedures are associated with risk to the patient including: illness, injury and death as well as high economic costs.

Frequently, PNs cannot be biopsied to determine if they are benign or malignant due to their size and/or location in the lung. However, PNs are connected to the circulatory system, and so if malignant, protein markers of cancer can enter the blood and provide a signal for determining if a PN is malignant or not.

Diagnostic methods that can replace or complement current diagnostic methods for patients presenting with PNs are needed to improve diagnostics, reduce costs and minimize invasive procedures and complications to patients. The present invention provides novel compositions, methods and kits for identifying protein markers to identify, diagnose, classify and monitor lung conditions, and particularly lung cancer. The present invention uses a blood-based multiplexed assay to distinguish benign pulmonary nodules from malignant pulmonary nodules to classify patients with or without

lung cancer. The present invention may be used in patients who present with symptoms of lung cancer, but do not have pulmonary nodules.

### SUMMARY

The present invention provides a method of determining the likelihood that a lung condition in a subject is cancer by measuring an abundance of a panel of proteins in a sample obtained from the subject; calculating a probability of cancer score based on the protein measurements and ruling out cancer for the subject if the score) is lower than a pre-determined score, wherein When cancer is ruled out the subject does not receive a treatment protocol. Treatment protocols include for example pulmonary function test (PFT), pulmonary imaging, a biopsy, a surgery, a chemotherapy, a radiotherapy, or any combination thereof. In some embodiments, the imaging is an x-ray, a chest computed tomography (CT) scan, or a positron emission tomography (PET) scan.

The present invention further provides a method of ruling in the likelihood of cancer for a subject by measuring an abundance of panel of proteins in a sample obtained from the subject, calculating a probability of cancer score based on the protein measurements and ruling in the likelihood of cancer for the subject if the score in step is higher than a pre-determined score

In another aspect, the invention further provides a method of determining the likelihood of the presence of a lung condition in a subject by measuring an abundance of panel of proteins in a sample obtained from the subject, calculating a probability of cancer score based on the protein measurements and concluding the presence of said lung condition if the score is equal or greater than a pre-determined score. The lung condition is lung cancer such as for example, non-small cell lung cancer (NSCLC). The subject at risk of developing lung cancer

The panel includes at least 4 proteins selected from ALDOA, FRIL, LG3BP, IBP3, LRP1, ISLR, TSP COIA1, GRP78, TETN, PRXD1 and CD14. Optionally, the panel further includes at least one protein selected from BGH3, COIA1, TETN, GRP78, PRDX, FIBA and GSLG1.

The subject has or is suspected of having a pulmonary nodule. The pulmonary nodule has a diameter of less than or equal to 3 cm. In one embodiment, the pulmonary nodule has a diameter of about 0.8 cm to 2.0 cm.

The score is calculated from a logistic regression model applied to the protein measurements. For example, the score is determined as  $P_s = 1/[1 + \exp(-\alpha - \sum_{i=1}^N \beta_i * \check{I}_{i,s})]$ , where  $\check{I}_{i,s}$  is logarithmically transformed and normalized intensity of transition i in said sample (s),  $\beta_i$  is the corresponding logistic regression coefficient,  $\alpha$  was a panel-specific constant, and N was the total number of transitions in said panel.

In various embodiments, the method of the present invention further comprises normalizing the protein measurements. For example, the protein measurements are normalized by one or more proteins selected from PEDF, MASP1, GELS, LUM, C163A and PTPRJ.

The biological sample such as for example tissue, blood, plasma, serum, whole blood, urine, saliva, genital secretion, cerebrospinal fluid, sweat and excreta.

In one aspect, the determining the likelihood of cancer is determined by the sensitivity, specificity, negative predictive value or positive predictive value associated with the score. The score determined has a negative predictive value (NPV) is at least about 80%.

The measuring step is performed by selected reaction monitoring mass spectrometry, using a compound that spe-

cifically binds the protein being detected or a peptide transition. In one embodiment, the compound that specifically binds to the protein being measured is an antibody or an aptamer.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a line graph showing area under the curve for a receiving operating curve for 15 protein LC-SRM-MS panels.

FIG. 2 shows six line graphs each showing area under the curve for a receiving operating curve for 15 protein LC-SRM-MS panels for different patient populations and for subjects with large and small PN

FIG. 3 is a graph showing variability among three studies used to evaluate 15 protein panels.

FIG. 4 is a line graph showing area under the curve for a receiving operating curve for a 15 protein LC-SRM-MS panel.

FIG. 5 shows three line graphs each showing area under the curve for a receiving operating curve for a 15 protein LC-SRM-MS panel for a different patient population.

FIG. 6 shows the results of a query of blood proteins used to identify lung cancer using the "Ingenuity"® program.

FIG. 7 is a bar diagram showing Pearson correlations for peptides from the same peptide, from the same protein and from different proteins.

FIG. 8 is a graph showing performance of the classifier on the training samples, validation samples and all samples combined.

FIG. 9 is a graph showing clinical and molecular factors.

FIG. 10 is a schematic showing the molecular network containing the 13 classifier proteins (green), 5 transcription factors (blue) and the three networks (orange lines) of lung cancer, response to oxidative stress and lung inflammation.

FIG. 11 is a graph depicting interpretation of classifier score in terms of risk

#### DETAILED DESCRIPTION

The disclosed invention derives from the surprising discovery, that in patients presenting with pulmonary nodule(s), protein markers in the blood exist that specifically identify and classify lung cancer. Accordingly the invention provides unique advantages to the patient associated with early detection of lung cancer in a patient, including increased life span, decreased morbidity and mortality, decreased exposure to radiation during screening and repeat screenings and a minimally invasive diagnostic model. Importantly, the methods of the invention allow for a patient to avoid invasive procedures.

The routine clinical use of chest computed tomography (CT) scans identifies millions of pulmonary nodules annually, of which only a small minority are malignant but contribute to the dismal 15% five-year survival rate for patients diagnosed with non-small cell lung cancer (NSCLC). The early diagnosis of lung cancer in patients with pulmonary nodules is a top priority, as decision-making based on clinical presentation, in conjunction with current non-invasive diagnostic options such as chest CT and positron emission tomography (PET) scans, and other invasive alternatives, has not altered the clinical outcomes of patients with Stage I NSCLC. The subgroup of pulmonary nodules between 8 mm and 20 mm in size is increasingly recognized as being "intermediate" relative to the lower rate of malignancies below 8 mm and the higher rate of malignancies above 20 mm [9]. Invasive sampling of the lung nodule by biopsy using transthoracic needle aspiration or bronchoscopy may provide a cytopathologic

diagnosis of NSCLC, but are also associated with both false-negative and non-diagnostic results. In summary, a key unmet clinical need for the management of pulmonary nodules is a non-invasive diagnostic test that discriminates between malignant and benign processes in patients with indeterminate pulmonary nodules (IPNs), especially between 8 mm and 20 mm in size.

The clinical decision to be more or less aggressive in treatment is based on risk factors, primarily nodule size, smoking history and age [9] in addition to imaging. As these are not conclusive, there is a great need for a molecular-based blood test that would be both non-invasive and provide complementary information to risk factors and imaging.

Accordingly, these and related embodiments will find uses in screening methods for lung conditions, and particularly lung cancer diagnostics. More importantly, the invention finds use in determining the clinical management of a patient. That is, the method of invention are useful in ruling in or ruling out a particular treatment protocol for an individual subject.

Cancer biology requires a molecular strategy to address the unmet medical need for an assessment of lung cancer risk. The field of diagnostic medicine has evolved with technology and assays that provide sensitive mechanisms for detection of changes in proteins. The methods described herein use a LC-SRM-MS technology for measuring the concentration of blood plasma proteins that are collectively changed in patients with a malignant PN. This protein signature is indicative of lung cancer. LC-SRM-MS is one method that provides for both quantification and identification of circulating proteins in plasma. Changes in protein expression levels, such as but not limited to signaling factors, growth factors, cleaved surface proteins and secreted proteins, can be detected using such a sensitive technology to assay cancer. Presented herein is a blood-based classification test to determine the likelihood that a patient presenting with a pulmonary nodule has a nodule that is benign or malignant. The present invention presents a classification algorithm that predicts the relative likelihood of the PN being benign or malignant.

More broadly, it is demonstrated that there are many variations on this invention that are also diagnostic tests for the likelihood that a PN is benign or malignant. These are variations on the panel of proteins, protein standards, measurement methodology and/or classification algorithm.

As disclosed herein, archival plasma samples from subjects presenting with PNs were analyzed for differential protein expression by mass spectrometry and the results were used to identify biomarker proteins and panels of biomarker proteins that are differentially expressed in conjunction with various lung conditions (cancer vs. non-cancer).

In one aspect of the invention, one hundred and sixty three panels were discovered that allow for the classification of PN as being benign or malignant. These panels include those listed on Table 1. In some embodiments the panel according to the invention includes measuring 1, 2, 3, 4, 5 or more proteins selected from ISLR, ALDOA, KIT, GRP78, AIFM1, CD14, COIA1, IBP3, TSP1, BGH3, TETN, FR1, LG3BP, GGH, PRDX1 or LRP1. In other embodiments the panel includes any panel or protein exemplified on Table 1. For example the panel includes ALDOA, GRP78, CD14, COIA1, IBP3, FRIL, LG3BP, and LRP1

TABLE 1

Identifier	Number	pAUC	Proteins						
	Proteins	Factor	ISLR	ALDOA	KIT	GRP78	AIFM1	CD14	COIA1
1	9	4.562	0	1	0	1	0	1	1
2	8	4.488	0	1	0	1	0	1	1
3	11	4.451	1	1	0	1	0	0	1
4	11	4.357	1	1	0	1	0	0	1
5	11	4.331	1	1	0	0	0	1	1
6	13	4.324	1	1	0	0	0	1	1
7	10	4.205	1	1	0	1	0	0	1
8	11	4.193	1	1	0	0	0	0	1
9	12	4.189	1	1	0	1	0	0	1
10	12	4.182	1	0	0	0	0	1	1
11	12	4.169	1	1	0	1	0	0	1
12	8	4.107	1	1	0	1	0	1	1
13	13	4.027	0	1	1	1	0	1	1
14	10	3.994	0	1	1	1	0	1	1
15	11	3.979	1	1	1	1	0	1	1
16	10	3.932	1	1	0	1	0	1	1
17	11	3.926	1	1	0	0	0	1	1
18	12	3.913	1	0	1	1	0	0	1
19	12	3.872	0	1	1	1	0	1	1
20	12	3.864	1	1	1	0	0	1	1
21	14	3.853	1	1	0	1	0	1	1
22	9	3.849	1	1	0	1	0	0	1
23	12	3.846	1	1	1	1	0	0	1
24	10	3.829	0	1	1	1	0	1	0
25	10	3.829	0	1	1	1	0	1	1
26	12	3.826	1	0	0	0	1	0	1
27	7	3.804	1	1	0	1	0	1	1
28	10	3.802	0	1	0	1	0	1	1
29	10	3.787	0	1	0	1	0	1	0
30	9	3.779	1	1	0	1	0	1	1
31	11	3.774	0	1	0	1	0	1	1
32	8	3.759	1	1	0	0	0	0	1
33	13	3.758	1	1	0	0	0	1	1
34	11	3.757	1	1	0	1	0	0	0
35	12	3.754	0	1	1	1	0	1	1
36	10	3.750	1	1	0	1	0	1	1
37	11	3.747	0	1	1	1	0	1	1
38	12	3.744	1	0	1	1	0	0	1
39	11	3.742	1	1	0	1	0	1	1
40	9	3.740	1	1	0	1	0	1	1
41	12	3.740	1	1	1	1	0	1	1
42	12	3.739	1	1	0	1	0	1	1
43	9	3.734	1	1	0	0	0	0	1
44	12	3.730	1	1	0	1	0	0	1
45	11	3.725	0	1	1	1	0	1	1
46	12	3.717	0	1	0	0	1	1	1
47	9	3.713	0	1	0	1	0	1	1
48	9	3.713	1	1	1	1	0	1	1
49	10	3.709	0	1	0	1	0	1	1
50	11	3.709	1	1	0	1	0	1	1
51	11	3.701	0	1	1	1	1	1	1
52	12	3.685	1	1	0	1	0	1	1
53	10	3.680	0	0	0	1	0	1	0
54	11	3.676	1	1	1	1	0	0	1
55	9	3.668	0	1	0	1	0	1	1
56	9	3.659	0	0	0	1	0	1	0
57	14	3.657	1	1	0	1	1	1	1
58	10	3.655	1	1	0	1	0	0	1
59	11	3.643	0	1	1	1	0	1	1
60	9	3.643	0	1	0	1	0	1	0
61	8	3.640	1	1	0	1	0	1	0
62	12	3.640	1	1	1	1	0	1	1
63	10	3.638	1	1	0	1	0	0	1
64	12	3.633	1	0	0	1	1	0	1
65	10	3.632	1	1	0	1	0	1	1
66	11	3.627	1	1	0	1	0	1	0
67	10	3.627	1	1	0	0	0	1	0
68	10	3.623	1	1	1	0	0	0	1
69	11	3.619	1	0	0	1	0	1	1
70	6	3.617	1	1	0	1	0	0	1
71	12	3.617	1	0	0	1	0	1	1
72	11	3.613	1	1	0	1	0	1	0
73	11	3.608	1	1	0	1	0	1	0
74	13	3.608	1	1	1	1	0	1	1
75	11	3.605	0	1	1	1	0	1	1
76	11	3.602	0	1	1	1	0	1	1

TABLE 1-continued

77	10	3.600	1	1	0	1	0	0	0
78	11	3.596	1	1	0	1	0	0	1
79	10	3.592	1	1	0	1	0	1	0
80	11	3.587	1	0	1	0	0	0	1
81	13	3.584	1	1	0	1	1	1	1
82	8	3.584	0	1	0	1	0	1	0
83	11	3.581	1	1	1	1	0	1	0
84	13	3.578	1	1	0	1	0	1	0
85	9	3.573	1	1	1	0	0	1	1
86	9	3.572	1	1	0	1	0	0	1
87	13	3.571	1	1	1	1	0	1	0
88	10	3.569	1	1	0	1	0	0	1
89	9	3.569	0	1	0	1	0	1	0
90	8	3.559	0	1	0	1	0	1	0
91	10	3.558	0	1	0	1	0	1	0
92	12	3.554	1	1	0	1	0	1	1
93	11	3.552	0	1	0	1	0	1	0
94	12	3.549	0	1	0	1	0	1	0
95	8	3.547	1	1	1	0	0	1	1
96	12	3.545	1	1	1	1	0	1	1
97	8	3.542	1	1	1	0	0	0	0
98	11	3.536	1	1	1	1	0	0	1
99	14	3.530	1	1	1	1	0	1	1
100	9	3.527	1	1	0	1	0	1	1
101	10	3.522	0	1	1	0	1	1	1
102	12	3.509	1	1	0	1	0	1	1
103	5	3.505	0	1	0	0	0	1	0
104	11	3.500	1	1	0	0	1	0	1
105	11	3.497	1	1	1	1	0	0	1
106	9	3.491	1	1	0	0	0	1	0
107	7	3.489	0	1	1	0	0	1	0
108	13	3.486	1	1	1	1	0	1	1
109	11	3.483	1	1	1	1	0	0	1
110	10	3.477	1	1	1	1	0	1	1
111	10	3.473	1	1	0	0	0	1	1
112	15	3.468	1	1	0	1	1	1	1
113	10	3.467	0	1	0	0	1	1	0
114	12	3.467	1	1	0	0	1	1	1
115	13	3.467	1	1	0	1	1	0	1
116	10	3.467	0	1	0	1	0	1	0
117	8	3.465	1	1	0	1	0	0	1
118	10	3.464	0	1	0	1	1	1	1
119	15	3.464	1	1	0	1	1	1	1
120	11	3.462	1	1	0	1	0	1	1
121	9	3.460	1	1	0	0	0	1	0
122	13	3.453	1	1	0	1	0	1	1
123	12	3.449	1	1	1	0	0	1	0
124	10	3.448	1	1	0	1	0	1	0
125	10	3.445	0	1	1	1	0	1	0
126	6	3.441	0	1	0	0	0	1	0
127	11	3.440	1	1	0	1	0	1	0
128	12	3.440	1	1	0	1	1	0	0
129	11	3.439	1	1	0	1	0	1	0
130	10	3.426	0	1	0	0	1	1	0
131	11	3.423	1	1	0	0	0	0	1
132	10	3.420	1	1	0	0	0	1	0
133	10	3.419	1	1	1	1	0	1	0
134	11	3.417	1	1	0	1	1	0	1
135	12	3.414	0	1	0	1	1	1	1
136	10	3.413	0	1	1	1	0	1	0
137	11	3.400	0	1	0	0	1	1	0
138	12	3.398	1	1	0	1	0	1	0
139	13	3.396	1	1	0	1	0	1	0
140	9	3.386	1	1	0	0	0	1	0
141	9	3.373	1	1	0	1	0	1	0
142	12	3.363	1	1	0	0	1	0	1
143	8	3.362	0	1	0	1	0	1	0
144	10	3.360	1	1	0	1	0	1	1
145	9	3.359	1	1	1	0	0	1	0
146	7	3.349	0	1	0	0	0	0	0
147	7	3.348	1	1	0	0	0	1	1
148	9	3.340	1	0	0	0	0	1	0
149	9	3.335	1	1	0	1	0	1	0
150	11	3.333	0	1	1	1	0	1	0
151	9	3.333	0	0	0	1	0	1	0
152	10	3.328	1	1	0	1	0	1	0
153	7	3.315	0	1	0	1	0	1	0
154	11	3.311	1	1	0	1	1	1	1
155	11	3.293	1	1	0	1	0	1	0
156	8	3.292	1	1	0	1	0	0	0

TABLE 1-continued

157	9	3.289	0	1	0	1	0	1	0
158	7	3.229	0	1	0	0	0	1	0
159	7	3.229	1	1	0	0	0	1	0
160	7	3.203	1	1	0	1	0	0	0
161	12	3.161	1	1	1	0	1	1	0
162	9	3.138	1	1	0	0	1	0	1
163	13	3.078	1	1	0	0	1	0	1

Identifier	IBP3	TSP1	BGH3	TETN	FRIL	LG3BP	GGH	PRDX1	LRP1
1	1	0	0	0	1	1	0	0	1
2	1	0	0	0	1	1	0	0	1
3	1	1	1	1	1	0	0	1	1
4	1	1	0	0	1	1	1	1	1
5	0	1	1	1	1	0	1	1	1
6	1	1	1	1	1	1	1	1	1
7	0	1	1	1	1	0	0	1	1
8	0	1	1	1	1	0	1	1	1
9	1	1	1	1	1	0	0	1	1
10	1	1	1	1	1	1	0	1	1
11	1	1	0	0	1	1	1	1	1
12	0	0	0	0	1	1	0	0	1
13	1	1	0	0	1	1	1	1	1
14	1	0	0	0	1	1	0	0	1
15	0	0	0	0	1	1	1	0	1
16	0	0	0	1	1	1	0	0	1
17	1	1	1	1	1	0	0	1	1
18	1	1	0	0	1	1	1	1	1
19	1	0	0	0	1	1	1	1	1
20	0	1	1	1	1	1	0	1	1
21	1	1	1	1	1	1	0	1	1
22	0	1	1	1	1	0	0	0	1
23	1	1	0	0	1	1	1	1	1
24	1	0	0	0	1	1	1	1	1
25	1	0	0	0	1	1	1	0	1
26	1	1	1	1	1	0	1	1	1
27	0	0	0	0	0	1	0	0	1
28	1	0	0	0	1	1	1	1	1
29	1	1	0	0	1	1	1	1	1
30	0	0	0	0	1	1	0	0	1
31	1	0	0	0	1	1	1	1	1
32	0	0	1	1	1	0	0	1	1
33	1	1	1	1	1	1	0	1	1
34	1	1	1	1	1	1	0	1	1
35	1	1	0	0	1	1	1	1	1
36	1	0	0	0	1	1	0	1	1
37	1	1	0	0	1	1	1	1	0
38	1	1	1	1	1	0	0	1	1
39	1	1	0	1	1	1	0	0	1
40	1	0	0	0	1	1	0	0	1
41	1	0	0	1	1	1	0	0	1
42	1	1	0	0	1	1	1	1	1
43	0	1	1	1	1	0	0	1	1
44	1	1	1	1	1	1	0	1	1
45	1	0	0	1	1	1	0	0	1
46	1	1	1	1	1	1	1	1	0
47	1	0	0	0	1	1	0	1	1
48	0	0	0	0	1	1	0	0	1
49	1	0	0	0	1	1	1	0	1
50	0	1	1	1	1	1	0	0	1
51	1	0	0	0	1	1	0	0	1
52	1	1	1	1	1	1	0	0	1
53	1	1	1	1	1	1	0	1	1
54	0	1	1	1	1	0	0	1	1
55	1	0	0	0	1	1	1	0	1
56	1	1	0	0	1	1	1	1	0
57	1	1	1	1	1	0	0	1	1
58	0	1	0	0	1	1	1	0	1
59	1	0	0	0	1	1	1	1	1
60	1	0	1	0	1	1	0	0	1
61	1	0	0	0	1	1	0	0	1
62	0	0	0	1	1	1	0	1	1
63	0	1	1	1	1	1	0	0	1
64	1	1	1	1	1	0	0	1	1
65	1	0	0	0	1	1	0	0	1
66	1	1	1	1	1	1	0	0	1
67	1	1	1	1	1	1	0	0	1
68	0	1	1	1	1	1	0	0	1
69	1	1	1	0	1	1	0	0	1
70	0	0	0	0	0	1	0	0	1

TABLE 1-continued

71	1	1	1	1	1	0	0	1	1
72	1	1	0	0	1	1	1	1	1
73	1	1	1	1	0	1	1	0	1
74	1	1	0	0	1	1	0	1	1
75	1	0	0	0	1	1	0	1	1
76	1	0	0	0	1	1	1	0	1
77	1	1	1	1	1	1	0	1	0
78	1	1	1	1	1	0	1	0	1
79	1	1	0	0	1	1	0	1	1
80	1	1	1	1	0	1	0	1	1
81	1	1	1	1	1	1	0	0	1
82	1	1	0	0	1	1	0	1	0
83	1	1	0	0	1	1	1	1	0
84	1	1	1	1	1	1	0	1	1
85	1	0	0	0	1	1	0	0	0
86	0	1	0	0	1	1	0	0	1
87	1	1	0	0	1	1	1	1	1
88	1	1	0	1	1	0	0	1	1
89	1	1	0	0	1	1	0	1	1
90	1	0	0	0	1	1	0	0	1
91	1	0	0	1	1	1	1	1	1
92	0	1	1	1	1	0	1	1	1
93	1	1	0	0	1	1	1	1	1
94	1	1	1	1	1	1	1	1	1
95	1	1	0	0	0	1	0	0	0
96	1	0	0	0	1	1	1	0	1
97	1	1	0	1	0	1	0	0	0
98	1	0	0	0	1	1	1	1	1
99	1	1	0	1	1	1	1	1	0
100	0	1	0	0	1	1	0	0	1
101	1	1	0	0	1	1	0	1	0
102	0	0	1	1	1	1	0	1	1
103	1	1	0	0	0	1	0	0	0
104	1	1	1	1	1	0	1	1	0
105	1	1	0	0	1	1	0	0	1
106	1	1	0	0	0	1	1	1	0
107	1	1	0	0	0	1	0	1	0
108	1	0	0	1	1	1	0	1	1
109	1	0	0	0	1	1	1	0	1
110	1	0	0	0	1	1	0	0	1
111	0	0	1	1	1	1	0	0	1
112	1	1	1	1	1	0	1	1	1
113	1	1	1	1	1	1	0	1	0
114	1	1	1	1	0	1	0	1	1
115	1	1	1	1	1	0	0	1	1
116	1	1	0	0	1	1	1	0	1
117	0	1	0	0	1	1	0	0	1
118	1	0	0	0	1	1	0	0	1
119	1	1	1	1	1	1	1	1	0
120	0	0	0	1	1	1	0	1	1
121	1	1	1	1	0	1	0	1	0
122	1	1	1	1	1	1	1	1	0
123	1	1	0	1	1	1	1	1	0
124	1	1	0	0	1	1	1	1	0
125	1	1	0	0	1	1	0	1	1
126	1	1	0	0	0	1	0	0	0
127	1	1	0	0	1	1	1	0	1
128	1	1	1	1	1	0	0	1	1
129	1	0	0	0	1	1	1	1	1
130	1	1	1	1	0	1	0	1	0
131	1	1	1	1	1	1	1	1	0
132	1	1	0	1	1	1	1	1	0
133	1	0	0	0	1	1	0	0	1
134	0	0	1	1	1	0	0	1	1
135	1	1	0	1	1	1	0	0	1
136	1	1	0	0	1	1	0	1	0
137	1	1	1	1	1	1	0	1	0
138	1	0	1	1	1	1	1	1	1
139	1	1	1	1	1	1	1	1	1
140	1	1	0	0	1	1	1	1	0
141	1	0	0	0	1	1	0	0	1
142	1	1	1	1	1	1	1	1	0
143	1	0	0	0	1	1	0	1	1
144	0	0	0	1	1	1	0	1	0
145	1	1	0	0	1	1	0	0	0
146	1	1	1	1	0	1	0	0	0
147	1	1	0	0	0	1	0	0	0
148	1	1	1	1	0	1	0	1	0
149	1	1	0	0	1	1	0	0	1
150	1	1	0	0	1	1	0	1	1

TABLE 1-continued

151	1	1	1	0	1	1	0	0	1
152	1	0	0	0	1	1	1	0	1
153	1	0	0	0	1	1	0	0	1
154	0	0	0	1	1	1	1	0	0
155	1	0	1	0	1	1	0	1	1
156	1	1	0	0	1	1	0	0	1
157	1	1	0	0	1	1	0	1	0
158	1	1	0	0	1	1	0	0	0
159	1	1	0	0	0	1	0	1	0
160	1	0	0	0	1	1	0	0	1
161	1	1	1	1	1	1	0	1	0
162	0	0	1	1	1	1	0	0	0
163	1	1	1	1	1	1	1	1	0

1 = in the panel;  
0 = not in the panel.

The one hundred best random panels of proteins out of the million generated are shown in Table 2.

TABLE 2

	Protein 1	Protein 2	Protein 3	Protein 4	Protein 5	Protein 6	Protein 7	Protein 8	Protein 9	Protein 10
1	IBP3	TSP1	CO6A3	PDIA3	SEM3G	SAA	6PGD	EF1A1	PRDX1	TERA
2	EPHB6	CNTN1	CLUS	IBP3	BGH3	6PGD	FRIL	LRP1	TBB3	ERO1A
3	PPIB	LG3BP	MDHC	DSG2	BST1	CD14	DESP	PRDX1	CDCP1	MMP9
4	TPIS	CO1A1	IBP3	GGH	ISLR	MMP2	AIFM1	DSG2	1433T	CBPB2
5	TPIS	IBP3	CH10	SEM3G	6PGD	FRIL	ICAM3	TERA	FINC	ERO1A
6	BGH3	ICAM1	MMP12	6PGD	CD14	EF1A1	HYOU1	PLXC1	PROF1	ERO1A
7	KIT	LG3BP	TPIS	IBP3	LDHB	GGH	TCPA	ISLR	CBPB2	EF1A1
8	LG3BP	IBP3	LDHB	TSP1	CRP	ZA2G	CD14	LRP1	PLIN2	ERO1A
9	CO1A1	TSP1	ISLR	TFR1	CBPB2	FRIL	LRP1	UGPA	PTPA	ERO1A
10	CO6A3	SEM3G	APOE	FRIL	ICAM3	PRDX1	EF2	HS90B	NCF4	PTPA
11	PPIB	LG3BP	CO1A1	APOA1	DSG2	APOE	CD14	PLXC1	NCF4	GSLG1
12	SODM	EPHB6	C163A	CO1A1	LDHB	TETN	1433T	CD14	PTPA	ERO1A
13	SODM	KPYM	IBP3	TSP1	BGH3	SEM3G	6PGD	CD14	RAP2B	EREG
14	EPHB6	ALDOA	MMP7	CO1A1	TIMP1	GRP78	MMP12	CBPB2	G3P	PTPA
15	KIT	TSP1	SCF	TIMP1	OSTP	PDIA3	GRP78	TNF12	PRDX1	PTPA
16	IBP2	LG3BP	GELS	HPT	FIBA	GGH	ICAM1	BST1	HYOU1	GSLG1
17	KIT	CD44	CH10	PEDF	ICAM1	6PGD	S10A1	ERO1A	GSTP1	MMP9
18	LG3BP	C163A	GGH	ERBB3	TETN	BGH3	ENOA	GDIR2	LRP1	ERO1A
19	SODM	KPYM	BGH3	FOLH1	6PGD	DESP	LRP1	TBA1B	ERO1A	GSTP1
20	CNTN1	TETN	ICAM1	K1C19	ZA2G	6PGD	EF2	RAN	ERO1A	GSTP1
21	GELS	ENPL	OSTP	PEDF	ICAM1	BST1	TNF12	GDIR2	LRP1	ERO1A
22	KIT	LDHA	IBP3	PEDF	DSG2	FOLH1	CD14	LRP1	UGPA	ERO1A
23	KIT	TSP1	ISLR	BGH3	COF1	PTPRJ	6PGD	LRP1	S10A6	MPRI
24	LG3BP	C163A	GGH	DSG2	ICAM1	6PGD	GDIR2	HYOU1	EREG	ERO1A
25	IBP2	C163A	ENPL	FIBA	BGH3	CERU	6PGD	LRP1	PRDX1	MMP9
26	LG3BP	C163A	TENX	PDIA3	SEM3G	BST1	VTNC	FRIL	PRDX1	ERO1A
27	ALDOA	CO1A1	TETN	1433T	CBPB2	CD14	G3P	CD59	ERO1A	MMP9
28	IBP3	TENX	CRP	TETN	MMP2	SEM3G	VTNC	CD14	PROF1	ERO1A
29	SODM	EPHB6	TPIS	TENX	ERBB3	SCF	TETN	FRIL	LRP1	ERO1A
30	LG3BP	IBP3	POSTN	DSG2	MDHM	1433Z	CD14	EF1A1	PLXC1	ERO1A
31	IBP2	LG3BP	CO1A1	CNTN1	IBP3	POSTN	TETN	BGH3	6PGD	ERO1A
32	PVR	TSP1	GGH	CYTB	AIFM1	ICAM1	MDHM	1433Z	6PGD	FRIL
33	LYOX	GELS	CO1A1	IBP3	AIFM1	ICAM1	FRIL	PRDX1	RAP2B	NCF4
34	KIT	AMPN	TETN	TNF12	6PGD	FRIL	LRP1	EF2	ERO1A	MMP9
35	LG3BP	GELS	CO1A1	CLUS	CALU	AIFM1	1433T	CD14	UGPA	S10A1
36	ALDOA	IBP3	TSP1	TETN	SEM3G	ICAM1	EF1A1	G3P	RAP2B	NCF4
37	ALDOA	CO1A1	CH10	TETN	PTPRJ	SEM3G	1433T	6PGD	FRIL	ERO1A
38	LG3BP	CO1A1	PLSL	FIBA	TENX	POSTN	CD14	LRP1	NCF4	ERO1A
39	LUM	IBP3	CH10	AIFM1	MDHM	6PGD	PLXC1	EF2	CD59	GSTP1
40	SODM	LG3BP	LUM	LDHA	MDHC	GGH	ICAM1	LRP1	TBA1B	ERO1A
41	LG3BP	CD44	IBP3	CALU	CERU	1433T	CD14	CLIC1	NCF4	ERO1A
42	LG3BP	TPIS	CO1A1	HPT	FIBA	AIFM1	1433Z	6PGD	CD14	EF2
43	ALDOA	CD44	MMP2	CD14	FRIL	PRDX1	RAN	NCF4	MPRI	PTPA
44	CO1A1	CLUS	OSTP	ICAM1	1433T	PLXC1	PTGIS	RAP2B	PTPA	GSTP1
45	KIT	LYOX	IBP3	GRP78	FOLH1	MASP1	CD14	LRP1	ERO1A	GSTP1
46	LG3BP	GGH	CRP	SCF	ICAM1	ZA2G	1433T	RAN	NCF4	ERO1A
47	LG3BP	C163A	BGH3	MMP2	GRP78	LRP1	RAN	ITA5	HS90B	PTPA
48	ALDOA	CLUS	TENX	ICAM1	K1C19	MASP1	6PGD	CBPB2	PRDX1	PTPA
49	IBP3	PDIA3	PEDF	FOLH1	ICAM1	NRP1	6PGD	UGPA	RAN	ERO1A
50	ENPL	FIBA	ISLR	SAA	6PGD	PRDX1	EF2	PLIN2	HS90B	GSLG1
51	LG3BP	CO1A1	CO6A3	GGH	ERBB3	FOLH1	ICAM1	RAN	CDCP1	ERO1A
52	GELS	ENPL	A1AG1	SCF	COF1	ICAM1	6PGD	RAP2B	EF2	HS90B
53	SODM	IBP2	CO1A1	CLUS	IBP3	ENPL	PLSL	TNF12	6PGD	ERO1A
54	KIT	MMP7	CO1A1	TSP1	CO6A3	GGH	PDIA3	ICAM1	LRP1	GSLG1

TABLE 2-continued

	Protein 1	Protein 2	Protein 3	Protein 4	Protein 5	Protein 6	Protein 7	Protein 8	Protein 9	Protein 10
55	ALDOA	COIA1	TSP1	CH10	NRP1	CD14	DESP	LRP1	CLIC1	ERO1A
56	C163A	GELS	CALU	A1AG1	AIFM1	DSG2	ICAM1	6PGD	RAP2B	NCF4
57	PPIB	LG3BP	IBP3	TSP1	PLSL	GRP78	FOLH1	6PGD	HYOU1	RAP2B
58	KIT	LG3BP	LUM	GELS	OSTP	ICAM1	CD14	EF1A1	NCF4	MMP9
59	KIT	PPIB	LG3BP	GELS	FOLH1	ICAM1	MASP1	GDIR2	ITA5	NCF4
60	IBP3	ENPL	ERBB3	BGH3	VTNC	6PGD	EF1A1	TBA1B	S10A6	HS90B
61	LG3BP	CLUS	IBP3	SCF	TCPA	ISLR	GRP78	6PGD	ERO1A	GSTP1
62	LG3BP	LEG1	GELS	GGH	TETN	ENOA	ICAM1	MASP1	FRIL	NCF4
63	LG3BP	CD44	TETN	BGH3	G3P	LRP1	PRDX1	CDCP1	PTPA	MMP9
64	CALU	ENPL	ICAM1	VTNC	FRIL	LRP1	PROF1	TBB3	GSLG1	ERO1A
65	PPIB	PLSL	TENX	A1AG1	COF1	6PGD	FRIL	LRP1	CLIC1	ERO1A
66	IBP2	IBP3	CERU	ENOA	6PGD	CD14	LRP1	PDGFB	ERO1A	GSTP1
67	COIA1	1433T	CD14	DESP	GDIR2	PLXC1	PROF1	RAP2B	RAN	ERO1A
68	LYOX	OSTP	TETN	SEM3G	ICAM1	ZA2G	FRIL	EREG	RAN	ERO1A
69	LG3BP	IBP3	TSP1	PEDF	FOLH1	MDHM	TNF12	NRP1	S10A6	RAP2B
70	KIT	ALDOA	LG3BP	COIA1	TSP1	A1AG1	BGH3	SEM3G	FOLH1	RAN
71	ALDOA	OSTP	BST1	CD14	G3P	PRDX1	PTGIS	FINC	PTPA	MMP9
72	EPHB6	TETN	PEDF	ICAM1	APOE	PROF1	UGPA	NCF4	GSLG1	PTPA
73	LG3BP	COIA1	ENPL	MMP2	1433T	EF1A1	LRP1	HS90B	GSLG1	ERO1A
74	KIT	IBP3	CYTB	MMP2	1433Z	6PGD	CLIC1	EF2	NCF4	PTPA
75	SODM	LYOX	IBP3	TETN	SEM3G	CD14	PRDX1	PTPA	ERO1A	GSTP1
76	SODM	KPYM	COIA1	MDHC	TCPA	CD14	FRIL	LRP1	EF2	ERO1A
77	PPIB	LG3BP	FIBA	GRP78	AIFM1	ICAM1	6PGD	NCF4	GSLG1	PTPA
78	LG3BP	C163A	PVR	MDHC	TETN	SEM3G	AIFM1	6PGD	EREG	ERO1A
79	GELS	ISLR	BGH3	DSG2	ICAM1	SAA	HYOU1	ICAM3	PTGIS	RAP2B
80	KPYM	TPIS	IBP3	TIMP1	GRP78	ICAM1	LRP1	TERA	ERO1A	MMP9
81	IBP3	HPT	TSP1	GRP78	SAA	MMP12	1433Z	6PGD	CD14	S10A6
82	TENX	A1AG1	ENOA	AIFM1	6PGD	CD14	FRIL	LRP1	RAP2B	CD59
83	ALDOA	KPYM	ISLR	TETN	BGH3	VTNC	LRP1	ITA5	PTPA	MMP9
84	SODM	TENX	ISLR	TETN	VTNC	6PGD	LRP1	EF2	ERO1A	MMP9
85	LG3BP	C163A	COIA1	FOLH1	CD14	LRP1	TBA1B	GSLG1	ERO1A	GSTP1
86	SODM	PVR	COIA1	ISLR	PDIA3	APOE	CD14	FRIL	LRP1	CDCP1
87	ALDOA	PEDF	ICAM1	6PGD	CD14	FINC	RAN	NCF4	GSLG1	PTPA
88	LG3BP	KPYM	GELS	COIA1	IBP3	CD14	EF1A1	PLIN2	HS90B	ERO1A
89	LG3BP	PVR	CLUS	TETN	COF1	SEM3G	DESP	EF2	HS90B	ERO1A
90	LG3BP	COIA1	FIBA	TETN	TFR1	ICAM1	MDHM	CD14	PLXC1	ERO1A
91	PPIB	LG3BP	GELS	CLUS	TENX	ICAM1	SAA	NCF4	PTPA	ERO1A
92	COIA1	TSP1	ISLR	BGH3	SAA	6PGD	LRP1	PROF1	EREG	ERO1A
93	CALU	FIBA	OSTP	ISLR	PDIA3	SEM3G	K1C19	6PGD	HYOU1	RAP2B
94	FIBA	CH10	GRP78	SEM3G	AIFM1	ICAM1	MDHM	FRIL	UGPA	GSTP1
95	COIA1	IBP3	PDIA3	ICAM1	K1C19	CD14	EF1A1	FRIL	PTGIS	PDGFB
96	LG3BP	C163A	COIA1	LDHA	1433T	1433Z	FRIL	LRP1	ERO1A	MMP9
97	LG3BP	GELS	COIA1	GRP78	SEM3G	FRIL	PLXC1	PROF1	S10A1	ERO1A
98	LG3BP	COIA1	ENPL	GRP78	AIFM1	ICAM1	1433Z	CD14	LRP1	ERO1A
99	COIA1	PLSL	NRP1	1433T	CD14	FRIL	LRP1	RAP2B	PDGFB	ERO1A
100	IBP2	COIA1	TETN	DSG2	FOLH1	1433T	CD14	FRIL	LRP1	ERO1A

Preferred panels for ruling in treatment for a subject include the panels listed on Table 3 and 4. In various other embodiments, the panels according to the invention include measuring at least 2, 3, 4, 5, 6, 7, or more of the proteins listed on Tables 2 and 3.

TABLE 3

Average (19)	Rule-out (20)	Rule-in (16)
ERO1A	ERO1A	ERO1A
6PGD	6PGD	6PGD
FRIL	FRIL	FRIL
GSTP1	GSTP1	GSTP1
COIA1	COIA1	COIA1
GGH	GGH	GGH
PRDX1	PRDX1	PRDX1
LRP1	CD14	SEM3G
ICAM1	LRP1	GRP78
CD14	LG3BP	TETN
LG3BP	PTPA	AIFM1
PTPA	ICAM1	TSP1
TETN	TSP1	MPRI
GRP78	IBP3	TNF12
AIFM1	FOLH1	MMP9
SEM3G	SODM	OSTP
BGH3	FIBA	

TABLE 3-continued

Average (19)	Rule-out (20)	Rule-in (16)
PDIA3	GSLG1	
FINC	RAP2B	
	C163A	

TABLE 4

Average (13)	Rule-out (13)	Rule-in (9)
LRP1	LRP1 (	LRP1
BGH3	COIA1	COIA1
COIA1	TETN	TETN
TETN	TSP1	TSP1
TSP1	ALDOA	ALDOA
PRDX1	GRP78	GRP78
PROF1	FRIL	FRIL
GRP78	LG3BP	APOE
FRIL	BGH3	TBB3
LG3BP	ISLR	
CD14	PRDX1	
GGH	FIBA	
AIFM1	GSLG1	

A preferred normalizer panel is listed in Table 5.

TABLE 5

Normalizer (6)
PEDF
MASP1
GELS
LUM
C163A
PTPRJ

The term “pulmonary nodules” (PNs) refers to lung lesions that can be visualized by radiographic techniques. A pulmonary nodule is any nodules less than or equal to three centimeters in diameter. In one example a pulmonary nodule has a diameter of about 0.8 cm to 2 cm.

The term “masses” or “pulmonary masses” refers to lung nodules that are greater than three centimeters maximal diameter.

The term “blood biopsy” refers to a diagnostic study of the blood to determine whether a patient presenting with a nodule has a condition that may be classified as either benign or malignant.

The term “acceptance criteria” refers to the set of criteria to which an assay, test, diagnostic or product should conform to be considered acceptable for its intended use. As used herein, acceptance criteria are a list of tests, references to analytical procedures, and appropriate measures, which are defined for an assay or product that will be used in a diagnostic. For example, the acceptance criteria for the classifier refers to a set of predetermined ranges of coefficients.

The term “average maximal AUC” refers to the methodology of calculating performance. For the present invention, in the process of defining the set of proteins that should be in a panel by forward or backwards selection proteins are removed or added one at a time. A plot can be generated with performance (AUC or partial AUC score on the Y axis and proteins on the X axis) the point which maximizes performance indicates the number and set of proteins the gives the best result.

The term “partial AUC factor or pAUC factor” is greater than expected by random prediction. At sensitivity=0.90 the pAUC factor is the trapezoidal area under the ROC curve from 0.9 to 1.0 Specificity/(0.1\*0.1/2).

The term “incremental information” refers to information that may be used with other diagnostic information to enhance diagnostic accuracy. Incremental information is independent of clinical factors such as including nodule size, age, or gender.

The term “score” or “scoring” refers to the refers to calculating a probability likelihood for a sample. For the present invention, values closer to 1.0 are used to represent the likelihood that a sample is cancer, values closer to 0.0 represent the likelihood that a sample is benign.

The term “robust” refers to a test or procedure that is not seriously disturbed by violations of the assumptions on which it is based. For the present invention, a robust test is a test wherein the proteins or transitions of the mass spectrometry chromatograms have been manually reviewed and are “generally” free of interfering signals

The term “coefficients” refers to the weight assigned to each protein used to in the logistic regression equation to score a sample.

In certain embodiments of the invention, it is contemplated that in terms of the logistic regression model of MC CV, the model coefficient and the coefficient of variation (CV) of each

protein’s model coefficient may increase or decrease, dependent upon the method (or model) of measurement of the protein classifier. For each of the listed proteins in the panels, there is about, at least, at least about, or at most about a 2-, 3-, 4-, 5-, 6-, 7-, 8-, 9-, or 10-, -fold or any range derivable therein for each of the coefficient and CV. Alternatively, it is contemplated that quantitative embodiments of the invention may be discussed in terms of as about, at least, at least about, or at most about 10, 20, 30, 40, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99% or more, or any range derivable therein.

The term “best team players” refers to the proteins that rank the best in the random panel selection algorithm, i.e., perform well on panels. When combined into a classifier these proteins can segregate cancer from benign samples. “Best team player” proteins is synonymous with “cooperative proteins”. The term “cooperative proteins” refers proteins that appear more frequently on high performing panels of proteins than expected by chance. This gives rise to a protein’s cooperative score which measures how (in)frequently it appears on high performing panels. For example, a protein with a cooperative score of 1.5 appears on high performing panels 1.5× more than would be expected by chance alone.

The term “classifying” as used herein with regard to a lung condition refers to the act of compiling and analyzing expression data for using statistical techniques to provide a classification to aid in diagnosis of a lung condition, particularly lung cancer.

The term “classifier” as used herein refers to an algorithm that discriminates between disease states with a predetermined level of statistical significance. A two-class classifier is an algorithm that uses data points from measurements from a sample and classifies the data into one of two groups. In certain embodiments, the data used in the classifier is the relative expression of proteins in a biological sample. Protein expression levels in a subject can be compared to levels in patients previously diagnosed as disease free or with a specified condition.

The “classifier” maximizes the probability of distinguishing a randomly selected cancer sample from a randomly selected benign sample, i.e., the AUC of ROC curve.

In addition to the classifier’s constituent proteins with differential expression, it may also include proteins with minimal or no biologic variation to enable assessment of variability, or the lack thereof, within or between clinical specimens; these proteins may be termed endogenous proteins and serve as internal controls for the other classifier proteins.

The term “normalization” or “normalizer” as used herein refers to the expression of a differential value in terms of a standard value to adjust for effects which arise from technical variation due to sample handling, sample preparation and mass spectrometry measurement rather than biological variation of protein concentration in a sample. For example, when measuring the expression of a differentially expressed protein, the absolute value for the expression of the protein can be expressed in terms of an absolute value for the expression of a standard protein that is substantially constant in expression. This prevents the technical variation of sample preparation and mass spectrometry measurement from impeding the measurement of protein concentration levels in the sample.

The term “condition” as used herein refers generally to a disease, event, or change in health status.

The term “treatment protocol” as used herein including further diagnostic testing typically performed to determine whether a pulmonary nodule is benign or malignant. Treat-

ment protocols include diagnostic tests typically used to diagnose pulmonary nodules or masses such as for example, CT scan, positron emission tomography (PET) scan, bronchoscopy or tissue biopsy. Treatment protocol as used herein is also meant to include therapeutic treatments typically used to treat malignant pulmonary nodules and/or lung cancer such as for example, chemotherapy, radiation or surgery.

The terms “diagnosis” and “diagnostics” also encompass the terms “prognosis” and “prognostics”, respectively, as well as the applications of such procedures over two or more time points to monitor the diagnosis and/or prognosis over time, and statistical modeling based thereupon. Furthermore the term diagnosis includes: a. prediction (determining if a patient will likely develop a hyperproliferative disease) b. prognosis (predicting whether a patient will likely have a better or worse outcome at a pre-selected time in the future) c. therapy selection d. therapeutic drug monitoring e. relapse monitoring.

In some embodiments, for example, classification of a biological sample as being derived from a subject with a lung condition may refer to the results and related reports generated by a laboratory, while diagnosis may refer to the act of a medical professional in using the classification to identify or verify the lung condition.

The term “providing” as used herein with regard to a biological sample refers to directly or indirectly obtaining the biological sample from a subject. For example, “providing” may refer to the act of directly obtaining the biological sample from a subject (e.g., by a blood draw, tissue biopsy, lavage and the like). Likewise, “providing” may refer to the act of indirectly obtaining the biological sample. For example, providing may refer to the act of a laboratory receiving the sample from the party that directly obtained the sample, or to the act of obtaining the sample from an archive.

As used herein, “lung cancer” preferably refers to cancers of the lung, but may include any disease or other disorder of the respiratory system of a human or other mammal. Respiratory neoplastic disorders include, for example small cell carcinoma or small cell lung cancer (SCLC), non-small cell carcinoma or non-small cell lung cancer (NSCLC), squamous cell carcinoma, adenocarcinoma, broncho-alveolar carcinoma, mixed pulmonary carcinoma, malignant pleural mesothelioma, undifferentiated large cell carcinoma, giant cell carcinoma, synchronous tumors, large cell neuroendocrine carcinoma, adenocarcinoma, undifferentiated carcinoma; and small cell carcinoma, including oat cell cancer, mixed small cell/large cell carcinoma, and combined small cell carcinoma; as well as adenoid cystic carcinoma, hamartomas, mucoepidermoid tumors, typical carcinoid lung tumors, atypical carcinoid lung tumors, peripheral carcinoid lung tumors, central carcinoid lung tumors, pleural mesotheliomas, and undifferentiated pulmonary carcinoma and cancers that originate outside the lungs such as secondary cancers that have metastasized to the lungs from other parts of the body. Lung cancers may be of any stage or grade. Preferably the term may be used to refer collectively to any dysplasia, hyperplasia, neoplasia, or metastasis in which the protein biomarkers expressed above normal levels as may be determined, for example, by comparison to adjacent healthy tissue.

Examples of non-cancerous lung condition include chronic obstructive pulmonary disease (COPD), benign tumors or masses of cells (e.g., hamartoma, fibroma, neurofibroma), granuloma, sarcoidosis, and infections caused by bacterial (e.g., tuberculosis) or fungal (e.g. histoplasmosis) pathogens. In certain embodiments, a lung condition may be associated with the appearance of radiographic PNs.

As used herein, “lung tissue”, and “lung cancer” refer to tissue or cancer, respectively, of the lungs themselves, as well as the tissue adjacent to and/or within the strata underlying the lungs and supporting structures such as the pleura, intercostal muscles, ribs, and other elements of the respiratory system. The respiratory system itself is taken in this context as representing nasal cavity, sinuses, pharynx, larynx, trachea, bronchi, lungs, lung lobes, aveoli, aveolar ducts, aveolar sacs, aveolar capillaries, bronchioles, respiratory bronchioles, visceral pleura, parietal pleura, pleural cavity, diaphragm, epiglottis, adenoids, tonsils, mouth and tongue, and the like. The tissue or cancer may be from a mammal and is preferably from a human, although monkeys, apes, cats, dogs, cows, horses and rabbits are within the scope of the present invention. The term “lung condition” as used herein refers to a disease, event, or change in health status relating to the lung, including for example lung cancer and various non-cancerous conditions.

“Accuracy” refers to the degree of conformity of a measured or calculated quantity (a test reported value) to its actual (or true) value. Clinical accuracy relates to the proportion of true outcomes (true positives (TP) or true negatives (TN) versus misclassified outcomes (false positives (FP) or false negatives (FN)), and may be stated as a sensitivity, specificity, positive predictive values (PPV) or negative predictive values (NPV), or as a likelihood, odds ratio, among other measures.

The term “biological sample” as used herein refers to any sample of biological origin potentially containing one or more biomarker proteins. Examples of biological samples include tissue, organs, or bodily fluids such as whole blood, plasma, serum, tissue, lavage or any other specimen used for detection of disease.

The term “subject” as used herein refers to a mammal, preferably a human.

The term “biomarker protein” as used herein refers to a polypeptide in a biological sample from a subject with a lung condition versus a biological sample from a control subject. A biomarker protein includes not only the polypeptide itself, but also minor variations thereof, including for example one or more amino acid substitutions or modifications such as glycosylation or phosphorylation.

The term “biomarker protein panel” as used herein refers to a plurality of biomarker proteins. In certain embodiments, the expression levels of the proteins in the panels can be correlated with the existence of a lung condition in a subject. In certain embodiments, biomarker protein panels comprise 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 60, 70, 80, 90 or 100 proteins. In certain embodiments, the biomarker proteins panels comprise from 100-125 proteins, 125-150 proteins, 150-200 proteins or more.

“Treating” or “treatment” as used herein with regard to a condition may refer to preventing the condition, slowing the onset or rate of development of the condition, reducing the risk of developing the condition, preventing or delaying the development of symptoms associated with the condition, reducing or ending symptoms associated with the condition, generating a complete or partial regression of the condition, or some combination thereof.

The term “ruling out” as used herein is meant that the subject is selected not to receive a treatment protocol.

The term “ruling-in” as used herein is meant that the subject is selected to receive a treatment protocol.

Biomarker levels may change due to treatment of the disease. The changes in biomarker levels may be measured by

the present invention. Changes in biomarker levels may be used to monitor the progression of disease or therapy.

“Altered”, “changed” or “significantly different” refer to a detectable change or difference from a reasonably comparable state, profile, measurement, or the like. One skilled in the art should be able to determine a reasonable measurable change. Such changes may be all or none. They may be incremental and need not be linear. They may be by orders of magnitude. A change may be an increase or decrease by 1%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100%, or more, or any value in between 0% and 100%. Alternatively the change may be 1-fold, 1.5-fold 2-fold, 3-fold, 4-fold, 5-fold or more, or any values in between 1-fold and five-fold. The change may be statistically significant with a p value of 0.1, 0.05, 0.001, or 0.0001.

Using the methods of the current invention, a clinical assessment of a patient is first performed. If there exists a higher likelihood for cancer, the clinician may rule in the disease which will require the pursuit of diagnostic testing options yielding data which increase and/or substantiate the likelihood of the diagnosis. “Rule in” of a disease requires a test with a high specificity.

“FN” is false negative, which for a disease state test means classifying a disease subject incorrectly as non-disease or normal.

“FP” is false positive, which for a disease state test means classifying a normal subject incorrectly as having disease.

The term “rule in” refers to a diagnostic test with high specificity that coupled with a clinical assessment indicates a higher likelihood for cancer. If the clinical assessment is a lower likelihood for cancer, the clinician may adopt a stance to rule out the disease, which will require diagnostic tests which yield data that decrease the likelihood of the diagnosis. “Rule out” requires a test with a high sensitivity.

The term “rule out” refers to a diagnostic test with high sensitivity that coupled with a clinical assessment indicates a lower likelihood for cancer.

The term “sensitivity of a test” refers to the probability that a patient with the disease will have a positive test result. This is derived from the number of patients with the disease who have a positive test result (true positive) divided by the total number of patients with the disease, including those with true positive results and those patients with the disease who have a negative result, i.e. false negative.

The term “specificity of a test” refers to the probability that a patient without the disease will have a negative test result. This is derived from the number of patients without the disease who have a negative test result (true negative) divided by all patients without the disease, including those with a true negative result and those patients without the disease who have a positive test result, e.g. false positive. While the sensitivity, specificity, true or false positive rate, and true or false negative rate of a test provide an indication of a test’s performance, e.g. relative to other tests, to make a clinical decision for an individual patient based on the test’s result, the clinician requires performance parameters of the test with respect to a given population.

The term “positive predictive value” (PPV) refers to the probability that a positive result correctly identifies a patient who has the disease, which is the number of true positives divided by the sum of true positives and false positives.

The term “negative predictive value” or “NPV” is calculated by  $TN/(TN+FN)$  or the true negative fraction of all negative test results. It also is inherently impacted by the prevalence of the disease and pre-test probability of the population intended to be tested.

The term “disease prevalence” refers to the number of all new and old cases of a disease or occurrences of an event during a particular period. Prevalence is expressed as a ratio in which the number of events is the numerator and the population at risk is the denominator.

The term disease incidence refers to a measure of the risk of developing some new condition within a specified period of time; the number of new cases during some time period, it is better expressed as a proportion or a rate with a denominator.

Lung cancer risk according to the “National Lung Screening Trial” is classified by age and smoking history. High risk—age  $\geq 55$  and  $\geq 30$  pack-years smoking history; Moderate risk—age  $\geq 50$  and  $\geq 20$  pack-years smoking history; Low risk—age  $\leq 50$  or  $\leq 20$  pack-years smoking history.

The term “negative predictive value” (NPV) refers to the probability that a negative test correctly identifies a patient without the disease, which is the number of true negatives divided by the sum of true negatives and false negatives. A positive result from a test with a sufficient PPV can be used to rule in the disease for a patient, while a negative result from a test with a sufficient NPV can be used to rule out the disease, if the disease prevalence for the given population, of which the patient can be considered a part, is known.

The clinician must decide on using a diagnostic test based on its intrinsic performance parameters, including sensitivity and specificity, and on its extrinsic performance parameters, such as positive predictive value and negative predictive value, which depend upon the disease’s prevalence in a given population.

Additional parameters which may influence clinical assessment of disease likelihood include the prior frequency and closeness of a patient to a known agent, e.g. exposure risk, that directly or indirectly is associated with disease causation, e.g. second hand smoke, radiation, etc., and also the radiographic appearance or characterization of the pulmonary nodule exclusive of size. A nodule’s description may include solid, semi-solid or ground glass which characterizes it based on the spectrum of relative gray scale density employed by the CT scan technology.

“Mass spectrometry” refers to a method comprising employing an ionization source to generate gas phase ions from an analyte presented on a sample presenting surface of a probe and detecting the gas phase ions with a mass spectrometer.

The technology liquid chromatography selected reaction monitoring mass spectrometry (LC-SRM-MS) was used to assay the expression levels of a cohort of 388 proteins in the blood to identify differences for individual proteins which may correlate with the absence or presence of the disease. The individual proteins have not only been implicated in lung cancer biology, but are also likely to be present in plasma based on their expression as membrane-anchored or secreted proteins. An analysis of epithelial and endothelial membranes of resected lung cancer tissues (including the subtypes of adenocarcinoma, squamous, and large cell) identified 217 tissue proteins. A review of the scientific literature with search terms relevant to lung cancer biology identified 319 proteins. There was an overlap of 148 proteins between proteins identified by cancer tissue analysis or literature review, yielding a total of 388 unique proteins as candidates. The majority of candidate proteins included in the multiplex LC-SRM-MS assay were discovered following proteomics analysis of secretory vesicle contents from fresh NSCLC resections and from adjacent non-malignant tissue. The secretory proteins reproducibly upregulated in the tumor tissue were identified and prioritized for inclusion in the LC-SRM-MS assay using extensive bioinformatic and literature

annotation. An additional set of proteins that were present in relevant literature was also added to the assay. In total, 388 proteins associated with lung cancer were prioritized for

SRM assay development. Of these, 371 candidate protein biomarkers were ultimately included in the assay. These are listed in Table 6, below.

TABLE 6

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
1433B_HUMAN	14-3-3 protein beta/alpha	YWHAB	Secreted, EPI	LungCancers	Cytoplasm. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Literature, Detection
1433E_HUMAN	14-3-3 protein epsilon	YWHAE	ENDO	LungCancers, Benign-Nodules	Cytoplasm (By similarity). Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Literature, Detection
1433S_HUMAN	14-3-3 protein sigma	SFN	Secreted, EPI	LungCancers	Cytoplasm. Nucleus (By similarity). Secreted. Note = May be secreted by a non-classical secretory pathway.	UniProt, Literature, Detection
1433T_HUMAN	14-3-3 protein theta	YWHAQ	EPI	LungCancers, Benign-Nodules	Cytoplasm. Note = In neurons, axonally transported to the nerve terminals.	Detection
1433Z_HUMAN	14-3-3 protein zeta/delta	YWHAZ	EPI	LungCancers, Benign-Nodules	Cytoplasm. Melanosome. Note = Located to stage I to stage IV melanosomes.	Detection
6PGD_HUMAN	6-phosphogluconate dehydrogenase, decarboxylating	PGD	EPI, ENDO		Cytoplasm (By similarity).	Detection
A1AG1_HUMAN	Alpha-1-acid glycoprotein 1	ORM1	EPI	Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
ABCD1_HUMAN	ATP-binding cassette sub-family D member 1	ABCD1	ENDO		Peroxisome membrane; Multi-pass membrane protein.	Detection, Prediction
ADA12_HUMAN	Disintegrin and metallo-proteinase domain-containing protein 12	ADAM12		LungCancers, Benign-Nodules, Symptoms	Isoform 1: Cell membrane; Single-pass type I membrane protein. Isoform 2: Secreted. Isoform 3: Secreted (Potential). Isoform 4: Secreted (Potential).	UniProt, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
ADML_HUMAN	ADM	ADM		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
AGR2_HUMAN	Anterior gradient protein 2 homolog	AGR2	EPI	LungCancers	Secreted. Endoplasmic reticulum (By similarity).	UniProt, Prediction
AIFM1_HUMAN	Apoptosis-inducing factor 1, mitochondrial	AIFM1	EPI, ENDO	LungCancers	Mitochondrion inter-membrane space. Nucleus. Note = Translocated to the nucleus upon induction of apoptosis.	Detection, Prediction
ALDOA_HUMAN	Fructose-bisphosphate aldolase A	ALDOA	Secreted, EPI	LungCancers, Symptoms		Literature, Detection
AMPN_HUMAN	Aminopeptidase N	ANPEP	EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Cell membrane; Single-pass type II membrane protein. Cytoplasm, cytosol (Potential). Note = A soluble form has also been detected.	UniProt, Detection
ANGP1_HUMAN	Angiopoietin-1	ANGPT1		LungCancers, Benign-Nodules	Secreted.	UniProt, Literature, Prediction
ANGP2_HUMAN	Angiopoietin-2	ANGPT2		LungCancers, Benign-Nodules	Secreted.	UniProt, Literature, Prediction
APOA1_HUMAN	Apolipoprotein A-I	APOA1		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
APOE_HUMAN	Apolipoprotein E	APOE	EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
ASM3B_HUMAN	Acid sphingomyelinase-like phosphodiesterase 3b	SMPDL3B	EPI, ENDO		Secreted (By similarity).	UniProt, Prediction
AT2A2_HUMAN	Sarcoplasmic/endoplasmic reticulum calcium ATPase 2	ATP2A2	EPI, ENDO	LungCancers, Benign-Nodules	Endoplasmic reticulum membrane; Multi-pass membrane protein. Sarcoplasmic reticulum membrane; Multi-pass membrane protein.	Detection
ATS1_HUMAN	A disintegrin and metalloproteinase with thrombospondin motifs 1	ADAMTS1		LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix (By similarity).	UniProt, Literature, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
ATS12_HUMAN	A disintegrin and metallo-proteinase with thrombospondin motifs 12	ADAMTS12		LungCancers	Secreted, extracellular space, extracellular matrix (By similarity).	UniProt, Detection, Prediction
ATS19_HUMAN	A disintegrin and metallo-proteinase with thrombospondin motifs 19	ADAMTS19		LungCancers	Secreted, extracellular space, extracellular matrix (By similarity).	UniProt, Prediction
BAGE1_HUMAN	B melanoma antigen 1	BAGE		LungCancers	Secreted (Potential).	UniProt, Prediction
BAGE2_HUMAN	B melanoma antigen 2	BAGE2		LungCancers	Secreted (Potential).	UniProt, Prediction
BAGE3_HUMAN	B melanoma antigen 3	BAGE3		LungCancers	Secreted (Potential).	UniProt, Prediction
BAGE4_HUMAN	B melanoma antigen 4	BAGE4		LungCancers	Secreted (Potential).	UniProt, Prediction
BAGE5_HUMAN	B melanoma antigen 5	BAGE5		LungCancers	Secreted (Potential).	UniProt, Prediction
BASP1_HUMAN	Brain acid soluble protein 1	BASP1	Secreted, EPI		Cell membrane; Lipid-anchor. Cell projection, growth cone. Note = Associated with the membranes of growth cones that form the tips of elongating axons.	Detection
BAX_HUMAN	Apoptosis regulator BAX	BAX	EPI	LungCancers, Benign-Nodules	Isoform Alpha: Mitochondrion membrane; Single-pass membrane protein. Cytoplasm. Note = Colocalizes with 14-3-3 proteins in the cytoplasm. Under stress conditions, redistributes to the mitochondrion membrane through the release from JNK-phosphorylated 14-3-3 proteins.    Isoform Beta: Cytoplasm.    Isoform Gamma: Cytoplasm.    Isoform Delta: Cytoplasm (Potential).	UniProt, Literature, Prediction
BDNF_HUMAN	Brain-derived neurotrophic factor	BDNF		Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
BGH3_HUMAN	Transforming growth factor-beta-induced protein igh3	TGFBI		LungCancers, Benign-Nodules	Secreted, extracellular space, extracellular matrix. Note = May be associated both with microfibrils and with the cell surface.	UniProt, Detection
BMP2_HUMAN	Bone morphogenetic protein 2	BMP2		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature
BST1_HUMAN	ADP-ribosyl cyclase 2	BST1	EPI	Symptoms	Cell membrane; Lipid-anchor, GPI-anchor.	Detection, Prediction
C163A_HUMAN	Scavenger receptor cysteine-rich type 1 protein M130	CD163	EPI	Symptoms	Soluble CD163: Secreted.  Cell membrane; Single-pass type I membrane protein. Note = Isoform 1 and isoform 2 show a lower surface expression when expressed in cells.	UniProt, Detection
C4BPA_HUMAN	C4b-binding protein alpha chain	C4BPA		LungCancers, Symptoms	Secreted.	UniProt, Detection, Prediction
CAH9_HUMAN	Carbonic anhydrase 9	CA9		LungCancers, Benign-Nodules, Symptoms	Nucleus. Nucleus, nucleolus. Cell membrane; Single-pass type I membrane protein. Cell projection, microvillus membrane; Single-pass type I membrane protein. Note = Found on the surface microvilli and in the nucleus, particularly in nucleolus.	UniProt
CALR_HUMAN	Calreticulin	CALR	EPI	Symptoms	Endoplasmic reticulum lumen. Cytoplasm, cytosol. Secreted, extracellular space, extracellular matrix. Cell surface. Note = Also found in cell surface (T cells), cytosol and extracellular	UniProt, Literature, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
CALU_HUMAN	Calumenin	CALU	EPI	Symptoms	matrix. Associated with the lytic granules in the cytolytic T-lymphocytes. Endoplasmic reticulum lumen. Secreted. Melanosome. Sarcoplasmic reticulum lumen (By similarity). Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	UniProt, Detection, Prediction
CALX_HUMAN	Calnexin	CANX	Secreted, EPI, ENDO	Benign-Nodules	Endoplasmic reticulum membrane; Single-pass type I membrane protein. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	UniProt, Literature, Detection
CAP7_HUMAN	Azurocidin	AZU1	EPI	Symptoms	Cytoplasmic granule. Note = Cytoplasmic granules of neutrophils.	Prediction
CATB_HUMAN	Cathepsin B	CTSB	Secreted	LungCancers	Lysosome. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Literature, Detection, Prediction
CATG_HUMAN	Cathepsin G	CTSG	Secreted, ENDO	Benign-Nodules	Cell surface.	Detection, Prediction
CBPB2_HUMAN	Carboxypeptidase B2	CPB2		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Detection, Prediction
CCL22_HUMAN	C-C motif chemokine 22	CCL22		LungCancers, Benign-Nodules	Secreted.	UniProt, Prediction
CD14_HUMAN	Monocyte differentiation antigen CD14	CD14	EPI	LungCancers, Benign-Nodules, Symptoms	Cell membrane; Lipid-anchor, GPI-anchor.	Literature, Detection, Prediction
CD24_HUMAN	Signal transducer CD24	CD24		LungCancers, Benign-Nodules	Cell membrane; Lipid-anchor, GPI-anchor.	Literature
CD2A2_HUMAN	Cyclin-dependent kinase inhibitor 2A, isoform 4	CDKN2A		LungCancers, Benign-Nodules	Cytoplasm. Nucleus.  Nucleus, nucleolus (By similarity).	Literature, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
CD38_HUMAN	ADP-ribosyl cyclase 1	CD38	EPI, ENDO	Symptoms	Membrane; Single-pass type II membrane protein.	UniProt, Literature
CD40L_HUMAN	CD40 ligand	CD40LG		LungCancers, Benign-Nodules, Symptoms	Cell membrane; Single-pass type II membrane protein.  CD40 ligand, soluble form: Secreted.	UniProt, Literature
CD44_HUMAN	CD44 antigen	CD44	EPI	LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature, Detection, Prediction
CD59_HUMAN	CD59 glycoprotein	CD59		LungCancers, Benign-Nodules, Symptoms	Cell membrane; Lipid-anchor, GPI-anchor. Secreted. Note = Soluble form found in a number of tissues.	UniProt, Literature, Detection, Prediction
CD97_HUMAN	CD97 antigen	CD97	EPI, ENDO	Symptoms	Cell membrane; Multi-pass membrane protein.  CD97 antigen sub-unit alpha: Secreted, extracellular space.	UniProt
CDCP1_HUMAN	CUB domain-containing protein 1	CDCP1		LungCancers	Isoform 1: Cell membrane; Single-pass membrane protein (Potential). Note = Shedding may also lead to a soluble peptide.  Isoform 3: Secreted.	UniProt, Prediction
CDK4_HUMAN	Cell division protein kinase 4	CDK4		LungCancers, Symptoms		Literature
CEAM5_HUMAN	Carcinoembryonic antigen related cell adhesion molecule 5	CEACAM5	EPI	LungCancers, Benign-Nodules, Symptoms	Cell membrane; Lipid-anchor, GPI-anchor.	Literature, Prediction
CEAM8_HUMAN	Carcinoembryonic antigen-related cell adhesion molecule 8	CEACAM8	EPI	LungCancers	Cell membrane; Lipid-anchor, GPI-anchor.	Detection, Prediction
CERU_HUMAN	Ceruloplasmin	CP	EPI	LungCancers, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
CH10_HUMAN	10 kDa heat shock protein, mitochondrial	HSPE1	ENDO	LungCancers	Mitochondrion matrix.	Literature, Detection, Prediction
CH60_HUMAN	60 kDa heat shock protein, mitochondrial	HSPD1	Secreted, EPI, ENDO	LungCancers, Symptoms	Mitochondrion matrix.	Literature, Detection

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
CKAP4_HUMAN	Cytoskeleton associated protein 4	CKAP4	EPI, ENDO	LungCancers	Endoplasmic reticulum-Golgi intermediate compartment membrane; Single-pass membrane protein (Potential).	UniProt
CL041_HUMAN	Uncharacterized protein C12orf41	C12orf41	ENDO			Prediction
CLCA1_HUMAN	Calcium-activated chloride channel regulator 1	CLCA1		LungCancers, Benign-Nodules	Secreted, extracellular space. Cell membrane; Peripheral membrane protein; Extracellular side. Note = Protein that remains attached to the plasma membrane appeared to be predominantly localized to microvilli.	UniProt, Prediction
CLIC1_HUMAN	Chloride intracellular channel protein 1	CLIC1	EPI		Nucleus. Nucleus membrane; Single-pass membrane protein (Probable). Cytoplasm. Cell membrane; Single-pass membrane protein (Probable). Note = Mostly in the nucleus including in the nuclear membrane. Small amount in the cytoplasm and the plasma membrane. Exists both as soluble cytoplasmic protein and as membrane protein with probably a single transmembrane domain.	UniProt, Literature, Detection
CLUS_HUMAN	Clusterin	CLU	EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
CMGA_HUMAN	Chromogranin-A	CHGA		LungCancers, Benign-Nodules	Secreted. Note = Neuro endocrine and endocrine secretory granules.	UniProt, Literature, Detection, Prediction
CNTN1_HUMAN	Contactin-1	CNTN1		LungCancers	Isoform 1: Cell membrane; Lipid-anchor, GPI-anchor; Extracellular side. Isoform 2: Cell membrane; Lipid-anchor, GPI-anchor; Extracellular side.	Detection, Prediction
CO4A1_HUMAN	Collagen alpha-1(IV) chain	COL4A1		LungCancers	Secreted, extracellular space, extracellular matrix, basement membrane.	UniProt, Detection, Prediction
CO5A2_HUMAN	Collagen alpha-2(V) chain	COL5A2		LungCancers	Secreted, extracellular space, extracellular matrix (By similarity).	UniProt, Detection, Prediction
CO6A3_HUMAN	Collagen alpha-3(VI) chain	COL6A3	Secreted	Symptoms	Secreted, extracellular space, extracellular matrix (By similarity).	UniProt, Detection, Prediction
COCA1_HUMAN	Collagen alpha-1(XII) chain	COL12A1	ENDO	LungCancers, Symptoms	Secreted, extracellular space, extracellular matrix (By similarity).	UniProt, Prediction
COF1_HUMAN	Cofilin-1	CFL1	Secreted, EPI	LungCancers, Benign-Nodules	Nucleus matrix. Cytoplasm, cytoskeleton. Note = Almost completely in nucleus in cells exposed to heat shock or 10% dimethyl sulfoxide.	Detection, Prediction
CO1A1_HUMAN	Collagen alpha-1(XVIII) chain	COL18A1		LungCancers, Benign-Nodules	Secreted, extracellular space, extracellular matrix (By similarity). Mitochondrion inner membrane.	UniProt, Literature, Detection, Prediction
COX5A_HUMAN	Cytochrome c oxidase subunit 5A, mitochondrial	COX5A	Secreted, ENDO			Prediction
CRP_HUMAN	C-reactive protein	CRP		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
CS051_HUMAN	UPF0470 protein C19orf51	C19orf51	ENDO			Prediction
CSF1_HUMAN	Macrophage colony-stimulating factor 1	CSF1		LungCancers, Benign-Nodules	Cell membrane; Single-pass membrane protein (By similarity).  Processed macrophage colony-	UniProt, Literature, Detection

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
CSF2_HUMAN	Granulocyte macrophage colony-stimulating factor	CSF2		LungCancers, Benign-Nodules	stimulating factor 1: Secreted, extracellular space (By similarity). Secreted.	UniProt, Literature, Prediction
CT085_HUMAN	Uncharacterized protein C20orf85	C20orf85		LungCancers, Benign-Nodules		Prediction
CTGF_HUMAN	Connective tissue growth factor	CTGF		LungCancers, Benign-Nodules	Secreted, extracellular space, extracellular matrix (By similarity). Secreted (By similarity).	UniProt, Literature, Detection, Prediction
CYR61_HUMAN	Protein CYR61	CYR61		LungCancers, Benign-Nodules	Secreted.	UniProt, Prediction
CYTA_HUMAN	Cystatin-A	CSTA		LungCancers	Cytoplasm.	Literature, Detection
CYTB_HUMAN	Cystatin-B	CSTB	Secreted		Cytoplasm. Nucleus.	Literature, Detection
DDX17_HUMAN	Probable ATP-dependent RNA helicase DDX17	DDX17	ENDO	LungCancers, Benign-Nodules	Nucleus.	Detection, Prediction
DEFB1_HUMAN	Beta-defensin 1	DEFB1		LungCancers, Benign-Nodules	Secreted.	UniProt, Prediction
DESP_HUMAN	Desmoplakin	DSP	EPI, ENDO	LungCancers	Cell junction, desmosome. Cytoplasm, cytoskeleton. Note = Inner most portion of the desmosomal plaque.	Detection
DFB4A_HUMAN	Beta-defensin 4A	DEFB4A		LungCancers, Benign-Nodules	Secreted.	UniProt
DHI1L_HUMAN	Hydroxysteroid 11-beta-dehydrogenase 1-like protein	HSD11B1L		LungCancers	Secreted (Potential).	UniProt, Prediction
DMBT1_HUMAN	Deleted in malignant brain tumors 1 protein	DMBT1		LungCancers, Benign-Nodules	Secreted (By similarity). Note = Some isoforms may be membrane-bound. Localized to the luminal aspect of crypt cells in the small intestine. In the colon, seen in the luminal aspect of surface epithelial cells. Formed in the ducts of von Ebner	UniProt, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
DMKN_HUMAN	Dermokine	DMKN		LungCancers	gland, and released into the fluid bathing the taste buds contained in the taste papillae (By similarity). Secreted.	UniProt, Detection, Prediction
DPP4_HUMAN	Dipeptidyl peptidase 4	DPP4	EPI	LungCancers, Benign-Nodules, Symptoms	Dipeptidyl peptidase 4 soluble form: Secreted.  Cell membrane; Single-pass type II membrane protein.	UniProt, Detection
DSG2_HUMAN	Desmoglein-2	DSG2	ENDO	Symptoms	Cell membrane; Single-pass type I membrane protein. Cell junction, desmosome.	UniProt, Detection
DDX39A_HUMAN	ATP-dependent RNA helicase DDX39A	DDX39A	EPI		Nucleus (By similarity).	Prediction
DDX39B_HUMAN	Spliceosome RNA helicase DDX39B	DDX39B	EPI		Nucleus. Nucleus speckle.	Prediction
DYRK2_HUMAN	Dual specificity tyrosine-phosphorylation-regulated kinase 2	DYRK2	ENDO	LungCancers	Cytoplasm. Nucleus. Note = Translocates into the nucleus following DNA damage.	Literature
EDN2_HUMAN	Endothelin-2	EDN2		LungCancers	Secreted.	UniProt, Prediction
EEF1A1_HUMAN	Elongation factor 1-alpha 1	EEF1A1	Secreted, EPI	LungCancers, Benign-Nodules	Cytoplasm.	Detection
EEF1D_HUMAN	Elongation factor 1-delta	EEF1D	Secreted, EPI	LungCancers		Prediction
EEF2_HUMAN	Elongation factor 2	EEF2	Secreted, EPI		Cytoplasm.	Literature, Detection
EGF_HUMAN	Pro-epidermal growth factor	EGF		LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature
EGFL6_HUMAN	Epidermal growth factor-like protein 6	EGFL6		LungCancers	Secreted, extracellular space, extracellular matrix, basement membrane (By similarity).	UniProt, Detection, Prediction
ENOA_HUMAN	Alpha-enolase	ENO1	Secreted, EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Cytoplasm. Cell membrane. Cytoplasm, myofibril, sarcomere, M-band. Note = Can translocate to the plasma membrane in	Literature, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
ENOG_HUMAN	Gamma-enolase	ENO2	EPI	LungCancers, Symptoms	either the homodimeric (alpha/alpha) or heterodimeric (alpha/gamma) form. ENO1 is localized to the M-band. Isoform MBP-1: Nucleus. Cytoplasm (By similarity). Cell membrane (By similarity). Note = Can translocate to the plasma membrane in	Literature, Detection, Prediction
ENOX2_HUMAN	Ecto-NOX disulfide thiol exchanger 2	ENOX2		LungCancers	either the homodimeric (alpha/alpha) or heterodimeric (alpha/gamma) form (By similarity). Cell membrane. Secreted, extracellular space. Note = Extracellular and plasma membrane-associated.	UniProt, Detection
ENPL_HUMAN	Endoplasmic	HSP90B1	Secreted, EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Endoplasmic reticulum lumen. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Literature, Detection, Prediction
EPHB6_HUMAN	Ephrin type-B receptor 6	EPHB6		LungCancers	Membrane; Single-pass type I membrane protein. Isoform 3: Secreted (Probable).	UniProt, Literature
EPOR_HUMAN	Erythropoietin receptor	EPOR		LungCancers, Benign-Nodules, Symptoms	Cell membrane; Single-pass type I membrane protein. Isoform EPOR-S: Secreted. Note = Secreted and located to the cell surface.	UniProt, Literature, Detection
ERBB3_HUMAN	Receptor tyrosine-protein kinase erbB-3	ERBB3		LungCancers, Benign-Nodules	Isoform 1: Cell membrane; Single-pass type I membrane	UniProt, Literature, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
EREG_HUMAN	Proepiregulin	EREG		LungCancers	protein.  Isoform 2: Secreted. Epiregulin: Secreted, extracellular space.  Proepiregulin: Cell membrane; Single-pass type I membrane protein.	UniProt
ERO1A_HUMAN	ERO1-like protein alpha	ERO1L	Secreted, EPI, ENDO	Symptoms	Endoplasmic reticulum membrane; Peripheral membrane protein; Luminal side. Note = The association with ERP44 is essential for its retention in the endoplasmic reticulum.	Prediction
ESM1_HUMAN	Endothelial cell-specific molecule 1	ESM1		LungCancers, Benign-Nodules	Secreted.	UniProt, Prediction
EZRI_HUMAN	Ezrin	EZR	Secreted	LungCancers, Benign-Nodules	Apical cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection. Cell projection, micro-villus membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Note = Localization to the apical membrane of parietal cells depends on the interaction with MPP5. Localizes to cell extensions and peripheral processes of astrocytes (By similarity). Micro-villar peripheral membrane protein (cytoplasmic side).	Literature, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
F10A1_HUMAN	Hsc70-interacting protein	ST13	EPI		Cytoplasm (By similarity).  Cytoplasm (Probable).	Detection, Prediction
FAM3C_HUMAN	Protein FAM3C	FAM3C	EPI, ENDO		Secreted (Potential).	UniProt, Detection
FAS_HUMAN	Fatty acid synthase	FASN	EPI	LungCancers, Benign-Nodules, Symptoms	Cytoplasm. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Literature, Detection
FCGR1_HUMAN	High affinity immunoglobulin gamma Fc receptor 1	FCGR1A	EPI	LungCancers, Benign-Nodules, Symptoms	Cell membrane; Single-pass type I membrane protein. Note = Stabilized at the cell membrane through interaction with FCER1G.	UniProt
FGF10_HUMAN	Fibroblast growth factor 10	FGF10		LungCancers	Secreted (Potential).	UniProt, Prediction
FGF2_HUMAN	Heparin-binding growth factor 2	FGF2		LungCancers, Benign-Nodules, Symptoms		Literature
FGF7_HUMAN	Keratinocyte growth factor	FGF7		LungCancers, Benign-Nodules	Secreted.	UniProt, Literature, Prediction
FGF9_HUMAN	Glia-activating factor	FGF9		LungCancers	Secreted.	UniProt, Literature, Prediction
FGFR2_HUMAN	Fibroblast growth factor receptor 2	FGFR2		LungCancers, Benign-Nodules	Cell membrane; Single-pass type I membrane protein.  Isoform 14: Secreted.  Isoform 19: Secreted.	UniProt, Literature, Prediction
FGFR3_HUMAN	Fibroblast growth factor receptor 3	FGFR3		LungCancers	Membrane; Single-pass type I membrane protein.	UniProt, Literature, Prediction
FGL2_HUMAN	Fibroblast growth factor 2	FGL2		Benign-Nodules, Symptoms	Secreted.	UniProt, Detection, Prediction
FHIT_HUMAN	Bis(5'-adenosyl)-triphosphatase	FHIT		LungCancers, Benign-Nodules, Symptoms	Cytoplasm.	Literature
FIBA_HUMAN	Fibrinogen alpha chain	FGA		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
FINC_HUMAN	Fibronectin	FN1	Secreted, EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix.	UniProt, Literature, Detection, Prediction
FKBP11_HUMAN	Peptidyl-prolyl cis-trans isomerase FKBP11	FKBP11	EPI, ENDO		Membrane; Single-pass membrane protein (Potential).	UniProt, Prediction
FOLH1_HUMAN	Glutamate carboxypeptidase 2	FOLH1	ENDO	LungCancers, Symptoms	Cell membrane; Single-pass type II membrane	UniProt, Literature

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
FOLR1_HUMAN	Folate receptor alpha	FOLR1		LungCancers	protein. Isoform PSMA': Cytoplasm. Cell membrane; Lipid-anchor, GPI-anchor. Secreted (Probable).	UniProt
FOXA2_HUMAN	Hepatocyte nuclear factor 3-beta	FOXA2		LungCancers	Nucleus.	Detection, Prediction
FP100_HUMAN	Fanconi anemia-associated protein of 100 kDa	C17orf70	ENDO	Symptoms	Nucleus.	Prediction
FRIH_HUMAN	Ferritin heavy chain	FTH1	EPI	LungCancers, Benign-Nodules		Literature, Detection, Prediction
FRIL_HUMAN	Ferritin light chain	FTL	Secreted, EPI, ENDO	Benign-Nodules, Symptoms		Literature, Detection
G3P_HUMAN	Glyceraldehyde-3-phosphate dehydrogenase	GAPDH	Secreted, EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Cytoplasm. Cytoplasm, perinuclear region. Membrane. Note = Postnuclear and Perinuclear regions.	Detection
G6PD_HUMAN	Glucose-6-phosphate 1-dehydrogenase	G6PD	Secreted, EPI	LungCancers, Symptoms		Literature, Detection
G6PI_HUMAN	Glucose-6-phosphate isomerase	GPI	Secreted, EPI	Symptoms	Cytoplasm. Secreted.	UniProt, Literature, Detection
GAS2L1_HUMAN	GAS2-like protein 1	GAS2L1	ENDO		Cytoplasm, cytoskeleton (Probable).	Prediction
GALT2_HUMAN	Polypeptide N-acetylgalactosaminyl-transferase 2	GALNT2	EPI, ENDO		Golgi apparatus, Golgi stack membrane; Single-pass type II membrane protein. Secreted. Note = Resides preferentially in the trans and medial parts of the Golgi stack. A secreted form also exists.	UniProt, Detection
GAS6_HUMAN	Growth arrest-specific protein 6	GAS6		LungCancers	Secreted.	UniProt, Detection, Prediction
GDIR2_HUMAN	Rho GDP-dissociation inhibitor 2	ARHGDI2	EPI		Cytoplasm.	Detection
GELS_HUMAN	Gelsolin	GSN		LungCancers, Benign-Nodules	Isoform 2: Cytoplasm, cytoskeleton. Isoform 1: Secreted.	UniProt, Literature, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
GGH_HUMAN	Gamma-glutamyl hydrolase	GGH		LungCancers	Secreted, extracellular space. Lysosome. Melanosome. Note = While its intracellular location is primarily the lysosome, most of the enzyme activity is secreted. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	UniProt, Detection, Prediction
GPC3_HUMAN	Glypican-3	GPC3		LungCancers, Symptoms	Cell membrane; Lipid-anchor; GPI-anchor; Extracellular side (By similarity).  Secreted glypican-3: Secreted, extracellular space (By similarity). Cytoplasm. Cytoplasmic granule membrane; Peripheral membrane protein; Cytoplasmic side. Note = Primarily cytosolic in the absence of calcium or magnesium ions. Relocates to granules and other membranes in response to elevated calcium and magnesium levels.	UniProt, Literature, Prediction
GRAN_HUMAN	Grancalcin	GCA	EPI		Cytoplasm. Cytoplasmic granule membrane; Peripheral membrane protein; Cytoplasmic side. Note = Primarily cytosolic in the absence of calcium or magnesium ions. Relocates to granules and other membranes in response to elevated calcium and magnesium levels.	Prediction
GREB1_HUMAN	Protein GREB1	GREB1	ENDO		Membrane; Single-pass membrane protein (Potential).	UniProt, Prediction
GREM1_HUMAN	Gremlin-1	GREM1		LungCancers, Benign-Nodules	Secreted (Probable).	UniProt, Prediction
GRP_HUMAN	Gastrin-releasing peptide	GRP		LungCancers, Symptoms	Secreted.	UniProt, Prediction
GRP78_HUMAN	78 kDa glucose-regulated protein	HSPA5	Secreted, EPI, ENDO	LungCancers, Benign-Nodules	Endoplasmic reticulum lumen. Melanosome. Note = Identified by mass spectrometry in melanosome	Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
GSLG1_HUMAN	Golgi apparatus protein 1	GLG1	EPI, ENDO	Benign-Nodules	fractions from stage I to stage IV. Golgi apparatus membrane; Single-pass type I membrane protein.	UniProt
GSTP1_HUMAN	Glutathione S-transferase P	GSTP1	Secreted	LungCancers, Benign-Nodules, Symptoms		Literature, Detection, Prediction
GTR1_HUMAN	Solute carrier family 2, facilitated glucose transporter member 1	SLC2A1	EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Cell membrane; Multi-pass membrane protein (By similarity). Melanosome. Note = Localizes primarily at the cell surface (By similarity). Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Literature
GTR3_HUMAN	Solute carrier family 2, facilitated glucose transporter member 3	SLC2A3	EPI		Membrane; Multi-pass membrane protein.	Detection
H2A1_HUMAN	Histone H2A type 1	HIST1H2AG	Secreted		Nucleus.	Detection, Prediction
H2A1B_HUMAN	Histone H2A type 1-B/E	HIST1H2AB	Secreted		Nucleus.	Detection, Prediction
H2A1C_HUMAN	Histone H2A type 1-C	HIST1H2AC	Secreted		Nucleus.	Literature, Detection, Prediction
H2A1D_HUMAN	Histone H2A type 1-D	HIST1H2AD	Secreted		Nucleus.	Detection, Prediction
HG2A_HUMAN	HLA class II histocompatibility antigen gamma chain	CD74		LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type II membrane protein (Potential).	UniProt, Literature
HGF_HUMAN	Hepatocyte growth factor	HGF		LungCancers, Benign-Nodules, Symptoms		Literature, Prediction
HMGA1_HUMAN	High mobility group protein HMG-I/HMG-Y	HMGA1		LungCancers, Benign-Nodules, Symptoms	Nucleus.	Literature
HPRT_HUMAN	Hypoxanthine guanine phosphoribosyltransferase	HPRT1	EPI		Cytoplasm.	Detection, Prediction
HPSE_HUMAN	Heparanase	HPSE		LungCancers, Benign-Nodules, Symptoms	Lysosome membrane; Peripheral membrane protein. Secreted. Note = Secreted, internalised and	UniProt, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
					transferred to late endosomes/lysosomes as a proheparanase. In lysosomes, it is processed into the active form, the heparanase. The uptake or internalisation of proheparanase is mediated by HSPGs. Heparin appears to be a competitor and retain proheparanase in the extracellular medium.	
HPT_HUMAN	Haptoglobin	HP		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
HS90A_HUMAN	Heat shock protein HSP 90-alpha	HSP90AA1	Secreted, EPI	LungCancers, Symptoms	Cytoplasm. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Literature, Detection
HS90B_HUMAN	Heat shock protein HSP 90-beta	HSP90AB1	Secreted, EPI	LungCancers	Cytoplasm. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Literature, Detection
HSPB1_HUMAN	Heat shock protein beta-1	HSPB1	Secreted, EPI	LungCancers, Benign-Nodules	Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, spindle. Note = Cytoplasmic in interphase cells. Colocalizes with mitotic spindles in mitotic cells. Translocates to the nucleus during heat shock.	Literature, Detection, Prediction
HTRA1_HUMAN	Serine protease HTRA1	HTRA1		LungCancers	Secreted.	UniProt, Prediction
HXK1_HUMAN	Hexokinase-1	HK1	ENDO	Symptoms	Mitochondrion outer membrane. Note = Its hydrophobic N-terminal sequence may be involved in	Literature, Detection

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
HYAL2_HUMAN	Hyaluronidase-2	HYAL2		LungCancers	membrane binding. Cell membrane; Lipid-anchor, GPI-anchor.	Prediction
HYOU1_HUMAN	Hypoxia up-regulated protein 1	HYOU1	EPI, ENDO	Symptoms	Endoplasmic reticulum lumen.	Detection
IBP2_HUMAN	Insulin-like growth factor-binding protein 2	IGFBP2		LungCancers	Secreted.	UniProt, Literature, Detection, Prediction
IBP3_HUMAN	Insulin-like growth factor-binding protein 3	IGFBP3		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
ICAM1_HUMAN	Intercellular adhesion molecule 1	ICAM1		LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature, Detection
ICAM3_HUMAN	Intercellular adhesion molecule 3	ICAM3	EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Detection
IDHP_HUMAN	Isocitrate dehydrogenase [NADP], mitochondrial	IDH2	Secreted, ENDO		Mitochondrion.	Prediction
IF4A1_HUMAN	Eukaryotic initiation factor 4A-I	EIF4A1	Secreted, EPI, ENDO			Detection, Prediction
IGF1_HUMAN	Insulin-like growth factor I	IGF1		LungCancers, Benign-Nodules, Symptoms	Secreted.  Secreted.	UniProt, Literature, Detection, Prediction
IKIP_HUMAN	Inhibitor of nuclear factor kappa-B kinase-interacting protein	IKIP	ENDO	Symptoms	Endoplasmic reticulum membrane; Single-pass membrane protein. Note = Isoform 4 deletion of the hydrophobic, or transmembrane region between AA 45-63 results in uniform distribution throughout the cell, suggesting that this region is responsible for endoplasmic reticulum localization.	UniProt, Prediction
IL18_HUMAN	Interleukin-18	IL18		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Prediction
IL19_HUMAN	Interleukin-19	IL19		LungCancers	Secreted.	UniProt, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
IL22_HUMAN	Interleukin-22	IL22		LungCancers, Benign-Nodules	Secreted.	UniProt, Prediction
IL32_HUMAN	Interleukin-32	IL32		LungCancers, Benign-Nodules	Secreted.	UniProt, Prediction
IL7_HUMAN	Interleukin-7	IL7		LungCancers, Benign-Nodules	Secreted.	UniProt, Literature, Prediction
IL8_HUMAN	Interleukin-8	IL8		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature
ILEU_HUMAN	Leukocyte elastase inhibitor	SERPINB1	Secreted, EPI		Cytoplasm (By similarity).	Detection, Prediction
ILK_HUMAN	Integrin-linked protein kinase	ILK	Secreted	LungCancers, Benign-Nodules, Symptoms	Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side.	Literature, Detection
INHBA_HUMAN	Inhibin beta A chain	INHBA		LungCancers, Benign-Nodules	Secreted.	UniProt, Literature, Prediction
ISLR_HUMAN	Immunoglobulin superfamily containing leucine-rich repeat protein	ISLR		LungCancers	Secreted (Potential).	UniProt, Detection, Prediction
ITA5_HUMAN	Integrin alpha-5	ITGA5	EPI	LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature, Detection
ITAM_HUMAN	Integrin alpha-M	ITGAM	EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature
K0090_HUMAN	Uncharacterized protein KIAA0090	KIAA0090	EPI	Symptoms	Membrane; Single-pass type I membrane protein (Potential).	UniProt, Prediction
K1C18_HUMAN	Keratin, type I cytoskeletal 18	KRT18	Secreted	LungCancers, Benign-Nodules	Cytoplasm, perinuclear region.	Literature, Detection, Prediction
K1C19_HUMAN	Keratin, type I cytoskeletal 19	KRT19		LungCancers, Benign-Nodules		Literature, Detection, Prediction
K2C8_HUMAN	Keratin, type II cytoskeletal 8	KRT8	EPI	LungCancers	Cytoplasm.	Literature, Detection
KIT_HUMAN	Mast/stem cell growth factor receptor	KIT		LungCancers	Membrane; Single-pass type I membrane protein.	UniProt, Literature, Detection
KITH_HUMAN	Thymidine kinase, cytosolic	TK1		LungCancers	Cytoplasm.	Literature, Prediction
KLK11_HUMAN	Kallikrein-11	KLK11		LungCancers	Secreted.	UniProt, Literature, Prediction
KLK13_HUMAN	Kallikrein-13	KLK13		LungCancers	Secreted (Probable).	UniProt, Literature, Detection, Prediction
KLK14_HUMAN	Kallikrein-14	KLK14		LungCancers, Symptoms	Secreted, extracellular space.	UniProt, Literature, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
KLK6_HUMAN	Kallikrein-6	KLK6		LungCancers, Benign-Nodules, Symptoms	Secreted. Nucleus, nucleolus. Cytoplasm. Mitochondrion. Microsome. Note = In brain, detected in the nucleus of glial cells and in the nucleus and cytoplasm of neurons. Detected in the mitochondrial and microsomal fractions of HEK-293 cells and released into the cytoplasm following cell stress.	UniProt, Literature, Detection, Prediction
KNG1_HUMAN	Kininogen-1	KNG1		LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space.	UniProt, Detection, Prediction
KPYM_HUMAN	Pyruvate kinase isozymes M1/M2	PKM2	Secreted, EPI	LungCancers, Symptoms	Cytoplasm. Nucleus. Note = Translocates to the nucleus in response to different apoptotic stimuli. Nuclear translocation is sufficient to induce cell death that is caspase independent, isoform-specific and independent of its enzymatic activity.	Literature, Detection
KRT35_HUMAN	Keratin, type I cuticular Ha5	KRT35	ENDO			Detection, Prediction
LAMB2_HUMAN	Laminin subunit beta-2	LAMB2	ENDO	LungCancers, Symptoms	Secreted, extracellular space, extracellular matrix, basement membrane. Note = S-laminin is concentrated in the synaptic cleft of the neuromuscular junction. Cytoplasm.	UniProt, Detection, Prediction
LDHA_HUMAN	L-lactate dehydrogenase A chain	LDHA	Secreted, EPI, ENDO	LungCancers	Cytoplasm.	Literature, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
LDHB_HUMAN	L-lactate dehydrogenase B chain	LDHB	EPI	LungCancers	Cytoplasm.	Detection, Prediction
LEG1_HUMAN	Galectin-1	LGALS1	Secreted	LungCancers	Secreted, extracellular space, extracellular matrix.	UniProt, Detection
LEG3_HUMAN	Galectin-3	LGALS3		LungCancers, Benign-Nodules	Nucleus. Note = Cytoplasmic in adenomas and carcinomas. May be secreted by a non-classical secretory pathway and associate with the cell surface.	Literature, Detection, Prediction
LEG9_HUMAN	Galectin-9	LGALS9	ENDO	Symptoms	Cytoplasm (By similarity). Secreted (By similarity). Note = May also be secreted by a non-classical secretory pathway (By similarity).	UniProt
LG3BP_HUMAN	Galectin-3-binding protein	LGALS3BP	Secreted	LungCancers, Benign-Nodules, Symptoms	Secreted. Secreted, extracellular space, extracellular matrix.	UniProt, Literature, Detection, Prediction
LPLC3_HUMAN	Long palate, lung and nasal epithelium carcinoma-associated protein 3	C20orf185		LungCancers	Secreted (By similarity). Cytoplasm. Note = According to Pub-Pub-Med: 12837268 it is cytoplasmic.	UniProt, Prediction
LPLC4_HUMAN	Long palate, lung and nasal epithelium carcinoma-associated protein 4	C20orf186		LungCancers	Secreted (By similarity). Cytoplasm.	UniProt, Prediction
LPPRC_HUMAN	Leucine-rich PPR motif-containing protein, mitochondrial	LRPPRC	Secreted, ENDO	LungCancers, Symptoms	Mitochondrion. Nucleus, nucleoplasm. Nucleus inner membrane. Nucleus outer membrane. Note = Seems to be predominantly mitochondrial.	Prediction
LRP1_HUMAN	Prolow-density lipoprotein receptor-related protein 1	LRP1	EPI	LungCancers, Symptoms	Low-density lipoprotein receptor-related protein 1 85 kDa subunit: Cell membrane; Single-pass type I membrane	UniProt, Detection

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
LUM_HUMAN	Lumican	LUM	Secreted, EPI	LungCancers, Benign-Nodules, Symptoms	protein. Membrane, coated pit. Low-density lipoprotein receptor-related protein 1 515 kDa subunit: Cell membrane; Peripheral membrane protein; Extracellular side. Membrane, coated pit. Low-density lipoprotein receptor-related protein 1 intracellular domain: Cytoplasm. Nucleus. Note = After cleavage, the intracellular domain (LRPICD) is detected both in the cytoplasm and in the nucleus.	UniProt, Detection, Prediction
LY6K_HUMAN	Lymphocyte antigen 6K	LY6K		LungCancers, Symptoms	Secreted. Cytoplasm. Cell membrane; Lipid-anchor, GPI-anchor (Potential).	UniProt, Prediction
LYAM2_HUMAN	E-selectin	SELE		LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature, Detection
LYAM3_HUMAN	P-selectin	SELP		LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature, Detection
LYOX_HUMAN	Protein-lysine 6-oxidase	LOX		LungCancers, Benign-Nodules	Secreted, extracellular space.	UniProt, Detection, Prediction
LYPD3_HUMAN	Ly6/PLAUR domain-containing protein 3	LYPD3		LungCancers	Cell membrane; Lipid-anchor, GPI-anchor.	Detection, Prediction
MAGA4_HUMAN	Melanoma-associated antigen 4	MAGEA4		LungCancers		Literature, Prediction
MASP1_HUMAN	Mannan-binding lectin serine protease 1	MASP1		LungCancers, Symptoms	Secreted.	UniProt, Detection, Prediction
MDHC_HUMAN	Malate dehydrogenase, cytoplasmic	MDH1	Secreted		Cytoplasm.	Literature, Detection, Prediction
MDHM_HUMAN	Malate dehydrogenase, mitochondrial	MDH2	ENDO	LungCancers	Mitochondrion matrix.	Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
MIF_HUMAN	Macrophage migration inhibitory factor	MIF	Secreted	LungCancers, Benign-Nodules, Symptoms	Secreted. Cytoplasm. Note = Does not have a cleavable signal sequence and is secreted via a specialized, non-classical pathway. Secreted by macrophages upon stimulation by bacterial lipopolysaccharide (LPS), or by M. tuberculosis antigens.	UniProt, Literature, Prediction
MLH1_HUMAN	DNA mismatch repair protein Mlh1	MLH1	ENDO	LungCancers, Benign-Nodules, Symptoms	Nucleus.	Literature
MMP1_HUMAN	Interstitial collagenase	MMP1		LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix (Probable).	UniProt, Literature, Prediction
MMP11_HUMAN	Stromelysin-3	MMP11		LungCancers, Symptoms	Secreted, extracellular space, extracellular matrix (Probable).	UniProt, Literature, Prediction
MMP12_HUMAN	Macrophage metalloelastase	MMP12		LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix (Probable).	UniProt, Literature, Prediction
MMP14_HUMAN	Matrix metallo-proteinase-14	MMP14	ENDO	LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein (Potential). Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	UniProt, Literature, Detection
MMP2_HUMAN	72 kDa type IV collagenase	MMP2		LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix (Probable).	UniProt, Literature, Detection, Prediction
MMP26_HUMAN	Matrix metallo-proteinase-26	MMP26		LungCancers	Secreted, extracellular space, extracellular matrix.	UniProt, Prediction
MMP7_HUMAN	Matrilysin	MMP7		LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix (Probable).	UniProt, Literature, Prediction
MMP9_HUMAN	Matrix metallo-proteinase-9	MMP9		LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix (Probable).	UniProt, Literature, Detection, Prediction
MOGS_HUMAN	Mannosyl-oligosaccharide glucosidase	MOGS	ENDO		Endoplasmic reticulum membrane; Single-	UniProt, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
MPRI_HUMAN	Cation-independent mannose-6-phosphate receptor	IGF2R	EPI, ENDO	LungCancers, Symptoms	pass type II membrane protein. Lysosome membrane; Single-pass type I membrane protein.	UniProt, Literature, Detection
MRP3_HUMAN	Canalicular multi-specific organic anion transporter 2	ABCC3	EPI	LungCancers	Membrane; Multi-pass membrane protein.	Literature, Detection
MUC1_HUMAN	Mucin-1	MUC1	EPI	LungCancers, Benign-Nodules, Symptoms	Apical cell membrane; Single-pass type I membrane protein. Note = Exclusively located in the apical domain of the plasma membrane of highly polarized epithelial cells. After endocytosis, internalized and recycled to the cell membrane. Located to microvilli and to the tips of long filopodial protusions.  Isoform 5: Secreted.  Isoform 7: Secreted.  Isoform 9: Secreted.  Mucin-1 subunit beta: Cell membrane. Cytoplasm. Nucleus. Note = On EGF and PDGFRB stimulation, transported to the nucleus through interaction with CTNNA1, a process which is stimulated by phosphorylation. On HRG stimulation, colocalizes with JUP/gamma-catenin at the nucleus.	UniProt, Literature, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
MUC16_HUMAN	Mucin-16	MUC16		LungCancers	Cell membrane; Single-pass type I membrane protein. Secreted, extracellular space. Note = May be liberated into the extracellular space following the phosphorylation of the intracellular C-terminus which induces the proteolytic cleavage and liberation of the extracellular domain.	UniProt, Detection
MUC4_HUMAN	Mucin-4	MUC4		LungCancers, Benign-Nodules	Membrane; Single-pass membrane protein (Potential). Secreted. Note = Isoforms lacking the Cys-rich region, EGF-like domains and transmembrane region are secreted. Secretion occurs by splicing or proteolytic processing.  Mucin-4 beta chain: Cell membrane; Single-pass membrane protein.  Mucin-4 alpha chain: creted.  Isoform 3: Cell membrane; Single-pass membrane protein.  Isoform 15: Secreted.	UniProt
MUC5B_HUMAN	Mucin-5B	MUC5B		LungCancers, Benign-Nodules	Secreted.	UniProt, Detection, Prediction
MUCL1_HUMAN	Mucin-like protein 1	MUCL1		LungCancers	Secreted (Probable). Membrane (Probable).	UniProt, Prediction
NAMPT_HUMAN	Nicotinamide phosphoribosyltransferase	NAMPT	EPI	LungCancers, Benign-Nodules, Symptoms	Cytoplasm (By similarity).	Literature, Detection

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
NAPSA_HUMAN	Napsin-A	NAPSA	Secreted	LungCancers		Prediction
NCF4_HUMAN	Neutrophil cytosol factor 4	NCF4	ENDO		Cytoplasm.	Prediction
NDKA_HUMAN	Nucleoside di-phosphate kinase A	NME1	Secreted	LungCancers, Benign-Nodules, Symptoms	Cytoplasm. Nucleus. Note = Cell-cycle dependent nuclear localization which can be induced by interaction with Epstein-barr viral proteins or by degradation of the SET complex by GzmA.	Literature, Detection
NDKB_HUMAN	Nucleoside di-phosphate kinase B	NME2	Secreted, EPI	Benign-Nodules	Cytoplasm. Nucleus. Note = Isoform 2 is mainly cytoplasmic and isoform 1 and isoform 2 are excluded from the nucleolus.	Literature, Detection
NDUS1_HUMAN	NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial	NDUFS1	Secreted, ENDO	Symptoms	Mitochondrion inner membrane.	Prediction
NEBL_HUMAN	Nebulette	NEBL	ENDO			Prediction
NEK4_HUMAN	Serine/threonine protein kinase Nek4	NEK4	ENDO	LungCancers	Nucleus (Probable).	Prediction
NET1_HUMAN	Netrin-1	NTN1		LungCancers, Benign-Nodules	Secreted, extracellular space, extracellular matrix (By similarity).	UniProt, Literature, Prediction
NEU2_HUMAN	Vasopressin neurophysin 2-copeptin	AVP		LungCancers, Symptoms	Secreted.	UniProt, Prediction
NGAL_HUMAN	Neutrophil gelatinase-associated lipocalin	LCN2	EPI	LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Detection, Prediction
NGLY1_HUMAN	Peptide-N(4)-(N-acetyl-beta-glucosaminyl)asparagine amidase	NGLY1	ENDO		Cytoplasm.	Detection, Prediction
NHRF1_HUMAN	Na(+)/H(+) exchange regulatory cofactor NHE-RF1	SLC9A3R1	EPI	Benign-Nodules	Endomembrane system; Peripheral membrane protein. Cell projection, filopodium. Cell projection, ruffle. Cell projection, microvillus.	Detection

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
					Note = Colocalizes with actin in microvilli-rich apical regions of the syncytiotrophoblast. Found in microvilli, ruffling membrane and filopodia of HeLa cells. Present in lipid rafts of T-cells. Cytoplasm.	
NIBAN_HUMAN	Protein Niban	FAM129A	EPI			Literature, Detection
NMU_HUMAN	Neuromedin-U	NMU		LungCancers	Secreted.	UniProt, Prediction
NRP1_HUMAN	Neuropilin-1	NRP1		LungCancers, Benign-Nodules, Symptoms	Cell membrane; Single-pass type I membrane protein. Isoform 1: Secreted. Isoform 2: Secreted (By similarity).	UniProt, Literature, Detection, Prediction
ODAM_HUMAN	Odontogenic ameloblast associated protein	ODAM		LungCancers	Secreted (By similarity).	UniProt, Prediction
OSTP_HUMAN	Osteopontin	SPP1		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
OVOS2_HUMAN	Ovostatin homolog 2	OVOS2	ENDO		Secreted (By similarity).	UniProt, Prediction
P5CS_HUMAN	Delta-1-pyrroline-5-carboxylate synthase	ALDH18A1	ENDO		Mitochondrion inner membrane.	Prediction
PA2GX_HUMAN	Group 10 secretory phospholipase A2	PLA2G10		Symptoms	Secreted.	UniProt
PAPP1_HUMAN	Pappalysin-1	PAPPA		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Prediction
PBIP1_HUMAN	Pre-B-cell leukemia transcription factor-interacting protein 1	PBXIP1	EPI		Cytoplasm, cytoskeleton. Nucleus. Note = Shuttles between the nucleus and the cytosol. Mainly localized in the cytoplasm, associated with microtubules. Detected in small amounts in the nucleus.	Prediction
PCBP1_HUMAN	Poly(rC)-binding protein 1	PCBP1	EPI, ENDO		Nucleus. Cytoplasm. Note = Loosely bound in the nucleus. May shuttle between the	Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
PCBP2_HUMAN	Poly(rC)-binding protein 2	PCBP2	EPI		nucleus and the cytoplasm. Nucleus. Cytoplasm. Note = Loosely bound in the nucleus. May shuttle between the nucleus and the cytoplasm.	Detection, Prediction
PCD15_HUMAN	Protocadherin-15	PCDH15	ENDO		Cell membrane; Single-pass type I membrane protein (By similarity). Isoform 3: Secreted.	UniProt, Detection
PCNA_HUMAN	Proliferating cell nuclear antigen	PCNA	EPI	LungCancers, Benign-Nodules, Symptoms	Nucleus.	Literature, Prediction
PCYOX_HUMAN	Preylcysteine oxidase 1	PCYOX1	Secreted	LungCancers, Symptoms	Lysosome.	Detection, Prediction
PDGFA_HUMAN	Platelet-derived growth factor subunit A	PDGFA		LungCancers	Secreted.	UniProt, Literature, Prediction
PDGFB_HUMAN	Platelet-derived growth factor subunit B	PDGFB		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
PDGFD_HUMAN	Platelet-derived growth factor D	PDGFD		LungCancers	Secreted.	UniProt, Prediction
PDIA3_HUMAN	Protein disulfide-isomerase A3	PDIA3	ENDO	LungCancers	Endoplasmic reticulum lumen (By similarity). Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Detection, Prediction
PDIA4_HUMAN	Protein disulfide-isomerase A4	PDIA4	Secreted, EPI, ENDO		Endoplasmic reticulum lumen. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Detection, Prediction
PDIA6_HUMAN	Protein disulfide-isomerase A6	PDIA6	Secreted, EPI, ENDO		Endoplasmic reticulum lumen (By similarity). Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
PECA1_HUMAN	Platelet endothelial cell adhesion molecule	PECAM1		LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature, Detection
PEDF_HUMAN	Pigment epithelium-derived factor	SERPINF1		LungCancers, Symptoms	Secreted. Melanosome. Note = Enriched in stage I melanosomes.	UniProt, Literature, Detection, Prediction
PERM_HUMAN	Myeloperoxidase	MPO	Secreted, EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Lysosome.	Literature, Detection, Prediction
PERP1_HUMAN	Plasma cell-induced resident endoplasmic reticulum protein	PACAP	EPI, ENDO		Secreted (Potential). Cytoplasm. Note = In (PubMed: 11350957) diffuse granular localization in the cytoplasm surrounding the nucleus.	UniProt, Detection, Prediction
PGAM1_HUMAN	Phosphoglycerate mutase 1	PGAM1	Secreted, EPI	LungCancers, Symptoms		Detection
PLAC1_HUMAN	Placenta-specific protein 1	PLAC1		LungCancers	Secreted (Probable).	UniProt, Prediction
PLACL_HUMAN	Placenta-specific 1-like protein	PLAC1L		LungCancers	Secreted (Potential).	UniProt, Prediction
PLIN2_HUMAN	Perilipin-2	ADFP	ENDO	LungCancers	Membrane; Peripheral membrane protein.	Prediction
PLIN3_HUMAN	Perilipin-3	M6PRBP1	EPI		Cytoplasm. Endosome membrane; Peripheral membrane protein; Cytoplasmic side (Potential). Lipid droplet (Potential). Note = Membrane associated on endosomes. Detected in the envelope and the core of lipid bodies and in lipid sails.	Detection, Prediction
PLOD1_HUMAN	Procollagen-lysine,2-oxoglutarate 5-dioxygenase 1	PLOD1	EPI, ENDO		Rough endoplasmic reticulum membrane; Peripheral membrane protein; Luminal side.	Prediction
PLOD2_HUMAN	Procollagen-lysine,2-oxoglutarate 5-dioxygenase 2	PLOD2	ENDO	Benign-Nodules, Symptoms	Rough endoplasmic reticulum membrane; Peripheral membrane protein; Luminal side.	Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
PLSL_HUMAN	Plastin-2	LCP1	Secreted, EPI	LungCancers	Cytoplasm, cytoskeleton. Cell junction. Cell projection. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side (By similarity). Note = Relocalizes to the immunological synapse between peripheral blood T lymphocytes and antibody-presenting cells in response to costimulation through TCR/CD3 and CD2 or CD28. Associated with the actin cytoskeleton at membrane ruffles (By similarity). Relocalizes to actin-rich cell projections upon serine phosphorylation.	Detection, Prediction
PLUNC_HUMAN	Protein Plunc	PLUNC		LungCancers, Benign-Nodules	Secreted (By similarity). Note = Found in the nasal mucus (By similarity). Apical side of airway epithelial cells. Detected in nasal mucus (By similarity).	UniProt, Prediction
PLXB3_HUMAN	Plexin-B3	PLXNB3	ENDO		Membrane; Single-pass type I membrane protein.	UniProt, Detection, Prediction
PLXC1_HUMAN	Plexin-C1	PLXNC1	EPI		Membrane; Single-pass type I membrane protein (Potential).	UniProt, Detection
POSTN_HUMAN	Periostin	POSTN	Secreted, ENDO	LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix.	UniProt, Literature, Detection, Prediction
PPAL_HUMAN	Lysosomal acid phosphatase	ACP2	EPI	Symptoms	Lysosome membrane; Single-pass membrane protein; Luminal side. Lysosome lumen.	UniProt, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
PPBT_HUMAN	Alkaline phosphatase, tissue-nonspecific isozyme	ALPL	EPI	LungCancers, Benign-Nodules, Symptoms	Note = The soluble form arises by proteolytic processing of the membrane-bound form. Cell membrane; Lipid-anchor, GPI-anchor.	Literature, Detection, Prediction
PPIB_HUMAN	Peptidyl-prolyl cis-trans isomerase B	PPIB	Secreted, EPI, ENDO		Endoplasmic reticulum lumen. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Detection, Prediction
PRDX1_HUMAN	Peroxiredoxin-1	PRDX1	EPI	LungCancers	Cytoplasm. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Detection, Prediction
PRDX4_HUMAN	Peroxiredoxin-4	PRDX4	Secreted, EPI, ENDO		Cytoplasm.	Literature, Detection, Prediction
PROF1_HUMAN	Profilin-1	PFN1	Secreted, EPI	LungCancers	Cytoplasm, cytoskeleton.	Detection
PRP31_HUMAN	U4/U6 small nuclear ribo-nucleo-protein Prp31	PRPF31	ENDO		Nucleus speckle. Nucleus, Cajal body. Note = Predominantly found in speckles and in Cajal bodies.	Prediction
PRSGA_HUMAN	26S protease regulatory subunit 6A	PSMC3	EPI	Benign-Nodules	Cytoplasm (Potential). Nucleus (Potential).	Detection
PSCA_HUMAN	Prostate stem cell antigen	PSCA		LungCancers	Cell membrane; Lipid-anchor, GPI-anchor.	Literature, Prediction
PTGIS_HUMAN	Prostacyclin synthase	PTGIS	EPI	LungCancers, Benign-Nodules	Endoplasmic reticulum membrane; Single-pass membrane protein.	UniProt, Detection, Prediction
PTPA_HUMAN	Serine/threonine-protein phosphatase 2A activator	PPP2R4	ENDO	Symptoms		Detection, Prediction
PTPRC_HUMAN	Receptor-type tyrosine-protein phosphatase C	PTPRC	Secreted, EPI, ENDO	LungCancers	Membrane; Single-pass type I membrane protein.	UniProt, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
PTPRJ_HUMAN	Receptor-type tyrosine-protein phosphatase eta	PTPRJ	EPI	LungCancers, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Detection, Prediction
PVR_HUMAN	Poliovirus receptor	PVR		Symptoms	Isoform Alpha: Cell membrane; Single-pass type I membrane protein.   Isoform Delta: Cell membrane; Single-pass type I membrane protein.   Isoform Beta: Secreted.   Isoform Gamma: Secreted.	UniProt, Detection, Prediction
RAB32_HUMAN	Ras-related protein Rab-32	RAB32	EPI		Mitochondrion.	Prediction
RAGE_HUMAN	Advanced glycosylation end product-specific receptor	AGER	Secreted	LungCancers, Benign-Nodules	Isoform 1: Cell membrane; Single-pass type I membrane protein.   Isoform 2: Secreted.	UniProt, Literature
RAN_HUMAN	GTP-binding nuclear protein Ran	RAN	Secreted, EPI	LungCancers, Benign-Nodules	Nucleus. Cytoplasm. Melanosome. Note = Becomes dispersed throughout the cytoplasm during mitosis. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Detection, Prediction
RAP2B_HUMAN	Ras-related protein Rap-2b	RAP2B	EPI		Cell membrane; Lipid-anchor; Cytoplasmic side (Potential).	Prediction
RAP2C_HUMAN	Ras-related protein Rap-2c	RAP2C	EPI		Cell membrane; Lipid-anchor; Cytoplasmic side (Potential).	Prediction
RCN3_HUMAN	Reticulocalbin-3	RCN3	EPI	Symptoms	Endoplasmic reticulum lumen (Potential).	Prediction
RL24_HUMAN	60S ribosomal protein L24	RPL24	EPI			Prediction
S10A1_HUMAN	Protein S100-A1	S100A1		Symptoms	Cytoplasm.	Literature, Prediction
S10A6_HUMAN	Protein S100-A6	S100A6	Secreted	LungCancers	Nucleus envelope. Cytoplasm.	Literature, Detection, Prediction
S10A7_HUMAN	Protein S100-A7	S100A7		LungCancers	Cytoplasm. Secreted. Note = Secreted by a non-	UniProt, Literature, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
SAA_HUMAN	Serum amyloid A protein	SAA1		Symptoms	classical secretory pathway. Secreted.	UniProt, Literature, Detection, Prediction
SCF_HUMAN	Kit ligand	KITLG		LungCancers, Symptoms	Isoform 1: Cell membrane; Single-pass type I membrane protein (By similarity). Secreted (By similarity). Note = Also exists as a secreted soluble form (isoform 1 only) (By similarity). Isoform 2: Cell membrane; Single-pass type I membrane protein (By similarity). Cytoplasm, cytoskeleton (By similarity).	UniProt, Literature
SDC1_HUMAN	Syndecan-1	SDC1		LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature, Detection
SEM3G_HUMAN	Semaphorin-3G	SEMA3G		LungCancers	Secreted (By similarity).	UniProt, Prediction
SEPR_HUMAN	Seprase	FAP	ENDO	Symptoms	Cell membrane; Single-pass type II membrane protein. Cell projection, lamellipodium membrane; Single-pass type II membrane protein. Cell projection, invadopodium membrane; Single-pass type II membrane protein. Note = Found in cell surface lamellipodia, invadopodia and on shed vesicles.	UniProt, Literature, Detection
SERPH_HUMAN	Serpin H1	SERPINH1	Secreted, EPI, ENDO	LungCancers, Benign-Nodules	Endoplasmic reticulum lumen.	Detection, Prediction
SFPA2_HUMAN	Pulmonary surfactant-associated protein A2	SFTPA2	Secreted	LungCancers, Benign-Nodules	Secreted, extracellular space, extracellular matrix.	UniProt, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
SFTA1_HUMAN	Pulmonary surfactant-associated protein A1	SFTA1	Secreted	LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, surface film. Secreted, extracellular space, extracellular matrix. Secreted, extracellular space, surface film.	UniProt, Prediction
SG3A2_HUMAN	Secretoglobulin family 3A member 2	SCGB3A2		LungCancers, Benign-Nodules	Secreted.	UniProt, Prediction
SGPL1_HUMAN	Sphingosine-1-phosphate lyase 1	SGPL1	ENDO		Endoplasmic reticulum membrane; Single-pass type III membrane protein.	UniProt, Prediction
SIAL_HUMAN	Bone sialoprotein 2	IBSP		LungCancers	Secreted.	UniProt, Literature, Prediction
SLPI_HUMAN	Antileukoproteinase	SLPI		LungCancers, Benign-Nodules	Secreted.	UniProt, Literature, Detection, Prediction
SMD3_HUMAN	Small nuclear ribonucleoprotein Sm D3	SNRPD3	Secreted	Benign-Nodules	Nucleus.	Prediction
SMS_HUMAN	Somato-statin	SST		LungCancers	Secreted.	UniProt, Literature, Prediction
SODM_HUMAN	Superoxide dismutase [Mn], mitochondrial	SOD2	Secreted	LungCancers, Benign-Nodules, Symptoms	Mitochondrion matrix.	Literature, Detection, Prediction
SORL1_HUMAN	Sortilin-related receptor	SORL1	EPI	LungCancers, Symptoms	Membrane; Single-pass type I membrane protein (Potential).	UniProt, Detection
SPB3_HUMAN	Serpin B3	SERPINB3		LungCancers, Benign-Nodules	Cytoplasm. Note = Seems to also be secreted in plasma by cancerous cells but at a low level.	Literature, Detection
SPB5_HUMAN	Serpin B5	SERPINB5		LungCancers	Secreted, extracellular space.	UniProt, Detection
SPON2_HUMAN	Spondin-2	SPON2		LungCancers, Benign-Nodules	Secreted, extracellular space, extracellular matrix (By similarity).	UniProt, Prediction
SPRC_HUMAN	SPARC	SPARC		LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix, basement membrane. Note = In or around the basement membrane.	UniProt, Literature, Detection, Prediction
SRC_HUMAN	Proto-oncogene tyrosine-protein kinase Src	SRC	ENDO	LungCancers, Benign-Nodules, Symptoms		Literature

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
SSRD_HUMAN	Translocon-associated protein subunit delta	SSR4	Secreted, ENDO		Endoplasmic reticulum membrane; Single-pass type I membrane protein.	UniProt, Prediction
STAT1_HUMAN	Signal transducer and activator of transcription 1-alpha/beta	STAT1	EPI	LungCancers, Benign-Nodules	Cytoplasm. Nucleus. Note = Translocated into the nucleus in response to IFN-gamma-induced tyrosine phosphorylation and dimerization.	Detection
STAT3_HUMAN	Signal transducer and activator of transcription 3	STAT3	ENDO	LungCancers, Benign-Nodules, Symptoms	Cytoplasm. Nucleus. Note = Shuttles between the nucleus and the cytoplasm. Constitutive nuclear presence is independent of tyrosine phosphorylation.	Prediction
STC1_HUMAN	Stanniocalcin-1	STC1		LungCancers, Symptoms	Secreted.	UniProt, Prediction
STT3A_HUMAN	Dolichyl-diphosphooligoprotein glycosyltransferase subunit STT3A	STT3A	EPI		Endoplasmic reticulum membrane; Multi-pass membrane protein.	Literature
TAGL_HUMAN	Transgelin	TAGLN	EPI	LungCancers	Cytoplasm (Probable). Nucleus.	Literature, Prediction
TARA_HUMAN	TRIO and F-actin-binding protein	TRIOBP	ENDO		Cytoplasm, cytoskeleton. Note = Localized to F-actin in a periodic pattern.	Detection, Prediction
TBA1B_HUMAN	Tubulin alpha-1B chain	TUBA1B	EPI	LungCancers		Detection
TBB2A_HUMAN	Tubulin beta-2A chain	TUBB2A	EPI	LungCancers, Benign-Nodules		Detection, Prediction
TBB3_HUMAN	Tubulin beta-3 chain	TUBB3	EPI	LungCancers, Benign-Nodules		Detection
TBB5_HUMAN	Tubulin beta chain	TUBB	EPI	LungCancers, Benign-Nodules		Detection
TCPA_HUMAN	T-complex protein 1 subunit alpha	TCP1	EPI		Cytoplasm.	Prediction
TCPD_HUMAN	T-complex protein 1 subunit delta	CCT4	EPI		Cytoplasm. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV.	Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
TCPQ_HUMAN	T-complex protein 1 subunit theta	CCT8	Secreted, EPI		Cytoplasm.	Prediction
TCPZ_HUMAN	T-complex protein 1 subunit zeta	CCT6A	Secreted, EPI		Cytoplasm.	Detection
TDRD3_HUMAN	Tudor domain-containing protein 3	TDRD3	ENDO		Cytoplasm. Nucleus. Note = Predominantly cytoplasmic. Associated with actively translating polyribosomes and with mRNA stress granules.	Prediction
TENA_HUMAN	Tenascin	TNC	ENDO	LungCancers, Benign-Nodules, Symptoms	Secreted, extracellular space, extracellular matrix.	UniProt, Literature, Detection
TENX_HUMAN	Tenascin-X	TNXB	ENDO	LungCancers, Symptoms	Secreted, extracellular space, extracellular matrix.	UniProt, Detection, Prediction
TERA_HUMAN	Transitional endoplasmic reticulum ATPase	VCP	EPI	LungCancers, Benign-Nodules	Cytoplasm, cytosol. Nucleus. Note = Present in the neuronal hyaline inclusion bodies specifically found in motor neurons from amyotrophic lateral sclerosis patients. Present in the Lewy bodies specifically found in neurons from Parkinson disease patients.	Detection
TETN_HUMAN	Tetranectin	CLEC3B		LungCancers	Secreted.	UniProt, Literature, Detection, Prediction
TF_HUMAN	Tissue factor	F3		LungCancers, Benign-Nodules, Symptoms	Membrane; Single-pass type I membrane protein.	UniProt, Literature
TFR1_HUMAN	Transferrin receptor protein 1	TFRC	Secreted, EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Cell membrane; Single-pass type II membrane protein. Melanosome. Note = Identified by mass spectrometry in melanosome fractions from stage I to stage IV. Transferrin receptor protein 1,	UniProt, Literature, Detection

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
TGFA_HUMAN	Protransforming growth factor alpha	TGFA		LungCancers, Benign-Nodules	serum form: Secreted. Transforming growth factor alpha: Secreted, extracellular space.   Protransforming growth factor alpha: Cell membrane; Single-pass type I membrane protein.	UniProt, Literature
THAS_HUMAN	Thromboxane-A synthase	TBXAS1	EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Membrane; Multi-pass membrane protein.	Prediction
THY1_HUMAN	Thy-1 membrane glycoprotein	THY1	EPI		Cell membrane; Lipid-anchor, GPI-anchor (By similarity).	Detection, Prediction
TIMP1_HUMAN	Metalloproteinase inhibitor 1	TIMP1		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
TIMP3_HUMAN	Metalloproteinase inhibitor 3	TIMP3		LungCancers, Benign-Nodules	Secreted, extracellular space, extracellular matrix.	UniProt, Literature, Prediction
TLL1_HUMAN	Tolloid-like protein 1	TLL1	ENDO		Secreted (Probable).	UniProt, Prediction
TNFSF12_HUMAN	Tumor necrosis factor ligand superfamily member 12	TNFSF12		LungCancers, Benign-Nodules	Cell membrane; Single-pass type II membrane protein.   Tumor necrosis factor ligand superfamily member 12, secreted form: Secreted.	UniProt
TNR6_HUMAN	Tumor necrosis factor receptor superfamily member 6	FAS		LungCancers, Benign-Nodules, Symptoms	Isoform 1: Cell membrane; Single-pass type I membrane protein.   Isoform 2: Secreted.   Isoform 3: Secreted.   Isoform 4: Secreted.   Isoform 5: Secreted.   Isoform 6: Secreted.	UniProt, Literature, Prediction
TPIS_HUMAN	Triosephosphate isomerase	TPI1	Secreted, EPI	Symptoms		Literature, Detection, Prediction
TRFL_HUMAN	Lactotransferrin	LTF	Secreted, EPI, ENDO	LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
TSP1_HUMAN	Thrombospondin-1	THBS 1		LungCancers, Benign-Nodules, Symptoms		Literature, Detection, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
TTHY_HUMAN	Transthyretin	TTR		LungCancers, Benign-Nodules	Secreted. Cytoplasm.	UniProt, Literature, Detection, Prediction
TYPH_HUMAN	Thymidine phosphorylase	TYMP	EPI	LungCancers, Benign-Nodules, Symptoms		Literature, Detection, Prediction
UGGG1_HUMAN	UDP-glucose:glyco protein glucosyl-transferase 1	UGGT1	Secreted, ENDO		Endoplasmic reticulum lumen. Endoplasmic reticulum-Golgi intermediate compartment.	Detection, Prediction
UGGG2_HUMAN	UDP-glucose:glyco protein glucosyl-transferase 2	UGGT2	ENDO		Endoplasmic reticulum lumen. Endoplasmic reticulum-Golgi intermediate compartment.	Prediction
UGPA_HUMAN	UTP--glucose-1-phosphate uridyl-dyl-yltransferase	UGP2	EPI	Symptoms	Cytoplasm.	Detection
UPAR_HUMAN	Urokinase plasminogen activator surface receptor	PLAUR		LungCancers, Benign-Nodules, Symptoms	Isoform 1: Cell membrane; Lipid-anchor, GPI-anchor. Isoform 2: Secreted (Probable).	UniProt, Literature, Prediction
UTER_HUMAN	Uteroglobin	SCGB1A1		LungCancers, Benign-Nodules, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
VA0D1_HUMAN	V-type proton ATPase subunit d1	ATP6V0D1	EPI			Prediction
VAV3_HUMAN	Guanine nucleotide exchange factor VAV3	VAV3	ENDO			Prediction
VEGFA_HUMAN	Vascular endothelial growth factor A	VEGFA		LungCancers, Benign-Nodules, Symptoms	Secreted. Note = VEGF 121 is acidic and freely secreted. VEGF165 is more basic, has heparin-binding properties and, although a significant proportion remains cell-associated, most is freely secreted. VEGF189 is very basic, it is cell-associated after secretion and is bound avidly by heparin and the	UniProt, Literature, Prediction

TABLE 6-continued

UniProt Protein	Protein Name	Gene Symbol	Sources of Tissue Biomarkers	Biomarkers in Literature	Subcellular Location (UniProt)	Evidence for Presence in Blood
VEGFC_HUMAN	Vascular endothelial growth factor C	VEGFC		LungCancers, Benign-Nodules	extracellular matrix, although it may be released as a soluble form by heparin, heparinase or plasmin. Secreted.	UniProt, Literature, Prediction
VEGFD_HUMAN	Vascular endothelial growth factor D	FIGF		LungCancers	Secreted.	UniProt, Literature, Prediction
VEGFR1_HUMAN	Vascular endothelial growth factor receptor 1	FLT1		LungCancers, Benign-Nodules, Symptoms	Isoform Flt1: Cell membrane; Single-pass type I membrane protein.   Isoform sFlt1: Secreted.	UniProt, Literature, Detection, Prediction
VTNC_HUMAN	Vitronectin	VTN	ENDO	Symptoms	Secreted, extracellular space.	UniProt, Literature, Detection, Prediction
VWC2_HUMAN	Bronin	VWC2		LungCancers	Secreted, extracellular space, extracellular matrix, basement membrane (By similarity).	UniProt, Prediction
WNT3A_HUMAN	Protein Wnt-3a	WNT3A		LungCancers, Symptoms	Secreted, extracellular space, extracellular matrix.	UniProt, Prediction
WT1_HUMAN	Wilms tumor protein	WT1		LungCancers, Benign-Nodules, Symptoms	Nucleus. Cytoplasm (By similarity). Note = Shuttles between nucleus and cytoplasm (By similarity).   Isoform 1: Nucleus speckle.   Isoform 4: Nucleus, nucleoplasm.	Literature, Prediction
ZA2G_HUMAN	Zinc-alpha-2-glycoprotein	AZGP 1		LungCancers, Symptoms	Secreted.	UniProt, Literature, Detection, Prediction
ZG16B_HUMAN	Zymogen granule protein 16 homolog B	ZG16B		LungCancers	Secreted (Potential).	UniProt, Prediction

190 of these candidate protein biomarkers were shown to be measured reproducibly in blood. A moderately powered multisite and unbiased study of 242 blood samples from patients with PN was designed to determine whether a statistically significant subpanel of proteins could be identified to distinguish benign and malignant nodules of sizes under 2 cm. The three sites contributing samples and clinical data to this study were the University of Laval, University of Pennsylvania and New York University.

In an embodiment of the invention, a panel of 15 proteins effectively distinguished between samples derived from patients with benign and malignant nodules less than 2 cm diameter.

Bioinformatic and biostatistical analyses were used first to identify individual proteins with statistically significant differential expression, and then using these proteins to derive one or more combinations of proteins or panels of proteins, which collectively demonstrated superior discriminatory per-

formance compared to any individual protein. Bioinformatic and biostatistical methods are used to derive coefficients (C) for each individual protein in the panel that reflects its relative expression level, i.e. increased or decreased, and its weight or importance with respect to the panel's net discriminatory ability, relative to the other proteins. The quantitative discriminatory ability of the panel can be expressed as a mathematical algorithm with a term for each of its constituent proteins being the product of its coefficient and the protein's plasma expression level (P) (as measured by LC-SRM-MS), e.g.  $C \times P$ , with an algorithm consisting of n proteins described as:  $C1 \times P1 + C2 \times P2 + C3 \times P3 + \dots + Cn \times Pn$ . An algorithm that discriminates between disease states with a predetermined level of statistical significance may be referred to a "disease classifier". In addition to the classifier's constituent proteins with differential expression, it may also include proteins with minimal or no biologic variation to enable assessment of variability, or the lack thereof, within or between clinical specimens; these proteins may be termed typical native proteins and serve as internal controls for the other classifier proteins.

In certain embodiments, expression levels are measured by MS. MS analyzes the mass spectrum produced by an ion after its production by the vaporization of its parent protein and its separation from other ions based on its mass-to-charge ratio. The most common modes of acquiring MS data are 1) full scan acquisition resulting in the typical total ion current plot (TIC), 2) selected ion monitoring (SIM), and 3) selected reaction monitoring (SRM).

In certain embodiments of the methods provided herein, biomarker protein expression levels are measured by LC-SRM-MS. LC-SRM-MS is a highly selective method of tandem mass spectrometry which has the potential to effectively filter out all molecules and contaminants except the desired analyte(s). This is particularly beneficial if the analysis sample is a complex mixture which may comprise several isobaric species within a defined analytical window. LC-SRM-MS methods may utilize a triple quadrupole mass spectrometer which, as is known in the art, includes three quadrupole rod sets. A first stage of mass selection is performed in the first quadrupole rod set, and the selectively transmitted ions are fragmented in the second quadrupole rod set. The resultant transition (product) ions are conveyed to the third quadrupole rod set, which performs a second stage of mass selection. The product ions transmitted through the third quadrupole rod set are measured by a detector, which generates a signal representative of the numbers of selectively transmitted product ions. The RF and DC potentials applied to the first and third quadrupoles are tuned to select (respectively) precursor and product ions that have m/z values lying within narrow specified ranges. By specifying the appropriate transitions (m/z values of precursor and product ions), a peptide corresponding to a targeted protein may be measured with high degrees of sensitivity and selectivity. Signal-to-noise ratio is superior to conventional tandem mass spectrometry (MS/MS) experiments, which select one mass window in the first quadrupole and then measure all generated transitions in the ion detector. LC-SRM-MS.

In certain embodiments, an SRM-MS assay for use in diagnosing or monitoring lung cancer as disclosed herein may utilize one or more peptides and/or peptide transitions derived from the proteins set forth in Table 6. In certain embodiments, the assay may utilize peptides and/or peptide transitions from 100 or more, 150 or more, 200 or more, 250 or more, 300 or more, 345 or more, or 371 or more biomarker proteins. In certain embodiments, two or more peptides may be utilized per biomarker proteins, and in certain of these

embodiments three or more of four or more peptides may be utilized. Similarly, in certain embodiments two or more transitions may be utilized per peptide, and in certain of these embodiments three or more; four or more; or five or more transitions may be utilized per peptide. In one embodiment, an LC-SRM-MS assay for use in diagnosing lung cancer may measure the intensity of five transitions that correspond to selected peptides associated with each biomarker protein. The achievable limit of quantification (LOQ) may be estimated for each peptide according to the observed signal intensities during this analysis. For examples, for sets of target proteins associated with lung cancer see Table 12.

The expression level of a biomarker protein can be measured using any suitable method known in the art, including but not limited to mass spectrometry (MS), reverse transcriptase-polymerase chain reaction (RT-PCR), microarray, serial analysis of gene expression (SAGE), gene expression analysis by massively parallel signature sequencing (MPSS), immunoassays (e.g., ELISA), immunohistochemistry (RIC), transcriptomics, and proteomics.

To evaluate the diagnostic performance of a particular set of peptide transitions, a ROC curve is generated for each significant transition.

An "ROC curve" as used herein refers to a plot of the true positive rate (sensitivity) against the false positive rate (specificity) for a binary classifier system as its discrimination threshold is varied. A ROC curve can be represented equivalently by plotting the fraction of true positives out of the positives (TPR=true positive rate) versus the fraction of false positives out of the negatives (FPR=false positive rate). Each point on the ROC curve represents a sensitivity/specificity pair corresponding to a particular decision threshold. FIGS. 7 and 9 provide a graphical representation of the functional relationship between the distribution of biomarker or biomarker panel sensitivity and specificity values in a cohort of diseased subjects and in a cohort of non-diseased subjects.

AUC represents the area under the ROC curve. The AUC is an overall indication of the diagnostic accuracy of 1) a biomarker or a panel of biomarkers and 2) a ROC curve. AUC is determined by the "trapezoidal rule." For a given curve, the data points are connected by straight line segments, perpendiculars are erected from the abscissa to each data point, and the sum of the areas of the triangles and trapezoids so constructed is computed. In certain embodiments of the methods provided herein, a biomarker protein has an AUC in the range of about 0.75 to 1.0. In certain of these embodiments, the AUC is in the range of about 0.8 to 0.8, 0.9 to 0.95, or 0.95 to 1.0.

The methods provided herein are minimally invasive and pose little or no risk of adverse effects. As such, they may be used to diagnose, monitor and provide clinical management of subjects who do not exhibit any symptoms of a lung condition and subjects classified as low risk for developing a lung condition. For example, the methods disclosed herein may be used to diagnose lung cancer in a subject who does not present with a PN and/or has not presented with a PN in the past, but who nonetheless deemed at risk of developing a PN and/or a lung condition. Similarly, the methods disclosed herein may be used as a strictly precautionary measure to diagnose healthy subjects who are classified as low risk for developing a lung condition.

The present invention provides a method of determining the likelihood that a lung condition in a subject is cancer by measuring an abundance of a panel of proteins in a sample obtained from the subject; calculating a probability of cancer score based on the protein measurements and ruling out cancer for the subject if the score) is lower than a pre-determined

score, wherein when cancer is ruled out the subject does not receive a treatment protocol. Treatment protocols include for example pulmonary function test (PFT), pulmonary imaging, a biopsy, a surgery, a chemotherapy, a radiotherapy, or any combination thereof. In some embodiments, the imaging is an x-ray, a chest computed tomography (CT) scan, or a positron emission tomography (PET) scan.

The present invention further provides a method of ruling in the likelihood of cancer for a subject by measuring an abundance of panel of proteins in a sample obtained from the subject, calculating a probability of cancer score based on the protein measurements and ruling in the likelihood of cancer for the subject if the score in step is higher than a pre-determined score

In another aspect the invention further provides a method of determining the likelihood of the presence of a lung condition in a subject by measuring an abundance of panel of proteins in a sample obtained from the subject, calculating a probability of cancer score based on the protein measurements and concluding the presence of said lung condition if the score is equal or greater than a pre-determined score. The lung condition is lung cancer such as for example, non-small cell lung cancer (NSCLC). The subject at risk of developing lung cancer

The panel includes at least 4 proteins selected from ALDOA, FRIL, LG3BP, IBP3, LRP1, ISLR, TSP COIA1, GRP78, TETN, PRXD1 and CD14. Optionally, the panel further includes at least one protein selected from BGH3, COIA1, TETN, GRP78, PRDX, FIBA and GSLG1.

The subject has or is suspected of having a pulmonary nodule. The pulmonary nodule has a diameter of less than or equal to 3 cm. In one embodiment, the pulmonary nodule has a diameter of about 0.8 cm to 2.0 cm.

The score is calculated from a logistic regression model applied to the protein measurements. For example, the score is determined as  $P_s = 1/[1 + \exp(-\alpha - \sum_{i=1}^N \beta_i * \tilde{I}_{i,s})]$ , where  $\tilde{I}_{i,s}$  is logarithmically transformed and normalized intensity of transition  $i$  in said sample ( $s$ ),  $\beta_i$  is the corresponding logistic regression coefficient,  $\alpha$  was a panel-specific constant, and  $N$  was the total number of transitions in said panel.

In various embodiments, the method of the present invention further comprises normalizing the protein measurements. For example, the protein measurements are normalized by one or more proteins selected from PEDF, MASP1, GELS, LUM, C163A and PTPRJ.

The biological sample such as for example tissue, blood, plasma, serum, whole blood, urine, saliva, genital secretion, cerebrospinal fluid, sweat and excreta.

In one aspect, the determining the likelihood of cancer is determined by the sensitivity, specificity, negative predictive value or positive predictive value associated with the score. The score determined has a negative predictive value (NPV) is at least about 80%.

The measuring step is performed by selected reaction monitoring mass spectrometry, using a compound that specifically binds the protein being detected or a peptide transition. In one embodiment, the compound that specifically binds to the protein being measured is an antibody or an aptamer.

In specific embodiments, the diagnostic methods disclosed herein are used to rule out a treatment protocol for a subject, measuring the abundance of a panel of proteins in a sample obtained from the subject, calculating a probability of cancer score based on the protein measurements and ruling out the treatment protocol for the subject if the score determined in the sample is lower than a pre-determined score. In some embodiments the panel contains at least 4 proteins selected

ALDOA, FRIL, LG3BP, IBP3, LRP1, ISLR, TSP, COIA1, GRP78, TETN, PRXD1 and CD14

Optionally, the panel further comprises one or more proteins selected from ERO1A, 6PGD, GSTP1, GGH, PRDX1, CD14, PTPA, ICAM1, FOLH1, SODM, FIBA, GSLG1, RAP2B, or C163A or one or more proteins selected from LRP1, COIA1, TSP1, ALDOA, GRP78, FRIL, LG3BP, BGH3, ISLR, PRDX1, FIBA, or GSLG. In preferred embodiments, the panel contains at least TSP1, LG3BP, LRP1, ALDOA, and COIA1. In more a preferred embodiment, the panel contains at least TSP1, LRP1, ALDOA and COIA1.

In specific embodiments, the diagnostic methods disclosed herein are used to rule in a treatment protocol for a subject by measuring the abundance of a panel of proteins in a sample obtained from the subject, calculating a probability of cancer score based on the protein measurements and ruling in the treatment protocol for the subject if the score determined in the sample is greater than a pre-determined score. In some embodiments the panel contains at least 4 proteins selected ALDOA, FRIL, LG3BP, IBP3, LRP1, ISLR or TSP1 or ALDOA, FRIL, LG3BP, IBP3, LRP1, ISLR, TSP COIA1, GRP78, TETN, PRXD1 and CD14. Optionally, the panel further comprises one or more proteins selected from ERO1A, 6PGD, GSTP1, COIA1, GGH, PRDX1, SEM3G, GRP78, TETN, AIFM1, MPRI, TNF12, MMP9 or OSTP or COALTETN, GRP78, APOE or TBB3.

In certain embodiments, the diagnostic methods disclosed herein can be used in combination with other clinical assessment methods, including for example various radiographic and/or invasive methods. Similarly, in certain embodiments, the diagnostic methods disclosed herein can be used to identify candidates for other clinical assessment methods, or to assess the likelihood that a subject will benefit from other clinical assessment methods.

The high abundance of certain proteins in a biological sample such as plasma or serum can hinder the ability to assay a protein of interest, particularly where the protein of interest is expressed at relatively low concentrations. Several methods are available to circumvent this issue, including enrichment, separation, and depletion. Enrichment uses an affinity agent to extract proteins from the sample by class, e.g., removal of glycosylated proteins by glyco-capture. Separation uses methods such as gel electrophoresis or isoelectric focusing to divide the sample into multiple fractions that largely do not overlap in protein content. Depletion typically uses affinity columns to remove the most abundant proteins in blood, such as albumin, by utilizing advanced technologies such as IgY14/Supermix (SigmaSt. Louis, Mo.) that enable the removal of the majority of the most abundant proteins.

In certain embodiments of the methods provided herein, a biological sample may be subjected to enrichment, separation, and/or depletion prior to assaying biomarker or putative biomarker protein expression levels. In certain of these embodiments, blood proteins may be initially processed by a glyco-capture method, which enriches for glycosylated proteins, allowing quantification assays to detect proteins in the high pg/ml to low ng/ml concentration range. Exemplary methods of glyco-capture are well known in the art (see, e.g., U.S. Pat. No. 7,183,188; U.S. Patent Appl. Publ. No. 2007/0099251; U.S. Patent Appl. Publ. No. 2007/0202539; U.S. Patent Appl. Publ. No. 2007/0269895; and U.S. Patent Appl. Publ. No. 2010/0279382). In other embodiments, blood proteins may be initially processed by a protein depletion method, which allows for detection of commonly obscured biomarkers in samples by removing abundant proteins. In one such embodiment, the protein depletion method is a GenWay depletion method.

In certain embodiments, a biomarker protein panel comprises two to 100 biomarker proteins. In certain of these embodiments, the panel comprises 2 to 5, 6 to 10, 11 to 15, 16 to 20, 21-25, 5 to 25, 26 to 30, 31 to 40, 41 to 50, 25 to 50, 51 to 75, 76 to 100, biomarker proteins. In certain embodiments, a biomarker protein panel comprises one or more subpanels of biomarker proteins that each comprise at least two biomarker proteins. For example, biomarker protein panel may comprise a first subpanel made up of biomarker proteins that are overexpressed in a particular lung condition and a second subpanel made up of biomarker proteins that are under-expressed in a particular lung condition.

In certain embodiments of the methods, compositions, and kits provided herein, a biomarker protein may be a protein that exhibits differential expression in conjunction with lung cancer. For example, in certain embodiments a biomarker protein may be one of the proteins associated with lung cancer set forth in Table 6.

In other embodiments, the diagnosis methods disclosed herein may be used to distinguish between two different lung conditions. For example, the methods may be used to classify a lung condition as malignant lung cancer versus benign lung cancer, NSCLC versus SCLC, or lung cancer versus non-cancer condition (e.g., inflammatory condition).

In certain embodiments, kits are provided for diagnosing a lung condition in a subject. These kits are used to detect expression levels of one or more biomarker proteins. Optionally, a kit may comprise instructions for use in the form of a label or a separate insert. The kits can contain reagents that specifically bind to proteins in the panels described, herein. These reagents can include antibodies. The kits can also contain reagents that specifically bind to mRNA expressing proteins in the panels described, herein. These reagents can include nucleotide probes. The kits can also include reagents for the detection of reagents that specifically bind to the proteins in the panels described herein. These reagents can include fluorophores.

The following examples are provided to better illustrate the claimed invention and are not to be interpreted as limiting the scope of the invention. To the extent that specific materials are mentioned, it is merely for purposes of illustration and is not

intended to limit the invention. One skilled in the art may develop equivalent means or reactants without the exercise of inventive capacity and without departing from the scope of the invention

EXAMPLES

Example 1

Identification of Lung Cancer Biomarker Proteins

A retrospective, case-control study design was used to identify biomarker proteins and panels thereof for diagnosing various lung diseases in pre-defined control and experimental groups. The first goal of these studies was to demonstrate statistically significant differential expression for individual proteins between control and experimental groups. The second goal is to identify a panel of proteins which all individually demonstrate statistically significant differential expression between control and experimental groups. This panel of proteins can then be used collectively to distinguish between dichotomous disease states.

Specific study comparisons may include 1) cancer vs. non-cancer, 2) small cell lung cancer versus non-small cell lung cancer (NSCLC), 3) cancer vs. inflammatory disease state (e.g., infectious granuloma), or 4) different nodule size, e.g., <10 mm versus ≥10 mm (alternatively using 10, 15 or 20 mm cut-offs depending upon sample distributions).

Data for each subject consisted of the following:

Archived plasma samples from subjects previously enrolled in Institute Review Board (IRB)-approved studies was used to identify biomarker proteins and biomarker panels for distinguishing lung malignancies from non-malignancies. Plasma samples were originally obtained by routine phlebotomy, aliquotted, and stored at -80° C. or lower. Sample preparation, assignment of subject identification codes, initial subject record entry, and specimen storage were performed as per IRB study protocols. Sample eligibility is based on clinical parameters, including the subject, PN, and clinical staging parameters. Parameters for inclusion and exclusion are set forth in Table 7.

TABLE 7

Inclusion Criteria	
Sample Inclusion Criteria	Sample eligibility will be based on clinical parameters, including the following subject, nodule and clinical staging parameters: Subject age ≥40 any smoking status, e.g. current, former, or never co-morbid conditions, e.g. COPD prior malignancy - only skin carcinomas - squamous or basal cell Nodule radiology size ≥4 mm and ≤30 mm solid, semi-solid or non-solid any spiculation or ground glass opacity pathology malignant - e.g. adenocarcinoma, squamous, or large cell benign - inflammatory (e.g. granulomatous, infectious) or non-inflammatory (e.g. hamartoma) confirmed by biopsy, surgery or stability of lung nodule for 2 years or more. Clinical stage Primary tumor: ≤T1 (e.g. 1A, 1B) Regional lymph nodes: N0 or N1 only Distant metastasis: M0 only

TABLE 7-continued

Inclusion Criteria	
Sample Exclusion Criteria	Subject prior malignancy within 5 years of lung nodule diagnosis Nodule size data unavailable for cancer or benign nodule, no pathology or follow-up CT data available Clinical stage Primary tumor: $\geq$ T2 Regional lymph nodes: $\geq$ N2 Distant metastasis: $\geq$ M1

15

The assignment of a sample to a control or experimental group, and its further stratification or matching to other samples within and between these groups, is dependent on various clinical data about the subject. This data includes, for example, demographic information such as age, gender, and clinical history (e.g., smoking status), co-morbid conditions, PN characterization, and pathologic interpretation of resected lesions and tissues (Table 8).

20

The study design and analytical plan prioritizes the control: experimental group pairings set forth in Table 9. Additional clinical and molecular insights may be gained by selective inclusion of phenotypes, e.g. effect of smoking, in the assignment of experimental and control groups. Demographic information available in the clinical database will enable further refinements in sample selection via the stratification or matching of samples in the case-control analyses with respect to clinical parameters, e.g., age and nodule size.

TABLE 8

	1. Enrollment Data a. Demographics - age, birth date, gender, ethnicity b. Measurements - Height (cm) and weight (kg) c. Smoking history - never, former, or current with pack-year estimation d. Medical history - details of co-morbid conditions, e.g. chronic obstructive pulmonary disease (COPD), inflammatory or autoimmune diseases, endocrine (diabetes), and cardiovascular e. Medication history - current medications, dosages and indications f. Radiographic data and nodule characteristics 1) nodule size in millimeters (width $\times$ height $\times$ length) 2) location, e.g. right or left and upper, lower or middle 3) quality, e.g. solid, semi-solid, ground glass, calcified, etc. 2. Diagnostic Evaluation Data a. Primary diagnosis and associated reports (clinical history, physical exam, and laboratory tests report) b. Pulmonary Function Tests (PFTs), if available c. Follow-up CT scans - subsequent nodule evaluations by chest CT d. PET scan e. Clinical Staging f. Biopsy procedures 1) FNA or TTNA 2) bronchoscopy with transbronchial or needle biopsy 3) surgical diagnostic procedures, e.g. VATS and/or thoracotomy 3. Radiology Report(s) 4. Pathology Report(s) 5. Blood Sample Collection Information 6. Reporting of Adverse Events a. AEs resulting from center's SOC, e.g. procedural morbidity.
Subject	demographics - e.g. age, gender, ethnicity smoking status - e.g. never-, former-("ex-") or current-smoker; pack-years clinical history - e.g. co-morbid conditions, e.g. COPD, infection
Nodule	size - e.g. planar (width $\times$ height $\times$ length) and volume dimensions appearance - e.g. calcifications, ground glass appearance, eccentricity
Pathology	primary lung vs. systemic disorder malignancy status - malignant vs. benign (vs. indeterminate) histopathology - e.g. small cell lung cancer (SCLC) vs. non-small cell lung cancer (NSCLC - adenocarcinoma, squamous carcinoma, large cell carcinoma); other types, e.g. hematologic, carcinoid, etc. immunologically quiescent, e.g. hamartoma, vs. inflammatory, e.g. granulomatous and/or infectious, e.g. fungal

TABLE 9

Assignment of Experimental and Control Groups to Achieve Proteomic Analysis Objectives			
Analysis	Objective	Experimental Group	Control Group
1	Differentiate cancer from benign lung nodule	A. Cancer nodule	Any non-malignant (benign) phenotype with nodule $\geq 4$ mm in diameter
2	Differentiate cancer from non-malignant (inflammatory, infectious) lung nodule	A. Cancer nodule	Non-malignant (non-benign) lung disorder, e.g. granulomatous (fungal) disease, with nodule

LC-SRM-MS is performed to identify and quantify various plasma proteins in the plasma samples. Prior to LC-SRM-MS analysis, each sample is depleted using IgY14/Supermix (Sigma) and then trypsin-digested. Samples from each control or experimental group are batched randomly and processed together on a QTrap 5500 instrument (AB SCIEX, Foster City, Calif.) for unbiased comparisons. Each sample analysis takes approximately 30 minutes. Peak areas for two transitions (native and heavy label) are collected and reported for all peptides and proteins. The data output for each protein analyzed by LC-SRM-MS typically yields four measurements consisting of two transition measurements from each of two peptides from the same protein. These measurements enable an inference of the relative abundance of the target protein, which will be used as its expression level in the bioinformatics and statistical analyses.

Identification of biomarker proteins having differential expression levels between the control and experimental groups yields one or more novel proteomic profiles. For example, biomarker proteins are identified with expression levels that differ in subjects with PNs who are diagnosed with NSCLC versus those without an NSCLC diagnosis, or in subjects with PNs who are diagnosed with NSCLC versus an inflammatory disorder. Panels of biomarker proteins are also identified which can collectively discriminate between dichotomous disease states.

Analyses may be (a priori) powered appropriately to control type 1 and type 2 errors at 0.05 and to detect inter-cohort differences of 25% per analyte. The diagnostic power of individual proteins is generally assessed to distinguish between two cohorts, assuming a one-sided paired non-parametric test is used. This provides a lower bound on the sample size required to demonstrate differential expression between experimental and control groups. Multiple testing effects apply for the identification of panels of proteins for assessing diagnostic efficacy, which requires larger sample sizes.

The sequence of steps for determining statistical significance for differential expression of an individual protein includes the following: 1) assessing and correlating the calibrated values of transitions of a single protein (a quality control measure); 2) comparing paired analysis of groups to control for other influences using the Mann-Whitney U-test (rank sum) to determine statistical significance; and 3) determining its significance based on a pre-defined significance threshold. Transitions within a protein that are not correlated across samples (e.g., Pearson correlation  $< 0.5$ ) will be deemed unreliable and excluded from the analysis.

Comparison of calibrated samples between two cohorts, e.g., cancer and non-cancer, requires pairing or matching

using a variety of clinical parameters such as nodule size, age and gender. Such pairing controls for the potential influence of these other parameters on the actual comparison goal, e.g. cancer and non-cancer. A non-parametric test such as the Mann-Whitney U-test (rank sum) will then be applied to measure the statistical difference between the groups. The resulting p value can be adjusted using multiple testing corrections such as the false discovery rate. Permutation tests can be used for further significance assessments.

Significance will be determined by the satisfaction of a pre-defined threshold, such as 0.05, to filter out assays, with the potential use of higher threshold values for additional filtering. An additional significance criterion is that two of three replicate assays must individually be significant in order for the assay, e.g., single protein, to be significant.

Panels of proteins that individually demonstrate statistically significant differential expression as defined above and which can collectively be used to distinguish dichotomous disease states are identified using statistical methods described herein. This requires developing multivariate classifiers and assessing sensitivity, specificity, and ROC AUC for panels. In addition, protein panels with optimal discriminatory performance, e.g., ROC AUC, are identified and may be sufficient for clinical use in discriminating disease states.

The sequence of steps for determining the statistical significance of the discriminatory ability of a panel of proteins includes 1) developing multivariate classifiers for protein panels, and 2) identifying a protein panel with optimal discriminatory performance, e.g. ROC AUC, for a set of disease states.

A multivariate classifier (e.g., majority rule) will be developed for protein panels, including single protein assays deemed to be significant. The sensitivity and specificity of each classifier will be determined and used to generate a receiver operating characteristics (ROC) curve and its AUC to assess a given panel's discriminatory performance for a specific comparison, e.g. cancer versus non-cancer.

#### Protocol

1. Review clinical data from a set of subjects presenting with lung disease.

2. Provide plasma samples from the subjects wherein the samples are either benign, cancerous, COPD or another lung disease.

3. Group the plasma samples that are benign or cancerous by PNs that are separated by size of the nodule.

4. Target a pool of 371 putative lung cancer biomarker proteins consisting of at least two peptides per protein and at least two LC-SRM-MS transitions per peptide. Measuring the LC-SRM-MS transitions in each specimen along with 5 synthetic internal standards consisting of 10 transitions to compare peptide transitions from the plasma to the synthetic internal standards by LC-SRM-MS mass spectroscopy.

5. Quantitate the intensity of each transition.

6. Normalize the quantitated transitions to internal standards to obtain a normalized intensity.

7. Review the measured peptide transitions for correlations from the same peptide, rejecting discordant transitions.

8. Generate an ROC for each transition by comparing cancerous with benign samples. (ROC compare specificity (true positive) to (1-sensitivity) false positive).

9. Define the AUC for each transition. (An AUC of 0.5 is a random classifier; 1.0 is a perfect classifier).

10. Determine an AUC cut-off point to determine transitions that are statistically significant.

## 111

11. Define the transitions that exceed the AUC cutoff point.
12. Combine all pairings of significant transitions.
13. Define a new AUC for each transition pair by means of logistical regression.
14. Repeat pairing combinations into triples, quad, etc.; defining a new AUC based upon the logistical regression of combined transitions until a panel of biomarker transitions with combined desired performance (sensitivity & specificity) have been achieved.
15. The panel of biomarker transitions is verified against previously unused set of plasma panels.

## Example 2

## Diagnosis/Classification of Lung Disease Using Biomarker Proteins

Plasma samples will be obtained from one or more subjects presenting with PNs to evaluate whether the subjects have a lung condition. The plasma samples will be depleted using IgY14/Supermix (Sigma) and optionally subjected to one or more rounds of enrichment and/or separation, and then trypsinized. The expression level of one or more biomarker proteins previously identified as differentially expressed in subjects with the lung condition will be measured using an LC-SRM-MS assay. The LC-SRM-MS assay will utilize two to five peptide transitions for each biomarker protein. For example, the assay may utilize one or more of the peptide transitions generated from any of the proteins listed in Table 6. Subjects will be classified as having the lung condition if one or more of the biomarker proteins exhibit expression levels that differ significantly from the pre-determined control expression level for that protein.

## Example 3

## Blood-Based Diagnostic Test to Determine the Likelihood that a Pulmonary Nodule (PN) is Benign or Malignant

A panel of 15 proteins was created where the concentration of these 15 proteins relative to the concentration of 6 protein standards is indicative of likelihood of cancer. The relative concentration of these 15 proteins to the 6 protein standards was measured using a mass spectrometry methodology. A classification algorithm is used to combine these relative concentrations into a relative likelihood of the PN being benign or malignant. Further it has been demonstrated that there are many variations on these panels that are also diagnostic tests for the likelihood that a PN is benign or malignant. Variations on the panel of proteins, protein standards, measurement methodology and/or classification algorithm are described herein.

## Study Design

A Single Reaction Monitoring (SRM) mass spectrometry (MS) assay was developed consisting of 1550 transitions from 345 lung cancer associated proteins. The SRM-MS assay and methodology is described above. The goal of this study was to develop a blood-based diagnostic for classifying PNs under 2 cm in size as benign or malignant. The study design appears in Table 10.

## 112

TABLE 10

	Study Design					
	Small (<2 cm)			large (>2 cm)		
	Laval	UPenn	NYU	Laval	UPenn	NYU
Benign	14	29	29	13	21	15
Malignant	14	29	29	13	21	15
Batches	1	2	2	1	2	1
	72 vs. 72 (94% power)			49 vs. 49 (74% power)		

The study consisted of 242 plasma samples from three sites (Laval, UPenn and NYU). The number of benign and malignant samples from each site are indicated in Table 10. The study consisted of 144 plasma samples from patients with PNs of size 2 cm or less and of 98 samples from patients with PNs of size larger than 2 cm. This resulted in an estimated power of 94% for discovering proteins with blood concentrations of 1.5 fold or more between benign and malignant cancer samples of size 2 cm or less. Power is 74% for PNs of size larger than 2 cm.

This study was a retrospective multisite study that was intended to derive protein biomarkers of lung cancer that are robust to site-to-site variation. The study included samples larger than 2 cm to ensure that proteins not detectable due to the limit of detection of the measurement technology (LC-SRM-MS) for tumors of size 2 cm or less could still be detected in tumors of size 2 cm or larger.

Samples from each site and in each size class (above and below 2 cm) were matched on nodule size, age and gender.

## Sample Analysis

Each sample was analyzed using the LC-SRM-MS measurement methodology as follows:

1. Samples were depleted of high abundance proteins using the IGy14 and Supermix depletion columns from Sigma-Aldrich.

2. Samples were digested using trypsin into tryptic peptides.

3. Samples were analyzed by LC-SRM-MS using a 30 minute gradient on a Waters nanoacuity LC system followed by SRM-MS analysis of the 1550 transitions on a AB-Sciex 5500 triple quad device.

4. Raw transition ion counts were obtained and recorded for each of the 1550 transitions.

It is important to note that matched samples were processed at each step either in parallel (steps 2 and 4) or back-to-back serially (steps 1 and 3). This minimizes analytical variation. Finally, steps 1 and 2 of the sample analysis are performed in batches of samples according to day of processing. There were five batches of 'small' samples and four batches of 'large' samples as denoted in Table 10.

## Protein Shortlist

A shortlist of 68 proteins reproducibly diagnostic across sites was derived as follows. Note that each protein can be measured by multiple transitions.

## Step 1: Normalization

Six proteins were identified that had a transition detected in all samples of the study and with low coefficient of variation. For each protein the transition with highest median intensity across samples was selected as the representative transition for the protein. These proteins and transitions are found in Table 11.

TABLE 11

Normalizing Factors		
Protein (Uniprot ID)	Peptide (Amino Acid Sequence) (SEQ ID NO: )	Transition (m/z)
CD44_HUMAN	YGFIEGHVVIPR (SEQ ID NO: 1)	272.2
TENX_HUMAN	YEVTVVSVR (SEQ ID NO: 2)	759.5
CLUS_HUMAN	ASSIIDELFQDR (SEQ ID NO: 3)	565.3
IBP3_HUMAN	FLNVLSPR (SEQ ID NO: 4)	685.4
GELS_HUMAN	TASDFITK (SEQ ID NO: 5)	710.4
MASPI_HUMAN	TGVITSPDFPNPYPK (SEQ ID NO: 6)	258.10

We refer to the transitions in Table 11 as normalizing factors (NFs). Each of the 1550 transitions were normalized by each of the six normalizing factors where the new intensity of a transition *t* in a sample *s* by NF *f*, denoted  $New(s,t,f)$ , is calculated as follows:

$$New(s,t,f) = Raw(s,t) * Median(f) / Raw(s,f)$$

where  $Raw(s,t)$  is the original intensity of transition *t* in sample *s*;  $Median(f)$  is the median intensity of the NF *f* across all samples; and  $Raw(s,f)$  is the original intensity of the NF *f* in sample *s*.

For each protein and normalized transition, the AUC of each batch was calculated. The NF that minimized the coefficient of variation across the 9 batches was selected as the NF

for that protein and for all transitions of that protein. Consequently, every protein (and all of its transitions) are now normalized by a single NF.

#### Step 2: Reproducible Diagnostic Proteins

For each normalized transition its AUC for each of the nine batches in the study is calculated as follows. If the transition is detected in fewer than half of the cancer samples and in fewer than half of the benign samples then the batch AUC is 'ND'. Otherwise, the batch AUC is calculated comparing the benign and cancer samples in the batch.

The batch AUC values are transformed into percentile AUC scores for each transition. That is, if a normalized transition is in the 82nd percentile of AUC scores for all transitions then it is assigned percentile AUC 0.82 for that batch.

Reproducible transitions are those satisfying at least one of the following criteria:

1. In at least four of the five small batches the percentile AUC is 75% or more (or 25% and less).
2. In at least three of the five small batches the percentile AUC is 80% or more (or 20% and less) AND the remaining percentile AUCs in the small batches are above 50% (below 50%).
3. In all five small batches the percentile AUC is above 50% (below 50%).
4. In at least three of the four large batches the percentile AUC is 85% or more (or 15% and less).
5. In at least three of the four large batches the percentile AUC is 80% or more (or 20% and less) AND the remaining percentile AUCs in the large batches are above 50% (below 50%).
6. In all four large batches the percentile AUC is above 50% (below 50%).

These criteria result in a list of 67 proteins with at least one transition satisfying one or more of the criteria. These proteins appear in Table 12.

TABLE 12

Protein (Uniprot)	Occurrence Across 131 Panels	Percentage Occurrence Across 131 Panels	Protein Names	Uniprot Accession No.
G3P_HUMAN	113	86%	Glyceraldehyde-3-phosphate dehydrogenase; Short name = GAPDH; Alternative name(s): Peptidyl-cysteine S-nitrosylase GAPDH	P04406
FRIL_HUMAN	107	82%	Recommended name: Ferritin light chain Short name = Ferritin L subunit	P02792
HYOU1_HUMAN	69	53%	Recommended name: Hypoxia up-regulated protein 1 Alternative name(s): 150 kDa oxygen-regulated protein Short name = ORP-150 170 kDa glucose-regulated protein Short name = GRP-170	Q9Y4L1
ALDOA_HUMAN	66	50%	Recommended name: Fructose-bisphosphate aldolase A EC = 4.1.2.13 Alternative name(s): Lung cancer antigen NY-LU-1 Muscle-type aldolase	P04075
HXK1_HUMAN	65	50%	Recommended name: Hexokinase-1 EC = 2.7.1.1 Alternative name(s): Brain form hexokinase Hexokinase type I Short name = HK I	P19367
APOE_HUMAN	63	48%	Recommended name: Apolipoprotein E Short name = Apo-E	P02649

TABLE 12-continued

Protein (Uniprot)	Occurrence Across 131 Panels	Percentage Occurrence Across 131 Panels	Protein Names	Uniprot Accession No.
TSP1_HUMAN	63	48%	Recommended name: Thrombospondin-1	P07996
FINC_HUMAN	62	47%	Recommended name: Fibronectin Short name = FN Alternative name(s): Cold-insoluble globulin Short name = CIG Cleaved into the following 4 chains: 1. Anastellin 2. Ugl-Y1 3. Ugl-Y2 4. Ugl-Y3	P02751
LRP1_HUMAN	58	44%	Recommended name: Prolow-density lipoprotein receptor-related protein 1 Short name = LRP-1 Alternative name(s): Alpha-2-macroglobulin receptor Short name = A2MR Apolipoprotein E receptor Short name = APOER CD_antigen = CD91 Cleaved into the following 3 chains: 1. Low-density lipoprotein receptor-related protein 1 85 kDa subunit Short name = LRP-85 2. Low-density lipoprotein receptor-related protein 1 515 kDa subunit Short name = LRP-515 3. Low-density lipoprotein receptor-related protein 1 intracellular domain Short name = LRPICD	
6PGD_HUMAN	50	38%	Recommended name: 6-phosphogluconate dehydrogenase, decarboxylating	P52209
S10A6_HUMAN	47	36%	Recommended name: Protein S100-A6 Alternative name(s): Calcyclin Growth factor-inducible protein 2A9 MLN 4 Prolactin receptor-associated protein Short name = PRA S100 calcium-binding protein A6	P06703
CALU_HUMAN	45	34%	Recommended name: Calumenin Alternative name(s): Crocalbin IEF SSP 9302	O43852
PRDX1_HUMAN	45	34%	Recommended name: Peroxiredoxin-1 EC = 1.11.1.15 Alternative name(s): Natural killer cell-enhancing factor A Short name = NKEF-A Proliferation-associated gene protein Short name = PAG Thioredoxin peroxidase 2 Thioredoxin-dependent peroxide reductase 2	Q06830
RAN_HUMAN	45	34%	Recommended name: GTP-binding nuclear protein Ran Alternative name(s): Androgen receptor-associated protein 24 GTPase Ran Ras-like protein TC4 Ras-related nuclear protein	P62826
CD14_HUMAN	43	33%	Recommended name: Monocyte differentiation antigen CD14 Alternative name(s): Myeloid cell-specific leucine-rich glycoprotein CD_antigen = CD14 Cleaved into the following 2 chains: 1. Monocyte differentiation antigen CD14,	P08571

TABLE 12-continued

Protein (Uniprot)	Occurrence Across 131 Panels	Percentage Occurrence Across 131 Panels	Protein Names	Uniprot Accession No.
AMPN_HUMAN	41	31%	urinary form 2. Monocyte differentiation antigen CD14, membrane-bound form Recommended name: Aminopeptidase N Short name = AP-N Short name = hAPN EC = 3.4.11.2 Alternative name(s): Alanyl aminopeptidase Aminopeptidase M Short name = AP-M Microsomal aminopeptidase Myeloid plasma membrane glycoprotein CD13 gp150 CD_antigen = CD13	P15144
GSLG1_HUMAN	36	27%	Recommended name: Golgi apparatus protein 1 Alternative name(s): CFR-1 Cysteine-rich fibroblast growth factor receptor E-selectin ligand 1 Short name = ESL-1 Golgi sialoglycoprotein MG-160	Q92896
1433Z_HUMAN	32	24%	Recommended name: 14-3-3 protein zeta/delta Alternative name(s): Protein kinase C inhibitor protein 1 Short name = KCIP-1	P63104
IBP3_HUMAN	31	24%	Recommended name: Insulin-like growth factor-binding protein 3 Short name = IBP-3 Short name = IGF-binding protein 3 Short name = IGFBP-3	P17936
ILK_HUMAN	31	24%	Recommended name: Integrin-linked protein kinase EC = 2.7.11.1 Alternative name(s): 59 kDa serine/threonine-protein kinase ILK-1 ILK-2 p59ILK	Q13418
LDHB_HUMAN	30	23%	Recommended name: L-lactate dehydrogenase B chain Short name = LDH-B EC = 1.1.1.27 Alternative name(s): LDH heart subunit Short name = LDH-H	P07195
MPRI_HUMAN	29	22%	Renal carcinoma antigen NY-REN-46 Recommended name: Cation-independent mannose-6-phosphate receptor Short name = CI Man-6-P receptor Short name = CI-MPR Short name = M6PR Alternative name(s): 300 kDa mannose 6-phosphate receptor Short name = MPR 300 Insulin-like growth factor 2 receptor Insulin-like growth factor II receptor Short name = IGF-II receptor M6P/IGF2 receptor Short name = M6P/IGF2R CD_antigen = CD222	P11717
PROF1_HUMAN	29	22%	Recommended name: Profilin-1 Alternative name(s): Profilin I	P07737
PEDF_HUMAN	28	21%	Recommended name: Pigment epithelium-derived factor Short name = PEDF Alternative name(s):	P36955

TABLE 12-continued

Protein (Uniprot)	Occurrence Across 131 Panels	Percentage Occurrence Across 131 Panels	Protein Names	Uniprot Accession No.
CLIC1_HUMAN	26	20%	Cell proliferation-inducing gene 35 protein EPC-1 Serpine F1 Recommended name: Chloride intracellular channel protein 1 Alternative name(s): Chloride channel ABP Nuclear chloride ion channel 27 Short name = NCC27 Regulatory nuclear chloride ion channel protein Short name = hRNCC	O00299
GRP78_HUMAN	25	19%	Recommended name: 78 kDa glucose-regulated protein Short name = GRP-78 Alternative name(s): Endoplasmic reticulum luminal Ca(2+)- binding protein grp78 Heat shock 70 kDa protein 5 Immunoglobulin heavy chain-binding protein Short name = BiP	P11021
CEAM8_HUMAN	24	18%	Recommended name: Carcinoembryonic antigen-related cell adhesion molecule 8 Alternative name(s): CD67 antigen Carcinoembryonic antigen CGM6 Non-specific cross-reacting antigen NCA- 95 CD_antigen = CD66b	P31997
VTNC_HUMAN	24	18%	Recommended name: Vitronectin Alternative name(s): S-protein Serum-spreading factor V75 Cleaved into the following 3 chains: 1. Vitronectin V65 subunit 2. Vitronectin V10 subunit 3. Somatomedin-B	P04004
CERU_HUMAN	22	17%	Recommended name: Ceruloplasmin EC = 1.16.3.1 Alternative name(s): Ferroxidase	P00450
DSG2_HUMAN	22	17%	Recommended name: Desmoglein-2 Alternative name(s): Cadherin family member 5 HDGC	Q14126
KIT_HUMAN	22	17%	Recommended name: Mast/stem cell growth factor receptor Kit Short name = SCFR EC = 2.7.10.1 Alternative name(s): Piebald trait protein Short name = PBT Proto-oncogene c-Kit Tyrosine-protein kinase Kit p145 c-kit v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog CD_antigen = CD117	P10721
TBB3_HUMAN	22	17%	Recommended name: Tubulin beta-3 chain Alternative name(s): Tubulin beta-4 chain Tubulin beta-III	Q13509
CH10_HUMAN	21	16%	Recommended name: 10 kDa heat shock protein, mitochondrial Short name = Hsp10 Alternative name(s): 10 kDa chaperonin Chaperonin 10	P61604

TABLE 12-continued

Protein (Uniprot)	Occurrence Across 131 Panels	Percentage Occurrence Across 131 Panels	Protein Names	Uniprot Accession No.
ISLR_HUMAN	21	16%	Short name = CPN10 Early-pregnancy factor Short name = EPF Immunoglobulin superfamily containing leucine-rich repeat protein	O14498
MASP1_HUMAN	21	16%	Recommended name: Mannan-binding lectin serine protease 1 EC = 3.4.21. - Alternative name(s): Complement factor MASP-3 Complement-activating component of Ra- reactive factor Mannose-binding lectin-associated serine protease 1 Short name = MASP-1 Mannose-binding protein-associated serine protease Ra-reactive factor serine protease p100 Short name = RaRF Serine protease 5 Cleaved into the following 2 chains: 1. Mannan-binding lectin serine protease 1 heavy chain 2. Mannan-binding lectin serine protease 1 light chain	P48740
ICAM3_HUMAN	20	15%	Recommended name: Intercellular adhesion molecule 3 Short name = ICAM-3 Alternative name(s): CDw50 ICAM-R CD_antigen = CD50	P32942
PTPRJ_HUMAN	20	15%	Recommended name: Receptor-type tyrosine-protein phosphatase eta Short name = Protein-tyrosine phosphatase eta Short name = R-PTP-eta EC = 3.1.3.48 Alternative name(s): Density-enhanced phosphatase 1 Short name = DEP-1 HPTP eta Protein-tyrosine phosphatase receptor type J Short name = R-PTP-J CD_antigen = CD148	Q12913
A1AG1_HUMAN	19	15%	Recommended name: Alpha-1-acid glycoprotein 1 Short name = AGP 1 Alternative name(s): Orosomucoid-1 Short name = OMD 1	P02763
CD59_HUMAN	18	14%	Recommended name: CD59 glycoprotein Alternative name(s): 1F5 antigen 20 kDa homologous restriction factor Short name = HRF-20 Short name = HRF20 MAC-inhibitory protein Short name = MAC-IP MEM43 antigen Membrane attack complex inhibition factor Short name = MACIF Membrane inhibitor of reactive lysis Short name = MIRL Protectin CD_antigen = CD59	P13987
MDHM_HUMAN	18	14%	Recommended name: Malate dehydrogenase, mitochondrial	P40926
PVR_HUMAN	18	14%	Recommended name: Poliovirus receptor Alternative name(s): Nectin-like protein 5	P15151

TABLE 12-continued

Protein (Uniprot)	Occurrence Across 131 Panels	Percentage Occurrence Across 131 Panels	Protein Names	Uniprot Accession No.
SEM3G_HUMAN	18	14%	Short name = NECL-5 CD_antigen = CD155 Recommended name: Semaphorin-3G Alternative name(s): Semaphorin sem2	Q9NS98
CO6A3_HUMAN	17	13%	Collagen alpha-3(VI) chain	P12111
MMP9_HUMAN	17	13%	Recommended name: Matrix metalloproteinase-9 Short name = MMP-9 EC = 3.4.24.35 Alternative name(s): 92 kDa gelatinase 92 kDa type IV collagenase Gelatinase B Short name = GELB Cleaved into the following 2 chains: 1.67 kDa matrix metalloproteinase-9 2.82 kDa matrix metalloproteinase-9	P14780
TETN_HUMAN	17	13%	Recommended name: Tetranectin Short name = TN Alternative name(s): C-type lectin domain family 3 member B Plasminogen kringle 4-binding protein	P05452
TNF12_HUMAN	17	13%	Recommended name: Tumor necrosis factor ligand superfamily member 12 Alternative name(s): APO3 ligand TNF-related weak inducer of apoptosis Short name = TWEAK Cleaved into the following 2 chains: 1. Tumor necrosis factor ligand superfamily member 12, membrane form 2. Tumor necrosis factor ligand superfamily member 12, secreted form	O43508
BST1_HUMAN	16	12%	Recommended name: ADP-ribosyl cyclase 2 EC = 3.2.2.5 Alternative name(s): Bone marrow stromal antigen 1 Short name = BST-1 Cyclic ADP-ribose hydrolase 2 Short name = cADPr hydrolase 2 CD_antigen = CD157	Q10588
CO1A1_HUMAN	16	12%	Recommended name: Collagen alpha-1(XVIII) chain Cleaved into the following chain: 1. Endostatin	P39060
CRP_HUMAN	16	12%	Recommended name: C-reactive protein Cleaved into the following chain: 1. C-reactive protein(1-205)	P02741
PLSL_HUMAN	16	12%	Recommended name: Plastin-2 Alternative name(s): L-plastin LC64P Lymphocyte cytosolic protein 1 Short name = LCP-1	P13796
BGH3_HUMAN	15	11%	Recommended name: Transforming growth factor-beta-induced protein ig-h3 Short name = Beta ig-h3 Alternative name(s): Kerato-epithelin RGD-containing collagen-associated protein Short name = RGD-CAP	Q15582
CD44_HUMAN	15	11%	Recommended name: CD44 antigen Alternative name(s): CDw44 Epican	P16070

TABLE 12-continued

Protein (Uniprot)	Occurrence Across131 Panels	Percentage Occurrence Across 131 Panels	Protein Names	Uniprot Accession No.
ENOA_HUMAN	15	11%	<p>Extracellular matrix receptor III Short name = ECMR-III GP90 lymphocyte homing/adhesion receptor HUTCH-I Heparan sulfate proteoglycan Hermes antigen Hyaluronate receptor Phagocytic glycoprotein 1 Short name = PGP-1 Phagocytic glycoprotein I Short name = PGP-I CD_antigen = CD44 Recommended name: Alpha-enolase EC = 4.2.1.11 Alternative name(s): 2-phospho-D-glycerate hydro-lyase C-myc promoter-binding protein Enolase 1 MBP-1 MPB-1 Non-neural enolase Short name = NNE Phosphopyruvate hydratase Plasminogen-binding protein</p>	P06733
LUM_HUMAN	15	11%		
SCF_HUMAN	15	11%	<p>Recommended name: Kit ligand Alternative name(s): Mast cell growth factor Short name = MGF Stem cell factor Short name = SCF c-Kit ligand Cleaved into the following chain: 1. Soluble KIT ligand Short name = sKITLG</p>	P21583
UGPA_HUMAN	15	11%	<p>Recommended name: UTP--glucose-1-phosphate uridylyltransferase EC = 2.7.7.9 Alternative name(s): UDP-glucose pyrophosphorylase Short name = UDPGP Short name = UGPase</p>	Q16851
ENPL_HUMAN	14	11%	<p>Recommended name: Endoplasmic Alternative name(s): 94 kDa glucose-regulated protein Short name = GRP-94 Heat shock protein 90 kDa beta member 1 Tumor rejection antigen 1 gp96 homolog</p>	P14625
GDIR2_HUMAN	14	11%	<p>Recommended name: Rho GDP-dissociation inhibitor 2 Short name = Rho GDI 2 Alternative name(s): Ly-GDI Rho-GDI beta</p>	P52566
GELS_HUMAN	14	11%	<p>Recommended name: Gelsolin Alternative name(s): AGEL Actin-depolymerizing factor Short name = ADF Brevin</p>	P06396
SODM_HUMAN	14	11%	<p>Recommended name: Superoxide dismutase [Mn], mitochondrial</p>	P04179
TPIS_HUMAN	14	11%	<p>Recommended name: Triosephosphate isomerase Short name = TIM EC = 5.3.1.1 Alternative name(s): Triose-phosphate isomerase</p>	P60174

TABLE 12-continued

Protein (Uniprot)	Occurrence Across 131 Panels	Percentage Occurrence Across 131 Panels	Protein Names	Uniprot Accession No.
TENA_HUMAN	13	10%	Recommended name: Tenascin Short name = TN Alternative name(s): Cytotactin GMEM GP 150-225 Glioma-associated-extracellular matrix antigen Hexabrachion JI Myotendinous antigen Neuronectin Tenascin-C Short name = TN-C	P24821
ZA2G_HUMAN	13	10%	Recommended name: Zinc-alpha-2-glycoprotein Short name = Zn-alpha-2-GP Short name = Zn-alpha-2-glycoprotein	P25311
LEG1_HUMAN	11	8%	Recommended name: Galectin-1 Short name = Gal-1 Alternative name(s): 14 kDa laminin-binding protein Short name = HLBP14 14 kDa lectin Beta-galactoside-binding lectin L-14-I Galaptin HBL HPL Lactose-binding lectin 1 Lectin galactoside-binding soluble 1 Putative MAPK-activating protein PM12 S-Lac lectin 1	P09382
FOLH1_HUMAN	9	7%	Recommended name: Glutamate carboxypeptidase 2 EC = 3.4.17.21 Alternative name(s): Cell growth-inhibiting gene 27 protein Folate hydrolase 1 Folylpoly-gamma-glutamate carboxypeptidase Short name = FGCP Glutamate carboxypeptidase II Short name = GCPII Membrane glutamate carboxypeptidase Short name = mGCP N-acetylated-alpha-linked acidic dipeptidase I Short name = NAALADase I Prostate-specific membrane antigen Short name = PSM Short name = PSMA Pteroylpoly-gamma-glutamate carboxypeptidase	Q04609
PLXC1_HUMAN	9	7%		
PTGIS_HUMAN	9	7%	Recommended name: Prostacyclin synthase EC = 5.3.99.4 Alternative name(s): Prostaglandin I2 synthase	Q16647

60

## Step 3: Significance and Occurrence

To find high performing panels, 10,000 trials were performed where on each trial the combined AUC of a random panel of 15 proteins selected from Table 12 was estimated. To calculate the combined AUC of each panel of 15 proteins, the highest intensity normalized transition was utilized. Logistic

65

regression was used to calculate the AUC of the panel of 15 across all small samples. 131 panels of 15 proteins had combined AUC above 0.80, as shown in FIG. 1. (The significance by study separated into small (<2.0 cm) and large (>2.0 cm) PN are shown in FIG. 2). The resilience of the panels persisted despite site based variation in the samples as shown in FIG. 3. The panels are listed in Table 13.

TABLE 13

AUC	P1	P2	P3	P4	P5	P6	P7	P8
0.8282	CD59	CALU	LDHB	ALDOA	DSG2	MDHM	TENA	6PGD
0.8255	CD59	TSP1	KIT	ISLR	ALDOA	DSG2	1433Z	CD14
0.8194	S10A6	ALDOA	PVR	TSP1	CD44	CH10	PEDF	APOE
0.8189	ALDOA	LEG1	CALU	LDHB	TETN	FOLH1	MASP1	1433Z
0.8187	PVR	CD59	CRP	ALDOA	GRP78	DSG2	6PGD	CD14
0.8171	AMPN	IBP3	CALU	CD44	BGH3	GRP78	1433Z	6PGD
0.8171	CALU	CH10	ALDOA	BST1	MDHM	VTNC	APOE	CD14
0.8165	LDHB	CO6A3	CD44	A1AG1	GRP78	DSG2	MDHM	VTNC
0.8163	TPIS	CD59	S10A6	CALU	ENPL	CH10	ALDOA	DSG2
0.8163	LEG1	AMPN	S10A6	CALU	ISLR	ENOA	VTNC	6PGD
0.8161	AMPN	S10A6	TSP1	MPRI	VTNC	LUM	6PGD	APOE
0.8159	ALDOA	AMPN	TSP1	BGH3	GRP78	PTPRJ	MASP1	CERU
0.8159	ALDOA	CO6A3	MPRI	SEM3G	CERU	LUM	APOE	CD14
0.8159	AMPN	CALU	ISLR	SODM	CERU	LUM	6PGD	APOE
0.8159	CALU	PEDF	CRP	GRP78	VTNC	1433Z	CD14	FRIL
0.8157	TPIS	LEG1	S10A6	LDHB	TSP1	ENPL	MDHM	6PGD
0.8155	CALU	CRP	ALDOA	SODM	SEM3G	1433Z	FRIL	G3P
0.8153	CALU	MPRI	ALDOA	PEDF	DSG2	CERU	APOE	G3P
0.814	LEG1	COIA1	AMPN	S10A6	TSP1	MPRI	PEDF	GRP78
0.8138	TSP1	KIT	CERU	6PGD	APOE	CD14	FRIL	G3P
0.8132	S10A6	COIA1	AMPN	TSP1	PEDF	ISLR	PTPRJ	CERU
0.8128	TPIS	LEG1	AMPN	S10A6	IBP3	CALU	DSG2	PTPRJ
0.8128	TPIS	AMPN	TSP1	PEDF	A1AG1	MPRI	ALDOA	VTNC
0.8124	ALDOA	CALU	LDHB	PLSL	PEDF	MASP1	6PGD	APOE
0.8124	AMPN	S10A6	TSP1	ENOA	GRP78	6PGD	APOE	FRIL
0.812	IBP3	TSP1	CRP	A1AG1	SCF	ALDOA	PEDF	DSG2
0.8106	COIA1	CALU	CD44	BGH3	ALDOA	TETN	BST1	LUM
0.8106	TSP1	PLSL	CRP	ALDOA	GRP78	MDHM	APOE	FRIL
0.8099	CD59	CALU	ENPL	CD44	ALDOA	TENA	6PGD	FRIL
0.8097	AMPN	S10A6	IBP3	A1AG1	MPRI	ALDOA	GRP78	FRIL
0.8093	ALDOA	S10A6	TSP1	ENPL	PEDF	A1AG1	GRP78	APOE
0.8093	PVR	IBP3	LDHB	SCF	TNF12	LUM	1433Z	FRIL
0.8093	CALU	LDHB	CO6A3	PEDF	CH10	BGH3	PTPRJ	ALDOA
0.8087	ALDOA	AMPN	ENPL	KIT	MPRI	GRP78	LUM	1433Z
0.8087	CD59	S10A6	IBP3	TSP1	ENPL	SODM	MDHM	6PGD
0.8083	ALDOA	AMPN	S10A6	IBP3	PLSL	CRP	SCF	MPRI
0.8081	PVR	IBP3	TSP1	CRP	ALDOA	SODM	MDHM	TNF12
0.8081	S10A6	LDHB	ENPL	PLSL	CH10	CERU	FRIL	G3P
0.8081	IBP3	LDHB	PEDF	MPRI	SEM3G	VTNC	APOE	CD14
0.8079	ALDOA	AMPN	CALU	PLSL	PEDF	CH10	MASP1	TNF12
0.8077	S10A6	IBP3	LDHB	MDHM	ZA2G	FRIL	G3P	HYOU1
0.8077	CD59	S10A6	LDHB	TSP1	CD44	ISLR	CERU	1433Z
0.8077	AMPN	CALU	LDHB	TSP1	PLSL	CD44	ALDOA	TETN
0.8075	TPIS	AMPN	S10A6	TSP1	CH10	COIA1	CERU	ZA2G
0.8073	CALU	PEDF	MPRI	ISLR	BGH3	ENOA	CERU	1433Z
0.8071	TPIS	CALU	CO6A3	KIT	DSG2	MASP1	6PGD	APOE
0.8071	LEG1	COIA1	TSP1	CD44	MPRI	ALDOA	FOLH1	TNF12
0.8065	AMPN	S10A6	CALU	CO6A3	TSP1	PLSL	KIT	MASP1
0.8063	S10A6	TSP1	A1AG1	BGH3	ZA2G	1433Z	FRIL	G3P
0.8063	CALU	KIT	ENOA	6PGD	APOE	CD14	G3P	ICAM3
0.8061	AMPN	MPRI	GRP78	DSG2	TENA	APOE	CD14	FRIL
0.8059	TPIS	IBP3	TSP1	PEDF	TNF12	1433Z	6PGD	APOE
0.8059	CALU	LDHB	PLSL	CRP	PEDF	SEM3G	MDHM	APOE
0.8058	ALDOA	TSP1	PLSL	CD44	KIT	CRP	ISLR	TNF12
0.8058	TPIS	TSP1	MPRI	ISLR	ALDOA	PEDF	GRP78	SEM3G
0.8054	ALDOA	S10A6	CALU	CRP	A1AG1	VTNC	TENA	ZA2G
0.8054	TPIS	CO6A3	TSP1	MPRI	DSG2	TNF12	FRIL	G3P
0.8054	CALU	LDHB	DSG2	1433Z	CD14	FRIL	G3P	HYOU1
0.805	CALU	MPRI	ENOA	FOLH1	LUM	ZA2G	APOE	CD14
0.8048	PVR	S10A6	IBP3	PEDF	ALDOA	BST1	MDHM	VTNC
0.8048	AMPN	CALU	CH10	DSG2	TNF12	CERU	6PGD	APOE
0.8046	ALDOA	LDHB	TSP1	KIT	ISLR	DSG2	MASP1	1433Z
0.8046	ALDOA	COIA1	CD59	IBP3	PTPRJ	SEM3G	CERU	CD14
0.8046	PVR	CD59	S10A6	PLSL	PEDF	CH10	SCF	BST1
0.8046	COIA1	IBP3	MASP1	DSG2	TENA	ZA2G	1433Z	APOE
0.8042	BGH3	CD59	CALU	LDHB	CO6A3	SODM	TENA	APOE
0.8042	IBP3	TSP1	ENPL	CH10	CD14	FRIL	G3P	HYOU1
0.8042	IBP3	TSP1	KIT	ZA2G	6PGD	APOE	CD14	FRIL
0.804	TPIS	BGH3	S10A6	LDHB	CO6A3	CH10	PEDF	TENA
0.804	CALU	LDHB	BGH3	TETN	FOLH1	TNF12	VTNC	FRIL
0.8038	TPIS	PVR	COIA1	CALU	SCF	MPRI	ALDOA	ENOA
0.8036	S10A6	TPIS	COIA1	CD59	CO6A3	TSP1	MPRI	ALDOA
0.8036	LEG1	CD59	AMPN	CALU	CH10	GRP78	SEM3G	TETN
0.8036	AMPN	S10A6	TSP1	ENPL	PEDF	SODM	FOLH1	6PGD
0.8036	S10A6	CALU	MASP1	A1AG1	MPRI	ALDOA	VTNC	TENA
0.8036	IBP3	CALU	PLSL	CD44	KIT	CERU	6PGD	CD14
0.8036	TSP1	PLSL	FOLH1	COIA1	TNF12	VTNC	6PGD	FRIL
0.8034	ALDOA	BGH3	CD59	TSP1	KIT	CH10	SODM	VTNC

TABLE 13-continued

0.8034	S10A6	CALU	LDHB	TSP1	GRP78	1433Z	6PGD	G3P	
0.8032	S10A6	CALU	TSP1	KIT	CH10	PEDF	GRP78	SEM3G	
0.8032	TSP1	MASP1	CRP	ALDOA	GRP78	TETN	TNF12	1433Z	
0.803	AMPN	TSP1	KIT	MPRI	SEM3G	TETN	DSG2	1433Z	
0.803	CALU	CO6A3	PLSL	A1AG1	ALDOA	GRP78	6PGD	APOE	
0.8028	COIA1	CD59	AMPN	TSP1	KIT	ISLR	ALDOA	MDHM	
0.8024	S10A6	CD44	SCF	MPRI	ISLR	ALDOA	APOE	FRIL	
0.8024	S10A6	TSP1	ALDOA	SODM	ENOA	BST1	FRIL	HYOU1	
0.8024	IBP3	TSP1	SCF	ALDOA	SODM	DSG2	VTNC	1433Z	
0.802	ALDOA	TSP1	PLSL	CD44	CH10	A1AG1	ENOA	TETN	
0.802	LEG1	CALU	LDHB	TSP1	CH10	ALDOA	MDHM	APOE	
0.802	CD59	IBP3	TSP1	A1AG1	MPRI	PTPRJ	6PGD	APOE	
0.802	IBP3	TSP1	CRP	BST1	TNF12	VTNC	1433Z	FRIL	
0.8018	LEG1	S10A6	IBP3	CALU	TSP1	MASP1	A1AG1	SCF	
0.8018	COIA1	CD59	AMPN	CALU	MASP1	BST1	VTNC	CERU	
0.8018	AMPN	ALDOA	SODM	GRP78	MDHM	VTNC	6PGD	FRIL	
0.8018	LDHB	CO6A3	ALDOA	SEM3G	DSG2	6PGD	APOE	FRIL	
0.8016	S10A6	LDHB	SCF	MPRI	ALDOA	PEDF	ENOA	SEM3G	
0.8016	LDHB	CO6A3	TSP1	1433Z	APOE	CD14	FRIL	G3P	
0.8014	ALDOA	PEDF	MPRI	ISLR	FOLH1	TNF12	MASP1	CERU	
0.8014	COIA1	PEDF	CRP	A1AG1	ENOA	CERU	FRIL	G3P	
0.8014	CD59	IBP3	TSP1	KIT	MASP1	ENOA	TNF12	CD14	
0.8014	LDHB	KIT	SCF	BGH3	SEM3G	VTNC	1433Z	FRIL	
0.8013	PVR	AMPN	LDHB	CD44	DSG2	TETN	MDHM	FRIL	
0.8013	S10A6	LDHB	TSP1	ISLR	LUM	G3P	HYOU1	ICAM3	
0.8013	CALU	A1AG1	MPRI	ALDOA	PEDF	DSG2	VTNC	ZA2G	
0.8013	TSP1	ENPL	KIT	SODM	SEM3G	DSG2	TETN	LUM	
0.8013	TSP1	PLSL	ISLR	ALDOA	ENOA	MDHM	APOE	G3P	
0.8011	ALDOA	AMPN	CO6A3	SEM3G	APOE	CD14	FRIL	G3P	
0.8011	TPIS	BGH3	AMPN	S10A6	CALU	LDHB	KIT	TENA	
0.8011	COIA1	IBP3	TSP1	A1AG1	TETN	DSG2	6PGD	FRIL	
0.8011	AMPN	S10A6	IBP3	CALU	KIT	SCF	ALDOA	APOE	
0.8011	IBP3	A1AG1	PEDF	SEM3G	MDHM	TNF12	VTNC	1433Z	
0.8009	ALDOA	BGH3	AMPN	LDHB	TSP1	PLSL	MPRI	ISLR	
0.8009	LEG1	COIA1	IBP3	CH10	MASP1	SCF	ALDOA	TNF12	
0.8009	AMPN	ENPL	ALDOA	TETN	FOLH1	BST1	ZA2G	6PGD	
0.8009	CALU	CO6A3	ENPL	ALDOA	GRP78	PTPRJ	VTNC	APOE	
0.8009	TSP1	CH10	PTPRJ	TETN	TNF12	VTNC	TENA	1433Z	
0.8007	CD59	S10A6	IBP3	CO6A3	TSP1	KIT	ISLR	GRP78	
0.8007	AMPN	TSP1	KIT	SCF	TETN	ZA2G	1433Z	6PGD	
0.8007	S10A6	IBP3	TSP1	CD44	PEDF	A1AG1	PTPRJ	SODM	
0.8007	CALU	CO6A3	TSP1	CH10	SCF	BGH3	ALDOA	ENOA	
0.8007	ENPL	CD44	MASP1	GRP78	1433Z	CD14	FRIL	G3P	
0.8005	TPIS	LEG1	LDHB	TSP1	MASP1	A1AG1	MPRI	ALDOA	
0.8005	PEDF	CRP	ISLR	ALDOA	GRP78	PTPRJ	ZA2G	6PGD	
0.8003	ALDOA	S10A6	CALU	CRP	BGH3	TETN	6PGD	CD14	
0.8003	AMPN	TSP1	A1AG1	MPRI	ISLR	ALDOA	MASP1	LUM	
0.8003	CO6A3	TSP1	SCF	MPRI	ISLR	FOLH1	1433Z	APOE	
0.8001	S10A6	IBP3	TSP1	KIT	TETN	COIA1	CERU	6PGD	
0.8001	S10A6	CALU	CH10	ISLR	ALDOA	SODM	PTPRJ	MDHM	
0.8001	IBP3	TSP1	ENPL	CH10	CRP	ISLR	ALDOA	SODM	
0.8001	IBP3	TSP1	PTPRJ	ALDOA	BST1	LUM	1433Z	APOE	
0.8001	LDHB	TSP1	MPRI	GRP78	SEM3G	LUM	ZA2G	FRIL	
		AUC	P9	P10	P11	P12	P13	P14	P15
		0.8282	APOE	FRIL	G3P	HYOU1	LRP1	RAN	HXK1
		0.8255	FRIL	HYOU1	LRP1	PROF1	TBB3	FINC	CEAM8
		0.8194	FRIL	G3P	HYOU1	LRP1	TBB3	CLIC1	RAN
		0.8189	APOE	G3P	HYOU1	PRDX1	PROF1	ILK	HXK1
		0.8187	FRIL	G3P	PRDX1	ILK	FINC	GSLG1	HXK1
		0.8171	CD14	FRIL	G3P	LRP1	TBB3	FINC	RAN
		0.8171	FRIL	G3P	ICAM3	PRDX1	PROF1	PVR	HXK1
		0.8165	1433Z	FRIL	G3P	S10A6	FINC	GSLG1	HXK1
		0.8163	6PGD	FRIL	G3P	HYOU1	ICAM3	PRDX1	FINC
		0.8163	APOE	G3P	LRP1	UGPA	RAN	CEAM8	HXK1
		0.8161	CD14	FRIL	G3P	LRP1	PROF1	RAN	CEAM8
		0.8159	6PGD	FRIL	G3P	HYOU1	LRP1	PRDX1	CEAM8
		0.8159	FRIL	G3P	LRP1	TBB3	FINC	GSLG1	HXK1
		0.8159	CD14	FRIL	G3P	PRDX1	CLIC1	ILK	HXK1
		0.8159	G3P	TBB3	ILK	GELS	FINC	RAN	GSLG1
		0.8157	APOE	FRIL	G3P	HYOU1	CLIC1	ILK	HXK1
		0.8155	HYOU1	LRP1	PRDX1	PROF1	FINC	RAN	GSLG1
		0.8153	HYOU1	PLXC1	PRDX1	ILK	CEAM8	HXK1	BST1
		0.814	CERU	FRIL	G3P	PLXC1	PRDX1	ILK	HXK1
		0.8138	HYOU1	PLXC1	RAN	CEAM8	HXK1	BST1	MMP9
		0.8132	6PGD	CD14	FRIL	HYOU1	FINC	GSLG1	BST1
		0.8128	BST1	6PGD	G3P	HYOU1	ILK	FINC	HXK1
		0.8128	1433Z	APOE	FRIL	G3P	LRP1	PTGIS	RAN
		0.8124	CD14	FRIL	G3P	GDIR2	FINC	GSLG1	HXK1

TABLE 13-continued

0.8124	GDIR2	LRP1	CLIC1	FINC	GSLG1	HXX1	BST1
0.812	1433Z	APOE	FRIL	LRP1	PRDX1	PROF1	FINC
0.8106	1433Z	6PGD	FRIL	G3P	HYOU1	PRDX1	CLIC1
0.8106	G3P	PRDX1	UGPA	ILK	CEAM8	GSLG1	HXX1
0.8099	G3P	HYOU1	PRDX1	PROF1	FINC	GSLG1	HXX1
0.8097	G3P	HYOU1	LRP1	PTGIS	ILK	FINC	MMP9
0.8093	CD14	FRIL	G3P	LRP1	PLXC1	CLIC1	GSLG1
0.8093	G3P	GDIR2	PRDX1	UGPA	CLIC1	FINC	HXX1
0.8093	SEM3G	MASP1	G3P	HYOU1	FINC	CEAM8	HXX1
0.8087	6PGD	CD14	FRIL	HYOU1	TBB3	CLIC1	FINC
0.8087	FRIL	G3P	HYOU1	LRP1	FINC	CEAM8	HXX1
0.8083	GRP78	CERU	CD14	FRIL	LRP1	FINC	CEAM8
0.8081	TENA	FRIL	G3P	HYOU1	PROF1	RAN	HXX1
0.8081	HYOU1	ICAM3	PLXC1	CLIC1	ILK	FINC	GSLG1
0.8081	FRIL	G3P	HYOU1	S10A6	CEAM8	GSLG1	HXX1
0.8079	LUM	6PGD	APOE	FRIL	HYOU1	RAN	HXX1
0.8077	LRP1	PTGIS	CLIC1	FINC	RAN	GSLG1	MMP9
0.8077	FRIL	G3P	HYOU1	LRP1	ILK	GSLG1	HXX1
0.8077	APOE	CD14	FRIL	G3P	LRP1	PRDX1	GSLG1
0.8075	6PGD	FRIL	G3P	LRP1	UGPA	ILK	HXX1
0.8073	6PGD	FRIL	G3P	HYOU1	LRP1	PRDX1	FINC
0.8071	CD14	FRIL	G3P	LRP1	AMPN	RAN	HXX1
0.8071	APOE	FRIL	HYOU1	LRP1	PTGIS	CLIC1	AMPN
0.8065	ALDOA	APOE	FRIL	G3P	TBB3	RAN	HXX1
0.8063	LRP1	PROF1	TBB3	UGPA	CLIC1	AMPN	RAN
0.8063	LRP1	PLXC1	PROF1	FINC	RAN	HXX1	MMP9
0.8061	G3P	LRP1	PLXC1	PROF1	PVR	FINC	CEAM8
0.8059	CD14	FRIL	G3P	LRP1	TBB3	RAN	GSLG1
0.8059	G3P	HYOU1	PRDX1	TBB3	ILK	RAN	HXX1
0.8058	APOE	CD14	FRIL	G3P	HYOU1	RAN	HXX1
0.8058	FRIL	G3P	HYOU1	PROF1	GELS	PVR	RAN
0.8054	6PGD	FRIL	G3P	HYOU1	ILK	GSLG1	HXX1
0.8054	HYOU1	ICAM3	PLXC1	TBB3	GELS	RAN	BST1
0.8054	PLXC1	PRDX1	PROF1	FINC	CEAM8	GSLG1	MMP9
0.805	G3P	HYOU1	ICAM3	PRDX1	UGPA	ILK	HXX1
0.8048	CD14	FRIL	G3P	HYOU1	PTGIS	FINC	RAN
0.8048	FRIL	G3P	LRP1	PRDX1	UGPA	RAN	CEAM8
0.8046	FRIL	G3P	GDIR2	HYOU1	RAN	GSLG1	HXX1
0.8046	FRIL	G3P	LRP1	PRDX1	FINC	GSLG1	MMP9
0.8046	FRIL	G3P	CLIC1	ILK	AMPN	FINC	HXX1
0.8046	CD14	FRIL	G3P	ICAM3	AMPN	FINC	HXX1
0.8042	G3P	HYOU1	S10A6	ILK	FINC	RAN	HXX1
0.8042	ICAM3	LRP1	PRDX1	PROF1	GELS	FINC	GSLG1
0.8042	GDIR2	HYOU1	LRP1	PRDX1	PROF1	CLIC1	HXX1
0.804	FRIL	G3P	HYOU1	LRP1	PRDX1	ILK	GSLG1
0.804	G3P	GDIR2	PRDX1	CLIC1	GELS	FINC	HXX1
0.8038	MASP1	APOE	FRIL	G3P	PRDX1	FINC	HXX1
0.8036	ENOA	6PGD	FRIL	G3P	GDIR2	LRP1	PRDX1
0.8036	APOE	G3P	HYOU1	ICAM3	RAN	CEAM8	HXX1
0.8036	APOE	FRIL	G3P	HYOU1	LRP1	HXX1	MMP9
0.8036	FRIL	G3P	PROF1	PTGIS	FINC	CEAM8	HXX1
0.8036	FRIL	G3P	HYOU1	PRDX1	FINC	CEAM8	HXX1
0.8036	G3P	LRP1	PRDX1	PROF1	GELS	FINC	RAN
0.8034	TENA	6PGD	G3P	HYOU1	LRP1	TBB3	ILK
0.8034	HYOU1	ICAM3	PROF1	ILK	GELS	AMPN	FINC
0.8032	MASP1	6PGD	CD14	FRIL	G3P	HYOU1	ILK
0.8032	APOE	CD14	G3P	HYOU1	PVR	RAN	HXX1
0.803	APOE	FRIL	G3P	TBB3	UGPA	PVR	RAN
0.803	CD14	FRIL	G3P	HYOU1	ICAM3	PRDX1	RAN
0.8028	CERU	LUM	ZA2G	APOE	FRIL	LRP1	MMP9
0.8024	G3P	HYOU1	PRDX1	GELS	FINC	CEAM8	HXX1
0.8024	LRP1	PROF1	CLIC1	GELS	FINC	CEAM8	GSLG1
0.8024	APOE	FRIL	G3P	LRP1	PRDX1	UGPA	PTPRJ
0.802	TENA	APOE	FRIL	G3P	TBB3	AMPN	GSLG1
0.802	FRIL	G3P	HYOU1	ILK	PVR	GSLG1	PTPRJ
0.802	FRIL	G3P	LRP1	ILK	RAN	CEAM8	MMP9
0.802	G3P	GDIR2	HYOU1	LRP1	PRDX1	TBB3	FINC
0.8018	ALDOA	SEM3G	VTNC	FRIL	G3P	LRP1	CLIC1
0.8018	6PGD	APOE	CD14	FRIL	HYOU1	PROF1	GSLG1
0.8018	G3P	HYOU1	LRP1	PTGIS	GELS	FINC	RAN
0.8018	G3P	HYOU1	ICAM3	PROF1	FINC	PTPRJ	HXX1
0.8016	APOE	FRIL	G3P	HYOU1	PRDX1	CLIC1	GSLG1
0.8016	HYOU1	PROF1	UGPA	CLIC1	RAN	CEAM8	PTPRJ
0.8014	6PGD	FRIL	G3P	HYOU1	PRDX1	FINC	HXX1
0.8014	GDIR2	LRP1	S10A6	GELS	FINC	GSLG1	HXX1
0.8014	FRIL	G3P	PRDX1	UGPA	FINC	PTPRJ	HXX1
0.8014	G3P	HYOU1	LRP1	PRDX1	PROF1	FINC	HXX1
0.8013	G3P	LRP1	PRDX1	ILK	FINC	HXX1	MMP9
0.8013	LRP1	PROF1	UGPA	ILK	FINC	PTPRJ	HXX1
0.8013	6PGD	FRIL	G3P	CLIC1	S10A6	ILK	PVR

TABLE 13-continued

0.8013	APOE	FRIL	G3P	HYOU1	CLIC1	RAN	HXX1
0.8013	GDIR2	LRP1	PTGIS	FINC	RAN	HXX1	MMP9
0.8011	GDIR2	HYOU1	ICAM3	PRDX1	FINC	HXX1	MMP9
0.8011	6PGD	APOE	G3P	LRP1	PROF1	GELS	MMP9
0.8011	GDIR2	HYOU1	LRP1	CLIC1	S10A6	PVR	GSLG1
0.8011	G3P	ICAM3	LRP1	GELS	FINC	RAN	CEAM8
0.8011	G3P	HYOU1	PRDX1	FINC	GSLG1	PTPRJ	HXX1
0.8009	APOE	FRIL	LRP1	PVR	FINC	RAN	PTPRJ
0.8009	CERU	APOE	CD14	FRIL	TBB3	ILK	FINC
0.8009	CD14	FRIL	CLIC1	S10A6	ILK	FINC	MMP9
0.8009	CD14	G3P	TBB3	CLIC1	GELS	RAN	HXX1
0.8009	6PGD	FRIL	G3P	HYOU1	RAN	HXX1	MMP9
0.8007	MDHM	CD14	FRIL	G3P	HYOU1	GSLG1	HXX1
0.8007	APOE	G3P	GDIR2	LRP1	PRDX1	TBB3	RAN
0.8007	CERU	APOE	FRIL	ICAM3	LRP1	UGPA	GSLG1
0.8007	TETN	LUM	APOE	FRIL	G3P	RAN	HXX1
0.8007	GDIR2	ICAM3	LRP1	PRDX1	PROF1	FINC	HXX1
0.8005	ENOA	FRIL	G3P	LRP1	UGPA	ILK	FINC
0.8005	G3P	HYOU1	PRDX1	TBB3	FINC	RAN	CEAM8
0.8003	FRIL	G3P	CLIC1	FINC	GSLG1	HXX1	MMP9
0.8003	6PGD	APOE	FRIL	ICAM3	TBB3	GSLG1	BST1
0.8003	G3P	HYOU1	ICAM3	PRDX1	UGPA	RAN	HXX1
0.8001	CD14	FRIL	G3P	PROF1	FINC	HXX1	MMP9
0.8001	VTNC	FRIL	G3P	CLIC1	ILK	AMPN	HXX1
0.8001	1433Z	G3P	HYOU1	LRP1	PRDX1	PROF1	CEAM8
0.8001	G3P	HYOU1	LRP1	PTGIS	TBB3	PVR	RAN
0.8001	G3P	ICAM3	PROF1	TBB3	FINC	RAN	GSLG1

To calculate the combined AUC of each panel of 15 proteins, the highest intensity normalized transition was utilized. Logistic regression was used to calculate the AUC of the panel of 15 across all small samples. 5 panels of 15 proteins had combined AUC above 0.80.

Finally, the frequency of each of the 67 proteins on the 131 panels listed in Table 13 is presented in Table 12 both as raw counts (column 2) and percentage (column 3). It is an important observation that the panel size of 15 was pre-selected to prove that there are diagnostic proteins and panels. Furthermore, there are numerous such panels. Smaller panels

selected from the list of 67 proteins can also be formed and can be generated using the same methods here.

Example 4

A Diagnostic Panel of 15 Proteins for Determining the Probability that a Blood Sample from a Patient with a PN of Size 2 cm or Less is Benign or Malignant

In Table 14 a logistic regression classifier trained on all small samples is presented.

TABLE 14

Protein	Transition	Transition column		Normalized By column	Normalized SEQ ID NO:	Logistic Regression Coefficient
		SEQ ID NO:	Normalized By			
ALDOA_HUMAN	ALQASALK_401.25_617.40	7	YGFIIEGHVVIPR_462.92_272.20	1	-1.96079	
BGH3_HUMAN	LTLPLNSVFK_658.40_804.50	8	YEVTVVSVR_526.29_759.50	2	2.21074	
CLIC1_HUMAN	LAALNPESNTAGLDIFAK_922.99_256.20	9	ASSIIEELFQDR_465.24_565.30	3	0.88028	
CO6A3_HUMAN	VAVVQYSR_518.77_767.40	10	ASSIIEELFQDR_465.24_565.30	3	-1.52046	
COIA1_HUMAN	AVGLAGTFR_446.26_721.40	11	YGFIIEGHVVIPR_462.92_272.20	1	-0.76786	
FINC_HUMAN	VPGTSTSATLTGLTR_487.94_446.30	12	FLNVLSPR_473.28_685.40	4	0.98842	
G3P_HUMAN	GALQNIIPASTGAAK_706.40_815.50	13	TASDFITK_441.73_710.40	5	0.58843	
ISLR_HUMAN	ALPGTPVASSQPR_640.85_841.50	14	FLNVLSPR_473.28_685.40	4	1.02005	
LRP1_HUMAN	TVLWPNGSLDIPAGR_855.00_400.20	15	YEVTVVSVR_526.29_759.50	2	-2.14383	
PRDX1_HUMAN	QITVNDLPVGR_606.30_428.30	16	YGFIIEGHVVIPR_462.92_272.20	1	-1.38044	
PROF1_HUMAN	STGGAPT FNVT VTK_690.40_503.80	17	TASDFITK_441.73_710.40	5	-1.78666	
PVR_HUMAN	SVDIWLRL_444.75_702.40	18	TASDFITK_441.73_710.40	5	2.26338	
TBB3_HUMAN	ISVYYNEASSHK_466.60_458.20	19	FLNVLSPR_473.28_685.40	4	-0.46786	
TETN_HUMAN	LDTLAQEVALLK_657.39_330.20	20	TASDFITK_441.73_710.40	5	-1.99972	
TPIS_HUMAN	VVFEQTK_425.74_652.30	21	YGFIIEGHVVIPR_462.92_272.20	1	2.65334	
Constant (C <sub>0</sub> )					21.9997	

The classifier has the structure

$$\text{Probability} = \frac{\exp(W)}{1 + \exp(W)}$$

$$W = C_0 + \sum_{i=1}^{15} C_i * P_i$$

Where  $C_0$  and  $C_i$  are logistic regression coefficients,  $P_i$  are logarithmically transformed normalized transition intensities. Samples are predicted as cancer if  $\text{Probability} \geq 0.5$  or as benign otherwise. In Table 14 the coefficients  $C_i$  appear in the sixth column,  $C_0$  in the last row, and the normalized transitions for each protein are defined by column 2 (protein transition) and column 4 (the normalizing factor).

The performance of this classifier, presented as a ROC plot, appears in FIG. 4. Overall AUC is 0.81. The performance can also be assessed by applying the classifier to each study site individually which yields the three ROC plots appearing in

FIG. 5. The resulting AUCs are 0.79, 0.88 and 0.78 for Laval, NYU and UPenn, respectively.

Example 5

The Program “Ingenuity”<sup>®</sup> was Used to Query the Blood Proteins that are Used to Identify Lung Cancer in Patients with Nodules that were Identified Using the Methods of the Present Invention

Using a subset of 35 proteins (Table 15) from the 67 proteins identified as a diagnostic panel (Table 13), a backward systems analysis was performed. Two networks were queried that are identified as cancer networks with the identified 35 proteins. The results show that the networks that have the highest percentage of “hits” when the proteins are queried that are found in the blood of patients down to the level of the nucleus are initiated by transcription factors that are regulated by either cigarette smoke or lung cancer among others. See also Table 16 and FIG. 6.

These results are further evidence that the proteins that were identified using the methods of the invention as diagnostic for lung cancer are prognostic and relevant.

TABLE 15

No.	Protein	Protein Name	Gene Symbol	Gene Name
1	6PGD_HUMAN	6-phosphogluconate dehydrogenase, decarboxylating	PGD	phosphogluconate dehydrogenase
2	AIFM1_HUMAN	Apoptosis-inducing factor 1, mitochondrial	AIFM1	apoptosis-inducing factor, mitochondrion-associated, 1
3	ALDOA_HUMAN	Fructose-bisphosphate aldolase A	ALDOA	aldolase A, fructose-bisphosphate
4	BGH3_HUMAN	Transforming growth factor-beta-induced protein ig-h3	TGFBI	transforming growth factor, beta-induced, 68 kDa
5	CD163A_HUMAN	Scavenger receptor cysteine-rich type 1 protein M130	CD163	CD163 molecule
6	CD14_HUMAN	Monocyte differentiation antigen CD14	CD14	CD14 molecule
7	COIA1_HUMAN	Collagen alpha-1(XVIII) chain	COL18A1	collagen, type XVIII, alpha 1
8	ERO1A_HUMAN	ERO1-like protein alpha	ERO1L	ERO1-like (S. cerevisiae)
9	FIBA_HUMAN	Fibrinogen alpha chain	FGA	fibrinogen alpha chain
10	FINC_HUMAN	Fibronectin	FN1	fibronectin 1
11	FOLH1_HUMAN	Glutamate carboxypeptidase 2	FOLH1	folate hydrolase (prostate-specific membrane antigen) 1
12	FRIL_HUMAN	Ferritin light chain	FTL	ferritin, light polypeptide
13	GELS_HUMAN	Gelsolin	GSN	gelsolin (amyloidosis, Finnish type)
14	GGH_HUMAN	Gamma-glutamyl hydrolase	GGH	gamma-glutamyl hydrolase (conjugase, folic acid-gammaglutamyl hydrolase)
15	GRP78_HUMAN	78 kDa glucose-regulated protein	HSPA5	heat shock 70 kDa protein 5 (glucose-regulated protein, 78 kDa)
16	GSLG1_HUMAN	Golgi apparatus protein 1	GLG1	golgi apparatus protein 1
17	GSTP1_HUMAN	Glutathione S-transferase P	GSTP1	glutathione S-transferase pi 1
18	IBP3_HUMAN	Insulin-like growth factor-binding protein 3	IGFBP3	insulin-like growth factor binding protein 3
19	ICAM1_HUMAN	Intercellular adhesion molecule 1	ICAM1	intercellular adhesion molecule 1
20	ISLR_HUMAN	Immunoglobulin superfamily containing leucine-rich repeat protein	ISLR	immunoglobulin superfamily containing leucine-rich repeat
21	LG3BP_HUMAN	Galectin-3-binding protein	LGALS3BP	lectin, galactoside-binding, soluble, 3 binding protein
22	LRP1_HUMAN	Prolow-density lipoprotein receptor-related protein 1	LRP1	low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)
23	LUM_HUMAN	Lumican	LUM	lumican
24	MASP1_HUMAN	Mannan-binding lectin serine protease 1	MASP1	mannan-binding lectin serine peptidase 1 (C4/C2 activating component of Ra-reactive factor)
25	PDIA3_HUMAN	Protein disulfide-isomerase A3	PDIA3	protein disulfide isomerase family A, member 3

TABLE 15-continued

No.	Protein	Protein Name	Gene Symbol	Gene Name
26	PEDF_HUMAN	Pigment epithelium-derived factor	SERPINF1	serpin peptidase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 1
27	PRDX1_HUMAN	Peroxiredoxin-1	PRDX1	peroxiredoxin 1
28	PROF1_HUMAN	Profilin-1	PFN1	profilin 1
29	PTPA_HUMAN	Serine/threonine-protein phosphatase 2A activator	PPP2R4	protein phosphatase 2A activator, regulatory subunit 4
30	PTPRJ_HUMAN	Receptor-type tyrosine-protein phosphatase eta	PTPRJ	protein tyrosine phosphatase, receptor type, J
31	RAP2B_HUMAN	Ras-related protein Rap-2b	RAP2B	RAP2B, member of RAS oncogene family
32	SEMA3G_HUMAN	Semaphorin-3G	SEMA3G	sema domain, immunoglobulin domain (Ig), short basic domain, secreted, (semaphorin) 3G
33	SODM_HUMAN	Superoxide dismutase [Mn], mitochondrial	SOD2	superoxide dismutase 2, mitochondrial
34	TETN_HUMAN	Tetranectin	CLEC3B	C-type lectin domain family 3, member B
35	TSP1_HUMAN	Thrombospondin-1	THBS1	thrombospondin 1

TABLE 16

Gene Name	Protein	Lung Cancer PubMed Associations	Sample Publications
NFE2L2 (NRF2)	nuclear factor (erythroid-derived 2)-like 2	92	Cigarette Smoking Blocks the Protective Expression of Nrf2/ARE Pathway . . . Molecular mechanisms for the regulation of Nrf2-mediated cell proliferation in non-small-cell lung cancers . . .
EGR-1	early growth response	38	Cigarette smoke-induced Egr-1 upregulates proinflammatory cytokines in pulmonary epithelial cells . . . EGR-1 regulates Ho-1 expression induced by cigarette smoke . . . Chronic hypoxia induces Egr-1 via activation of ERK1/2 and contributes to pulmonary vascular remodeling. Early growth response-1 induces and enhances vascular endothelial growth factor-A expression in lung cancer cells . . .

## Example 6

## Cooperative Proteins for Diagnosing Pulmonary Nodules

To achieve unbiased discovery of cooperative proteins, selected reaction monitoring (SRM) mass spectrometry (Ad-  
dona, Abbatiello et al. 2009) was utilized. SRM is a form of  
mass spectrometry that monitors predetermined and highly  
specific mass products of particularly informative (proteo-  
typic) peptides of selected proteins. These peptides are re-  
cognized as specific transitions in mass spectra. SRM pos-  
sesses the following required features that other technologies,  
notably antibody-based technologies, do not possess:

Highly multiplexed SRM assays can be rapidly and cost-  
effectively developed for tens or hundreds of proteins.

The assays developed are for proteins of one's choice and  
are not restricted to a catalogue of pre-existing assays.  
Furthermore, the assays can be developed for specific  
regions of a protein, such as the extracellular portion of  
a transmembrane protein on the cell surface of a tumor  
cell, or for a specific isoform.

SRM technology can be used from discovery to clinical  
testing. Peptide ionization, the foundation of mass spec-

45 trometry, is remarkably reproducible. Using a single  
technology platform avoids the common problem of  
translating an assay from one technology platform to  
another.

SRM has been used for clinical testing of small molecule  
analytes for many years, and recently in the development of  
biologically relevant assays [10].

Labeled and unlabeled SRM peptides are commercially  
available, together with an open-source library and data  
repository of mass spectra for design and conduct of SRM  
analyses. Exceptional public resources exist to accelerate  
assay development including the PeptideAtlas [11] and the  
Plasma Proteome Project [12, 13], the SRM Atlas and PAS-  
SEL, the PeptideAtlas SRM Experimental Library ([www-  
.systemsbio.org/passel](http://www.systemsbio.org/passel)).

Two SRM strategies that enhance technical performance  
were introduced. First, large scale SRM assay development  
introduces the possibility of monitoring false signals. Using  
an extension of expression correlation techniques [14], the  
rate of false signal monitoring was reduced to below 3%. This  
is comparable and complementary to the approach used by  
mProphet (Reiter, Rinner et al. 2011).

Second, a panel of endogenous proteins was used for nor-  
malization. However, whereas these proteins are typically

selected as “housekeeping” proteins (Lange, Picotti et al. 2008), proteins that were strong normalizers for the technology platform were identified. That is, proteins that monitored the effects of technical variation so that it could be controlled effectively. This resulted, for example, in the reduction of technical variation due to sample depletion of high abundance proteins from 23.8% to 9.0%. The benefits of endogenous signal normalization has been previously discussed (Price, Trent et al. 2007).

The final component of the strategy was to carefully design the discovery and validation studies using emerging best practices. Specifically, the cases (malignant nodules) and controls (benign nodules) were pairwise matched on age, nodule size, gender and participating clinical site. This ensures that the candidate markers discovered are not markers of age or variations in sample collection from site to site. The studies were well-powered, included multiple sites, a new site participated in the validation study, and importantly, were designed to address the intended use of the test. The careful selection and matching of samples resulted in an exceptionally valuable feature of the classifier. The classifier generates a score that is independent of nodule size and smoking status. As these are currently used risk factors for clinical management of IPNs, the classifier is a complementary molecular tool for use in the diagnosis of IPNs.

#### Selection of Biomarker Candidates for Assay Development

To identify lung cancer biomarkers in blood that originate from lung tumor cells, resected lung tumors and distal normal tissue of the same lobe were obtained. Plasma membranes were isolated from both endothelial and epithelial cells and analyzed by tandem mass spectrometry to identify cell surface proteins over expressed on tumor cells. Similarly, Golgi apparatus were isolated to identify over-secreted proteins from tumor cells. Proteins with evidence of being present in blood or secreted were prioritized resulting in a set of 217 proteins. See Example 7: Materials and Methods for details.

To ensure other viable lung cancer biomarkers were not overlooked, a literature search was performed and manually curated for lung cancer markers. As above, proteins with evidence of being present in blood or secreted were prioritized. This resulted in a set of 319 proteins. See Example 7: Materials and Methods for details.

The tissue (217) and literature (319) candidates overlapped by 148 proteins resulting in a final candidate list of 388 protein candidates. See Example 7: Materials and Methods.

#### Development of SRM Assays

SRM assays for the 388 proteins were developed using standard synthetic peptide techniques (See Example 7: Materials and Methods). Of the 388 candidates, SRM assays were successfully developed for 371 candidates. The 371 SRM assays were applied to benign and lung cancer plasma samples to evaluate detection rate in blood. 190 (51% success rate) of the SRM assays were detected. This success rate compares favorably to similar attempts to develop large scale SRM assays for detection of cancer markers in plasma.

Recently 182 SRM assays for general cancer markers were developed from 1172 candidates (16% success rate) [15]. Despite focusing only on lung cancer markers, the 3-fold increase in efficiency is likely due to sourcing candidates from cancer tissues with prior evidence of presence in blood. Those proteins of the 371 that were previously detected by mass spectrometry in blood had a 64% success rate of detection in blood whereas those without had a 35% success rate. Of the 190 proteins detected in blood, 114 were derived from the tissue-sourced candidates and 167 derived from the literature-sourced candidates (91 protein overlap). See Example 7: Materials and Methods and Table 6.

Typically, SRM assays are manually curated to ensure assays are monitoring the intended peptide. However, this becomes unfeasible for large scale SRM assays such as this 371 protein assay. More recently, computational tools such as mProphet (Reiter, Rinner et al. 2011) enable automated qualification of SRM assays. A complementary strategy to mProphet was introduced that does not require customization for each dataset set. It utilizes correlation techniques (Kearney, Butler et al. 2008) to confirm the identity of protein transitions with high confidence. In FIG. 7 a histogram of the Pearson correlations between every pair of transitions in the assay is presented. The correlation between a pair of transitions is obtained from their expression profiles over all 143 samples in the discovery study detailed below. As expected, transitions from the same peptide are highly correlated. Similarly, transitions from different peptide fragments of the same protein are also highly correlated. In contrast, transitions from different proteins are not highly correlated and enables a statistical analysis of the quality of a protein’s SRM assay. For example, if the correlation of transitions from two peptides from the same protein is above 0.5 then there is less than a 3% probability that the assay is false. See Example 7: Materials and Methods.

#### Classifier Discovery

A summary of the 143 samples used for classifier discovery appears in Table 17. Samples were obtained from three sites to avoid overfitting to a single site. Participating sites were Laval (Institut Universitaire de Cardiologie et de Pneumologie de Quebec), NYU (New York University) and UPenn (University of Pennsylvania). Samples were also selected to be representative of the intended use population in terms of nodule size (diameter), age and smoking status.

Benign and cancer samples were paired by matching on age, gender, site and nodule size (benign and cancer samples were required to have a nodule identified radiologically). The benign and cancer samples display a bias in smoking (pack years), however, the majority of benign and cancer samples were current or past smokers. In comparing malignant and benign samples, the intent was to find proteins that were markers of lung cancer; not markers of age, nodule size or differences in site sample collection. Note that cancer samples were pathologically confirmed and benign samples were either pathologically confirmed or radiologically confirmed (no tumor growth demonstrated over two years of CT scan surveillance).

TABLE 17

Clinical data summaries and demographic analysis for discovery and validation sets.						
	Discovery			Validation		
	Cancer	Benign	P value	Cancer	Benign	P value
Sample (total)	72	71		52	52	

TABLE 17-continued

Clinical data summaries and demographic analysis for discovery and validation sets.							
		Discovery			Validation		
		Cancer	Benign	P value	Cancer	Benign	P value
Sample (Center)	Laval	14	14	1.00†	13	12	0.89†
	NYU	29	28		6	9	
	UPenn	29	29		14	13	
	Vanderbilt	0	0		19	18	
Sample (Gender)	Male	29	28	1.00†	25	27	0.85†
	Female	43	43		27	25	
Sample (Smoking History)	Never	5	19	0.006†	3	15	0.006†
	Past	60	44		38	29	
	Current	6	6		11	7	
Age	No data	1	2		0	1	
	Median (quartile range)	65 (59-72)	64 (52-71)	0.46‡	63 (60-73)	62 (56-67)	0.03‡
Nodule Size (mm)	Median (quartile range)	13 (10-16)	13 (10-18)	0.69‡	16 (13-20)	15 (12-22)	0.68‡
	Pack-year§ (quartile range)	37 (20-52)	20 (0-40)	0.001‡	40 (19-50)	27 (0-50)	0.09‡

†Based on Fisher's exact test.

‡Based on Mann-Whitney test.

§No data (cancer, benign): Discovery (4, 6), Validation (2, 3)

The processing of samples was conducted in batches. Each batch contained a set of randomly selected cancer-benign pairs and three plasma standards, included for calibration and quality control purposes.

All plasma samples were immunodepleted, trypsin digested and analyzed by reverse phase HPLC-SRM-MS. Protein transitions were normalized using an endogenous protein panel. The normalization procedure was designed to reduce overall variability, but in particular, the variability introduced by the depletion step. Overall technical variability was reduced from 32.3% to 25.1% and technical variability due to depletion was reduced from 23.8% to 9.0%. Details of the sample analysis and normalization procedure are available in Example 7: Materials and Methods.

To assess panels of proteins, they were fit to a logistic regression model. Logistic regression was chosen to avoid the overfitting that can occur with non-linear models, especially when the number of variables measured (transitions) is similar or larger than the number of samples in the study. The

performance of a panel was measured by partial area under the curve (AUC) with sensitivity fixed at 90% (McClish 1989). Partial AUC correlates to high NPV performance while maximizing ROR.

To derive the 13 protein classifier, four criteria were used: The protein must have transitions that are reliably detected above noise across samples in the study.

The protein must be highly cooperative.

The protein must have transitions that are robust (high signal to noise, no interference, etc.)

The protein's coefficient within the logistic regression model must have low variability during cross validation, that is, it must be stable.

Details of how each of these criteria were applied appear in Example 7: Materials and Methods.

Finally, the 13 protein classifier was trained to a logistic regression model by Monte Carlo cross validation (MCCV) with a hold out rate of 20% and 20,000 iterations. The thirteen proteins for the rule-out classifier are listed in Table 18 along with their highest intensity transition and model coefficient.

TABLE 18

The 13 protein classifier.			
Protein	Transition	SEQ ID NO	Coefficient
Constant( $\alpha$ )			36.16
LRP1_HUMAN	TVLWPNGLSLDIPAGR_855.00_400.20	15	-1.59
BGH3_HUMAN	LTL LAPLNSVFK_658.40_804.50	8	1.73
COIA1_HUMAN	AVGLAGTFR_446.26_721.40	11	-1.56
TETN_HUMAN	LDTLAQEVALLK_657.39_330.20	20	-1.79
TSP1_HUMAN	GFLLLASLR_495.31_559.40	22	0.53
ALDOA_HUMAN	ALQASALK_401.25_617.40	7	-0.80
GRP78_HUMAN	TWNDPVQQDIK_715.85_260.20	23	1.41
ISLR_HUMAN	ALPGTPVASSQPR_640.85_841.50	14	1.40
FRIL_HUMAN	LGPEAGLGEYLFER_804.40_913.40	24	0.39
LG3BP_HUMAN	VEIFYR_413.73_598.30	25	-0.58
PRDX1_HUMAN	QITVNDLPVGR_606.30_428.30	16	-0.34
FIBA_HUMAN	NSLFEYQK_514.76_714.30	26	0.31
GSLG1_HUMAN	IIIQESALDYR_660.86_338.20	27	-0.70

## Validation of the Rule-Out Classifier

52 cancer and 52 benign samples (see Table 17) were used to validate the performance of the 13 protein classifier. All samples were independent of the discovery samples, in addition, over 36% of the validation samples were sourced from a new fourth site (Vanderbilt University). Samples were selected to be consistent with intended use and matched in terms of gender, clinical site and nodule size. We note a slight age bias, which is due to 5 benign samples from young patients. Anticipating a NPV of 90%, the 95% confidence interval is  $\pm 5\%$ .

At this point we refer to the 13 protein classifier trained on 143 samples the Discovery classifier. However, once validation is completed, to find the optimal coefficients for the classifier, it was retrained on all 247 samples (discovery and validation sets) as this is most predictive of future performance. We refer to this classifier as the Final classifier. The coefficients of the Final classifier appear in Table 21.

The performance of the Discovery and Final classifiers is summarized in FIG. 8. Reported are the NPV and ROR for the Discovery classifier when applied to the discovery set, the validation set. The NPV and ROR for the Final classifier are reported for all samples and also for all samples restricted to nodule size 8 mm to 20 mm (191 samples).

NPV and ROR are each reported as a fraction from 0 to 1. Similarly, the classifier produces a score between 0 and 1, which is the probability of cancer predicted by the classifier.

The discovery and validation curves for NPV and ROR are similar with the discovery curves superior as expected. This demonstrates the reproducibility of performance on an independent set of samples. A Discovery classifier rule out threshold of 0.40 achieves NPV of 96% and 90%, whereas ROR is 33% and 23%, for the discovery samples and the validation samples, respectively. Final classifier rule threshold of 0.60 achieves NPV of 91% and 90%, whereas ROR is 45% and 43%, for all samples and all samples restricted to be 8 mm-20 mm, respectively.

## Applications of the Classifier

FIG. 9 presents the application of the final classifier to all 247 samples from the discovery and validation sets. The intent of FIG. 9 is to contrast the clinical risk factors of smoking (measured in pack years) and nodule size (proportional to the size of each circle) to the classifier score assigned to each sample.

First, note the density of cancer samples with high classifier scores. The classifier has been designed to detect a cancer signature in blood with high sensitivity. As a consequence, to the left of the rule out threshold (0.60) there are very few (<10%) cancer samples, assuming cancer prevalence of 25% [16, 17].

Third is the observation that nodule size does not appear to increase with the classifier score. Both large and small nodules are spread across the classifier score spectrum. Similarly, although there are a few very heavy smokers with very high classifier scores, increased smoking does not seem to increase with classifier score. To quantify this observation the correlation between the classifier score and nodule size, smoking and age were calculated and appear in Table 19. In all cases there is no significant relationship between the classifier score and the risk factors. The one exception is a weak correlation between benign classifier scores and benign ages. However, this correlation is so weak that the classifier score increases by only 0.04 every 10 years.

TABLE 19

Correlation between classifier scores and clinical risk factors.			
	Age	Nodule Size	Smoking
Benign	0.25	-0.06	0.11
Cancer	0.01	-0.01	0.06

This lack of correlation has clinical utility. It implies that the classifier provides molecular information about the disease status of an IPN that is incremental upon risk factors such as nodule size and smoking status. Consequently, it is a clinical tool for physicians to make more informed decisions around the clinical management of an IPN.

To visual how this might be accomplished, we demonstrate how the cancer probability score generated by the classifier can be related to cancer risk (see FIG. 11)

At a given classifier score, some percentage of all cancer nodules will have a smaller score. This is the sensitivity of the classifier. For example, at classifier score 0.8, 47% of cancer patients have a lower score, at classifier score 0.7, 28% of cancer patients have a lower score, at classifier score 0.5, only 9% are lower and finally at score 0.25, only 4% are lower. This enables a physician to interpret a patient's classifier score in terms of relative risk.

## The Molecular Foundations of the Classifier

The goal was to identify the molecular signature of a malignant pulmonary nodule by selecting proteins that were the cooperative, robustly detected by SRM and stable within the classifier. How well associated with lung cancer is the derived classifier? Is there a molecular foundation for the perturbation of these 13 proteins in blood? And finally, how unique is the classifier among other possible protein combinations?

To answer these questions the 13 proteins of the classifier were submitted for pathway analysis using IPA (Ingenuity Systems, www.ingenuity.com). The first step was to work from outside the cell inwards to identify the transcription factors most likely to cause a modulation of these 13 proteins. The five most significant were FOS, NRF2, AHR, HD and MYC. FOS is common to many forms of cancer. However, NRF2 and AHR are associated with lung cancer, response to oxidative stress and lung inflammation. MYC is associated with lung cancer and response to oxidative stress while HD is associated with lung inflammation and response to oxidative stress.

The 13 classifier proteins are also highly specific to these three networks (lung cancer, response to oxidative stress and lung inflammation). This is summarized in FIG. 10 where the classifier proteins (green), transcription factors (blue) and the three merged networks (orange) are depicted. Only ISLR is not connected through these three lung specific networks to the other proteins, although it is connected through cancer networks not specific to cancer. In summary, the modulation of the 13 classifier proteins can be tracked back to a few transcription factors specific to lung cancer, lung inflammation and oxidative stress networks.

To address the question of classifier uniqueness, every classifier from the 21 robust and cooperative proteins was formed (Table 20). Due to the computational overhead, these classifiers could not be fully trained by Monte Carlo cross validation, consequently, only estimates of their performance could be obtained. Five high performing alternative classifiers were identified and then fully trained. The classifier and the five high performing alternatives appear in Table 20. The frequency of each protein appears in the tally column, in particular, the first 11 proteins appear in 4 out of the 6 clas-

sifiers. These 11 proteins have significantly higher cooperative scores than the remaining proteins. By this analysis it appears that there is a core group of proteins that form the blood signature of a malignant nodule.

TABLE 20

The classifier and the high performing alternatives; coefficients for proteins on the respective panels are shown.								
Protein	Classifier	Panel 110424	Panel 130972	Panel 126748	Panel 109919	Panel 60767	Protein Tally	Cooperative Score
Constant	36.16	27.72	27.69	23.47	21.32	23.17	—	—
ALDOA	-0.8	-0.67	-0.87	-0.83	-0.64	-0.68	6	1.3
COIA1	-1.56	-1.04	-1.68	-1.37	-0.94	-1.2	6	3.7
TSP1	0.53	0.53	0.39	0.42	0.47	0.41	6	1.8
FRIL	0.39	0.45	0.39	0.41	0.41	0.41	6	2.8
LRP1	-1.59	-0.84	-1.32	1.15	-0.84	-0.87	6	4.0
GRP78	1.41	1.14	1.31	-0.34	0.78	0.6	6	1.4
ISLR	1.4	1.03	1.08	0.75	0.74		5	1.4
IBP3		-0.23	-0.21	-0.38	-0.33	-0.54	5	3.4
TETN	-1.79	-1.23	-1.99	-1.26			4	2.5
PRDX1	-0.34	-0.38			-0.36	-0.4	4	1.5
LG3BP	-0.58		-0.61		-0.38	-0.48	4	4.3
CD14			0.99	1.08		1.4	3	4.0
BGH3	1.73		1.67	-0.83			3	1.8
KIT					-0.31	-0.56	3	1.4
GGH					0.44	0.52	3	1.3
AIFM1			-0.51				1	1.4
FIBA	0.31						1	1.1
GSLG1	-0.7						1	1.2
ENPL							0	1.1
EF1A1							0	1.2
TENX							0	1.1

This result suggests that there is a core group of proteins that define a high performance classifier, but alternative panels exist. However, changes in panel membership affect the tradeoff between NPV and ROR.

### Example 7

#### Materials and Methods

##### Assay Development Candidates Sourced from Tissue

Patient samples obtained from fresh lung tumor resections were collected from Centre Hospitalier de l'Université de Montreal and McGill University Health Centre under IRB approval and with informed patient consent. Samples were obtained from the tumor as well as from distal normal tissue in the same lung lobe. Plasma membranes of each pair of samples were then isolated from the epithelial cells of 30 patients (19 adenocarcinoma, 6 squamous, 5 large cell carcinoma) and endothelial cells of 38 patients (13 adenocarcinoma, 18 squamous, 7 large cell carcinoma) using immune-affinity protocols. Golgi apparatus were isolated from each pair of samples from 33 patients (18 adenocarcinoma, 14 squamous, 1 adenosquamous) using isopycnic centrifugation followed by ammonium carbonate extraction. Plasma membrane isolations and Golgi isolations were then analyzed by tandem mass spectrometry to identify proteins overexpressed in lung cancer tissue over normal tissue, for both plasma membranes and Golgi.

##### Assay Development Candidates Sourced from Literature

Candidate lung cancer biomarkers were identified from two public and one commercial database: Entrez (www.ncbi.nlm.nih.gov/books/NBK3836), UniProt (www.uni-

prot.org) and NextBio (www.nextbio.com). Terminologies were predefined for the database queries which were automated using PERL scripts. The mining was carried out on May 6, 2010 (UniProt), May 17, 2010 (Entrez) and Jul. 8,

2010 (NextBio), respectively. Biomarkers were then assembled and mapped to UniProt identifiers.

##### Evidence of Presence in Blood

The tissue-sourced and literature-source biomarker candidates were required to have evidence of presence in blood. For evidence by mass spectrometry detection, three datasets were used. HUPO9504 contains 9504 human proteins identified by tandem mass spectrometry [13]. HUPO889, a higher confidence subset of HUPO9504, contains 889 human proteins [18]. The PeptideAtlas (November 2009 build) was also used. A biomarker candidate was marked as previously detected if it contained at least one HUPO889, or at least two HUPO9504 peptides, or at least two PeptideAtlas peptides.

In addition to direct evidence of detection in blood by mass spectrometry, annotation as secreted proteins or as single-pass membrane proteins [19] were also accepted as evidence of presence in blood. Furthermore, proteins in UniProt or designation as plasma proteins three programs for predicting whether or not a protein is secreted into the blood were used. These programs were TMHMM [20], SignalP [21] and SecretomeP [22]. A protein was predicted as secreted if TMHMM predicted the protein had one transmembrane domain and SignalP predicted the transmembrane domain was cleaved; or TMHMM predicted the protein had no transmembrane domain and either SignalP or SecretomeP predicted the protein was secreted.

##### SRM Assay Development

SRM assays for 388 targeted proteins were developed based on synthetic peptides, using a protocol similar to those described in the literature [15, 23, 24]. Up to five SRM suitable peptides per protein were identified from public sources such as the PeptideAtlas, Human Plasma Proteome Database

or by proteotypic prediction tools [25] and synthesized. SRM triggered MS/MS spectra were collected on an ABSciex 5500 QTrap for both doubly and triply charged precursor ions. The obtained MS/MS spectra were assigned to individual peptides using MAS-COT (cutoff score  $\geq 15$ ) [26]. Up to four transitions per precursor ion were selected for optimization. The resulting corresponding optimal retention time, declustering potential and collision energy were assembled for all transitions. Optimal transitions were measured on a mixture of all synthetic peptides, a pooled sample of benign patients and a pooled sample of cancer patients. Transitions were analyzed in batches, each containing up to 1750 transitions. Both biological samples were immuno-depleted and digested by trypsin and were analyzed on an ABSciex 5500 QTrap coupled with a reversed-phase (RP) high-performance liquid chromatography (HPLC) system. The obtained SRM data were manually reviewed to select the two best peptides per protein and the two best transitions per peptide. Transitions having interference with other transitions were not selected. Ratios between intensities of the two best transitions of peptides in the synthetic peptide mixture were also used to assess the specificity of the transitions in the biological samples. The intensity ratio was considered as an important metric defining the SRM assays.

#### Processing of Plasma Samples

Plasma samples were sequentially depleted of high- and medium-abundance proteins using immuno-depletion columns packed with the IgY14-Supermix resin from Sigma. The depleted plasma samples were then denatured, digested by trypsin and desalted. Peptide samples were separated using a capillary reversed-phase LC column (Thermo BioBasic 18 KAPPA; column dimensions: 320  $\mu\text{m} \times 150$  mm; particle size: 5  $\mu\text{m}$ ; pore size: 300  $\text{\AA}$ ) and a nano-HPLC system (nanoACQUITY, Waters Inc.). The mobile phases were (A) 0.2% formic acid in water and (B) 0.2% formic acid in acetonitrile. The samples were injected (8  $\mu\text{l}$ ) and separated using a linear gradient (98% A to 70% A over 19 minutes, 5  $\mu\text{l}/\text{minute}$ ). Peptides were eluted directly into the electrospray source of the mass spectrometer (5500 QTrap LC/MS/MS, AB Sciex) operating in scheduled SRM positive-ion mode (Q1 resolution: unit; Q3 resolution: unit; detection window: 180 seconds; cycle time: 1.5 seconds). Transition intensities were then integrated by software MultiQuant (AB Sciex). An intensity threshold of 10,000 was used to filter out noisy data and undetected transitions.

#### Plasma Samples Used for Discovery and Validation Studies

Aliquots of plasma samples were provided by the Institut Universitaire de Cardiologie et de Pneumologie de Quebec (IUCPQ, Hospital Laval), New York University, the University of Pennsylvania, and Vanderbilt University (see Table 17). Subjects were enrolled in clinical studies previously approved by their Ethics Review Board (ERB) or Institutional Review Boards (IRB), respectively. In addition, plasma samples were provided by study investigators after review and approval of the sponsor's study protocol by the respective institution's IRB as required. Sample eligibility for the proteomic analysis was based on the satisfaction of the study inclusion and exclusion criteria, including the subject's demographic information, the subject's corresponding lung nodule radiographic characterization by chest computed tomography (CT), and the histopathology of the lung nodule obtained at the time of diagnostic surgical resection. Cancer samples had a histopathologic diagnosis of either non-small cell lung cancer (NSCLC), including adenocarcinoma, squamous cell, large cell, or bronchoalveolar cell carcinoma and a radiographic nodule of 30 mm or smaller. Benign samples,

including granulomas, hamartomas and scar tissue, were also required to have a radiographic nodule of 30 mm or smaller and either histopathologic confirmation of being non-malignant or radiological confirmation in alignment with clinical guidelines. To ensure the accuracy of the clinical data, independent monitoring and verification of the clinical data associated with both the subject and lung nodule were performed in accordance with the guidance established by the Health Insurance Portability and Accountability Act (HIPAA) of 1996 to ensure subject privacy.

#### Study Design

The objective of the study design was to eliminate clinical and technical bias. Clinically, cancer and benign samples were paired so that they were from the same site, same gender, nodule sizes within 10 mm, age within 10 years, and smoking history within 20 pack years. Up to 15 pairs of matched cancer and benign samples per batch were assigned iteratively to processing batches until no statistical bias was demonstrable based on age, gender or nodule size.

Paired samples within each processing batch were further randomly and repeatedly assigned to positions within the processing batch, until the absolute values of the corresponding Pearson correlation coefficients between position and gender, nodule size, and age were less than 0.1. Afterwards, each pair of cancer and benign samples was randomized to their relative positions. To provide a control for sample batching, three 200  $\mu\text{l}$  aliquots of a pooled human plasma standard (HPS) (Bioreclamation, Hicksville, N.Y.) were positioned at the beginning, middle and end of each processing batch, respectively. Samples within a batch were analyzed together.

#### Logistic Regression Model

The logistic regression classification method [27] was used to combine a panel of transitions into a classifier and to calculate a classification probability score between 0 and 1 for each sample. The probability score ( $P_s$ ) of a sample was determined as  $P_s = 1/[1 + \exp(-\alpha - \sum_{i=1}^N \beta_i * \hat{I}_{i,s})]$ , where  $\hat{I}_{i,s}$  was the logarithmically transformed (base 2), normalized intensity of transition  $i$  in sample  $s$ ,  $\beta_i$  was the corresponding logistic regression coefficient,  $\alpha$  was a classifier-specific constant, and  $N$  was the total number of transitions in the classifier. A sample was classified as benign if  $P_s$  was less than a decision threshold. The decision threshold can be increased or decreased depending on the desired NPV. To define the classifier, the panel of transitions (i.e. proteins), their coefficients, the normalization transitions, classifier coefficient  $\alpha$  and the decision threshold must be learned (i.e. trained) from the discovery study and then confirmed using the validation study.

#### Discovery of the Rule-Out Classifier

A summary of the 143 samples used for classifier discovery appears in Table 17 and processed as described above.

Protein transitions were normalized as described above. Transitions that were not detected in at least 50% of the cancer samples or 50% of the benign samples were eliminated leaving 117 transitions for further consideration. Missing values for these transitions were replaced by half the minimum detected value over all samples for that transition.

The next step was finding the set of most cooperative proteins. The cooperative score of a protein is the number of high performing panels it participates in divided by the number of such panels it could appear on by chance alone. Hence, a cooperative score above 1 is good, and a score below 1 is not. The cooperative score for each protein is estimated by the following procedure:

One million random panels of 10 proteins each, selected from the 117 candidates, were generated. Each panel of 10 proteins was trained using the Monte Carlo cross validation

(MCCV) method with a 20% hold-off rate and one hundred sample permutations per panel) to fit a logistic regression model and its performance assessed by partial AUC [28].

By generating such a large number of panels, we sample the space of classifiers sufficiently well to find some high performers by chance. The one hundred best random panels (see Table 2) out of the million generated were kept and for

each of the 117 proteins we determined how frequently each occurred on these top panels. Of the 117 proteins, 36 had frequency more than expected by chance, after endogenous normalizers were removed. (Table 22) The expected number of panels on which a protein would appear by chance is  $100 \cdot 10 / 117 = 8.33$ . The cooperative score for a protein is the number of panels it appears on divided by 8.33.

TABLE 21

Category	Protein (UniProt)	Official Gene Name	Cooperative Score	Partial AUC	Coefficient CV	Transition
Classifier	TSP1_HUMAN	THBS1	1.8	0.25	0.24	GFLLLASLR_495.31_559.40
Classifier	COIA1_HUMAN	COL18A1	3.7	0.16	0.25	AVGLAG-TFR_446.26_721.40
Classifier	ISLR_HUMAN	ISLR	1.4	0.32	0.25	ALPGTPVASS-QPR_640.85_841.50
Classifier	TETN_HUMAN	CLEC3B	2.5	0.26	0.26	LDTLAQE-VALLK_657.39_330.20
Classifier	FRIL_HUMAN	FTL	2.8	0.31	0.26	LGG-PEAGLGEYLFER_804.40_913.40
Classifier	GRP78_HUMAN	HSPA5	1.4	0.27	0.27	TWNDPSPVQQDIK_715.85_260.20
Classifier	ALDOA_HUMAN	ALDOA	1.3	0.26	0.28	ALQASALK_401.25_617.40
Classifier	BGH3_HUMAN	TGFBI	1.8	0.21	0.28	LTLAPLNSVFK_658.40_804.50
Classifier	LG3BP_HUMAN	LGALS3BP	4.3	0.29	0.29	VE-IFYR_413.73_598.30
Classifier	LRP1_HUMAN	LRP1	4.0	0.13	0.32	TVLWPNGLSLDIPA GR_855.00_400.20
Classifier	FI-BA_HUMAN	FGA	1.1	0.31	0.35	NSLFEYQK_514.76_714.30
Classifier	PRDX1_HUMAN	PRDX1	1.5	0.32	0.37	QITVNDLPVGR_606.30_428.30
Classifier	GSLG1_HUMAN	GLG1	1.2	0.34	0.45	IIIQESALDYR_660.86_338.20
Robust	KIT_HUMAN	KIT	1.4	0.33	0.46	
Robust	CD14_HUMAN	CD14	4.0	0.33	0.48	
Robust	EF1A1_HUMAN	EEF1A1	1.2	0.32	0.56	
Robust	TENX_HUMAN	TNXB	1.1	0.30	0.56	
Robust	AIFM1_HUMAN	AIFM1	1.4	0.32	0.70	
Robust	GGH_HUMAN	GGH	1.3	0.32	0.81	
Robust	IBP3_HUMAN	IGFBP3	3.4	0.32	1.82	
Robust	ENPL_HUMAN	HSP90B1	1.1	0.29	5.90	
Non-Robust	ERO1A_HUMAN	ERO1L	6.2			
Non-Robust	6PGD_HUMAN	PGD	4.3			
Non-Robust	ICAM1_HUMAN	ICAM1	3.9			
Non-Robust	PTPA_HUMAN	PPP2R4	2.1			
Non-Robust	NCF4_HUMAN	NCF4	2.0			
Non-Robust	SEM3G_HUMAN	SEMA3G	1.9			
Non-Robust	1433T_HUMAN	YWHAQ	1.5			
Non-Robust	RAP2B_HUMAN	RAP2B	1.5			
Non-Robust	MMP9_HUMAN	MMP9	1.4			
Non-Robust	FOLH1_HUMAN	FOLH1	1.3			
Non-Robust	GSTP1_HUMAN	GSTP1	1.3			
Non-Robust	EF2_HUMAN	EEF2	1.3			
Non-Robust	RAN_HUMAN	RAN	1.2			
Non-Robust	SODM_HUMAN	SOD2	1.2			

TABLE 21-continued

Non- Robust	DSG2_HUMAN	DSG2	1.1			
Category	SEQ ID NO	Coefficient (Discovery) alpha = 36.16	Coefficient (Final) alpha = 26.25	Tissue Candidate	Predicted Concen- tration (ng/ml)	
Classifier	22	0.53	0.44		510	
Classifier	11	-1.56	-0.91		35	
Classifier	14	1.40	0.83		—	
Classifier	20	-1.79	-1.02		58000	
Classifier	24	0.39	0.17	Secreted, Epi, Endo	12	
Classifier	23	1.41	0.55	Secreted, Epi, Endo	100	
Classifier	7	-0.80	-0.26	Secreted, Epi	250	
Classifier	8	1.73	0.54	Epi	140	
Classifier	25	-0.58	-0.21	Secreted	440	
Classifier	15	-1.59	-0.83	Epi	20	
Classifier	26	0.31	0.13		130000	
Classifier	16	-0.34	-0.26	Epi	60	
Classifier	27	-0.70	-0.44	Epi, Endo	—	
Robust					8.2	
Robust				Epi	420	
Robust				Secreted, Epi	61	
Robust				Endo	70	
Robust				Epi, Endo	1.4	
Robust					250	
Robust					5700	
Robust				Secreted, Epi, Endo	88	
Non- Robust				Secreted, Epi, Endo	—	
Non- Robust				Epi, Endo	29	
Non- Robust					71	
Non- Robust				Endo	3.3	
Non- Robust				Endo	—	
Non- Robust					—	
Non- Robust				Epi	180	
Non- Robust				Epi	—	
Non- Robust					28	
Non- Robust					—	
Non- Robust				Endo	32	
Non- Robust				Secreted, Epi	30	
Non- Robust				Secreted, Epi	4.6	
Non- Robust				Secreted	7.1	
Non- Robust				Endo	2.7	

The 36 most cooperative proteins are listed in Table 22.

TABLE 22

Category	Protein (UniProt)	Official Gene Name	Coopera- tive Score	Partial AUC	Coeffi- cient CV	Transition
Classifier	TSP1_HUMAN	THBS1	1.8	0.25	0.24	GFLLLASLR_495.31_559.40
Classifier	COL1A1_HUMAN	COL1&A1	3.7	0.16	0.25	AVGLAGTFR_446.26_721.40
Classifier	ISLR_HUMAN	ISLR	1.4	0.32	0.25	ALPGTPVASS- QPR_640.85_841.50
Classifier	TETN_HUMAN	CLEC3B	2.5	0.26	0.26	LDTLAQE- VALLK_657.39_330.20

TABLE 22-continued

Classifier	FRIL_HUMAN	FTL	2.8	0.31	0.26	LGG- PEAGLGEYLFER_804.40_913. 40
Classifier	GRP78_HUMAN	HSPA5	1.4	0.27	0.27	TWNDPSVQQDIK_715.85_26 0.20
Classifier	ALDOA_HUMAN	ALDOA	1.3	0.26	0.28	ALQASALK_401.25_617.40
Classifier	BGH3_HUMAN	TGFBI	1.8	0.21	0.28	LTLAPLNSVFK_658.40_804. 50
Classifier	LG3BP_HUMAN	LGALS3BP	4.3	0.29	0.29	VEIFYR_413.73_598.30
Classifier	LRP1_HUMAN	LRP1	4.0	0.13	0.32	TVLWPNGLSLDIPAGR_855.0 0_400.20
Classifier	FI- BA_HUMAN	FGA	1.1	0.31	0.35	NSLFEYQK_514.76_714.30
Classifier	PRDX1_HUMAN	PRDX1	1.5	0.32	0.37	QITVNDLPVGR_606.30_428. 30
Classifier	GSLG1_HUMAN	GLG1	1.2	0.34	0.45	IIIQESALDYR_660.86_338.20
Robust	KIT_HUMAN	KIT	1.4	0.33	0.46	
Robust	CD14_HUMAN	CD14	4.0	0.33	0.48	
Robust	EF1A1_HUMAN	EEF1A1	1.2	0.32	0.56	
Robust	TENX_HUMAN	TNXB	1.1	0.30	0.56	
Robust	AIFM1_HUMAN	AIFM1	1.4	0.32	0.70	
Robust	GGH_HUMAN	GGH	1.3	0.32	0.81	
Robust	IBP3_HUMAN	IGFBP3	3.4	0.32	1.82	
Robust	ENPL_HUMAN	HSP90B1	1.1	0.29	5.90	
Non-Robust	ERO1A_HUMAN	ERO1L	6.2			
Non-Robust	6PGD_HUMAN	PGD	4.3			
Non-Robust	ICAM1_HUMAN	ICAM1	3.9			
Non-Robust	PTPA_HUMAN	PPP2R4	2.1			
Non-Robust	NCF4_HUMAN	NCF4	2.0			
Non-Robust	SEM3G_HUMAN	SE- MA3G	1.9			
Non-Robust	1433T_HUMAN	YWHAQ	1.5			
Non-Robust	RAP2B_HUMAN	RAP2B	1.5			
Non-Robust	MMP9_HUMAN	MMP9	1.4			
Non-Robust	FOLH1_HUMAN	FOLH1	1.3			
Non-Robust	GSTP1_HUMAN	GSTP1	1.3			
Non-Robust	EF2_HUMAN	EEF2	1.3			
Non-Robust	RAN_HUMAN	RAN	1.2			
Non-Robust	SODM_HUMAN	SOD2	1.2			
Non-Robust	DSG2_HUMAN	DSG2	1.1			

Category	SEQ ID NO	Coefficient (Discovery) alpha = 36.16	Coefficient (Final) alpha = 26.25	Tissue Candidate	Predicted Concentration (ng/ml)
Classifier	22	0.53	0.44		510
Classifier	11	-1.56	-0.91		35
Classifier	14	1.40	0.83		—
Classifier	20	-1.79	-1.02		58000
Classifier	24	0.39	0.17	Secreted, Epi, Endo	12
Classifier	23	1.41	0.55	Secreted, Epi, Endo	100
Classifier	7	-0.80	-0.26	Secreted, Epi	250
Classifier	8	1.73	0.54	Epi	140
Classifier	25	-0.58	-0.21	Secreted	440
Classifier	15	-1.59	-0.83	Epi	20
Classifier	26	0.31	0.13		130000
Classifier	16	-0.34	-0.26	Epi	60
Classifier	27	-0.70	-0.44	Epi, Endo	—
Robust					8.2
Robust				Epi	420
Robust				Secreted, Epi	61
Robust				Endo	70
Robust				Epi, Endo	1.4
Robust					250
Robust					5700
Robust				Secreted, Epi, Endo	88
Non-Robust				Secreted, Epi, Endo	—
Non-Robust				Epi, Endo	29
Non-Robust					71
Non-Robust				Endo	3.3
Non-Robust				Endo	—
Non-Robust					—
Non-Robust				Epi	180
Non-Robust				Epi	—

TABLE 22-continued

Non-Robust		28
Non-Robust		—
Non-Robust	Endo	32
Non-Robust	Secreted,	30
	Epi	
Non-Robust	Secreted,	4.6
	Epi	
Non-Robust	Secreted	7.1
Non-Robust	Endo	2.7

The set of 36 cooperative proteins was further reduced to a set of 21 proteins by manually reviewing raw SRM data and eliminating proteins that did not have robust SRM transitions due to low signal to noise or interference.

Proteins were iteratively eliminated from the set of 21 proteins until a classifier with the optimal partial AUC was obtained. The criteria for elimination was coefficient stability. In a logistic regression model each protein has a coefficient. In the process of training the model the coefficient for each protein is determined. When this is performed using cross validation (MCCV), hundreds of coefficient estimates for each protein are derived. The variability of these coefficients is an estimate of the stability of the protein. At each step the proteins were trained using MCCV (hold out rate 20%, ten thousand sample permutations per panel) to a logistic regression model and their stability measured. The least stable protein was eliminated. This process continued until a 13 protein classifier with optimal partial AUC was reached.

Finally, the 13 protein classifier was trained to a logistic regression model by MCCV (hold out rate 20%, twenty thousand sample permutations). The thirteen proteins for the rule-out classifier are listed in Table 18 along with their highest intensity transition and model coefficient.

#### Selection of a Decision Threshold

Assuming the cancer prevalence of lung nodules is *prev*, the performance of a classifier (NPV and ROR) on the patient population with lung nodules was calculated from sensitivity (*sens*) and specificity (*spec*) as follows:

$$NPV = \frac{(1 - prev) * spec}{prev * (1 - sens) + (1 - prev) * spec}, \quad (1)$$

$$PPV = \frac{prev * sens}{prev * sens + (1 - prev) * (1 - spec)}, \quad (2)$$

$$ROR = prev * (1 - sens) + (1 - prev) * spec. \quad (3)$$

The threshold separating calls for cancer or benign samples was then selected as the probability score with  $NPV \geq 90\%$  and  $ROR \geq 20\%$ . As we expect the classifier's performance measured on the discovery set to be an overestimate, the threshold is selected to be a range, as performance will usually degrade on an independent validation set.

#### Validation of the Rule-Out Classifier

52 cancer and 52 benign samples (see Table 17) were used to validate the performance of the 13 protein classifier. Half of the samples were placed in pre-determined processing batches analyzed immediately after the discovery samples and the other half of samples were analyzed at a later date. This introduced variability one would expect in practice. More specifically, the three HPS samples run in each processing batch were utilized as external calibrators. Details on HPS calibration are described below.

#### Calibration by HPS Samples

For label-free MS approach, variation on signal intensity between different experiments is expected. To reduce this variation, we utilized HPS samples as an external standard and calibrated the intensity between the discovery and validation studies. Assume that  $\check{I}_{i,s}$  is the logarithmically transformed (base 2), normalized intensity of transition *i* in sample *s*,  $\check{I}_{i,dis}$  and  $\check{I}_{i,val}$  are the corresponding median values of HPS samples in the discovery and the validation studies, respectively. Then the HPS corrected intensity is

$$\check{I}_{i,s} = \check{I}_{i,s} - \check{I}_{i,val} + \check{I}_{i,dis}$$

Consequently, assume that the probability for cancer of a clinical sample in the validation study is predicted as *prob* by the classifier. Then the HPS corrected probability of cancer of the clinical sample is calculated as follows:

$$probability_{corrected} = \frac{1}{1 + e^{-S_{corrected}}}$$

where

$$S_{corrected} = S - S_{HPS,val} + S_{HPS,dis}$$

and

$$S = \ln \frac{prob}{1 - prob}.$$

Here  $S_{HPS,dis}$  and  $S_{HPS,val}$  were the median value of *S* of all HPS samples in the discovery and validation studies, respectively.

#### Statistical Analysis

All statistical analyses were performed with Stata, R and/or MatLab.

#### Depletion Column Drift

We observed an increase of signal intensity as more and more samples were depleted by the same column. We used transition intensity in HPS samples to quantify this technical variability. Assuming  $I_{i,s}$  was the intensity of transition *i* in a HPS sample *s*, the drift of the sample was defined as

$$drift_s = \text{median} \left( \frac{I_{i,s} - \hat{I}_i}{\hat{I}_i} \right),$$

where  $\hat{I}_i$  was the mean value of  $I_{i,s}$  among all HPS samples that were depleted by the same column and the median was taken over all detected transitions in the sample. Then the drift of the column was defined as

$$drift_{col} = \text{median}(drift_s > 0) - \text{median}(drift_s < 0).$$

Here the median was taken over all HPS samples depleted by the column. If no sample drift was greater or less than zero,

the corresponding median was taken as 0. The median column drift was the median of drifts of all depletion columns used in the study.

#### Identification of Endogenous Normalizing Proteins

The following criteria were used to identify a transition as a normalizer:

Possessed the highest median intensity of all transitions from the same protein.

Detected in all samples.

Ranked high in reducing median technical CV (median CV of transition intensities that were measured on HPS samples) as a normalizer.

Ranked high in reducing median column drift that was observed in sample depletion.

Possessed low median technical CV and low median biological CV (median CV of transition intensities that were measured on clinical samples).

Six transitions were selected and appear in Table 23.

TABLE 23

Panel of endogenous normalizers.				
Normalizer	Transition	SEQ ID NO	Median Technical CV (%)	Median Column Drift (%)
PEDF_HUMAN	LQSLFDSPDFSK_692.34_593.30	28	25.8	6.8
MASP1_HUMAN	TGVTSPDFPNPYPK_816.92_258.10	6	26.5	18.3
GELS_HUMAN	TASDFITK_441.73_710.40	5	27.1	16.8
LUM_HUMAN	SLEDLQLTHNK_433.23_499.30	29	27.1	16.1
C163A_HUMAN	INPASLDK_429.24_630.30	30	26.6	14.6
PTPRJ_HUMAN	VITEPIPVSDLR_669.89_896.50	31	27.2	18.2
	Normalization by Panel of Transitions		25.1	9.0
	Without Normalization		32.3	23.8

#### Data Normalization

A panel of six normalization transitions (see Table 23) were used to normalize raw SRM data for two purposes: (A) to reduce sample-to-sample intensity variations within same study and (B) to reduce intensity variations between different studies. For the first purpose, a scaling factor was calculated for each sample so that the intensities of the six normalization transitions of the sample were aligned with the corresponding median intensities of all HGS samples. Assuming that  $N_{i,s}$  is the intensity of a normalization transition  $i$  in sample  $s$  and  $\hat{N}_i$  the corresponding median intensity of all HGS samples, then the scaling factor for sample  $s$  is given by  $\hat{S}_s/S_s$ , where

$$S_s = \text{median} \left( \frac{N_{1,s}}{\hat{N}_1}, \frac{N_{2,s}}{\hat{N}_2}, \dots, \frac{N_{6,s}}{\hat{N}_6} \right)$$

is the median of the intensity ratios and  $\hat{S}$  is the median of  $S_s$  over all samples in the study. For the second purpose, a scaling factor was calculated between the discovery and the validation studies so that the median intensities of the six normalization transitions of all HGS samples in the validation study were comparable with the corresponding values in the discovery study. Assuming that the median intensities of all HGS samples in the two studies are  $\hat{N}_{i,dis}$  and  $\hat{N}_{i,val}$ , respectively, the scaling factor for the validation study is given by

$$R = \text{median} \left( \frac{\hat{N}_{1,dis}}{\hat{N}_{1,val}}, \frac{\hat{N}_{2,dis}}{\hat{N}_{2,val}}, \dots, \frac{\hat{N}_{6,dis}}{\hat{N}_{6,val}} \right)$$

Finally, for each transition of each sample, its normalized intensity was calculated as

$$\tilde{I}_{i,s} = I_{i,s} * R * \hat{S}_s / S_s$$

where  $I_{i,s}$  was the raw intensity.

#### Isolation of Membrane Proteins from Tissues

Endothelial plasma membrane proteins were isolated from normal and tumor lung tissue samples that were obtained from fresh lung resections. Briefly, tissues were washed in buffer and homogenates were prepared by disrupting the tissues with a Polytron. Homogenates were filtered through a 180- $\mu\text{m}$  mesh and filtrates were centrifuged at 900 $\times$ g for 10 min, at 4° C. Supernatants were centrifuged on top of a 50% (w:v) sucrose cushion at 218,000 $\times$ g for 60 min at 4° C. to pellet the membranes. Pellets were resuspended and treated with micrococcal nuclease. Membranes from endothelial cells were incubated with a combination of anti-thrombomodulin, anti-ACE, anti-CD34 and anti-CD144 antibodies,

and then centrifuged on top of a 50% (w:v) sucrose cushion at 280,000 $\times$ g for 60 min at 4° C. After pellets were resuspended, endothelial cell plasma membranes were isolated using MACS microbeads, treated with potassium iodide to remove cytoplasmic peripheral proteins.

Epithelial plasma membrane proteins from normal and tumor lung tissue samples were isolated from fresh lung resections. Tissues were washed and homogenates as described above for endothelial plasma membrane proteins preparation. Membranes from epithelial cells were labeled with a combination of anti-ESA, anti-CEA, anti-CD66c and anti-EMA antibodies, and then centrifuged on top of a 50% (w:v) sucrose cushion at 218,000 $\times$ g for 60 min at 4° C. Epithelial cell plasma membranes were isolated using MACS microbeads and the eluate was centrifuged at 337,000 $\times$ g for 30 minutes at 4° C. over a 33% (w:v) sucrose cushion. After removing the supernatant and sucrose cushion, the pellet was resuspended in Laemmli/Urea/DTT.

#### Isolation of Secreted Proteins from Tissues

Secreted proteins were isolated from normal and tumor lung tissue samples that were isolated from fresh lung resections. Tissues were washed and homogenized using a Polytron homogenization. The density of the homogenates was adjusted to 1.4 M with concentrated sucrose prior to isolating the secretory vesicles by isopycnic centrifugation at 100,000 $\times$ g for 2 hr at 4° C. on a 0.8 and 1.2 M discontinuous sucrose gradient. Vesicles concentrating at the 0.8/1.2 M interface were collected and further incubated for 25 minutes with 0.5 M KCl (final concentration) to remove loosely bound peripheral proteins. Vesicles were recuperated by ultracentrifugation at 150,000 $\times$ g for one hour at 4° C. and then opened with 100 mM ammonium carbonate pH 11.0 for 30 minutes at

4° C. Secreted proteins were recovered in the supernatant following a 1-hour ultracentrifugation at 150,000×g at 4° C.

#### Preparation of IgY14-SuperMix Immunoaffinity Columns

Immunoaffinity columns were prepared in-house using a slurry containing a 2:1 ratio of IgY14 and SuperMix immunoaffinity resins, respectively (Sigma Aldrich). Briefly, a slurry (10 ml, 50%) of mixed immunoaffinity resins was added to a glass chromatography column (Tricorn, GE Healthcare) and the resin was allowed to settle under gravity flow, resulting in a 5 ml resin volume in the column. The column was capped and placed on an Agilent 1100 series HPLC system for further packing (20 minutes, 0.15M ammonium bicarbonate, 2 ml/min). The performance of each column used in the study was then assessed by replicate injections of aliquots of HPS sample. Column performance was assessed prior to beginning immunoaffinity separation of each batch of clinical samples.

#### IgY14-Sumermix Immunoaffinity Chromatography

Plasma samples (60 µl) were diluted (0.15M ammonium bicarbonate, 1:2 v/v, respectively) and filtered (0.2 µm Acro-Prep 96-well filter plate, Pall Life Sciences) prior to immunoaffinity separation. Dilute plasma (90 µl) was separated on the IgY14-SuperMix column connected to an Agilent 1100 series HPLC system using a three buffers (loading/washing: 0.15M ammonium bicarbonate; stripping/elution: 0.1M glycine, pH 2.5; neutralization: 0.01M Tris-HCl, 0.15M NaCl, pH 7.4) with a load-wash-elute-neutralization-re-equilibration cycle (36 minutes total time). The unbound and bound fractions were monitored using a UV absorbance (280 nm) and were baseline resolved after separation. Only the unbound fraction containing the low abundance proteins was collected for downstream processing and analysis. Unbound fractions were lyophilized prior to enzymatic digestion.

#### Enzymatic Digestion of Low Abundance Proteins

Low abundance proteins were reconstituted under mild denaturing conditions (200 µl of 1:1 0.1M ammonium bicarbonate/trifluoroethanol v/v) and allowed to incubate (30 minutes, room temperature, orbital shaker). Samples were then diluted (800 µl of 0.1M ammonium bicarbonate) and digested with trypsin (Princeton Separations; 0.4 µg trypsin per sample, 37° C., 16 hours). Digested samples were lyophilized prior to solid-phase extraction.

#### Solid-Phase Extraction

Solid phase extraction was used to reduce salt and buffer contents in the samples prior to mass spectrometry. The lyophilized samples containing tryptic peptides were reconstituted (350 µl 0.01M ammonium bicarbonate) and allowed to incubate (15 minutes, room temperature, orbital shaker). A reducing agent was then added to the samples (30 µl 0.05M TCEP) and the samples were incubated (60 minutes, room temperature). Dilute acid and a low percentage of organic solvent (375 µl 90% water/10% acetonitrile/0.2% trifluoroacetic acid) were added to optimize the solid phase extraction of peptides. The extraction plate (Empore C18, 3M Bioanalytical Technologies) was conditioned according to manufacturer protocol. Samples were loaded onto the solid phase extraction plate, washed (500 µl 95% water/5% acetonitrile/0.1% trifluoroacetic acid) and eluted (200 µl 52% water/48% acetonitrile/0.1% trifluoroacetic acid) into a collection plate. The eluate was split into two equal aliquots and each aliquot was taken to dryness in a vacuum concentrator. One aliquot was used immediately for mass spectrometry, while the other was stored (-80° C.) and used as needed. Samples were reconstituted (12 µl 90% water/10% acetonitrile/0.2% formic acid) just prior to LC-SRM MS analysis.

#### Inclusion and Exclusion Criteria

Plasma samples were eligible for the studies if they were (A) obtained in EDTA tubes, (B) obtained from subjects previously enrolled in IRB-approved studies at the participating institutions, and (C) archived, e.g. labeled, aliquotted and frozen, as stipulated by the study protocols. The samples must also satisfy the following inclusion and exclusion criteria:

##### 1) Inclusion Criteria:

2) Sample eligibility was based on clinical parameters, including the following subject, nodule and clinical staging parameters:

##### a) Subject

i) age ≥ 40

ii) any smoking status, e.g. current, former, or never

iii) co-morbid conditions, e.g. COPD

iv) prior malignancy with a minimum of 5 years in clinical remission

v) prior history of skin carcinomas—squamous or basal cell

##### b) Nodule

i) Radiology

(1) size ≥ 4 mm and ≤ 70 mm (up to Stage 2B eligible)

(2) any spiculation or ground glass opacity

ii) pathology

(1) malignant—adenocarcinoma, squamous, or large cell

(2) benign—inflammatory (e.g. granulomatous, infectious) or non-inflammatory (e.g. hamartoma)

##### c) Clinical stage

i) Primary tumor: ≤ T2 (e.g. 1A, 1B, 2A and 2B)

ii) Regional lymph nodes: N0 or N1 only

iii) Distant metastasis: M0 only

##### 3) Exclusion Criteria

a) Subject: prior malignancy within 5 years of IPN diagnosis

b) Nodule:

i) size data unavailable

ii) for cancer or benign SPNs, no pathology data available

iii) pathology—small cell lung cancer

c) Clinical stage

i) Primary tumor: ≥ T3

ii) Regional lymph nodes: ≥ N2

iii) Distant metastasis: ≥ M1

#### Power Analysis for the Discovery Study

The power analysis for the discovery study was based on the following assumptions: 1) The overall false positive rate ( $\alpha$ ) was set to 0.05. 2) Šidák correction for multiple testing was used to calculate the effective  $\alpha_{eff}$  for testing 200 proteins, i.e.,

$$\alpha_{eff} = 1 - \sqrt[200]{1 - \alpha}.$$

3) The effective sample size was reduced by a factor of 0.864 to account for the larger sample requirement for the Mann-Whitney test than for the t-test. 4) The overall coefficient of variation was set to 0.43 based on a previous experience. 5) The power ( $1 - \beta$ ) of the study was calculated based on the formula for the two-sample, two-sided t-test, using effective  $\alpha_{eff}$  and effective sample size. The power for the discovery study was tabulated in Table 24 by the sample size per cohort and the detectable fold difference between control and disease samples.

TABLE 24

Cohort size required to detect protein fold changes with a given probability.				
Cohort Size	Detectable Protein Fold Difference			
	1.25	1.5	1.75	2
20	0.011	0.112	0.368	0.653
30	0.025	0.277	0.698	0.925
40	0.051	0.495	0.905	0.992
50	0.088	0.687	0.977	0.999
60	0.129	0.812	0.994	1
70	0.183	0.902	0.999	1
80	0.244	0.953	1	1
90	0.302	0.977	1	1
100	0.369	0.99	1	1

Power Analysis for the Validation Study

Sufficient cancer and benign samples are needed in the validation study to confirm the performance of the rule-out classifier obtained from the discovery study. We are interested in obtaining the 95% confidence intervals (CIs) on NPV and ROR for the rule-out classifier. Using the Equations in the Selection of a Decision Threshold section herein, one can derive sensitivity (sens) and specificity (spec) as functions of NPV and ROR, i.e.,

$$\text{sens} = 1 - \text{ROR} * (1 - \text{NPV}) / \text{prev},$$

$$\text{spec} = \text{ROR} * \text{NPV} / (1 - \text{prev}),$$

where prev is the cancer prevalence in the intended use population. Assume that the validation study contains  $N_C$  cancer samples and  $N_B$  benign samples. Based on binomial distribution, variances of sensitivity and specificity are given by

$$\text{var}(\text{sens}) = \text{sens} * (1 - \text{sens}) / N_C$$

$$\text{var}(\text{spec}) = \text{spec} * (1 - \text{spec}) / N_B$$

Using the Equations in the Selection of a Decision Threshold section herein, the corresponding variances of NPV and ROR can be derived under the large-sample, normal-distribution approximation as

$$\text{var}(\text{NPV}) = \text{NPV}^2 (1 - \text{NPV})^2 \left[ \frac{\text{var}(\text{sens})}{(1 - \text{sens})^2} + \frac{\text{var}(\text{spec})}{\text{spec}^2} \right],$$

$$\text{var}(\text{ROR}) = \text{prev}^2 * \text{var}(\text{sens}) + (1 - \text{prev})^2 * \text{var}(\text{spec}).$$

The two-sided 95% CIs of NPV and ROR are then given by  $\pm z_{\alpha/2} \sqrt{\text{var}(\text{NPV})}$  and  $\pm z_{\alpha/2} \sqrt{\text{var}(\text{ROR})}$ , respectively, where  $z_{\alpha/2} = 1.959964$  is the 97.5% quantile of the normal distribution. The anticipated 95% CIs for the validation study were tabulated in Table 24 by the sample size ( $N_C = N_B = N$ ) per cohort.

TABLE 24

The 95% confidence interval (CI) of NPV as a function of cohort size. The corresponding 95% CI of ROR is also listed. The prevalence was set at 28.5%. The expected NPV and ROR were set to values in the discovery study, i.e., 90% and 52%, respectively.		
Cohort Size	95% CI of NPV (± %)	95% CI of ROR (± %)
10	12.5	22.1
20	8.8	15.7

TABLE 24-continued

The 95% confidence interval (CI) of NPV as a function of cohort size. The corresponding 95% CI of ROR is also listed. The prevalence was set at 28.5%. The expected NPV and ROR were set to values in the discovery study, i.e., 90% and 52%, respectively.		
Cohort Size	95% CI of NPV (± %)	95% CI of ROR (± %)
30	7.2	12.8
40	6.2	11.1
50	5.6	9.9
60	5.1	9.0
70	4.7	8.4
80	4.4	7.8
90	4.2	7.4
100	3.9	7.0
150	3.2	5.7
200	2.8	5.0

Calculation of Q-Values of Peptide and Protein Assays

To determine the false positive assay rate the q-values of peptide SRM assays were calculated as follows. Using the distribution of Pearson correlations between transitions from different proteins as the null distribution (FIG. 7), an empirical p-value was assigned to a pair of transitions from the same peptide, detected in at least five common samples otherwise a value of 'NA' is assigned. The empirical p-value was converted to a q-value using the "qvalue" package in Bioconductor ([www.bioconductor.org/packages/release/bioc/html/qvalue.html](http://www.bioconductor.org/packages/release/bioc/html/qvalue.html)). Peptide q-values were below 0.05 for all SRM assays presented in Table 6.

The q-values of protein SRM assays were calculated in the same way except Pearson correlations of individual proteins were calculated as those between two transitions from different peptides of the protein. For proteins not having two peptides detected in five or more common samples, their q-values could not be properly evaluated and were assigned 'NA'.

Impact of Categorical Confounding Factors

TABLE 25

Impact of categorical confounding factors on classifier score.					
		Cancer	p-value	Benign	p-value
Gender	# Female	70	0.786*	68	0.387*
	Median score	0.701		0.570	
	(quartile range)	(0.642-0.788)		(0.390-0.70)	
# Male	Median score	54		55	
	(quartile range)	0.736		0.621	
	(quartile range)	(0.628-0.802)		(0.459-0.723)	
Smoking Status	# Never	8	0.435**	34	0.365**
	Median score	0.664		0.554	
	(quartile range)	(0.648-0.707)		(0.452-0.687)	
# Past	Median score	98		73	
	(quartile range)	0.703		0.586	
	(quartile range)	(0.618-0.802)		(0.428-0.716)	
# Current	Median score	17		13	
	(quartile range)	0.749		0.638	
	(quartile range)	(0.657-0.789)		(0.619-0.728)	

\*p-value by Mann-Whitney test

\*\*p-value by Kruskal-Wallis test

TABLE 26

Impact of continuous confounding factors on classifier score.				
		Correlation	Coefficient of linear fit (95% CI)	p-value
Age	All	0.198	0.003 (0.001-0.005)	0.002
	Cancer	0.012	0.000 (-0.003-0.003)	0.893
	Benign	0.248	0.004 (0.001-0.007)	0.006
Nodule size	All	-0.057	-0.002 (-0.005-0.002)	0.372
	Cancer	-0.013	0.000 (-0.005-0.004)	0.889
	Benign	-0.055	-0.001 (-0.006-0.003)	0.542
Pack-year	All	0.154	0.001 (0.00-0.002)	0.019
	Cancer	0.060	0.000 (-0.001-0.001)	0.520
	Benign	0.108	0.001 (0.00-0.002)	0.254

## REFERENCES

- Albert & Russell Am Fam Physician 80:827-831 (2009)
- Gould et al. Chest 132:108S-130S (2007)
- Kitteringham et al. J Chromatogr B Analyt Technol Biomed Life Sci 877:1229-1239 (2009)
- Lange et al. Mol Syst Biol 4:222 (2008)
- Lehtio & De Petris J Proteomics 73:1851-1863 (2010)
- MacMahon et al. Radiology 237:395-400 (2005)
- Makawita Clin Chem 56:212-222 (2010)
- Ocak et al. Proc Am Thorac Soc 6:159-170 (2009)
- Ost, D. E. and M. K. Gould, *Decision making in patients with pulmonary nodules*. Am J Respir Crit Care Med, 2012. 185(4): p. 363-72.
- Cima, I., et al., *Cancer genetics-guided discovery of serum biomarker signatures for diagnosis and prognosis of prostate cancer*. Proc Natl Acad Sci USA, 2011. 108(8): p. 3342-7.
- Desiere, F., et al., *The PeptideAtlas project*. Nucleic Acids Res, 2006. 34 (Database issue): p. D655-8.
- Farrar, T., et al., *A high-confidence human plasma proteome reference set with estimated concentrations in PeptideAtlas*. Mol Cell Proteomics, 2011. 10(9): p. M110006353.

- Omenn, G. S., et al., *Overview of the HUPO Plasma Proteome Project: results from the pilot phase with 35 collaborating laboratories and multiple analytical groups, generating a core dataset of 3020 proteins and a publicly-available database*. Proteomics, 2005. 5(13): p. 3226-45.
- Kearney, P., et al., *Protein identification and Peptide expression resolver: harmonizing protein identification with protein expression data*. J Proteome Res, 2008. 7(1): p. 234-44.
- Huttenhain, R., et al., *Reproducible quantification of cancer-associated proteins in body fluids using targeted proteomics*. Sci Transl Med, 2012. 4(142): p. 142ra94.
- Henschke, C. I., et al., *CT screening for lung cancer: suspiciousness of nodules according to size on baseline scans*. Radiology, 2004. 231(1): p. 164-8.
- Henschke, C. I., et al., *Early Lung Cancer Action Project: overall design and findings from baseline screening*. Lancet, 1999. 354(9173): p. 99-105.
- States, D. J., et al., *Challenges in deriving high-confidence protein identifications from data gathered by a HUPO plasma proteome collaborative study*. Nat Biotechnol, 2006. 24(3): p. 333-8.
- Polanski, M. and N. L. Anderson, *A list of candidate cancer biomarkers for targeted proteomics*. Biomark Insights, 2007. 1: p. 1-48.
- Krogh, A., et al., *Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes*. J Mol Biol, 2001. 305(3): p. 567-80.
- Bendtsen, J. D., et al., *Improved prediction of signal peptides: SignalP 3.0*. J Mol Biol, 2004. 340(4): p. 783-95.
- Bendtsen, J. D., et al., *Feature-based prediction of non-classical and leaderless protein secretion*. Protein Eng Des Sel, 2004. 17(4): p. 349-56.
- Lange, V., et al., *Selected reaction monitoring for quantitative proteomics: a tutorial*. Mol Syst Biol, 2008. 4: p. 222.
- Picotti, P., et al., *High-throughput generation of selected reaction-monitoring assays for proteins and proteomes*. Nat Methods, 2010. 7(1): p. 43-6.
- Mallik, P., et al., *Computational prediction of proteotypic peptides for quantitative proteomics*. Nat Biotechnol, 2007. 25(1): p. 125-31.
- Perkins, D. N., et al., *Probability-based protein identification by searching sequence databases using mass spectrometry data*. Electrophoresis, 1999. 20(18): p. 3551-67.
- Hastie, T., R. Tibshirani, and J. H. Friedman, *The elements of statistical learning: data mining, inference, and prediction: with 200 full-color illustrations*. Springer series in statistics. 2001, New York: Springer. xvi, 533 p.
- McClish, D. K., *Analyzing a portion of the ROC curve*. Med Decis Making, 1989. 9(3): p. 190-5.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 31

<210> SEQ ID NO 1

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Tyr Gly Phe Ile Glu Gly His Val Val Ile Pro Arg  
1 5 10

<210> SEQ ID NO 2

<211> LENGTH: 9

-continued

---

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Tyr Glu Val Thr Val Val Ser Val Arg  
1 5

<210> SEQ ID NO 3

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Ala Ser Ser Ile Ile Asp Glu Leu Phe Gln Asp Arg  
1 5 10

<210> SEQ ID NO 4

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Phe Leu Asn Val Leu Ser Pro Arg  
1 5

<210> SEQ ID NO 5

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Thr Ala Ser Asp Phe Ile Thr Lys  
1 5

<210> SEQ ID NO 6

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Thr Gly Val Ile Thr Ser Pro Asp Phe Pro Asn Pro Tyr Pro Lys  
1 5 10 15

<210> SEQ ID NO 7

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Ala Leu Gln Ala Ser Ala Leu Lys  
1 5

<210> SEQ ID NO 8

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Leu Thr Leu Leu Ala Pro Leu Asn Ser Val Phe Lys  
1 5 10

<210> SEQ ID NO 9

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

-continued

&lt;400&gt; SEQUENCE: 9

Leu Ala Ala Leu Asn Pro Glu Ser Asn Thr Ala Gly Leu Asp Ile Phe  
 1 5 10 15

Ala Lys

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 9

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 10

Val Ala Val Val Gln Tyr Ser Asp Arg  
 1 5

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 9

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 11

Ala Val Gly Leu Ala Gly Thr Phe Arg  
 1 5

&lt;210&gt; SEQ ID NO 12

&lt;211&gt; LENGTH: 15

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 12

Val Pro Gly Thr Ser Thr Ser Ala Thr Leu Thr Gly Leu Thr Arg  
 1 5 10 15

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 15

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 13

Gly Ala Leu Gln Asn Ile Ile Pro Ala Ser Thr Gly Ala Ala Lys  
 1 5 10 15

&lt;210&gt; SEQ ID NO 14

&lt;211&gt; LENGTH: 13

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 14

Ala Leu Pro Gly Thr Pro Val Ala Ser Ser Gln Pro Arg  
 1 5 10

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 16

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 15

Thr Val Leu Trp Pro Asn Gly Leu Ser Leu Asp Ile Pro Ala Gly Arg  
 1 5 10 15

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 11

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

-continued

---

&lt;400&gt; SEQUENCE: 16

Gln Ile Thr Val Asn Asp Leu Pro Val Gly Arg  
1                   5                   10

&lt;210&gt; SEQ ID NO 17

&lt;211&gt; LENGTH: 14

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 17

Ser Thr Gly Gly Ala Pro Thr Phe Asn Val Thr Val Thr Lys  
1                   5                   10

&lt;210&gt; SEQ ID NO 18

&lt;211&gt; LENGTH: 7

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 18

Ser Val Asp Ile Trp Leu Arg  
1                   5

&lt;210&gt; SEQ ID NO 19

&lt;211&gt; LENGTH: 12

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 19

Ile Ser Val Tyr Tyr Asn Glu Ala Ser Ser His Lys  
1                   5                   10

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 12

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 20

Leu Asp Thr Leu Ala Gln Glu Val Ala Leu Leu Lys  
1                   5                   10

&lt;210&gt; SEQ ID NO 21

&lt;211&gt; LENGTH: 7

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 21

Val Val Phe Glu Gln Thr Lys  
1                   5

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 9

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 22

Gly Phe Leu Leu Leu Ala Ser Leu Arg  
1                   5

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 12

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 23

-continued

Thr Trp Asn Asp Pro Ser Val Gln Gln Asp Ile Lys  
 1 5 10

<210> SEQ ID NO 24  
 <211> LENGTH: 15  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

Leu Gly Gly Pro Glu Ala Gly Leu Gly Glu Tyr Leu Phe Glu Arg  
 1 5 10 15

<210> SEQ ID NO 25  
 <211> LENGTH: 6  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

Val Glu Ile Phe Tyr Arg  
 1 5

<210> SEQ ID NO 26  
 <211> LENGTH: 8  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

Asn Ser Leu Phe Glu Tyr Gln Lys  
 1 5

<210> SEQ ID NO 27  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

Ile Ile Ile Gln Glu Ser Ala Leu Asp Tyr Arg  
 1 5 10

<210> SEQ ID NO 28  
 <211> LENGTH: 12  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

Leu Gln Ser Leu Phe Asp Ser Pro Asp Phe Ser Lys  
 1 5 10

<210> SEQ ID NO 29  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

Ser Leu Glu Asp Leu Gln Leu Thr His Asn Lys  
 1 5 10

<210> SEQ ID NO 30  
 <211> LENGTH: 8  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

Ile Asn Pro Ala Ser Leu Asp Lys

-continued

---

```

1           5

<210> SEQ ID NO 31
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

Val Ile Thr Glu Pro Ile Pro Val Ser Asp Leu Arg
1           5           10

```

---

What is claimed is:

1. A method of determining the likelihood that a pulmonary nodule in a subject is not lung cancer, comprising:

(a) contacting a blood sample obtained from the subject with a proteolytic enzyme to produce peptide fragments from a panel of proteins present in the blood sample, wherein the panel comprises ALDOA, FRIL, LG3BP, TSP1, COIA1, PEDE, MASP1, GELS, LUM, C163A and PTPRJ;

(b) combining the produced peptide fragments from the panel from step (a) with labeled, synthetic peptide fragments which correspond to the produced peptide fragments from the panel;

(c) performing selected reaction monitoring mass spectrometry to measure the abundance of the peptide fragments from step (b);

(d) calculating a probability of lung cancer score based on the peptide fragment measurements of step (c); and

(e) ruling out lung cancer for the subject if the score in step (d) is lower than a pre-determined score.

2. The method of claim 1, wherein said panel further comprises at least one protein selected from the group consisting of IBP3, LRP1, ISLR, GRP78, TETN, PRDX1, CD14, BGH3, FIBA and GSLG1.

3. The method of claim 1, wherein when lung cancer is ruled out the subject does not receive a treatment protocol.

4. The method of claim 3, wherein said treatment protocol is a pulmonary function test (PFT), pulmonary imaging, a biopsy, a surgery, a chemotherapy, a radiotherapy, or any combination thereof.

5. The method of claim 4, where said imaging is an x-ray, a chest computed tomography (CT) scan, or a positron emission tomography (PET) scan.

15 6. The method of claim 1, wherein said pulmonary nodule has a diameter of less than or equal to 3 cm.

7. The method of claim 1, wherein said pulmonary nodule has a diameter of about 0.8 cm to 2.0 cm.

20 8. The method of claim 1, wherein said score is calculated from a logistic regression model applied to the peptide fragment measurements.

9. The method of claim 1, wherein said score is determined as  $P_s = 1/[1 + \exp(-\alpha - \sum_{i=1}^N \beta_i * \check{I}_{i,s})]$ , where  $\check{I}_{i,s}$  is logarithmically transformed and normalized intensity of transition  $i$  in said sample ( $s$ ),  $\beta_i$  is the corresponding logistic regression coefficient,  $\alpha$  was a panel-specific constant, and  $N$  was the total number of transitions in said panel.

30 10. The method of claim 1, wherein the determining the likelihood that a pulmonary nodule is not lung cancer is determined by the sensitivity, specificity, negative predictive value or positive predictive value associated with the score.

11. The method of claim 1, wherein said score determined in step (d) has a negative predictive value (NPV) of at least about 80%.

35 12. The method of claim 1, wherein the selected reaction monitoring mass spectrometry is performed using a compound that specifically binds the peptide fragments being detected.

40 13. The method of claim 12, wherein the compound that specifically binds to the peptide fragments being measured is an antibody or an aptamer.

14. The method of claim 1, wherein the subject is at risk of developing lung cancer.

45 15. The method of claim 1, wherein the proteolytic enzyme is trypsin.

\* \* \* \* \*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 9,201,044 B2  
APPLICATION NO. : 13/724823  
DATED : December 1, 2015  
INVENTOR(S) : Paul Edward Kearney et al.

Page 1 of 1


It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In the claims

Claim 9, Column 176, line 23:

Please replace  $P_s = 1/[1 + \exp(-\alpha - \sum_{i=1}^N \beta_i * \tilde{I}_{i,s})]$  with  $P_s = 1/[1 + \exp(-\alpha - \sum_{i=1}^N \beta_i * \tilde{I}_{i,s})]$

Signed and Sealed this  
Nineteenth Day of April, 2016



Michelle K. Lee  
Director of the United States Patent and Trademark Office

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 9,201,044 B2  
APPLICATION NO. : 13/724823  
DATED : December 1, 2015  
INVENTOR(S) : Paul Edward Kearney et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In the Claims


Claim 9, Column 156, Line number 36-37:

“mined as  $P_s = 1 / [1 + \exp(-\alpha - \sum_{i=1}^N \beta_i * \tilde{I}_{i,s})]$ , where is logarithmically transformed and normalized intensity of transition in”

Should be replaced by:

-- mined as  $P_s = 1 / [1 + \exp(-\alpha - \sum_{i=1}^N \beta_i * \tilde{I}_{i,s})]$ , where  $\tilde{I}_{i,s}$  is logarithmically transformed and normalized intensity of transition  $i$  in --

Signed and Sealed this  
Twenty-sixth Day of September, 2017



Joseph Matal

*Performing the Functions and Duties of the  
Under Secretary of Commerce for Intellectual Property and  
Director of the United States Patent and Trademark Office*

专利名称(译)	用于诊断肺癌的组合物，方法和试剂盒		
公开(公告)号	<a href="#">US9201044</a>	公开(公告)日	2015-12-01
申请号	US13/724823	申请日	2012-12-21
当前申请(专利权)人(译)	集成诊断，INC.		
[标]发明人	KEARNEY PAUL EDWARD FANG KENNETH CHARLES LI XIAO JUN HAYWARD CLIVE		
发明人	KEARNEY, PAUL EDWARD FANG, KENNETH CHARLES LI, XIAO-JUN HAYWARD, CLIVE		
IPC分类号	G01N33/53 G01N27/62 G01N33/574 G06F19/18		
CPC分类号	G01N27/62 G01N33/57423 G06F19/18 G01N2800/60 G01N33/6848 G16B20/00 G16B40/00		
优先权	61/578712 2011-12-21 US 61/589920 2012-01-24 US 61/725153 2012-11-12 US 61/676859 2012-07-27 US		
其他公开文献	US20130217057A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

提供了用于鉴定在具有第一肺病症的受试者中与健康受试者或具有第二肺病症的受试者中表现出差异表达的生物标志物蛋白的方法。还提供了包含这些生物标志物蛋白的组合物和使用这些生物标志物蛋白或其组以诊断，分类和监测各种肺病症的方法。本文提供的方法和组合物可用于诊断或分类受试者患有肺癌或非癌症，并区分不同类型的癌症（例如，恶性与良性，SCLC与NSCLC）。




FIG. 1