



US 20100317034A1

(19) **United States**  
(12) **Patent Application Publication**  
Switzer et al.

(10) **Pub. No.: US 2010/0317034 A1**  
(43) **Pub. Date: Dec. 16, 2010**

(54) **PRIMATE T-LYMPHOTROPIC VIRUSES**

continuation-in-part of application No. PCT/US06/05869, filed on Feb. 21, 2006.

(75) Inventors: **William M. Switzer**, Stone Mountain, GA (US); **Walid Heneine**, Atlanta, GA (US); **Thomas M. Folks**, Helotes, TX (US); **Nathan D. Wolfe**, Los Angeles, CA (US); **Donald S. Burke**, Pittsburgh, PA (US); **Eitel Mpoudi Ngole**, Yaounde (CM)

(60) Provisional application No. 60/654,484, filed on Feb. 21, 2005.

**Publication Classification**

(51) **Int. Cl.**  
**G01N 33/53** (2006.01)  
**C07K 14/005** (2006.01)  
**C07K 16/08** (2006.01)

Correspondence Address:  
**KLARQUIST SPARKMAN, LLP**  
121 S.W. SALMON STREET, SUITE 1600  
PORTLAND, OR 97204 (US)

(52) **U.S. Cl. .... 435/7.21; 530/350; 436/501; 530/389.4**

(57) **ABSTRACT**

(73) Assignees: **The Government of the United States of America as represented by the Secretary of the Department of Health and Human Services, Centers for Disease Control and Prevention,; and Johns Hopkins University**

Disclosed are compositions and methods related to the isolation and identification of the primate T-lymphotropic viruses, HTLV-3 and HTLV-4. The diversity of HTLVs was investigated among central Africans reporting contact with NHP blood and body fluids through hunting, butchering, and keeping primate pets. Herein it is shown that this population is infected with a variety of HTLVs, including two retroviruses; HTLV-4 is the first member of a novel phylogenetic lineage that is distinct from all known HTLVs and STLVs; HTLV-3 falls within the genetic diversity of STLV-3, a group that has not previously been seen in humans. The present disclosure also relates to vectors and vaccines for use in humans against infection and disease. The disclosure further relates to a variety of bioassays and kits for the detection and diagnosis of infection with and diseases caused by HTLV-3 and HTLV-4 and related viruses.

(21) Appl. No.: **12/829,125**

(22) Filed: **Jul. 1, 2010**

**Related U.S. Application Data**

(63) Continuation of application No. 11/678,596, filed on Feb. 24, 2007, now Pat. No. 7,794,998, which is a




Fig. 1




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E





FIG. 3







FIG. 5A

TGTCGATGATGAGCCCCGAGACGGGTCAACAACCACCGCTAGAGGACAATA  
 GCTGAGTCAACCCGCTGAGAACCGTCTCACACCGGATTGTGCCCAAAAAGAACA  
 CCGGGCTCTGACGCTCTCCTCCCTAC **21 R**  
 ATTTCCATGTTTGCCTAA **21 R** GCTCTGACGATAACCCTAAAAAATTTGACTAGCAA  
 ATAAAGAACCCCTGGGCCCTATAAAAGGGGAGAGCAACCTAAAAATGGGATCCCTT **U3 R**  
**poly (A) signal** TATA box  
 TTCTGCACCTCGCCAAACCCCTCCTCTGGCCACGGTCCGACTTTGGTCATTCCTGCCT **cap**  
 ACCTGAATCGCCGCTTCGGGATCGAGCCATCCCTCTTCTATTTGGTGGCACTTCGC  
 GCACTCCGGCCCTCCACTCGGTAAGATC **sd-LTR** CCACTGGGICGAGCTAGGCCATCACC  
 CCTGGCCGCTCCCTGGAGCTCTCTCGCGGGCTCTTAAGGTTGCTCCCCCTCAG  
 CAAAGGCCCCAGGGCTTCTCTACT **R U5** CCTTGTTCAAGTCTCTTTTGGGGTGG  
 ACCTAAATCGAAAGTAGCACTTCGCTGTICAGCAGGAGGCTTGGCCCCAGGGCCA  
 GCGCCTGTAAAGTTACCCAGCTCGGAGTTGGTCTCTAGAGAATCAGGGCTAAAG  
 CTGCTAGCCCTAGGAAAGAAGGCAACAGGTGGGGCTCGTCCGGGATTGATCACC **PBS**  
 TCCTGTATTGGCCCTTCCCTGTCTGAAAGCC

FIG. 5B



**FIG. 6**

HTLV-3(2026ND)

NLS

MAHFPGFGQSLLYGYFVYVFGDCVQADWCPTISGGLCSARLHRHALLATKPEHQITWDPIDGRVVSSALQY 70

CBP/P300 Binding

LIPRLPSFPTQRTTRTLKVLTPPTTAATPKIPPSFFHAVKKHTPFRNNCLELTLGEQLPAMSFDPGLRP 140

NES

QNIYTMWGSSVCLYLYQLSPMTWPLIPHVIFCHPEQLGAFLTRVPTKRLEELLYKIFLSTGAIILPE 210

NCFPTTLFQPTRAPAVQA?WHTGLLPCQKEIATPGLIWFTDGSMPISGPCPKEGQPSLVVQSSTFIFQQ 280

CR2 binding

EDZ

FQTKASHPAFLLSHKLIHYSSFHSLLHLLFEEYTTIPFSLLEFNEKGANVDDDEPRDGSQPPARGQIAE<sup>SPV</sup> 350


SEQ ID NO: 50


**FIG. 7**


ERRRRCREELEERKROKKERRQQLDCIDMLGFEGFCDLEGYIDFRESQQRAGCEES


Arginine rich Leucine Zipper

SEQ ID NO: 84












FIG. 10 (page 1 of 4)

TGACAGGGACAACGACCCTCTCCAGGGGCGACAGCAAGCCCCAAGGACA  
AAACTAGCAGGGACTAGTCATCAGCCAAAAAGGTCAACTGTCTCACACAAAT  
AAGGATCCGAAGGTTCTGACGTCCCAGCCCAGECTCAAAAACCAGGAAATCCA  
TAGAAATGCACCTCGGCCCTACCCACTTCCCCTATCATGAAAAACAAAGGCT  
GTGACGACTACCCCTTCCCAAAAAAATTTGCTTAAACCATCAATAAAGACA  
GCCTAGCCTATAATAAGCATGAGGATGGTTCAGGAGGGGGCTCGCTCTCTTC  
CGATCGCCCTGCTCACTCGAGTGTCCATCTCCTGGTCAATCAGTTGAGACGC  
CGCCGCTCGCCGTCTCCTGGTTGTCCACCTCTGAACCACCCCTTGGGTAA  
GTCCCCCTTGGTCCGAGCTTGGCTACGGTTTCTGTAGTCCCTCCCAGGGAAG  
TCTCCGAGACTGCCAAGCCTCTGCTTGC AAGGCTACGGCCCTCCACCCCTCT  
TCCCGTCCGTGTTAATCTCTTCGCGCAACC GAAAAACGAAAAGCCCTCCAG  
CTCTCTTGGCCCGGGCCAGGCCTGAGCCGCGGGCCGACCACCTTAAAGCC  
CGCTGFACTCAAACCCCTCCGGGAGGGGCCCTTACAGTAGGCGCCCGTCCC  
CCCGGGGAAAACATAACAAGTGGGGGCTCGTCCGGGATCTGTTCCGCTCTCGC  
CGTTCACCCCTCCCACTATGGGTACAGCCACACATCCAGTCCCGTCCCTAA  
GGCCCCAGGGGCTCTCCACCACCACTGGCTTAAATTTCTGCAGGCGGCTT  
ACCGCTGCAACCTGGACCCTCCGAATTCGATTTTACCAGTTAAGACGATTC  
CTTAAGCTAGCGCTCCAACCCCAAGTCTGGTTAAACCCCTATCGATTAATCCCT  
CCTAGCCGCCCTAATCCCAAGGGGTACCCCGGTCCGGGTGACCGAGATCGTT  
AATATCTCTCCGCGCTCATCCACCCCCAGCGCCCGGGCAATTTCCATGCC  
CACGGCCACCGGCCCGGCCCTGCCCCCAAGCTCAGGAGGCGCACACGCC  
CCCCCTTATGCGGAGCCTGCTGCGCTCCAGTGCCTTCCCATTATGCACTCCCA  
CGGGGCCCTTCGAGCCACCGCCCTGGCAGATGAAAGACTTACAAGCCATT  
AAACAAGAAATTAGCACTCAGCTCCCGGCACTCTCAATTTATGCATACCA  
TTCGACTTGCCATCCAGCAGTTTGACCTACGGCTAAAGATCTACATGATCTT  
TTGCAGTACTTGTGCTCGTCCCTTATTGTCTCCCTTACCACCAACAGCTACA  
AGCACTCATTGTGGAGGCAGAAACCCGAGGGTTGACAGGTTACAATCCATATG  
GCAGGGCCCCCTCCGGGTACAAGCAAAACAACCCCGCCAGCAAGGCCTCCAG  
AGAGAATACCAAAGTCTTTGGCTGBCCGCCTTTGCGGCCCTGCTGGTAACA  
CCCGAGATCCTTCTGGGCCGCAATATTGCAAGGCCTCGAGGAACCTTATTGT  
GCTTTGTAGAGCGCTCAATGCGGCCCTCGATAATGGTCTACCTGAAGGCA  
CACC AAAAGGAACCCATCCTGCGGCCGCTGCCATACTCCAATGCCAAACAAAGA  
ATGCCAGAAACTCCTCAGGCGCGGGGCCATACCAACAGTCCCCTTGGCGAA  
ATGCTCCGAGCCTGTCAAGCTTGGACACCAAAAGGATAAGACCAAAAGTTCTAG  
TAGTTCAGCCCCGTAAAACCCCTCCAACACAACCGTGCTTCCGGTGTGGAAA  
GGTGGGACACTGGAGCCGAGACTGCACTCAGCCTCGCCCCCTCCGGGGCC  
TGCCCCCTATGTCAGGACCATCCCACTGGAAGCGAGATTGCCCCAGCTAA  
AAACCCCGCCGGAGGCAGAAGAACCCCTCCTAGCGGATTTGCCCTGCCCTCT  
CCCGGAGGAAAAAACTCCECAGGGGGGAGAACTAGTCTCCCCCCGACCC  
GGTAAACGTGCCCTTCCCTGCTTCCCCTTGTCTCCTATGGCAGGCCCAACAATC  
TCTCTCAATAATTAAGTTTCTTCTTCGATCGCCACCCCTGGCATCACAGG  
CGCTCCTGGACACCGGAGCCGECTCACTGTCATGCCCCAGGTTTTGGCTCGG  
GGGCTCACGGACCTCCAGGACACCACCATTTCTGGGGGCCGGCGGTA AAAACC  
ACTCCAGTTTAAACTCCTACGGTGTCCGGTACATGTATACTTGGCCCTCCGT  
AGGGCTCCCGTGTCCCTTCCCTCATGTCTAATTGACACCAAGAATGAGTGGAC  
CATCATCGGCCGGGACGTCTCGACCAATGCCAGGGGCCCTTACTTACCG

FIG. 10 (page 2 of 4)

GAGGACCTCCGGCCCCGACCCAGTTATCCCGGTGACCACCCCTGCAGTCA  
TCGGCTTAGAACATCTTCCAGAGCCCCAGAGGTCAGCCAGTTTCCTTTAAAC  
CTGAACGCCCTCCAGGCCCTAATAGACCTGGTCTCCAAGGCACTGGAGGCTGG  
CCATAICGAACCTTACTCTGGACCAGGCAACAACCCAGTTFCTTCCCTGTFAAAA  
AAACCAACGGCAAGTGGCGATTTATCCATGACCTCAGGGCCACTAATGCCAT  
CACCCTACCCCTGGCTCGCCCTCCCCTGGCCCCCTTGATCTTACCAGCCTGC  
CACAGGCCCTTGGCCCATCTTCAGACCATCGATCTCACGGACCGCTTCTTCCAG  
ATFCCCTCCCAAAGCGATTCCAGCCCTACTTCGCCCTTACCATCCCCAGCC  
ATTAAATCATGGGCCCTGGGAGCAGGTACGCTTGGACAGTCTTCCCAAGGC  
TTCAAAAACAGCCCCACGCTCTTTGAGCAACAGCTGGCCAGCGTACTAGGCC  
CAGCCCGAAAAGCCTTCCCACATCCGTTCATCGTCCAATACATGGACGACAT  
CCTCTTGGCATGCCCCCTCCAGCAGAACFAGATCAGCTGGCCACCCCTTACCG  
CACAGCTATTGTCCTCTCATGGTCTCCAGTTFCCCAGGAAAAAACCCAACGC  
ACCCAGGAAAAATAACACTTCTGGGCCAAATCATACATCCAGATCACATCA  
CCTATGAAACCACCCCCACCATCCCCATFAAGGCACACTGGACCCCTGACTGA  
ACTGCAAACCCCTCTGGGGGAGCTCCAGTGGGTCTCCAAGGGGACTCCTGTC  
CTCCGAGAACACCTTCACTGTCTTACTCAGCCTTGAGAGGTCTCAAAGACCC  
CCGGGACACTATCACCCCTCGTTCATCTCACCTCCACGCTCTCCACAACATTC  
AGCAAAGCCCTGCATCACAATTGCCGCGGTCCCTTACTCTACGCTCCCCCTC  
CTTGGCTCATCTTCTCAGTCCATCCGGCACGACCTCAGTCTCTTCCAGAC  
AAATCATAAATGGCCCTAGTCTGGCTCCACGCCCCCATCCCCGACCAGC  
CTATGCCCTGGGGGCACATACTCGCTGCACTGTACTTACCCTTGACAAGTA  
TGCCTTGACGACTATGGCCAACTATGCAAATCATTCCATCATAACATGTCCA  
CCCAGGCCCTACACGATTTCTGTAATAAATTCCTCTCACCCAGCGTCCGATA  
TTAATTCACCACATGCATCGTTCTGTGATCTGGGCAGACAGCCACCGGGAC  
CCTGGCGAACCCCTTACAACCTCCCGGCCCTTCTCCGGGAACCCCAAGCTCCTC  
AGGCCGTGATTTTCCCTATCCCCAGTGGTTATAGATCAGGCCCTTGTCTGTT  
CTCTGATGGGTCTCCCCAAAAGGCCGCTATGTAATTTGGGACAAGGTCAATTC  
TCAGCCAGCGGTCCGGTCCCCCTGCCCCCCCATGCCAATAACTCAGCACAAAA  
GGGGGAATTAGTCGGACTCCTCTTGGGCTTGCAAGCCGCACAGCCCTGGCCA  
TCCCTTAACATTTTCTAGACTCAAAGTTCTCATCCGGTACTTCCAGTCCCTC  
GCTTCCGGGGCCCTTCCAAGGATCATCCACACACCACCGTCTCCAGGCGTCCCT  
GCCACACTCTCCAGGGCAAGGTCTGTGTATCTCCACCACACCCCGCAGCCAC  
ACCCAATGGCTGATCCCATCTCGACCCCTCAATGAATATACCGACTCTCTCAT  
TGTGCCCCCGTAACCCCTTGAAGCCTGAGGGCCCTCCATGCCCTCACCCACT  
GCAACCAACAGGCCCTCGTTFCCACGGAGCCACCCCTGCACAGGCTAAGCA  
ACTCGTGCAGGCCCTGCCGACCTGTCAAATCATTAAACCCCTCAACACCACATG  
CCGCGTGGCCACATCCGCGCGGCCACTTCCCAAACCACACATGGCAAGGAG  
ATGTCACCCACCTTAAGCACAAACGGACCCGATACTGCCTCCACGCTGGGT  
GGATACCTTCTCAGGTGCGGTATCTTGTGTCTGCAAAAAGAAAAGAACTAGC  
AGCGACCTTATCAAACCCCTCTACATGCCATCTCCGTGCTAGGCAAGCCCTT  
CTCTGTAAACACGGACAATGGACCCGCTTACCTTCTCAGGAGTTCACGAAT  
TCTGTACCACCTCTGCATCAAACACTCCACCATAATFCCCTACAATCCGACA  
AGTTACAGGCCCTGGTGGAGCGCACAAATGGCATTCTCAAAGACTACTATACA  
AATATTTCTAGACCACCCCTGACCTCCCCCTAGAAAAGCGCGGTTTCAAAGGCT  
CTCTGGACCATTAACCATTTAAATGTCATGCGCCCTGTGGTAAGACTCGGTG


FIG. 10 (page 3 of 4)

GCAGCTCCATCACACCCCCCCCCTGCCTCCTATTTCCGAGTCCATAACAAACCA  
CTCCACCCAGGCTACATTGGTACTATTACAAAACCCCTGGACTTACCAACCA  
GCGATGGAAAGGGCCCGTACAATCTCTCCAGGAAGCAGCAGGAGCAGCTCTC  
CTTCAAGTCAGTGACGGCTCGCCCCAGTGGATCCCTTGGCGGCTCCTGAAGA  
AGACTGTATGCCCCAAAACCCGACGACCCCGAACCCGCAGGGGCACGTCCGAAA  
CAGACCACCAACACCATGGGTAACGTACTCTTCTTAACTTTATTGGCCACCCF  
GGGCATCCCAGTACTTCAGGCCAGCCGGGTGTACAAATCACGGTAGGTATCTCC  
TCCTACCACTCCAGCCCCGTGCAGCCCAGCCCAGCCCTTATGTACCTGGGCCCT  
CGACCTTGTGTCCATCACFAAGGACCAGCTCCCTCTACCCCCCTGCCAAAAAC  
TGATCACCTATTCCAACACCACAAGAUCTACTCCCTGTATCTCTCCACAC  
TGGGTACAAAAGCCACTCCGCCGGGGCTTGGATACTACTCAGCCTCCCTACT  
CTGATCCTTGTCTCCCTACAATGTCCCTACCTAGGAAGTCAATCATGGACTTGC  
CCCTATACTGGCCCTGTCTCGAGCCCAACTTGGAGATTCCTCACAGATGTAAA  
TTTCACCCAAGAAGTCAGCCGTGTCTCCCTAAAACTTCATTTCTCCAAATGTG  
GTTCCTCCTTAACTCTGTAAATAGATGCCCCCGGTTACGATCCCGCTGTGGTAC  
CTCACATCCGAGCCTACTCAGGAACCCCAACCCCTCCGECCTAGTCAGCG  
ACTCAGACCTAGAGCATGTCTGACTCCTTCGGCCCTCCTGGGCCCTCCAAGATG  
CTGACCCCTCATCCACCTAACCTTGCAGAGCACCAACTATTCCCTGTATGGTCTG  
TATTGACCGCGCCAGCCTCTCTTCCCTGGCACGTATTATACACTCCCAACATCT  
CTAGTAATGCCCCCTCAAAAACCCATCGTCCGCCCTTCCCTTGGCCATCCGCC  
CCCGGACCACAGCCCTTCCCCTGGACCCATTGCTATCAACCACAGGTGCAAG  
CTGTAACCACCGCAAAGTGCAATAATTCATCATACTTCCCCCATTTTCTCTC  
TCTCCCTTGCCTGGTGGCCCTCTCACTAGGGCGACGCCGGGCCGTCCAGTGGC  
GGTCTGGCTCGTTTCCGCTTTGGCCGCAGGGACAGGAATAGCAGGAGGTGT  
ACCGGGTCCCTATCCCTGGCCCTCCAGTAGAAGTCTCCTGTCCGAAGTGGACA  
AGGATATTTCCACCTCACACGGGCCATTGTAAAAAACCAACCAAAACATTCT  
TCGAGTGGCCCAATATGCCGCCAAAACAGGGCGAGGGTTAGACCTCCTGTTC  
TGGGAACAAGGGGGGCTGTGTAAAGCGATACAAGAACAATGCTGCTTCTCA  
ACATCAGCAATACCCATAATTCAGTCTTACAAGAGCGACCCCTCTAGAAAC  
TCGGGTAACACTACTGGATGGGGCTTAAATTGGGATCTAGGACTCTCCAGTGG  
GCCCCGTGAGGCTCTCCAGACTGGTATTACCCTTTTGGCCCTCTCTGTTAATC  
ATCATCCTCGGGCCCTGCATTATTCGCCAGCTGCAAGCCCTCCCCAGAGGT  
ACAGCAGCGACCTGACCAGTACCCCTCCTCAACCCCTGAGACCCCTTATAAT  
AACTCCGCCAATACACCCAACAGGTCCCATGGTTGACCCCTCTACCGTTCAC  
CCACCCGCACTCCGCTAGACCTGACGAGTCCCCCATATGTCCAAAGTCTGTT  
CCAAGCCAGCTGATAACCGAAATAATTCCTTAAGTTATGGTTACATTCCTCC  
TCCAGATCCTTCCCTTCTCTAATAACATCAATATAGCCTTGCAACAAGTC  
ACAATACCCCTCAAACCCAGCAGGTCCATGCACTCCGTTGTTGATGACGC  
GCTCTCTCTCCTTGGCCTTCTCTCCCTCTCCTGCAATCGCTCCCTCCGCCGC  
GCTCTCTTCTCCTCTCTCTCGCAGGAGCCGCTGAATCTCCGCCCTGCTCGTCC  
ACCAGGGCCCTCAGGCGAGACTTCCGGGTACCATCATTTGGCGCCTCCCGACC  
CCAGGGGGCGGCCCTTTCGCGCACGACGAGCCCGCTACCAGGCATCTCCTC  
TGGTGTGAGACCTTCTTTGCCCGATCCTCTGATGATAACCCCTAAAAAATT  
CTATAAAAAAATCCCCGTIATTTTTTCAGCCCACTTCCCAGGATTCGGGCAG  
AGCCTCCTCTATGGATACCCCGTCTATGTGTTTGGCGATTGTGTTCAAGCCGA  
TTGGTGGCCCATCTCCGGTGGATTATGCTCCCCCGCCTACATCGCCACGCC

FIG. 10 (page 4 of 4)

TCC TGG CCA CCT G C C C C G A G C A C C A G A T C A C C T G G G A C C C C A T C G A T G G A C G  
A G T T G T C G G C T G C C T C T C C A A T A C C T T A T C C C T C G C C T C C C C T C C T T C C C C A C  
C C A A C G A A E C T U C A A G A C C C T C A A A G T C C T T A C C C C A C C A A C C A C T C C T G T C  
A C C C C C A A G G T T C C A C C C T C C T T C T T T C A G T C C G T G C G G A G G C A C A G C C C E T A  
C C G C A A C G G A T G T C T T G A A A C A A C C C T T G G A G A G C A G C T C C C C T C C C T T G C A  
T T C C T G A G C C A G G C C T C A G G C C C A A A A C G T C T A C A C C A T C T G G G G A A A G A  
C C A T A G T G T G T C T A T A C A T C T A C C A G C T G T C C C C T C C C A T G A C C T G G C C C C T C  
A T T C C C C A T G T C A T A T T T T G C A A C C C C A G G C A G C T T G G C C G T T T T C T A A G C A A  
T G T G C C C C C C A A G C G A T F A G A A G A A C T C C T C T A C A A A C T T T A T C T A C A C A C C G  
G E G C C A T A A T C A T C C T G C C G G A A G A C G C C C T G C C T A C C A C C C T A T T T C A G C C T  
G T T C G A G C A C C C T G T G T C C A A A C T A C C T G G A A C A C A G G A C T T C T C C C A T A C C  
A G C C A A A C C T G A C T A C C C C T G G C C T G A T A T G G A C C T T T A A T G A T G G G T C T C C T  
A T G A T T T C A G G A C C T T G C C C T A A G G C A G G G C A G C C A T C C T T G G T A G T A C A G T  
C C T C A C T A C T A A T C T T C G A G A G A T T T C A A A C C A A A G C C T A T C A T C C C C T T T A C  
C T C C T C T C C C A C C A A T T G A T A C A G T A T T C C T C C T T C C A T C A C C T C T A C T F A C T C  
T T T G A T G A A T A T A C T A C T A T C C C C T T C T C T A C T A T T T A A G G A A A A A G A G G G  
A G A T G A C A G G G A C A A C G A C C C T C T C C C A G G G G C G A C A G C A A G C C C C C A A G G  
A C A A A A C T A G C A G G G A C T A G T C A T C A G C C A A A A A G G T C A A C T G T C T C A C A C A  
A A T A A G G A T C C G A A G G T T C T G A C G T C C C A G C C C A G C C T C A A A A C C A G G A A A T  
C C A T A G A A A T G C A C C T C G C C C T T A C C C A C T T C C C C T A T C A T G A A A A A C A A A G  
G C T G T G A C G A C T A C C C C C T T C C C C A A A A A A T T T G C T T A A A C C A T C A A T A A A G  
A C A G C C T A G C C T A T A T A A G C A T G A G G A T G G T T C A G G A G G G G G C T C G C T C T C T  
T G C E G A T C G C C C T G C T C A C C T C G A G T G T C C A T C T C C T G G T C A A T C A G T T G A G A  
C G C C G C C G G C T G C C G G T C C C T G G T T G T C G C A C C T C C T G A A C C A C C C C T T G G G  
T A A G T C C C C C C T T G G T C C G A G C T T G G C T A C A G T T T C T G T A G T C G C T C C C A G G G  
A A G T C T C C G A G A C T G C C C A A G C C T C T G C T T G C A A G G C T A C G G C C C T C C A C C C  
C T C T C C G C G T C C G T G T T A A T C T C T T C G C G C C A A C C G A A A C G A A A G C G C C T C  
C A G C T C T T T G G C C C G G G C C A G G C C T G A G C C G C G C G G G C G C A C C A C C T T A A  
G C C C C G T G T A C T C A A A C C C C T C C G G G A G G G G C C C T T T A C A G T A G G C G C C C G T  
C C C C C C G G G G A A A C A T A C A

FIG. 11



# Identification of Novel HTLVs in Cameroon




FIG. 12

*PTLV-3 and PTLV-4 are ancient lineages*




FIG. 13

## PRIMATE T-LYMPHOTROPIC VIRUSES

### CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This is a continuation of U.S. patent application Ser. No. 11/678,596, filed Feb. 24, 2007, which is a continuation-in-part of International Application No. PCT/US2006/005869, filed Feb. 21, 2006, which claims the benefit of U.S. Provisional Application No. 60/654,484, filed on Feb. 21, 2005, now expired. Each of these prior applications is incorporated by herein by reference.

### ACKNOWLEDGEMENT OF GOVERNMENT SUPPORT

[0002] Aspects of this invention were made with United States government support. Therefore, the government has certain rights in the invention.

### FIELD

[0003] Disclosed are compositions and methods related to the isolation and identification of the primate T-lymphotropic viruses, HTLV-3 and HTLV-4. The present disclosure also relates to vectors and vaccines for use in humans against infection and disease.

### BACKGROUND

[0004] Primate T-lymphotropic viruses (PTLVs) are diverse deltaretroviruses, composed of 3 distinct species (PTLV-1, -2, -3) which by conventional nomenclature are named 'STLV' (simian T-lymphotropic virus) when found in non-human primates (NHPs) and 'HTLV' (human T-lymphotropic virus) when found in humans, regardless of suspected zoonotic origin (Mahieux et al., 1998; Salemi et al. 1999; Slattery et al., 1999; Courgnaud et al., 2004). Like HIV, HTLV has the potential to cause disease and circulate globally in humans sexually, from mother-to-child, and by exposure to contaminated blood from transfusions and intravenous drug use. HTLV-1 causes adult T-cell leukemia and HTLV-1-associated myelopathy/tropical spastic paraparesis (HAM/TSP) and other inflammatory diseases (Gessain & Mahieux 2000) and HTLV-2 has been associated with a neurologic disease similar to HAM/TSP (Araujo & Hall 2004). There has been no evidence to date of STLVs crossing into people occupationally exposed to NHPs in laboratories and primate centers, as has been documented with other primate retroviruses, including simian immunodeficiency virus (SIV) (Khabbaz et al., 1994), simian foamy virus (SFV) (Switzer et al., 2004, Heneine et al., 1998), and simian type D retrovirus (Lerche et al. 2001). Nevertheless, ongoing zoonotic transmission of STLV to widespread human populations naturally exposed to NHPs through hunting or butchering, similar to that recently reported for SFV in African hunters (Wolfe et al., 2004b), would be of particular public health significance due to the transmissible and pathogenic nature of this group of viruses among humans. HTLV outside of the PTLV-1 and PTLV-2 groups has not previously been documented (Busch et al. 2000; VanDamme et al. 1997; Salemi et al. 1999; Slattery et al. 1999).

### SUMMARY

[0005] Disclosed herein are compositions and methods that include the full and partial nucleic acid sequences of primate

T-lymphotropic viruses known as HTLV-3 and HTLV-4, including viral fragments. These viruses are useful as reagents for the screening of human populations for the prevalence of such viruses. The disclosed viruses also can serve as vectors in gene therapy because the viruses appear to not be transmitted from humans to other humans. Additionally, the disclosed viruses can be used as reagents in pathogenicity studies of these and related viruses. Moreover, the sequences of the primate T-lymphotropic viruses described herein can be used as probes to detect virus in biological samples. Vectors are disclosed that express the HTLV-3 and HTLV-4 nucleic acid sequences, and include, but are not limited to, prokaryotic, eukaryotic and viral vectors. The disclosed viruses also can be used as live recombinant virus vaccines. Additionally, the disclosed viruses can be used as replicating viral systems to kill live dividing cells, either in vitro or in vivo.

[0006] The present disclosure also includes the isolation and characterization of primate T-lymphotropic viruses, HTLV-3 and HTLV-4, that are believed to have been transmitted from non-human primates to humans at some point in the past. The primate T-lymphotropic viruses described herein do not appear to be readily transmitted from human to human, and can be used in protocols for diagnosing primate T-lymphotropic virus infections, and as vectors in gene therapy procedures.

[0007] Compositions are provided that include live replicating retroviral vectors, wherein the vector is derived from a primate T-lymphotropic virus, and wherein the vector also includes a nucleic acid that encodes a primate T-lymphotropic virus peptide, polypeptide, or protein, or a fragment of a primate T-lymphotropic virus peptide, polypeptide, or protein. For example, the vector can be derived from an HTLV-3 or HTLV-4 virus. Thus, in one aspect, a composition is provided that includes live replicating primate T-lymphotropic virus vectors.

[0008] Also provided is a method of treating a subject with a condition, wherein the condition is a viral infection, bacterial infection, parasitic infection, proliferative disorder (e.g., cancer), or a condition associated with a genetic or autoimmune disorder. The method includes administering to the subject a live replicating viral vector, wherein the immunizing construct is specific for the condition.

[0009] Also provided is a method of preventing a condition in a subject, wherein the condition can be a viral infection, bacterial infection, parasitic infection, proliferative disorder, or a condition associated with a genetic or autoimmune disorder. The method includes administering to the subject a live replicating viral vector, wherein the antigen-encoding nucleic acid is specific for the condition. Also provided are methods of using the vectors, isolated viruses, and/or infectious clones described herein for making viral infection models and using models to study diseases and potential treatments, as well as the models themselves.

[0010] Also disclosed are methods and compositions for detecting primate T-lymphotropic virus or a protein encoded therein in biological fluids. The disclosure also encompasses antibodies specific for the primate T-lymphotropic virus and antibodies that inhibit the binding of antibodies specific for the primate T-lymphotropic virus. These antibodies can be polyclonal antibodies or monoclonal antibodies, which also includes fragments of any type of antibody. Thus, disclosed are antibodies to HTLV-3 or HTLV-4. The antibodies specific for the primate T-lymphotropic virus can be used in diagnos-

tic kits to detect the presence and quantity of primate T-lymphotropic virus in biological fluids or in organs from nonhuman primates for xenotransplantation. For example, an HTLV-3 antibody can be used in a diagnostic kit to detect HTLV-3. Antibodies specific for primate T-lymphotropic virus may also be administered to a human or animal to passively immunize the human or animal against primate T-lymphotropic virus, thereby reducing infection, for instance after accidental exposure to nonhuman primate bodily fluids.

**[0011]** Other embodiments of the disclosure are methods and kits for detecting the presence and quantity of antibodies that bind primate T-lymphotropic virus, for example in body fluids. Such kits can be used for the detection of primate T-lymphotropic virus itself, or for the detection of antibodies to the primate T-lymphotropic virus, and also can be used to monitor the blood supply for the presence of primate T-lymphotropic virus. The disclosed kits include, for example, a kit for the detection of antibodies to HTLV-3 or HTLV-4.

**[0012]** Also included in the disclosure are recombinant live virus vaccines. The virus of the present disclosure has areas of its genome that make it useful for the insertion of exogenous genes. The inserted gene(s) can code for any protein for which vaccination or gene therapy is desired. A useful aspect of such recombinant live viruses is that the recombinant HTLV-3 or HTLV-4 does not cause disease in the host organism. The recombinant live virus vaccines of the present disclosure are a safe way to provide antigen to the immune system.

**[0013]** Accordingly, provided is a composition comprising a primate T-lymphotropic virus, or a fragment of the viral gene or the encoded protein. An example of the disclosed primate T-lymphotropic virus includes, but is not limited to HTLV-3 and HTLV-4. Also provided is a method of detecting a primate T-lymphotropic virus, such as HTLV-3 or HTLV-4.

**[0014]** Also provided are methods and compositions for detecting the presence and amount of primate T-lymphotropic virus in a body fluid or organ. Further embodiments are compositions and methods for treating genetic and physiologic disorders using gene therapy techniques that include the primate T-lymphotropic virus of the present disclosure as a vector for nucleic acid sequences and antisense sequences.

**[0015]** Further embodiments include providing compositions and methods useful for manipulating the expression of genes, providing vaccines, providing compositions and methods for treating viral infections in humans or animals, providing compositions and methods that are effective in treating genetic diseases, and providing a method of treating microbial infections in humans or animals. Yet still other embodiments include providing for treatments of conditions that are caused in part by rapidly dividing cellular growth, providing live recombinant virus vaccines, and providing diagnostic tools such as antibodies or antigens for the monitoring of the blood supply or organ and tissue donation for the presence of primate T-lymphotropic virus.

**[0016]** These and other features and advantages will become apparent after a review of the following detailed description of the disclosed embodiments and the appended claims.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0017]** The accompanying drawings illustrate several embodiments and, together with the description, illustrate the disclosed compositions and methods.

**[0018]** FIG. 1 is a digital image showing the Western blot serological pattern of Human T-cell lymphotropic virus (HTLV) infected African hunters. HTLV classification based on phylogenetic analyses is provided above specimen names. Reactivity to HTLV-specific proteins is indicated on left.

**[0019]** FIG. 2 shows the phylogenetic relationships of (a) primate T-cell lymphotropic virus polymerase (a) PTLV pol (662-bp), (b) PTLV env (297-bp), (c) PTLV tax (730-bp), (d) PTLV-3 long terminal repeat (LTR) (398-bp), and (e) PTLV pol-env-tax region (5258-bp) sequences by neighbor joining analysis. Sequences generated in the current study are noted with boxes. Nonhuman primate taxon codes are provided in the Methods portion of the Examples section of the specification. Support for the branching order was determined by 1,000 bootstrap replicates, and only values 60% or greater are shown. Branch lengths are proportional to the evolutionary distance (scale bar) between the taxa.

**[0020]** FIG. 3 shows the phylogenetic relationships of PTLV type 1 LTR (377-bp) sequences by neighbour joining analysis. Sequences generated in the current study are noted with boxes. Nonhuman primate taxon codes are provided in the Methods portion of the Examples section of the specification. Support for the branching order was determined by 1,000 bootstrap replicates and only values 60% or greater are shown. Branch lengths are proportional to the evolutionary distance (scale bar) between the taxa.

**[0021]** FIG. 4 shows the strategy for PCR-amplifying the entire HTLV-3 (2026ND) genome. Small proviral sequences were first amplified in each major gene region and the long terminal repeat (stippled bars) using generic primers as described in the Methods portion of the Examples section of the specification. The complete proviral sequence was then obtained by using PCR primers located within each major gene region by genome walking as indicated with arrows and orange bars. The typical HTLV-1 genomic organization is provided for reference.

**[0022]** FIG. 5(a) shows the nucleotide sequence of the HTLV-3 (2026ND) LTR and pre-gag region (nucleotides 1-755 of SEQ ID NO: 36). The U3-R-U5 locations (vertical lines), the approximate cap site (cap), the polyadenylation signal, TATA box, the predicted splice donor site (sd-LTR), and two 21-bp repeats are indicated. In the R and U5 regions, the predicted Rex core elements and nuclear riboprotein A1 binding sites are underlined. The pre-gag region and primer binding site (PBS, underlined) are in italics. FIG. 5(b) shows the plot of predicted RNA stem loop secondary structure of HTLV-3(2026ND) LTR region (nucleotides 421-464 of SEQ ID NO: 36). Position of the Rex responsive element (RexRE) core is indicated.

**[0023]** FIG. 6 shows the amino acid sequence of HTLV-3 Tax (SEQ ID NO: 50). Shown in boxes are known functional motifs: NLS, nuclear localization signal; (CBP)/P300, cAMP response element (CREB) binding protein; NES, nuclear export signal; CR2, C-terminal transcriptional activating domain binding; PDZ.

**[0024]** FIG. 7 shows the amino acid sequence of a basic leucine zipper (bZIP) transcription factor from HTLV-3 (SEQ ID NO: 84). Arginine rich and leucine zipper regions of the bZIP protein are boxed.

**[0025]** FIG. 8 shows the phylogenetic relationship of HTLV-3(2026ND) to other PTLVs (a) entire genome sans long terminal repeat (LTR), (b) gag, (c), polymerase (pol), and (d) envelope (env). Sequences generated in the current study are shown in boxes. Support for the branching order

was determined by 1,000 bootstrap replicates; only values of 60% or more are shown. Branch lengths are proportional to the evolutionary distance (scale bar) between the taxa.

**[0026]** FIG. 9 shows the estimated divergence dates for the most recent common ancestor of HTLV-3(2026ND) and other PTLVs. Divergence dates are provided for each major node of a neighbour-joining tree rooted with PTLV-1 as the outgroup; estimates are provided as ranges using as calibration points 40,000 and 60,000 years ago (YA) as the separation of the Melanesian HTLV-1 (MEL5) sequence from other PTLV-1 strains. Bootstrap analysis of 1000 replicates is shown on the tree branches; only values >60% are shown.

**[0027]** FIG. 10 (which includes four pages) shows the full-length genomic sequence of HTLV-4(1863LE) (SEQ ID NO: 81).

**[0028]** FIG. 11 shows the plot of predicted RNA stem loop secondary structure of the HTLV-4(1863LE) LTR region. Position of the Rex responsive element (RexRE) core is indicated (nucleotides 425-466 of SEQ ID NO: 81).

**[0029]** FIG. 12 shows the phylogenetic relationships of PTLV full-length genomic sequences, including full-length genomic HTLV-3 and HTLV-4. These findings confirm the genetic relationships found earlier that were based on smaller sequences. Four major phylogroups were inferred with very high bootstrap support. Nonhuman primate taxon codes are provided in the Methods portion of the Examples section of the specification. Support for the branching order was determined by 1,000 bootstrap replicates and only values 60% or greater are shown. Branch lengths are proportional to the evolutionary distance (scale bar) between the taxa.

**[0030]** FIG. 13 shows the estimated divergence dates for the most recent common ancestor of HTLV-3(2026ND), HTLV-4(1863LE) and other PTLVs. Divergence dates are provided for each major node of a neighbor-joining tree rooted with PTLV-1 as the outgroup; estimates are provided as ranges using as calibration points 40,000 and 60,000 years ago (YA) as the separation of the Melanesian HTLV-1 (MEL5) sequence from other PTLV-1 strains. Using the bovine leukemia virus (BLV) as an outgroup, a substitution rate of  $8.6 \times 10^{-7}$  to  $1.3 \times 10^{-6}$  substitutions/site/year for PTLV was inferred which is 3 logs lower than that seen in HIV, confirming the genetic stability of these deltaretroviruses. Bootstrap analysis of 1,000 replicates is shown on the tree branches; only values >60% are shown.

#### DETAILED DESCRIPTION

**[0031]** Before the present compounds, compositions, articles, devices, and/or methods are disclosed and described, it is to be understood that they are not limited to specific synthetic methods or specific recombinant biotechnology methods unless otherwise specified, or to particular reagents unless otherwise specified, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

##### I. Terms

**[0032]** As used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a pharmaceutical carrier” includes mixtures of two or more such carriers, and the like.

**[0033]** Ranges can be expressed herein as from “about” one particular value, and/or to “about” another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent “about,” it will be understood that the particular value forms another embodiment. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint, and independently of the other endpoint. It is also understood that there are a number of values disclosed herein, and that each value is also herein disclosed as “about” that particular value in addition to the value itself. For example, if the value “10” is disclosed, then “about 10” is also disclosed. It is also understood that when a value is disclosed that “less than or equal to” the value, “greater than or equal to the value” and possible ranges between values are also disclosed, as appropriately understood by the skilled artisan. For example, if the value “10” is disclosed the “less than or equal to 10” as well as “greater than or equal to 10” is also disclosed. It is also understood that throughout the document, data are provided in a number of different formats, and that this data, represents endpoints and starting points, and ranges for any combination of the data points. For example, if a particular data point “10” and a particular data point 15 are disclosed, it is understood that greater than, greater than or equal to, less than, less than or equal to, and equal to 10 and 15 are considered disclosed as well as between 10 and 15.

**[0034]** “Optional” or “optionally” means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances where it does not.

**[0035]** “Primers” are a subset of probes which are capable of supporting some type of enzymatic manipulation and which can hybridize with a target nucleic acid such that the enzymatic manipulation can occur. A primer can be made from any combination of nucleotides or nucleotide derivatives or analogs available in the art which do not interfere with the enzymatic manipulation.

**[0036]** “Probes” are molecules capable of interacting with a target nucleic acid, typically in a sequence specific manner, for example through hybridization. The hybridization of nucleic acids is well understood in the art and discussed herein. Typically a probe can be made from any combination of nucleotides or nucleotide derivatives or analogs available in the art.

**[0037]** Depending on context, the term “virus” is understood to include the infectious viral particle or the nucleic acid contained therein, or both.

**[0038]** Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this disclosure pertains. The references disclosed are also individually and specifically incorporated by reference herein for the material contained in them that is discussed in the sentence in which the reference is relied upon.

##### II. Compositions

**[0039]** Disclosed are the components to be used to prepare the disclosed compositions as well as the compositions themselves to be used within the methods disclosed herein. These and other materials are disclosed herein, and it is understood that when combinations, subsets, interactions, groups, etc. of

these materials are disclosed that while specific reference of each various individual and collective combinations and permutation of these compounds may not be explicitly disclosed, each is specifically contemplated and described herein. For example, if a particular HTLV-3 or HTLV-4 or gene of the HTLV-3 or HTLV-4 such as gag, pol, env, LTR, rex, and tax is disclosed and discussed and a number of modifications that can be made are discussed, specifically contemplated is each and every combination and permutation of HTLV-3 or HTLV-4 or genes of the HTLV-3 or HTLV-4 such as gag, pol, env, LTR, rex, and tax and the modifications that are possible unless specifically indicated to the contrary. Thus, if a class of molecules A, B, and C are disclosed as well as a class of molecules D, E, and F and an example of a combination molecule, A-D is disclosed, then even if each is not individually recited each is individually and collectively contemplated meaning combinations, A-E, A-F, B-D, B-E, B-F, C-D, C-E, and C-F are considered disclosed. Likewise, any subset or combination of these is also disclosed. Thus, for example, the sub-group of A-E, B-F, and C-E would be considered disclosed. This concept applies to all aspects of this application including, but not limited to, steps in methods of making and using the disclosed compositions. Thus, if there are a variety of additional steps that can be performed it is understood that each of these additional steps can be performed with any specific embodiment or combination of embodiments of the disclosed methods.

**[0040]** Furthermore, although the disclosed nucleic acid sequences are represented as DNA sequences, it is understood that the equivalent RNA sequences also are contemplated. For instance, if a DNA sequence contains a thymine, it is understood that a uracil also can be substituted.

**[0041]** Disclosed herein are compositions relating to primate T-lymphotropic viruses HTLV-3 (SEQ ID NO: 36) and HTLV-4 (SEQ ID NOs: 53 and 81). It is understood and herein contemplated that the compositions of the disclosure can comprise the entire HTLV-3 or HTLV-4 virus nucleic acid sequence. It is also understood that the disclosed compositions can comprise proteins of the disclosed primate T-lymphotropic viruses or fragments of the disclosed proteins. For example, specifically disclosed and herein contemplated are compositions comprising SEQ ID NOs: 1, 3, 5, 35, 45, 47, 49, 51, and 52, or any combination thereof. Also disclosed are compositions comprising SEQ ID NOs: 2, 4, 6, 59, 61, and 63 or any combination thereof. Also disclosed are compositions comprising SEQ ID NOs: 37, 40, 44, 46, 48, and 50 or any combination thereof. Also disclosed are compositions comprising SEQ ID NOs: 54, 57, 58, 60, and 62 or any combination thereof. Also disclosed are compositions comprising fragments of the disclosed proteins. Thus, for example are compositions comprising SEQ ID NOs: 38, 39, 41, 42, and 43 or any combination thereof. Also disclosed are compositions comprising SEQ ID NOs: 55 and 56. It is understood and herein contemplated that any of the disclosed proteins can be used in combination with any of the protein fragments in the compositions disclosed herein. Thus, for example, disclosed herein are compositions comprising SEQ ID NOs: 37, 38, 39, 40, 41, 42, 43, 44, 46, 48, and 50 or any combination thereof. Also disclosed are SEQ ID NOs: 54, 55, 56, 57, 58, 60, and 62 or any combination thereof. SEQ ID NOs 1-6, 35, and 45 can be used for all the molecular biological techniques known to those skilled in the art. Such uses include, but are not limited to, generation of probes and vectors containing the sequences, antisense sequences derived from such sequences,

and proteins synthesized using the sequences. RNA and other nucleic acid derivatives are contemplated by the present disclosure.

**[0042]** It is understood that there are known viruses in the art that based on certain genomic or sequence similarity or taxonomically related to the viruses disclosed herein. It is also understood that the known viruses in the art thought related taxonomically do not encode the specific viruses disclosed herein. Thus specifically disclosed and herein contemplated are isolated primate T-lymphotropic viruses having a pol gene that has less than 63.5% identity to the pol gene of HTLV-1, HTLV-2, STLV-2, and STLV-3, for example, HTLV-4. Also disclosed are isolated primate T-lymphotropic viruses having a gag gene that has less than 69% identity to the gag gene of HTLV-1, HTLV-2, STLV-2, and STLV-3, for example, HTLV-3. Also disclosed are isolated primate T-lymphotropic viruses having a pol gene that has less than 62% identity to the pol gene of HTLV-1, HTLV-2, STLV-2, and less than 86% identity to the pol gene of STLV-3, for example, HTLV-3. Similarly, the disclosed viruses can be distinguished based on the genes encoded by the disclosed viruses, and specifically the identity of said genes to the corresponding genes of known viruses. Thus, specifically disclosed are isolated primate T-lymphotropic viruses having a LTR that has less than 41% identity to the LTR of HTLV-1, HTLV-2 and STLV-3. Also disclosed are isolated primate T-lymphotropic viruses having at least 92.8% identity to the nucleic acid SEQ ID NO: 1.

**[0043]** Also disclosed are isolated primate T-lymphotropic virus having at least 92.5% identity to the nucleic acid SEQ ID NO: 3. Also disclosed are primate T-lymphotropic viruses having at least 94.2% identity to the nucleic acid SEQ ID NO: 5. Also disclosed are primate T-lymphotropic viruses having at least 91.5% identity to the nucleic acid SEQ ID NO: 35. Also disclosed are isolated primate T-lymphotropic viruses having at least 92.8% identity to the nucleic acid SEQ ID NO: 1, at least 92.5% identity to the nucleic acid SEQ ID NO: 3, and at least 94.2% identity to the nucleic acid SEQ ID NO: 5.

**[0044]** HTLV-4 is a unique delta primate T-lymphotropic virus that is distinct from all known PTLV lineages with 29-34.4% and 18.3-25% nucleotide divergence in the conserved pol and tax genes, respectively, a range of divergence similar to that between PTLV-1, PTLV-2, and PTLV-3. This virus formed a separate phylogenetic lineage with a long branch length and significant bootstrap support in both the pol (FIG. 2a) and tax (FIG. 2c) trees. Identical topologies were obtained by using maximum likelihood analysis. Phylogenetic analyses combined with GenBank blast searches show that this is the only known virus in this group. For these reasons, this virus, which was designated HTLV-4, qualifies as the first member of a group in the deltaretrovirus genus. Following the guidelines of the International Committee on Taxonomy of Viruses and pending formal classification, primate T-lymphotropic virus 4 (PTLV-4) was proposed as the name for this species, and PTLV-4(1863LE) as the prototype strain. Due to the classification of the virus within the family retroviridae, certain sequence similarity is expected to exist with known retroviruses. It is understood that the known viruses in the art thought to be related taxonomically do not encode the specific viruses disclosed herein. Thus, specifically disclosed and herein contemplated are isolated primate T-lymphotropic viruses having at least 71.5% identity to the nucleic acid SEQ ID NO: 2. Also disclosed are isolated primate T-lymphotropic viruses having at least 73.5% identity to the nucleic acid SEQ ID NO: 4. Also disclosed are isolated

primate T-lymphotropic viruses having at least 82% identity to the nucleic acid SEQ ID NO: 6. Also disclosed are isolated primate T-lymphotropic viruses having at least 71.5% identity to the nucleic acid SEQ ID NO: 2, at least 73.5% identity to the nucleic acid SEQ ID NO: 4, and at least 82% identity to the nucleic acid SEQ ID NO: 6.

**[0045]** Knowing the sequence for HTLV-3 and/or HTLV-4 allows for various uses of the virus and viral sequences. The env gene of HTLV-3 and/or HTLV-4 is necessary for primate T-lymphotropic virus entry into animal cells. The genes of the present disclosure are effective in permitting infection of cells in a human host. Thus, for example, the env gene is used for uptake of foreign DNA by a wide range of human cells. There has long been a need for vectors for getting foreign nucleic acids into cells, both in vivo and in vitro. The introduction of foreign or exogenous nucleic acids into cells has been a technological hurdle for many gene therapy applications and has now been solved by the virus and sequences herein disclosed. The env sequences can be used with any vector known to those skilled in the art, and with any other genetic sequences of choice, to allow for entry of the nucleic acids into the cells.

**[0046]** The recent advent of technology, and advances in the understanding of the structure and function of many genes makes it possible to selectively turn off or modify the activity of a given gene. Alteration of gene activity can be accomplished many ways. For example, oligonucleotides that are complementary to certain gene messages or viral sequences, known as "antisense" compounds, have been shown to have an inhibitory effect against viruses. By creating an antisense compound that hybridizes with the targeted RNA message of cells or viruses the translation of the message into protein can be interrupted or prevented. In this fashion gene activity can be modulated.

**[0047]** The ability to deactivate specific genes provides great therapeutic benefits. For example, it is possible to fight viral diseases with antisense molecules that seek out and destroy viral gene products. In tissue culture, antisense oligonucleotides have inhibited infections by herpes-viruses, influenza viruses and the human immunodeficiency virus that causes AIDS. It is also possible to target antisense oligonucleotides against mutated oncogenes. Antisense technology also can be used to regulate growth and development. However, in order for the gene therapy to work, antisense sequences must be delivered across cellular plasma membranes to the cytosol.

**[0048]** Gene activity is also modified using sense DNA in a technique known as gene therapy. Defective genes are replaced or supplemented by the administration of "good" or normal genes that are not subject to the defect. Instead of being defective, the gene may have been deleted, thus replacement therapy would provide a copy of the gene for use by the cell. The administered normal genes can either insert into a chromosome or may be present as extracellular DNA and can be used to produce normal RNA, leading to production of the normal gene product. In this fashion gene defects and deficiencies in the production of a gene product may be corrected.

**[0049]** Still further gene therapy has the potential to augment the normal genetic complement of a cell. For example, one way to combat HIV is to introduce into an infected person's T cells a gene that makes the cells resistant to HIV infection. This form of gene therapy is sometimes called "intracellular immunization." Genetic material such as a polynucleotide sequence may be administered to a mammal

in a viral vector to elicit an immune response against the gene product of the administered nucleic acid sequence. Such gene vaccines elicit an immune response in the following manner. First, the viral vector containing the nucleic acid sequence is administered to a human or animal. Next, the administered sequence is expressed to form a gene product within the human or animal. The gene product inside the human or animal is recognized as foreign material and the immune system of the human or animal mounts an immunological response against the gene product. The viruses disclosed herein can be used as viral vectors to provide the foreign nucleic acid sequences to the intracellular metabolic processes.

**[0050]** Additionally, gene therapy can be used as a method of delivering drugs in vivo. For example, if genes that code for therapeutic compounds can be delivered to endothelial cells, the gene products would have facilitated access to the blood stream. Additionally, cells could be infected with a retroviral vector such as the present disclosure carrying nucleic acid sequences coding for pharmaceutical agents that prevent infection from occurring in the retrovirally infected cells.

**[0051]** The primate T-lymphotropic viruses of the present disclosure can also be used as a safe and effective vaccine agent. Genetic sequences for immunogenic proteins from a variety of infectious agents can be incorporated into the primate T-lymphotropic virus RNA. Once inside a cell, the gene product is expressed and releases the immunizing peptide to the body's immune system. In another method, the disclosed viruses can be used to immunize the body against cell markers found on cancer or tumor cells. The genetic sequence of the cancer cell marker is incorporated into the primate T-lymphotropic virus RNA, and after infection with the virus, the expressed gene product stimulates the immune system. The subject's immune system is used to remove the cancerous cells, obviating the need for chemotherapeutic methods.

**[0052]** Such treatment with HTLV-3 or HTLV-4 can be used for any condition in which rapidly dividing cells provide an aspect of the pathology of the condition. One such condition is the presence of uncontrolled angiogenesis within the body. Angiogenesis dependent diseases are well known in the art and are caused in part by the rapid growth of blood vessels. Another such condition is cancer or tumor growth. Cancer or tumors include both solid tumors and other types. Infection with the virus of the present disclosure, which can cause no disease and does not affect the host systemically, is an improvement over currently known treatments that involved systemically administered agents. Such chemotherapeutic agents kill rapidly dividing cells but also cause trauma to the entire person. The dosages of such chemotherapeutic agents must be titrated between killing the cancer and killing the subject.

**[0053]** In contrast, the cancer treatments disclosed are not as harmful to the subject. The virus can either be administered systemically or injected in situ into the tumor. The infected cells are killed and tumor growth is stopped. The virus may be administered in one treatment or in a series of treatments.

**[0054]** The HTLV-3 or HTLV-4 of the present disclosure can be recombinantly modified to be selective for cellular receptors on the tumor to make the virus even more specifically targeted to just those cells. Additionally, the virus may have altered promoter regions that can be selectively activated to cause a productive infection. The combination of different levels of control of the virus, both natural and recombinantly-produced, are contemplated herein. A virus can be made

specific for attachment to only certain types of cellular receptors, for those cells that are dividing, and will only undergo replication if another exogenous promoter factor is present. Viral infection by two or more individually defective viruses, that require factors or promoters supplied by other primate T-lymphotropic viruses or any type of virus, can provide for many levels of control of infection or treatment of specific conditions.

**[0055]** The virus may be administered to the host, for cancer treatment, gene therapy or vaccination by any methods known to those skilled in the art. Such methods include but are not limited to injection, inhalation, ingestion, topical administration and implantation. The virus may be killed or live, depending on the treatment considered.

**[0056]** The antibodies disclosed herein can be used to detect the presence of the disclosed viruses or viral particles. These antibodies can be used in diagnostic or screening kits to assess the presence of the virus. Additionally, the antibodies can be used to screen organs from nonhuman primates that may be used in humans. For instance, detection of the presence of a virus that is transmitted from nonhuman primates to humans is crucial in providing virus-free organs for transplantation.

**[0057]** It is believed that the virus of the present disclosure, comprising the isolates from HTLV-3, is the first definitive isolation of an STLV-3-like primate T-lymphotropic virus from persons exposed to nonhuman primates. This belief is supported by the epidemiology data, the PCR and sequencing data and the serology data and the absence of such reports in the literature. It is understood that HIV-1 and HIV-2 used to be called HTLV-III and HTLV-IV before it was known they were different types of viruses. Additionally, the virus of the present disclosure comprising the isolates from HTLV-4, are a new species in the delta primate T-lymphotropic viruses.

### III. Vectors

**[0058]** Disclosed are live replicating human primate T-lymphotropic virus vectors suitable for human use comprising an immunizing construct, wherein the immunizing construct is inserted in nontranslated region between env and tax/rev. The disclosed immunizing construct can be an antigen-encoding nucleic acid.

**[0059]** Where reference is made to "antigen"-encoding nucleic acid, it is understood that in the context of the disclosure antigens encoded by the antigen-encoding nucleic acid can include but are not limited to immunogenic or non-immunogenic peptides, polypeptides, proteins, enzymes, cytokines. These antigens can be non-human exogenous antigenic sequences from viruses, bacteria, or parasites. The antigens can also be antigenic endogenous human or human derived sequences from a condition such as a cancer. Also, peptides encoded by the antigen-encoding nucleic acid can include non-antigenic sequences for the purposes of gene therapy.

**[0060]** In another embodiment of the present disclosure, sequences of the disclosed primate T-lymphotropic viruses can be used for other molecular biological applications. Regions of the gag gene are important in packaging genetic material. For example, the gag sequence or regions of the sequence are incorporated into other vectors and direct the packaging of the resultant genetic material for the particular application desired, such as packaging recombinant sequences to make altered infectious virions. Regions of the pol gene are known to be critical for the stable integration of foreign/viral DNA into the host genome. Vectors comprising

the pol gene sequences can be used to integrate any DNA into a genome. The primate T-lymphotropic virus and sequences of the present disclosure infect human cells, and thus, these sequences are used with other foreign or exogenous sequences in humans in methods, including, but not limited to, entry into cells, packaging, and insertion into the genome. Additionally, methods of using the disclosed primate T-lymphotropic virus and other sequences of the present disclosure are not limited to human cells, but all cells that allow for infection or entry of the nucleic acids.

**[0061]** The present disclosure is directed to compositions and methods comprising new primate T-lymphotropic viruses, HTLV-3 and/or HTLV-4, particularly compositions and methods for the sequences of the viral genome. The virus was obtained from humans. The new virus of the present disclosure is an excellent vector for gene therapy and for vaccination purposes. Additionally, the antibodies or other detection methods for detecting the new virus are important in detecting the presence of this and related viruses for xenotransplantation. In addition, the disclosed primate T-lymphotropic viruses can be used as reagents in pathogenicity studies of these and related viruses. Moreover, the sequences of the disclosed primate T-lymphotropic viruses can be used as probes to detect virus in biological samples. Vectors include but are not limited to prokaryotic, eukaryotic and viral vectors.

**[0062]** Many new useful technologies have been developed that use viral vectors and form the basis of medical therapies. Examples of such technologies include, but are not limited to, gene replacement, antisense gene therapy, in situ drug delivery, treatment of cancer or infectious agents, and vaccine therapy. However, to be successful, these technologies require an effective means for the delivery of the genetic information across cellular membranes.

**[0063]** It is well-known in the art that vaccinations can be used prophylactically for the prevention of infections as well as therapeutically for the treatment of ongoing conditions. Such infections or conditions can be but are not limited to viral infections. Thus, also disclosed are vectors, wherein the antigen-encoding nucleic acid is an antigen from a virus. The viral antigen is selected from the group of viruses consisting of Herpes simplex virus type-1, Herpes simplex virus type-2, Cytomegalovirus, Epstein-Barr virus, Varicella-zoster virus, Human herpesvirus 6, Human herpesvirus 7, Human herpesvirus 8, Variola virus, Vesicular stomatitis virus, Hepatitis A virus, Hepatitis B virus, Hepatitis C virus, Hepatitis D virus, Hepatitis E virus, SARS, Rhinovirus, Coronavirus, Influenza virus A, Influenza virus B, Measles virus, Polyomavirus, Human Papillomavirus, Respiratory syncytial virus, Adenovirus, Coxsackie virus, Dengue virus, Mumps virus, Poliovirus, Rabies virus, Rous sarcoma virus, Yellow fever virus, Ebola virus, Marburg virus, Lassa fever virus, Eastern Equine Encephalitis virus, Japanese Encephalitis virus, St. Louis Encephalitis virus, Murray Valley fever virus, West Nile virus, Rift Valley fever virus, Rotavirus A, Rotavirus B, Rotavirus C, Sindbis virus, Simian Immunodeficiency virus, Human T-lymphotropic virus type-1, Human T-lymphotropic virus type-2, Primate T-lymphotropic virus, Hantavirus, Rubella virus, Simian Immunodeficiency virus, Human Immunodeficiency virus type-1, Human Immunodeficiency virus type-2, and Simian Immunodeficiency virus (SW). Also disclosed are vectors, wherein the antigen-encoding nucleic

acid is SIV-GAG. The art is replete with examples of viral antigens whose sequences and methods of obtaining them are well known.

[0064] Vaccinations are also known for the prevention of bacterial infections. Additionally, antibiotics are well-known in the art for the treatment of various bacterial infections. Herein contemplated and disclosed are vectors, wherein the antigen-encoding nucleic acid is an antigen from a bacterium. The bacterial antigen is selected from the group consisting of *M. tuberculosis*, *M. bovis*, *M. bovis* strain BCG, BCG substrains, *M. avium*, *M. intracellulare*, *M. africanum*, *M. kansasii*, *M. marinum*, *M. ulcerans*, *M. avium* subspecies *paratuberculosis*, *Nocardia asteroides*, other *Nocardia* species, *Legionella pneumophila*, other *Legionella* species, *Salmonella typhi*, other *Salmonella* species, *Shigella* species, *Yersinia pestis*, *Pasteurella haemolytica*, *Pasteurella multocida*, other *Pasteurella* species, *Actinobacillus pleuropneumoniae*, *Listeria monocytogenes*, *Listeria ivanovii*, *Brucella abortus*, other *Brucella* species, *Cowdria ruminantium*, *Chlamydia pneumoniae*, *Chlamydia trachomatis*, *Chlamydia psittaci*, *Coxiella burnetti*, other *Rickettsial* species, *Ehrlichia* species, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Bacillus anthracis*, *Escherichia coli*, *Vibrio cholerae*, *Campylobacter* species, *Neisseria meningitidis*, *Neisseria gonorrhoea*, *Pseudomonas aeruginosa*, other *Pseudomonas* species, *Haemophilus influenzae*, *Haemophilus ducreyi*, other *Hemophilus* species, *Clostridium tetani*, other *Clostridium* species, *Yersinia enterocolitica*, and other *Yersinia* species. The art is replete with examples of bacterial antigens whose sequences and methods of obtaining them are well known.

[0065] Vaccinations are also known for the prevention of fungal infections. Additionally, antibiotics are well-known in the art for the treatment of various fungal infections. Herein contemplated and disclosed are vectors, wherein the antigen-encoding nucleic acid is an antigen from a fungus. The fungal antigen can be selected from the group consisting of *Candida albicans*, *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Aspergillus fumigatus*, *Coccidioides immitis*, *Paracoccidioides brasiliensis*, *Blastomyces dermatitidis*, *Pneumocystis carinii*, *Penicillium marneffi*, and *Alternaria alternata*.

[0066] The vectors of the disclosure are not limited to fungi, bacteria, and viruses. Also disclosed are vectors, wherein the antigen-encoding nucleic acid is an antigen from a parasite. The parasitic antigen can be selected from the group consisting of *Toxoplasma gondii*, *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium malariae*, other *Plasmodium* species, *Trypanosoma brucei*, *Trypanosoma cruzi*, *Leishmania major*, other *Leishmania* species, *Schistosoma mansoni*, other *Schistosoma* species, and *Entamoeba histolytica*. The art is replete with examples of parasitic antigens whose sequences and methods of obtaining them are well known.

[0067] There are instances wherein it is advantageous to administer the vector of the disclosure in a pharmaceutical composition that comprises other vaccines. Pharmaceutical compositions comprising multiple vaccines can be for therapeutic or prophylactic purposes. Examples of such compositions include the mumps, measles, rubella (MMR) vaccine, and vaccines against *M. tuberculosis*, *M. bovis*, *M. bovis* strain BCG, BCG substrains, *M. avium*, *M. intracellulare*, *M. africanum*, *M. kansasii*, *M. marinum*, *M. ulcerans*, *M. avium* subspecies *paratuberculosis*, *Nocardia asteroides*, other *Nocardia* species, *Legionella pneumophila*, other *Legionella*

species, *Salmonella typhi*, other *Salmonella* species, *Shigella* species, *Yersinia pestis*, *Pasteurella haemolytica*, *Pasteurella multocida*, other *Pasteurella* species, *Actinobacillus pleuropneumoniae*, *Listeria monocytogenes*, *Listeria ivanovii*, *Brucella abortus*, other *Brucella* species, *Cowdria ruminantium*, *Chlamydia pneumoniae*, *Chlamydia trachomatis*, *Chlamydia psittaci*, *Coxiella burnetti*, other *Rickettsial* species, *Ehrlichia* species, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Bacillus anthracis*, *Escherichia coli*, *Vibrio cholerae*, *Campylobacter* species, *Neisseria meningitidis*, *Neisseria gonorrhoea*, *Pseudomonas aeruginosa*, other *Pseudomonas* species, *Haemophilus influenzae*, *Haemophilus ducreyi*, other *Hemophilus* species, *Clostridium tetani*, other *Clostridium* species, *Yersinia enterocolitica*, and other *Yersinia* species. Specifically contemplated and disclosed are pharmaceutical compositions comprising the vector of the disclosure and one or more additional vaccines. Also disclosed are instances in which the vector comprises more than one antigen-encoding nucleic acid. In such a situation, the vector will produce each antigen encoded in the vector as a separate antigen.

[0068] There are instances in which a disclosed vector alone may not be suitable for a given purpose (e.g., a kit designed to screen potential drugs for the treatment of a condition, such kit intended for use in laboratories without the capabilities to transfect a cell-line with the vector). In such cases, cells previously transfected with the vector of the disclosure are needed. Thus, also disclosed are cells comprising the disclosed vectors.

[0069] In one embodiment, the antigen-encoding nucleic acid can encode a non-antigenic sequence of DNA. This sequence provides a functional copy of a disrupted, mutated, disregulated or deleted gene. Examples of nucleic acids encoding proteins that play a role in genetic disorders are known in the literature relating to genetic disorders. Methods of making these cells are described and exemplified herein and in the art.

[0070] The ability to detect the presence of a construct can be a desirable feature of any vector. As such, vectors often contain a marker to show that the construct of interest has been delivered to the subject (e.g., in a cell), and once delivered, is being expressed. A marker can take the form of a gene that is detectable when expressed. Thus, also disclosed are vectors further comprising a reporter gene. One example of a reporter gene is green fluorescence protein (GFP).

#### IV. Delivery of the Compositions to Cells

[0071] There are a number of compositions and methods which can be used to deliver nucleic acids to cells, either in vitro or in vivo. These methods and compositions can largely be broken down into two classes: viral based delivery systems and non-viral based delivery systems. For example, the nucleic acids can be delivered through a number of direct delivery systems, such as electroporation, lipofection, calcium phosphate precipitation, plasmids, viral vectors, viral nucleic acids, phage nucleic acids, phages, cosmids, or via transfer of genetic material in cells or carriers such as cationic liposomes. Appropriate means for transfection, including viral vectors, chemical transfectants, or physico-mechanical methods such as electroporation and direct diffusion of DNA, are described by, for example, Wolff, J. A., et al., *Science*, 247, 1465-1468, (1990); and Wolff, J. A. *Nature*, 352, 815-818, (1991). Such methods are well known in the art and readily

adaptable for use with the compositions and methods described herein. In certain cases, the methods will be modified to specifically function with large DNA molecules. Further, these methods can be used to target certain diseases and cell populations by using the targeting characteristics of the carrier.

#### V. Nucleic Acid Based Delivery Systems

**[0072]** Transfer vectors can be any nucleotide construction used to deliver genes into cells (e.g., a plasmid), or as part of a general strategy to deliver genes (e.g., as part of recombinant retrovirus or adenovirus; Ram et al. *Cancer Res.* 53:83-88, (1993)).

**[0073]** As used herein, plasmid or viral vectors are agents that transport nucleic acids into the cell without degradation, and include a promoter yielding expression of the gene in the cells into which it is delivered. In some embodiments the vectors are derived from either a virus or specifically a retrovirus. Viral vectors can include for example, for example, HTLV-1, HTLV-2, HTLV-3, HTLV-4, Adenovirus, Adeno-associated virus, Herpes virus, Vaccinia virus, Polio virus, AIDS virus, neuronal trophic virus, Sindbis and other RNA viruses, including these viruses with the HIV backbone. Also preferred are any viral families which share the properties of these viruses which make them suitable for use as vectors. Retroviruses include Murine Maloney Leukemia virus, MMLV, and retroviruses that express the desirable properties of MMLV as a vector. Retroviral vectors are able to carry a larger genetic payload (e.g., a transgene or marker gene) than other viral vectors, and for this reason are commonly used vectors. However, they are not as useful in non-proliferating cells. Adenovirus vectors are relatively stable and easy to work with, have high titers, can be delivered in aerosol formulation, and can transfect non-dividing cells. Pox viral vectors are large, have several sites for inserting genes, are thermostable, and can be stored at room temperature. A preferred embodiment is a viral vector which has been engineered so as to suppress the immune response of the host organism, elicited by the viral antigens. Preferred vectors of this type will carry coding regions for Interleukin 8 or 10.

**[0074]** Viral vectors can have higher transaction (ability to introduce genes) abilities than chemical or physical methods to introduce genes into cells. Typically, viral vectors contain nonstructural early genes, structural late genes, an RNA polymerase III transcript, inverted terminal repeats necessary for replication and encapsidation, and promoters to control the transcription and replication of the viral genome. When engineered as vectors, viruses typically have one or more of the early genes removed and a gene or gene/promoter cassette is inserted into the viral genome in place of the removed viral DNA. Constructs of this type can carry up to about 8 kb of foreign genetic material. The necessary functions of the removed early genes are typically supplied by cell lines which have been engineered to express the gene products of the early genes in trans.

#### VI. Retroviral Vectors

**[0075]** Primate T-lymphotropic viruses are retroviruses. A retrovirus is an animal virus belonging to the virus family of Retroviridae, including any types, subfamilies, genus, or tropisms. Retroviral vectors, in general, are described by Verma, I. M., Retroviral vectors for gene transfer, In *Microbiology*-1985, American Society for Microbiology, pp. 229-232,

Washington, (1985), which is incorporated by reference herein. Examples of methods for using retroviral vectors for gene therapy are described in U.S. Pat. Nos. 4,868,116 and 4,980,286; PCT applications WO 90/02806 and WO 89/07136; and Mulligan, (*Science* 260:926-932 (1993)); the teachings of which are incorporated herein by reference. Although the present primate T-lymphotropic virus vector is unique, the methods described for using other types of viral vectors can be useful in certain contexts. See for example U.S. Pat. No. 5,646,032, which is incorporated herein for its teaching of those methods.

**[0076]** A retrovirus is essentially a package which has packed into it nucleic acid cargo. The nucleic acid cargo carries with it a packaging signal, which ensures that the replicated daughter molecules will be efficiently packaged within the package coat. In addition to the package signal, there are a number of molecules which are needed in cis, for the replication, and packaging of the replicated virus. Typically a retroviral genome contains the gag, pol, and env genes which are involved in the making of the protein coat. It is the gag, pol, and env genes which are typically replaced by the foreign DNA that is to be transferred to the target cell. Retrovirus vectors typically contain a packaging signal for incorporation into the package coat, a sequence which signals the start of the gag transcription unit, elements necessary for reverse transcription, including a primer binding site to bind the tRNA primer of reverse transcription, terminal repeat sequences that guide the switch of RNA strands during DNA synthesis, a purine rich sequence 5' to the 3' LTR that serves as the priming site for the synthesis of the second strand of DNA synthesis, and specific sequences near the ends of the LTRs that enable the insertion of the DNA state of the retrovirus to insert into the host genome. The removal of the gag, pol, and env genes allows for large fragments of foreign sequence to be inserted into the viral genome, become reverse transcribed, and upon replication, be packaged into a new retroviral particle. This amount of nucleic acid is sufficient for the delivery of many genes depending on the size of each transcript. It is preferable to include either positive or negative selectable markers along with other genes in the insert.

**[0077]** Since the replication machinery and packaging proteins in most retroviral vectors have been removed (gag, pol, and env), the vectors are typically generated by placing them into a packaging cell line. A packaging cell line is a cell line which has been transfected or transformed with a retrovirus that contains the replication and packaging machinery, but lacks any packaging signal. When the vector carrying the DNA of choice is transfected into these cell lines, the vector containing the gene of interest is replicated and packaged into new retroviral particles, by the machinery provided in cis by the helper cell. The genomes for the machinery are not packaged because they lack the necessary signals.

**[0078]** A packaging cell line is a cell line which has been transfected or transformed with a retrovirus that contains the replication and packaging machinery, but lacks any packaging signal. When the vector carrying the DNA of choice is transfected into these cell lines, the vector containing the gene of interest is replicated and packaged into new retroviral particles, by the machinery provided in cis by the helper cell. The genomes for the machinery are not packaged because they lack the necessary signals.

**[0079]** It is also understood that the pX region can be used to construct a vector. The pX region is located between the end of env and the beginning of Tax and contains small ORFs,

hence this is another good region for insertion of foreign DNA in an HTLV genome based vector.

**[0080]** Disclosed are methods of detecting the expression of the disclosed vectors comprising using a first antibody to the antigen to measure protein expression in a quantitative or qualitative way, and further comprising detecting the first antibody directly via a colorimetric measurement produced through the use of a substrate and a conjugated antibody or indirectly via a first antibody to the antigen, which in turn is bound by a second antibody that is conjugated and will result in a colorimetric measurement when combined with a substrate.

**[0081]** Also disclosed are methods wherein the antigen is detected by placing an aliquot of the disclosed vector in a lane on a gel and probing the gel for the antigen.

**[0082]** Some methods are methods of detecting the expression of the disclosed vector using a fluorescently labeled first antibody specific for the antigen and visualizing the antigen using a flow cytometer, fluorescence microscope, or chemiluminescence. In some embodiments, the first antibody is not fluorescently labeled, but a target for a second antibody with a fluorescent label.

**[0083]** Also disclosed are methods of detecting the expression of a disclosed vector comprising using cytolytic killing assay to assess activity, and methods of detecting the vector that further include obtaining a sample from a subject comprising a tissue biopsy or removal of blood or bone marrow.

#### VII. Non-Nucleic Acid Based Systems

**[0084]** The disclosed compositions can be delivered to the target cells in a variety of ways. For example, the compositions can be delivered through electroporation, or through lipofection, or through calcium phosphate precipitation. The delivery mechanism chosen will depend in part on the type of cell targeted and whether the delivery is occurring for example in vivo or in vitro. In the methods described above which include the administration and uptake of exogenous DNA into the cells of a subject (e.g., gene transduction or transfection), delivery of the compositions to cells can be via a variety of mechanisms. As one example, delivery can be via a liposome, using commercially available liposome preparations such as LIPOFECTIN, LIPOFECTAMINE (GIBCO-BRL, Inc., Gaithersburg, Md.), SUPERFECT (Qiagen, Inc. Hilden, Germany) and TRANSFECTAM (Promega Biotec, Inc., Madison, Wis.), as well as other liposomes developed according to procedures standard in the art. In addition, the disclosed nucleic acid or vector can be delivered in vivo by electroporation, the technology for which is available from Genetronics, Inc. (San Diego, Calif.) as well as by means of a SONOPORATION machine (ImaRx Pharmaceutical Corp., Tucson, Ariz.).

**[0085]** The materials may be in solution or suspension (for example, incorporated into microparticles, liposomes, or cells). These may be targeted to a particular cell type via antibodies, receptors, or receptor ligands. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Senter et al., *Bioconjugate Chem.* 2:447-451, (1991); Bagshawe, K. D., *Br. J. Cancer* 60:275-281 (1989); Bagshawe et al., *Br. J. Cancer* 58:700-703 (1988); Senter et al., *Bioconjugate Chem.* 4:3-9 (1993); Battelli et al., *Cancer Immunol. Immunother.* 35:421-425 (1992); Pietersz and McKenzie, *Immunolog. Reviews* 129:57-80 (1992); and Roffler et al., *Biochem. Pharmacol.* 42:2062-2065 (1991)). These techniques can be used for a variety of

other specific cell types. Vehicles such as "stealth" and other antibody conjugated liposomes (including lipid mediated drug targeting to colonic carcinoma), receptor mediated targeting of DNA through cell specific ligands, lymphocyte directed tumor targeting, and highly specific therapeutic retroviral targeting of murine glioma cells in vivo. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Hughes et al., *Cancer Research* 49:6214-6220 (1989); and Litzinger and Huang, *Biochimica et Biophysica Acta* 1104:179-187 (1992)). In general, receptors are involved in pathways of endocytosis, either constitutive or ligand induced. These receptors cluster in clathrin-coated pits, enter the cell via clathrin-coated vesicles, pass through an acidified endosome in which the receptors are sorted, and then either recycle to the cell surface, become stored intracellularly, or are degraded in lysosomes. The internalization pathways serve a variety of functions, such as nutrient uptake, removal of activated proteins, clearance of macromolecules, opportunistic entry of viruses and toxins, dissociation and degradation of ligand, and receptor-level regulation. Many receptors follow more than one intracellular pathway, depending on the cell type, receptor concentration, type of ligand, ligand valency, and ligand concentration. Molecular and cellular mechanisms of receptor-mediated endocytosis has been reviewed (Brown and Greene, *DNA and Cell Biology* 10(6):399-409 (1991)).

**[0086]** Nucleic acids that are delivered to cells which are to be integrated into the host cell genome, typically contain integration sequences. These sequences are often viral related sequences, particularly when viral based systems are used. These viral integration systems can also be incorporated into nucleic acids which are to be delivered using a non-nucleic acid based system of deliver, such as a liposome, so that the nucleic acid contained in the delivery system can be come integrated into the host genome.

**[0087]** Other general techniques for integration into the host genome include, for example, systems designed to promote homologous recombination with the host genome. These systems typically rely on sequence flanking the nucleic acid to be expressed that has enough homology with a target sequence within the host cell genome that recombination between the vector nucleic acid and the target nucleic acid takes place, causing the delivered nucleic acid to be integrated into the host genome. These systems and the methods necessary to promote homologous recombination are known to those of skill in the art.

#### VIII. In Vivo/Ex Vivo Methods

**[0088]** As described above, the compositions can be administered in a pharmaceutically acceptable carrier and can be delivered to the subject's cells in vivo and/or ex vivo by a variety of mechanisms well known in the art (e.g., uptake of naked DNA, liposome fusion, intramuscular injection of DNA via a gene gun, endocytosis and the like).

**[0089]** If ex vivo methods are employed, cells or tissues can be removed and maintained outside the body according to standard protocols well known in the art. The compositions can be introduced into the cells via any gene transfer mechanism, such as, for example, calcium phosphate mediated gene delivery, electroporation, microinjection or proteoliposomes. The transduced cells can then be infused (e.g., in a pharmaceutically acceptable carrier) or homotopically transplanted back into the subject per standard methods for the cell or

tissue type. Standard methods are known for transplantation or infusion of various cells into a subject.

#### IX. Expression Systems

**[0090]** The nucleic acids that are delivered to cells typically contain expression controlling systems. For example, the inserted genes in viral and retroviral systems usually contain promoters, and/or enhancers to help control the expression of the desired gene product. A promoter is generally a sequence or sequences of DNA that function when in a relatively fixed location in regard to the transcription start site. A promoter contains core elements required for basic interaction of RNA polymerase and transcription factors, and may contain upstream elements and response elements.

#### X. Viral Promoters and Enhancers

**[0091]** Preferred promoters controlling transcription from vectors in mammalian host cells may be obtained from various sources, for example, the genomes of viruses such as: polyoma, Simian Virus 40 (SV40), adenovirus, retroviruses, hepatitis-B virus and most preferably cytomegalovirus, or from heterologous mammalian promoters, e.g. beta actin promoter. The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment which also contains the SV40 viral origin of replication (Fiers et al., *Nature*, 273:113 (1978)). The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment (Greenway et al., *Gene* 18:355-360 (1982)). Of course, promoters from the host cell or related species also are useful herein. Such preferred promoters are in the LTRs of HTLV.

**[0092]** Enhancer generally refers to a sequence of DNA that functions at no fixed distance from the transcription start site and can be either 5' (Laimins et al., *Proc. Natl. Acad. Sci.* 78:993 (1981)) or 3' (Lusky et al., *Mol. Cell. Bio.* 3:1108 (1983)) to the transcription unit. Furthermore, enhancers can be within an intron (Banerji et al., *Cell* 33:729 (1983)), as well as within the coding sequence itself (Osborne et al., *Mol. Cell. Bio.* 4:1293 (1984)). They are usually between 10 and 300 by in length, and they function in cis. Enhancers function to increase transcription from nearby promoters. Enhancers also often contain response elements that mediate the regulation of transcription. Promoters can also contain response elements that mediate the regulation of transcription. Enhancers often determine the regulation of expression of a gene. While many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, fetoprotein and insulin), typically one will use an enhancer from a eukaryotic cell virus for general expression. Preferred examples are the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

**[0093]** The promoter and/or enhancer may be specifically activated, for instance by light or specific chemical events which trigger their function. Systems can be regulated by reagents such as tetracycline and dexamethasone. There are also ways to enhance viral vector gene expression by exposure to irradiation, such as gamma irradiation, or alkylating chemotherapy drugs.

**[0094]** In certain embodiments the promoter and/or enhancer region can act as a constitutive promoter and/or enhancer to maximize expression of the region of the tran-

scription unit to be transcribed. In certain constructs the promoter and/or enhancer region be active in all eukaryotic cell types, even if it is only expressed in a particular type of cell at a particular time. A preferred promoter of this type is the CMV promoter (650 bases). Other preferred promoters are SV40 promoters, cytomegalovirus (full length promoter), and retroviral vector LTR.

**[0095]** It has been shown that all specific regulatory elements can be cloned and used to construct expression vectors that are selectively expressed in specific cell types such as melanoma cells. The glial fibrillary acetic protein (GFAP) promoter has been used to selectively express genes in cells of glial origin.

**[0096]** Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human or nucleated cells) may also contain sequences necessary for the termination of transcription which may affect mRNA expression. These regions are transcribed as polyadenylated segments in the untranslated portion of the mRNA encoding tissue factor protein. The 3' untranslated regions also include transcription termination sites. It is preferred that the transcription unit also contains a polyadenylation region. One benefit of this region is that it increases the likelihood that the transcribed unit will be processed and transported like mRNA. The identification and use of polyadenylation signals in expression constructs is well established. It is preferred that homologous polyadenylation signals be used in the transgene constructs. In certain transcription units, the polyadenylation region is derived from the SV40 early polyadenylation signal and consists of about 400 bases. It is also preferred that the transcribed units contain other standard sequences alone or in combination with the above sequences improve expression from, or stability of, the construct.

#### XI. Markers

**[0097]** The viral vectors can include nucleic acid sequence encoding a marker product. This marker product is used to determine if the gene has been delivered to the cell and once delivered is being expressed. Preferred marker genes are the *E. Coli lacZ* gene, which encodes  $\beta$ -galactosidase, and green fluorescent protein.

**[0098]** In some embodiments the marker is a selectable marker. Examples of suitable selectable markers for mammalian cells are dihydrofolate reductase (DHFR), thymidine kinase, neomycin, neomycin analog G418, hygromycin, and puromycin. When such selectable markers are successfully transferred into a mammalian host cell, the transformed mammalian host cell can survive if placed under selective pressure. There are two widely used distinct categories of selective regimes. The first category is based on a cell's metabolism and the use of a mutant cell line which lacks the ability to grow independent of a supplemented media. Two examples are: CHO DHFR-cells and mouse LTK-cells. These cells lack the ability to grow without the addition of such nutrients as thymidine or hypoxanthine. Because these cells lack certain genes necessary for a complete nucleotide synthesis pathway, they cannot survive unless the missing nucleotides are provided in a supplemented media. An alternative to supplementing the media is to introduce an intact DHFR or TK gene into cells lacking the respective genes, thus altering their growth requirements. Individual cells which were not transformed with the DHFR or TK gene will not be capable of survival in non-supplemented media.

**[0099]** The second category is dominant selection which refers to a selection scheme used in any cell type and does not require the use of a mutant cell line. These schemes typically use a drug to arrest growth of a host cell. Those cells that have a novel gene would express a protein conveying drug resistance and would survive the selection. Examples of such dominant selection use the drugs neomycin, (Southern & Berg, *J. Molec. Appl. Genet.* 1:327 (1982)), mycophenolic acid, (Mulligan & Berg *Science* 209:1422 (1980)) or hygromycin, (Sugden et al., *Mol. Cell. Biol.* 5:410-413 (1985)). The three examples employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug G418 or neomycin (geneticin), xgpt (mycophenolic acid) or hygromycin, respectively. Others include the neomycin analog G418 and puromycin.

## XII. Sequence Similarities

**[0100]** It is understood that as discussed herein the use of the terms homology and identity mean the same thing as similarity. Thus, for example, if the use of the word homology is used between two non-natural sequences it is understood that this is not necessarily indicating an evolutionary relationship between these two sequences, but rather is looking at the similarity or relatedness between their nucleic acid sequences. Many of the methods for determining homology between two evolutionarily related molecules are routinely applied to any two or more nucleic acids or proteins for the purpose of measuring sequence similarity regardless of whether they are evolutionarily related or not.

**[0101]** In general, it is understood that one way to define any known variants and derivatives or those that might arise, of the disclosed genes and proteins herein, is through defining the variants and derivatives in terms of homology to specific known sequences. This identity of particular sequences disclosed herein is also discussed elsewhere herein. In general, variants of genes and proteins herein disclosed typically have at least, about 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 percent homology to the stated sequence or the native sequence. Those of skill in the art readily understand how to determine the homology of two proteins or nucleic acids, such as genes. For example, the homology can be calculated after aligning the two sequences so that the homology is at its highest level.

**[0102]** Another method of calculating homology can be performed by published algorithms. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith & Waterman *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by inspection.

**[0103]** The same types of homology can be obtained for nucleic acids by for example the algorithms disclosed in Zuker *Science* 244:48-52, 1989, Jaeger et al. *Proc. Natl. Acad. Sci. USA* 86:7706-7710, 1989, and Jaeger et al. *Methods Enzymol.* 183:281-306, 1989, which are herein incorporated by reference for at least material related to nucleic acid alignment. It is understood that any of the methods typically can be used and that in certain instances the results of these

various methods may differ, but the skilled artisan understands if identity is found with at least one of these methods, the sequences would be said to have the stated identity, and be disclosed herein.

**[0104]** For example, as used herein, a sequence recited as having a particular percent homology to another sequence refers to sequences that have the recited homology as calculated by any one or more of the calculation methods described above. For example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using the Zuker calculation method even if the first sequence does not have 80 percent homology to the second sequence as calculated by any of the other calculation methods. As another example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using both the Zuker calculation method and the Pearson and Lipman calculation method even if the first sequence does not have 80 percent homology to the second sequence as calculated by the Smith & Waterman calculation method, the Needleman & Wunsch calculation method, the Jaeger calculation methods, or any of the other calculation methods. As yet another example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using each of calculation methods (although, in practice, the different calculation methods will often result in different calculated homology percentages).

## XIII. Nucleic Acids

**[0105]** There are a variety of molecules disclosed herein that are nucleic acid based, including for example the nucleic acids that encode HTLV-3 or HTLV-4 (e.g., SEQ ID NOs: 36, 53, and 81). The disclosed nucleic acids are made up of, for example, nucleotides, nucleotide analogs, or nucleotide substitutes. Non-limiting examples of these and other molecules are discussed herein. It is understood that for example, when a vector is expressed in a cell, the expressed mRNA will typically be made up of A, C, G, and U. Likewise, it is understood that if, for example, an antisense molecule is introduced into a cell or cell environment through for example exogenous delivery, it is advantageous that the antisense molecule be made up of nucleotide analogs that reduce the degradation of the antisense molecule in the cellular environment.

## XIV. Nucleotides and Related Molecules

**[0106]** A nucleotide is a molecule that contains a base moiety, a sugar moiety and a phosphate moiety. Nucleotides can be linked together through their phosphate moieties and sugar moieties creating an internucleoside linkage. The base moiety of a nucleotide can be adenin-9-yl (A), cytosin-1-yl (C), guanin-9-yl (G), uracil-1-yl (U), and thymine-1-yl (T). The sugar moiety of a nucleotide is a ribose or a deoxyribose. The phosphate moiety of a nucleotide is pentavalent phosphate. A non-limiting example of a nucleotide would be 3'-AMP (3'-adenosine monophosphate) or 5'-GMP (5'-guanosine monophosphate). There are many varieties of these types of molecules available in the art and available herein.

**[0107]** A nucleotide analog is a nucleotide which contains some type of modification to either the base, sugar, or phos-

phate moieties. Modifications to nucleotides are well known in the art and would include for example, 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, and 2-aminoadenine as well as modifications at the sugar or phosphate moieties. There are many varieties of these types of molecules available in the art and available herein.

**[0108]** Nucleotide substitutes are molecules having similar functional properties to nucleotides, but which do not contain a phosphate moiety, such as peptide nucleic acid (PNA). Nucleotide substitutes are molecules that will recognize nucleic acids in a Watson-Crick or Hoogsteen manner, but which are linked together through a moiety other than a phosphate moiety. Nucleotide substitutes are able to conform to a double helix type structure when interacting with the appropriate target nucleic acid. There are many varieties of these types of molecules available in the art and available herein.

**[0109]** It is also possible to link other types of molecules (conjugates) to nucleotides or nucleotide analogs to enhance for example, cellular uptake. Conjugates can be chemically linked to the nucleotide or nucleotide analogs. Such conjugates include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86:6553-6556). There are many varieties of these types of molecules available in the art and available herein.

**[0110]** A Watson-Crick interaction is at least one interaction with the Watson-Crick face of a nucleotide, nucleotide analog, or nucleotide substitute. The Watson-Crick face of a nucleotide, nucleotide analog, or nucleotide substitute includes the C2, N1, and C6 positions of a purine based nucleotide, nucleotide analog, or nucleotide substitute and the C2, N3, C4 positions of a pyrimidine based nucleotide, nucleotide analog, or nucleotide substitute.

**[0111]** A Hoogsteen interaction is the interaction that takes place on the Hoogsteen face of a nucleotide or nucleotide analog, which is exposed in the major groove of duplex DNA. The Hoogsteen face includes the N7 position and reactive groups (NH<sub>2</sub> or O) at the C6 position of purine nucleotides.

#### XV. Sequences

**[0112]** There are a variety of sequences related to the protein molecules, for example the protein coding regions gag, pol, env, tax, rex, and protease (pro) genes and noncoding regions such as the LTR of HTLV-3 and HTLV-4, or any of the nucleic acids disclosed herein for making HTLV-3 or HTLV-4, all of which are encoded by nucleic acids or are nucleic acids. The sequences for the human analogs of these genes, as well as other analogs, and alleles of these genes, and splice variants and other types of variants, are available in a variety of protein and gene databases, including GenBank. Those sequences available at the time of filing this application at GenBank are herein incorporated by reference in their entireties as well as for individual subsequences contained therein. GenBank can be accessed at <http://www.ncbi.nih.gov/entrez/query.fcgi>. Those of skill in the art understand how to resolve sequence discrepancies and differences and to adjust the compositions and methods relating to a particular sequence to other related sequences. Primers and/or probes can be designed for any given sequence given the information disclosed herein and known in the art.

#### XVI. Primers and Probes

**[0113]** Disclosed are compositions including primers and probes, which are capable of interacting with the disclosed

nucleic acids, such as the HTLV-3 or HTLV-4 as disclosed herein. In certain embodiments the primers are used to support nucleic acid (DNA, RNA, etc.) amplification reactions. Thus, for example, disclosed herein are primers wherein the primer comprises SEQ ID NOs: 7 and 8, SEQ ID NOs: 11 and 12, SEQ ID NOs: 15 and 16, SEQ ID NOs: 23 and 24, SEQ ID NOs: 27 and 28, SEQ ID NOs: 31 and 32, SEQ ID NOs: 69 and 70, SEQ ID NOs: 73 and 74, SEQ ID NOs: 77 and 78, SEQ ID NOs: 9 and 10, SEQ ID NOs: 13 and 14, SEQ ID NOs: 17 and 18, SEQ ID NOs: 25 and 26, SEQ ID NOs: 29 and 30, SEQ ID NOs: 33 and 34, SEQ ID NOs: 64 and 65, SEQ ID NOs: 71 and 72, SEQ ID NOs: 75 and 76, and SEQ ID NOs: 79 and 80. Typically the primers will be capable of being extended in a sequence specific manner. Extension of a primer in a sequence specific manner includes any methods wherein the sequence and/or composition of the nucleic acid molecule to which the primer is hybridized or otherwise associated directs or influences the composition or sequence of the product produced by the extension of the primer. Extension of the primer in a sequence specific manner therefore includes, but is not limited to, PCR, DNA sequencing, DNA extension, DNA polymerization, RNA transcription, or reverse transcription. Techniques and conditions that amplify the primer in a sequence specific manner are preferred. In certain embodiments the primers are used for the DNA amplification reactions, such as PCR or direct sequencing. Thus, herein are disclosed primer pairs used in conjunction with a second nested set of primers pairs. For example, disclosed herein are PCR amplification methods comprising a first primer pair and a second primer pair, wherein the second primer pair is internal to the first primer pair and wherein the first primer pair is selected from the group consisting of SEQ ID NOs: 7 and 8, SEQ ID NOs: 11 and 12, SEQ ID NOs: 15 and 16, SEQ ID NOs: 23 and 24, SEQ ID NOs: 27 and 28, SEQ ID NOs: 31 and 32, SEQ ID NOs: 69 and 70, SEQ ID NOs: 73 and 74, and SEQ ID NOs: 77 and 78, wherein the second set of primers is selected from the group consisting of SEQ ID NOs: 9 and 10, SEQ ID NOs: 13 and 14, SEQ ID NOs: 17 and 18, SEQ ID NOs: 25 and 26, SEQ ID NOs: 29 and 30, SEQ ID NOs: 33 and 34, SEQ ID NOs: 71 and 72, SEQ ID NOs: 75 and 76, and SEQ ID NOs: 79 and 80. It is understood that in certain embodiments the primers can also be extended using non-enzymatic techniques, where for example, the nucleotides or oligonucleotides used to extend the primer are modified such that they will chemically react to extend the primer in a sequence specific manner. Typically, the disclosed primers hybridize with the disclosed nucleic acids or region of the nucleic acids or they hybridize with the complement of the nucleic acids or complement of a region of the nucleic acids.

#### XVII. Functional Nucleic Acids

**[0114]** Functional nucleic acids are nucleic acid molecules that have a specific function, such as binding a target molecule or catalyzing a specific reaction. Functional nucleic acid molecules can be divided into the following categories, which are not meant to be limiting. For example; functional nucleic acids include antisense molecules, aptamers, ribozymes, triplex forming molecules, and external guide sequences. The functional nucleic acid molecules can act as effectors, inhibitors, modulators, and stimulators of a specific activity possessed by a target molecule, or the functional nucleic acid molecules can possess a de novo activity independent of any other molecules.

Functional nucleic acid molecules can interact with any macromolecule, such as DNA, RNA, polypeptides, or carbohydrate chains. Thus, functional nucleic acids can interact with the mRNA of any of the disclosed nucleic acids, such as the pol, tax, env, gag, rex and pro genes and non-coding regions such as the LTR of HTLV-3 and HTLV-4, or the nucleic acids used for the generation of HTLV-3 and HTLV-4, or the genomic DNA of any of the disclosed viruses, such as HTLV-3 and HTLV-4, or they can interact with the polypeptide encoded by any of the disclosed nucleic acids, such as pol, tax, rex, env, gag, or pro genes of HTLV-3 and HTLV-4, or the nucleic acids used for the generation of pol, tax, rex, env, gag, or LTR proteins of HTLV-3 and HTLV-4. Often functional nucleic acids are designed to interact with other nucleic acids based on sequence homology between the target molecule and the functional nucleic acid molecule. In other situations, the specific recognition between the functional nucleic acid molecule and the target molecule is not based on sequence homology between the functional nucleic acid molecule and the target molecule, but rather is based on the formation of tertiary structure that allows specific recognition to take place.

#### XVIII. Protein Variants

**[0115]** As discussed herein, there are numerous disclosed variants of the HTLV-3 proteins encoded herein, such as gag (SEQ ID NO: 40), pol (SEQ ID NO: 44), env (SEQ ID NO: 37), tax (SEQ ID NO: 50), rex (SEQ ID NO: 48), protease (SEQ ID NO: 46), and non-coding regions such as the LTR, and HTLV-4 proteins encoded herein, such as gag, pol (SEQ ID NO: 57), env (SEQ ID NO: 54), tax (SEQ ID NO: 62), rex (SEQ ID NO: 60), protease (SEQ ID NO: 58) and non-coding regions such as the LTR. In addition, to the known functional HTLV-3 and HTLV-4 strain variants there are derivatives of the HTLV-3 and HTLV-4 gag, pol, tax, rex, and env, LTR proteins that also function in the disclosed methods and compositions. Protein variants and derivatives are well understood to those of skill in the art and in can involve amino acid sequence modifications. For example, amino acid sequence modifications typically fall into one or more of three classes: substitutional, insertional or deletional variants. Insertions include amino and/or carboxyl terminal fusions as well as intrasequence insertions of single or multiple amino acid residues. Insertions ordinarily will be smaller insertions than those of amino or carboxyl terminal fusions, for example, on the order of one to four residues. Immunogenic fusion protein derivatives, such as those described in the examples, are made by fusing a polypeptide sufficiently large to confer immunogenicity to the target sequence by cross-linking in vitro or by recombinant cell culture transformed with DNA encoding the fusion. Deletions are characterized by the removal of one or more amino acid residues from the protein sequence. Typically, no more than about from 2 to 6 residues are deleted at any one site within the protein molecule. These variants ordinarily are prepared by site specific mutagenesis of nucleotides in the DNA encoding the protein, thereby producing DNA encoding the variant, and thereafter expressing the DNA in recombinant cell culture. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, for example M13 primer mutagenesis and PCR mutagenesis. Amino acid substitutions are typically of single residues, but can occur at a number of different locations at once; insertions usually will be on the order of about from 1 to 10 amino acid residues; and deletions

will range about from 1 to 30 residues. Deletions or insertions preferably are made in adjacent pairs, e.g., a deletion of 2 residues or insertion of 2 residues. Substitutions, deletions, insertions or any combination thereof may be combined to arrive at a final construct. The mutations must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. Substitutional variants are those in which at least one residue has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the following Tables 1 and 2 and are referred to as conservative substitutions.

TABLE 1

Amino Acid Abbreviations	
Amino Acid	Abbreviation
alanine	Ala A
alloseleucine	Alle
arginine	Arg R
asparagine	Asn N
aspartic acid	Asp D
cysteine	Cys C
glutamic acid	Glu E
glutamine	Gln K
glycine	Gly G
histidine	His H
isoleucine	Ile I
leucine	Leu L
lysine	Lys K
phenylalanine	Phe F
proline	Pro P
pyroglutamic acid	Glu
serine	Ser S
threonine	Thr T
tyrosine	Tyr Y
tryptophan	Trp W
valine	Val V

TABLE 2

Amino Acid Substitutions Original Residue & Exemplary Conservative Substitutions (others are known in the art)
Ala, ser
Arg, lys, gln
Asn, gln, his
Asp, glu
Cys, ser
Gln, asn, lys
Glu, asp
Gly, pro
His, asn, gln
Ile, leu, val
Leu, ile, val
Lys, arg, gln,
Met, Leu, ile
Phe, met, leu, tyr
Ser, thr
Thr, ser
Trp, tyr
Tyr, trp, phe
Val, ile, leu

**[0116]** Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative than those in Table 2, e.g., selecting residues that differ more significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the sub-

stitution, for example as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site or (c) the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in the protein properties will be those in which (a) a hydrophilic residue, e.g. seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine, in this case, (e) by increasing the number of sites for sulfation and/or glycosylation.

[0117] For example, the replacement of one amino acid residue with another that is biologically and/or chemically similar is known to those skilled in the art as a conservative substitution. For example, a conservative substitution would be replacing one hydrophobic residue with another, or one polar residue with another. The substitutions include combinations such as, for example, Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe, Tyr. Such conservatively substituted variations of each explicitly disclosed sequence are included within the mosaic polypeptides provided herein.

[0118] Substitutional or deletional mutagenesis can be employed to insert sites for N-glycosylation (Asn-X-Thr/Ser) or O-glycosylation (Ser or Thr). Deletions of cysteine or other labile residues also may be desirable. Deletions or substitutions of potential proteolysis sites, e.g. Arg, is accomplished for example by deleting one of the basic residues or substituting one by glutaminyl or histidyl residues.

[0119] Certain post-translational derivatizations are the result of the action of recombinant host cells on the expressed polypeptide. Glutaminyl and asparaginyl residues are frequently post-translationally deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Other post-translational modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the o-amino groups of lysine, arginine, and histidine side chains (T.E. Creighton, *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco pp 79-86 [1983]), acetylation of the N-terminal amine and, in some instances, amidation of the C-terminal carboxyl.

[0120] It is understood that one way to define the variants and derivatives of the disclosed proteins herein is through defining the variants and derivatives in terms of homology/identity to specific known sequences. For example, SEQ ID NO: 1 sets forth a particular sequence of HTLV-3 pol protein and SEQ ID NO: 2 sets forth a particular sequence of a HTLV-4 pol protein. Specifically disclosed are variants of these and other proteins herein disclosed which have at least, 70% or 75% or 80% or 85% or 90% or 95% homology or any amount of homology in between to the stated sequence. Those of skill in the art readily understand how to determine the homology of two proteins. For example, the homology can be calculated after aligning the two sequences so that the homology is at its highest level.

[0121] Another way of calculating homology can be performed by published algorithms. Optimal alignment of sequences for comparison may be conducted by the local

homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by inspection.

[0122] The same types of homology can be obtained for nucleic acids by for example the algorithms disclosed in Zuker, M. *Science* 244:48-52, 1989, Jaeger et al. *Proc. Natl. Acad. Sci. USA* 86:7706-7710, 1989, Jaeger et al. *Methods Enzymol.* 183:281-306, 1989 which are herein incorporated by reference for at least material related to nucleic acid alignment.

[0123] It is understood that the description of conservative mutations and homology can be combined together in any combination, such as embodiments that have at least 70%, 80%, 85%, 90%, 92%, 95%, 97% or more homology to a particular sequence wherein the variants are conservative mutations.

[0124] As this specification discusses various proteins and protein sequences it is understood that the nucleic acids that can encode those protein sequences are also disclosed. This would include all degenerate sequences related to a specific protein sequence, e.g. all nucleic acids having a sequence that encodes one particular protein sequence as well as all nucleic acids, including degenerate nucleic acids, encoding the disclosed variants and derivatives of the protein sequences. Thus, while each particular nucleic acid sequence may not be written out herein, it is understood that each and every sequence is in fact disclosed and described herein through the disclosed protein sequence. For example, one of the many nucleic acid sequences that can encode the protein sequence set forth in SEQ ID NO: 44 is set forth in SEQ ID NO: 1. In addition, for example, disclosed are conservative derivatives of SEQ ID NO: 44.

[0125] It is understood that there are numerous amino acid and peptide analogs which can be incorporated into the disclosed compositions. For example, there are numerous D amino acids or amino acids which have a different functional substituent than the amino acids shown in Table 1 and Table 2. The opposite stereo isomers of naturally occurring peptides are disclosed, as well as the stereo isomers of peptide analogs. These amino acids can readily be incorporated into polypeptide chains by charging tRNA molecules with the amino acid of choice and engineering genetic constructs that utilize, for example, amber codons, to insert the analog amino acid into a peptide chain in a site specific way (Thorson et al., *Methods in Molec. Biol.* 77:43-73 (1991), Zoller, *Current Opinion in Biotechnology*, 3:348-354 (1992); Ibba, *Biotechnology & Genetic Engineering Reviews* 13:197-216 (1995), Cahill et al., *TIBS*, 14(10):400-403 (1989); Benner, *TIB Tech*, 12:158-163 (1994); Ibba and Hennecke, *Bio/technology*, 12:678-682 (1994) all of which are herein incorporated by reference at least for material related to amino acid analogs).

[0126] Molecules can be produced that resemble peptides, but which are not connected via a natural peptide linkage. For example, linkages for amino acids or amino acid analogs can include  $\text{CH}_2\text{NH}-$ ,  $-\text{CH}_2\text{S}-$ ,  $-\text{CH}_2-\text{CH}_2-\text{CH}=\text{CH}-$  (cis and trans),  $-\text{COCH}_2-\text{CH}(\text{OH})\text{CH}_2-$ , and  $-\text{CHH}_2\text{SO}-$ . (These and others can be found in Spatola in *Chemistry and Biochemistry of Amino Acids, Peptides, and*

*Proteins*, B. Weinstein, eds., Marcel Dekker, New York, p. 267 (1983); Spatola, *Vega Data* (March 1983), Vol. 1, Issue 3, *Peptide Backbone Modifications* (general review); Morley, *Trends Pharm Sci* (1980) pp. 463-468; Hudson et al., *Int J Pept Prot Res* 14:177-185 (1979) ( $-\text{CH}_2\text{NH}-$ ,  $\text{CH}_2\text{CH}_2-$ ); Spatola et al. *Life Sci* 38:1243-1249 (1986) ( $-\text{CH}_2\text{H}_2-\text{S}$ ); Hann *J. Chem. Soc Perkin Trans.* 1307-314 (1982) ( $-\text{CH}-\text{CH}-$ , cis and trans); Almquist et al. *J. Med. Chem.* 23:1392-1398 (1980) ( $-\text{COCH}_2-$ ); Jennings-White et al. *Tetrahedron Lett* 23:2533 (1982) ( $-\text{COCH}_2-$ ); Szelke et al. *European Appln*, EP 45665 CA (1982): 97:39405 (1982) ( $-\text{CH}(\text{OH})\text{CH}_2-$ ); Holladay et al. *Tetrahedron. Lett* 24:4401-4404 (1983) ( $-\text{C}(\text{OH})\text{CH}_2-$ ); and Hruby *Life Sci* 31:189-199 (1982) ( $-\text{CH}_2-\text{S}-$ ); each of which is incorporated herein by reference. A particularly preferred non-peptide linkage is  $-\text{CH}_2\text{NH}-$ . It is understood that peptide analogs can have more than one atom between the bond atoms, such as b-alanine, g-aminobutyric acid, and the like.

**[0127]** Amino acid analogs and analogs and peptide analogs often have enhanced or desirable properties, such as, more economical production, greater chemical stability, enhanced pharmacological properties (half-life, absorption, potency, efficacy, etc.), altered specificity (e.g., a broad-spectrum of biological activities), reduced antigenicity, and others.

**[0128]** D-amino acids can be used to generate more stable peptides, because D amino acids are not recognized by peptidases and such. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) can be used to generate more stable peptides. Cysteine residues can be used to cyclize or attach two or more peptides together. This can be beneficial to constrain peptides into particular conformations (Rizo and Gierasch *Ann. Rev. Biochem.* 61:387 (1992), incorporated herein by reference).

#### XIX. Pharmaceutical Carriers/Delivery of Pharmaceutical Products

**[0129]** As described above, the compositions can be administered in vivo in a pharmaceutically acceptable carrier. By "pharmaceutically acceptable" is meant a material that is not biologically or otherwise undesirable, e.g., the material may be administered to a subject, along with the nucleic acid or vector, without causing any undesirable biological effects or interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained. The carrier would naturally be selected to minimize any degradation of the active ingredient and to minimize any adverse side effects in the subject, as would be well known to one of skill in the art.

**[0130]** The compositions may be administered orally, parenterally (e.g., intravenously), by intramuscular injection, by intraperitoneal injection, transdermally, extracorporeally, topically or the like, including topical intranasal administration or administration by inhalant. As used herein, "topical intranasal administration" means delivery of the compositions into the nose and nasal passages through one or both of the nares and can comprise delivery by a spraying mechanism or droplet mechanism, or through aerosolization of the nucleic acid or vector. Administration of the compositions by inhalant can be through the nose or mouth via delivery by a spraying or droplet mechanism. Delivery can also be directly to any area of the respiratory system (e.g., lungs) via intubation. The exact amount of the compositions required will vary

from subject to subject, depending on the species, age, weight and general condition of the subject, the severity of the allergic disorder being treated, the particular nucleic acid or vector used, its mode of administration and the like. Thus, it is not possible to specify an exact amount for every composition. However, an appropriate amount can be determined by one of ordinary skill in the art using only routine experimentation given the teachings herein.

**[0131]** Parenteral administration of the composition, if used, is generally characterized by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution of suspension in liquid prior to injection, or as emulsions. A more recently revised approach for parenteral administration involves use of a slow release or sustained release system such that a constant dosage is maintained. See, e.g., U.S. Pat. No. 3,610,795, which is incorporated by reference herein.

**[0132]** The materials may be in solution, suspension (for example, incorporated into microparticles, liposomes, or cells). These may be targeted to a particular cell type via antibodies, receptors, or receptor ligands. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Senter et al., *Bioconjugate Chem.* 2:447-451 (1991); Bagshawe *Br. J. Cancer* 60:275-281 (1989); Bagshawe et al., *Br. J. Cancer* 58:700-703 (1988); Senter et al., *Bioconjugate Chem.* 4:3-9 (1993); Battelli et al., *Cancer Immunol. Immunother.* 35:421-425 (1992); Pietersz and McKenzie, *Immunolog. Reviews* 129:57-80 (1992); and Roffler et al., *Biochem. Pharmacol.* 42:2062-2065 (1991)). Vehicles such as "stealth" and other antibody conjugated liposomes (including lipid mediated drug targeting to colonic carcinoma), receptor mediated targeting of DNA through cell specific ligands, lymphocyte directed tumor targeting, and highly specific therapeutic retroviral targeting of murine glioma cells in vivo. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Hughes et al., *Cancer Research* 49:6214-6220 (1989); and Litzinger and Huang, *Biochimica et Biophysica Acta* 1104:179-187 (1992)). In general, receptors are involved in pathways of endocytosis, either constitutive or ligand induced. These receptors cluster in clathrin-coated pits, enter the cell via clathrin-coated vesicles, pass through an acidified endosome in which the receptors are sorted, and then either recycle to the cell surface, become stored intracellularly, or are degraded in lysosomes. The internalization pathways serve a variety of functions, such as nutrient uptake, removal of activated proteins, clearance of macromolecules, opportunistic entry of viruses and toxins, dissociation and degradation of ligand, and receptor-level regulation. Many receptors follow more than one intracellular pathway, depending on the cell type, receptor concentration, type of ligand, ligand valency, and ligand concentration. Molecular and cellular mechanisms of receptor-mediated endocytosis has been reviewed (Brown and Greene, *DNA and Cell Biology* 10:6, 399-409 (1991)).

#### XX. Pharmaceutically Acceptable Carriers

**[0133]** The compositions, including antibodies, can be used therapeutically in combination with a pharmaceutically acceptable carrier. Suitable carriers and their formulations are described in *Remington: The Science and Practice of Pharmacy* (19th ed.) ed. A. R. Gennaro, Mack Publishing Company, Easton, Pa. 1995. Typically, an appropriate amount of a pharmaceutically-acceptable salt is used in the formulation to

render the formulation isotonic. Examples of the pharmaceutically-acceptable carrier include, but are not limited to, saline, Ringer's solution and dextrose solution. The pH of the solution is preferably from about 5 to about 8, and more preferably from about 7 to about 7.5. Further carriers include sustained release preparations such as semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, liposomes or microparticles. It will be apparent to those persons skilled in the art that certain carriers may be more preferable, depending upon, for instance, the route of administration and concentration of composition being administered.

**[0134]** Pharmaceutical carriers are known to those skilled in the art. These most typically would be standard carriers for administration of drugs to humans, including solutions such as sterile water, saline, and buffered solutions at physiological pH. The compositions can be administered intramuscularly or subcutaneously. Other compounds will be administered according to standard procedures used by those skilled in the art.

**[0135]** Pharmaceutical compositions may include carriers, thickeners, diluents, buffers, preservatives, surface active agents and the like in addition to the molecule of choice. Pharmaceutical compositions may also include one or more active ingredients such as antimicrobial agents, antiinflammatory agents, anesthetics, and the like.

**[0136]** The pharmaceutical composition may be administered in a number of ways depending on whether local or systemic treatment is desired, and on the area to be treated. Administration may be topically (including ophthalmically, vaginally, rectally, intranasally), orally, by inhalation, or parenterally, for example by intravenous drip, subcutaneous, intraperitoneal or intramuscular injection. The disclosed antibodies can be administered intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, or transdermally.

**[0137]** Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

**[0138]** Formulations for topical administration may include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

**[0139]** Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets, or tablets. Thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders may be desirable.

**[0140]** Some of the compositions may potentially be administered as a pharmaceutically acceptable acid- or base-addition salt, formed by reaction with inorganic acids such as

hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide, potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines.

#### XXI. Therapeutic Uses

**[0141]** Effective dosages and schedules for administering the compositions may be determined empirically, and making such determinations is within the skill in the art. The dosage ranges for the administration of the compositions are those large enough to produce the desired therapeutic or prophylactic effect. The dosage should not be so large as to cause adverse side effects, such as unwanted cross-reactions, anaphylactic reactions, and the like. Generally, the dosage will vary with the age, condition, sex and extent of the disease in the patient, route of administration, or whether other drugs are included in the regimen, and can be determined by one of skill in the art. The dosage can be adjusted by the individual physician in the event of any counterindications. Dosage can vary, and can be administered in one or more dose administrations daily, for one or several days. Guidance can be found in the literature for appropriate dosages for given classes of pharmaceutical products. For example, guidance in selecting appropriate doses for antibodies can be found in the literature on therapeutic uses of antibodies, e.g., *Handbook of Monoclonal Antibodies*, Ferrone et al., eds., Noyes Publications, Park Ridge, N.J., (1985) ch. 22 and pp. 303-357; Smith et al., *Antibodies in Human Diagnosis and Therapy*, Haber et al., eds., Raven Press, New York (1977) pp. 365-389. A typical daily dosage of the antibody used alone might range from about 1 µg/kg to up to 100 mg/kg of body weight or more per day, depending on the factors mentioned above.

**[0142]** Following administration of a disclosed composition, such as an antibody, for treating, inhibiting, or preventing an HTLV-3 or HTLV-4 infection, the efficacy of the therapeutic antibody can be assessed in various ways well known to the skilled practitioner. For instance, one of ordinary skill in the art will understand that a composition, such as an antibody disclosed herein, is efficacious in treating or inhibiting an HTLV-3 or HTLV-4 infection in a subject by observing that the composition reduces viral load or prevents a further increase in HTLV-3 or HTLV-4 viral load. Techniques used to measure the response of HTLV-3 or HTLV-4-infected subject to treatment with an antibody include determining whether the treatment partially or completely inhibits the appearance of the virus in the blood or other body fluid.

**[0143]** Other molecules that interact with HTLV-3 or HTLV-4 (or the proteins encoded by those virus nucleic acid sequences) can be used that do not have a specific pharmaceutical function, but which may be used for tracking changes within cellular chromosomes or for the delivery of diagnostic tools, for example can be delivered in ways similar to those described for the pharmaceutical products. The disclosed compositions and methods can also be used, for example, as tools to isolate and test new drug candidates for a variety of primate T-lymphotropic virus related diseases.

#### XXII. Treatment and Prevention Methods

**[0144]** By "treating" is meant an improvement in or abatement of the disease state (e.g., viral infection, bacterial infec-

tion, parasitic infection, cancer, genetic disorder, or autoimmune disease) is observed and/or detected upon or after administration of a substance of the present disclosure to a subject. Treatment can range from a positive change in a symptom or symptoms of the disease to complete amelioration of the disease (e.g., viral infection, bacterial infection, parasitic infection, or cancer) (e.g., reduction in severity, intensity, or duration of disease, alteration of clinical parameters indicative of the subject's condition, relief of discomfort or increased or enhanced function), as detected by art-known techniques. The methods of the present disclosure can be utilized, for instance, to prevent or treat a viral infection, bacterial infection, parasitic infection, or cancer. One of skill in the art would recognize that this viral infection, bacterial infection, parasitic infection, or cancer can include conditions characterized by the presence of a foreign pathogen or abnormal cell growth. Clinical symptoms will depend on the particular condition and are easily recognizable by those skilled in the art of treating the specific condition. Treatment methods can include, but are not limited to therapeutic vaccinations. Thus, disclosed are methods of treating a subject with a condition comprising administering to the vector or other composition disclosed herein.

**[0145]** Also disclosed are methods wherein the condition being treated or prevented is a viral infection. The viral infection can be selected from the list of viruses consisting of Herpes simplex virus type-1, Herpes simplex virus type-2, Cytomegalovirus, Epstein-Barr virus, Varicella-zoster virus, Human herpesvirus 6, Human herpesvirus 7, Human herpesvirus 8, Variola virus, Vesicular stomatitis virus, Hepatitis A virus, Hepatitis B virus, Hepatitis C virus, Hepatitis D virus, Hepatitis E virus, Rhinovirus, SARS, Coronavirus, Influenza virus A, Influenza virus B, Measles virus, Polyomavirus, Human Papillomavirus, Respiratory syncytial virus, Adenovirus, Coxsackie virus, Dengue virus, Mumps virus, Poliovirus, Rabies virus, Rous sarcoma virus, Yellow fever virus, Ebola virus, Marburg virus, Lassa fever virus, Eastern Equine Encephalitis virus, Japanese Encephalitis virus, St. Louis Encephalitis virus, Murray Valley fever virus, West Nile virus, Rift Valley fever virus, Rotavirus A, Rotavirus B, Rotavirus C, Sindbis virus, Simian Immunodeficiency virus, Human T-lymphotropic virus type-1, Human T-lymphotropic virus type-2, Primate T-lymphotropic virus, Hantavirus, Rubella virus, Simian Immunodeficiency virus, Human Immunodeficiency virus type-1, and Human Immunodeficiency virus type-2.

**[0146]** Also disclosed are methods wherein the condition being treated or prevented is a bacterial infection. The bacterial infection can be selected from the list of bacterium consisting of *M. tuberculosis*, *M. bovis*, *M. bovis* strain BCG, BCG substrains, *M. avium*, *M. intracellulare*, *M. africanum*, *M. kansasii*, *M. marinum*, *M. ulcerans*, *M. avium* subspecies *paratuberculosis*, *Nocardia asteroides*, other *Nocardia* species, *Legionella pneumophila*, other *Legionella* species, *Salmonella typhi*, other *Salmonella* species, *Shigella* species, *Yersinia pestis*, *Pasteurella haemolytica*, *Pasteurella multocida*, other *Pasteurella* species, *Actinobacillus pleuropneumoniae*, *Listeria monocytogenes*, *Listeria ivanovii*, *Brucella abortus*, other *Brucella* species, *Cowdria ruminantium*, *Chlamydia pneumoniae*, *Chlamydia trachomatis*, *Chlamydia psittaci*, *Coxiella burnetii*, other *Rickettsial* species, *Ehrlichia* species, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Bacillus anthracis*, *Escherichia coli*, *Vibrio cholerae*,

*Campylobacter* species, *Neisseria meningitidis*, *Neisseria gonorrhoea*, *Pseudomonas aeruginosa*, other *Pseudomonas* species, *Haemophilus influenzae*, *Haemophilus ducreyi*, other *Haemophilus* species, *Clostridium tetani*, other *Clostridium* species, *Yersinia enterocolitica*, and other *Yersinia* species.

**[0147]** Also disclosed are methods wherein the antigen-encoding nucleic acid is an antigen from a bacterium. The bacterial antigen can be selected from the group consisting of *M. tuberculosis*, *M. bovis*, *M. bovis* strain BCG, BCG substrains, *M. avium*, *M. intracellulare*, *M. africanum*, *M. kansasii*, *M. marinum*, *M. ulcerans*, *M. avium* subspecies *paratuberculosis*, *Nocardia asteroides*, other *Nocardia* species, *Legionella pneumophila*, other *Legionella* species, *Salmonella typhi*, other *Salmonella* species, *Shigella* species, *Yersinia pestis*, *Pasteurella haemolytica*, *Pasteurella multocida*, other *Pasteurella* species, *Actinobacillus pleuropneumoniae*, *Listeria monocytogenes*, *Listeria ivanovii*, *Brucella abortus*, other *Brucella* species, *Cowdria ruminantium*, *Chlamydia pneumoniae*, *Chlamydia trachomatis*, *Chlamydia psittaci*, *Coxiella burnetii*, other *Rickettsial* species, *Ehrlichia* species, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Bacillus anthracis*, *Escherichia coli*, *Vibrio cholerae*, *Campylobacter* species, *Neisseria meningitidis*, *Neisseria gonorrhoea*, *Pseudomonas aeruginosa*, other *Pseudomonas* species, *Haemophilus influenzae*, *Haemophilus ducreyi*, other *Hemophilus* species, *Clostridium tetani*, other *Clostridium* species, *Yersinia enterocolitica*, and other *Yersinia* species.

**[0148]** Also disclosed are methods wherein the condition being treated or prevented is a fungal infection. The fungal infection can be selected from the list of fungus consisting of *Candida albicans*, *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Aspergillus fumigatus*, *Coccidioides immitis*, *Paracoccidioides brasiliensis*, *Blastomyces dermatitidis*, *Pneumocystis carinii*, *Penicillium marneffi*, and *Alternaria alternata*.

**[0149]** Also disclosed are methods wherein the condition being treated is a parasitic infection. The parasitic infection can be selected from the list of parasites consisting of *Toxoplasma Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium malariae*, other *Plasmodium* species, *Trypanosoma brucei*, *Trypanosoma cruzi*, *Leishmania major*, other *Leishmania* species, *Schistosoma mansoni*, other *Schistosoma* species, and *Entamoeba histolytica*.

**[0150]** In addition, the disclosed vectors and vector containing compositions can be used to treat any disease where uncontrolled cellular proliferation occurs, such as a cancer. A non-limiting list of different types of cancers that can be treated with the disclosed compositions is as follows: lymphomas (including Hodgkin's and non-Hodgkin's, B cell lymphoma, and T cell lymphoma), mycosis fungoides, leukemias (including myeloid leukemia), carcinomas, carcinomas of solid tissues, squamous cell carcinomas, adenocarcinomas, sarcomas, gliomas, high grade gliomas, bladder cancer, brain cancer, nervous system cancer, head and neck cancer, blastomas, neuroblastomas, plasmacytomas, histiocytomas, melanomas, adenomas, hypoxic tumors, myelomas, AIDS-related lymphomas or sarcomas, kidney cancer, lung cancers such as small cell lung cancer and non-small cell lung cancer, neuroblastoma/glioblastoma, ovarian cancer, pancreatic cancer, prostate cancer, skin cancer, liver cancer, melanoma, squamous cell carcinomas of the mouth, throat, larynx, and lung, large bowel cancer, hematopoietic cancers; testicu-

lar cancer; colon and rectal cancers, prostatic cancer, or pancreatic cancer, cervical cancer, cervical carcinoma, breast cancer, and epithelial cancer, renal cancer, genitourinary cancer, pulmonary cancer, esophageal carcinoma, metastatic cancers, or cancers in general.

**[0151]** Also disclosed are methods wherein the antigen-encoding nucleic acid is a tumor antigen. The tumor antigen can be selected from the list consisting of human epithelial cell mucin (Muc-1; a 20 amino acid core repeat for Muc-1 glycoprotein, present on breast cancer cells and pancreatic cancer cells), the Ha-ms oncogene product, p53, carcino-embryonic antigen (CEA), the raf oncogene product, gp100/pme117, GD2, GD3, GM2, TF, sTn, MAGE-1, MAGE-3, BAGE, GAGE, tyrosinase, gp75, Melan-A/Mart-1, gp100, HER2/neu, EBV-LMP 1 & 2, HPV-F4, 6, 7, prostate-specific antigen (PSA), HPV-16, MUM, alpha-fetoprotein (AFP), C017-1A, GA733, gp72, p53, the ras oncogene product, HPV E7, Wilm's tumor antigen-1, telomerase, and melanoma gangliosides.

**[0152]** Disclosed are methods of treating a condition in a subject comprising administering to the subject the vector of the disclosure, wherein the condition is due to a mutated, disregulated, disrupted, or deleted gene; autoimmunity; or inflammatory diseases, including but not limited to cystic fibrosis, asthma, multiple sclerosis, muscular dystrophy, diabetes, tay-sachs, spinobifida, cerebral palsy, Parkinson's disease, Lou Gehrig's disease, Alzheimer's, systemic lupus erythematosus, hemophilia, Addison's disease, Cushing's disease.

**[0153]** By "preventing" is meant that after administration of a substance of the present disclosure to a subject, the subject does not develop the symptoms of the viral, bacterial, or parasitic infection, and/or does not develop the viral, bacterial, or parasitic infection. "Preventing" or "prevention" can also refer to the ultimate reduction of an infection, condition, or symptoms of an infection, or condition relative to infections or conditions in subjects that do not receive the substance. Methods of prevention can include, but are not limited to prophylactic vaccination. As such, disclosed are methods of preventing an infection in a subject comprising administering to the subject the vector of the disclosure.

**[0154]** Also disclosed are methods of the disclosure, wherein the infection prevented is a fungal infection or the antigen-encoding nucleic acid is an antigen from a fungus. The fungal infection or antigen can be selected from the list of *Candida albicans*, *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Aspergillus fumigatus*, *Coccidioides immitis*, *Paracoccidioides brasiliensis*, *Blastomyces dermatitidis*, *Pneumocystis carinii*, *Penicillium marneffi*, and *Alternaria alternata*.

**[0155]** Also disclosed are methods of the disclosure, wherein the antigen-encoding nucleic acid is an antigen from a parasite. The parasitic antigen can be selected from the group consisting of *Toxoplasma gondii*, *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium malariae*, other *Plasmodium* species, *Trypanosoma brucei*, *Trypanosoma cruzi*, *Leishmania major*, other *Leishmania* species, *Schistosoma mansoni*, other *Schistosoma* species, and *Entamoeba histolytica*.

**[0156]** Also disclosed are methods of the disclosure, wherein the subject is a horse, cow, pig, dog, cat, mouse, monkey, human, or a cell isolated from such an animal.

### XXIII. Screening Methods

**[0157]** Disclosed herein are methods of identifying new primate T-lymphotropic viruses comprising: a) contacting a

nucleic acid using a first set of primers and a second set of primers internal to the first set of primers, wherein the first set of primers is SEQ ID NOs: 19 and 20, and wherein the second set of primers is SEQ ID NOs: 21 and 22 under conditions that permit primer extension; b) identifying any amplified nucleic acid; and c) comparing the sequence to known primate T-lymphotropic viral sequences, wherein a sequence divergence greater than 5% indicates a new virus.

**[0158]** Also disclosed are methods of identifying new primate T-lymphotropic viruses comprising: a) contacting a nucleic acid using a first set of primers and a second set of primers internal to the first set of primers, wherein the first set of primers is selected from the group of primers pairs consisting of SEQ ID NOs: 7 and 8, SEQ ID NOs: 11 and 12, SEQ ID NOs: 15 and 16, SEQ ID NOs: 23 and 24, SEQ ID NOs: 27 and 28, SEQ ID NOs: 31 and 32, SEQ ID NOs: 69 and 70, SEQ ID NOs: 73 and 74, and SEQ ID NOs: 77 and 78, wherein the second set of primers is selected from the group consisting of SEQ ID NOs: 9 and 10, SEQ ID NOs: 13 and 14, SEQ ID NOs: 17 and 18, SEQ ID NOs: 25 and 26, SEQ ID NOs: 29 and 30, SEQ ID NOs: 33 and 34, SEQ ID NOs: 71 and 72, SEQ ID NOs: 75 and 76, and SEQ ID NOs: 79 and 80; b) identifying any amplified nucleic acid; and c) comparing the sequence to known primate T-lymphotropic viral sequences, wherein sequence divergence greater than 5% indicates a new virus.

**[0159]** It is also understood that the disclosed methods of identifying a new primate T-lymphotropic virus can be achieved using non-nested PCR techniques such as real-time PCR. Thus, for example, specifically disclosed are methods of identifying new primate T-lymphotropic viruses comprising a) contacting a nucleic acid using a set of primers, wherein the set of primers is selected from the set of primers consisting of SEQ ID NOs: 19 and 20, SEQ ID NOs: 21 and 22, SEQ ID NOs: 7 and 8, SEQ ID NOs: 11 and 12, SEQ ID NOs: 15 and 16, SEQ ID NOs: 23 and 24, SEQ ID NOs: 27 and 28, SEQ ID NOs: 31 and 32, SEQ ID NOs: 69 and 70, SEQ ID NOs: 73 and 74, SEQ ID NOs: 77 and 78, SEQ ID NOs: 9 and 10, SEQ ID NOs: 13 and 14, SEQ ID NOs: 17 and 18, SEQ ID NOs: 25 and 26, SEQ ID NOs: 29 and 30, SEQ ID NOs: 33 and 34, SEQ ID NOs: 71 and 72, SEQ ID NOs: 75 and 76, SEQ ID NOs: 79 and 80, and SEQ ID NOs: 64 and 65; b) identifying any amplified nucleic acid; and c) comparing the sequence to known primate T-lymphotropic viral sequences, wherein sequence divergence greater than 5% indicates a new virus. Also disclosed are identification methods wherein the method is a real-time PCR method.

**[0160]** Furthermore, the disclosed methods can be used in conjunction with probes to detect the presence of amplification product. Specifically disclosed are fluorescently labeled probes that can be used to detect the amplification product of the disclosed methods. For example, a fluorescent probe, can comprise  
 TTCCCCAAGGCTTCAAAAACAGC-  
 CCCACGC (SEQ ID NO: 66).

**[0161]** The surface antigen (SU) and transmembrane regions of env can be used serologically for the identification and differentiation of PTLVs (the type specific peptides MTA-1 and K55 are in SU; likewise the p24 region of gag can be used for the serological identification of PTLV). Thus, disclosed herein are methods of identifying a PTLV comprising contacting a nucleic acid with a set of primers specific for the surface antigen or transmembrane regions of env and identifying any amplified nucleic acid.

**[0162]** In addition, the disclosed peptides, polypeptides, proteins and protein fragments can be used to generate antibodies that can be used to identify new and known primate T-lymphotropic viruses. Specifically disclosed are methods of identifying the presence of a primate T-lymphotropic virus in a subject comprising taking a tissue sample from the subject and contacting the sample with an antibody directed to an HTLV-3 or HTLV-4 peptide, polypeptide, protein, or protein fragment, wherein the peptide, polypeptide, protein, or protein fragment can be SEQ ID NO: 37, 38, 39, 40, 41, 42, 43, 44, 46, 48, 50, 54, 55, 56, 57, 58, 60, 62, 67, or 68, or the polypeptide, protein, or protein fragment encoded by the nucleic acid of SEQ ID NO: 1, 2, 3, 4, 5, 6, 35, 36, 45, 47, 49, 51, 52, 53, 59, 61, 63, or 81, and wherein binding of the antibody to the sample indicates the presence of a new or known primate T-lymphotropic virus. The disclosed methods also can be used to identify new primate T-lymphotropic viruses as well as detect all primate T-lymphotropic viruses or a group of particular primate T-lymphotropic viruses. Those of skill in the art will know which antibodies to use to accomplish their detection goal. For example, to detect more than one of the known HTLV viruses (HTLV-1, 2, and 3, or HTLV-1, 2, and 4) one can use type specific peptide of HTLV-1 and HTLV-2 such as SEQ ID NO: 67 and 68.

**[0163]** Also provided is a method of screening a substance for effectiveness in treating or reducing the severity of the condition (e.g., HTLV-3 or HTLV-4 infection) comprising: a) obtaining an animal having the condition or characteristic (e.g., symptom) of the condition; b) administering the substance to an animal having one or more characteristics of the condition; and assaying the animal for an effect on the condition, thereby identifying a substance effective in reducing the condition. The ability of a substance to reduce the severity of a condition can be determined by evaluating the histological and/or clinical manifestations of the condition before and after administration of the substance of interest, and quantitating the degree of reduction of the histological and/or clinical manifestations of the condition. The animal in which the condition or characteristic (e.g., symptom) of the condition is produced can be any mammal, and can include but is not limited to mouse, rat, guinea pig, hamster, rabbit, cat, dog, goat, monkey, and chimpanzee. The condition or characteristic (e.g., symptom) of the condition can be produced in the animal by any method known in the art. For example, HTLV-3 or HTLV-4 can be produced by introducing into the animal (e.g., a chimpanzee infected with HTLV-3 or HTLV-4 or rhesus macaques or nemestrina macaques infected with an HTLV-3 or HTLV-4 env on an SIV backbone. Pullium et al., *J. Infectious Dis.* 183:1023, 2001) an infectious amount of HTLV-3 or HTLV-4.

**[0164]** The present disclosure also provides a method of screening for a substance effective in preventing the condition (e.g., HTLV-3 or HTLV-4 infection) comprising: a) administering the substance to an animal susceptible to the condition; b) subjecting the animal to treatment that will induce the condition or characteristic (e.g., symptom) of the condition; and c) assaying cells from the animal for an change in immune responses as compared to an the immune responses in a control animal having the condition in the absence of the substance identifies a substance that is effective in preventing the condition.

**[0165]** Also provided is a model for use in screening for substances effective in treating or preventing a disease com-

prising an animal capable of manifesting a characteristic of the disease, wherein the animal has been administered the vector of the disclosure.

**[0166]** Further embodiments are methods of making a model of HTLV-3 or HTLV-4 infection, comprising obtaining an animal capable of manifesting a characteristic of the disease, and administering to said animal one of the vectors disclosed herein that encodes an antigen associate with the disease. Also disclosed is a method of screening for a substance effective in treating a disease associated with an immunizing construct, the method comprising: a) administering the substance to the model of the disclosure; and b) assaying for an change in the course of the disease as compared to an the course of the disease in a control subject. An improvement in the course of the disease in the presence of the substance identifies a substance that is effective in treating the disease.

**[0167]** Still other embodiments are methods of screening for a substance effective in preventing a disease associated with an immunizing construct, the method comprising: a) administering one of the vectors disclosed herein to a subject; b) subjecting the subject to treatment that will induce the disease or characteristic (e.g., symptom) of the disease; and c) assaying for an change in the course of the disease as compared to an the course of the disease in a control subject. An improvement in the course of the disease in the presence of the substance identifies a substance that is effective in preventing the disease.

**[0168]** Yet still other embodiments are methods of screening for a substance effective in treating a disease associated with an immunizing construct, the methods comprising: a) subjecting a subject to treatment that induces the disease or characteristic (e.g., symptom) of the disease; b) administering to the subject one of the vectors disclosed herein; and c) assaying for a change in the course of the disease as compared to an the course of the disease in a control subject. An improvement in the course of the disease in the presence of the substance identifies a substance that is effective in treating the disease.

#### XXIV. Methods of Using the Disclosed Compositions as Research Tools

**[0169]** The disclosed compositions can also be used diagnostic tools related to primate T-lymphotropic diseases such as HTLV-3 and HTLV-4.

#### XXV. Methods of Making the Compositions

**[0170]** The compositions disclosed herein and the compositions necessary to perform the disclosed methods can be made using any method known to those of skill in the art for that particular reagent or compound unless otherwise specifically noted.

#### XXVI. Processes for Making the Compositions

**[0171]** Disclosed are processes for making the disclosed compositions as well as making the intermediates leading to the compositions. For example, disclosed are nucleic acids in SEQ ID NOs: 1-6, 35, 36, 45, 53, and 81. There are a variety of methods that can be used for making these compositions, such as synthetic chemical methods and standard molecular biology methods.

**[0172]** In some embodiments, a nucleic acid molecule is produced by the process of linking in an operative way a nucleic acid comprising the sequence set forth in SEQ ID



TABLE 3-continued

Nonhuman primate exposures for human T-lymphotropic virus (HTLV)-infected central African hunters																
ID	Site	HTLV	Nearest PTLV	Sex	Age	Technique	NHP Exposure									
							Hunting			Butcher			Pet			Reported
							m	c	g	m	c	g	m	c	g	
2656	ND	HTLV-1	Group G - Central West Africa	m	65	G	x			x				x		
1259	NG	HTLV-1	Group D - Mandrill clade	m	71	g, s	x	x						bitten/scratched by wild animal		
2810	YI	HTLV-1	Group G - Central West Africa	m	55	S	x			x						

\* m = monkey, c = chimpanzee, g = gorilla.

†, PTLV, primate T lymphotropic virus; STLV, simian T-lymphotropic virus

TABLE 4

Nucleotide and Amino Acid Percent Identities <sup>1</sup>					
	HTLV-1 (ATK)	HTLV-2 (MoT)	STLV-2 (PP1664)	STLV-3 (TGE2117)	HTLV-3 (2026ND)
HTLV-3 <sub>2026ND</sub>					
Genome (8917-bp)	61.6	62.9	62.6	87.0	—
LTR(697-bp)	48.7	43.7	41.4	86.7	—
gag (1268-bp)	69.3 (83.2)	69.4 (80.5)	70.6 (80.7)	87.5 (96.0)	—
pro (534-bp)	59.7 (62.6)	59.2 (66.7)	59.4 (59.3)	84.3 (88.1)	—
pol (2670-bp)	62.2 (66.2)	63.9 (71.2)	63.5 (69.9)	86.2 (93.1)	—
env (1476-bp)	65.9 (73.8)	69.0 (78.2)	67.1 (77.4)	87.8 (95.7)	—
tax (1053-bp)	76.3 (81.4)	75.1 (83.4)	74.4 (80.4)	91.2 (97.4)	—
rex (549-bp)	76.9 (61.9)	76.3 (60.6)	75.8 (63.5)	87.6 (89.6)	—
pX (699-bp)	43.3	50.5	49.8	85.6	—
HTLV-4 <sub>1863LE</sub>					
Genome (5320-bp)	64.0	72.2	71.4	66.2	66.1
pro (273-bp) <sup>2</sup>	71.4 (55.6)	79.5 (28.1)	79.5 (36.0)	71.8 (29.2)	73.3 (31.7)
pol (2549-bp) <sup>2</sup>	63.6 (68.7)	71.4 (80.1)	71.0 (79.7)	65.2 (71.7)	64.8 (71.6)
env (1458-bp)	65.8 (75.9)	73.1 (85.3)	72.0 (85.5)	67.2 (78.8)	68.5 (79.4)
tax (765-bp) <sup>2</sup>	77.4 (85.1)	81.7 (92.6)	79.4 (92.9)	75.2 (86.7)	75.0 (86.3)
rex (512-bp)	76.0 (63.9)	79.5 (74.1)	80.7 (68.8)	72.5 (57.7)	72.7 (59.4)
pX (559-bp)	46.1	60.8	59.9	53.6	51.3

<sup>1</sup>Amino acid identities are in parentheses.

<sup>2</sup>Only partial sequences are available

**[0177]** Most notable of the findings was the discovery of a human virus that is distinct from all known PTLV lineages with 26-34% and 18-25% nucleotide divergence in the conserved pol and tax genes, respectively, a range of nucleotide divergence similar to that seen between HTLV-1, HTLV-2, and STLV-3 (Meertens et al. 2002; Table 4). This virus formed a separate phylogenetic lineage with a long branch length and significant bootstrap support in both the pol (FIG. 2a: pol tree) and tax trees. Phylogenetic analyses combined with GenBank blast searches show that this is the only known virus in this novel group. For these reasons this virus, which is designated HTLV-4, qualifies as the first member of a novel species in the deltaretrovirus genus. Following ICTV guidelines (van Regenmortel 2000) and pending formal classification, it is proposed that primate T-lymphotropic virus 4 (PTLV-4) be the name for this species, with PTLV-4(1863LE) as the prototype strain. HTLV-4 was found in a 48 year old male hunter (1863LE) from the southern forests of Cameroon who had an HTLV-2-like WB result and reported hunting

monkeys, chimpanzees, and gorillas, and also being bitten and scratched by a wild animal, although the animal causing the injury was not specified.

**[0178]** Also documented, with significant phylogenetic bootstrap support, is the first evidence of human infection within the PTLV-3 group (FIG. 2a: pol tree). This virus, which is designated HTLV-3, clusters with STLV-3 viruses present in West African NHPs as expected (FIG. 2d: LTR tree). HTLV-3 was found in a 63 year old male (2026ND) from the southern forests of Cameroon who had an HTLV-1-like WB result and who reported hunting and butchering of monkeys. The fact that this virus falls within the diversity of a group of STLVs first identified in 1994 (Goubau et al 1994) without evidence of a human counterpart to date, indicates that this infection was most likely acquired zoonotically through exposure to the blood or body fluids of a hunted NHP from this region (Courgnaud et al. 2004).

**[0179]** In addition, broad diversity of HTLV-1 viruses was also found in this collection. Of the 11 HTLV-1 sequences,

two did not fall within any of the known HTLV-1 subtypes but clustered clearly within a clade that included only STLV-1 from central and west Africa (FIG. 2d: LTR tree). One of these viruses clustered with STLV-1 from monkeys in Cameroon and was from a 65 year-old male (2656ND) from the southern forest zone of Cameroon. He reported hunting and butchering of monkeys and kept a gorilla as a pet (Table 3). The second virus clustered with STLV-1 recently identified in chimpanzees and red colobus monkeys (Leendertz et al. 2004) and was from a 55 year old male (2810YI) who reported hunting and butchering of monkeys (Table 3). The presence of these viruses in hunters, seen previously only among NHPs, indicates that these persons were infected zoonotically. This distinct clade is referred to as HTLV-1 subtype G. Three subjects (1259NG, 1127MO, 1842LE) from different villages were found to have HTLV-1 subtype D, viruses known to infect geographically overlapping populations of humans and mandrills in central Africa (FIG. 3 LTR tree (Mahieux et al. 1998). Two of the three viruses were found in hunters (Table 3), providing indirect evidence of cross-species transmission between humans and mandrills within subtype D and supporting further the claims of cross-species transmission of this subtype (Mahieux et al. 1998). These results are consistent with SFV infection from mandrills that was documented previously in this population (Wolfe et al 2004) and indicate that the frequent hunting of mandrills may explain the widespread transmission of mandrill retroviruses. Five persons (979MO, 1380MV, 1443MV, 1503MV, 1537MV) were infected with HTLV-1 subtype B viruses, which are known to be endemic among humans in central Africa and which are believed to have originated from STLV-1 in this region (Mahieux, R. et al 1997, Gessain, A. & Mahieux, R 2000; FIG. 3: LTR tree). Thus, these five new subtype B viruses may have been acquired either zoonotically from STLV-1-infected primates or from human-to-human transmission, or both.

**[0180]** Notably, a 71 year old female (1443MV) who reported butchering gorillas was found to be infected with a virus most closely related to STLV-1 found in two gorillas from Cameroon (Nerrienet 2004, Courgnaud et al 2004), although without significant bootstrap support (FIG. 3 LTR). Interestingly, person 1503MV is also WB positive for SFV (Wolfe et al 2004), indicating that zoonotic transmission in an individual is not limited to a single retrovirus and providing a biological setting for viral recombination and altered pathogenicity and transmissibility of these viruses. One person (2472LE) was infected with the HTLV-1 subtype A virus, a clade consisting of sequences from only globally disseminated HTLV-1 and thus this infection was most likely acquired through human-to-human transmission. DNA samples from the remaining 73 persons with reactive WB results were all negative by the generic PCR assay for tax sequences and four other sequences specific for each PTLV clade, including HTLV-4. The results demonstrate that HTLV diversity is far greater than previously understood. The data indicate that contact with the blood and body fluids of NHPs is a major factor in the emergence of novel HTLVs, which are known to be transmissible among humans and have the potential to cause disease. Because the hunting and butchering of wild NHPs is widespread throughout central Africa (Bowen-Jones & Pendry 1999) and STLVs are known to be highly prevalent among hunted NHPs (Courgnaud et al. 2004), it is suspected that zoonotic transmission of STLV is not a restricted risk. Since blood banks in central Africa do not generally screen for HTLV, further spread of these viruses

among central Africans may be facilitated by blood donations from infected persons. That HTLV-4 represents a previously unrecognized virus being transmitted between humans indicates that more substantial screening for this virus in central African populations is needed. The finding that both HTLV-4 and HTLV-3 are serologically indistinguishable from HTLV-1 and HTLV-2 in current assays can explain why these viruses have not been previously identified, and highlight the importance of improved diagnostic assays. The increasing evidence that primate hunting is associated with the emergence of a range of simian retroviruses (Wolfe et al. 2004b) calls for increased surveillance and follow-up of individuals exposed to the blood and body fluids of wild NHPs, and for effective strategies to control the hunting of NHPs.

## Methods

### Ethical Approvals

**[0181]** Studies were conducted in the context of a community-based HIV prevention campaign designed to provide information using Cameroonian educators and counselors and therefore to decrease transmission. Participation in the study was completely voluntary. The study protocol was approved by the Johns Hopkins Committee for Human Research, the Cameroon National Ethical Review Board, and the HIV Tri-Services Secondary Review Board. Questionnaires and matching samples were anonymized by removing all personal identifiers to provide an unlinked study population.

### Sample Preparation and Serology

**[0182]** Blood was collected from participants, transported to a central laboratory, processed into plasma and PBMC aliquots and stored at  $-80^{\circ}$  C. Initial screening for HTLV antibodies in serum and plasma samples was performed by using the Vironostika HTLV-1/2 microelisa system (Organon-Teknika, Durham, N.C.) following the manufacturer's instructions. Reactive samples were then tested in a WB test (HTLV Blot 2.4, Genelabs Diagnostics, Singapore) that contains disrupted HTLV-1 virions, a gp21 recombinant protein (GD21) common to both HTLV-1 and HTLV-2, and two HTLV-type specific recombinant envelope (Env) peptides, MTA-1 and K55, which allow serological differentiation of HTLV-1 and HTLV-2, respectively. Samples with reactivity to the Gag (p24) and Env (GD21) proteins were considered seropositive. Seropositive samples with reactivity to MTA-1 or K55 were considered HTLV-1-like or HTLV-2-like, respectively. Samples with reactivity to either p24 or GD21 alone or in combination with other HTLV proteins (FIG. 1) were considered indeterminate.

### PCR and Sequence Analysis

**[0183]** DNA was prepared from uncultured PBMCs and its integrity was confirmed by  $\beta$ -actin PCR as previously described. All DNA preparation and PCR assays were performed in a laboratory where only human samples are processed and tested following recommended precautions to prevent contamination. DNA samples were first screened with a generic PTLV tax PCR assay capable of detecting 222-bp sequences from each of the three major PTLV groups (Busch et al. 2000, van Dooren et al. 2004). Sequence analysis of this tax sequence provided broad genetic classification into each PTLV group. Phylogenetic resolution within the PTLV-1 and

PTLV-3 groups was done using LTR sequences as described previously (van Dooren et al. 2004, Meertens et al. 2001). A portion of the 3' HTLV-1 LTR from selected samples (1259NG, 1127MO, 1842LE, and 2810YI) was amplified by nested PCR using external primers 5VLTRext 5' AACCA-CCATTCCTCCCCATG 3' (SEQ ID NO: 19; Meertens et al. 2001) and 1MNDR1 5'GTCGTGAATGAAAGG-GAAAGGGGT 3' (SEQ ID NO: 20; Meertens et al. 2001), and the internal primers Enh780 5' TGACGACAACCCCT-CACCTCAA 3' (SEQ ID NO: 21; Meertens et al. 2001) and 1MNDR25' AGGGGTGGAACCTTCGATCTGTAA 3'(SEQ ID NO: 22; Meertens et al. 2001). The tax (577-bp) and polymerase (pol) (709-bp) sequences of HTLV-3 and HTLV-4 were amplified by nested PCR using primers designed from conserved PTLV regions. The external and internal tax primers are PTLVTPG 5'T(C/T)ACCT(G/A)GGACCCCATCGATGGACG 3' (SEQ ID NO: 7) and PGTAXR15' GAIGA(T/C)TGI A(C/G)TAC(T/C)AAA-GATGGCTG 3' (SEQ ID NO: 8) and PH2Rrev 5' CCTATC-CCTCGICTCCCTC CTT 3' (SEQ ID NO: 9) and PGTAXR25' TTIGGG(T/C)AIGGICCGG AAATCAT 3'(SEQ ID NO: 10), respectively. The external and internal pol primers are PGPOLFI 5' C(T/G)TTAAACCGA(A/G)CGCCT CCAGGC 3' (SEQ ID NO: 11) and PGPOLR1GG (T/C)(A/G)TGIA (A/G)CCA(A/G)(A/G)CIAG(T/G)GG CCA 3' (SEQ ID NO: 12) and PGPOLF2 5' AC(T/C)TGGT (C/T)(C/T) (G/C)(G/C)A(A/G)GGCCCTGGAGG 3' (SEQ ID NO: 13) and PGPOLR25' G(A/G)(T/C)(A/G)GGIGTIC CTTTIGAGACCCA 3'(SEQ ID NO: 14), respectively. Inosines (I) and wobble bases (N/N) were used to accommodate areas of heterogeneity (Table 5).

[0184] Additional diagnostic PCR with PTLV-specific primers was carried out on samples with negative results for the generic 222-bp tax fragments. Assays described previously were used for PTLV-1 env and STLV-3 LTR (van Dooren et al. 2004) and HTLV-2 env (Switzer et al. 1995). For HTLV-4, a new nested PCR assay was developed based on the

HTLV-4 tax sequence using the external primers 1863TF1 5' CTCCTTCTTTCAGTCCCGTGCAGG 3' (SEQ ID NO: 15) and 1863TR15' GGGGTAGTCAGGTTTGGCTGGTAT 3' (SEQ ID NO: 16) and the internal primers 1863TF2 5' CCTACCGCAACGGATGTCTTGTAAA 3' (SEQ ID NO: 17) and 1863TR25' TATGGCGCC GGTGTGATGATAAAG 3' (SEQ ID NO: 18) and standard conditions to generate a 275-bp fragment. Percent nucleotide divergence was calculated using the Gap program in the Genetic Computer Group's Wisconsin package. Sequences were aligned using the Clustal W program, gaps were removed, and distance-based trees were generated by using the Kimura two-parameter model in conjunction with the NJ method in the MEGA program (version 2.1) as described elsewhere (van Dooren et al. 2004). 1000 bootstrap replicates were used to test the reliability of the final topology of the trees.

#### Primate Taxonomic Nomenclature

[0185] Nomenclature used herein was as described. NHPs were coded using the first letter of the genus and the first two letters of the species names with their house names or codes within parentheses. Cmo=*Cercopithecus mona* (Mona monkey), Cne=*C. neglectus* (De Brazza's guenon), Cmi=*C. mitis* (Sykes's monkey), Cni=*C. nictitans* (greater spot-nosed guenon), Cae=*Chlorocebus* species (African green monkey), Cpo=*C. pogonias* (crowned monkey), Cto=*Cercocebus torquatus* (red-capped mangabey), Cag=*Cercocebus agilis* (agile mangabey), Mog=*Miopithecus ogouensis* (talapoin monkey), Ani=*Allenopithecus nigrrpyridis* (Allen's swamp monkey), Msp=*Mandrillus sphinx* (mandrill (mnd)), Pan=*Papio anubis* (olive baboon (bab)), Pcy=*P. cynocephalus* (yellow baboon), Pha=*P. hamadryas* (sacred baboon), Ppu=*P. ursinus* (chacma baboon), Ppa=*P. papio* (Guinea baboon), Pba=*Ptilocolobus badius* (red colobus monkey), Mto=*Macaca tonkeana* (Celebes macaque), Ptr=*Pan troglodytes* (chimpanzee), Ppn=*Pan paniscus* (bonobo), Ggo=*Gorilla gorilla* (western lowland gorilla).

TABLE 5

Sequences of primers used for amplifying partial tax, envelope (env), polymerase (pol) and LTR regions of primate T-cell lymphotropic viruses

Name	Primer sequence <sup>a</sup> (5' to 3')	SEQ ID NO.	Location <sup>b</sup>	Expected PCR product size (bp)	Annealing temp (° C.), No. of cycles
PH1F	TTGTCATCAGCCCACTTCCCAGG	(SEQ ID NO: 23)	tax, 7243-7262, outer		
PH2R	AAGGAGGGGAGTCGAGGGATAAGG	(SEQ ID NO: 24)	tax, 7478-7455, outer	236	50, 40
PH2F	CCCAGGITTCCGGCAAGCCITCT	(SEQ ID NO: 25)	tax, 7257-7280, inner		
PH2R <sup>c</sup>	AAGGAGGGGAGTCGAGGGATAAGG	(SEQ ID NO: 26)	tax, 7478-7455, inner	222	50, 40
PTLVTPG	T(C/T)ACCT(G/A)GGACCCCATCGATGGACG	(SEQ ID NO: 7)	tax, 7480-7504, outer		
PGTAXR1	GAIGA(T/C)TGIA(C/G)TAC(T/C)AAAGATGGCTG	(SEQ ID NO: 8)	tax, 8140-8115, outer	660	45, 40

TABLE 5-continued

Sequences of primers used for amplifying partial tax, envelope (env), polymerase (pol) and LTR regions of primate T-cell lymphotropic viruses					
Name	Primer sequence <sup>a</sup> (5' to 3')	SEQ ID NO.	Location <sup>b</sup>	Expected PCR product size (bp)	Annealing temp (° C.), No. of cycles
PH2Rrev	CCTTATCCCTCGICTCCCCTCCTT	(SEQ ID NO: 9)	tax, 7529-7552, inner		
PGTAXR2	TTIGGG (T/C) AIGGICCGAAATCAT	(SEQ ID NO: 10)	tax, 8106-8085, inner	577	45, 40
PGPOLF1	C (T/G) TTAAACCIGA (A/G) CGCCTCCAGGC	(SEQ ID NO: 11)	pol, 2611-2634, outer		
PGPOLR1	GG (T/C) (A/G) TGIA (A/G) CCA (A/G) (A/G) CIAG (T/G) GGCCA	(SEQ ID NO: 12)	pol, 3598-3575, outer	987	45, 40
PGPOLF2	AC (T/C) TGGT (C/T) (C/T) (G/C) (G/C) A (A/G) GGCCCTGGAGG	(SEQ ID NO: 13)	pol, 2643-2666, inner		
PGPOLR2	G (A/G) (T/C) (A/G) GGIGTICCTTTIGAGACCCA	(SEQ ID NO: 14)	pol, 3352-3329, inner	709	45, 40
PGENVF1	TGGATCCCGTGG (A/C) GI (C/T) TCCTIAA	(SEQ ID NO: 27)	env, 5114-5136, outer		
PGENVR1	GT (A/G) TAIG (C/G) (A/G) (C/G) AIGTCCAIG (A/C) (T/C) TGG	(SEQ ID NO: 28)	env, 5576-5552, outer	462	45, 40
PGENVF2	AIAGACC (T/A) (C/T) CAAC (A/T) CCATGGGTAA	(SEQ ID NO: 29)	env, 5186-5209, inner		
PGENVR2	G (A/C) (T/C) TGGCAICCIA (A/G) GTAIGGGCA	(SEQ ID NO: 30)	env, 5557-5535, inner	371	45, 40
GPLTRF1	(G/A) CCACCAICTIGIGGACAAATAGCTGA	(SEQ ID NO: 31)	LTR, 8256-8282, outer		
GPLTRR2	C (C/T) GGGCCAAGCCTCGCTGCAGGCA	(SEQ ID NO: 32)	LTR, 8830-8807, outer	575	45, 40
GPLTRF2	ACCIIGGCTCTGACGTCTCTCCCT	(SEQ ID NO: 33)	LTR, 8333-8356, inner		
GPLTRR2	GGCAGIAGAAGTGCTACTTTTCGAT	(SEQ ID NO: 34)	LTR, 8810-8787, inner	478	45, 40

<sup>a</sup>Inosines and wobble nucleotides were included in the primers to accommodate sequence heterogeneity.

<sup>b</sup>The positions of the pol, env, and tax primers are given according to human T-cell lymphotropic virus type 1 (strain ATK); the LTR primer positions are given according to the simian T-cell lymphotropic virus type 3 (strain PH969) genome.

<sup>c</sup>The primer PH2R is used with PH2F in a semi-nested PCR.

#### Nucleotide Sequence Accession Numbers

**[0186]** The GenBank accession numbers for the 28 new HTLV sequences include AY818406 and AY818433.

#### Example 2

Ancient Origin and Molecular Features of the Human T-Lymphotropic Virus Type 3 Revealed by Complete Genome Analysis

Comparison of the HTLV-3(2026ND) Proviral Genome with Prototypical PTLVs

**[0187]** Using a combination of primers designed from small sequences obtained in each of the three major genes of PTLV and the LTR region, the complete genome of HTLV-3

(2026ND) was successfully generated as depicted in FIG. 4. Sequence analysis of the overlapping regions, followed by comparison with the genetic structure of other PTLVs, demonstrated that the complete proviral genome of HTLV-3 (2026ND) is 8917-bp. Despite being genetically equidistant from HTLV-1 and HTLV-2, the genomic structure of HTLV-3(2026ND) was similar to that of other PTLVs and included the structural, enzymatic, and regulatory proteins all flanked by long terminal repeats (LTRs). Comparison of HTLV-3 (2026ND) with prototypical PTLV genomes demonstrates that this new human virus is equidistant from the PTLV-1 (62% identity) and PTLV-2 (63% identity) groups across the genome. The results also confirm that HTLV-3 has the closest nucleotide and protein sequence identity to STLV-3 (87-92% identity; Table 6).

TABLE 6

Percent Nucleotide and Amino Acid Identity of HTLV-3(2026ND) with other PTLV Prototypes <sup>1</sup>							
	HTLV-1 (ATK)	HTLV-2 (MoT)	STLV-2 (PP1664)	STLV-3 (PH969)	STLV-3 (PPAF3)	STLV-3 (CTO604)	STLV-3 (NG409)
Genome	61.6	62.9	62.6	86.7	92.0	88.4	90.6
LTR	48.7	43.7	41.4	86.2	91.1	86.9	86.9
gag	69.3 (83.2)	69.4 (80.5)	70.6 (80.7)	86.4 (95.5)	91.3 (97.6)	89.4 (96.2)	90.6 (96.7)
p19	(74.4)	(68.3)	(67.2)	(95.9)	(95.9)	(95.9)	(94.3)
p24	(90.1)	(90.1)	(90.6)	(98.1)	(99.1)	(98.6)	(99.1)
p15	(78.0)	(73.8)	(72.6)	(88.4)	(96.5)	(90.7)	(94.2)
pro	59.7 (62.6)	59.2 (66.7)	59.4 (59.3)	83.3 (87.0)	88.8 (91.5)	85.0 (89.3)	88.0 (90.4)
pol	62.2 (66.2)	63.9 (71.2)	63.5 (69.9)	86.1 (92.7)	92.6 (94.9)	88.4 (92.9)	92.0 (92.9)
env	65.9 (73.8)	69.0 (78.2)	67.1 (77.4)	88.1 (95.1)	92.3 (95.1)	88.4 (94.3)	91.2 (95.3)
SU <sup>2</sup>	(68.4)	(70.7)	(69.7)	(92.7)	(97.1)	(92.4)	(94.0)
TM <sup>2</sup>	(83.5)	(91.6)	(91.0)	(99.4)	(98.9)	(97.8)	(97.8)
rex	76.9 (61.9)	76.3 (60.6)	75.8 (63.5)	87.1 (88.5)	90.9 (94.5)	88.5 (94.0)	88.3 (92.3)
tax	75.4 (81.4)	73.1 (83.4)	72.3 (80.4)	90.2 (97.4)	94.0 (98.3)	91.4 (96.6)	92.8 (96.9)

<sup>1</sup>amino acid identity in parentheses; strain names given in parentheses below PTLV designation

<sup>2</sup>SU, surface protein; TM, transmembrane protein

**[0188]** The most genetic divergence between the PTLV groups was seen in the LTR region (52-59%) while the highest inter-group identity was observed in the highly conserved regulatory genes, tax and rex (72-77%). Interestingly, within the PTLV-3 group, HTLV-3(2026ND), which was identified in a hunter from Cameroon, was unique but shared the most overall sequence identity to STLV-3(PPAF3) (92%) from a Senegalese baboon instead of STLV-3(CTO604) (88.4%) identified in red-capped mangabeys, also from Cameroon. This relationship is highlighted further by comparison of HTLV-3(2026ND) with all available full-length STLV-3 genomes in similarity plot analysis where the highest identity was seen in the highly conserved tax gene. As seen within other PTLV groups, there was no clear evidence of genetic recombination of HTLV-3(2026ND) with STLV-3 or PTLV-1 and PTLV-2 proviral sequences by using bootscanning analysis. HTLV-3(2026ND) was not compared to the recently reported second strain of HTLV-3 because only two short sequences were available at GenBank and in these region this virus has been shown to be nearly identical to STLV-3(CTO604) (Callatini et al. (2005)*Retrovirology*. 2:30).

#### Organization of the LTR and Pre-Gag Region

**[0189]** As with STLV-3, the HTLV-3(2026ND) LTR (697-bp) was smaller than that of HTLV-1 (756-bp) and HTLV-2 (764-bp), by having two and not three of the 21-bp transcription regulatory repeat sequences in the U3 region (FIG. 5a; Meertens and Gessain. (2003)*J. Virol.* 77:782-789; Meertens et al. (2002)*J. Virol.* 76:259-268; Van Brussel et al. (1997)*J. Virol.* 71:5464-5472; Van Dooren et al., (2004)*J. Gen. Virol.* 85:507-519). Other regulatory motifs such as the polyadenylation signal, TATA box, and cap site were all conserved in the HTLV-3(2026ND) LTR (FIG. 5a). By secondary structure analysis of the LTR RNA sequence, a stable stem loop structure from nucleotides 421-464 (FIG. 5b) was also observed similar to that shown to be essential for Rex-responsiveness control of viral expression in both HTLV-1 and HTLV-2.

#### Analysis of the Genomic Structure of HTLV-3(2026ND)

**[0190]** Translation of predicted protein open reading frames (ORFs) across the viral genome identified all major Gag, Pol, Pro (protease), and Env proteins, as well as the

regulatory proteins, Tax and Rex. Translation of the overlapping gag and pro and pro and pol ORFs occurs by one or more successive-1 ribosomal frameshifts that align the different ORFs. The conserved slippage nucleotide sequence 6(A)-8 nt-6(G)-11nt-6(C) is present in the Gag-Pro overlap, while a point mutation in the Pro-Pol overlap slippage sequence (GT-TAAAC (SEQ ID NO: 82) compared to TTAAAC (SEQ ID NO: 83) in HTLV-1 and HTLV-2) was observed in HTLV-3(2026ND) but the asparagine codon (AAC) crucial for the slippage mechanism was unaffected.

**[0191]** The structural and group specific precursor Gag protein consisted of 422 amino acids (aa) that is predicted to be cleaved into the three core proteins p19 (matrix), p24 (capsid), and p15 (nucleocapsid) similar to HTLV-1, HTLV-2, and STLV-3. Across PTLVs, Gag was one of the most conserved proteins with identities ranging from 81% and 83% for HTLV-1 and PTLV-2, to 95% for STLV-3 supporting the observed cross-reactivity seen with PTLV-3 antisera in Western blot assays using HTLV-1 antigens. Within Gag, the capsid protein showed greater than 90% identity to HTLV-1, while the matrix and nucleocapsid proteins were more divergent sharing less than 78% identity to PTLV-1 and PTLV-2 indicating their potential use in serologic assays for discriminating the three major PTLV groups.

**[0192]** The predicted size of the Env polyprotein is 491 aa, which is slightly shorter than that found in STLV-3s (313 aa versus 314 and 315 for STLV-3(PH969) and STLV-3(CTO604) due to sequence variation at the carboxy terminus of the surface (SU) protein. In contrast, the transmembrane (TM) protein (178 aa) was highly conserved across all PTLVs supporting further the use of the recombinant HTLV-1 GD21 protein spiked onto WB strips for the identification of divergent PTLVs. Despite the weak reactivity of anti-HTLV-3(2026ND) antibodies to the HTLV-1 type specific SU peptide (MTA-1; Wolfe et al. (2005)*Proc. Natl. Acad. Sci. USA.* 102:7994-7999) spiked onto WB strips, there was only 70.8% identity of MTA-1 to HTLV-3(2026ND), which is similar to the 68.8% identity of MTA-1 to HTLV-2, demonstrating no clear correlation of WB profile and predicted SU sequence.

**[0193]** The HTLV-1 and HTLV-2 Tax proteins (Tax1 and Tax2, respectively) transactivate initiation of viral replication from the promoter in the 5' LTR and are thus essential for viral expression (Feuer and Green. (2005)*Oncogene*. 24:5996-

6004). Tax1 and Tax2 have also been shown to be important for T-cell immortalization, while the HTLV-3 Tax (Tax3) has not yet been characterized (Feuer and Green. (2005) *Oncogene*. 24:5996-6004). Hence, the Tax3 sequences were compared with those of prototypic HTLV-1, PTLV-2, and STLV-3s to determine if motifs associated with these functional characteristics are preserved. Alignment of predicted Tax3 sequences shows excellent conservation of the critical functional regions, including the nuclear localization signal (NLS), cAMP response element (CREB) binding protein (CBP)/p300 binding motifs, and nuclear export signal (NES; HTLV-3 Tax is shown in FIG. 6). The C-terminal transcriptional activating domain (CR2), essential for CBP/p300 binding, was also very conserved except for a single IN to F mutation at position five of the motif compared to HTLV-1 and PTLV-2, respectively. However, this single amino acid change in the STLV-3 Tax has recently been shown in transient transfection assays to have no deleterious effect on viral transactivation (Chevalier et al. (2005) *AIDS Res. Hum. Retrovir.* 21:513 (Abs. P174)). Since the predicted CR2 domain is conserved in Tax3, similar transactivation activity can be seen with HTLV-3.

**[0194]** Interestingly, although these important functional motifs are highly conserved in PTLV, phenotypic differences of HTLV-1 and HTLV-2 Tax proteins have been observed leading to speculation, that these differences account for the different pathologies associated with both HTLVs (Feuer and Green. (2005) *Oncogene*. 24:5996-6004). Recently, the C-terminus of Tax1, and not Tax2, has been shown to contain a conserved PDZ domain present in cellular proteins involved in signal transduction and induction of the IL-2-independent growth required for T-cell transformation (Rousset et al. (1998) *Oncogene*. 6:643-654; Tsubata et al. (2005) *Retrovirology*. 2:46). The presence of a PDZ domain in PTLV-1 and its absence in PTLV-2 indicates a potential role of this motif in the phenotypic differences of the two viral groups. The consensus PDZ domain has been defined as S/TXV-COOH, where the first amino acid is serine or threonine, X is any amino acid, followed by valine and the carboxy terminus. Examination of the PTLV-3 Tax sequences showed that both HTLV-3 and STLV-3 have predicted PDZ domains with the consensus sequence S(P/S)V compared to T(E/D)V in PTLV-1 (the HTLV-3 PDZ domain is shown in FIG. 6).

**[0195]** Besides Tax and Rex, two additional ORFs coding for four proteins (p27<sup>I</sup>, p12<sup>I</sup>, p30<sup>II</sup>, and p13<sup>II</sup> where I and II denote ORFI and ORFII, respectively) have been identified in the pX region of HTLV-1 (FIG. 4) and are important in viral infectivity and replication, T-cell activation, and cellular gene expression (Bindhu et al. (2004) *Front. Biosc.* 9:2556-2576). Analysis of the pX region of HTLV-3(2026ND) revealed a total of four putative ORFs (named I-IV, respectively) coding for 96, 122, 72, and 118 aa in length. While both ORFIII (72 aa) and ORFIV (118 aa) shared identity to the ORFII of STLV-3 and HTLV-1 and STLV-2/HTLV-2, respectively, and each contained two PXXP motifs, only ORF III was leucine rich like that seen in the leucine zipper motifs of ORFI p12<sup>I</sup> (Bindhu et al. (2004) *Front. Biosc.* 9:2556-2576). However, ORFIII did not share any sequence homology with p12<sup>I</sup> and both ORFI and ORFII shared only weak sequence identity to miscellaneous cellular proteins available at GenBank. Interestingly, 22 of 28 (79%) amino acids in ORFIV (pos 64-91) were identical among the ORFIIs of all PTLVs indicating a conserved functionality of this motif.

**[0196]** A protein termed the HTLV-1 basic leucine zipper (bZIP) factor (HBZ) was recently identified in translation of the complementary strand of the viral RNA genome between the env and tax/rex genes (Gaudray et al. (2002) *J. Virol.* 76:12813-12822). Although originally reported to be exclusive to PTLV-1 (Gaudray et al. (2002) *J. Virol.* 76:12813-12822), HBZ is conserved among PTLVs, including HTLV-3(2026ND) (HTLV-3 HBZ is shown in FIG. 7), demonstrating further the potential importance of this protein in viral replication and oncogenesis. The carboxy terminus of the HBZ ORF contains a 21 aa arginine rich region that is relatively conserved in PTLV and known cellular bZIP transcription factors, followed by a leucine zipper region possessing five or four conserved leucine heptads in HTLV-1 and all other PTLVs, respectively. PTLV-1 has 5 leucine heptads similar to that found in mammalian bZIP proteins, while PTLV-1 and PTLV-2 have four leucine heptads followed by leucine octet. Of all PTLVs with full length genomes available at GenBank, only HTLV-2(MoT) did not have the full complement of leucine heptads but was limited to the initial three leucine motifs due to a one nucleotide deletion at position 6823 causing a frameshift in the predicted HBZ sequence.

#### Phylogenetic Analysis

**[0197]** The genetic relationship of HTLV-3(2026ND) to PTLV-3 was confirmed by using aligned full-length prototype sequences excluding the LTR region (FIG. 8a). Phylogenetic analysis inferred three major PTLV groups with very high bootstrap support (100%) with HTLV-1, HTLV-2 and HTLV-3 each clustering in separate clades (FIG. 8a). Within the PTLV-3 phylogroup, HTLV-3(2026ND) formed a separate lineage but clustered with high bootstrap support with STLV-3s from west central Africa (strains CTO604, CTO-NG409, and PPA-F3) indicating a possible primate origin for this human infection in this geographic region. The relationship of HTLV-3 to STLV-3 was supported further by phylogenetic inference of identical tree topologies using an alignment of each major gene region (FIG. 8b-8d). The phylogenetic stability seen across the PTLV genome also demonstrates further the absence of major recombination events occurring in PTLV despite evidence of dual infections in humans and primates (Courgnaud et al. (2004) *J. Virol.* 78:4700-4709), compared to other retroviruses such as HIV which undergo frequent recombination.

#### Dating the Origin of HTLV-3(2026ND) and Other PTLVs

**[0198]** The finding of HTLVs in three distinct clades indicates an ancient, independent evolution of these viruses. Hence, additional molecular analyses was undertaken in order estimate the divergence times of the PTLV lineages. Although others have reported finding a clock-like behavior of STLV-3 sequences (Meertens and Gessain. (2003) *J. Virol.* 77:782-789; Meertens et al. (2002) *J. Virol.* 76:259-268; Meertens et al. (2003) *J. Gen. Virol.* 84:2723-2727), these results were not confirmed and instead found that PTLVs evolved at different rates by using an alignment of full-length PTLV genomes sans LTR sequences. However, reliable retrovirus divergence times can be obtained by using nonparametric rate smoothing of the sequences to relax the stringency of a clock assumption followed by time calibration of the tree using a value of 40,000-60,000 YA for the origin of the Melanesian HTLV-1 (Sanderson (2003) *Bioinformatics*.

19:301-2; Switzer et al. (2005) *Nature*. 434:376-380; Van Dooren et al. (2004) *J. Gen. Virol.* 85:507-519). By using these dates and methods, the mean evolutionary rate for PTLV was estimated to be  $1.12 \times 10^{-6}$  (confidence interval  $6.82 \times 10^{-7}$  to  $1.56 \times 10^{-6}$ ) substitutions/site/year, respectively, which is consistent with rates determined previously both with and without enforcing a molecular clock (Lemey et al. (2005) *Infect. Gen. Evol.* 5:291-298; Meertens and Gessain. (2003) *J. Virol.* 77:782-789; Meertens et al. (2002) *J. Virol.* 76:259-268; Meertens et al. (2003) *J. Gen. Virol.* 84:2723-2727; Salemi et al. (2000) *Mol. Biol. Evol.* 17:374-386; Van Dooren et al. (2004) *J. Gen. Virol.* 85:507-519). The mean evolutionary rate for HTLV-3(2026ND) is estimated to be  $9.94 \times 10^{-7}$  (confidence interval  $6.04 \times 10^{-7}$  to  $1.38 \times 10^{-6}$ ). The PTLV ancestor was estimated to have originated about 630,000-947,000 YA confirming an archaic evolution of the primate deltaretroviruses (FIG. 9; Salemi et al. (2000) *Mol. Biol. Evol.* 17:374-386). The separation of PTLV-1 and PTLV-2 occurred about 579,077-867,458 YA, while HTLV-2 and STL-2 diverged around 191,621-286,730 YA (FIG. 9). The origin of all PTLV-3s was estimated to be between 63,294-94,700 YA with the ancestor of HTLV-3(2026ND) occurring about 36,087-54,067 YA (FIG. 9) indicating an ancient origin of this virus in humans. Alternatively, HTLV-3 may represent a recent zoonoses from a primate infected with a very old, divergent STL-3. However, if HTLV-3 is an old human infection, then it appeared during the same period as the ancestor of both HTLV-1 and HTLV-2 (40,000-60,000 and 28,800-43,392 YA, respectively) and may have also spread to become endemic in specific populations yet to be identified.

#### Discussion

**[0199]** The complete nucleotide sequence and genomic characterization of the first HTLV-3 that is clearly distinct from all STL-3s and is genetically equidistant to HTLV-1 and HTLV-2 is described herein. HTLV-3(2026ND) is also unique from the second HTLV-3(Py143) reported recently in a Bakola pygmy from Cameroon since the latter strain is nearly identical to STL-3 found in a red-capped mangabey, based on the limited sequence data available for this virus (Callatini et al. (2005) *Retrovirology*. 2:30). Although HTLV-1 and HTLV-2 are pathogenic and have spread globally to become endemic in different human populations, little is known about the epidemiology of HTLV-3 infection. However, detailed, comparative sequence analyses of viral genomes can help provide important molecular clues to the origin, evolution, and public health importance of novel human infections.

**[0200]** Like other PTLVs, HTLV-3(2026ND) is genetically stable and its slow evolutionary rate, combined with estimates of known human migrations, can then be used to infer divergence times for HTLV. The finding that the predecessor of HTLV-3(2026ND) originated over 30 millennia ago, an age which is estimated that the ancestors of both HTLV-1 and -2 to have appeared, combined with the wide geographic distribution of STLVs and the recent finding of another HTLV-3 in an African pygmy (Callatini et al. (2005) *Retrovirology*. 2:30; Gessain and Mahieux. (2000) *Bull. Soc. Pathol. Exot.* 93:163-171; Meertens and Gessain. (2003) *J. Virol.* 77:782-789; Meertens et al. (2002) *J. Virol.* 76:259-268; Meertens et al. (2003) *J. Gen. Virol.* 84:2723-2727; Takemura et al. (2002) *J. Virol.* 76:1642-1648; Van Dooren et al. (2004) *J. Gen. Virol.* 85:507-519), collectively indicate that HTLV-3 infection be more frequent than previously understood. In addition, the

archaic age of the ancestral HTLVs and the recent finding of STL-3-like infections in African hunters collectively imply that cross-species transmission of STLVs to humans is both an ancient and contemporary phenomenon coupled to behavior that exposes humans to nonhuman primates. The ancient origin of HTLV contrasts with that reported for HIV, which is believed to have only crossed over into humans from SW-infected NHPs within the last century, and indicates a long period of viral evolution and adaptation in humans possibly resulting in the observed lower pathogenicity for HTLV compared to HIV (Hahn et al. (2000) *Science* 287:607-614; Sharp et al. (2000) *Biochem Soc Trans.* 28:275-282).

**[0201]** Screening for HTLV-3 can be facilitated by the application of diagnostic serologic and molecular assays based on the sequences reported here. For example, the data show that the Gag matrix and nucleocapsid regions and the envelope surface protein are relatively conserved within PTLV-3 but are divergent from PTLV-1 and PTLV-2 and can thus be used to differentiate the three PTLV groups with serological methods.

**[0202]** At the molecular level, examination of the genomic structure showed that the enzymatic, regulatory, and structural proteins are well preserved in HTLV-3(2026ND), including conserved functional motifs in Tax important for viral expression and T-cell proliferation. The finding of a PDZ domain in the Tax protein of HTLV-3(2026ND), like that seen in HTLV-1 but not HTLV-2 (Feuer and Green. (2005) *Oncogene*. 24:5996-6004), which has been shown to be important in cellular signal transduction and T-cell transformation (Rousset et al. (1998) *Oncogene*. 6:643-654; Tsubata et al. (2005) *Retrovirology*. 2:46), indicates that the HTLV-3 Tax is more phenotypically similar to HTLV-1 than HTLV-2. The high amino acid identity of the PTLV-3 Tax proteins combined with the ability of STL-3 to transform human cells in vitro indicates that the HTLV-3 Tax functions similarly (Goubau et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:2848-2852).

**[0203]** In contrast to the tax gene, the HTLV-3(2026ND) LTR has only two of the three conserved promoters identified in HTLV-1 and HTLV-2 that are responsible for basal viral transcription levels and like STL-3 is missing the TATA-distal 21-bp repeat element (Meertens and Gessain. (2003) *J. Virol.* 77:782-789; Meertens et al. (2002) *J. Virol.* 76:259-268; Meertens et al. (2003) *J. Gen. Virol.* 84:2723-2727; Van Brussel et al. (1997) *J. Virol.* 7:5464-5472; Van Dooren et al. (2004) *J. Gen. Virol.* 85:507-519). All of the remaining functional elements in the LTR were conserved, including the stem loop structure necessary for Rex responsive control of viral expression in HTLV-1 and -2.

**[0204]** Recently, a HBZ protein was identified in translation of the complementary strand of the viral RNA genome between the env and tax/rex genes (Gaudray et al. (2002) *J. Virol.* 76:12813-12822). Protein translation on the minus-strand RNA is a unique feature of HTLV-1 not previously seen in retroviruses. HBZ was shown to be involved in the negative regulation of viral replication (Gaudray et al. (2002) *J. Virol.* 76:12813-12822). The more recent finding of HBZ mRNA expression in ATL patients indicates a role of HBZ mRNA in the survival of leukemic cells in vivo and in HTLV-1-associated oncogenesis (Satou et al. (2006) *Proc. Natl. Acad. Sci. USA*. 103:720-725). Although originally reported to be exclusive to PTLV-1 (Gaudray et al. (2002) *J. Virol.* 76:12813-12822), HBZ is conserved among PTLVs, including HTLV-3(2026ND), demonstrating further the potential

importance of this protein in viral replication and oncogenesis. Of all PTLVs with full length genomes available at GenBank, only HTLV-2(MoT) did not have the full complement of leucine heptads in the leucine zipper due to a frameshift mutation in the predicted HBZ sequence.

**[0205]** In summary, disclosed herein, HTLV-3(2026ND) is genetically stable and has an ancient origin. HTLV-3(2026ND) genomic structure is relatively conserved and contains many of the functional motifs important for the viral expression and pathology associated with HTLVs.

## Materials and Methods

### DNA Preparation and PCR-Based Genome Walking

**[0206]** DNA was prepared from uncultured PBMCs available from person 2026ND identified in the original PTLV surveillance study in Cameroon reported in detail elsewhere (Wolfe et al. (2005) *Proc. Natl. Acad. Sci. USA.* 102:7994-7999). DNA integrity was confirmed by  $\beta$ -actin polymerase chain reaction (PCR) as previously described (Wolfe et al. (2005) *Proc. Natl. Acad. Sci. USA.* 102:7994-7999). All DNA preparation and PCR assays were performed in a laboratory where only human specimens are processed and tested according to recommended precautions to prevent contamination. To obtain the full-length genomic sequence of HTLV-3 small regions of each major coding region were PCR-amplified by using nested PCR and degenerate PTLV primers. The tax (577-bp) and polymerase (pol) (709-bp) sequences were amplified by using primers and conditions provided elsewhere (Wolfe et al. (2005) *Proc. Natl. Acad. Sci. USA.* 102:7994-7999). Envelope (env) (371-bp) sequences were amplified by using standard PCR conditions with a 45° C. annealing temperature and the external and internal primers PGENVF1 5' TGGATCCCGTGG(A/C)GI(C/T)TCCTIAA 3' (SEQ ID NO: 27) and PGENVR1 5' GT(A/G)TAIG(C/G)(A/G)(C/G)AIGTCCAIG(A/C)(T/C)TGG 3' (SEQ ID NO: 28) and PFENVF2 5' AIAGACC(T/A)(C/T)CAAC(A/T)CCATGGGTAA 3' (SEQ ID NO: 29) and PGENVR2 5' G(A/C)(T/C)TGGCAICCA(A/G)GTAIGGGCA 3' (SEQ ID NO: 30), respectively. A 398-bp fragment of the long terminal repeat (LTR) was obtained by using conserved STLV-3 primers as previously reported (Wolfe et al. (2005) *Proc. Natl. Acad. Sci. USA.* 102:7994-7999).

**[0207]** HTLV-3(2026ND)-specific primers were then designed from sequences obtained in each of the four viral regions described above and were used in nested, long-template PCRs to fill in the gaps in the genome as depicted in FIG. 4 by using an expand high fidelity kit containing both Taq and Tgo DNA polymerases (Roche). The external and internal primer sequences for the LTR-pol and pol-env fragments are 2026LF1 5' GGTAAGATCCCACTGGGTCGAGC 3'(SEQ ID NO: 69) and 2026PR1 5' GAAGCCAGGTCTCGGGT-GACG 3'(SEQ ID NO: 70) and 2026LF2 5' CGCTCCCCTG-GAGCTCTCTCG 3'(SEQ ID NO: 71) and 2026PR25' GCCACTTCCCATTGGGCTTTTGTACGG 3' (SEQ ID NO: 72) and 2026 PF3 5' GCTCTCACCGATAAAGTAA-CAAACG 3' (SEQ ID NO: 73) and 2026ER15' GGTAG-GAAGAGGCTCCTATGAACAG 3' (SEQ ID NO: 74) and 2026 PF2 5' CAGGACTGCATAACATACGAGACCCTCC 3' (SEQ ID NO: 75) and 2026ER35' CCTATGAACAGGGT-GCATCGACTGGG 3' (SEQ ID NO: 76), respectively. The external and internal primer sequences used to obtain about 3 kb of the 3' end of the genome (env-tax-LTR) are 2026EF1 5'

CCCTAAGCCCCCATGTCCAGAC 3'(SEQ ID NO: 77) and 2026LR15' CGAGAGAGCTCCAGGGGAGCG 3' (SEQ ID NO: 78) and 2026EF3 5' CCTACTCCCTGTATGTATTC-CCCCATTGG 3' (SEQ ID NO: 79) and 2026LR25' GCTC-GACCCAGTGGGATCTTACCGAGTGG 3' (SEQ ID NO: 80), respectively.

**[0208]** PCR products were revealed on 1.5% agarose gels stained with ethidium bromide, purified with a QIAQUICK™ PCR purification kit (Qiagen) and sequenced in both directions with a BIGDYE™ terminator cycle kit and automated sequencers (Applied Biosystems). Selected PCR products were also cloned into the pCR4-TOPO vector using the TOPO TA Cloning kit (Invitrogen) and recombinant plasmid DNA was prepared using the Qiagen plasmid purification kit prior to automated sequencing.

### Sequence and Phylogenetic Analysis

**[0209]** Percent nucleotide divergence was calculated by using the GAP program in the Genetic Computer Group's (GCG) Wisconsin package (Thompson et al. (1994) *Nucleic Acids Res.* 22:4673-4680). LTR RNA secondary structure was determined using the program RNAstructure v4.2 (Mathews et al. (1999) *J. Mol. Biol.* 288:911-940). Sequences were aligned by using the Clustal W program (Womble (2000) *Methods Mol. Biol.* 132:3-22), gaps were removed, and distance-based trees were generated by using the Kimura two-parameter model in conjunction with the neighbor-joining (NJ) method in the MEGA program (version 2.1) and maximum likelihood (ML) analysis in the PAUP\* program as described in detail elsewhere (Switzer et al. (2005) *Nature.* 434:376-380; Wolfe et al. (2005) *Proc. Natl. Acad. Sci. USA.* 102:7994-7999). The reliability of the final topology of the trees was tested with 1,000 bootstrap replicates. Comparison of full-length PTLV genomes available at GenBank was done using HTLV-3(2026ND) as the query sequence and the F84 (ML) model and a transition/transversion ratio of 2.0 implemented in the program SimPlot (Lole et al. (1999) *J. Virol.* 73:152-160).

**[0210]** For dating of HTLV-3(2026ND), full-length genomes from prototypical PTLVs available at GenBank were aligned with HTLV-3(2026ND) by using Clustal W, gaps were removed, and minor adjustments in the alignment were made manually. LTR sequences were excluded from the analysis since this region does not align accurately in PTLVs. The best fitting evolutionary model for the aligned sequences was determined with Modeltest v3.6 (Posada and Crandall. (1998) *Bioinformatics.* 14:817-818). The general time-reversible model, allowing six different substitution rate categories, with gamma-distributed rate heterogeneity (1.9724) and an estimated proportion of invariable sites (0.3687), was determined to be the best fit to the data. Little substitution saturation was observed in the 7213-bp alignment ( $P < 0.0001$ ) as determined with the DAMBE program, and was therefore satisfactory for use in phylogenetic analyses. Likewise, using the best-fitting evolutionary model defined above, good phylogenetic signal in the alignment was also found with likelihood mapping analysis using the program Tree-Puzzle v5.2.

**[0211]** The molecular clock hypothesis, or constant rate of evolution, was tested by using the likelihood ratio test with the likelihoods for the ML and clock-like ML trees obtained in PAUP\*. The clock was tested with the best-fitting evolutionary model estimated in Modeltest, and ML trees were constructed in PAUP\* starting from the NJ tree that is iteratively optimized using two consecutive heuristic searches

with nearest neighbor interchange followed by a final heuristic search with the tree-bisection-reconnection algorithm. To adjust for rate heterogeneity among different PTLV taxa, clock-like ML trees were then transformed into ultrametric trees using the nonparametric rate smoothing (NPRS) algorithm in the program TreeEdit (v1.0a10 carbon) (Sanderson (2003) *Bioinformatics*. 19:301-2). The branches of the NPRS tree were then scaled by using a divergence time of 40,000-60,000 years ago (ya) for the Melanesian HTLV-1mel lineage based on genetic and archaeological evidence of when the ancestors of indigenous Melanesians and Australians migrated from Southeast Asia (Lemey et al. (2005) *Infect. Gen. Evol.* 5:291-298; Salemi et al. (2000) *Mol. Biol. Evol.* 17:374-386; Salemi et al. (1999) *AIDS Rev.* 1:131-139). Variance in age estimates (branch lengths) was determined in PAUP\* with 100 bootstrap repetitions by enforcing topological constraints and using a heuristic search without branch swapping on the clock-like ML tree. Branch lengths in all 100 trees were calibrated as before and average divergence times and confidence intervals ( $\alpha=0.05$ ) were calculated in Excel. The evolutionary rate was estimated based on a known divergence time point of 40,000-60,000 ya and on the branch length of the ML clock-like tree according to the formula: evolutionary rate (r)=branch length (bl)/divergence time (t) (Van Dooren et al. (2004) *J. Gen. Virol.* 85:507-519).

#### Nucleotide Sequence Accession Number

**[0212]** The HTLV-3(2026ND) proviral sequence has the GenBank accession number DQ093792.

#### Example 3

##### Generation and Analysis of the Human T-Lymphotropic Virus Type 4 Complete Genome

**[0213]** The full-length genomic sequence of HTLV-4 (SEQ ID NO: 81) shown in FIG. 10 was obtained substantially as

**[0214]** HTLV-3(2026ND)-specific primers were then designed from sequences obtained in each of the four viral regions described above (tax, pol, env, and LTR), and used in nested, long-template PCRs to fill in the gaps in the genome using an expand high fidelity kit containing both Taq and Tgo DNA polymerases (Roche). PCR products were revealed on 1.5% agarose gels stained with ethidium bromide, purified with a Qiaquick™ PCR purification kit (Qiagen) and sequenced in both directions with a BigDye™ terminator cycle kit and automated sequencers (Applied Biosystems). Selected PCR products were also cloned into the pCR4-TOPO vector using the TOPO TA Cloning kit (Invitrogen) and recombinant plasmid DNA was prepared using the Qiagen plasmid purification kit prior to automated sequencing.

**[0215]** Percent nucleotide divergence was calculated by using the GAP program in the Genetic Computer Group's (GCG) Wisconsin package (Thompson et al. (1994) *Nucleic Acids Res.* 22:4673-4680). LTR RNA secondary structure was determined using the program RNAstructure v4.2 (Mathews et al. (1999) *J. Mol. Biol.* 288:911-940). Sequences were aligned by using the Clustal W program (Womble (2000) *Methods Mol. Biol.* 132:3-22), gaps were removed, and distance-based trees were generated by using the Kimura two-parameter model in conjunction with the neighbor-joining (NJ) method in the MEGA program (version 2.1) and maximum likelihood (ML) analysis in the PAUP\* program as described in detail elsewhere (Switzer et al. (2005) *Nature*. 434:376-380; Wolfe et al. (2005) *Proc. Natl. Acad. Sci. USA*. 102:7994-7999). The reliability of the final topology of the trees was tested with 1,000 bootstrap replicates. Table 7 shows a comparison of the genetic identity of the HTLV-3 and HTLV-4 full-length genomes with other PTLV prototypes. The stem loop structure necessary for Rex responsive control of viral expression in HTLV-1 and -2 was retained in HTLV-4(1863LE), and is shown in FIG. 11.

TABLE 7

Genetic Identity of HTLV-3 and HTLV-4 Genomes with other PTLV Prototypes (strain)									
	HTLV-1 (ATK)	STLV-1 (Tan)	HTLV-2 (MoT)	STLV-2 (PP1664)	STLV-3 (TGE2117)	STLV-3 (CTO604)	HTLV-3 (2026ND)	HTLV-3 (Pyl43)	HTLV-4 (1863LE)
HTLV-3									
2026ND	61.6	61.6	62.9	62.6	87.1	88.4	—	88.5	63.2
Pyl43	62.3	62.1	63.1	63.2	87.7	99.2	88.5	—	63.0
HTLV-4									
1863LE	62.0	62.0	70.7	70.8	63.5	63.1	63.2	63.0	—

described above in Example 2 for the identification of the HTLV-3 full-length genomic sequence. Briefly, DNA was prepared from uncultured PBMCs available from a subject identified in the original PTLV surveillance study in Cameroon reported in detail elsewhere (Wolfe et al. (2005) *Proc. Natl. Acad. Sci. USA*. 102:7994-7999). DNA integrity was confirmed by  $\beta$ -actin polymerase chain reaction (PCR) as previously described (Wolfe et al. (2005) *Proc. Natl. Acad. Sci. USA*. 102:7994-7999). To obtain the full-length genomic sequence of HTLV-3, small regions of each major coding region were PCR-amplified using nested PCR and degenerate PTLV primers.

**[0216]** For dating of HTLV-4(1863LE), full-length genomes from prototypical PTLVs available in GenBank were aligned with HTLV-3(2026ND) and HTLV-4(1863LE) essentially as described in Example 2 using Clustal W. The analysis, shown in FIG. 12, again inferred four major phylogroups with very high bootstrap support, confirming the genetic relationships that were based on smaller sequences. Both HTLV-3s again clustered with STLV-3s supporting a primate origin for these viruses. HTLV-4 again formed a new lineage distinct from PTLV-1, PTLV-2, and PTLV-3. However, the primate origin of HTLV-4 was less clear since there is not yet a known simian counterpart for this virus. These results also indicated the absence of genetic recombination in

PTLVs, which is a common mechanism that leads to increased genetic diversity of HIV.

**[0217]** The finding of HTLVs in four distinct clades indicated an ancient, independent evolution of these viruses. Thus, additional molecular analyses were performed to estimate the divergence times of the PTLV lineages. FIG. 13 shows the estimated divergence dates for the most recent common ancestor of HTLV-3(2026ND), HTLV-4(1863LE) and other PTLVs. Using the bovine leukemia virus (BLV) as an outgroup, a substitution rate of  $8.6 \times 10^{-7}$  to  $1.3 \times 10^{-6}$  substitutions/site/year for PTLV was inferred which is 3 logs lower than that seen in HIV, confirming the genetic stability of these deltaretroviruses.

**[0218]** Using these substitution rates, molecular dating inferred an ancient origin for PTLVs hundreds of thousands of years ago with the most recent common ancestor for each HTLV group ranging from 30,000 years ago for HTLV-2 to 456,000 years ago for HTLV-4. This finding contrasts with the more recent origin of HIV-1, which has been estimated to have occurred within the last century.

unique or similar to those of other HTLVs were identified (see Table 8). First, the genomes of HTLV-3 and HTLV-4 are shorter than HTLV-1 and HTLV-2 by having only two of three Tax response elements in the LTRs. However, the loss of this distal TRE has been shown to not significantly affect HTLV expression. In addition, only two TREs are present in STLV-3 and STLV-2 suggesting this difference is not a result of adaptation to a new host. Likewise, the finding of AP-1 and c-Myb transcription factors in place of the HTLV-3 or HTLV-4 LTRs is also not unique but are also present in STLV-3.

**[0222]** Overall, the HTLV-3 Tax protein contains many of the functional motifs important for viral expression and leukemogenesis attributed to HTLV-1 Tax. Detailed in vitro analysis confirmed that the HTLV-3 Tax was similar in function to the HTLV-1 Tax protein, suggesting a pathogenic potential in HTLV-3-infected persons like that observed in HTLV-1. The HTLV-3(Py143) genome is also shorter by a 366-bp deletion in the pX region that disrupts the HBZ reading frame suggesting a loss of Tax suppression and T-cell proliferation believed to be associated with this gene.

TABLE 8

Unique Genetic Features of HTLV Prototypes: HTLV-3 is more similar to HTLV-1					
	HTLV-1 (ATK)	HTLV-2 (MoT)	HTLV-3 (2026ND)*	HTLV-3 (Py143)**	HTLV-4 (1863LE)
Genome (bp)	9068	8952	8917	8553	8791
LTR (bp)	756	764	697	695	696
# LTR TREs	3	3	2 <sup>1</sup>	2 <sup>1</sup>	2 <sup>1</sup>
Other LTR TFs	—	—	AP-1	c-Myb	c-Myb
Tax transactivates	Yes	Yes	Yes	Yes	Yes
Tax localization	Nucleus	Cytoplasm		Nucleus	
Tax p53 inhibition	Yes			Yes	
PDZ BD in Tax	Yes	No	Yes	Yes	No
HBZ	Yes	No <sup>2</sup>	Yes	No <sup>3</sup>	Yes

<sup>1</sup>missing distal TRE

<sup>2</sup>HBZ is present in other HTLV-2

<sup>3</sup>366-bp deletion in pX

\*Switzer et al. J Virol. 2006; 80: 7427-38.

\*\*Calattini et al. J Virol. 2006; 80: 9876-88.

**[0219]** The inferred ancient origin for HTLV-3 and HTLV-4 indicates that exposure to these viruses may have been occurring for millennia, and thus these viruses may be more prevalent than currently known. Alternatively, HTLV-3 and HTLV-4 may represent more recent infections with highly divergent STLVs that have yet to be identified. This is probably the case for the HTLV-3(Py143) strain, since the high genetic identity of this virus to STLV-3RCM is similar to that seen in transmission pairs. Expanded surveillance of both humans and primates is warranted.

**[0220]** Changes in the molecular structure and genetic sequences of viruses has been proposed to play a role in the increased transmissibility and pathogenicity of viruses following cross-species transmission and adaptation to a new host. Thus, the genetic structure and sequences of HTLV-3 and HTLV-4 were examined to determine if the genome was intact and if important functional motifs involved in viral expression and HTLV-1-induced leukemogenesis are preserved. The Tax proteins of HTLV-3 were also characterized using in vitro assays to determine if motifs involved in Tax-mediated leukemogenesis were present and functioning.

**[0221]** While all structural and enzymatic proteins of both HTLV-3s and HTLV-4 were intact, features that are either

## REFERENCES

- [0223]** Araujo & Hall, Human T-lymphotropic virus type II and neurological disease. *Ann. Neurol.* 56, 10-19 (2004).
- [0224]** Barnhart et al., 1997. Function of the human T-cell leukemia virus type 1 21-base-pair repeats in basal transcription. *J. Virol.* 71:337-344.
- [0225]** Bindhu et al., 2004. Role of accessory proteins of HTLV-1 in viral replication, T cell activation, and cellular gene expression. *Front. Biosc.* 9:2556-2576.
- [0226]** Bowen-Jones & Pendry, The threat to primates and other mammals from the bushmeat trade in Africa, and how this threat could be diminished. *Oryx* 33, 233-246 (1999).
- [0227]** Busch et al., Absence of evidence of infections with divergent simian T-lymphotropic viruses in US blood donors with seroindeterminate human T-lymphotropic results. *Transfusion* 40, 443-449 (2000).
- [0228]** Callatini et al., 2005. Discovery of a new human T-cell lymphotropic virus (HTLV-3) in Central Africa. *Retrovirology*. 2:30.
- [0229]** Chevalier et al., 2005. Molecular characterization of the Tax protein from the highly divergent simian T-cell lymphotropic virus type 3 strain. *AIDS Res. Hum. Retrovir.* 21:513 (Abs. P174).

- [0230] Courgnaud et al., Simian T-cell leukemia virus (STLV) infection in wild primate populations in Cameroon: evidence for dual STLV type 1 and type 3 infection in agile mangabeys (*Cercocebus agilis*). *J. Virol.* 78, 4700-4709 (2004).
- [0231] Digilio et al., 1997. The simian T-lymphotropic/leukemia virus from *Pan paniscus* belongs to the type 2 family and infects Asian macaques. *J. Virol.* 71:3684-3692.
- [0232] Feuer and Green. 2005. Comparative biology of human T-cell lymphotropic virus type 1 (HTLV-1) and HTLV-2. *Oncogene.* 24:5996-6004.
- [0233] Gaudray et al., 2002. The complementary strand of the human T-cell leukemia virus type 1 RNA genome encodes a bZIP transcription factor that down-regulates viral transcription. *J. Virol.* 76:12813-12822.
- [0234] Gessain & Mahieux, Epidemiology, origin and genetic diversity of HTLV-1 retrovirus and STLV-1 simian affiliated retrovirus. *Bull. Soc. Pathol. Exot.* 93, 163-171 (2000).
- [0235] Goubau et al., A primate T-lymphotropic virus, PTLV-L, different from human T-lymphotropic viruses types I and II, in a wild-caught baboon (*Papio hamadryas*). *Proc. Natl. Acad. Sci. USA* 91, 2848-2852 (1994).
- [0236] Groves *Primate Taxonomy* (Smithsonian Institution Press, Washington, D.C., 2001).
- [0237] Hahn et al., 2000. AIDS as a zoonosis: scientific and public health implications. *Science* 287:607-614.
- [0238] Heneine et al., Identification of a human population infected with simian foamy viruses. *Nat. Med.* 4, 403-407 (1998).
- [0239] Khabbaz et al., Brief report: infection of a laboratory worker with simian immunodeficiency virus. *N. Engl. J. Med.* 330, 172-177 (1994).
- [0240] Leendertz et al., High variety of different simian T-cell leukemia virus type 1 strains in chimpanzees (*Pan troglodytes verus*) of Tai National Park, Cote d'Ivoire. *J. Virol.* 78, 4352-4356 (2004).
- [0241] Lemey et al., 2005. A Bayesian statistical analysis of human T-cell lymphotropic virus evolutionary rates. *Infect. Gen. Eva* 5:291-298.
- [0242] Lerche et al., Evidence of infection with simian type D retrovirus in persons occupationally exposed to nonhuman primates. *J. Virol.* 75, 1783-1789 (2001).
- [0243] Lole et al., 1999. Full-length human immunodeficiency virus type 1 genomes from subtype C-infected seroconverters in India, with evidence of intersubtype recombination. *J. Virol.* 73:152-160.
- [0244] Mahieux et al., Simian T-cell lymphotropic virus type 1 from *Mandrillus sphinx* as a simian counterpart of human T-cell lymphotropic virus type 1 subtype D. *J. Virol.* 72, 10316-10322 (1998).
- [0245] Mahieux et al., Molecular epidemiology of 58 new African human T-cell leukemia virus type 1 (HTLV-1) strains: identification of a new and distinct molecular subtypes in central Africa and in pygmies. *J. Virol.* 71, 1317-1333 (1997).
- [0246] Mahieux et al., 2000. Human T-cell lymphotropic virus type 1 gag indeterminate western blot patterns in Central Africa: relationship to *Plasmodium falciparum* infection. *J. Clin. Microbiol.* 38:4049-4057.
- [0247] Mathews et al., 1999. Expanded sequence dependence of thermodynamic parameters improves prediction of RNA secondary structure. *J. Mol. Biol.* 288:911-940.
- [0248] Meertens and Gessain. 2003. Divergent simian T-cell lymphotropic virus type 3 (STLV-3) in wild-caught *Papio hamadryas papio* from Senegal: widespread distribution of STLV-3 in Africa. *J. Virol.* 77:782-789.
- [0249] Meertens, et al., Complete sequence of a novel highly divergent simian T-cell lymphotropic virus from wild-caught red-capped mangabeys (*Cercocebus torquatus*) from Cameroon: a new primate T-lymphotropic virus type 3 subtype. *J. Virol.* 76, 259-268 (2002).
- [0250] Meertens et al., A. Molecular and Phylogenetic Analyses of 16 Novel Simian T Cell Leukemia Virus Type 1 from Africa: Close Relationship of STLV-1 from *Allenopithecus nigroviridis* to HTLV-1 Subtype B Strains. *Virology* 287, 275-285 (2001).
- [0251] Meertens et al., 2003. A novel, divergent simian T-cell lymphotropic virus type 3 in a wild-caught red-capped mangabey (*Cercocebus torquatus torquatus*) from Nigeria. *J. Gen. Virol.* 84:2723-2727.
- [0252] Milner-Gulland et al., Wild meat: the bigger picture. *TREE* 18, 351-357 (2003).
- [0253] Nerrienet et al., Simian T cell leukemia virus type I subtype B in a wild-caught gorilla (*Gorilla gorilla gorilla*) and chimpanzee (*Pan troglodytes vellerosus*) from Cameroon. *J. Gen. Virol.* 85: 25-9 (2004).
- [0254] Posada and Crandall. 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics.* 14:817-818.
- [0255] Rousset et al., 1998. The C-terminus of the HTLV-1 Tax oncoprotein mediates interaction with the PDZ domain of cellular proteins. *Oncogene.* 6:643-654.
- [0256] Salemi et al., Tempo and mode of human and simian T-lymphotropic virus (HTLV/STLV) evolution revealed by analyses of full-genome sequences. *Mol. Biol. Evol.* 17, 374-386 (2000).
- [0257] Salemi et al., A.M. Origin and evolution of human and simian T-cell lymphotropic viruses. *AIDS Rev.* 1, 131-139 (1999).
- [0258] Sanderson 2003. r8s: inferring absolute rates of molecular evolution and divergence times in the absence of a molecular clock. *Bioinformatics.* 19:301-2.
- [0259] Satou et al., 2006 HTLV-1 basic leucine zipper factor gene mRNA supports proliferation of adult T cell leukemia cells. *Proc. Natl. Acad. Sci. USA.* 103:720-725.
- [0260] Sharp et al., 2000. Origins and evolution of AIDS viruses: estimating the time-scale. *Biochem Soc Trans.* 28:275-282.
- [0261] Slattery et al., Genomic evolution, patterns of global dissemination, and interspecies transmission of human and simian T-cell leukemia/lymphotropic viruses. *Genome Res.* 9, 525-549 (1999).
- [0262] Switzer M et al., 2005. Ancient co-speciation of simian foamy viruses and primates. *Nature.* 434:376-380.
- [0263] Switzer et al., Phylogenetic relationship and geographic distribution of multiple human T-cell lymphotropic virus type II subtypes. *J. Virol.* 69, 621-632 (1995).
- [0264] Takemura et al., 2002. High prevalence of simian T-lymphotropic virus type L in wild Ethiopian baboons. *J. Virol.* 76:1642-1648.
- [0265] Thompson et al., 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22:4673-4680.

- [0266] Tsubata et al., 2005. PDZ domain-binding motif of human T-cell leukemia virus type 1 Tax oncoprotein is essential for the interleukin 2 independent growth induction of a T-cell line. *Retrovirol.* 2:46.
- [0267] Van Brussel et al., 1998. The simian T-lymphotropic virus STLV-PP1664 from Pan paniscus is distinctly related to HTLV-2 but differs in genomic organization. *Virology.* 243:366-379.
- [0268] Van Brussel et al., 1997. Complete nucleotide sequence of the new simian T-lymphotropic virus, STLV-PH969 from a Hamadryas baboon, and unusual features of its long terminal repeat. *J. Virol.* 7:5464-5472.
- [0269] Van Dooren et al., Evidence for a second simian T-cell lymphotropic virus type 3 in *Cercopithecus nictitans* from Cameroon. *J. Virol.* 2001 December; 75(23):11939-41.
- [0270] Van Dooren et al., Identification in gelada baboons (*Theropithecus gelada*) of a distinct simian T-cell lymphotropic virus type 3 with a broad range of Western blot reactivity. *J. Gen. Virol.* 85, 507-519 (2004).
- [0271] van Regenmortel et al., Seventh Report of the International Committee on Taxonomy of Viruses. (Academic Press, San Diego, Wien, New York, 2000). (online version <http://www.virustaxonomyonline.com/>)
- [0272] Vandamme et al., Use of a generic polymerase chain reaction assay detecting human T-lymphotropic virus (HTLV) types I, II and divergent simian strains in the evaluation of individuals with indeterminate HTLV serology. *J. Med. Virol.* 52, 1-7 (1997).
- [0273] Wolfe et al., 2005. Emergence of unique primate T-lymphotropic viruses among central African bushmeat hunters. *Proc. Natl. Acad. Sci. USA.* 102:7994-7999.
- [0274] Wolfe et al., 2004. Exposure to nonhuman primates in rural Cameroon. *Emerg. Infect. Dis.* 10, 2094 (2004).
- [0275] Wolfe et al., Naturally acquired simian retrovirus infections in central African hunters. *Lancet* 363, 932-937 (2004b).
- [0276] Wolfe et al., Simian retroviral infections in human beings. *Lancet* 364, 139-140 (2004c).
- [0277] Womble 2000. GCG: The Wisconsin Package of sequence analysis programs. *Methods Mol. Biol.* 132:3-22.
- [0278] Yamashita et al., Molecular Epidemiology of HTLV-I in the world. *J. Acquir Immune Defic Syndr Hum Retrovirol.* 13: S124-31 (1996).
- Sequences
- [0279] SEQ ID NO: 1 (HTLV-3 pol) (pos 2407-5076)
- [0280] SEQ ID NO: 2 (HTLV-4 pol) (pos 3-2549)
- [0281] SEQ ID NO: 3 (HTLV-3 env) (pos 5069-6544)
- [0282] SEQ ID NO: 4 (HTLV-4 env) (pos 2542-3999)
- [0283] SEQ ID NO: 5 (HTLV-3 tax)
- [0284] SEQ ID NO: 6 (HTLV-4 tax)
- [0285] SEQ ID NO: 7 (PTLVTPG)
- [0286] SEQ ID NO: 8 (PGTAXR1)
- [0287] SEQ ID NO: 9 (PH2Rrev)
- [0288] SEQ ID NO: 10 (PGTAXR2)
- [0289] SEQ ID NO: 11 (PGPOLF1)
- [0290] SEQ ID NO: 12 (PGPOLR1)
- [0291] SEQ ID NO: 13 (PGPOLF2)
- [0292] SEQ ID NO: 14 (PGPOLR2)
- [0293] SEQ ID NO: 15 (1863TF1)
- [0294] SEQ ID NO: 16 (1863TR1)
- [0295] SEQ ID NO: 17 (1863TF2)
- [0296] SEQ ID NO: 18 (1863TR2)
- [0297] SEQ ID NO: 19 (5VLTRExt)
- [0298] SEQ ID NO: 20 (1MNDR1)
- [0299] SEQ ID NO: 21 (Enh280)
- [0300] SEQ ID NO: 22 (1MNDR2)
- [0301] SEQ ID NO: 23 (PH1F)
- [0302] SEQ ID NO: 24 (PH2R)
- [0303] SEQ ID NO: 25 (PH2F)
- [0304] SEQ ID NO: 26 (PH2R)
- [0305] SEQ ID NO: 27 (PGENVF1)
- [0306] SEQ ID NO: 28 (PGENVR1)
- [0307] SEQ ID NO: 29 (PGENVF2)
- [0308] SEQ ID NO: 30 (PGENVR2)
- [0309] SEQ ID NO: 31 (GPLTRF1)
- [0310] SEQ ID NO: 32 (GPLTRR1)
- [0311] SEQ ID NO: 33 (GPLTRF2)
- [0312] SEQ ID NO: 34 (GPLTRR2)
- [0313] SEQ ID NO: 35 (HTLV-3 gag) (pos 756-2023)
- [0314] SEQ ID NO: 36 (HTLV-3 Complete genome: 2026ND.seq (8917 bp))
- [0315] SEQ ID NO: 37 (HTLV-3 env amino acid)
- [0316] SEQ ID NO: 38 (HTLV-3 env surface antigen (SU)=aa 1-315)
- [0317] SEQ ID NO: 39 (HTLV-3 env transmembrane=aa 316-491)
- [0318] SEQ ID NO: 40 (HTLV-3 gag amino acid)
- [0319] SEQ ID NO: 41 (HTLV-3 gag p15=aa 337-422)
- [0320] SEQ ID NO: 42 (HTLV-3 gag p19=aa 1-123)
- [0321] SEQ ID NO: 43 (HTLV-3 gag p24=aa 124-336)
- [0322] SEQ ID NO: 44 (HTLV-3 pol) amino acid
- [0323] SEQ ID NO: 45 (HTLV-3 LTR) (pos 1-697 & 8221-8917)
- [0324] SEQ ID NO: 46 (HTLV-3 pro) amino acid
- [0325] SEQ ID NO: 47 (HTLV-3 pro) (pos 1976-2509)
- [0326] SEQ ID NO: 48 (HTLV-3 rex) amino acid
- [0327] SEQ ID NO: 49 (HTLV-3 rex)(pos 5010-5071. 7245-7730)
- [0328] SEQ ID NO: 50 (HTLV-3 tax) amino acid
- [0329] SEQ ID NO: 51 (HTLV-3 tax) (pos5069-5071. 7244-8293)
- [0330] SEQ ID NO: 52 (HTLV-3 pX (pos 6545-7243)
- [0331] SEQ ID NO: 53 (HTLV-4 pol-env-tax region)
- [0332] SEQ ID NO: 54 (HTLV-4 env) amino acid
- [0333] SEQ ID NO: 55 (HTLV-4 env surface antigen (SU)=aa 1-307)
- [0334] SEQ ID NO: 56 (HTLV-4 env transmembrane=aa 308-485)
- [0335] SEQ ID NO: 57 (HTLV-4 pot) amino acid
- [0336] SEQ ID NO: 58 (HTLV-4 pro) amino acid
- [0337] SEQ ID NO: 59 (HTLV-4 pro) (pos 1-273)
- [0338] SEQ ID NO: 60 (HTLV-4 rex) amino acid
- [0339] SEQ ID NO: 61 (HTLV-4 rex) (pos 2483-2545. 4560-5009)
- [0340] SEQ ID NO: 62 (HTLV-4 tax) amino acid
- [0341] SEQ ID NO: 63 (HTLV-4-pX) (pos. 4000-4558)
- [0342] SEQ ID NO: 64 1863 PF1
- [0343] SEQ ID NO: 65 1863PR2
- [0344] SEQ ID NO: 66 1863PP2 FAM (fluorescent labeled probe)
- [0345] SEQ ID NO: 67 region of HTLV-3 where type specific peptides of HTLV-1 and HTLV-2 are located
- [0346] SEQ ID NO: 68 region of HTLV-4 where type specific peptides of HTLV-1 and HTLV-2 are located
- [0347] SEQ ID NO: 69 2026LF1
- [0348] SEQ ID NO: 70 2026PR1

[0349]	SEQ ID NO: 71 2026LF2	[0356]	SEQ ID NO: 78 2026LR1
[0350]	SEQ ID NO: 72 2026PR2	[0357]	SEQ ID NO: 79 2026EF3
[0351]	SEQ ID NO: 73 2026 PF3	[0358]	SEQ ID NO: 80 2026LR2
[0352]	SEQ ID NO: 74 2026ER1	[0359]	SEQ ID NO: 81 (HTLV-4(1863LE) Complete genome)
[0353]	SEQ ID NO: 75 2026 PF2	[0360]	SEQ ID NO: 82 (GTAAAC)
[0354]	SEQ ID NO: 76 2026ER3	[0361]	SEQ ID NO: 83 (TTAAAC)
[0355]	SEQ ID NO: 77 2026EF1	[0362]	SEQ ID NO: 84 HTLV-3 HBZ

---

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 84

<210> SEQ ID NO 1

<211> LENGTH: 2670

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 1

```

ggtcctccct gtctagcac ccaagcttat cggattagag caccttcccc cgccccaga    60
agttctctcag ttcccgtaa accagagcgc ctccaggctc tgactgacct ggtttccagg    120
gccttgagg ctaaacatat agaacctat caaggaccag gcaataacc tatcttcccc    180
gtcaaaaagc ccaatgggaa gtggcggttt attcatgatc tccgggccac aaactccgctc    240
acccgagacc tggcttcacc gtcccccgcc cctcggacc ttaccagtct gcccgaaggc    300
ctccccacatc ttcggacaat tgaccttact gacgccttct tccaaatccc gctaccaacc    360
atTTTTcagc cgtactttgc attcactctc ccccaacoga acaactatgg tcccgaacc    420
agatactctt ggagagtact accccagggg ttcaaaaata gtccaacttt atttgaacag    480
caactctccc atatacttac cctgtgctgg aaaaacttcc ctaattccct tattatacaa    540
tatatggatg acattctact ggccagcccc gcccccgcg agctagctgc tctcaccgat    600
aaagtaacaa acgccttaac aaagggaaggc ctgcccctat ctccgaaaa gactcaggcc    660
actcctggtc ccatacattt tcttgacaaa gtcatatccc aggactgcat aacatacgag    720
accctcccat ccatcaatgt aaaatccacc tggctactgg cagaactaca gtccatgcta    780
ggagaattac aatgggtctc taaaggcacc cctgtctctc gctcctctct acaccagctc    840
tatctcgctc tccgaggcca tctgaccctc cgggatacta taaaattaac ctcaatacag    900
gtacaagctc taagaactat tcagaaggcc ttgaccctaa actgcccgaag tagactagta    960
aatcagctgc ccatcttggc ccttataatg ctccggccta caggcaccac ggcagtctc    1020
ttccaaacaa aacaaaagtg gccactcgtc tggttacaca ccctcaccg ggtaccagc    1080
ttgccccatc ggggacaact attggccaac gctgtcatca tcttagacaa atactcacta    1140
caactatg gccaaagtat caaatcattt catcataaca tctctaatca agcctcact    1200
tactacttac atacatctga ccaatccagc gtagccatcc tattacaaca ctcacacaga    1260
tttcacaatc tcggggcgca gccatcaggg ccatggagaa gcctcttaca aatgccccag    1320
atTTTccaaa acatcgatgt cctaaggcct ccttttaoca tctcgctgt agtcatcaac    1380
catgccccct gtctcttttc ggacgggtct gcctccaagg cagcatttat catctgggat    1440
agacaagtca tccaccaaca ggtcctctcc ctgcccctaa cctgttcagc tcaagcgggg    1500

```

-continued

---

```

gaactatttg gcctgttggc aggactacaa aagtctcaac cctgggtggc actaaatata 1560
tttctagatt caaaatttct tctcgggcac ctcaggcgaa tggcattagg ggcctttcca 1620
ggaccatcca cccagtgcca attacacaca cagcttctcc cctgctgca gggaaagact 1680
gtctatgtgc accacgtaag gagtcacact cttctgcaag accctatata cgccttaat 1740
gaagccacag atgccctcat gcttgccct ctgttacctc tcgacccac aactctccac 1800
cagctcacc actgtaacc ttacgccta cgtaacctg gagccactgc cctggaagcc 1860
catgcaattg tgcaggcatg tcacacatgc aaggctcatca acccagagg acggttacct 1920
cagggatata ttctcagagg ccatgcccct aatgatatac ggcaagggga cgtcactcac 1980
ctccagtaca agcgatacaa atattgcctg ctggtttggg tagatactta ctccggggcg 2040
gtctctgtgt cctgcccggc caaggagacc ggctccgact gtgtcgcctc gttacttctg 2100
gccatctcca ttctagaaa accacaaaa atcaacaccg ataacggagc tgcgtatctc 2160
tccaagagt tccaacagtt ttgcaactca cttgccatta aacattcaac ccacattccc 2220
tacaaccca ccagctctgg cttgggtgaa agaacaaatg gtattcttaa aactttgatt 2280
tccaaatata tcttagaaa tcaccacctg ccctggaga cagccgtctc caaatccctc 2340
tggactataa accatcttaa tgcctctccc tcgtgcaaaa agactcggtg gcagctacat 2400
cagggccagc cctgctctcc cgtccctgaa gacacactcc caccacacac atcaccaaaa 2460
tggattatt ataaaaattc tggctttacc aactcaaggt ggagtgggcc tgtacaatcc 2520
ctcaaggaag cagcaggagc ggctctcacc ccagtaggtg gaagctatct ctggatcccg 2580
tggcgcctcc taaaaggggg tatatgccc agaccgaaa gcagcgcagc cgtcgacca 2640
aaaaccagag accatcaact ccatgggtaa 2670

```

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 2549

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

&lt;400&gt; SEQUENCE: 2

```

ctggccatat cgaaccttac tctggaccag gcaacaaccc agttttccct gttaaaaaac 60
ccaacggcaa gtggcgattt atccatgacc tcagggccac taatgccatc accactacc 120
ttgctcggcc ctccccggc cccctgacc ttaccagcct gccacaggcc ttgcccac 180
ttcagaccat cgatctcagc gacgctttct tccagattcc cctccaaaag cgattccagc 240
cctacttcgc ctttaccatc cccagccat taaatcatgg gcctgggagc aggtacgctt 300
ggacagtcct tcccaaggc ttcaaaaaca gcccacgct ctttgagcaa cagctggcca 360
gcgtactagg cccagcccga aaagccttcc ccacatccgt catcgtccaa tacatggagc 420
acatcctctt ggcattgccc tcccagcagc aactagatca gctggccacc cttaccgcac 480
agctattgtc ctctcatggt ctcccagttt cccagaaaa aacccaacgc accccaggaa 540
aaatacactt cctgggcca atcatacatc cagatcacat cacctatgaa accacccca 600
ccatccccat taaggcacac tggaccctga ctgaaactgca aacctcctg ggggagctcc 660
agtgggtctc caaggggact cctgtcctcc gagaacacct tcaactgtctc tactcagcct 720
tgagaggtct caaagacccc cgggacacta tcacctctg tcatcctcac ctccagctc 780

```

-continued

---

```

tccacaacat tcagcaagcc ctgcatcaca attgcecgcg tgccttgac tetaagctcc 840
ccctccttgg cctcatcttc ctcagtcctat ccggcacgac ctcagtcctc tccagacaa 900
atcataaatg gccctagtc tggctccacg cccccatcc cccgaccagc ctatgccct 960
gggggcacat actgcctgc actgtactta cccttgacaa gtatgccttg cagcactatg 1020
gccaaactatg caaatcattc catcataaca tgtccacca ggccctacac gatttcgtaa 1080
aaaattctc tcaccccagc gtgcctat taattacca catgcatcgg ttctgtgatc 1140
tgggcagaca gccaccggga ccctggcgaa ccctcttaca actcccgcc cttctccggg 1200
aaccacagct cctcaggcct gcattttccc tatccccagt ggttatagat caggccctt 1260
gtctgttctc tgatgggtct ccccaaaagg ccgcctatgt aatttgggac aaggtcattc 1320
tcagccagcg gtcggtcccc ctgccccccc atgccaataa ctcagcacia aagggggaat 1380
tagtcggact cctcttgggc ttgcaagccg cacagccctg gccatcctt aacattttcc 1440
tagactcaaa gttcctcctc cggtaacctc agtccctcgc ttccggggcc tccaaggat 1500
catccacaca ccaccgtctc caggcgtccc tgcccacact cctccagggc aaggctcgtg 1560
atctccacca caaccgcagc cacaccat tgctgatcc catctcgacc ctcaatgaat 1620
ataccgactc tctcattgtc gcccccgtaa ccccttgaa gctgagggc ctccatgcc 1680
tcaccactg caaccaacg gccctcgttt cccacggagc caccctgca caggctaagc 1740
aactcgtgca ggctgcccgc acctgtcaaa tcattaacco tcaacaccac atgcccgtg 1800
gccacatccg ccgcccgcac ttcccaaac acacatggca aggagatgc acccacctta 1860
agcacaacg gaaccgatac tgcctccacg tctgggtgga taccttctca ggtgcggtat 1920
cttgtgtctg caaaaagaaa gaaactagca ggcacctat caaaacctc ctacatgcca 1980
tctccgtgct aggcaagccc ttctctgtta acacggacaa tggaccgct tacctttctc 2040
aggagtcca cgaattctgt accaccctc gcatcaaa caaccaccat attcctaca 2100
atccgacaag ttcaggcctg gtggagcgca caaatggcat tctcaagaca ctactataca 2160
aatatttct agaccacct gacctcccc tagaaagcgc ggttcaaa gctctctgga 2220
ccattaacca tttaaatgtc atgcgccct gtggaagac tcggtggcag ctccatcaca 2280
cccccccct gctcctatt tccgagtcca tacaaccac tcccaccagg ctacattggt 2340
actattaca aacctctgga cttaccaacc agcgatggaa agggcccga caatctctcc 2400
aggaagcagc aggagcagct ctcttcaag tcagtgcagg ctgccccag tggatccct 2460
ggcggctcct gaagaagact gtatgccaa aaccgacga ccccgaccc gcagggcagc 2520
tcgaaacaga ccaccaacac catgggtaa 2549

```

&lt;210&gt; SEQ ID NO 3

&lt;211&gt; LENGTH: 1476

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

&lt;400&gt; SEQUENCE: 3

```

atgggtaagt tctgccttta tttctgtctt atttacatac tctctctgc ctctctggc 60
aatcccagtc gatgcacct gttcatagga gcctcttctt accactccga cccctgegg 120

```

-continued

---

```

tccgatcacc cgcggtgtac ctggagactc gacctatfff ctctcacaag agatcaaage 180
ctaagccccc catgtccaga cttagttact tactcacagt atcataggcc ctactccctg 240
tatgtattcc cccattggat aaccaaaacct aaccgtcgag gcctagggta ctattctgct 300
tcctactcag acccctgccc tatacaatgc ccttacttag gatgccagtc atggacatgt 360
ccttatacag gcccggtgtc cagcccacat tggaaatact cctccgatct taattttacc 420
caagaagtat catccatctc cctacacttg cacttttcca aatgtgggtc ctcattctcc 480
tttctactag acgcaccagg atatgatcca gtgtggttcc tctcctccca ggccacacag 540
gccccaccca cgccctcccc tctcatacag gactcagatc tocaacatat cctagaacct 600
tccateccct ggagctccaa aatcctcaat ctcatcctcc ttaccctaaa aagctctaata 660
tattcttgca tggctctgtg tgaccgctcc agcctatctt cgtggcatgt tctatatgac 720
ccacttaaag ccccaatcc acccgacccc aaagcccagt ctattctgcg accctcccta 780
gccattcccc ccagtaatgt caccocgcca tttccttgga cccattgcta tcgccccctt 840
ctacaggcca tctcctcgga aactgcaat aactccgtag tactgcccc cttttccctg 900
tccccacttc ctaacgtctc cagaccccga aagcgcggag cagtcccat tgccatatgg 960
ctagtgtccg cctagcggc cggaaccggg atagctggcg gagttaccgg ctccctgtct 1020
ctggcctcca gtaaaagcct actgcgcgaa gtagaccagg atatagatca cctgacgcag 1080
gcgattgtaa aaaacatga taataccctt cgggttgctc aatacgcagc ccaaaatcgc 1140
cgtggcctag atttactttt ctgggaacaa gggggccttt gtaaggccat ccaagaacaa 1200
tgttgcttcc ttaatatcag caacacccat gtgtccgtcc tcacggaaag acccccccta 1260
gaaaagagag taatcactgg ctggggactc aactgggacc tcgggctctc ccaatgggcc 1320
cgagaggccc ttcagacagg tataaacctt ttagcccttt ttctcctcct catcgttgta 1380
ggaccctgcg tcatacgcca gctacaggcc ctcccttccc gtctgcagca tcgcagtcag 1440
ccctactccc ttctcaatta tgaaccaaac ttataa 1476

```

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 1458

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

&lt;400&gt; SEQUENCE: 4

```

atgggtaacg tactcttctt aactttattg gccaccctgg gcacccagc acttcaggcc 60
agccggtgta caatcacggt aggtatctcc tctaccact ccagcccctg cagcccagcc 120
cagcctttat gtacctgggc cctcgacctt gtgtccatca ctaaggacca gctcctctac 180
ccccctgcc aaaacctgat cacctattcc aactaccaca agacctactc cctgatatcc 240
ttcccacact gggtaaaaa gccactccgc cgggggcttg gatactactc agcctctac 300
tctgatcctt gctccctaca atgtccctac ctagggaagtc aatcatggac ttgccctat 360
actggcccctg tctcgagccc aacttgaga ttctccacag atgtaaattt cacccaagaa 420
gtcagccgtg tctccctaaa acttcatttc tccaaatgtg gttcctcctt aactctggtta 480
atagatgccc ccggttacga tccgctgtgg tactccacat ccgagcctac tcaggaaccc 540
ccaacccctc cgccactagt cagcgactca gacctagagc atgtcctgac tcctteggcc 600

```

-continued

---

```

tcttgggct ccaagatgct gacctcacc cacctaacct tgcagagcac caactattcc 660
tgtatggtct gtattgaccg cgccagcctc tcttctggc acgtattata cactcccaac 720
atctctagta atgccccctc aaaaccacc gtccgccctt cccttgccct atccgccccg 780
cgaccacagc cttccccctg gaccattgc tatcaaccac aggtgcaagc tgtaaccacc 840
gcaaaagtca ataattccat catacttccc ccattttctc tctctccctt gcttgggtgc 900
cctctcacta ggccagcccg ggccgtccc gtggcggtct ggctcgttc cgtttggcc 960
gcagggacag gaatagcagg aggtgtcacc ggtccttat ccctggcctc cagtagaagt 1020
ctctgtccg aagtggacaa ggatatttc cacctcacac gggccattgt aaaaaaccac 1080
caaacattc ttgagtgcc ccaatatgcc gcccaaaaca ggcgagggtt agacctcctg 1140
ttctgggaac aaggggggct gtgtaaagcg atacaagaac aatgtgctt cctcaacac 1200
agcaataccc atatttcagt cttacaagag cgacccccctc tagaaactcg ggtaactact 1260
ggatggggct taaattggga tctaggactc tcccagtggt cccgtgagge tctccagact 1320
ggtattacc ttttggccct ccttctgta atcatcatcc tggggcctg cattattcgc 1380
cagctgcaag cctccccca gaggetacag cagcgacctg accagtacc tctcctaac 1440
cctgagacc cttataa 1458

```

&lt;210&gt; SEQ ID NO 5

&lt;211&gt; LENGTH: 1053

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

&lt;400&gt; SEQUENCE: 5

```

atggccatt tcccagggtt cggacagagc cttctctacg ggtaccctgt ctacgttttc 60
ggcgactgtg tacaggccga ttggtgcccc atttctgggg ggctttgttc cgctcggtca 120
caccgccatg ccctactggc cactgtcccc gaacatcaga ttacctggga ccccatcgat 180
ggacgggttg tcagctcagc tctacaatac cttatccctc gactccccctc cttccccacc 240
cagagaacta cccgcacctc caaggttctc acccccccaa ccactgctgc gacccccaaag 300
attcctccat cttcttcca cgccgttaa aaacacacc cctccgaaa caattgcctt 360
gaactcacc tgggagagca gttgccagcc atgtccttcc cgaacctgg gctccgacce 420
caaaacatct acaccatgtg ggaagctcc gttgtgtgcc ttaactcta tcagctctcc 480
cccccatga cctggcctct aatcccgcct gttatattct gccatcctga gcagcttggga 540
gccttctca cccgagtccc tacaaaaga ttagaagaac tctgtataa gatattttta 600
agcacagggg cgataatcat cctgctgaa aactgttttc caaccacct gttccaacc 660
accgcgcgc ccgcggtgca gggccccctg cacacaggcc tgetcccgtg tcaaaaggaa 720
attgctacc ccgggtcat ttggactttc actgatggca gcccctgat tccggccct 780
tgccccaaag aaggacagcc atctttagta gtacaatcat ctacattat ctttcaacaa 840
ttccaaacca aggccagtca ccccgtttc ctcttctccc aaaaactaat ccaactctcc 900
tcttttcatt cctccacct cctctttgag gaatatacaa ctatcccctt ttctctactt 960
tttaatgaaa aaggggcaaa tgtcgtgat gatgagcccc gagacgggtc acaaccacca 1020

```

-continued

---

gctagaggac aaatagctga gtcacccgctc tga 1053

<210> SEQ ID NO 6  
 <211> LENGTH: 765  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

&lt;400&gt; SEQUENCE: 6

```

atggcccact tcccaggatt cgggcagagc ctctctatg gataccccgt ctatgtgttt    60
ggcgattgtg ttcaagccga ttggtgcccc atctccggtg gattatgctc cccccgcta    120
catcgccacg cctctctggc cacctgcccc gagcaccaga tcacctggga ccccatcgat    180
ggacgagtgt tgggctcgcc tctccaatac cttatccctc gctcccctc cttccccacc    240
caacgaacct ccaagacct caaagtcctt accccaccaa cactcctgt ccccccaag    300
gttccacct cettetttca gtcctgctgg aggcacagcc cctaccgcaa cggatgtctt    360
gaaacaacc ttggagagca gctcccctcc cttgcatttc ctgagccagg cctcaggccc    420
caaacgtct acaccatctg gggaaagacc atagtgtgtc tatacatcta ccagctgtcc    480
cctccatga cctggcccct cattcccct gtcataatgtt gcaaccccag gcagcttggc    540
gcttttctaa gcaatgtgccc cccaagcga ttagaagaac tctctacaa actttatcta    600
cacaccggcg ccaataatc cctgcccggaa gacgcctgc ctaccacct atttcagcct    660
gttcgagcac cctgtgtcca aactacctgg aacacaggac ttctcccata ccagccaaac    720
ctgactaccc ctggcctgat atggaccttt aatgatgggt ctctt                    765

```

<210> SEQ ID NO 7  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

&lt;400&gt; SEQUENCE: 7

tyacctrugga ccccatcgat ggacg 25

<210> SEQ ID NO 8  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)...(26)  
 <223> OTHER INFORMATION: n = inosine

&lt;400&gt; SEQUENCE: 8

gangaytгна stacyaaaga tggctg 26

<210> SEQ ID NO 9  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;



-continued

---

<211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
           note=synthetic construct  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)...(24)  
 <223> OTHER INFORMATION: n = inosine  
  
 <400> SEQUENCE: 14  
  
 gryrgngn gtn cctttngaga ccca 24  
  
 <210> SEQ ID NO 15  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
           note=synthetic construct  
  
 <400> SEQUENCE: 15  
  
 ctccttcttt cagtcggtgc ggag 24  
  
 <210> SEQ ID NO 16  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
           note=synthetic construct  
  
 <400> SEQUENCE: 16  
  
 ggggtagtca ggtttggctg gtat 24  
  
 <210> SEQ ID NO 17  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
           note=synthetic construct  
  
 <400> SEQUENCE: 17  
  
 cctaccgcaa cggatgtctt gaaa 24  
  
 <210> SEQ ID NO 18  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
           note=synthetic construct  
  
 <400> SEQUENCE: 18  
  
 tatggcgccg gtgtgatgat aaag 24  
  
 <210> SEQ ID NO 19  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
           note=synthetic construct

---

-continued

---

<400> SEQUENCE: 19  
aaccacccat ttctcccga tg 22

<210> SEQ ID NO 20  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 20  
gtcgtgaatg aaagggaaaag ggg 24

<210> SEQ ID NO 21  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 21  
tgacgacaac cctcacctc aa 22

<210> SEQ ID NO 22  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 22  
aggggtggaa ctttcgatct gtaa 24

<210> SEQ ID NO 23  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 23  
ttgtcatcag cccacttccc agg 23

<210> SEQ ID NO 24  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 24  
aaggagggga gtcgaggat aagg 24

<210> SEQ ID NO 25  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;

-continued

---

note=synthetic construct

<400> SEQUENCE: 25

cccaggtttc gggcaaagcc ttct 24

<210> SEQ ID NO 26

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 26

aaggagggga gtcgaggat aagg 24

<210> SEQ ID NO 27

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)...(23)

<223> OTHER INFORMATION: n = inosine

<400> SEQUENCE: 27

tggatcccgt gmgnytcct naa 23

<210> SEQ ID NO 28

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)...(24)

<223> OTHER INFORMATION: n = inosine

<400> SEQUENCE: 28

gtrtangrs angtccangm ytg 24

<210> SEQ ID NO 29

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)...(24)

<223> OTHER INFORMATION: n = inosine

<400> SEQUENCE: 29

anagaccwyc aacwccatgg gtaa 24

<210> SEQ ID NO 30

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

---

```

<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(23)
<223> OTHER INFORMATION: n = inosine

<400> SEQUENCE: 30

gmytggcanc cnargtangg gca 23

<210> SEQ ID NO 31
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(27)
<223> OTHER INFORMATION: n = inosine

<400> SEQUENCE: 31

rccaccanct nngggacaaa tagctga 27

<210> SEQ ID NO 32
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

<400> SEQUENCE: 32

cygggccaag cctcgctgca ggca 24

<210> SEQ ID NO 33
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(24)
<223> OTHER INFORMATION: n = inosine

<400> SEQUENCE: 33

accnnggctc tgacgtctct ccct 24

<210> SEQ ID NO 34
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(24)
<223> OTHER INFORMATION: n = inosine

<400> SEQUENCE: 34

ggcagnagaa gtgctacttt cgat 24

```

-continued

---

```

<210> SEQ ID NO 35
<211> LENGTH: 1268
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
      note=synthetic construct

<400> SEQUENCE: 35
atgggaaaga cttatagctc cccaataaac cctatcccca aagccccaaa ggggctagca    60
attcaccact ggetgaactt cctccaggct gcgtaccgac tgcagccagg gccttctgaa    120
ttcgatttcc accagttacg aaagtttcta aaacttgcta ttaagacccc ggtatggtta    180
aatcccatta attactctgt cctcgccgga ctcaccccaa aaaactaccc cggcagggtt    240
catgaaatag tggccatcct aattcaagag acccctgcac gggaggcgcc cccgtcagct    300
ccgctagcag aggacctca aaagcctcca ccctatcccg agcaggcgca ggaggcatct    360
cagtcctcc  coactctca ccccatggg gcccagccg ctcacggcc ctggcaaatg    420
aaggatctcc aggctattaa acaggaagtc agctcttccg cccctggtag ccccagttc    480
atgcagacta tacgcttggc tgtccagcaa ttgatccca cagcaaaaga tctccacgat    540
ctcctacagt acctgtgctc ttccttagtt gcctcctgc accatcagca acttgagacc    600
ctcatagctc agggcgaaac ccaaggtata acaggatata acccctggc cggcccctta    660
cgaatacagg ccaacaatcc aaatcaacaa gggctccgaa aagaatatca gaacctgtgg    720
ttatcggcct tttccgccct cccggggaac accaaggacc ccacctgggc agctatcctc    780
cagggacctg aagaaccctt tggctctttt gtagaaagac tcaatgtggc tttagataat    840
ggccttcccg aaggaaaccc caaagatcca atccttaggt ccctcgcta ttcaaatgct    900
aacaaggagt gccaaaaact cctacaggcc cgaggacaaa ccaacagccc gctaggggaa    960
atgctcaggg cctgccaaac ttggacgccc cgagataaaa acaaaatact aatggtacaa   1020
cctaaaaaaa ctctcccc  gaaccagcca tgcttccgct gcgggcaagt aggtcattgg   1080
agcagagatt gtaaacagcc tcggccccct ccgggcccc gcccgtgtg tcaggatccc   1140
accactgga agcgggactg cccacagtta aaaacagata ccagagacag cgaggaccta   1200
ctcctagacc tgccctgtga agcaccat  gtccgggaac gaaaaaac  ctcagggggg   1260
gaggatta                                     1268

```

```

<210> SEQ ID NO 36
<211> LENGTH: 8917
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
      note=synthetic construct

<400> SEQUENCE: 36
tgtcgatgat gatgagcccc gagacgggtc acaaccacca gctagaggac aaatagctga    60
gtcacccgtc tgagaaccgt ctcacaccgg gattgtgcc aaaaagaaca ccggggctct    120
gacgtctctc cctaccctgg ctcccgaaa aaacaaaaa ccaccattt cctcatgttt    180
gcctaaagct ctgacgataa ccctaaaaa ttgactagc aaataaagaa ccctgggccc    240
tataaaaggg gagagcaacc taaaatggg atcccttttc tgcaacctgc caaccctcc    300
tctggccaag gtccgacttt ggtcattcct gcctacctga atcgccgctt cgggatcgag    360

```

-continued

---

ccatccctct	tctatttgg	ggcaacttgc	gcactccgc	gccttccact	cgtaagatc	420
ccactgggtc	gagctaggcc	atcaccctg	ggccgctccc	ctggagctct	ctcgcgccc	480
tcttaagggt	gtccccctc	agcaaaaggc	ccagggttt	ctctacttcc	ttgtttcaag	540
tctctttctt	tggcggctga	cctaaatcga	aagtagcact	tctgctgtca	gcagcgaggc	600
ttggcccagg	gccagcgctt	gtaagggtac	ccagctcgga	gttgggtctc	tagagaaatca	660
gggctaaagc	tgctagcctt	aggaagaag	gcaaacaggt	gggggctcgt	ccgggattga	720
tcacctctct	gtatttgccc	ttccctgtcg	aagccatggg	aaagacttat	agctccccc	780
taaaccttat	ccccaaagcc	ccaaaggggc	tagcaattca	ccactggctg	aacttcctcc	840
aggctgcgta	ccgactgcag	ccagggcctt	ctgaattcga	ttccaccag	ttacgaaagt	900
ttctaaaact	tgtattaag	accccggtat	ggttaaatcc	cattaattac	tctgtcctcg	960
ccggactcat	ccccaaaaac	taccocggca	gggttcatga	aatagtggtc	atcctaattc	1020
aagagacccc	tgccagggag	gcgccccgt	cagctccgct	agcagaggac	cctcaaaagc	1080
ctccacccta	tcccagcag	gcgcaggagg	catctcagtg	cctccccatc	cttcaccccc	1140
atggggcccc	agccctcat	ggccctggc	aatgaagga	tctccaggct	attaacagg	1200
aagtcagctc	ttccgcccct	ggtagcccc	agttcatgca	gactatacgc	ttggctgtcc	1260
agcaatttga	tcccacagca	aaagatctcc	acgatctcct	acagtactcg	tgctcttctt	1320
tagttgctc	cctgcaccat	cagcaacttg	agaccctcat	agctcaggcg	gaaacccaag	1380
gtataacagg	atataacccc	ctggccggcc	ccttacgaat	acaggccaac	aatccaaatc	1440
aacaagggct	ccgaaaagaa	tatcagaacc	tgtggttato	ggccttttcc	gcctccccgg	1500
ggaacaccaa	ggacccccac	tgggcagcta	tcctccaggg	acctgaagaa	ccctttggct	1560
cttttgtaga	aagactcaat	gtggctttag	ataatggcct	tcccgaagga	acccccaaag	1620
atccaatcct	taggtccctc	gcctattcaa	atgctaacaa	ggagtgcaca	aaactcctac	1680
aggcccagg	acaaaccaac	agcccgttag	gggaaatgct	cagggcctgc	caaacttgg	1740
cgccccgaga	taaaaaaaa	atactaattg	tacaacctaa	aaaaactcct	ccccggaacc	1800
agccatgctt	ccgctgcggg	caagtaggtc	attggagcag	agattgtaaa	cagcctcggc	1860
cccctccggg	cccctgcccc	gtgtgtcagg	atcccacca	ctggaagcgg	gactgcccac	1920
agttaaaaac	agataccaga	gacagcgagg	acctactcct	agacctgccc	tgtgaagcac	1980
ccaatgtcgc	ggaacgaaaa	aactcctcag	ggggggagga	ttagcctccc	ccggaacct	2040
actccccctt	atacctttgt	cccagcagaa	gcagcctacc	ctgcatatcc	aggtatcgtt	2100
ttccaacacc	ccccctgtta	gcgttcaggc	gctcctcgac	actggagcag	acatcactgt	2160
cctcccggcc	tgcttatgcc	ctcccgatc	caacctccag	gacacctg	tcctaggtgc	2220
aggcgggcca	agtaccaaca	agtttaaaat	cctgcctcgt	ccagtcata	tccacttgcc	2280
ttttcgaagg	cagccgggtga	ccctaaccgc	ttgcctaatt	gatattaaca	accagtggtc	2340
catattaggg	cgagatgccc	tacaacaatg	tcaaagtctc	ctctatctgg	ctgaccaacc	2400
ctctaaggtc	ctccctgtcc	tagcacccaa	gcttatcgga	ttagagcacc	ttccccgcc	2460
cccagaagtc	tctcagttcc	cgtaaaacca	gagcgcctcc	aggctctgac	tgacctggtt	2520
tccagggcc	tggaggctaa	acatatagaa	ccctatcaag	gaccagcaa	taaccctatc	2580
ttcccctca	aaaagcccaa	tgggaagtgg	cggtttatcc	atgatctcgc	ggccacaaac	2640

-continued

---

tccgtcacc gagacotggc ttcaccgtcc cccggccctc cggaccttac cagtctgccc	2700
caaggcctcc cacatcttcg gacaattgac cttactgacg ccttcttcca aatcccgcta	2760
ccaaccattt ttcagcgtga ctttgcatte actctcccc aaccgaacaa ctatgggtccc	2820
ggaaccagat actcttggag agtactaccc caggggttca aaaatagtcc aactttattt	2880
gaacagcaac tctcccatat acttaccocct gtgcggaaaa cctttcctaa ttcccttatt	2940
atacaatata tggatgacat tctactggcc agccccgccc cggcgagct agctgctctc	3000
accgataaag taacaaacgc ctttaacaaag gaaggcctgc ccctatctcc ggaaaagact	3060
caggccactc ctgggtcccat acattttctt ggacaagtca tatcccagga ctgcataaca	3120
tacgagaccc toccatccat caatgtaaaa tccacctggt cactggcaga actacagtcc	3180
atgctaggag aattacaatg ggtctctaaa ggcacccctg tctctcgctc ctctctaac	3240
cagctctatc tgcctctccg aggccatcgt gaccctcggg atactataaa attaacctca	3300
atacaggtac aagctetaag aactattcag aaggccttga cctaaactg cgaagtaga	3360
ctagtaaatc agctgcccct cttggccctt ataatgctcc ggctacagg caccacggca	3420
gtcctcttcc aaacaaaaca aaagtggcca ctctctgggt tacacacccc tcaccggct	3480
accagcttgc gcccatgggg acaactattg gccaacgctg tcatcatcct agacaaatac	3540
tcactacaac actatggcca agtatgcaaa tcatttcate ataacatctc taatcaagcc	3600
ctcacttact acttacatac atctgaccaa tccagcgtag ccatcctatt acaaacctca	3660
cacagatttc acaatctcgg ggccagccca tcagggccat ggagaagcct cttacaaatg	3720
ccccagattt tccaaaacat cgatgtccta aggcctcctt ttaccatctc gcctgtagtc	3780
atcaacctg cccctgtct cttttcggac gggctctgct ccaaggcagc atttatcatc	3840
tgggatagac aagtcataca ccaacaggtc ctctccctgc cctcaacctg ttcagctcaa	3900
gccccgggaa tattttggct gttggcagga ctacaaaagt ctcaacctg ggtggcacta	3960
aatatatttc tagattcaaa atttcttate gggcaacctc ggcaatggc attaggggccc	4020
tttccaggac catccacca gtgcgaatta cacacacagc ttctccccct gctgcaggga	4080
aagactgtct atgtgcacca cgtaaggagt cacactcttc tgcaagacc tatatcccgc	4140
cttaatgaag ccacagatgc cctcatgctt gccctctgtt taccctcga cccacaact	4200
ctccaccagc tcaccactg taacccttac gccctacgta accatggagc cactgcctct	4260
gaagcccctg caattgtgca ggcattgca acatgcaagg tcatcaaccc acagggacgg	4320
ttaccccagg gatatacttc tgcaggccat gccctaagc atatctggca aggggagctc	4380
actcacctcc agtacaagcg atacaaatat tgctgctgg tttgggtaga tacttactcc	4440
ggggcggtct ctgtgtcctg ccggcgcaag gagaccggct ccgactgtgt cgctctgta	4500
cttgtggcca tctccattct aggaaaacca caaaacatca acaccgataa cggagctgcg	4560
tatctctccc aagagtcca acagttttgc aactcacttg ccattaaaca ttcaaccac	4620
attccctaca accccaccag ctctggcttg gttgaagaa caaatggtat tcttaaaact	4680
ttgatttcca aatatctct agacaatcac cactgcccc tggagacagc cgtctccaaa	4740
tccctctgga ctataaacca tcttaagtgc ctccctcgt gccaaaagac tcgggtggcag	4800
ctacatcagg cccagccct gccctccgct cctgaagaca cactcccacc ccacacatca	4860
ccaaaatggt attattataa aattcctggt cttaccaact caagggtggag tgggcctgta	4920

---

-continued

---

caatccctca aggaagcagc aggagcggct ctcatcccag taggtggaag ctatctctgg 4980  
atcccgtggc gctcctaaa aaggggtata tgcccaagac cggaaagcag cgcagccgtc 5040  
gacccaaaaa ccagagacca tcaactccat gggttaagttc tgcctttatt tctgtcttat 5100  
ttacatactc ttctctgcct cctctggcaa tcccagtcga tgcaccctgt tcataggagc 5160  
ctcttcttac cactccgacc cctgcgggtc cgatcaccgc cgggtgacct ggagactcga 5220  
cctatcttct ctcaacaagag atcaaaagcct aagccccca tgtccagact tagttactta 5280  
ctcacagtat cataggccct actccctgta tgtattcccc cattggataa ccaaacctaa 5340  
ccgtcgaggc ctaggttact attctgcttc ctactcagac ccctgcgcta tacaatgcc 5400  
ttacttagga tgcagtcgat ggacatgtcc ttatacaggc cgggtgtcca gccacattg 5460  
gaaatactcc tccgatctta attttaccba agaagtatca tccatctccc tacacttgca 5520  
cttttccaaa tgtgggtcct cattctcctt tctactagac gcaccaggat atgatccagt 5580  
gtggttcttc tctcccagg ccacacaggc cccacccagc cctgccctc tcatacagga 5640  
ctcagatctc caacatatcc tagaaccttc catcccctgg agtccaaaa tctcaatct 5700  
catctcctt accctaaaaa gctctaatta ttcttgcatt gtctgtgttg accgctccag 5760  
cctatctctg tggcatgttc tatatgaccc acttaaagcc cccaatccac ccgaccccaa 5820  
agcccagtct attctcgcag cctccctagc cattcccgcg agtaagtca ccccgccatt 5880  
tccttgacc cattgtatc gcccccctt acagggcctc tctcggaac actgcaataa 5940  
ctccgtagta ctgccccct tttccctgtc cccacttctt aacgtctcca gaccccgaaa 6000  
gcgccgagca gtcccattg ccatatggct agtgtccgc ctagcggcgc gcaccggtat 6060  
agctggcggg gttaccggct ccctgtctct ggccctcagc aaaagcctac tgcgcgaagt 6120  
agaccaggat atagatcacc tgacgcagcc gattgtaaaa aaccatgata atatcctctg 6180  
ggttgtctca tacgcagccc aaaatcgccg tggcctagat ttacttttct gggaaacaagg 6240  
gggcctttgt aaggccatcc aagaacaatg ttgcttctt aatatcagca acaccatgt 6300  
gtccgtctc caggaagac cccccctaga aaagagagta atcactggct ggggactcaa 6360  
ctgggacctc gggctctccc aatgggccc agaggccctt cagacaggta taactcttt 6420  
agcccttttt ctccctccta tgcctttagg accctgcgct ataccgccgc tacaggccct 6480  
cccttcccgt ctgcagcctc gcagtcagcc ctactccctt ctcaattatg aaaccaactt 6540  
ataacagatc tgtacctcc tgtagcagga ggetatggct ctgcctcta ctagacacc 6600  
aagtacagca taactctgaa gaatcccctt cgatgtcgac gccctggccc caacagtcca 6660  
tataccaaaa gtattctctt aaagattcct cgcagcctgc gogtagctgc tggctctccc 6720  
gtcccaaaaa gtctatatag ccctctagta agtcacaaaa cccctcgaac cccaacatgt 6780  
ctatacagtc cagttgctgt cgccttctt ttttctgctt ctctctctc tccagctctt 6840  
cggggcacct tctccgagc tcttctttt ttttctgctc tgcacaataa ctccagctt 6900  
gtctctgctc ctgagcaagg tcatccagcc gactcttcca ataaccagg tcttactgct 6960  
tagatcttaa gggccgtccc cggggtcgtt tgccattccc ctgaagcatg tccatttgat 7020  
ccctacctga tctctcacat aagttaaaca aagttccac aggtgtaaga ggctcctctg 7080  
cagtcaacac cggcggctcc agactccgag atcgggaagt caaactgcct ccagaagtag 7140  
aaatgcagga atataccaca ggcacagttc ctgggattgc agtctccggg gctaggacag 7200

-continued

---

```

gcatctgcct aaagtaacct acaaaagttt tattcccttg tcagcccatt tcccaggttt 7260
cggacagagc cttctctacg ggtaccctgt ctacgttttc ggcgactgtg tacaggccga 7320
ttggtgcccc atttctgggg ggctttgttc cgctcggcta caccgccatg ccctactggc 7380
cacgtgcccc gaacatcaga ttacctggga ccccatcgat ggacgcgttg tcagctcagc 7440
tctacaatac cttatccctc gactcccctc cttcccacc cagagaacta cccgcaccct 7500
caaggttctc acccccccaa ccactgctgc gacccccaaag attcctccat ccttcttcca 7560
cgccgttaaa aaacacaccc ccttcgaaa caattgcctt gaactcacc tgggagagca 7620
gttgccagcc atgtccttcc ccgaccctgg gctccgaccc caaaacatct acaccatgtg 7680
gggaagctcc gttgtgtgct tttacctata tcagctctcc ccccccata cctggcctct 7740
aatcccgcat gttatattct gccatcctga gcagcttggg gccttcctca cccgagtccc 7800
taccaaaaga ttagaagaac tcctgtataa gatattttta agcacagggg cgataatcat 7860
cctgcctgaa aactgttttc caaccacct gttccaaccc acccgcgccg ccgcggtgca 7920
ggccccctgg cacacaggcc tgctcccgtg tcaaaaggaa attgctaccc ccgggctcat 7980
ttggactttc actgatggca gcccatgat ttccggccct tgcccacaag aaggacagcc 8040
atcttttaga gtacaatcat ctacatttat ctttcaacaa ttccaaacca aggccagtca 8100
ccccgcttcc ctctgttccc acaactaat ccactactcc tcttttcatt cctccacct 8160
cctctttgag gaatatacaa ctatcccctt ttctctactt tttaatgaaa aaggggcaaa 8220
tgtcgtgat gatgagcccc gagacgggtc acaaccacca gctagaggac aaatagctga 8280
gtcaccgctc tgagaaccgt ctacaccgg gattgtgccc aaaaagaaca ccggggctct 8340
gacgtctctc cctaccctgg ctcccgaaa aaacaaaaa ccacccattt cctcatgttt 8400
gcctaaagct ctgacgataa ccctaaaaaa ttgactagc aaataaagaa cctggggccc 8460
tataaaaggg gagagcaacc taaaaatggg atcccttttc tgcacctcgc caaccctcc 8520
tctggccacg gtccgacttt ggtcattcct gcctacctga atcgccgctt cgggatcgag 8580
ccatccctct tctatttggg ggcacttcgc gactccgcc gccttccact cggtaaagatc 8640
ccactgggtc gagctaggcc atcaccctg gcccgctccc ctggagctct ctgcgcggc 8700
tcttaagggt gctccccctc agcaaaaggc ccagggcttt ctctacttcc ttgtttcaag 8760
tctctttctt tggcggctga cctaaatcga aagtagcact tctgctgtca gcagcgaggc 8820
ttgcccagg gccagcgctt gtaaggttac ccagctcgga gttgggtctc tagagaatca 8880
gggctaaagc tgctagccct aggaaagaag gcaaaaca 8917

```

```

<210> SEQ ID NO 37
<211> LENGTH: 491
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

```

```

<400> SEQUENCE: 37

```

```

Met Gly Lys Phe Cys Leu Tyr Phe Cys Leu Ile Tyr Ile Leu Phe Ser
1           5           10          15
Ala Ser Ser Gly Asn Pro Ser Arg Cys Thr Leu Phe Ile Gly Ala Ser
           20          25          30

```

-continued

---

Ser Tyr His Ser Asp Pro Cys Gly Ser Asp His Pro Arg Cys Thr Trp  
           35                          40                          45  
 Arg Leu Asp Leu Phe Ser Leu Thr Arg Asp Gln Ser Leu Ser Pro Pro  
   50                          55                          60  
 Cys Pro Asp Leu Val Thr Tyr Ser Gln Tyr His Arg Pro Tyr Ser Leu  
   65                          70                          75                          80  
 Tyr Val Phe Pro His Trp Ile Thr Lys Pro Asn Arg Arg Gly Leu Gly  
                           85                          90                          95  
 Tyr Tyr Ser Ala Ser Tyr Ser Asp Pro Cys Ala Ile Gln Cys Pro Tyr  
                   100                          105                          110  
 Leu Gly Cys Gln Ser Trp Thr Cys Pro Tyr Thr Gly Pro Val Ser Ser  
           115                          120                          125  
 Pro His Trp Lys Tyr Ser Ser Asp Leu Asn Phe Thr Gln Glu Val Ser  
   130                          135                          140  
 Ser Ile Ser Leu His Leu His Phe Ser Lys Cys Gly Ser Ser Phe Ser  
   145                          150                          155                          160  
 Phe Leu Leu Asp Ala Pro Gly Tyr Asp Pro Val Trp Phe Leu Ser Ser  
           165                          170                          175  
 Gln Ala Thr Gln Ala Pro Pro Thr Pro Ala Pro Leu Ile Gln Asp Ser  
                   180                          185                          190  
 Asp Leu Gln His Ile Leu Glu Pro Ser Ile Pro Trp Ser Ser Lys Ile  
   195                          200                          205  
 Leu Asn Leu Ile Leu Leu Thr Leu Lys Ser Ser Asn Tyr Ser Cys Met  
   210                          215                          220  
 Val Cys Val Asp Arg Ser Ser Leu Ser Ser Trp His Val Leu Tyr Asp  
   225                          230                          235                          240  
 Pro Leu Lys Ala Pro Asn Pro Pro Asp Pro Lys Ala Gln Ser Ile Leu  
           245                          250                          255  
 Arg Pro Ser Leu Ala Ile Pro Ala Ser Asn Val Thr Pro Pro Phe Pro  
           260                          265                          270  
 Trp Thr His Cys Tyr Arg Pro Leu Leu Gln Ala Ile Ser Ser Glu His  
   275                          280                          285  
 Cys Asn Asn Ser Val Val Leu Pro Pro Phe Ser Leu Ser Pro Leu Pro  
   290                          295                          300  
 Asn Val Ser Arg Pro Arg Lys Arg Arg Ala Val Pro Ile Ala Ile Trp  
   305                          310                          315                          320  
 Leu Val Ser Ala Leu Ala Ala Gly Thr Gly Ile Ala Gly Gly Val Thr  
           325                          330                          335  
 Gly Ser Leu Ser Leu Ala Ser Ser Lys Ser Leu Leu Arg Glu Val Asp  
           340                          345                          350  
 Gln Asp Ile Asp His Leu Thr Gln Ala Ile Val Lys Asn His Asp Asn  
   355                          360                          365  
 Ile Leu Arg Val Ala Gln Tyr Ala Ala Gln Asn Arg Arg Gly Leu Asp  
   370                          375                          380  
 Leu Leu Phe Trp Glu Gln Gly Gly Leu Cys Lys Ala Ile Gln Glu Gln  
   385                          390                          395                          400  
 Cys Cys Phe Leu Asn Ile Ser Asn Thr His Val Ser Val Leu Gln Glu  
           405                          410                          415  
 Arg Pro Pro Leu Glu Lys Arg Val Ile Thr Gly Trp Gly Leu Asn Trp  
   420                          425                          430  
 Asp Leu Gly Leu Ser Gln Trp Ala Arg Glu Ala Leu Gln Thr Gly Ile

-continued

---

```

      435              440              445
Thr Leu Leu Ala Leu Phe Leu Leu Leu Ile Val Val Gly Pro Cys Val
  450              455              460

Ile Arg Gln Leu Gln Ala Leu Pro Ser Arg Leu Gln His Arg Ser Gln
  465              470              475              480

Pro Tyr Ser Leu Leu Asn Tyr Glu Thr Asn Leu
      485              490

<210> SEQ ID NO 38
<211> LENGTH: 315
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
      note=synthetic construct

<400> SEQUENCE: 38
Met Gly Lys Phe Cys Leu Tyr Phe Cys Leu Ile Tyr Ile Leu Phe Ser
  1              5              10              15
Ala Ser Ser Gly Asn Pro Ser Arg Cys Thr Leu Phe Ile Gly Ala Ser
      20              25              30
Ser Tyr His Ser Asp Pro Cys Gly Ser Asp His Pro Arg Cys Thr Trp
      35              40              45
Arg Leu Asp Leu Phe Ser Leu Thr Arg Asp Gln Ser Leu Ser Pro Pro
      50              55              60
Cys Pro Asp Leu Val Thr Tyr Ser Gln Tyr His Arg Pro Tyr Ser Leu
      65              70              75              80
Tyr Val Phe Pro His Trp Ile Thr Lys Pro Asn Arg Arg Gly Leu Gly
      85              90              95
Tyr Tyr Ser Ala Ser Tyr Ser Asp Pro Cys Ala Ile Gln Cys Pro Tyr
      100             105             110
Leu Gly Cys Gln Ser Trp Thr Cys Pro Tyr Thr Gly Pro Val Ser Ser
      115             120             125
Pro His Trp Lys Tyr Ser Ser Asp Leu Asn Phe Thr Gln Glu Val Ser
      130             135             140
Ser Ile Ser Leu His Leu His Phe Ser Lys Cys Gly Ser Ser Phe Ser
      145             150             155             160
Phe Leu Leu Asp Ala Pro Gly Tyr Asp Pro Val Trp Phe Leu Ser Ser
      165             170             175
Gln Ala Thr Gln Ala Pro Pro Thr Pro Ala Pro Leu Ile Gln Asp Ser
      180             185             190
Asp Leu Gln His Ile Leu Glu Pro Ser Ile Pro Trp Ser Ser Lys Ile
      195             200             205
Leu Asn Leu Ile Leu Leu Thr Leu Lys Ser Ser Asn Tyr Ser Cys Met
      210             215             220
Val Cys Val Asp Arg Ser Ser Leu Ser Ser Trp His Val Leu Tyr Asp
      225             230             235             240
Pro Leu Lys Ala Pro Asn Pro Pro Asp Pro Lys Ala Gln Ser Ile Leu
      245             250             255
Arg Pro Ser Leu Ala Ile Pro Ala Ser Asn Val Thr Pro Pro Phe Pro
      260             265             270
Trp Thr His Cys Tyr Arg Pro Leu Leu Gln Ala Ile Ser Ser Glu His
      275             280             285

```

-continued

---

Cys Asn Asn Ser Val Val Leu Pro Pro Phe Ser Leu Ser Pro Leu Pro  
290 295 300

Asn Val Ser Arg Pro Arg Lys Arg Arg Ala Val  
305 310 315

<210> SEQ ID NO 39  
 <211> LENGTH: 176  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 39

Pro Ile Ala Ile Trp Leu Val Ser Ala Leu Ala Ala Gly Thr Gly Ile  
1 5 10 15

Ala Gly Gly Val Thr Gly Ser Leu Ser Leu Ala Ser Ser Lys Ser Leu  
20 25 30

Leu Arg Glu Val Asp Gln Asp Ile Asp His Leu Thr Gln Ala Ile Val  
35 40 45

Lys Asn His Asp Asn Ile Leu Arg Val Ala Gln Tyr Ala Ala Gln Asn  
50 55 60

Arg Arg Gly Leu Asp Leu Leu Phe Trp Glu Gln Gly Gly Leu Cys Lys  
65 70 75 80

Ala Ile Gln Glu Gln Cys Cys Phe Leu Asn Ile Ser Asn Thr His Val  
85 90 95

Ser Val Leu Gln Glu Arg Pro Pro Leu Glu Lys Arg Val Ile Thr Gly  
100 105 110

Trp Gly Leu Asn Trp Asp Leu Gly Leu Ser Gln Trp Ala Arg Glu Ala  
115 120 125

Leu Gln Thr Gly Ile Thr Leu Leu Ala Leu Phe Leu Leu Ile Val  
130 135 140

Val Gly Pro Cys Val Ile Arg Gln Leu Gln Ala Leu Pro Ser Arg Leu  
145 150 155 160

Gln His Arg Ser Gln Pro Tyr Ser Leu Leu Asn Tyr Glu Thr Asn Leu  
165 170 175

<210> SEQ ID NO 40  
 <211> LENGTH: 422  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 40

Met Gly Lys Thr Tyr Ser Ser Pro Ile Asn Pro Ile Pro Lys Ala Pro  
1 5 10 15

Lys Gly Leu Ala Ile His His Trp Leu Asn Phe Leu Gln Ala Ala Tyr  
20 25 30

Arg Leu Gln Pro Gly Pro Ser Glu Phe Asp Phe His Gln Leu Arg Lys  
35 40 45

Phe Leu Lys Leu Ala Ile Lys Thr Pro Val Trp Leu Asn Pro Ile Asn  
50 55 60

Tyr Ser Val Leu Ala Gly Leu Ile Pro Lys Asn Tyr Pro Gly Arg Val  
65 70 75 80

-continued

---

```

His Glu Ile Val Ala Ile Leu Ile Gln Glu Thr Pro Ala Arg Glu Ala
      85                               90                               95

Pro Pro Ser Ala Pro Leu Ala Glu Asp Pro Gln Lys Pro Pro Pro Tyr
      100                               105                               110

Pro Glu Gln Ala Gln Glu Ala Ser Gln Cys Leu Pro Ile Leu His Pro
      115                               120                               125

His Gly Ala Pro Ala Ala His Arg Pro Trp Gln Met Lys Asp Leu Gln
      130                               135                               140

Ala Ile Lys Gln Glu Val Ser Ser Ser Ala Pro Gly Ser Pro Gln Phe
      145                               150                               155                               160

Met Gln Thr Ile Arg Leu Ala Val Gln Gln Phe Asp Pro Thr Ala Lys
      165                               170                               175

Asp Leu His Asp Leu Leu Gln Tyr Leu Cys Ser Ser Leu Val Ala Ser
      180                               185                               190

Leu His His Gln Gln Leu Glu Thr Leu Ile Ala Gln Ala Glu Thr Gln
      195                               200                               205

Gly Ile Thr Gly Tyr Asn Pro Leu Ala Gly Pro Leu Arg Ile Gln Ala
      210                               215                               220

Asn Asn Pro Asn Gln Gln Gly Leu Arg Lys Glu Tyr Gln Asn Leu Trp
      225                               230                               235                               240

Leu Ser Ala Phe Ser Ala Leu Pro Gly Asn Thr Lys Asp Pro Thr Trp
      245                               250                               255

Ala Ala Ile Leu Gln Gly Pro Glu Glu Pro Phe Gly Ser Phe Val Glu
      260                               265                               270

Arg Leu Asn Val Ala Leu Asp Asn Gly Leu Pro Glu Gly Thr Pro Lys
      275                               280                               285

Asp Pro Ile Leu Arg Ser Leu Ala Tyr Ser Asn Ala Asn Lys Glu Cys
      290                               295                               300

Gln Lys Leu Leu Gln Ala Arg Gly Gln Thr Asn Ser Pro Leu Gly Glu
      305                               310                               315                               320

Met Leu Arg Ala Cys Gln Thr Trp Thr Pro Arg Asp Lys Asn Lys Ile
      325                               330                               335

Leu Met Val Gln Pro Lys Lys Thr Pro Pro Pro Asn Gln Pro Cys Phe
      340                               345                               350

Arg Cys Gly Gln Val Gly His Trp Ser Arg Asp Cys Lys Gln Pro Arg
      355                               360                               365

Pro Pro Pro Gly Pro Cys Pro Val Cys Gln Asp Pro Thr His Trp Lys
      370                               375                               380

Arg Asp Cys Pro Gln Leu Lys Thr Asp Thr Arg Asp Ser Glu Asp Leu
      385                               390                               395                               400

Leu Leu Asp Leu Pro Cys Glu Ala Pro Asn Val Arg Glu Arg Lys Asn
      405                               410                               415

Ser Ser Gly Gly Glu Asp
      420

```

&lt;210&gt; SEQ ID NO 41

&lt;211&gt; LENGTH: 86

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

&lt;400&gt; SEQUENCE: 41

-continued

---

Leu Met Val Gln Pro Lys Lys Thr Pro Pro Pro Asn Gln Pro Cys Phe  
 1 5 10 15

Arg Cys Gly Gln Val Gly His Trp Ser Arg Asp Cys Lys Gln Pro Arg  
 20 25 30

Pro Pro Pro Gly Pro Cys Pro Val Cys Gln Asp Pro Thr His Trp Lys  
 35 40 45

Arg Asp Cys Pro Gln Leu Lys Thr Asp Thr Arg Asp Ser Glu Asp Leu  
 50 55 60

Leu Leu Asp Leu Pro Cys Glu Ala Pro Asn Val Arg Glu Arg Lys Asn  
 65 70 75 80

Ser Ser Gly Gly Glu Asp  
 85

<210> SEQ ID NO 42  
 <211> LENGTH: 123  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 42

Met Gly Lys Thr Tyr Ser Ser Pro Ile Asn Pro Ile Pro Lys Ala Pro  
 1 5 10 15

Lys Gly Leu Ala Ile His His Trp Leu Asn Phe Leu Gln Ala Ala Tyr  
 20 25 30

Arg Leu Gln Pro Gly Pro Ser Glu Phe Asp Phe His Gln Leu Arg Lys  
 35 40 45

Phe Leu Lys Leu Ala Ile Lys Thr Pro Val Trp Leu Asn Pro Ile Asn  
 50 55 60

Tyr Ser Val Leu Ala Gly Leu Ile Pro Lys Asn Tyr Pro Gly Arg Val  
 65 70 75 80

His Glu Ile Val Ala Ile Leu Ile Gln Glu Thr Pro Ala Arg Glu Ala  
 85 90 95

Pro Pro Ser Ala Pro Leu Ala Glu Asp Pro Gln Lys Pro Pro Pro Tyr  
 100 105 110

Pro Glu Gln Ala Gln Glu Ala Ser Gln Cys Leu  
 115 120

<210> SEQ ID NO 43  
 <211> LENGTH: 213  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 43

Pro Ile Leu His Pro His Gly Ala Pro Ala Ala His Arg Pro Trp Gln  
 1 5 10 15

Met Lys Asp Leu Gln Ala Ile Lys Gln Glu Val Ser Ser Ser Ala Pro  
 20 25 30

Gly Ser Pro Gln Phe Met Gln Thr Ile Arg Leu Ala Val Gln Gln Phe  
 35 40 45

Asp Pro Thr Ala Lys Asp Leu His Asp Leu Leu Gln Tyr Leu Cys Ser  
 50 55 60

-continued

---

```

Ser Leu Val Ala Ser Leu His His Gln Gln Leu Glu Thr Leu Ile Ala
65                               70                               75                               80

Gln Ala Glu Thr Gln Gly Ile Thr Gly Tyr Asn Pro Leu Ala Gly Pro
85                               90                               95

Leu Arg Ile Gln Ala Asn Asn Pro Asn Gln Gln Gly Leu Arg Lys Glu
100                              105                              110

Tyr Gln Asn Leu Trp Leu Ser Ala Phe Ser Ala Leu Pro Gly Asn Thr
115                              120                              125

Lys Asp Pro Thr Trp Ala Ala Ile Leu Gln Gly Pro Glu Glu Pro Phe
130                              135                              140

Gly Ser Phe Val Glu Arg Leu Asn Val Ala Leu Asp Asn Gly Leu Pro
145                              150                              155                              160

Glu Gly Thr Pro Lys Asp Pro Ile Leu Arg Ser Leu Ala Tyr Ser Asn
165                              170                              175

Ala Asn Lys Glu Cys Gln Lys Leu Leu Gln Ala Arg Gly Gln Thr Asn
180                              185                              190

Ser Pro Leu Gly Glu Met Leu Arg Ala Cys Gln Thr Trp Thr Pro Arg
195                              200                              205

Asp Lys Asn Lys Ile
210

```

```

<210> SEQ ID NO 44
<211> LENGTH: 889
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

```

```

<400> SEQUENCE: 44

```

```

Gly Pro Pro Cys Pro Ser Thr Gln Ala Tyr Arg Ile Arg Ala Pro Ser
1                               5                               10                               15

Pro Ala Pro Arg Ser Leu Ser Val Pro Val Lys Pro Glu Arg Leu Gln
20                              25                              30

Ala Leu Thr Asp Leu Val Ser Arg Ala Leu Glu Ala Lys His Ile Glu
35                              40                              45

Pro Tyr Gln Gly Pro Gly Asn Asn Pro Ile Phe Pro Val Lys Lys Pro
50                              55                              60

Asn Gly Lys Trp Arg Phe Ile His Asp Leu Arg Ala Thr Asn Ser Val
65                              70                              75                              80

Thr Arg Asp Leu Ala Ser Pro Ser Pro Gly Pro Pro Asp Leu Thr Ser
85                              90                              95

Leu Pro Gln Gly Leu Pro His Leu Arg Thr Ile Asp Leu Thr Asp Ala
100                             105                             110

Phe Phe Gln Ile Pro Leu Pro Thr Ile Phe Gln Pro Tyr Phe Ala Phe
115                             120                             125

Thr Leu Pro Gln Pro Asn Asn Tyr Gly Pro Gly Thr Arg Tyr Ser Trp
130                             135                             140

Arg Val Leu Pro Gln Gly Phe Lys Asn Ser Pro Thr Leu Phe Glu Gln
145                             150                             155                             160

Gln Leu Ser His Ile Leu Thr Pro Val Arg Lys Thr Phe Pro Asn Ser
165                             170                             175

Leu Ile Ile Gln Tyr Met Asp Asp Ile Leu Leu Ala Ser Pro Ala Pro

```

-continued

	180							185						190			
Gly	Glu	Leu	Ala	Ala	Leu	Thr	Asp	Lys	Val	Thr	Asn	Ala	Leu	Thr	Lys		
	195						200					205					
Glu	Gly	Leu	Pro	Leu	Ser	Pro	Glu	Lys	Thr	Gln	Ala	Thr	Pro	Gly	Pro		
	210					215					220						
Ile	His	Phe	Leu	Gly	Gln	Val	Ile	Ser	Gln	Asp	Cys	Ile	Thr	Tyr	Glu		
225				230					235						240		
Thr	Leu	Pro	Ser	Ile	Asn	Val	Lys	Ser	Thr	Trp	Ser	Leu	Ala	Glu	Leu		
			245					250						255			
Gln	Ser	Met	Leu	Gly	Glu	Leu	Gln	Trp	Val	Ser	Lys	Gly	Thr	Pro	Val		
		260						265					270				
Leu	Arg	Ser	Ser	Leu	His	Gln	Leu	Tyr	Leu	Ala	Leu	Arg	Gly	His	Arg		
	275						280					285					
Asp	Pro	Arg	Asp	Thr	Ile	Lys	Leu	Thr	Ser	Ile	Gln	Val	Gln	Ala	Leu		
	290					295					300						
Arg	Thr	Ile	Gln	Lys	Ala	Leu	Thr	Leu	Asn	Cys	Arg	Ser	Arg	Leu	Val		
305				310					315						320		
Asn	Gln	Leu	Pro	Ile	Leu	Ala	Leu	Ile	Met	Leu	Arg	Pro	Thr	Gly	Thr		
			325						330					335			
Thr	Ala	Val	Leu	Phe	Gln	Thr	Lys	Gln	Lys	Trp	Pro	Leu	Val	Trp	Leu		
			340					345					350				
His	Thr	Pro	His	Pro	Ala	Thr	Ser	Leu	Arg	Pro	Trp	Gly	Gln	Leu	Leu		
	355						360					365					
Ala	Asn	Ala	Val	Ile	Ile	Leu	Asp	Lys	Tyr	Ser	Leu	Gln	His	Tyr	Gly		
	370					375					380						
Gln	Val	Cys	Lys	Ser	Phe	His	His	Asn	Ile	Ser	Asn	Gln	Ala	Leu	Thr		
385					390					395					400		
Tyr	Tyr	Leu	His	Thr	Ser	Asp	Gln	Ser	Ser	Val	Ala	Ile	Leu	Leu	Gln		
			405					410						415			
His	Ser	His	Arg	Phe	His	Asn	Leu	Gly	Ala	Gln	Pro	Ser	Gly	Pro	Trp		
	420							425					430				
Arg	Ser	Leu	Leu	Gln	Met	Pro	Gln	Ile	Phe	Gln	Asn	Ile	Asp	Val	Leu		
	435						440					445					
Arg	Pro	Pro	Phe	Thr	Ile	Ser	Pro	Val	Val	Ile	Asn	His	Ala	Pro	Cys		
	450					455					460						
Leu	Phe	Ser	Asp	Gly	Ser	Ala	Ser	Lys	Ala	Ala	Phe	Ile	Ile	Trp	Asp		
465					470				475						480		
Arg	Gln	Val	Ile	His	Gln	Gln	Val	Leu	Ser	Leu	Pro	Ser	Thr	Cys	Ser		
			485						490					495			
Ala	Gln	Ala	Gly	Glu	Leu	Phe	Gly	Leu	Leu	Ala	Gly	Leu	Gln	Lys	Ser		
			500					505					510				
Gln	Pro	Trp	Val	Ala	Leu	Asn	Ile	Phe	Leu	Asp	Ser	Lys	Phe	Leu	Ile		
	515						520					525					
Gly	His	Leu	Arg	Arg	Met	Ala	Leu	Gly	Ala	Phe	Pro	Gly	Pro	Ser	Thr		
	530					535					540						
Gln	Cys	Glu	Leu	His	Thr	Gln	Leu	Leu	Pro	Leu	Leu	Gln	Gly	Lys	Thr		
545					550				555						560		
Val	Tyr	Val	His	His	Val	Arg	Ser	His	Thr	Leu	Leu	Gln	Asp	Pro	Ile		
			565						570					575			
Ser	Arg	Leu	Asn	Glu	Ala	Thr	Asp	Ala	Leu	Met	Leu	Ala	Pro	Leu	Leu		
			580					585					590				

-continued

---

Pro Leu Asp Pro Thr Thr Leu His Gln Leu Thr His Cys Asn Pro Tyr  
 595 600 605

Ala Leu Arg Asn His Gly Ala Thr Ala Ser Glu Ala His Ala Ile Val  
 610 615 620

Gln Ala Cys His Thr Cys Lys Val Ile Asn Pro Gln Gly Arg Leu Pro  
 625 630 635 640

Gln Gly Tyr Ile Arg Arg Gly His Ala Pro Asn Asp Ile Trp Gln Gly  
 645 650 655

Asp Val Thr His Leu Gln Tyr Lys Arg Tyr Lys Tyr Cys Leu Leu Val  
 660 665 670

Trp Val Asp Thr Tyr Ser Gly Ala Val Ser Val Ser Cys Arg Arg Lys  
 675 680 685

Glu Thr Gly Ser Asp Cys Val Ala Ser Leu Leu Val Ala Ile Ser Ile  
 690 695 700

Leu Gly Lys Pro Gln Asn Ile Asn Thr Asp Asn Gly Ala Ala Tyr Leu  
 705 710 715 720

Ser Gln Glu Phe Gln Gln Phe Cys Asn Ser Leu Ala Ile Lys His Ser  
 725 730 735

Thr His Ile Pro Tyr Asn Pro Thr Ser Ser Gly Leu Val Glu Arg Thr  
 740 745 750

Asn Gly Ile Leu Lys Thr Leu Ile Ser Lys Tyr Leu Leu Asp Asn His  
 755 760 765

His Leu Pro Leu Glu Thr Ala Val Ser Lys Ser Leu Trp Thr Ile Asn  
 770 775 780

His Leu Asn Val Leu Pro Ser Cys Gln Lys Thr Arg Trp Gln Leu His  
 785 790 795 800

Gln Ala Gln Pro Leu Pro Pro Val Pro Glu Asp Thr Leu Pro Pro His  
 805 810 815

Thr Ser Pro Lys Trp Tyr Tyr Tyr Lys Ile Pro Gly Leu Thr Asn Ser  
 820 825 830

Arg Trp Ser Gly Pro Val Gln Ser Leu Lys Glu Ala Ala Gly Ala Ala  
 835 840 845

Leu Ile Pro Val Gly Gly Ser Tyr Leu Trp Ile Pro Trp Arg Leu Leu  
 850 855 860

Lys Arg Gly Ile Cys Pro Arg Pro Glu Ser Ser Ala Ala Val Asp Pro  
 865 870 875 880

Lys Thr Arg Asp His Gln Leu His Gly  
 885

<210> SEQ ID NO 45  
 <211> LENGTH: 697  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 45

tgtcgatgat gatgagcccc gagacgggtc acaaccacca gctagaggac aaatagctga 60

gtcaccggc tgagaaccgt ctacaccgg gattgtgccc aaaaagaaca cggggctct 120

gacgtctctc cctaccctgg ctcccgaaa aaacaaaaa ccaccattt cctcatgttt 180

gcctaaagct ctgacgataa ccctaaaaa tttgactagc aaataaagaa cctggggccc 240

-continued

---

```

tataaaaggg gagagcaacc taaaaatggg atcccttttc tgcacctcgc caaccctccc 300
tctggccaag gtcgcatttt ggtcattcct gcctacctga atcgccgctt cgggatcgag 360
ccatccctct tctatttggg ggcacttcgc gcaactccgc gccttccact cggttaagatc 420
ccactgggtc gagctaggcc atcacccctg ggccgctccc ctggagctct ctgcgcgggc 480
tcttaagggt gctccccctc agcaaagggc ccagggcttt ctctacttcc ttgtttcaag 540
tctctttctt tggcggtcga cctaaatcga aagtagcact tctgctgtca gcagcgaggc 600
ttggcccagg gccagcgctt gtaaggttac ccagctcgga gttgggtctc tagagaatca 660
gggctaaagc tgctagccct aggaaagaag gcaaaca 697

```

```

<210> SEQ ID NO 46
<211> LENGTH: 177
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

```

```

<400> SEQUENCE: 46

```

```

Ser Thr Gln Cys Pro Gly Thr Lys Lys Leu Leu Arg Gly Gly Gly Leu
1          5          10          15
Ala Ser Pro Arg Thr Ile Leu Pro Leu Ile Pro Leu Ser Gln Gln Lys
20          25          30
Gln Pro Thr Leu His Ile Gln Val Ser Phe Ser Asn Thr Pro Pro Val
35          40          45
Ser Val Gln Ala Leu Leu Asp Thr Gly Ala Asp Ile Thr Val Leu Pro
50          55          60
Ala Cys Leu Cys Pro Pro Asp Ser Asn Leu Gln Asp Thr Thr Val Leu
65          70          75          80
Gly Ala Gly Gly Pro Ser Thr Asn Lys Phe Lys Ile Leu Pro Cys Pro
85          90          95
Val His Ile His Leu Pro Phe Arg Arg Gln Pro Val Thr Leu Thr Ala
100         105         110
Cys Leu Ile Asp Ile Asn Asn Gln Trp Thr Ile Leu Gly Arg Asp Ala
115         120         125
Leu Gln Gln Cys Gln Ser Ser Leu Tyr Leu Ala Asp Gln Pro Ser Lys
130         135         140
Val Leu Pro Val Leu Ala Pro Lys Leu Ile Gly Leu Glu His Leu Pro
145         150         155         160
Pro Pro Pro Glu Val Ser Gln Phe Pro Leu Asn Gln Ser Ala Ser Arg
165         170         175
Leu

```

```

<210> SEQ ID NO 47
<211> LENGTH: 534
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

```

```

<400> SEQUENCE: 47

```

```

agcacccaat gtcgggaac gaaaaaac ctcagggggg gaggattagc ctcccccca 60

```

-continued

---

```

accatactcc cccttataacc tttgtcccag cagaagcagc ctaccctgca tatccaggta 120
tcgttttcca acaccccccc tgttagcggt caggcgctcc togacactgg agcagacatc 180
actgtcctcc cggcctgctt atgcctccc gattccaacc tccaggacac cactgtccta 240
gggtgcaggcg ggccaagtac caacaagttt aaaatcctgc cctgtccagt ccatatccac 300
ttgccttttc gaaggcagcc ggtgacctta accgcttgcc taattgatat taacaaccag 360
tggaccatat tagggcgaga tgcctacaa caatgtcaaa gttcctcta tctggctgac 420
caaccctcta aggtcctccc tgtcctagca cccaagctta tcggattaga gcaccttccc 480
ccgccccccag aagtctctca gttcccgtta aaccagagcg cctccaggct ctga 534

```

```

<210> SEQ ID NO 48
<211> LENGTH: 182
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

```

```

<400> SEQUENCE: 48

```

```

Met Pro Lys Thr Arg Lys Gln Arg Ser Arg Arg Pro Lys Asn Gln Arg
1          5          10          15
Pro Ser Thr Pro Trp Pro Ile Ser Gln Val Ser Asp Arg Ala Phe Ser
          20          25          30
Thr Gly Thr Leu Ser Thr Phe Ser Ala Thr Val Tyr Arg Pro Ile Gly
          35          40          45
Ala Pro Phe Leu Gly Gly Phe Val Pro Leu Gly Tyr Thr Ala Met Pro
          50          55          60
Tyr Trp Pro Arg Ala Pro Asn Ile Arg Leu Pro Gly Thr Pro Ser Met
65          70          75          80
Asp Ala Leu Ser Ala Gln Leu Tyr Asn Thr Leu Ser Leu Asp Ser Pro
          85          90          95
Pro Ser Pro Pro Arg Glu Leu Pro Ala Pro Ser Arg Phe Ser Pro Pro
          100         105         110
Gln Pro Leu Leu Arg Pro Pro Arg Phe Leu His Pro Ser Ser Thr Pro
          115         120         125
Leu Lys Asn Thr Pro Pro Ser Glu Thr Ile Ala Leu Asn Ser Pro Trp
          130         135         140
Glu Ser Ser Cys Gln Pro Cys Pro Ser Pro Thr Leu Gly Ser Asp Pro
145         150         155         160
Lys Thr Ser Thr Pro Cys Gly Glu Ala Pro Leu Cys Ala Phe Thr Ser
          165         170         175
Ile Ser Ser Pro Pro Pro
          180

```

```

<210> SEQ ID NO 49
<211> LENGTH: 549
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

```

```

<400> SEQUENCE: 49

```

```

atgcccaaga cccgaagca gcgcagcgt cgacccaaaa accagagacc atcaactcca 60

```

-continued

```

tggcccattt cccaggtttc ggacagagcc ttctctacgg gtaccctgtc tacgttttcg 120
gcgactgtgt acaggccgat tgggtgcccc tttctggggg gctttgttcc gctcggctac 180
accgccatgc cctactggcc acgtgccccg aacatcagat tacctgggac cccatcgatg 240
gacgcgttgt cagctcagct ctacaatacc ttatccctcg actccctcc ttecccaccc 300
agagaactac ccgaccctc aaggttctca ccccccaac cactgctgcg accccaaga 360
ttctccatc cttcttccac gccgttaaaa aacacacccc cttcogaaac aattgccttg 420
aactcacctt gggagagcag ttgccagcca tgccttccc cgaccctggg ctccgacccc 480
aaaacatcta caccatgtgg ggaagctcgg ttgtgtgcct ttaccttat cagctctccc 540
cccccatga 549

```

```

<210> SEQ ID NO 50
<211> LENGTH: 350
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

```

```

<400> SEQUENCE: 50

```

```

Met Ala His Phe Pro Gly Phe Gly Gln Ser Leu Leu Tyr Gly Tyr Pro
1           5           10           15
Val Tyr Val Phe Gly Asp Cys Val Gln Ala Asp Trp Cys Pro Ile Ser
20          25          30
Gly Gly Leu Cys Ser Ala Arg Leu His Arg His Ala Leu Leu Ala Thr
35          40          45
Cys Pro Glu His Gln Ile Thr Trp Asp Pro Ile Asp Gly Arg Val Val
50          55          60
Ser Ser Ala Leu Gln Tyr Leu Ile Pro Arg Leu Pro Ser Phe Pro Thr
65          70          75          80
Gln Arg Thr Thr Arg Thr Leu Lys Val Leu Thr Pro Pro Thr Thr Ala
85          90          95
Ala Thr Pro Lys Ile Pro Pro Ser Phe Phe His Ala Val Lys Lys His
100         105         110
Thr Pro Phe Arg Asn Asn Cys Leu Glu Leu Thr Leu Gly Glu Gln Leu
115         120         125
Pro Ala Met Ser Phe Pro Asp Pro Gly Leu Arg Pro Gln Asn Ile Tyr
130         135         140
Thr Met Trp Gly Ser Ser Val Val Cys Leu Tyr Leu Tyr Gln Leu Ser
145         150         155         160
Pro Pro Met Thr Trp Pro Leu Ile Pro His Val Ile Phe Cys His Pro
165         170         175
Glu Gln Leu Gly Ala Phe Leu Thr Arg Val Pro Thr Lys Arg Leu Glu
180         185         190
Glu Leu Leu Tyr Lys Ile Phe Leu Ser Thr Gly Ala Ile Ile Ile Leu
195         200         205
Pro Glu Asn Cys Phe Pro Thr Thr Leu Phe Gln Pro Thr Arg Ala Pro
210         215         220
Ala Val Gln Ala Pro Trp His Thr Gly Leu Leu Pro Cys Gln Lys Glu
225         230         235         240
Ile Ala Thr Pro Gly Leu Ile Trp Thr Phe Thr Asp Gly Ser Pro Met
245         250         255

```

-continued

---

Ile Ser Gly Pro Cys Pro Lys Glu Gly Gln Pro Ser Leu Val Val Gln  
 260 265 270

Ser Ser Thr Phe Ile Phe Gln Gln Phe Gln Thr Lys Ala Ser His Pro  
 275 280 285

Ala Phe Leu Leu Ser His Lys Leu Ile His Tyr Ser Ser Phe His Ser  
 290 295 300

Leu His Leu Leu Phe Glu Glu Tyr Thr Thr Ile Pro Phe Ser Leu Leu  
 305 310 315 320

Phe Asn Glu Lys Gly Ala Asn Val Asp Asp Asp Glu Pro Arg Asp Gly  
 325 330 335

Ser Gln Pro Pro Ala Arg Gly Gln Ile Ala Glu Ser Pro Val  
 340 345 350

<210> SEQ ID NO 51  
 <211> LENGTH: 1053  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 51

```

atggccatt tcccaggtt cggacagagc cttctctacg ggtaccctgt ctacgttttc    60
ggcgactgtg tacaggccga ttggtgcccc atttctgggg ggetttgttc cgctcggeta    120
caccgccatg ccctactggc cacgtgcccc gaacatcaga ttacctggga ccccatcgat    180
ggacgcgttg tcagctcagc tctacaatac cttatccctc gactcccctc cttcccacc    240
cagagaacta cccgcacct caaggttctc acccccctca ccaactgctgc gaccccctag    300
attctccat ccttcttcca cgccgttaaa aaacacaccc ccttcgaaa caattgcctt    360
gaactcacc tgggagagca gttgccagcc atgtccttcc cggacctgg gctccgacc    420
caaaacatct acaccatgtg gggaaactcc gttgtgtgcc ttaacctta tcagctctcc    480
cccccatga cctggcctct aatcccgcct gttatattct gccatcctga gcagcttgg    540
gccttctca cccgagtccc taccaaacga ttagaagaac tcctgtataa gatattttta    600
agcacagggg cgataatcat cctgctgaa aactgttttc caaccacct gttccaaccc    660
accgcgcgc cccgggtgca ggccccctgg cacacaggcc tgctcccgtg tcaaaaggaa    720
attgctaccc cccggctcat ttggacttcc actgatggca gcccctgat tccgcccct    780
tgccccaaag aaggacagcc atctttagta gtacaatcat ctacatttat ctttcaacaa    840
ttccaaacca aggccagtca ccccgtttc ctcttgctcc acaaactaat ccactactcc    900
tcttttcatt cctccacct cctctttgag gaatatacaa ctatcccctt ttcttactt    960
ttaaataaaa aaggggcaaa tgctgatgat gatgagcccc gagacgggtc acaaccacca    1020
gctagaggac aaatagctga gtcaccctgc tga                                1053
    
```

<210> SEQ ID NO 52  
 <211> LENGTH: 699  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 52

-continued

---

```

cagatctgct acctcctgta gcaggaggct atggetctcg cctetaactag acaccaagt    60
acagcataat cctgaagaat ccccttcgat gtcgacgccc tggccccaac agtccatata    120
ccaaaagtat tcctctaaag attcctcgca gcctgcgctg agetgctggc tctcccgtc    180
caaaaagtct atatagccct ctagtaagtc aaaaacccc tcgaaccca acatgtctat    240
acagtccagt tgctgtcgcc ttcctttttt ctgcctcttc ctctcctcca gctcttcgag    300
gcacctcttc cgacgctctt cctttttttt tcggttctgc caataactca gcagttgctc    360
ctgctcctga gcaaggctac ccagccgact cttccaataa cccaggtcct tactgctaga    420
tcctaagggc cgtccccggg gtcgtttgcc attcccctga agcatgtcca ttgatacct    480
acctgatctc tcacataagt ttaacaaagt ttccacaggt gtaaggagct cctctgcagt    540
caacaccggc ggtcccagac tccgagatcg ggaagtcaaa ctgcctccag aagtagaaat    600
gcaggaatat accacaggca cagttcctgg gattgcagtc tccggggcta ggacaggcat    660
ctgcctaaag taacctacaa aagttttatt cccttgtea    699

```

&lt;210&gt; SEQ ID NO 53

&lt;211&gt; LENGTH: 5320

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

&lt;400&gt; SEQUENCE: 53

```

ctggccatat cgaaccttac tctggaccag gcaacaaccc agttttccct gttaaaaaac    60
ccaacggcaa gtggcgattt atccatgacc tcagggccac taatgccatc accactaccc    120
ttgcctcgcc ctccccggc ccccttgatc ttaccagcct gccacaggcc ttgccccatc    180
ttcagaccaa cgatctcagc gacgctttct tccagattcc cctcccaaag cgattccagc    240
cctacttcgc ctttaccatc cccagccat taaatcatgg gcctgggagc aggtacgctt    300
ggacagtcct tcccgaaggc ttcaaaaaa gccccacgct ctttgagcaa cagctggcca    360
gcgtactagg cccagcccca aaagccttcc ccacatccgt catcgtccaa tacatggagc    420
acatcctctt ggcagcctcc tcccagcagc aactagatca gctggccacc cttaccgcac    480
agctattgtc ctctcatggt ctcccagttt cccaggaaaa aacccaacgc accccaggaa    540
aaatacaact cctggggcca atcatacatc cagatcacat cacctatgaa accaccccca    600
ccatccccat taaggcacac tggaccctga ctgaactgca aacctcctg ggggagctcc    660
agtgggtctc caaggggact cctgtcctcc gagaacacct tcaactgtctc tactcagcct    720
tgagaggtct caaagacccc cgggacaacta tcacctctcg tcaatcctac ctccacgctc    780
tccacaacat tcagcaagcc ctgcatcaca attgcccggg tcgccttgac tctacgctcc    840
ccctccttgg cctcatcttc ctacgtccat ccggcagcag ctacgtcttc tccagacaa    900
atcataaatg gccctagtc tggctccacg cccccatcc cccgaccagc ctatgccct    960
gggggacat actgcctgc actgtaacta cccttgacaa gtatgccttg cagcactatg    1020
gccaactatg caaatcattc catcataaca tgtccacca ggccctacac gatttcgtaa    1080
aaaattcttc tcacccagc gtcgccatat taattcacca catgcacggg ttctgtgatc    1140
tgggcagaca gccaccggga ccctggcgaa ccctcttaca actccggcc cttctccggg    1200

```

-continued

---

aaccccagct cctcaggcct gcattttccc tatccccagt gggtatagat caggcccctt	1260
gtctgttctc tgatgggtct ccccaaaagg ccgcctatgt aatttgggac aaggctatc	1320
tcagccagcg gtcggteccc ctgccccccc atgcccaataa ctcagcacia aagggggaat	1380
tagtcggact cctcttgggc ttgcaagccg cacagccctg gccatccctt aacattttcc	1440
tagactcaaa gttcctcatc cggtagctcc agtccctcgc ttccggggcc ttccaaggat	1500
catccacaca ccaccgtctc caggcgtccc tgcccacact cctccagggc aaggtcgtgt	1560
atctccacca cacccgcagc cacaccaat tgctgatcc catctcgacc ctcaatgaat	1620
ataccgactc tctcattgtc gccccgtaa cccccctgaa gcctgagggc ctccatgccc	1680
tcaccactg caaccaacgc gccctcgttt cccacggagc caccctgca caggctaagc	1740
aactcgtgca ggctgcccgc acctgtcaaa tcattaaccc tcaacaccac atgccgctg	1800
gccacatccg ccggcgccac ttcccaaac acacatggca aggagatgc acccacctta	1860
agcaciaacg gaccgatac tgctccacg tctgggtgga taccttctca ggtgcggtat	1920
cttgtgtctg caaaaagaaa gaaactagca gcgacctat caaaacctc ctacatgcca	1980
tctcctgctc aggcaagccc ttctctgtta acacggacaa tggaccctc tacctttctc	2040
aggagtcca cgaattctgt accaccctct gcatacaaca ctccaccat attccctaca	2100
atccgacaag ttcaggcctg gtggagcgc caaatggcat tctcaagaca ctactataca	2160
aatatttctc agaccacctc gacctcccc tagaaagcgc ggtttcaaag gctctctgga	2220
ccattaacca tttaaatgct atgcccctc gtggtaagac tcgggtggcag ctccatcaca	2280
cccccccct gcctctatt tccgagtcca tacaaccac tcccaccagg ctacattggt	2340
actattaca aacccctgga cttaccaacc agcagtgga aggcccgtc caatctctcc	2400
aggaaagcagc aggagcagct ctcttcaag tcagtgacgg ctgccccag tggatccctt	2460
ggcggctcct gaagaagact gtatgccaa aaccgacga cccgaacc gcagggeacg	2520
tcgaaacaga ccaccaacac catgggtaac gtactcttct taactttatt ggccacctg	2580
ggcatcccag tacttcaggc cagccggtgt acaatcacgg taggtatctc ctctaccac	2640
tccagcccct gcagcccagc ccagccttta tgtacctggg ccctcgacct tgtgtccatc	2700
actaaggacc agctctctc cccccctgc caaaacctga tcacctatc caactaccac	2760
aagacctact cctgtatct cttcccacac tgggtacaaa agccactccg ccgggggctt	2820
ggatactact cagcctccta ctctgatcct tgctccctac aatgtccta cctaggaagt	2880
caatcatgga cttgcccta tactggcctc gtctcgagcc caacttgag attctccaca	2940
gatgtaaatt tcaccaaga agtcagccgt gtctccctaa aacttcattt ctccaaatgt	3000
ggttcctcct taactctgtt aatagatgcc ccgggttacg atccgctgtg gtacctcaca	3060
tccgagccta ctcaggaacc cccaacctc ccgcccactag tcagcgactc agacctagag	3120
catgtctga ctctctcggc ctctcgggccc tccaagatgc tgacctcat ccacctaac	3180
ttgcagagca ccaactatc ctgtatggc tgtattgacc gcgccagcct ctctcctgg	3240
cacgtattat acactcccaa catctctagt aatgccccct caaaacctc cgtccgccct	3300
tccttgccc tatccgccc gcgaccacag ccttcccct ggaccattg ctatcaacca	3360
cagggtcaag ctgtaaccac cgcaagtgc aataattcca tcatactcc cccattttct	3420
ctctctcct tgctcgtgct ccctctcact aggcgagcc gggccgtccc agtggcggctc	3480

-continued

---

```

tggctcgttt ccgctttggc cgcagggaca ggaatagcag gaggtgtcac cgggtcctta 3540
tccctggcct ccagtagaag tctcctgtcc gaagtggaca aggatatttc ccacctcaca 3600
cgggccattg taaaaaacca ccaaaacatt ctctgagtgg cccaatatgc cgcccaaac 3660
aggcgagggg tagacctctt gttctgggaa caaggggggc tgtgtaaagc gatacaagaa 3720
caatgctgct tctcaacat cagcaatacc catatttcag tcttacaaga gcgacccct 3780
ctagaaactc gggtaactac tggatggggc ttaaattggg atctaggact ctcccagtgg 3840
gcccgtgagg ctctccagac tggattacc cttttggccc tcttctgtt aatcatcatc 3900
ctcgggcctt gcattattcg ccagctgcaa gccctcccc agaggctaca gcagcgacct 3960
gaccagtacc ctctctcaa ccctgagacc cttttataat aactccgcca atacacccaa 4020
caggteccca tggttgacct ctctaccgtt caccaccgg cactccgcta gacctgacga 4080
gtcccccat atgtccaaag tctgttccaa gccagctgat aaccgaaata attctcctaa 4140
gttatggtta cattctctct ccagatcctt cctttccttc tctaatacat caatatagcc 4200
ttgcaacaag tcacaatacc cctcaaaccc cagcaggctc atgcacttcc gttgtgtgatg 4260
acgcgectct ctctctttgc gcttctctc cctctctctc aatcgctccc tccgcccgc 4320
ctccttttcc tctgtttctc gcaggagccg ctgaatctcc gctgctcgt ccaccagggc 4380
cctcaggcga gacttccggg taccatcatt ggcgctccc gacccagggg ggcggccttt 4440
gcgcgcacga cgagcggccc taccaggeat ctctctggt gttgagacct tctttgcccg 4500
atcctctgat gataaccccc taaaaaatc tataaaaaat tcccgttat tttttcagc 4560
ccacttccca ggattcgggc agagcctctt ctatggatac cccgtctatg tgtttggcga 4620
ttgtgttcaa gcgattggt gcccctctc cgggtggatta tgcctcccc gectacatcg 4680
ccacgcctc ctggccacct gcccggagca ccagatcacc tgggacccca tcatggagc 4740
agttgtcggc tcgctctcc aataccttat cctcgcctc cctccttcc ccaccaacg 4800
aacctcaag accctcaaag tccttaccac accaaccact cctgtcacc ccaaggttcc 4860
accctcctc tttcagtcg tgccggaggca cagcccctac cgcaacggat gtcttgaaac 4920
aaccttggg gagcagctcc cctccttgc atttctgag ccaggcctca ggccecaaaa 4980
cgtctacacc atctggggaa agaccatagt gtgtctatac atctaccagc tgtcccctcc 5040
catgacctgg cccctcattc cccatgtcat attttgcaac cccaggcagc ttggcgcttt 5100
tctaaagcaat gtgccccca agcgattaga agaactcctc taaaaacttt atctacacac 5160
cggcgccata atcctctgc cggaagacgc cctgcctacc accctatttc agcctgttcg 5220
agcacctgt gtccaaacta cctggaacac aggacttctc ccataccagc caaacctgac 5280
taccctggc ctgatatgga cctttaatga tgggtctct 5320

```

&lt;210&gt; SEQ ID NO 54

&lt;211&gt; LENGTH: 485

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

&lt;400&gt; SEQUENCE: 54

```

Met Gly Asn Val Leu Phe Leu Thr Leu Leu Ala Thr Leu Gly Ile Pro
1           5           10           15

```

-continued

---

Val	Leu	Gln	Ala	Ser	Arg	Cys	Thr	Ile	Thr	Val	Gly	Ile	Ser	Ser	Tyr
			20					25					30		
His	Ser	Ser	Pro	Cys	Ser	Pro	Ala	Gln	Pro	Leu	Cys	Thr	Trp	Ala	Leu
		35					40					45			
Asp	Leu	Val	Ser	Ile	Thr	Lys	Asp	Gln	Leu	Leu	Tyr	Pro	Pro	Cys	Gln
	50					55					60				
Asn	Leu	Ile	Thr	Tyr	Ser	Asn	Tyr	His	Lys	Thr	Tyr	Ser	Leu	Tyr	Leu
	65				70					75					80
Phe	Pro	His	Trp	Val	Gln	Lys	Pro	Leu	Arg	Arg	Gly	Leu	Gly	Tyr	Tyr
				85					90					95	
Ser	Ala	Ser	Tyr	Ser	Asp	Pro	Cys	Ser	Leu	Gln	Cys	Pro	Tyr	Leu	Gly
			100					105					110		
Ser	Gln	Ser	Trp	Thr	Cys	Pro	Tyr	Thr	Gly	Pro	Val	Ser	Ser	Pro	Thr
		115					120						125		
Trp	Arg	Phe	Ser	Thr	Asp	Val	Asn	Phe	Thr	Gln	Glu	Val	Ser	Arg	Val
	130					135					140				
Ser	Leu	Lys	Leu	His	Phe	Ser	Lys	Cys	Gly	Ser	Ser	Leu	Thr	Leu	Leu
	145				150					155					160
Ile	Asp	Ala	Pro	Gly	Tyr	Asp	Pro	Leu	Trp	Tyr	Leu	Thr	Ser	Glu	Pro
				165					170					175	
Thr	Gln	Glu	Pro	Pro	Thr	Pro	Pro	Pro	Leu	Val	Ser	Asp	Ser	Asp	Leu
			180					185						190	
Glu	His	Val	Leu	Thr	Pro	Ser	Ala	Ser	Trp	Ala	Ser	Lys	Met	Leu	Thr
		195					200						205		
Leu	Ile	His	Leu	Thr	Leu	Gln	Ser	Thr	Asn	Tyr	Ser	Cys	Met	Val	Cys
	210					215						220			
Ile	Asp	Arg	Ala	Ser	Leu	Ser	Ser	Trp	His	Val	Leu	Tyr	Thr	Pro	Asn
	225				230					235					240
Ile	Ser	Ser	Asn	Ala	Pro	Ser	Lys	Pro	Ile	Val	Arg	Pro	Ser	Leu	Ala
			245					250						255	
Leu	Ser	Ala	Pro	Arg	Pro	Gln	Pro	Phe	Pro	Trp	Thr	His	Cys	Tyr	Gln
		260						265						270	
Pro	Gln	Val	Gln	Ala	Val	Thr	Thr	Ala	Lys	Cys	Asn	Asn	Ser	Ile	Ile
		275					280						285		
Leu	Pro	Pro	Phe	Ser	Leu	Ser	Pro	Leu	Pro	Gly	Ala	Pro	Leu	Thr	Arg
	290					295					300				
Arg	Arg	Arg	Ala	Val	Pro	Val	Ala	Val	Trp	Leu	Val	Ser	Ala	Leu	Ala
	305				310					315					320
Ala	Gly	Thr	Gly	Ile	Ala	Gly	Gly	Val	Thr	Gly	Ser	Leu	Ser	Leu	Ala
			325						330					335	
Ser	Ser	Arg	Ser	Leu	Leu	Ser	Glu	Val	Asp	Lys	Asp	Ile	Ser	His	Leu
			340					345						350	
Thr	Arg	Ala	Ile	Val	Lys	Asn	His	Gln	Asn	Ile	Leu	Arg	Val	Ala	Gln
		355					360						365		
Tyr	Ala	Ala	Gln	Asn	Arg	Arg	Gly	Leu	Asp	Leu	Leu	Phe	Trp	Glu	Gln
	370					375						380			
Gly	Gly	Leu	Cys	Lys	Ala	Ile	Gln	Glu	Gln	Cys	Cys	Phe	Leu	Asn	Ile
	385				390						395				400
Ser	Asn	Thr	His	Ile	Ser	Val	Leu	Gln	Glu	Arg	Pro	Pro	Leu	Glu	Thr
			405						410					415	
Arg	Val	Thr	Thr	Gly	Trp	Gly	Leu	Asn	Trp	Asp	Leu	Gly	Leu	Ser	Gln

-continued

---

```

                420                425                430
Trp Ala Arg Glu Ala Leu Gln Thr Gly Ile Thr Leu Leu Ala Leu Leu
      435                440                445

Leu Leu Ile Ile Ile Leu Gly Pro Cys Ile Ile Arg Gln Leu Gln Ala
      450                455                460

Leu Pro Gln Arg Leu Gln Gln Arg Pro Asp Gln Tyr Pro Leu Leu Asn
      465                470                475                480

Pro Glu Thr Pro Leu
      485

<210> SEQ ID NO 55
<211> LENGTH: 307
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
      note=synthetic construct

<400> SEQUENCE: 55
Met Gly Asn Val Leu Phe Leu Thr Leu Leu Ala Thr Leu Gly Ile Pro
 1                5                10                15
Val Leu Gln Ala Ser Arg Cys Thr Ile Thr Val Gly Ile Ser Ser Tyr
 20                25                30
His Ser Ser Pro Cys Ser Pro Ala Gln Pro Leu Cys Thr Trp Ala Leu
 35                40                45
Asp Leu Val Ser Ile Thr Lys Asp Gln Leu Leu Tyr Pro Pro Cys Gln
 50                55                60
Asn Leu Ile Thr Tyr Ser Asn Tyr His Lys Thr Tyr Ser Leu Tyr Leu
 65                70                75                80
Phe Pro His Trp Val Gln Lys Pro Leu Arg Arg Gly Leu Gly Tyr Tyr
 85                90                95
Ser Ala Ser Tyr Ser Asp Pro Cys Ser Leu Gln Cys Pro Tyr Leu Gly
 100               105               110
Ser Gln Ser Trp Thr Cys Pro Tyr Thr Gly Pro Val Ser Ser Pro Thr
 115               120               125
Trp Arg Phe Ser Thr Asp Val Asn Phe Thr Gln Glu Val Ser Arg Val
 130               135               140
Ser Leu Lys Leu His Phe Ser Lys Cys Gly Ser Ser Leu Thr Leu Leu
 145               150               155               160
Ile Asp Ala Pro Gly Tyr Asp Pro Leu Trp Tyr Leu Thr Ser Glu Pro
 165               170               175
Thr Gln Glu Pro Pro Thr Pro Pro Pro Leu Val Ser Asp Ser Asp Leu
 180               185               190
Glu His Val Leu Thr Pro Ser Ala Ser Trp Ala Ser Lys Met Leu Thr
 195               200               205
Leu Ile His Leu Thr Leu Gln Ser Thr Asn Tyr Ser Cys Met Val Cys
 210               215               220
Ile Asp Arg Ala Ser Leu Ser Ser Trp His Val Leu Tyr Thr Pro Asn
 225               230               235               240
Ile Ser Ser Asn Ala Pro Ser Lys Pro Ile Val Arg Pro Ser Leu Ala
 245               250               255
Leu Ser Ala Pro Arg Pro Gln Pro Phe Pro Trp Thr His Cys Tyr Gln
 260               265               270

```

-continued

---

Pro Gln Val Gln Ala Val Thr Thr Ala Lys Cys Asn Asn Ser Ile Ile  
 275 280 285

Leu Pro Pro Phe Ser Leu Ser Pro Leu Pro Gly Ala Pro Leu Thr Arg  
 290 295 300

Arg Arg Arg  
 305

<210> SEQ ID NO 56  
 <211> LENGTH: 178  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 56

Ala Val Pro Val Ala Val Trp Leu Val Ser Ala Leu Ala Ala Gly Thr  
 1 5 10 15

Gly Ile Ala Gly Gly Val Thr Gly Ser Leu Ser Leu Ala Ser Ser Arg  
 20 25 30

Ser Leu Leu Ser Glu Val Asp Lys Asp Ile Ser His Leu Thr Arg Ala  
 35 40 45

Ile Val Lys Asn His Gln Asn Ile Leu Arg Val Ala Gln Tyr Ala Ala  
 50 55 60

Gln Asn Arg Arg Gly Leu Asp Leu Leu Phe Trp Glu Gln Gly Gly Leu  
 65 70 75 80

Cys Lys Ala Ile Gln Glu Gln Cys Cys Phe Leu Asn Ile Ser Asn Thr  
 85 90 95

His Ile Ser Val Leu Gln Glu Arg Pro Pro Leu Glu Thr Arg Val Thr  
 100 105 110

Thr Gly Trp Gly Leu Asn Trp Asp Leu Gly Leu Ser Gln Trp Ala Arg  
 115 120 125

Glu Ala Leu Gln Thr Gly Ile Thr Leu Leu Ala Leu Leu Leu Ile  
 130 135 140

Ile Ile Leu Gly Pro Cys Ile Ile Arg Gln Leu Gln Ala Leu Pro Gln  
 145 150 155 160

Arg Leu Gln Gln Arg Pro Asp Gln Tyr Pro Leu Leu Asn Pro Glu Thr  
 165 170 175

Pro Leu

<210> SEQ ID NO 57  
 <211> LENGTH: 848  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 57

Gly His Ile Glu Pro Tyr Ser Gly Pro Gly Asn Asn Pro Val Phe Pro  
 1 5 10 15

Val Lys Lys Pro Asn Gly Lys Trp Arg Phe Ile His Asp Leu Arg Ala  
 20 25 30

Thr Asn Ala Ile Thr Thr Thr Leu Ala Ser Pro Ser Pro Gly Pro Pro  
 35 40 45

Asp Leu Thr Ser Leu Pro Gln Ala Leu Pro His Leu Gln Thr Ile Asp

-continued

50					55					60					
Leu	Thr	Asp	Ala	Phe	Phe	Gln	Ile	Pro	Leu	Pro	Lys	Arg	Phe	Gln	Pro
65					70					75					80
Tyr	Phe	Ala	Phe	Thr	Ile	Pro	Gln	Pro	Leu	Asn	His	Gly	Pro	Gly	Ser
				85					90					95	
Arg	Tyr	Ala	Trp	Thr	Val	Leu	Pro	Gln	Gly	Phe	Lys	Asn	Ser	Pro	Thr
			100					105						110	
Leu	Phe	Glu	Gln	Gln	Leu	Ala	Ser	Val	Leu	Gly	Pro	Ala	Arg	Lys	Ala
		115					120							125	
Phe	Pro	Thr	Ser	Val	Ile	Val	Gln	Tyr	Met	Asp	Asp	Ile	Leu	Leu	Ala
	130					135					140				
Cys	Pro	Ser	Gln	His	Glu	Leu	Asp	Gln	Leu	Ala	Thr	Leu	Thr	Ala	Gln
145				150						155					160
Leu	Leu	Ser	Ser	His	Gly	Leu	Pro	Val	Ser	Gln	Glu	Lys	Thr	Gln	Arg
				165					170					175	
Thr	Pro	Gly	Lys	Ile	His	Phe	Leu	Gly	Gln	Ile	Ile	His	Pro	Asp	His
			180					185						190	
Ile	Thr	Tyr	Glu	Thr	Thr	Pro	Thr	Ile	Pro	Ile	Lys	Ala	His	Trp	Thr
		195					200						205		
Leu	Thr	Glu	Leu	Gln	Thr	Leu	Leu	Gly	Glu	Leu	Gln	Trp	Val	Ser	Lys
	210					215					220				
Gly	Thr	Pro	Val	Leu	Arg	Glu	His	Leu	His	Cys	Leu	Tyr	Ser	Ala	Leu
225				230							235				240
Arg	Gly	Leu	Lys	Asp	Pro	Arg	Asp	Thr	Ile	Thr	Leu	Arg	His	Pro	His
				245					250					255	
Leu	His	Ala	Leu	His	Asn	Ile	Gln	Gln	Ala	Leu	His	His	Asn	Cys	Arg
		260						265						270	
Gly	Arg	Leu	Asp	Ser	Thr	Leu	Pro	Leu	Leu	Gly	Leu	Ile	Phe	Leu	Ser
		275					280						285		
Pro	Ser	Gly	Thr	Thr	Ser	Val	Leu	Phe	Gln	Thr	Asn	His	Lys	Trp	Pro
		290				295					300				
Leu	Val	Trp	Leu	His	Ala	Pro	His	Pro	Pro	Thr	Ser	Leu	Cys	Pro	Trp
305				310						315				320	
Gly	His	Ile	Leu	Ala	Cys	Thr	Val	Leu	Thr	Leu	Asp	Lys	Tyr	Ala	Leu
			325						330					335	
Gln	His	Tyr	Gly	Gln	Leu	Cys	Lys	Ser	Phe	His	His	Asn	Met	Ser	Thr
			340						345					350	
Gln	Ala	Leu	His	Asp	Phe	Val	Lys	Asn	Ser	Ser	His	Pro	Ser	Val	Ala
		355					360						365		
Ile	Leu	Ile	His	His	Met	His	Arg	Phe	Cys	Asp	Leu	Gly	Arg	Gln	Pro
	370					375						380			
Pro	Gly	Pro	Trp	Arg	Thr	Leu	Leu	Gln	Leu	Pro	Ala	Leu	Leu	Arg	Glu
385				390							395			400	
Pro	Gln	Leu	Leu	Arg	Pro	Ala	Phe	Ser	Leu	Ser	Pro	Val	Val	Ile	Asp
				405					410					415	
Gln	Ala	Pro	Cys	Leu	Phe	Ser	Asp	Gly	Ser	Pro	Gln	Lys	Ala	Ala	Tyr
			420					425						430	
Val	Ile	Trp	Asp	Lys	Val	Ile	Leu	Ser	Gln	Arg	Ser	Val	Pro	Leu	Pro
		435					440						445		
Pro	His	Ala	Asn	Asn	Ser	Ala	Gln	Lys	Gly	Glu	Leu	Val	Gly	Leu	Leu
		450				455							460		

-continued

---

Leu Gly Leu Gln Ala Ala Gln Pro Trp Pro Ser Leu Asn Ile Phe Leu  
 465 470 475 480  
 Asp Ser Lys Phe Leu Ile Arg Tyr Leu Gln Ser Leu Ala Ser Gly Ala  
 485 490 495  
 Phe Gln Gly Ser Ser Thr His His Arg Leu Gln Ala Ser Leu Pro Thr  
 500 505 510  
 Leu Leu Gln Gly Lys Val Val Tyr Leu His His Thr Arg Ser His Thr  
 515 520 525  
 Gln Leu Pro Asp Pro Ile Ser Thr Leu Asn Glu Tyr Thr Asp Ser Leu  
 530 535 540  
 Ile Val Ala Pro Val Thr Pro Leu Lys Pro Glu Gly Leu His Ala Leu  
 545 550 555 560  
 Thr His Cys Asn Gln Gln Ala Leu Val Ser His Gly Ala Thr Pro Ala  
 565 570 575  
 Gln Ala Lys Gln Leu Val Gln Ala Cys Arg Thr Cys Gln Ile Ile Asn  
 580 585 590  
 Pro Gln His His Met Pro Arg Gly His Ile Arg Arg Gly His Phe Pro  
 595 600 605  
 Asn His Thr Trp Gln Gly Asp Val Thr His Leu Lys His Lys Arg Thr  
 610 615 620  
 Arg Tyr Cys Leu His Val Trp Val Asp Thr Phe Ser Gly Ala Val Ser  
 625 630 635 640  
 Cys Val Cys Lys Lys Lys Glu Thr Ser Ser Asp Leu Ile Lys Thr Leu  
 645 650 655  
 Leu His Ala Ile Ser Val Leu Gly Lys Pro Phe Ser Val Asn Thr Asp  
 660 665 670  
 Asn Gly Pro Ala Tyr Leu Ser Gln Glu Phe His Glu Phe Cys Thr Thr  
 675 680 685  
 Leu Cys Ile Lys His Ser Thr His Ile Pro Tyr Asn Pro Thr Ser Ser  
 690 695 700  
 Gly Leu Val Glu Arg Thr Asn Gly Ile Leu Lys Thr Leu Leu Tyr Lys  
 705 710 715 720  
 Tyr Phe Leu Asp His Pro Asp Leu Pro Leu Glu Ser Ala Val Ser Lys  
 725 730 735  
 Ala Leu Trp Thr Ile Asn His Leu Asn Val Met Arg Pro Cys Gly Lys  
 740 745 750  
 Thr Arg Trp Gln Leu His His Thr Pro Pro Leu Pro Pro Ile Ser Glu  
 755 760 765  
 Ser Ile Gln Thr Thr Pro Thr Arg Leu His Trp Tyr Tyr Tyr Lys Thr  
 770 775 780  
 Pro Gly Leu Thr Asn Gln Arg Trp Lys Gly Pro Val Gln Ser Leu Gln  
 785 790 795 800  
 Glu Ala Ala Gly Ala Ala Leu Leu Gln Val Ser Asp Gly Ser Pro Gln  
 805 810 815  
 Trp Ile Pro Trp Arg Leu Leu Lys Lys Thr Val Cys Pro Lys Pro Asp  
 820 825 830  
 Asp Pro Glu Pro Ala Gly His Val Glu Thr Asp His Gln His His Gly  
 835 840 845

&lt;210&gt; SEQ ID NO 58

&lt;211&gt; LENGTH: 90

-continued

---

```

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
      note=synthetic construct

<400> SEQUENCE: 58

Leu Ala Ile Ser Asn Leu Thr Leu Asp Gln Ala Thr Thr Gln Phe Ser
1          5          10          15

Leu Leu Lys Asn Pro Thr Ala Ser Gly Asp Leu Ser Met Thr Ser Gly
20          25          30

Pro Leu Met Pro Ser Pro Leu Pro Leu Pro Arg Pro Pro Pro Ala Pro
35          40          45

Leu Ile Leu Pro Ala Cys His Arg Pro Cys Pro Ile Phe Arg Pro Ser
50          55          60

Ile Ser Arg Thr Leu Ser Ser Arg Phe Pro Ser Gln Ser Asp Ser Ser
65          70          75          80

Pro Thr Ser Pro Leu Pro Ser Pro Ser His
85          90

```

```

<210> SEQ ID NO 59
<211> LENGTH: 273
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
      note=synthetic construct

```

```

<400> SEQUENCE: 59

ctggccatat cgaaccttac tctggaccag gcaacaaccc agttttccct gttaaaaaac   60
ccaacggcaa gtggcgattt atccatgacc tcaggggcac taatgccatc accactaccc   120
ttgcctcgcc ctcccccgcc ccccctgata ttaccagcct gccacaggcc ttgccccatc   180
ttcagaccat cgatctcagc gaagctttct tccagattcc cctcccaaag cgattccagc   240
cctacttcgc ctttaccatc cccagccat taa                               273

```

```

<210> SEQ ID NO 60
<211> LENGTH: 170
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial
      Sequence;note=synthetic construct

```

```

<400> SEQUENCE: 60

Met Pro Lys Thr Arg Arg Pro Arg Thr Arg Arg Ala Arg Arg Asn Arg
1          5          10          15

Pro Pro Thr Pro Trp Pro Thr Ser Gln Asp Ser Gly Arg Ala Ser Ser
20          25          30

Met Asp Thr Pro Ser Met Cys Leu Ala Ile Val Phe Lys Pro Ile Gly
35          40          45

Ala Pro Ser Pro Val Asp Tyr Ala Pro Pro Ala Tyr Ile Ala Thr Pro
50          55          60

Ser Trp Pro Pro Ala Pro Ser Thr Arg Ser Pro Gly Thr Pro Ser Met
65          70          75          80

Asp Glu Leu Ser Ala Arg Leu Ser Asn Thr Leu Ser Leu Ala Ser Pro
85          90          95

```

-continued

---

Pro Ser Pro Pro Asn Glu Pro Pro Arg Pro Ser Lys Ser Leu Pro His  
 100 105 110

Gln Pro Leu Leu Ser Pro Pro Arg Phe His Pro Pro Ser Phe Ser Pro  
 115 120 125

Cys Gly Gly Thr Ala Pro Thr Ala Thr Asp Val Leu Lys Gln Pro Leu  
 130 135 140

Glu Ser Ser Ser Pro Pro Leu His Phe Leu Ser Gln Ala Ser Gly Pro  
 145 150 155 160

Lys Thr Ser Thr Pro Ser Gly Glu Arg Pro  
 165 170

<210> SEQ ID NO 61  
 <211> LENGTH: 512  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 61

atgcccaaaa cccgacgacc ccgaaccgc agggcacgtc gaaacagacc accaacacca 60  
 tgcccacttc ccaggattcg ggcagagcct cctctatgga taccocgtct atgtgtttgg 120  
 cgattgtgtt caagccgatt ggtgcccat ctccggtgga ttatgctccc cccgcctaca 180  
 tgcaccgccc ctctggcca cctgcccga gcaccagatc acctgggacc ccatcgatgg 240  
 acgagttgtc ggetgcctc tccaatacct tatccctcgc ctcccctcct tccccacca 300  
 acgaacctcc aagacctca aagtccttac cccaccaacc actcctgtca ccccaaggt 360  
 tccacctcc tctttcagt ccgtgaggag gcacagcccc taccgcaacg gatgtttga 420  
 aacaacctt ggagagcagc tcccctcct tgcatttct gagccaggcc tcaggcccca 480  
 aaactgtac accatctggg gaaagacat ag 512

<210> SEQ ID NO 62  
 <211> LENGTH: 255  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 62

Met Ala His Phe Pro Gly Phe Gly Gln Ser Leu Leu Tyr Gly Tyr Pro  
 1 5 10 15

Val Tyr Val Phe Gly Asp Cys Val Gln Ala Asp Trp Cys Pro Ile Ser  
 20 25 30

Gly Gly Leu Cys Ser Pro Arg Leu His Arg His Ala Leu Leu Ala Thr  
 35 40 45

Cys Pro Glu His Gln Ile Thr Trp Asp Pro Ile Asp Gly Arg Val Val  
 50 55 60

Gly Ser Pro Leu Gln Tyr Leu Ile Pro Arg Leu Pro Ser Phe Pro Thr  
 65 70 75 80

Gln Arg Thr Ser Lys Thr Leu Lys Val Leu Thr Pro Pro Thr Thr Pro  
 85 90 95

Val Thr Pro Lys Val Pro Pro Ser Phe Phe Gln Ser Val Arg Arg His  
 100 105 110

-continued

---

Ser Pro Tyr Arg Asn Gly Cys Leu Glu Thr Thr Leu Gly Glu Gln Leu  
 115 120 125

Pro Ser Leu Ala Phe Pro Glu Pro Gly Leu Arg Pro Gln Asn Val Tyr  
 130 135 140

Thr Ile Trp Gly Lys Thr Ile Val Cys Leu Tyr Ile Tyr Gln Leu Ser  
 145 150 155 160

Pro Pro Met Thr Trp Pro Leu Ile Pro His Val Ile Phe Cys Asn Pro  
 165 170 175

Arg Gln Leu Gly Ala Phe Leu Ser Asn Val Pro Pro Lys Arg Leu Glu  
 180 185 190

Glu Leu Leu Tyr Lys Leu Tyr Leu His Thr Gly Ala Ile Ile Ile Leu  
 195 200 205

Pro Glu Asp Ala Leu Pro Thr Thr Leu Phe Gln Pro Val Arg Ala Pro  
 210 215 220

Cys Val Gln Thr Thr Trp Asn Thr Gly Leu Leu Pro Tyr Gln Pro Asn  
 225 230 235 240

Leu Thr Thr Pro Gly Leu Ile Trp Thr Phe Asn Asp Gly Ser Pro  
 245 250 255

<210> SEQ ID NO 63  
 <211> LENGTH: 559  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 63

taactcggcc aatacaccca acagggtccc atgggtgacc cctctaccgt tcaccacccc 60  
 gcactcggct agacctgacg agtcccccca tatgtccaaa gtctgttcca agccagctga 120  
 taaccgaaat aatttccta agttatggtt acattcctcc tccagatcct tcctttcctt 180  
 ctctaataca tcaatatagc cttgcaacaa gtcacaatac cctcacaacc ccagcaggtc 240  
 catgcacttc cgttgttgat gacgogcctc tctctccttg cgettcctct cctctcctg 300  
 caatcgctcc ctccgcccgg cctccttttc ctctgtttct cgcaggagcc gctgaatctc 360  
 cgctgctcg tccaccaggg ccctcaggcg agacttccgg gtaccatcat tggcgcctcc 420  
 cgaccccagg gggcggcctt tgcgcgacg acgagcggc ctaccaggca tctcctctgg 480  
 tgttgagacc ttctttgccc gatcctctga tgataacccc ctaaaaaatt ctataaaaa 540  
 ttccccgtta tttttttca 559

<210> SEQ ID NO 64  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 64

gattcccctc ccaaagcgat 20

<210> SEQ ID NO 65  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence

-continued

---

<220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 65

tgacggatgt ggggaaggct 20

<210> SEQ ID NO 66  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 66

ttccccaagg cttcaaaaac agccccacgc 30

<210> SEQ ID NO 67  
 <211> LENGTH: 50  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 67

Ser Phe Ser Phe Leu Leu Asp Ala Pro Gly Tyr Asp Pro Val Trp Phe  
 1 5 10 15

Leu Ser Ser Gln Ala Thr Gln Ala Pro Pro Thr Pro Ala Pro Leu Ile  
 20 25 30

Gln Asp Ser Asp Leu Gln His Ile Leu Glu Pro Ser Ile Pro Trp Ser  
 35 40 45

Ser Lys  
 50

<210> SEQ ID NO 68  
 <211> LENGTH: 48  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 68

Thr Leu Leu Ile Asp Ala Pro Gly Tyr Asp Pro Leu Trp Tyr Leu Thr  
 1 5 10 15

Ser Glu Pro Thr Gln Glu Pro Pro Thr Pro Pro Pro Leu Val Ser Asp  
 20 25 30

Ser Asp Leu Glu His Val Leu Thr Pro Ser Ala Ser Trp Ala Ser Lys  
 35 40 45

<210> SEQ ID NO 69  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence;  
 note=synthetic construct

<400> SEQUENCE: 69

ggtaagatcc cactgggtcg agc 23

---

-continued

---

<210> SEQ ID NO 70  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct  
  
<400> SEQUENCE: 70  
  
gaagccaggt ctcggtgac g 21

<210> SEQ ID NO 71  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct  
  
<400> SEQUENCE: 71  
  
cgctcccctg gagctctctc g 21

<210> SEQ ID NO 72  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct  
  
<400> SEQUENCE: 72  
  
gccacttccc attgggcttt ttgacgg 27

<210> SEQ ID NO 73  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct  
  
<400> SEQUENCE: 73  
  
gctctcaccg ataaagtaac aaacg 25

<210> SEQ ID NO 74  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct  
  
<400> SEQUENCE: 74  
  
ggtaggaaga ggctcctatg aacag 25

<210> SEQ ID NO 75  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct  
  
<400> SEQUENCE: 75

---

-continued

---

caggactgca taacatacga gaccctcc 28

<210> SEQ ID NO 76  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 76

cctatgaaca gggatcatcg actggg 26

<210> SEQ ID NO 77  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 77

cctaagcccc ccatgtccag ac 22

<210> SEQ ID NO 78  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 78

cgagagagct ccaggggagc g 21

<210> SEQ ID NO 79  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 79

cctactccct gtatgtattc cccattgg 29

<210> SEQ ID NO 80  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

<400> SEQUENCE: 80

gctcgacca gtgggatctt accgagtgg 29

<210> SEQ ID NO 81  
<211> LENGTH: 8791  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence;  
note=synthetic construct

---

-continued

---

<400> SEQUENCE: 81

tgacagggac aacgaccctc tcccaggggc gacagcaagc cccaaggac aaaactagca 60  
gggactagtc atcagccaaa aaggtcaact gtctcacaca aataaggatc cgaaggttct 120  
gacgtcccag ccagcctca aaaccaggaa atccatagaa atgcacctcg cccttaccca 180  
cttcccctat catgaaaaac aaaggctgtg acgactacce ccttcccacaa aaaatttgct 240  
taaaccatca ataaagacag cctagcctat ataagcatga ggatggttca ggagggggct 300  
cgctctcttg ccgatcgccc tgctcacctc gagtgtccat ctctgtgtca atcagttgag 360  
acgcgcgcgg ctgccggtct cctggtgtgc gcacctcctg aaccaccctt tgggtaagtc 420  
cccccttggc ccgagcttgg ctacgggttc tgtagtcgct cccagggaaag tctccgagac 480  
tgcccaagcc tctgcttgca aggtacggc cctccacccc tcttccgct cctgtttaat 540  
ctcttcgctc caaccgaaaa cgaaagcgc tccagctctc ttggcccggg gccaggcctg 600  
agccgcgcgg gcgcaccacc ttaagccgc tgtactcaaa cccctccggg aggggcccctt 660  
tacagtaggc gcccgctccc ccgggggaaa catacaagtg ggggctcgtc cgggatctgt 720  
tccgctctcg ccgttcccc cctcccacta tgggtcagac ccacacatcc agtcccgtcc 780  
ctaaggcccc cagggggctc tccaccacc actggcttaa tttcctgcag gcggcttacc 840  
gctgcaacc tggaccctcc gaattcgatt ttcaccagtt aagacgattc cttaaagctag 900  
cgctccaaac ccagctctgg ttaaacccta tgcattactc cctcctagcc ggctaatcc 960  
ccaaggggta ccccggtcgg gtgaccgaga tctgtaatat cctcctccgc gctcatccac 1020  
ccccagcgc ccggcaatt tccatgcca cggccaccgg cccggcccct gccccccagc 1080  
ctcaggaggc gcacacgccc ccccctatg cggagcctgc tgcgctccag tgccttccca 1140  
ttatgcactc ccacggggcc ccctcgagcc accgcccctg gcagatgaaa gacttacaag 1200  
ccattaaaca agaaattagc acctcagctc ccggcagtc tcaatttatg cataccattc 1260  
gacttgccat ccagcagttt gaccctacgg ctaaagatct acatgatctt ttgcagtact 1320  
tgtgctcgtc ccttattgtc tcccttcacc accaacagct acaagcactc attgtggagg 1380  
cagaaaccgc aggggtgaca ggttacaatc ctatggcagg gccctccgg gtacaagcaa 1440  
acaaccgcc ccagcaaggc ctccagagag aataccaaag tctttgctg gccgccttg 1500  
cggccctgcc tggtaacacc cgagatcctt cctgggcgc aatattgcaa ggctcgagg 1560  
aaccttattg tgctttgta gagegcctca atgcggccct cgataatggt ctacctgaag 1620  
gcacacccaa ggaaccatc ctgcggccgc tggcatactc caatgccaac aaagaatgcc 1680  
agaaactcct tcaggcgcgg ggcatacca acagtcccct tggcgaaatg ctccgagcct 1740  
gtcaggcttg gacaccaaag gataagacca aagttctagt agttcagccc cgtaaaacct 1800  
ctccaacaca accgtgcttc cgggtgtggaa aggtgggaca ctggagccga gactgcactc 1860  
agcctegccc cctcggggg ccctgcccc tatgtcagga cccatcccac tggagcgag 1920  
attgccccca gctaaaaacc ccgcccggag cagaagaacc cctcctagcg gatttgctg 1980  
cccttctccc ggagaaaaa aactcccag ggggggagaa ctagtctccc ccgaccccgg 2040  
taactgctc tcctgcttc ccctgtctc cctatggcag gcccaacaat ctctcctcaa 2100  
tattaaagtt tcttctctc atcgcacc cctggcatca caggcgtcc tggacaccgg 2160  
agccggcctc actgtcatgc cccaggtttt ggctcggggg ctcacggacc tccaggacac 2220

-continued

---

caccattctg ggggocggcg gtaaaaccca ctcccagttt aaactcctac ggtgtccggg	2280
acatgtatac ttgcccttcc gtagggctcc cgtgtccctt ccctcatgtc taattgacac	2340
caagaatgag tggaccatca tcggccggga cgtcctgcag caatgccagg gggcccttta	2400
cttaccggag gacctccggg ccccgaccca gttatccccg gtgaccaccc ctgcagtcac	2460
cggcttagaa catcttccag agccccaga ggtcagccag tttcctttaa acctgaacgc	2520
ctccaggccc taatagacct ggtctccaag gcaactggagg ctggccatat cgaaccttac	2580
tctggaccag gcaacaaccc agttttccct gttaaaaaac ccaacggcaa gtggcgattt	2640
atccatgacc tcaggggcac taatgccatc accactaccc ttgcctcgcc ctccccggc	2700
ccccctgac ttaccagcct gccacaggcc ttgccccatc ttcagaccat cgatctcacg	2760
gacgctttct tcagattcc cctcccaaag cgattccagc cctacttcgc ctttaccatc	2820
ccccagccat taaatcatgg gcctgggagc aggtacgctt ggacagtcct tccccaggc	2880
ttcaaaaaca gccccacgct ctttgagcaa cagctggcca gcgtactagg cccagcccga	2940
aaagccttcc ccacatccgt catcgtccaa tacatggacg acatcctctt ggcattgccc	3000
tcccagcacg aactagatca gctggccacc cttaccgcac agctattgtc ctctcatggt	3060
ctcccagttt cccaggaaaa aacccaaagc accccaggaa aaatacactt cctggggccaa	3120
atcatacatc cagatcacat cacctatgaa accacccccca ccatacccat taaggcacac	3180
tggacctga ctgaactgca aacctcctg ggggagctcc agtgggtctc caaggggact	3240
cctgtcctcc gagaacacct tcaactgtct tactcagcct tgagaggctc caaagacccc	3300
cgggacacta tcaccttctg tcactctcac ctccacgctc tccacaacat tcagcaagcc	3360
ctgcatacaca attgcgcggg tcgccttgac tctacgctcc ccctcctgg cctcatcttc	3420
ctcagtcacat ccggcacgac ctcagtcctc ttccagacaa atcataaatg gccctagtc	3480
tggctccacg cccccatcc cccgaccagc ctatgcccct gggggcacat actcgcctgc	3540
actgtactta cccttgacaa gtatgccttg cagcaactat gccaaactat caaatcattc	3600
catcataaca tgtccaccca ggcctacac gatttcgtaa aaaattcctc tcaccccagc	3660
gtcggccatc taattcacca catgcacggg ttctgtgacg tgggcagaca gccaccggga	3720
ccctggcgaa cctccttaca actcccggcc cttctccggg aaccacagct cctcaggcct	3780
gcattttccc tatcccaggt gggttatagat caggcccctt gtctgttctc tgatgggtct	3840
ccccaaaagg ccgcctatgt aatttgggac aaggctcatc tcagccagcg gtcgggtccc	3900
ctgccccccc atgccaataa ctcagcacia aagggggaat tagtccgact cctcttgggc	3960
ttgcaagccg cacagccctg gccatccctt aacattttcc tagactcaaa gttcctcatc	4020
cggtagctcc agtccctcgc ttccggggcc ttccaaggat catccacaca ccaccgtctc	4080
caggcgtccc tgcccacact cctccagggc aaggctcgtg atctccacca caccgcagc	4140
cacacccaat tgctgatcc catctcgacc ctcaatgaat ataccgactc tctcattgtc	4200
gcccccgtaa ccccctgaa gcctgagggc ctccatgccc tcaccactg caaccaacag	4260
gcctcgttt cccacggagc caccctgca caggctaage aactcgtgca ggcctgccc	4320
acctgtcaaa tcattaaccc tcaacaccac atgccgctg gccacatccg ccgcgccac	4380
ttccaaaacc acacatggca aggagatgct acccacctta agcacaacg gaccggatac	4440
tgctccacg tctgggtgga taccttctca ggtgcggtat cttgtgtctg caaaaagaaa	4500

-continued

---

gaaactagca gcgacottat caaaaccctc ctacatgcca tctccgtgct aggcaagccc	4560
ttctctgtta acacgggaaa tggaccgcgt tacctttctc aggagttcca cgaattctgt	4620
accaccctct gcatcaaaaca ctccaccocat attccctaca atccgacaag ttcaggcctg	4680
gtggagcgca caaatggcat tctcaagaca ctactataca aatatttctc agaccaccct	4740
gacctcccc tagaaagcgc ggtttcaaag gctctctgga ccattaacca tttaaatgtc	4800
atgcgcccct gtggaagac tcggtggcag ctccatcaca cccccccct gcctcctatt	4860
tccgagtcca tacaaccacc tcccaccagg ctacattggt actattacaa aaccctgga	4920
cttaccaacc agcgatggaa agggcccgtc caatctctcc aggaagcagc aggagcagct	4980
ctccttaag tcagtacggc ctcccccag tggatccctt ggcggctcct gaagaagact	5040
gtatgcccac aaccggacga ccccgaacct gcagggcacg tcgaacaga ccaccaaac	5100
catgggtaac gtactcttct taactttatt ggccaccctg ggcatcccag tacttcaggc	5160
cagccgggtg acaatcacgg taggtatctc ctctaccac tccagcccct gcagcccagc	5220
ccagccttta tgtacctggg ccctcgacct tgtgtccatc actaaggacc agctcctcta	5280
ccccccctgc caaaacctga tcacctatc caactaccac aagaactact ccctgtatct	5340
cttcccacac tgggtacaaa agccactccg ccgggggctt ggatactact cagcctccta	5400
ctctgatcct tgcctcctac aatgtcccta cctaggaagt caatcatgga cttgccccta	5460
tactggccct gtctcgagcc caacttggag attctccaca gatgtaaatt tcaccaaga	5520
agtcagccgt gtctccctaa aacttcattt ctccaaatgt ggctcctcct taactctgtt	5580
aatagatgcc ccgggttacg atccgctgtg gtacctcaca tccgagccta ctcaggaacc	5640
cccaaccctc ccgcccactg tcagcgactc agacctagag catgtcctga ctccctcggc	5700
ctcctgggcc tccaagatgc tgacctcat ccacctaacc ttgcagagca ccaactatc	5760
ctgtatggtc tgtattgacc gcgcccagcct ctcttctcgg cacgtattat acactcccaa	5820
catctctagt aatgccccct caaaaccocat cgtccgcccct tcccttgccc tatccgcccc	5880
gcgaccacag cccttccccct ggaccattg ctatcaacca caggtgcaag ctgtaaccac	5940
cgcaaagtgc aataattcca tcatacttcc cccattttct ctctctcctc tgctcgggtc	6000
ccctctcact aggcgacgcc gggccgctcc agtggcggtc tggctcgttt ccgctttggc	6060
cgcagggaca ggaatagcag gaggtgtcac cgggtcctta tccctggcct ccagtagaag	6120
tctcctgtcc gaagtggaca aggatatttc ccacctcaca cgggccattg taaaaacca	6180
ccaaaacatt cttcgagtgg cccaatatgc cgcccaaac aggcgagggt tagacctcct	6240
gttctgggaa caaggggggc tgtgtaaagc gatacaagaa caatgctgct tcctcaacat	6300
cagcaatacc catatttcag tcttacaaga gcgaccccct ctagaacctc gggtaactac	6360
tggatggggc ttaaattggg atctaggact ctcccagtg gcccgtgagg ctctccagac	6420
tggattacc cttttggccc tccctctggt aatcatcatc ctccggccct gcattattcg	6480
ccagctgcaa gccctcccc agaggetaca gcagcgacct gaccagtacc ctctcctcaa	6540
ccctgagacc cctttataat aactccgcca atacaccaa caggtcccca tggttgacc	6600
ctctaccgtt cacccaccg cactccgcta gacctgacga gtcccccat atgtccaaag	6660
tctgttccaa gccagctgat aaccgaaata attctcctaa gttatggtta cattcctcct	6720
ccagatcctt ccttctctc tctaatacat caatatagcc ttgcaacaag tcacaatacc	6780

-continued

---

```

cctcaaaacc cagcaggtcc atgcaactcc gttggtgatg acgcgcctct ctctcettgc 6840
gcttcctctc cctctcctgc aatcgctccc tccgccgcgc ctctttttcc tctgtttctc 6900
gcaggagccg ctgaatctcc gcctgctcgt ccaccagggc cctcaggcga gacttcgggg 6960
taccatcatt ggcgcctccc gaccccaggg ggccgccttt gcgcgcacga cgagcgcgc 7020
taccaggcat ctctctcgtt gttgagacct tctttgcccg atcctctgat gataaccccc 7080
taaaaaatc tataaaaaat tccccttat tttttcagc ccacttcca ggattcgggc 7140
agagcctcct ctatggatac cccgtctatg tgtttggcga ttgtgttcaa gccgattggt 7200
gccccatctc cggtgattta tgctccccc gctacatcg ccacgcctc ctggccacct 7260
gccccgagca ccagatcacc tgggacccca tcgatggacg agttgtcggc tcgcctctcc 7320
aataccttat cctcgcctc ccctcctcc ccaccaacg aacctccaag acctcaaag 7380
tccttcccc accaaccact cctgtcacc ccaaggttcc accctcctc tttcagtcgg 7440
tgccgaggca cagccctacc cgcaacggat gtcttgaaac aaccttggga gagcagctcc 7500
cctcccttgc atttctgag ccaggcctca ggccccaaaa cgtctacacc atctggggaa 7560
agaccatagt gtgtctatac atctaccagc tgtcccctcc catgacctgg cccctcattc 7620
cccatgtcat attttgcaac ccaggcagc ttggcgcttt tctaagcaat gtgccccca 7680
agcgattaga agaactcctc tacaactttt atctacacac cggcgccata atcatcctgc 7740
cggaagagc cctgcctacc acctatttc agcctgttcg agcaccctgt gtccaaacta 7800
cctggaacac aggacttctc ccataccagc caaacctgac taccctggc ctgatatgga 7860
cctttaatga tgggtctcct atgatttcag gacctgccc taaggcaggg cagccatcct 7920
tggtagtaca gtctcacta ctaactctcg agagatttca aaccaaagcc tatcatcct 7980
cttacctcct ctcccaccaa ttgatacagt attcctcctt ccatacctc tacttactct 8040
ttgatgaata tactactatc cccttctctc tactatttaa ggaaaaagag ggagatgaca 8100
gggacaacga ccctctcca gggcgacag caagcccca aggacaaaac tagcagggac 8160
tagtcatcag caaaaaaggt caactgtctc acacaaataa ggatccgaag gttctgacgt 8220
cccagcccag cctcaaaacc aggaaatcca tagaaatgca cctcgcctt acccacttcc 8280
cctatcatga aaaaacaaag ctgtgacgac taccctctc ccaaaaaat ttgcttaaac 8340
catcaataaa gacagcctag cctatataag catgaggatg gttcaggagg gggctcgcctc 8400
tcttgccgat cgccctgctc acctogagtg tccatctcct ggccaatcag ttgagacgcc 8460
gccggctgcc ggtctcctgg ttgtgcacc tctgaacca ccccttgggt aagtccccc 8520
ttggtccgag cttggctaca gtttctgtag tcgctcccag ggaagtctcc gagactgcc 8580
aagcctctgc ttgcaaggct acggccctcc acccctctc cgcgtccgtg ttaatctctt 8640
cgcgccaaac gaaaacgaaa gcgcctccag ctctcttggc cggggccag gcctgagccg 8700
cggggcgca ccaccttaag cccgctgtac taaaaccct cggggagggg ccttttacag 8760
taggcgcccg tcccccggg ggaacatac a 8791

```

&lt;210&gt; SEQ ID NO 82

&lt;211&gt; LENGTH: 7

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence;

-continued

---

```

note=synthetic construct

<400> SEQUENCE: 82

gttaaac                                     7

<210> SEQ ID NO 83
<211> LENGTH: 7
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

<400> SEQUENCE: 83

tttaaac                                     7

<210> SEQ ID NO 84
<211> LENGTH: 61
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence;
note=synthetic construct

<400> SEQUENCE: 84

Glu Arg Arg Arg Arg Cys Arg Glu Glu Leu Glu Glu Arg Lys Arg Gln
1           5           10           15

Lys Lys Glu Arg Arg Gln Gln Leu Asp Cys Ile Asp Met Leu Gly Phe
           20           25           30

Glu Gly Phe Cys Asp Leu Leu Glu Gly Tyr Ile Asp Phe Leu Glu Arg
           35           40           45

Glu Ser Gln Gln Leu Arg Ala Gly Cys Glu Glu Ser Leu
           50           55           60

```

---

1. An isolated primate T-lymphotropic virus (PTLV) polypeptide, wherein the polypeptide is a HTLV-3 or HTLV-4 gag, pol, env, tax, rex or pro protein selected from the group consisting of

- (i) a HTLV-3 gag protein encoded by a nucleic acid sequence at least 91.5% identical to SEQ ID NO: 35;
- (ii) a HTLV-3 pol protein encoded by a nucleic acid sequence at least 92.8% identical to SEQ ID NO: 1;
- (iii) a HTLV-3 env protein encoded by a nucleic acid sequence at least 92.5% identical to SEQ ID NO: 3;
- (iv) a HTLV-3 tax protein encoded by a nucleic acid sequence at least 94.2% identical to SEQ ID NO: 5;
- (v) a HTLV-3 rex protein encoded by a nucleic acid sequence comprising SEQ ID NO: 49;
- (vi) a HTLV-3 pro protein encoded by a nucleic acid sequence comprising SEQ ID NO: 47;
- (vii) a HTLV-4 gag protein encoded by a nucleic acid sequence comprising nucleotides 750-2024 of SEQ ID NO: 81;
- (viii) a HTLV-4 pol protein encoded by a nucleic acid sequence at least 71.5% identical to SEQ ID NO: 2;
- (ix) a HTLV-4 env protein encoded by a nucleic acid sequence at least 73.5% identical to SEQ ID NO: 4;
- (x) a HTLV-4 tax protein encoded by a nucleic acid sequence at least 82% identical to SEQ ID NO: 6;

(xi) a HTLV-4 rex protein encoded by a nucleic acid sequence comprising SEQ ID NO: 61; and

(xii) a HTLV-4 pro protein encoded by a nucleic acid sequence comprising SEQ ID NO: 59.

2. The isolated PTLV polypeptide of claim 1, wherein the protein is encoded by a nucleic acid sequence comprising SEQ ID NO: 35; SEQ ID NO: 1; SEQ ID NO: 3; SEQ ID NO: 5; SEQ ID NO: 49; SEQ ID NO: 47; nucleotides 750-2024 of SEQ ID NO: 81; SEQ ID NO: 2; SEQ ID NO: 4; SEQ ID NO: 6; SEQ ID NO: 61 or SEQ ID NO: 59.

3. The isolated PTLV polypeptide of claim 1, wherein the protein comprises:

- (i) an amino acid sequence having at least 95% homology to the amino acid sequence set forth as SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 37, SEQ ID NO: 50, SEQ ID NO: 48, SEQ ID NO: 46, SEQ ID NO: 57, SEQ ID NO: 54, SEQ ID NO: 62, SEQ ID NO: 60, or SEQ ID NO: 58;
- (ii) a conservative substitution of 1 or 2 amino acids of the amino acid sequence set forth as one of SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 37, SEQ ID NO: 50, SEQ ID NO: 48, SEQ ID NO: 46, SEQ ID NO: 57, SEQ ID NO: 54, SEQ ID NO: 62, SEQ ID NO: 60, or SEQ ID NO: 58; or
- (iii) a deletion of 2 to 6 amino acids of the amino acid sequence set forth as one of SEQ ID NO: 40, SEQ ID

- NO: 44, SEQ ID NO: 37, SEQ ID NO: 50, SEQ ID NO: 48, SEQ ID NO: 46, SEQ ID NO: 57, SEQ ID NO: 54, SEQ ID NO: 62, SEQ ID NO: 60, or SEQ ID NO: 58.
4. The isolated PTLV polypeptide of claim 1, wherein the amino acid sequence of the protein comprises SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 37, SEQ ID NO: 50, SEQ ID NO: 48, SEQ ID NO: 46, SEQ ID NO: 57, SEQ ID NO: 54, SEQ ID NO: 62, SEQ ID NO: 60, or SEQ ID NO: 58.
5. The isolated PTLV polypeptide of claim 1, wherein the amino acid sequence of the protein consists of SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 37, SEQ ID NO: 50, SEQ ID NO: 48, SEQ ID NO: 46, SEQ ID NO: 57, SEQ ID NO: 54, SEQ ID NO: 62, SEQ ID NO: 60, or SEQ ID NO: 58.
6. A method for detecting antibodies to a PTLV in a biological sample, comprising contacting the biological sample with the polypeptide of claim 1, wherein binding of antibodies from the biological sample to the polypeptide indicates the presence of antibodies to the PTLV in the biological sample.
7. The method of claim 6, wherein the PTLV is HTLV-3 or HTLV-4.
8. The method of claim 6, wherein the biological sample comprises a body fluid sample.
9. An isolated antibody that specifically binds the polypeptide of claim 1.
10. A method of detecting the presence of a PTLV in a subject, comprising:
- (i) contacting a sample from the subject with the isolated antibody of claim 10; and
  - (ii) detecting binding of the antibody to the sample, wherein binding of the antibody to the sample indicates the presence of PTLV in the subject.
11. The method of claim 10, further comprising obtaining a sample from the subject prior to step (i).
12. The method of claim 10, wherein the sample is a tissue sample.
13. The method of claim 10, wherein the PTLV is HTLV-3 or HTLV-4.
14. A method of detecting the presence of a PTLV in a subject comprising:
- (i) contacting a sample from the subject with an antibody directed to the polypeptide of claim 1; and
  - (ii) detecting binding of the antibody to the sample, wherein binding of the antibody to the sample indicates the presence of PTLV in the subject.
15. The method of claim 14, further comprising obtaining a sample from the subject prior to step (i).
16. The method of claim 14, wherein the sample is a tissue sample.
17. The method of claim 14, wherein the PTLV is HTLV-3 or HTLV-4.
18. A kit for detecting the presence of antibodies that bind a PTLV in a sample, wherein the kit comprises at least one polypeptide of claim 1.
19. The kit of claim 18, wherein the PTLV is HTLV-3 or HTLV-4.

\* \* \* \* \*

专利名称(译)	灵长类动物t-嗜淋巴细胞病毒		
公开(公告)号	<a href="#">US20100317034A1</a>	公开(公告)日	2010-12-16
申请号	US12/829125	申请日	2010-07-01
[标]申请(专利权)人(译)	美国证券交易委员会的政府 卫生与人类服务美分DEPT美国疾病控制与预防 与约翰霍普金斯UNIV		
申请(专利权)人(译)	美利坚合众国政府是代表的司 卫生和人类服务部, 疾病控制和预防中心, 与约翰霍普金斯大学		
当前申请(专利权)人(译)	美利坚合众国政府是代表的司 卫生和人类服务部, 疾病控制和预防中心, 与约翰霍普金斯大学		
[标]发明人	SWITZER WILLIAM M HENEINE WALID FOLKS THOMAS M WOLFE NATHAN D BURKE DONALD S NGOLE EITEL MPOUDI		
发明人	SWITZER, WILLIAM M. HENEINE, WALID FOLKS, THOMAS M. WOLFE, NATHAN D. BURKE, DONALD S. NGOLE, EITEL MPOUDI		
IPC分类号	G01N33/53 C07K14/005 C07K16/08		
CPC分类号	A61K2039/5256 C07K14/005 C12N15/86 C12Q1/70 C12N2740/14022 C12N2740/14043 C12Q1/701 C12N2740/14021 A61P37/00		
优先权	60/654484 2005-02-21 US		
其他公开文献	US8541221		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

#### 摘要(译)

公开了与灵长类T-嗜淋巴细胞病毒HTLV-3和HTLV-4的分离和鉴定有关的组合物和方法。HTLV的多样性在非洲中部地区进行了调查, 报告通过狩猎, 屠宰和饲养灵长类动物来接触NHP血液和体液。在此, 显示该群体感染了多种HTLV, 包括两种逆转录病毒; HTLV-4是新型系统发育谱系的第一个成员, 与所有已知的HTLV和STLV不同; HTLV-3属于STLV-3的遗传多样性, STLV-3是一种以前在人类中未见过的组。本公开还涉及用于人类抗感染和疾病的载体和疫苗。本公开还涉及用于检测和诊断由HTLV-3和HTLV-4及相关病毒引起的疾病和疾病的各种生物测定和试剂盒。

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100