

US008617831B2

(12) **United States Patent**
Bottaro et al.

(10) **Patent No.:** **US 8,617,831 B2**
(45) **Date of Patent:** **Dec. 31, 2013**

(54) **METHODS FOR DIAGNOSING AND MONITORING THE PROGRESSION OF CANCER BY MEASURING SOLUBLE C-MET ECTODOMAIN**

(71) Applicants: **Donald P. Bottaro**, Kensington, MD (US); **Gagani P. Athauda**, Miramar, FL (US); **Teresa Lynn Burgess**, Ventura, CA (US)

(72) Inventors: **Donald P. Bottaro**, Kensington, MD (US); **Gagani P. Athauda**, Miramar, FL (US); **Teresa Lynn Burgess**, Ventura, CA (US)

(73) Assignees: **The United States of America, as Represented by the Secretary, Department of Health and Human Services**, Washington, DC (US); **Amgen, Inc.**, Thousand Oaks, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/627,595**

(22) Filed: **Sep. 26, 2012**

(65) **Prior Publication Data**
US 2013/0029352 A1 Jan. 31, 2013

Related U.S. Application Data

(60) Continuation of application No. 13/053,560, filed on Mar. 22, 2011, now Pat. No. 8,304,199, which is a division of application No. 12/093,012, filed as application No. PCT/US2006/043654 on Nov. 8, 2006, now Pat. No. 7,964,365.

(60) Provisional application No. 60/734,993, filed on Nov. 8, 2005, provisional application No. 60/780,626, filed on Mar. 9, 2006.

(51) **Int. Cl.**
G01N 33/53 (2006.01)

(52) **U.S. Cl.**
USPC **435/7.2; 435/4**

(58) **Field of Classification Search**
USPC **435/7.2, 4**
See application file for complete search history.

(56) **References Cited**

FOREIGN PATENT DOCUMENTS

WO	WO 2005/026361	3/2005
WO	WO 2006/056766	6/2006
WO	WO 2007/016367	2/2007

OTHER PUBLICATIONS

Aebi et al., "cDNA structures and regulation of two interferon-induced human Mx proteins," *Mol. Cell Biol.*, Nov. 1989, 9(11), 5062-5072.

Athauda et al., "c-Met ectodomain shedding rate correlates with malignant potential," *Clinical Cancer Research: An Official Journal of the American Association for Cancer Research*, Jul. 2006, 12(14 Pt. 1), 4154-4162.

Bellinzona et al., "Biopolymer-mediated suramin chemotherapy in the treatment of experimental brain tumours," *Acta Oncologica*, 2004, 43(3), 259-263.

Birchmeier et al., "Developmental roles of HGF/SF and its receptor, the c-Met tyrosine kinase," *Trends in Cell Biology*, Oct. 1998, 8(10), 404-410.

Birchmeier et al., "Met, metastasis motility and more," *Nat. Rev. Mol. Cell Biol.*, Dec. 2003, 4(12), 915-925.

Birchmeier et al., "Role of HGF/SF and c-Met in morphogenesis and metastasis of epithelial cells," *Ciba Found. Symp.*, 1997, 212, 230-240.

Blobel, C.P., "ADAMs: Key Components in EGFR Signalling and Development," *Nat. Rev. Mol. Cell Biol.*, Jan. 2005, 6(1), 32-43.

Burgess et al., "Fully Human Monoclonal Antibodies to Hepatocyte Growth Factor with Therapeutic Potential against Hepatocyte Growth Factor/c-Met-Dependent Human Tumors," *Cancer Res.*, Feb. 2006, 66(3), 1721-1729.

Castagnino et al., "Induction of tissue inhibitor of metalloproteinase-3 is a delayed early cellular response to hepatocyte growth factor," *Oncogene*, Jul. 1998, 17(4), 481-492.

Cheng et al., "Overexpression of c-met as a prognostic indicator for transitional cell carcinoma of the urinary bladder: a comparison with p53 nuclear accumulation," *J. Clin. Oncol.*, Mar. 2002, 20(6), 1544-1550.

Christensen et al., "A Selective Small molecule inhibitor of c-Met kinases inhibits c-Met-dependent phenotypes in vitro and exhibits cytoreductive antitumor activity in vivo," *Cancer Research*, Nov. 2003, 63(21), 7345-7355.

Christensen et al., "c-Met as a target for human cancer and characterization of inhibitors for therapeutic intervention," *Cancer Letters*, Jul. 2005, 225(1), 1-26.

Chu et al., "Genetic abnormalities specifically associated with varying metastatic potential of prostate cancer cell lines as detected by comparative genomic hybridization," *Cancer Genet Cytogenet*, Jun. 2001, 127(2), 161-167.

Comoglio et al., "Scatter Factors and invasive growth," *Seminars in Cancer Biology*, Apr. 2001, 11(2), 153-165.

Crepaldi et al., "Generation of a truncated hepatocyte growth factor receptor in the endoplasmic reticulum," *J. Biol. Chem.*, Jan. 1994, 269(3), 1750-1755.

Dharmawardana et al., "Soluble c-Met ectodomain detection in urologic malignancies," *Proceedings of the American Association for Cancer Research Annual Meeting*, Apr. 2005, 46, p. 656.

Dunsmore et al., "Mechanisms of hepatocyte growth factor stimulation of keratinocyte metalloproteinase production," *J. Biol. Chem.*, Oct. 1996, 271(40), 24576-24582.

Funakoshi et al., "Hepatocyte growth factor: from diagnosis to clinical applications," *Clin. Chim. Acta*, Jan. 2003, 327(1-2), 1-23.

Furge et al., "Met receptor tyrosine kinase: enhanced signaling through adapter proteins," *Oncogene*, Nov. 2000, 19(49), 5582-5589.

Galvani et al., "Suramin modulates cellular levels of hepatocyte growth factor receptor by inducing shedding of a soluble form," *Biochem. Pharmacol.*, Sep. 1995, 50(7), 959-966.

Huh et al., "Hepatocyte growth factor/c-met signaling pathway is required for efficient liver regeneration and repair," *PNAS*, Mar. 2004, 101(13), 4477-4482.

(Continued)

Primary Examiner — Bin Shen

(74) *Attorney, Agent, or Firm* — Woodcock Washburn LLP

(57) **ABSTRACT**

Methods for measuring c-Met levels in urine and blood samples are provided. Methods for diagnosis and prognosis evaluation for cancer are also provided.

16 Claims, 13 Drawing Sheets

(56)

References Cited

OTHER PUBLICATIONS

- Jeffers et al., "Degradation of the Met tyrosine kinase receptor by the ubiquitin-proteasome pathway," *Mol. Cell Biol.*, Feb. 1997, 17(2), 799-808.
- Jeffers et al., "Enhanced tumorigenicity and invasion-metastasis by hepatocyte growth factor/scatter factor-met signalling in human cells concomitant with induction of the urokinase proteolysis network," *Mol. Cell Biol.*, Mar. 1996, 16(3), 1115-1125.
- Jin et al., "Early treatment with hepatocyte growth factor improves cardiac function in experimental heart failure induced by myocardial infarction," *J. Pharmacol. Exp. Ther.*, Feb. 2003, 304(2), 654-660.
- Joseph et al., "Expression of Scatter factor in human bladder carcinoma," *J. of the National Cancer Institute*, Mar. 1995, 87(5), 372-377.
- Komada et al. "Proteolytic processing of the hepatocyte growth factor/scatter factor receptor by furin," *FEBS Letters*, Aug. 1993, 328(1-2), 25-29.
- Koochekpour et al., Met and hepatocyte growth factor/scatter factor expression in human gliomas, *Cancer Res.*, Dec. 1997, 57(23), 5391-5398.
- Leone et al., "Transfection of human nm23-H1 into the human MDA-MB-435 breast carcinoma cell line: effects on tumor metastatic potential, colonization and enzymatic activity," *Oncogene*, Sep. 1993, 8(9), 2325-2333.
- Liu, Y., "Hepatocyte growth factor in kidney fibrosis: therapeutic potential and mechanisms of action," *Am. J. Phys. Renal Physiol.*, Jul. 2004, 287(1), F7-16.
- Masson et al., "Tumor necrosis factor receptor-associated periodic syndrome (TRAPS): definition, semiology, prognosis, pathogenesis, treatment, and place relative to other periodic joint diseases," *Joint Bone Sprine*, Jul. 2004, 71(4), 284-290.
- Matsumoto et al., "Hepatocyte growth factor: renotropic role and potential therapeutics for renal diseases," *Kidney International*, Jun. 2001, 59(6), 2023-2028.
- Michalopoulos et al., "Liver regeneration," *Science*, Apr. 1997, 276(5309), 60-66.
- Nakamura et al., "Myocardial protection from ischemia/reperfusion injury by endogenous and exogenous HGF," *J. Clin. Invest.*, Dec. 2000, 106(12), 1511-1519.
- Nath et al., "Shedding of c-Met is regulated by crosstalk between a G-protein couple receptor and the EGF receptor and is mediated by a TIMP-3 sensitive metalloproteinase," *J. Cell. Sci.*, Mar. 2001, 114(Pt. 6), 1213-1220.
- Parr et al., "Expression of hepatocyte growth factor/scatter factor, its activator, inhibitors and the c-Met receptor in human cancer cells," *International Journal of Oncology*, Editorial Academy of the International Journal of Oncology, Oct. 2001, 19(4), 847-863.
- Pennacchietti et al., "Hypoxia promotes invasive growth by transcriptional activation of the met protooncogene," *Cancer Cell*, Apr. 2003, 3(4), 347-361.
- Pisters et al., "c-Met proto-oncogene expression in benign and malignant human prostate tissues," *J. of Urology*, Jul. 1995, 154(1), 293-298.
- Prat et al., "C-terminal truncated forms of Met, the hepatocyte growth factor receptor," *Mol. Cell Biol.*, Dec. 1991, 11(12), 5954-5962.
- Rosario et al., "How to make tubes: signalling by the met receptor tyrosine kinase," *Trends Cell Biol.*, Jun. 2003, 13(6), 328-335.
- Rosen et al., "Urinary and tissue levels of scatter factor in transitional cell carcinoma of bladder," *J. Urol.*, Jan. 1997, 157(1), 72-78.
- Salerno et al., "Inhibition of signal transduction by the nm23 metastasis suppressor: Possible mechanisms," *Clin. Exp. Metastasis*, 2003, 20(1), 3-10.
- Santer et al., "Malignant MCF10CA1 cell lines derived from premalignant human breast epithelial MCF10AT cells," *Breast Cancer Res. Treat.*, Jan. 2001, 65(2), 101-110.
- Schmidt et al., "Scatter factor/hepatocyte growth factor is essential for liver development," *Nature*, Feb. 1995, 373(6516), 699-702.
- Stahl et al., "Functional and biophysical characterization of recombinant human hepatocyte growth factor isoforms produced in *Escherichia coli*," *Biochem J.*, Sep. 1997, 326(Pt. 3), 763-772.
- Tang et al., "TGF-beta switches from tumor suppressor to prometastatic factor in a model of breast cancer progression," *J. Clinical Invest.*, Oct. 2003, 112(7), 1116-1124.
- Uehara et al., "Placental defect and embryonic lethality in mice lacking hepatocyte growth factor/scatter factor," *Nature*, Feb. 1995, 373, 702-705.
- Wajih et al., "Vascular origin of a soluble truncated form of the hepatocyte growth factor receptor (c-Met)," *Circulation Research*, Grune and Stratton, Jan. 2002, 90(1), 46-52.
- Wallenius et al., "Overexpression of the hepatocyte growth factor (HGF) receptor (MET) and presence of a truncated and activated intracellular HGF receptor fragment in locally aggressive/malignant human musculoskeletal tumors," *Am. J. Pathol.*, Mar. 2000, 156(3), 821-829.
- Wang et al., "Alteration of APC, c-met and p53 genes in tumor tissue and serum of patients with gastric cancers," *J. of Surgical Research*, Aug. 2004, 120(2), 242-248.
- Yu et al., "Frequency of TPR-MET rearrangement in patients with gastric carcinoma and in first-degree relatives," *Cancer*, Apr. 2000, 88(8), 1801-1806.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

Creatinine normalized c-Met values of human patient urine samples




Figure 7

Creatinine normalized c-Met mean values of human patient urine samples




Figure 8

Creatinine normalized c-Met values over "No Malignancy" group





Figure 9

c-Met Extracellular Domain



PRIOR ART

Figure 10

Full length c-Met AA sequence

1 MKAPAVLAPG IIVLLETFIVQ RSNGECKEAL AKSEMNVNMK YQLPNFIAET PIONVILHEH HIFLGATNYI
71 YVLNEEDLQK VAEYKTPGVL EHPDQFPQD CSSKANLGG VKKDNINMAL VVDYYDDQL ISCGSVNRGT
141 QRHVFPNH TADIQSEVHC IFSPQIEEPS QCPCVVSAL GAKVLSSVKD RFINFFVGT INSSYFPDHP
211 LHSISVRLK ETKDGFMLT DQSYLDVLPF FRDSYPIKYV HAFESNNFY FLTVQRETID AQTHTRIIR
281 FCSINSGLHS YMEMPLCIL TEKRKRSTK KEVFNILQAA YVSKPGAQLA ROIGASLNDD ILFGVFAQSK
351 PDSAEEMDRS AMCAFFIKYV NDFFNKIVNK NNVRCLOHFY GENHEHCFNR TLLRNSGCE ARRDEYRTEF
421 TTALQRVDLF MGQFSEVILT SISTFIKGD L TIANLGTSEG REMQVVVRS GFSTPHVNF LLD SHPVSPDEV
491 IVEHILNQNG YTLVITGKKI TKIPLNGLGC RHEQSCSQCL SAPPFVQCGW CHDKCVRSEE CLSGTWTQQI
561 CLPAIXKVFP NSAPLEGGTR LTICGWDFG RRNKFDLKK TRVLLGNESC TLLSESTMN TLKCTVGPAM
631 NKHFMSIII SNGHGTQYS TFSVDPVLI SISPKYGPMA GGTLLTLTGN YLNSGNSRHI SIGGKTCTLK
701 SVSNSILECY TPAQTISTEF AVKLKIDLAN RETSIFSYRE DFIVYEIHT KSFISTWKE PLNIVSFLFC
771 FASGGSTIG VGNLNSVSV PRMVINVHEA GRNFTVACQH RSNSEIICCT TPSLQQLNLQ LPLKTKAFFM
841 LDGILSKYFD LIYVHNPVK PEEKPVMISM GNEVLEIKG NDIDPEAVKG EVLKVGNKSC ENIHLHSEAV
911 LCTVPNDLLK INSEINIEWK QALSSTVLGK VIVQPDQNF TGLIAGVVSIS TALLLLLGFF IWLKRRKQIK
981 DLGSELVRYD ARVTFPHDR LVSARSVSPT TEMVSNESVD YRATFPEDQF PNSSQNGSCR QVQYPLTDM
1051 PILTSGSDSI SSPLLQNTVH IDLSALNPEL VQAVQHVIG PSSLIVHFNE VIGRGHFGCV YHGTLIDNDG
1121 KKIHCAYKSL NRITDIGEVS QFLTEGIIMK DFHPNVLSL LGICLRSEGS PIVVLPYMKH GDIRNFIRNE
1191 TENPTVKDLI CFGLQVAKAM KYLASKKVVH RDLAARNCL DEKFTVKVAD FGLARDMYDK EYSVHNKTFG
1261 AKLPVKWML ESLQTQKFTT KSDVWSFGV LWEIWMTRGAP PYPDVNTFFI TVYLLQGRRL LQPEYCPDPL
1331 YEVMKCKWHP KAEMRPSFSE LVSRSIAIFS TFIGEHIYHV NATYVNVKCV APYPSLLSSE DNADDEVDFR
1401 PASFWEYS

PRIOR ART

Figure 11




Figure 12





Figure 13

**METHODS FOR DIAGNOSING AND
MONITORING THE PROGRESSION OF
CANCER BY MEASURING SOLUBLE C-MET
ECTODOMAIN**

CROSS REFERENCE TO RELATED
APPLICATIONS

This application is a continuation of U.S. application Ser. No. 13/053,560, filed Mar. 22, 2011, now U.S. Pat. No. 8,304,199, issued Nov. 6, 2012, which is a division of U.S. application Ser. No. 12/093,012, filed May 8, 2008, now U.S. Pat. No. 7,964,365, issued Jun. 21, 2011, which is the U.S. national stage of International Application Number PCT/US2006/043654, filed Nov. 8, 2006, which claims the benefit of U.S. Provisional Application Ser. No. 60/780,626, filed Mar. 9, 2006, and U.S. Provisional Application Ser. No. 60/734,993, filed Nov. 8, 2005, each of which is incorporated herein by reference in its entirety.

TECHNICAL FIELD

The present invention relates generally to the field of cancer and more specifically to cancer diagnosis and prognosis.

BACKGROUND

Many transmembrane proteins are proteolytically released from the cell surface by a process known as ectodomain shedding. Shedding occurs under normal physiologic conditions and can be increased in certain pathologies. Among the many receptors for which ectodomain shedding has been demonstrated is c-Met, the hepatocyte growth factor ("HGF") receptor tyrosine kinase (Komada et al., *FEBS Lett* 1993; 328:25-9, Wajih et al., *Circ Res* 2002; 90:46-52). HGF is a pleiotropic heparin-binding protein identified and isolated based on observations of its mitogenic activity on hepatocytes and epithelial cells, and independently identified and isolated based on observations of its ability to stimulate cell motility (scatter factor or SF). (Comoglio et al., *Semin Cancer Biol* 2001; 11:153-65, Funakoshi et al., *Clin Chim Acta* 2003; 327:1-23). HGF is typically produced by cells of mesenchymal origin and acts in a paracrine manner on a variety of cellular targets including epithelial and endothelial cells, hematopoietic cells, neurons and melanocytes during embryonic development and throughout adulthood, in normal and pathological processes (Birchmeier et al., *Nat Rev Mol Cell Biol* 2003; 4:915-25). HGF is essential for embryonic development, where it is involved in somite migration, limb bud and limb skeletal muscle formation, placenta formation (Schmidt et al., *Nature* 1995; 373:699-702, Uehara et al., *Nature* 1995:373:702-5) and later in organogenesis (Rosario et al., *Trends Cell Biol* 2003; 13:328-35), in neural development (Birchmeier et al., *Trends Cell Biol* 1998:8:404-10) and in tissue repair and regeneration (Jin et al., *J Pharmacol Exp Ther* 2003; 304:654-60, Huh et al., *Proc Natl Acad Sci USA* 2004; 101:4477-82). While the role of HGF in adult homeostasis is not yet completely understood, a growing body of evidence suggests that it is an endogenous tissue protective factor for several major organs and has potent antifibrotic activity (Liu, *Am J Physiol Renal Physiol* 2004; 287:F7-16).

The MET oncogene was isolated from a human osteogenic sarcoma cell line that had been chemically mutagenized in vitro. Transforming activity was due to a DNA rearrangement where sequences from the TPR (translocated promoter region) locus on chromosome 1 fused to sequences from the MET locus on chromosome 7 (TPR-MET) (Furge et al.,

Oncogene 2000; 19:5582-9). This rearrangement has been found in patients with gastric carcinoma (Yu et al., *Cancer* 2000; 88:1801-6). Isolation of the full-length MET proto-oncogene coding sequence revealed structural features of a membrane spanning receptor tyrosine kinase (Furge et al., *Oncogene* 2000; 19:5582-9). The identification of HGF as the natural ligand for c-Met and the identity of SF and HGF united a collection of findings demonstrating that a single receptor transduced multiple biological activities including motility, proliferation, survival and branching morphogenesis (Birchmeier et al., *Nat Rev Mol Cell Biol* 2003; 4:915-25). Activation of the c-Met intrinsic tyrosine kinase (TK) activity was required for all of these activities. Consistent with its relationship with HGF, c-Met is widely expressed early in development, deletion of the gene is embryonic lethal in mice, and widespread expression persists throughout adulthood (Birchmeier et al., id). Both HGF and c-Met are upregulated after kidney, liver or heart injury, suggestive of a general mechanism of protection against tissue damage, as well as one of tissue repair and regeneration (Matsumoto et al., *Kidney Int* 2001; 59:2023-38, Michalopoulos et al., *Science* 1997; 276:60-6, Nakamura et al., *J Clin Invest* 2000; 106:1511-9).

HGF and c-Met are implicated in a wide variety of human malignancies including colon, gastric, bladder, breast, kidney, liver, lung, head and neck, thyroid and prostate, but also sarcomas, hematological malignancies, melanoma and central nervous system (CNS) tumors (Birchmeier et al., *Nat Rev Mol Cell Biol* 2003; 4:915-25, Birchmeier et al., *Ciba Found Symp* 1997; 212:230-40). Through paracrine signaling, overexpression of ligand and/or receptor, autocrine loop formation and/or receptor mutation and gene rearrangement, this signaling pathway can enhance tumor cell growth, proliferation, survival, motility and invasion. Inappropriate c-Met signaling in disease can resemble, at least in part, developmental transitions between epithelial and mesenchymal cell types normally regulated by HGF. Among the many genes upregulated in response to activation of this pathway is that of the receptor itself, creating the potential for c-Met overexpression in otherwise normal target cells through persistent ligand stimulation; consistent with this, c-Met overexpression is widely observed in cancers of epithelial origin where paracrine delivery of HGF results in dysregulated signaling, whereas cells of mesenchymal origin that normally express HGF often acquire c-Met expression, and several sarcomas display autocrine c-Met signaling (Furge et al., *Oncogene* 2000; 19:5582-9). Importantly, the c-Met pathway activates a program of cell dissociation and increased cell motility coupled with increased protease production that has been shown to promote cellular invasion through extracellular matrices, and that closely resembles tumor metastasis in vivo (Birchmeier et al., *Ciba Found Symp* 71p 1997; 212:230-40). In addition, pathway activation in vascular cells stimulates tumor angiogenesis, facilitating tumor growth for cancers that are growth limited by hypoxia, and promoting tumor metastasis. Hypoxia alone upregulates c-Met expression and enhances HGF/SF signaling in cultured cells and mouse tumor models (Pennacchietti et al., *Cancer Cell* 2003; 3:347-61).

Early diagnosis is a key strategy in cancer treatment. Although it is known that c-Met is overexpressed in certain cancers, a need exists for a screening test for c-Met that is sensitive, cost-efficient, and can be used for diagnosis, determining stage of disease, prognosis, and/or assessing the efficacy of therapeutic intervention. The invention is directed to this and other ends.

The present invention generally relates to methods for diagnosing cancer in a patient or methods for identifying a patient having an increased risk of developing cancer. The methods provide for analysis of patient samples to identify the presence of tumor cells in the patient due to the observation that tumor cells expressing c-Met shed more c-Met ectodomain than their normal tissue counterparts, independent of changes in overall c-Met expression levels, and that this shedding is enhanced with increasingly malignant phenotype. Moreover, the methods for diagnosis provide that c-Met ectodomain shedding can be quantitated in cell lysates, culture supernatants, and biological samples, such as blood and urine, and that c-Met levels are measurable before tumors become palpable yet correlate directly with tumor volume. Thus, it was concluded that for a variety of cancers, c-Met ectodomain shedding provides a reliable and practical indicator of, among other things, malignant potential and overall tumor burden.

A method for identifying a patient having an increased risk of developing cancer which comprises the steps of measuring an amount of soluble c-Met ectodomain in a biological sample of the patient; and detecting an increased concentration of soluble c-Met ectodomain in the biological sample when compared to the concentration of soluble c-Met ectodomain in a corresponding biological sample from a control population, wherein an increased concentration of soluble c-Met ectodomain identifies the patient as having an increased risk of developing cancer. The biological sample includes, but is not limited to, blood plasma or urine. The concentration of soluble c-Met ectodomain in a urine sample and a blood plasma sample is measured and the combined measurement can be used to determine if the patient has the increased risk of developing cancer. An increased concentration of soluble c-Met ectodomain can identify the patient as having an increased risk of developing metastatic cancer.

In one aspect, a soluble c-Met ectodomain level of about 3500 pg c-Met per mg urine creatinine or greater is indicative of an increased risk of developing cancer in the patient. In a further aspect, a soluble c-Met ectodomain level of about 4900 pg c-Met per mg urine creatinine or greater is indicative of an increased risk of developing cancer in the patient. In a further aspect, a soluble c-Met ectodomain level of about 3800 pg c-Met per mg urine creatinine or greater is indicative of an increased risk of developing cancer in the patient. In a detailed aspect, a soluble c-Met ectodomain level of about 4200 pg c-Met per mg urine creatinine or greater is indicative of an increased risk of developing cancer in the patient. In a detailed aspect, a soluble c-Met ectodomain level of about 4500 pg c-Met per mg urine creatinine or greater is indicative of an increased risk of developing cancer in the patient. The cancer diagnosis can be a cancer of the urinary tract. The cancer includes, but is not limited to, kidney cancer, renal cancer, prostate cancer, bladder cancer, renal cell carcinoma, or breast cancer.

A method for monitoring the progression of cancer in a patient is provided which comprises the steps of measuring an amount of soluble c-Met ectodomain in a first biological sample at a first time point, measuring an amount of soluble c-Met ectodomain in a second biological sample at a second time point, and determining progression of cancer in the patient based upon a change in amount of soluble c-Met ectodomain in the first biological sample compared to the second biological sample. The amount of soluble c-Met ectodomain in a urine sample or blood plasma sample, or a

combination thereof, is measured and the combined measurement is used to determine the progression of cancer.

A method for measuring the response to cancer therapy in a patient is provided which comprises the steps of measuring an amount of soluble c-Met ectodomain in a biological sample of the patient at a first time point, measuring an amount of soluble c-Met ectodomain in a second biological sample of the patient at a second time point, and determining response to cancer therapy in the patient based upon the change in amount of soluble c-Met ectodomain in the first biological sample compared to the second biological sample. In one aspect, a decrease in the amount of soluble c-Met ectodomain in the first biological sample compared to the second biological sample indicates a positive response to cancer therapy in the patient. The cancer therapy can be an anti-cancer drug, for example, a chemotherapeutic agent, ionizing radiation therapy or hormone ablation therapy.

A method of assaying for an amount of c-Met ectodomain present in a biological sample from a human patient, said method is provided which comprises assaying the biological sample to determine an amount of c-Met ectodomain present in the biological sample from the patient, and comparing the amount of c-Met ectodomain present in the biological sample from the patient to an amount of c-Met ectodomain found in a biological sample of a control population, wherein an increase in the amount of c-Met ectodomain found in the sample of the patient compared to the amount of c-Met ectodomain found in the sample of the control population is indicative of the patient having cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. c-Met shedding in a cultured cell model of breast tumor progression. 1A. Reducing SDS-PAGE and immunoblot analysis of the c-Met expression in four mammary cell lines; p145 is the intact c-Met beta subunit. In this model, M2, M3 and M4 represent derivatives of the normal breast cell line M1 with increasingly malignant phenotype. The normal mammary epithelial cell line B5/589 was used as a positive control for c-Met expression, and lysates were immunoblotted for beta actin to confirm equal sample loading. 1B. Reducing SDS-PAGE and immunoblot analysis of c-Met ectodomain shedding by cell lines M1, M2, M3 and M4. Culture supernatant from B5/589, as well as a purified recombinant c-Met ectodomain IgG Fc fusion protein (358-MT, R&D Systems) were used as positive controls of c-Met shedding and ectodomain recognition, respectively. Molecular masses of predominant c-Met ectodomain fragments are indicated in kDa at left.

FIG. 2. Quantitation of c-Met shedding using an electrochemiluminescent two site immunoassay. 2A. Standard curve of purified recombinant c-Met ectodomain-IgG fusion protein (358-MT). Mean values from quadruplicate samples are shown as log [soluble c-Met] in ng/ml vs. log [signal intensity] in relative units. Standard deviations are smaller than the symbol size. 2B. Analysis of sample quenching in A431 human epidermoid carcinoma cells. Cells were serum deprived for 24 h and soluble c-Met ectodomain was measured in the presence and absence of added 358-MT. Mean values of quadruplicate samples from 358-MT (358MT, open bar), A431 conditioned medium (A431, dark gray bar), and A431 conditioned medium+358-MT (obs, light gray bar) are shown +/- standard deviation. The expected sum of 358-MT and A431 conditioned medium sample signals is also shown (exp, dark-gray/open bar). 2C. Soluble c-Met in 24 h B5/589 conditioned media (B5, open bar) was filtered using a 10 kDa cutoff molecular sieve and soluble c-Met in the filtrate was

measured (YM10, dark gray bar). 24 h B5/589 conditioned medium (open bar) was immunodepleted using an ectodomain-specific monoclonal antibody (mAb DO24, Upstate Biotechnology; light gray bar). 2D. HGF binding (ng/ml) was measured in samples containing equivalent amounts of a c-Met Fc fusion protein (open squares) and c-Met ectodomain present in B5/589 conditioned medium (black circles), as a function of the concentration of HGF added (ng/ml). Error bars represent standard deviation; where no bars are seen, the deviation is smaller than the symbol size.

FIG. 3. Quantitation of c-Met expression and shedding in a cultured cell model of breast tumor progression. Cell surface and soluble c-Met levels were measured in the MCF10A derived cell lines M1 (open bars), M2 (light gray bars), M3 (dark gray bars) and M4 (black bars). Non-ionic detergent extractable c-Met is expressed as pg/ug cell protein (Cellular c-Met, top panel); c-Met ectodomain present in 24 h cell conditioned medium is shown expressed as pg/ug cell protein (Soluble c-Met, 2nd panel) and after normalization to detergent extractable c-Met (Soluble/Cellular, 3rd panel). The percentage of detergent extractable c-Met that became soluble in 24 h is shown in the bottom panel.

FIG. 4. Quantitation of c-Met expression and shedding in paired cell line models of cancer progression. Cell line pairs, each derived from a single patient, were compared for levels of cellular (top panel) and soluble (2nd panel) c-Met. These data were used to calculate the amount of soluble receptor shed after 24 h corrected for cellular receptor (3rd panel) and the amount of soluble receptor produced in 24 h as a percentage of cellular receptor (bottom panel). Cell lines derived from normal tissue (open bars) corresponding to the tissue of tumor origin were paired with tumor-derived cell lines (light gray bars) and other tumor derived cell lines (dark gray bars) were paired with cell lines derived from corresponding metastatic lesions (black bars). CRL7636 is a normal skin cell line; CRL7637 is a skin melanoma cell line; HTB125 is a normal mammary gland cell line; HTB126 is a mammary gland ductal carcinoma cell line; CCL228 is a colorectal adenocarcinoma cell line; CCL227 is a lymph node metastasis of colorectal adenocarcinoma cell line; UOK124 is a renal cell carcinoma cell line; and UOK124 LN is a lymph node metastasis of renal cell carcinoma cell line.

FIG. 5. Genetically modified cultured cell models of cancer progression. 5A. The human prostate cancer derived cell line PC3 (open bars), a derived metastatic variant PC3M (black bars) and PC3M transfected with an ADxA gene expression plasmid (+MxA, gray bars) were compared for levels of cellular (top panel) and soluble (2nd panel) c-Met. These data were used to calculate the amount of soluble receptor shed after 24 h corrected for cellular receptor (3rd panel) and the amount of soluble receptor produced in 24 h as a percentage of cellular receptor (bottom panel). 5B. The human breast cancer derived cell line C100 (black bars) and a C100 derivative cell line cloned after transfection with an NM23 gene expression plasmid (H1.177, gray bars) were compared for levels of cellular and soluble c-Met as in Panel A.

FIG. 6. c-Met shedding in mice bearing human tumor cell xenografts. Soluble human c-Met concentrations in plasma (ng/ml, upper panels) or urine samples (pg/ml, lower panels) obtained from mice bearing subcutaneous UOK261 human bladder carcinoma (left panels) or U-87 MG human glioblastoma (right panels) cell xenografts plotted against the corresponding tumor volume (mm³) in each mouse at the time of sample procurement. Values represent the mean of triplicate measurements; standard deviations are smaller than the symbol size. Each line and corresponding R² value represent a

best fit linear regression analysis forced through the origin and were performed using GraphPad Prism software.

FIG. 7. Creatinine normalized c-Met values of human patient urine samples. FIG. 7 provides a column scattergram illustrating the c-Met values in human patient urine sample. The Y axis shows c-Met pg/mg creatinine. The following acronyms are used on the X (Category) axis: HLRCC refers to Hereditary Leiomyomatosis and Renal Cell Carcinoma (inherited kidney cancer syndrome); BPH refers to Benign Prostatic Hypertrophy (possibly prelude to prostate cancer); CaP refers to Prostate Cancer; CaB refers to Bladder Cancer; and PRC refers to Papillary Renal Cancer, sporadic, not inherited. Each data point is shown as a symbol. The horizontal line represents the median. Statistical analysis was not performed for this data set.

FIG. 8. Creatinine normalized c-Met mean values of human patient urine samples. FIG. 8 provides a bar graph illustrating the mean values of c-Met in the patient populations. The Y axis shows c-Met pg/mg creatinine. The following acronyms are used on the X (Category) axis: HLRCC refers to Hereditary Leiomyomatosis and Renal Cell Carcinoma (inherited kidney cancer syndrome); BPH refers to Benign Prostatic Hypertrophy (possibly prelude to prostate cancer); CaP refers to Prostate Cancer; CaB refers to Bladder Cancer; and PRC refers to Papillary Renal Cancer, sporadic, not inherited. Statistical analysis was not performed for this data set.

FIG. 9. Creatinine normalized c-Met values over "No Malignancy" group. FIG. 9 provides a bar graph illustrating normalized c-Met values over "No Malignancy group". The Y shows fold increase in c-Met. The following acronyms are used on the X (Category) axis: HLRCC refers to Hereditary Leiomyomatosis and Renal Cell Carcinoma (inherited kidney cancer syndrome); BPH refers to Benign Prostatic Hypertrophy (possibly prelude to prostate cancer); CaP refers to Prostate Cancer; CaB refers to Bladder Cancer; and PRC refers to Papillary Renal Cancer, sporadic, not inherited. Statistical analysis was not performed for this data set.

FIG. 10: Sequence for c-Met extracellular domain. FIG. 10 provides the sequence for the c-met extracellular domain, SEQ ID NO: 1 showing the functional domains. The c-Met ectodomain comprises residues 25 to 932 of the c-Met extracellular domain.

FIG. 11: c-Met full length sequence. FIG. 11 provides the full length c-Met sequence, SEQ ID NO:2.

FIG. 12: c-Met in normal patients versus patients with bladder cancer. FIG. 12 provides a box and whisker graph illustrating c-Met shedding in a group of normal patients versus patients with bladder cancer. The box extends from the 25th percentile to the 75th percentile, with a horizontal line at the median (50th percentile). Whiskers extend down to the smallest value and up the largest.

FIG. 13: Immunodepletion of plasma samples.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

Overview

The present inventors have observed, after analyzing patient samples, that tumor cells expressing c-Met shed more c-Met ectodomain than their normal tissue counterparts, independent of changes in overall c-Met expression levels, and that this shedding is enhanced with increasingly malignant phenotype. Moreover, the present inventors have observed, after further analysis, that c-Met ectodomain shedding can be quantitated in cell lysates, culture supernatants, and biological samples, such as blood and urine, and that

c-Met levels are measurable before tumors become palpable yet correlate directly with tumor volume. Thus, it was concluded that for a variety of cancers, c-Met ectodomain shedding provides a reliable and practical indicator of, among other things, malignant potential and overall tumor burden. The observation and conclusion that c-Met can be detected in urine and provides a reliable and practical indicator of malignant potential and overall tumor burden was particularly surprising given the large size of the c-Met ectodomain, the low expression levels of c-Met, the acidic pH of urine, and the protease content in urine.

This invention provides, inter alia, methods and assays for diagnosing cancer in a subject, identifying a patient having an increased risk of developing cancer, monitoring the progression of cancer in a subject, measuring a clinical parameter related to cancer in subject, and measuring the therapeutic response to a cancer drug. In a particularly preferred embodiment, the present invention provides methods and assays for diagnosing bladder cancer in a subject, monitoring the progression of bladder cancer in a subject, measuring a clinical parameter related to bladder cancer in subject, and measuring the therapeutic response to a cancer drug for the treatment of bladder cancer.

“Cancer” refers to any of a number of diseases that are characterized by uncontrolled, abnormal proliferation of cells, as well as any of a number of characteristic structural and/or molecular features. A “cancerous cell” is understood as a cell having specific structural properties, lacking differentiation and in many instances, being capable of invasion and metastasis, see DeVita, V. et al. (eds.), 2001, *Cancer Principles and Practice of Oncology*, 6th. Ed., Lippincott Williams & Wilkins, Philadelphia, Pa.). The term cancer includes, for example, cancers of the female reproductive organs including, for example, ovarian cancer, cervical cancer and uterine cancer; lung cancer; breast cancer; renal cell carcinoma; Hodgkin’s lymphoma; Non-Hodgkin’s lymphoma; cancers of the genitourinary system including, for example, kidney cancer, prostate cancer, bladder cancer, and urethral cancer; cancers of the head and neck; liver cancer; cancers of the gastrointestinal system including, for example, stomach cancer, esophageal cancer, small bowel cancer or colon cancer; cancers of the biliary tree; pancreatic cancer; cancers of the male reproductive system including, for example, testicular cancer; Gestational trophoblastic disease; cancers of the endocrine system including, for example, thyroid cancer, parathyroid cancer, adrenal gland cancer, carcinoid tumors, insulinomas and PNET tumors; sarcomas, including, for example, Ewing’s sarcoma, osteosarcoma, liposarcoma, leiomyosarcoma, and rhabdomyosarcoma; mesotheliomas; cancers of the skin; melanomas; cancers of the central nervous system; pediatric cancers; and cancers of the hematopoietic system including, for example, all forms of leukemia, myelodysplastic syndromes, myeloproliferative disorders and multiple myeloma. Cancers referred to in the present methods include those wherein the c-Met signaling pathway is overexpressed, accompanied by increased protease production and/or contributes to malignancy. Exemplary cancers include, for example urological cancers, such as bladder cancer; carcinomas, such as bladder, breast, cervical, cholangiocarcinoma, colorectal, oesophageal, gastric, head and neck, kidney, liver, lung, nasopharyngeal, ovarian, pancreas/gall bladder, prostate and thyroid carcinomas; musculoskeletal carcinomas, such as, osteosarcoma, synovial sarcoma, and rhabdomyosarcoma; soft tissue sarcomas, such as, MFH/fibrosarcoma, leiomyosarcoma, and kaposi’s sarcoma; haematopietic malignancies, such as, multiple myeloma, lymphomas, adult T-cell leukemia, acute myelogenous leu-

kemia, and chronic myeloid leukemia; and other neoplasms, such as glioblastomas, astrocytomas, melanoma, mesothelioma, and Wilms’ tumor (Birchmeier et al., *Nat Rev Mol Cell Bio* 2003 4(12):912-925.

In certain embodiments, the present invention provides methods of diagnosing cancer in a patient or methods for identifying a patient having an increased risk of developing cancer. Diagnosis, as used herein, includes not only the initial identification of a cancer associated with c-met ectodomain shedding in a patient but confirmatory testing, or screening in patients who have previously been identified as having or likely to have a cancer associated with c-met ectodomain shedding. The methods include the steps of providing a biological sample from the patient, measuring the amount of soluble c-Met ectodomain in the biological sample, preferably a urine and/or blood plasma sample, and determining if the patient has a greater likelihood of cancer based on the amount of c-Met measured. A patient has a greater likelihood of having cancer when he has a c-Met level that is greater than the mean c-Met level in a non-cancerous population.

In certain embodiments, a patient having a greater likelihood of cancer will have an amount of c-Met present in a biological sample, preferably a urine and/or plasma sample, that is at least two times the mean amount of c-Met present in a corresponding sample from a non-cancerous control population, at least three times the mean amount of c-Met present in a corresponding sample from a non-cancerous control population, at least four times the mean amount of c-Met present in a corresponding sample from a non-cancerous control population, or at least seven times the mean amount of c-Met present in a corresponding sample from a non-cancerous control population.

The present inventors have analyzed patient samples and after analysis concluded that the mean c-Met levels in a non-cancerous control population (n=30) is 4193 mg c-Met per mg urine creatinine with a standard error of 335. Further statistical analysis has shown that the lower 95% confidence level is 3508 mg c-Met per mg urine creatinine and the higher 95% confidence level is 4878 mg c-Met per mg urine creatinine. Accordingly, in certain embodiments of the present invention, a measurement of c-Met of about 3500 or greater, about 3800 or greater, about 4200 or greater, about 4500 or greater, or about 4900 c-Met per mg urine creatinine or greater, in a urine sample from a patient will be indicative of a greater likelihood of cancer in the patient. Similarly, a measurement of c-Met of less than about 3500 c-Met per mg urine creatinine will indicate that the patient has a smaller risk than the general population of having cancer.

The present inventors have analyzed the patient data and concluded that the mean c-Met levels in a population (n=12) having bladder cancer is 10,600 mg c-Met per mg urine creatinine with a standard error of 2678. Further statistical analysis has shown that the lower 95% confidence level is 4709 mg c-Met per mg urine creatinine and the higher lower 95% confidence level is 16,500 mg c-Met per mg urine creatinine. Accordingly, in certain embodiments of the present invention, a measurement of c-Met of about 4700 or greater, about 6000 or greater, about 7000 or greater or about 8000 or greater, c-Met per mg urine creatinine or greater, in a urine sample from a patient, will be indicative of a greater likelihood of having cancer, in particular bladder cancer

A human sample value, obtained in triplicate or greater replicates, with a mean above the upper limit of the normal range (defined by the upper 95% Confidence Interval, or 4878 pg c-Met/mg Creatinine), and determined to be significantly different (as defined using the Student’s t-test or other appropriate statistical analysis) from the normal mean presented

herein (4193 pg c-Met/mg Creatinine), indicates a greater likelihood of having cancer or other hyperplastic disease, in particular bladder cancer.

A patient having a greater likelihood of having cancer can then be further screened to determine whether the subject does, in fact, have cancer. These screening methods include, for example, conventional work-ups to determine risk factors that are associated with cancer, a history and physical or neurological exam with particular attention to obvious lesions; palpable masses; ulcerations; swelling or enlargement of any masses or organs; erosion of bone; laterality; size and number of palpable lymph nodes; vision changes, focal deficit, tumor impingement on a specific nerve or structure; evidence of increased intracranial pressure; evidence of obstructive hydrocephalus. A diagnosis of cancer can be confirmed, for example, by imaging tests such as X-rays, nuclear scans and/or biopsies.

The methods of the invention rely, in part, on measuring the amount of c-Met in a sample. The c-met protooncogene is transmembrane protein that is derived from a 170 kDa precursor. The mature c-met protein is composed of a 50 kDa α subunit that is linked by 2 disulfide bonds to a 145 kDa β subunit. The c-Met ectodomain comprises residues 25 to 932 of the c-Met extracellular domain, SEQ ID NO: 1, as shown in FIG. 11.

As used herein, the phrase "measuring the amount of c-Met or soluble c-Met ectodomain" means any direct or indirect quantitative assay for c-Met ectodomain, including fragments thereof.

Biological sample refers to a biological tissue or biological fluid from a patient or human patient. The biological tissue can be assayed for c-Met ectodomain as a detergent extract of the tissue. Biological tissue from a patient can be extracted with non-ionic detergent buffer solutions as described herein for preparing cell extracts. The tissue can be minced or ground (at 4° C. or on ice) to maximize surface area available for extraction, and generally the extraction (also on ice) is performed for a much longer period (2-3 hours) as opposed to a few minutes for cultured cells.

A variety of biological samples can be useful in practicing the methods of the invention including, for example, blood, serum, plasma, urine, salivary fluid, ascite fluid and the like. A preferred biological sample is urine.

Such assays provide valuable means of monitoring the status of neoplastic diseases. In addition to improving prognostication, knowledge of the disease status allows the attending physician to select the most appropriate therapy for the individual patient. For example, patients with a high likelihood of relapse can be treated rigorously, usually involving systemic chemotherapy and/or radiation therapy. When there is a lesser likelihood of relapse, less aggressive therapies can be chosen. Because of the severe patient distress caused by the more aggressive therapy regimens, it would be desirable to distinguish with a high degree of certainty those patients requiring such aggressive therapies. See for example, U.S. Pat. No. 5,674,753, incorporated herein by reference in its entirety.

The present methods are useful for screening a wide variety of neoplastic diseases, including both solid tumors and hematopoietic cancers. Exemplary neoplastic diseases include carcinomas, such as adenocarcinomas and melanomas, and squamous cell carcinomas of the head and neck; mesodermal tumors, such as neuroblastomas and retinoblastomas; sarcomas, such as osteosarcomas, Ewing's sarcoma, and various leukemias; and lymphomas. Of particular interest are gynecologic cancers including ovarian, cervical, vaginal, endometrial and vulval cancers; gastrointestinal cancer, such

as, stomach, colon and esophageal cancers; urinary tract cancer, such as, bladder and kidney cancers; skin cancer; liver cancer; prostate cancer; lung cancer; and breast cancer. Of still further particular interest are gynecologic cancers; breast cancer; urinary tract cancers, especially bladder cancer; lung cancer; gastrointestinal cancer, such as, stomach, colon and esophageal cancers; and liver cancer. Even further of particular interest are gynecologic cancers and breast cancer. Tumors which are known to overexpress intact c-Met may be good candidates for target neoplastic diseases for the assays of this invention, that is, such tumors may release into body fluids the c-Met ectodomain protein at elevated levels above normal. Particularly well studied are tumors of the breast and adenocarcinomas of the vulva which have been confirmed to overexpress c-Met. Preferably, in regard to tumors of the lung and cervix, squamous cell carcinomas have been particularly associated with elevated levels of c-Met. c-Met measurements have been determined to be markers for breast cancer and to provide valuable information diagnostically and prognostically, in correlation with other markers, such as, hormone receptor measurements, for example, with estrogen receptors (ERs) and progesterone receptors (PRs).

The body fluids that are of particular interest in assaying for the c-Met ectodomain protein according to the methods of this invention include blood, serum, semen, breast exudate, saliva, sputum, urine, cytosols, plasma, ascites, pleural effusions, amniotic fluid, bladder washes, bronchioalveolar lavages, and cerebrospinal fluid. Blood, serum, plasma, and urine are useful for screening according to the methods of this invention. The assays of this invention may also be useful in detecting and/or quantitating said c-Met ectodomain protein in tissue extracts.

From a knowledge of the structure of the external domain of the c-Met, a number of monoclonal or polyclonal antibodies can be generated that specifically recognize that protein. Because said c-Met ectodomain protein is found to exist freely in the biological fluids of mammals, it is possible to detect and/or quantitate the levels of that protein. Utilizing current antibody detection techniques that can quantitate the binding of monoclonal antibodies, made specifically to epitopes on the external domain of the c-Met, in particular to epitopes on the N-terminal, one can determine the amount of said c-Met ectodomain protein in the fluids of cancer patients.

Such an assay can be used to detect tumors, quantitate their growth, and help in the diagnosis and prognosis of the human disease. The assays can also be used to detect the presence of cancer metastasis, as well as confirm the absence or removal of all tumor tissue following surgery, cancer chemotherapy and/or radiation therapy. It can further be used to monitor cancer chemotherapy and tumor reappearance. Representative immunoassays involve the use of monoclonal or polyclonal antibodies which can be appropriately labeled to detect and/or quantitate said c-Met ectodomain protein in the body fluids of mammals.

Diagnosis is based on analyzing the sample for amount of c-Met and comparing it to a reference value, where the reference value serves to assist in differentiating those with cancer or a certain stage of cancer from other individuals. Statistical analysis routines can be performed to develop predictive models for identification of the different clinical parameters described herein. Measuring an "amount" in a sample means quantifying c-Met by determining, for example, the relative or absolute amount of protein.

A variety of assays for detecting c-Met are known in the art and include direct and indirect assays for c-Met protein. Immunoassays, including radioimmunoassays, enzyme-linked immunoassays and two-antibody sandwich assays as

described further below, are useful in the methods of the invention. Furthermore, monoclonal and polyclonal anti-c-Met antibodies useful in immunoassays can be readily obtained from a variety of sources.

A variety of immunoassay formats, including competitive and non-competitive immunoassay formats, antigen capture assays and two-antibody sandwich assays also are useful the methods of the invention (Self and Cook, *Curr. Opin. Biotechnol.* 7:60-65 (1996)). In one embodiment, a method of the invention relies on one or more antigen capture assays. In an antigen capture assay, antibody is bound to a solid phase, and sample is added such that c-Met antigen is bound by the antibody. After unbound proteins are removed by washing, the amount of bound antigen can be quantitated, if desired, using, for example, a radioassay (Harlow and Lane, *Antibodies A Laboratory Manual* Cold Spring Harbor Laboratory: New York, 1988)). Immunoassays can be performed under conditions of antibody excess, or as antigen competitions, to quantitate the amount of antigen and, thus, determine a level of c-Met.

Enzyme-linked immunosorbent assays (ELISAs) can be useful in certain methods of the invention. An enzyme such as horseradish peroxidase (HRP), alkaline phosphatase (AP), β -galactosidase or urease can be linked, for example, to an anti-c-Met antibody or to a secondary antibody for use in a method of the invention. A horseradish-peroxidase detection system can be used, for example, with the chromogenic substrate tetramethylbenzidine (TMB), which yields a soluble product in the presence of hydrogen peroxide that is detectable at 450 nm. Other convenient enzyme-linked systems include, for example, the alkaline phosphatase detection system, which can be used with the chromogenic substrate p-nitrophenyl phosphate to yield a soluble product readily detectable at 405 nm. Similarly, a β -galactosidase detection system can be used with the chromogenic substrate o-nitrophenyl- β -D-galactopyranoside (ONPG) to yield a soluble product detectable at 410 nm, or a urease detection system can be used with a substrate such as urea-bromocresol purple (Sigma Immunochemicals, St. Louis, Mo.). Useful enzyme-linked primary and secondary antibodies can be obtained from a number of commercial sources such as Jackson Immuno-Research (West Grove, Pa.) as described further below.

In certain preferred embodiments, c-Met ectodomain is detected and measured using chemiluminescent detection. For example in certain embodiments, c-Met ectodomain specific antibodies are used to capture c-Met ectodomain present in the biological sample and a antibody specific for the c-Met ectodomain specific antibodies and labeled with an chemiluminescent label is used to detect the c-Met ectodomain present in the sample. Any chemiluminescent label and detection system can be used in the present methods. Chemiluminescent secondary antibodies can be obtained commercially from various sources such as Amersham. Methods of detecting chemiluminescent secondary antibodies are known in the art and are not discussed herein in detail.

Fluorescent detection also can be useful for detecting c-Met in certain methods of the invention. Useful fluorochromes include, without limitation, DAPI, fluorescein, Hoechst 33258, R-phycoerythrin, B-phycoerythrin, R-phycoerythrin, rhodamine, Texas red and lissamine. Fluorescein or rhodamine labeled α 2-MG-, HA-, TIMP-1- or YKL-40-specific binding agents such as anti- α 2-MG, anti-HA, anti-TIMP-1, or anti-YKL-40 antibodies, or fluorescein- or rhodamine-labeled secondary antibodies can be useful in the invention. Useful fluorescent antibodies can be obtained commercially, for example, from Tago Immunologicals (Burlingame, Calif.) as described further below.

Radioimmunoassays (RIAs) also can be useful in certain methods of the invention. Such assays are well known in the art. Radioimmunoassays can be performed, for example, with ^{125}I -labeled primary or secondary antibody (Harlow and Lane, supra, 1988).

A signal from a detectable reagent can be analyzed, for example, using a spectrophotometer to detect color from a chromogenic substrate; a radiation counter to detect radiation, such as a gamma counter for detection of ^{125}I ; or a fluorometer to detect fluorescence in the presence of light of a certain wavelength. Where an enzyme-linked assay is used, quantitative analysis of the amount of c-Met can be performed using a spectrophotometer such as an EMAX Microplate Reader (Molecular Devices; Menlo Park, Calif.) in accordance with the manufacturer's instructions. The assays of the invention can be automated or performed robotically, if desired, and that the signal from multiple samples can be detected simultaneously.

The methods of the invention also encompass the use of capillary electrophoresis based immunoassays (CEIA), which can be automated, if desired. Immunoassays also can be used in conjunction with laser-induced fluorescence as described, for example, in Schmalzing and Nashabeh, *Electrophoresis* 18:2184-93 (1997), and Bao, *J. Chromatogr. B. Biomed. Sci.* 699:463-80 (1997). Liposome immunoassays, such as flow-injection liposome immunoassays and liposome immunosensors, also can be used to detect c-Met or to determine a level of c-Met according to certain methods of the invention (Rongen et al., *J. Immunol. Methods* 204:105-133 (1997)).

Sandwich enzyme immunoassays also can be useful in certain methods of the invention. In a two-antibody sandwich assay, a first antibody is bound to a solid support, and the antigen is allowed to bind to the first antibody. The amount of c-Met is quantitated by measuring the amount of a second antibody that binds to it.

Quantitative western blotting also can be used to detect c-Met or to determine a level of c-Met in a method of the invention. Western blots can be quantitated by well known methods such as scanning densitometry. As an example, protein samples are electrophoresed on 10% SDS-PAGE Laemmli gels. Primary murine monoclonal antibodies, for example, against c-Met are reacted with the blot, and antibody binding confirmed to be linear using a preliminary slot blot experiment. Goat anti-mouse horseradish peroxidase-coupled antibodies (BioRad) are used as the secondary antibody, and signal detection performed using chemiluminescence, for example, with the Renaissance chemiluminescence kit (New England Nuclear; Boston, Mass.) according to the manufacturer's instructions. Autoradiographs of the blots are analyzed using a scanning densitometer (Molecular Dynamics; Sunnyvale, Calif.) and normalized to a positive control. Values are reported, for example, as a ratio between the actual value to the positive control (densitometric index). Such methods are well known in the art as described, for example, in Parra et al., *J. Vasc. Surg.* 28:669-675 (1998).

As described herein above, immunoassays including, for example, enzyme-linked immunosorbent assays, radioimmunoassays and quantitative western analysis, can be useful in the diagnostic methods of the invention. Such assays typically rely on one or more antibodies, for example, anti-c-Met antibodies. As used herein, the term "antibody" is used in its broadest sense to include polyclonal and monoclonal antibodies, as well as polypeptide fragments of antibodies that retain binding activity for c-Met of at least about $1 \times 10^5 \text{M}^{-1}$. One skilled in the art understands that antibody fragments

such as anti-c-Met antibody fragments and including Fab, F(ab')₂ and Fv fragments can retain binding activity for c-Met and, thus, are included within the definition of the term antibody as used herein. Methods of preparing monoclonal and polyclonal antibodies are routine in the art, as described, for example, in Harlow and Lane, supra, 1988.

The term antibody, as used herein, also encompasses non-naturally occurring antibodies and fragments containing, at a minimum, one V_H and one V_L domain, such as chimeric antibodies, humanized antibodies and single chain Fv fragments (scFv) that specifically bind c-Met. Such non-naturally occurring antibodies can be constructed using solid phase peptide synthesis, produced recombinantly or obtained, for example, by screening combinatorial libraries consisting of variable heavy chains and variable light chains as described by Borrebaeck (Ed.), *Antibody Engineering* (Second edition) New York: Oxford University Press (1995). A variety of useful anti-c-Met monoclonal and polyclonal antibodies are well known in the art and, in many cases, are commercially available.

The present invention also provides, inter alia, novel methods for diagnosis and prognosis evaluation for cancer.

In one aspect, the amount of soluble c-Met in a biological sample is determined in different patient samples for which either diagnosis or prognosis information is desired, to provide profiles. An profile of a particular sample is essentially a "fingerprint" of the state of the sample. A normal state may be distinguished from a cancerous state, and within cancerous states, different prognosis states (good or poor long term survival prospects, for example) can be determined. Diagnosis may be done or confirmed by comparing patient samples with the known profiles. By assessing the evolution of c-Met levels at different times during disease progression, the stage of disease can be determined as well as the likely prognosis.

A principle of diagnostic testing is the correlation of the results of a procedure with particular clinical parameters. The correlation necessarily involves a comparison between two or more groups distinguished by the clinical parameter. A clinical parameter could be, for example, presence or absence of disease, risk of disease, stage of disease, severity of disease, class of disease or response to treatment of disease. Accordingly, the diagnostician uses this correlation to qualify the status of a subject with respect to the clinical parameter. That is, the diagnostician uses the results of a procedure on a subject to classify or diagnose a subject status with respect to a clinical parameter, the confidence of the diagnosis/classification being related to the classifying or splitting power of the signs or symptoms used in the test.

The methods described herein for quantifying soluble c-met in a biological sample provides information which can be correlated with pathological conditions, predisposition to disease, therapeutic monitoring, risk stratification, among others. Although the data generated from the methods of this invention is suited for manual review and analysis, data processing using high-speed computers can be utilized as well.

The present methods are particularly useful for diagnosing conditions, evaluating whether certain drugs will have a desired effect, and determining prognoses. The present methods can be used for early detection of cancer as well as for the optimization of treatment protocols and analysis of biopsy samples. The present methods can also be used to optimize chemotherapy through assessment of the effect that the chemotherapy has on c-Met levels.

In certain embodiments, the invention provides methods for monitoring the progression of cancer in a patient. The method comprises the steps of providing a first biological sample from the patient, preferably a urine and/or blood

plasma sample, measuring the amount of soluble c-Met in the first biological sample at a first time point, providing a second biological sample from the patient, measuring the amount of soluble c-Met in the second biological sample at a second time point, and determining progression of the disease state in the patient based upon the change in the amount of soluble c-Met in the biological sample. In certain embodiments, the amount of soluble c-Met ectodomain in a urine and blood plasma sample will be measured at a first time point and the amount of soluble c-Met ectodomain in a second urine and blood plasma sample will be measured at a second time point and the combined measurement will be used to determine progression of the cancer in the patient. By measuring the amount of c-Met in a patient sample over time, a clinician will be able to determine whether the cancer has, for example, regressed and whether the subject has been effectively treated. A clinician can therefore utilize these measurement for tailoring treatment appropriately. A subject whose cancer has regressed after treatment with an anti-cancer agent will have less detectable c-Met than he did before the treatment. Similarly, a subject whose cancer has remained stable during treatment will have similar levels of c-Met as he did before treatment, and a subject whose cancer has worsened will have increased c-Met levels.

In certain embodiments, the present invention provides methods for measuring the response to cancer therapy comprising the steps of providing a first biological sample, preferably a urine and/or blood plasma sample, measuring the amount of soluble c-Met in the first biological sample at a first time point, providing a second biological sample from the patient, measuring the amount of soluble c-Met in the second biological sample at a second time point, and determining response in the patient based upon the change in the amount of soluble c-Met present in the biological sample. In certain embodiments, the amount of soluble c-Met ectodomain in a urine and blood plasma sample will be measured at a first time point and the amount of soluble c-Met ectodomain in a second urine and blood plasma sample will be measured at a second time point and the combined measurement will be used to determine patient response. The subject may be a positive responder, poor responder, or non-responder. For use herein, a positive responder, is a subject who positively responds to treatment, i.e., a subject who experiences success in amelioration of the cancer, including any objective or subjective parameter such as abatement; remission; diminishing of symptoms or making the condition more tolerable to the patient; slowing in the rate of degeneration or decline; making the final point of degeneration less debilitating; or improving a subject's physical or mental well-being. A positive responder is one in which any toxic or detrimental side effects of the biologically active agent is outweighed in clinical terms by therapeutically beneficial effects. A non-responder is a subject who doesn't respond to the treatment or doesn't respond to a satisfactory level. A poor responder is a subject who responds to treatment but not at the level of the positive responder.

In certain embodiments, the cancer therapy will include the administration of an anti-cancer drug. A clinician or other suitable professional can use the information regarding a subject's likely response level to certain anti-cancer drugs to determine an appropriate treatment regimen for the patient. In certain embodiments, increasing dosages can be provided for those patients that are indicated to be poor responders or non-responders. In certain embodiments, patients that are indicated to be poor responders, or non-responders will receive a different class of drugs or therapy. In certain

embodiments, the cancer therapy will be ionizing radiation, hormone ablation therapy, and the like.

Generally, the amount of soluble c-Met ectodomain present in the sample of a patient with premalignant cancer will be less than the amount of soluble c-Met ectodomain present in the sample of a patient with a low grade cancer, and the amount of soluble c-Met ectodomain present in the sample of a patient with a low grade cancer will be less than the amount of soluble c-Met ectodomain present in the sample of a patient with a high grade metastatic cancer.

Anti-cancer drugs are known in the art and include, for example chemotherapeutic agents, alone, or in combination with, radiation treatment, surgical treatment, or treatments using biological or immunomodulatory agents. Chemotherapeutic drugs useful in treating cancer include alkylating agents, antimetabolites, natural products, hormones and antagonists (reviewed in B. A. Chabner and D. L. Longo Eds. *Cancer Chemotherapy and Biotherapy*, 3rd Edition, 2001). These include, for example, nitrogen mustards, including, for example, mechlorethamine, cyclophosphamide, ifosfamide, melphalan, and chlorambucil; ethylenimines and methylmelamines, including, for example, hexamethylmelamine and thiotepa; alkyl sulfonates, including, for example, busulfan, carmustine, lomustine, semustine, and streptozocin; triazines, including, for example, dacarbazine and temozolamide, folic acid analogs, including, for example, methotrexate and trimetrexate 5-fluoropyrimidines including, for example, fluorouracil, floxuridine, fltorafur, capecitabine, and eniluracil, cytidine analogs, including cytarabine; 5-azacytidine, gemcitabine, purine analogs and related inhibitors, including, for example, mercaptopurine, thioguanine, fludarabine, cladribine, and pentostatin; vinca alkaloids, including, for example, vinblastine, and vincristine; taxanes including paclitaxel and docetaxel, topoisomerase II inhibitors, including, for example, etoposide, amsacrine and teniposide; topoisomerase I targeting agents including, including, for example, camptothecin, topotecan, irinotecan, and karenitecin, antibiotics, including, for example, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicamycin, and mitomycin; enzymes, including, for example, L-Asparaginase; biological response modifiers, including, for example, IL-2, interferon-alfa IL-1, IL-2, IL-4, IL-12, tumor necrosis factor and macrophage colony stimulating factor, platinum coordination complexes, including, for example, cisplatin, oxaloplatin, and carboplatin; anthracenediones, including, for example, mitoxantrone; thalidomide and derivatives including, for example, revemid, proteasome inhibitors including, for example, bortezomib, substituted ureas, including, for example, hydroxyurea; methylhydrazine derivatives, including, for example, procarbazine; adrenocortical suppressants, including, for example, mitotane and aminoglutethimide; adrenocorticosteroids, including, for example, prednisone; progestins and dexamethasone, including, for example, hydroxyprogesterone caproate, medroxyprogesterone acetate, and megestrol acetate; estrogens, including, for example, diethylstilbestrol and ethinyl estradiol; antiestrogens, including, for example, tamoxifen; androgens, including, for example, testosterone propionate and fluoxymesterone; antiandrogens, including, for example, flutamide; and gonadotropin releasing hormone analogs, including, for example, leuprolide, aromatase inhibitors including, for example, anastrozole (brand name Arimidex®), exemestane (brand name Aromasin®), and letrozole (brand name Femara®); antibodies directed against cell surface molecules including, for example, rituximab, trastuzumab, CAMPATH, cetuximab and bevacizumab, including antibodies conjugated to toxins, including, for example, gem-

tuzumab, and antibodies conjugated to radioisotopes including, for example, ibritumomab; anti-cancer antibodies that have been humanized to avoid the development of human antimouse antibodies; small molecule tyrosine kinase inhibitors including, for example, gleevec and iressa (reviewed in Noble et al., *Science*. 2004; 303:1800-5); faranesyl transferase inhibitors including, for example, R115777 (tipifarnib, Zarnestra®), SCH66336 (lonafarnib, Sarasar®.) and BMS-214662, including formulations of chemotherapy drugs including, for example, liposomal formulations, including arsenic trioxide, including cancer differentiating agents including, for example, all trans retinoic acid, including cancer treatments of any kind that are expected to reduce tumor growth, tumor invasiveness, tumor metastasis or overall tumor burden. Methods of administering chemotherapeutic agents for treating cancer are known in the art. (Goodman and Gilman's *The Pharmacological Basis of Therapeutics*, Ninth Edition).

The term "cancer therapy" refers to cancer treatment of any kind expected to reduce tumor growth rate, tumor invasiveness, tumor metastasis, or overall tumor burden, including for example, the administration of anti-cancer agents, ionizing radiation therapy, hormone ablation therapy, surgical intervention, and the like.

The invention provides a number of methods, reagents, and compounds that can be used for the diagnosis and prognosis of cancer. It is to be understood that this invention is not limited to particular methods, reagents, compounds, compositions, or biological systems, which can, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting. As used in this specification and the appended claims, the singular forms "a", "an", and "the" include plural referents unless the content clearly dictates otherwise.

About" as used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass variations of $\pm 20\%$ or $\pm 10\%$, more preferably $\pm 5\%$ even more preferably $\pm 1\%$, and still more preferably $\pm 0.1\%$ from the specified value, as such variations are appropriate to perform the disclosed methods.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although any methods and materials similar or equivalent to those described herein can be used in the practice for testing of the present invention, the preferred materials and methods are described herein.

EXAMPLES

Reagents and Cell Culture

Full length purified recombinant human hepatocyte growth factor (HGF) protein was obtained from R&D Systems (294-HG, Minneapolis, Minn.). Antibodies against c-Met were obtained from R&D Systems (Minneapolis, Minn.) or Upstate Biotechnology (DO-24 anti-c-Met mAb; Lake Placid, N.Y.) as noted below.

The following human normal/tumor and primary tumor/metastasis cell line pairs derived from single individuals were obtained from ATCC and maintained according to ATCC recommendations: CRL7636 (normal skin) and CRL7637 (skin melanoma), HTB125 (normal mammary gland) and HTB126 (mammary gland ductal carcinoma), and CCL228 (colorectal adenocarcinoma) and CCL227 (lymph node metastasis of colorectal adenocarcinoma). The human cell line pair UOK124 (renal cell carcinoma) and UOK124 LN

(lymph node metastasis of renal cell carcinoma), as well as UOK261 (bladder cancer) were developed at the Urologic Oncology Branch, National Cancer Institute, National Institutes of Health, Bethesda, Md. 20892 USA; Athauda et al., *Clin. Cancer Res.* 12: 4154-4162, 2006). UOK124, UOK124 LN, UOK261, A431 (epidermoid carcinoma), U-87 MG (glioblastoma), C100 (breast carcinoma), H1.177 (C100 transfected with Nm23), PC3 (prostate carcinoma), PC3M (PC3 derived metastatic variant) and PC3M MxA (PC3M transfected with MxA) cells were maintained in DMEM supplemented with 10% fetal bovine serum, antibiotics and antimycotics. B5/589 human mammary epithelial cells were cultured as previously described (Stahl et al., *Biochem J* 1997; 326:763-72). The MCF-10A derived breast epithelial cell lines 141-M4 (MCF-10A or M1, MCF10AT1k.c12 or M2, MCF-10CA1h or M3 and MCF-10CA1a.c11 or M4) (Santner et al., *Breast Cancer Res Treat* 2001; 65:101-10) were obtained from the Barbara Ann Karmanos Cancer Institute (Detroit, Mich.) and were maintained in DMEM/F-12 supplemented with 5% horse serum, 0.029M sodium bicarbonate, 10 mM HEPES, penicillin and streptomycin at 37° C. 5% CO₂. This culture medium was supplemented with 10 µg/ml insulin, 20 ng/ml EGF, 0.5 µg/ml hydrocortisone and 100 ng/ml cholera toxin for the M1 and M2 cell lines.

SDS-PAGE and Immunoblotting

Analysis of cellular and soluble c-Met by immunoprecipitation and immunoblotting was performed as described previously (Stahl et al., *Biochem J* 1997; 326:763-72). Samples for soluble c-Met analysis were obtained by harvesting conditioned media (cell culture supernatants) prior to detergent extraction; conditioned media was subjected to high speed centrifugation and 0.2 micron filtration to remove cells and debris. To obtain cellular c-Met samples, intact cells were serum deprived for 16 or 24 h as noted, lysed in cold buffer containing non-ionic detergents and protease and phosphatase inhibitors and cleared by high speed centrifugation. After immunoprecipitation of clarified detergent extracts for 2 h on ice, immunocomplexes were captured using immobilized protein-G (GammaBind G-Agarose, Pharmacia), washed, eluted with SDS sample buffer and subjected to SDS-PAGE and electrophoretic transfer to PVDF membranes (Immobilon P, Millipore). Immunodetection was performed by conventional methods using enhanced chemiluminescence (ECL Plus, Amersham).

Electrochemiluminescence Immunoassays

Streptavidin coated 96-well plates designed specifically for use in a Meso Scale Discovery (MSD) Sector 2400 Imager were first blocked with I-Block solution (300 µl/well, 1 h; Applied Biosystems catalog #A1300) that had been prepared according to the manufacturers instructions. Wells were then washed 3 times with PBS (150 µl/well).

For c-Met ectodomain assays, a biotin-tagged, affinity purified c-Met ectodomain specific capture antibody (R&D BAF 358 diluted in 0.5% BSA in PBS) was added to each well (5 µg/ml, 25 µl/well) for 1 h with shaking. Wells were washed 3 times with PBS before adding samples or standards (R&D 358-MT recombinant c-Met ectodomain-IgG-Fc fusion protein in PBS+0.5% BSA). Standards (100 µl/well) were added to generate a curve from 0.01 ng/ml to 100 ng/ml in semi-log increments for 1 h with shaking. Samples for soluble c-Met analysis were obtained by harvesting conditioned media (cell culture supernatants) prior to detergent extraction; conditioned media was subjected to high speed centrifugation and 0.22 micron filtration to remove cells and debris, then stored at -80° C. prior to c-Met quantitation. In some cases conditioned media was concentrated using Centricon YM-10 microconcentration units (Millipore, Inc.) prior to analysis,

as described in the text, or subjected to a single round of immunodepletion using the human c-Met ectodomain specific monoclonal antibody DO24 (Upstate Biotechnology) followed by antibody capture with protein G agarose. Wells were washed 3 times with PBS before adding detection antibody (R&D AF 276 labeled with MSD Sulfoltag diluted in 0.5% BSA in PBS) at 1 µg/ml, 25 µl/well for 1 h with shaking. Wells were then washed 4 times with PBS before adding MSD Read Buffer T with surfactant (150 µl/well) and then read immediately in a MSD Sector 2400 Imager.

For cellular c-Met assays, intact cells at 80% confluence were serum deprived for 24 h, washed twice with cold PBS and then lysed in cold buffer containing non-ionic detergents and protease and phosphatase inhibitors. Detergent extracts were clarified by high speed centrifugation and applied to 96-well plates as described for soluble c-Met ectodomain samples, above.

c-Met assays can be performed on biological sample, e.g., biological tissue or biological fluid from a patient or human patient. The biological tissue can be assayed for c-Met ectodomain as a detergent extract of the tissue. Biological tissue from a patient can be extracted with non-ionic detergent buffer solutions as described herein for preparing cell extracts. The tissue can be minced or ground (at 4° C. or on ice) to maximize surface area available for extraction, and generally the extraction (also on ice) is performed for a much longer period (2-3 hours) as opposed to a few minutes for cultured cells. See, for example, U.S. Pat. No. 5,674,753.

For HGF immunoassays, I-Block treated plates were coated as described above with an affinity purified HGF specific capture antibody (R&D MAB 694 diluted in 0.5% BSA in PBS) that had been biotin labeled. Wells were washed as described above before adding samples or standards (R&D 294-HG recombinant HGF protein in PBS+0.5% BSA); standards were added to generate a curve from 0.03 ng/ml to 30 ng/ml in semi-log increments. Wells were washed before adding detection antibody (R&D AF-294 labeled with MSD Sulfoltag diluted in 0.5% BSA in PBS) at 1 µg/ml, 25 µl/well. Wells were then washed 4 times with PBS before adding Read Buffer T and reading in a MSD Sector 2400 Imager.

All samples were measured in quadruplicate unless otherwise noted. Mean values from negative control wells were subtracted from all other raw values and a standard curve was then constructed by plotting signal intensity against 358-MT c-Met Fc fusion protein or recombinant HGF protein standard concentration. A non-linear regression curve fitting algorithm (Microsoft Excel or GraphPad Prism software) was used to generate an equation from which sample values for c-Met or HGF concentration were derived from mean signal intensity values. Mean values among groups were compared for statistically significant differences using unpaired t-test (paired human cell lines) or analysis of variance (ANOVA; MCF10A derived cell lines); R² and P values are presented in the text and figure legends.

Human Tumor Xenografts in Mice

Subconfluent UOK261 and U-87 MG cells were trypsinized, washed two times in PBS to remove serum and then resuspended in HBSS at a concentration of 1×10⁷ cells/ml. One hundred microliters containing 1×10⁶ cells were injected subcutaneously into the right flank of 12-week-old male SCID/beige mice (Taconic Inc., Germantown, N.Y.). Tumor growth was monitored weekly by caliper measurements and tumor volume was calculated based on the formula (length.times.length.times.width)/6. Blood was obtained via the tail vein or retromandibular perforation; samples were collected in the presence of citrate as an anticoagulant were centrifuged at 1500 rpm, 4° C. to remove cells and plasma

samples were stored at -80°C . prior to analysis. At each of three weekly time points 4 mice were sampled per group. Samples were analyzed for soluble c-Met using the electrochemiluminescence immunoassay described above; values shown represent the mean of triplicate measurements. In FIG. 6, standard deviations are smaller than the symbol size in all cases. Mouse urine sample pH was adjusted to pH 7.5 with Trizma-HCl, 2M, pH 8.0 (Sigma), then centrifuged at 1500 rpm for 5 minutes to remove cells and debris. Pooled SCID/beige normal mouse plasma was used as a diluent for standard curves and was obtained in bulk from Taconic Inc. Pooled urine from normal SCID/beige mice was pH-corrected and used as a diluent for standard curves run in assays to measure c-Met concentration in mouse urine samples. R^2 and P values from linear regression analysis of mouse data are presented in the text and figure legends.

c-Met Shedding by Cultured Cells: Characterization and Assay Development

c-Met ectodomain shedding was examined in a cultured cell model of breast cancer progression where successive derivatives of the parent cell line show increasing malignancy (Santner et al., *Breast Cancer Res Treat* 2001; 65:101-10). MCF-10A (M1) is a spontaneously immortalized normal breast epithelial cell line which was transfected with activated Hras and xenografted in mice to obtain the premalignant MCF10AT1k.c12 (M2) cell line. Subsequent passages in mice and single cell cloning facilitated the isolation of cell lines that produced tumors with the phenotypic characteristics a low grade carcinoma (MCF-10CA1h or M3) and a high grade metastatic carcinoma (MCF-10CA1a.c11 or M4). In contrast to many carcinoma derived cell lines where c-Met is overexpressed, cell surface c-Met expression among the four cultured cell lines appeared to decrease with increasingly malignant phenotype (FIG. 1A). Nonetheless, analysis of c-Met shedding over 16 h by immunoblotting with ectodomain-specific monoclonal antibodies showed progressively higher ectodomain levels from normal cells to those displaying a metastatic phenotype (FIG. 1B). c-Met ectodomain fragments of approximate molecular masses 75, 85 and 100 kDa were the predominant species observed, similar to those observed in the normal human mammary epithelial cell line B5/589 (FIG. 1B) and to the predominant species present in cell culture supernatants and human plasma as reported previously [3-7]. These c-Met reactive protein bands were detectable with several antibodies against the c-Met ectodomain, but not with polyclonal antisera raised against a peptide corresponding to the carboxyl terminal c-Met sequence (data not shown).

To better characterize c-Met shedding across a range of cell lines and biological samples, a two site immunoassay was developed to provide greater sensitivity, higher throughput and more precise quantitation than immunoblotting. Distinct human c-Met specific antibodies are used for capture and detection; capture antibodies were biotin labeled for use with streptavidin coated multiwell plates and the amount of detection antibody bound was measured by electrochemiluminescence generated using ruthenium chelates in the presence of sacrificial redox co-reactants. The optimized c-Met immunoassay has a threshold of detection of 750 fg for the purified recombinant c-Met ectodomain protein and a dynamic range of four log units (FIG. 2A). Sample quenching or hypersensitivity that could occur in complex biological samples was investigated by adding known amounts of recombinant c-Met ectodomain to cultured cell supernatants from B5/589 mammary epithelial cells (data not shown) or A431 epidermoid carcinoma cells (FIG. 2B), both of which contained shed c-Met ectodomain as determined by immunoblotting. Agree-

ment between expected and observed c-Met concentrations indicated the absence of any interference (FIG. 2B). Ultrafiltration of B5/589 conditioned medium through a membrane with a 10 kDa cutoff removed all of the c-Met signal, consistent with the ectodomain molecular masses observed by immunoblotting (FIG. 2C). The c-Met selectivity of the immunoassay was tested by subjecting B5/589 conditioned medium to immunodepletion using a monoclonal antibody against c-Met ectodomain (DO-24) that was distinct from either capture or detection antibody used in the immunoassay. As shown in FIG. 2C, a single round of immunodepletion resulted in an 80% loss of signal.

To determine whether shed c-Met ectodomain fragments could act as competitive antagonists of ligand binding by cell surface receptors, the two site c-Met immunoassay was adapted to analyze the HGF binding capacity of c-Met fragments cultured cells conditioned medium. A purified recombinant full length c-Met ectodomain-IgG Fc fusion protein was used as positive control for HGF binding. The amount of c-Met ectodomain present in B5/589 conditioned medium was determined from a c-Met-Fc standard curve and assays were designed such that equal amounts of c-Met ectodomain in B5/589 conditioned medium and in the c-Met Fc fusion protein preparation were captured in replicate wells. Purified recombinant HGF was added to two sets of wells at concentrations of 10, 100 and 300 ng/ml; one set of wells was then detected using anti-c-Met and the other with anti-HGF. An HGF standard curve was created using the same HGF detection antibody and an HGF-specific capture antibody. No change in c-Met detection was observed in c-Met Fc containing wells that had bound HGF, indicating that bound HGF did not interfere with c-Met capture or detection (data not shown). As shown in FIG. 2D, the c-Met Fc fusion protein bound HGF in direct proportion to the amount added, whereas no HGF binding was detected for the c-Met fragment present in B5/589 conditioned medium. These results, consistent with prior studies (Wajih et al., *Circ Res* 2002; 90:46-52), show that the predominant c-Met ectodomain fragments found in B5/589 cell culture supernatants have lost meaningful HGF binding capability. In light of the observation that the B5/589 c-Met ectodomain fragments are very similar in size and distribution to those shed by a variety of cultured cell lines as well as to what has been found in human plasma (Wajih et al., *Circ Res* 2002; 90:46-52), c-Met shedding does not yield a soluble receptor fragment that could compete efficiently for ligand binding by intact cellular c-Met.

c-Met Shedding Correlates with Malignancy in Cultured Cell Cancer Models

Quantitative analysis of c-Met shedding by the MCF 10A derived breast cancer cell lines that were analyzed previously by immunoblotting showed excellent agreement in the trends among the cell lines for both cellular c-Met expression and c-Met ectodomain shedding (FIG. 1B and FIG. 3, upper panels). The external recombinant protein standard in the immunoassay allows results to be expressed in absolute terms, e.g. receptor concentration or number per total cellular protein or per cell. This enables realistic comparisons between successive experiments, different cell lines or with other biological samples. Manipulation of the numerical results also offers insight into trends in shedding relevant to molecular mechanism, such as determining shedding rate per cellular receptor (FIG. 3, 3rd panel) or the percentage of cellular receptor shed per time interval (FIG. 3, bottom panel). In the MCF10A derived model of breast cancer progression, the steady state cellular c-Met expression level is progressively and significantly lower at each step of increasingly malignant phenotype (ANOVA $R^2=0.998$; $P<0.0001$). In contrast, c-Met shedding

is significantly increased with the change from premalignant (M2) to malignant (M3) phenotype (FIG. 3, 2nd panel, t-test $P < 0.001$). This trend is maintained in the metastatic (M4) cell line (FIG. 3, 2nd and 3rd panels) and shedding in M4 is nearly four-fold higher than its normal counterpart M1 when expressed as a function of available cellular c-Met (FIG. 3, bottom panel; 4 group ANOVA $R^2 > 0.995$; $P < 0.0001$). The observed trends in expression and shedding are consistent with significant and progressive increases in proteolytic activity characteristic of advancing breast cancer.

Cellular and soluble c-Met levels in a series of paired human cell lines, each obtained from a single cancer patient, derived from tumor vs. corresponding normal tissue or, primary tumor vs. distant metastatic lesion were measured. Specifically, the paired cell lines were derived from normal skin (CRL7636) and skin melanoma, (CRL7637), normal mammary gland (HTB125) and mammary gland ductal carcinoma (HTB126), colorectal adenocarcinoma (CCL228) and lymph node metastasis of colorectal adenocarcinoma (CCL227), and renal cell carcinoma (UOK124) and lymph node metastasis of renal cell carcinoma (UOK124 LN).

All four cell line pairs exhibited one overall trend similar to that displayed by the MCF10A model: significantly greater c-Met shedding rate with more malignant phenotype (FIG. 4, 2nd, 3rd and 4th panels). In contrast, cellular c-Met levels varied widely among the paired lines (FIG. 4, top panel). Relative to the normal skin derived cell line CRL7636, cellular c-Met was 60% lower than in the paired melanoma derived cell line CRL7637 (top panel), while soluble c-Met concentration was 60% greater (2nd panel), contributing to a 4-fold greater c-Met shedding rate per cell protein (3rd panel) and per receptor (4th panel) correlating with the acquisition of malignancy. These findings suggest that, similar to the MCF10A derived cell lines, significantly increased proteolytic activity was the primary cause of the observed trend. In contrast, in the cell line pair derived from normal mammary gland and ductal carcinoma (HTB125 and HTB126), acquisition of malignancy correlated with 2.5 fold greater cellular c-Met level (top panel) and 4-fold greater soluble c-Met level (2nd panel) producing a 2.5 fold greater c-Met shedding rate per cell protein (3rd panel). This trend represented only a modest increase in the percentage of cellular c-Met shed in 24 h (bottom panel), suggesting that unlike the MCF10A derived breast cancer model, receptor overexpression contributed significantly to the increased receptor shedding observed with malignant transformation. This difference in underlying mechanism of increased c-Met shedding between two breast cancer cell models could reflect the oncogenic heterogeneity in breast cancer that is becoming more broadly appreciated.

Similar in trend to HTB125 and HTB126, but remarkable in magnitude, the cell line derived from a lymph node metastasis of colorectal carcinoma (CCL227) showed 40-fold greater cellular c-Met expression relative to the primary tumor derived cell line (CCL228) and 75-fold greater rate of c-Met shedding per total cell protein (FIG. 4, upper two panels). Correcting these values for receptor protein yielded a 2.7 fold greater shedding rate in the metastasis-derived line relative to the primary tumor line (FIG. 4, 3rd panel). The renal cell carcinoma derived cell line UOK124 and corresponding lymph node metastasis derived line UOK124 LN showed little difference in cellular c-Met expression (top panel), while soluble c-Met level was 60% greater in UOK124 LN (2nd panel), contributing to a 3-fold greater c-Met shedding rate per cellular receptor (3rd panel) and a doubling of the percentage of cellular receptor shed in 24 h (bottom panel). Thus, similar to the trends observed for

the MCF10A derived breast carcinoma model and the skin melanoma cell line pair, significantly greater proteolytic activity was likely to be responsible for the greater c-Met shedding rate that correlated with increased malignancy.

In addition to cultured normal/tumor or tumor/metastasis cell line pairs, cellular and soluble c-Met levels in two genetically engineered models representing reconstitution of metastasis suppressor genes in aggressively malignant prostate and breast tumor derived cell lines were examined. The prostate cancer cell line PC3 is tumorigenic in mice but not metastatic; PC3M is a PC3 derived cell line that is aggressively tumorigenic and metastatic (Chu et al., *Cancer Genet Cytogenet* 2001; 127:161-7.) This phenotypic difference was exploited to identify genes whose loss could contribute to metastasis in prostate cancer, leading to the identification of the MxA gene as a suppressor of metastasis (Aebi et al., *Mol Cell Biol* 1989; 9:5062-72 and unpublished observations). Upon reconstitution of MxA expression in PC3M, the aggressive metastatic phenotype of cultured cell xenografts in mice is reverted to that of the parental cell line (unpublished observations). Cellular c-Met expression in PC3M was nearly double that of PC3; restoration of MxA expression was associated with an even greater cellular c-Met expression level (FIG. 5A, top panel). Shedding by PC3M was 50% greater than that of the parental cell line, consistent with the overall trend of increased shedding with increasing malignancy observed in other cell models (FIG. 5A, 2nd panel), and apparently driven by increased c-Met expression. While the absolute level of c-Met shedding in PC3M and PC3M MxA transfectants was similar, a lower the rate of c-Met shedding per cellular receptor was detected in MxA transfectants (FIG. 5A, 3rd and 4th panels). These results suggest that MxA expression had no significant effect on c-Met expression, but suppressed the shedding mechanism to a level comparable to that displayed by the non-metastatic PC3 cell line.

In contrast to the PC3 based model of prostate cancer metastasis, a dramatic effect on cellular c-Met expression was exhibited by the metastasis suppressor gene Nm23 in a genetically engineered cell model of breast cancer progression (FIG. 5B). In this model, the breast cancer cell line C100, which aggressively forms primary tumors and metastases in mouse xenografts, was reverted to a non-metastatic phenotype upon transfection with Nm23 (H1.177) (Leone et al., *Oncogene* 1993; 8:2325-33). Correlated with the loss of metastatic phenotype was 70% lower cellular c-Met expression in H1.177 relative to the parental cell line (FIG. 5B, top panel) and 75% lower soluble c-Met production (FIG. 5B, 2nd panel). A 20% lower level of c-Met shedding per cellular receptor was also associated with Nm23 expression (FIG. 5B, 3rd and 4th panels), suggesting that shedding was attenuated primarily through the suppression of c-Met expression by Nm23.

c-Met Shedding Correlates with Tumor Burden in Tumor Xenograft Mouse Models

It was observed that cultured tumor cells expressing c-Met tended to shed more c-Met ectodomain than their normal tissue counterparts, independent of changes in overall c-Met expression level, and that this tendency was enhanced with increasingly malignant phenotype. Two different human tumor cell lines were injected subcutaneously with known c-Met shedding rates (data not shown) into immunocompromised (SCID/Beige) mice and thereafter measured tumor volume and plasma soluble c-Met levels at weekly intervals. The cell line UOK261 was derived from a human bladder carcinoma and displayed a relatively high level of soluble c-Met in culture, whereas the cell line U-87 MG was derived from a human glioblastoma and displayed a lower level of

c-Met shedding in culture. c-Met signaling is suspected of playing an important role in the progression of both cancers represented by these models [Burgess et al., *Cancer Res* 2006; 66: In Press; Rosen et al., *J Urol* 1997; 157:72-8; Cheng et al., *J Clin Oncol* 2002; 20:1544-50.] Note that the antibodies used in the immunoassay described here do not cross-react with mouse c-Met, thus the assay was conducted in the absence of any normal soluble murine c-Met background; any c-Met detected originated from the human tumor xenografts. Pooled plasma samples obtained from the same strain of mice was used as a diluent for the recombinant c-Met ectodomain-Fc standard curve, so that the absolute soluble c-Met values obtained could be related directly to those obtained from cultured cell experiments, as well as to future animal studies.

Remarkably, soluble human c-Met was easily detected in plasma samples obtained from several mice receiving UOK261 xenografts more than a week before subcutaneous tumors became palpable (data not shown). Non-palpable tumors up to 5.0 mm³ in volume are difficult to detect radiologically and represent an early but nonetheless clinically relevant stage of tumorigenesis. The soluble c-Met fragments measured in the immunoassay were similar in size to the fragments found in cultured cell conditioned media, as determined by SDS-PAGE and immunoblotting (data not shown). With increasing tumor mass, each mouse showed significant weekly increases in plasma c-Met levels ($P < 0.05$; data not shown); these data were pooled and plotted as plasma c-Met against tumor volume (FIG. 6, left upper panel). Regression analysis demonstrated a direct linear relationship between circulating soluble c-Met concentration and tumor burden (FIG. 6, left upper panel, $R^2 = 0.944$; $n = 4$ animals). Consistent with the lower level of c-Met shedding by cultured U-87 MG cells, soluble c-Met was not detected in plasma samples from mice receiving the U-87 MG glioblastoma xenografts before the tumors became palpable. Nonetheless, the smallest measurable tumors were each associated with plasma c-Met levels that were well above the threshold of detection, and each mouse showed significantly increasing c-Met concentrations measured at weekly intervals ($P < 0.05$; data not shown). Regression analysis of the pooled data plotted as c-Met concentration against tumor volume also supported a direct linear soluble c-Met/tumor burden relationship (FIG. 6, right upper panel, $R^2 = 0.933$; $n = 4$ animals).

As shown in the left lower panel of FIG. 6, subcutaneous UOK261 xenografts produced urinary soluble c-Met concentrations that were detectable at very small tumor sizes and that increased linearly with tumor burden over weekly intervals ($R^2 = 0.922$, $P < 0.01$, $n = 4$ animals). Urinary soluble c-Met concentrations were typically less than 1/1000th of corresponding plasma c-Met concentrations (FIG. 6, left lower panel). Urine samples from mice receiving U-87 MG xenografts also contained soluble c-Met at early stages of tumor growth that increased significantly with time in each mouse ($P < 0.05$; data not shown), and group data showed a similar direct linear relationship between c-Met concentration and overall tumor burden (FIG. 6, right lower panel; $R^2 = 0.953$, $n = 4$ animals).

These results demonstrate that soluble c-Met produced by tumor derived cultured cell xenografts can be detected in plasma at very early stages of tumorigenesis, and that urinary c-Met concentration is a reliable indicator of the soluble c-Met level in plasma. These data further support the overall fidelity of c-Met shedding as an index of malignant phenotype, tumor progression and tumor burden in c-Met expressing models of oncogenesis. The simplicity and sensitivity of the assay described here make it amenable to high throughput screening.

c-Met Shedding in a Group of Normal Patients and Patients with Cancers Delineated in FIGS. 7-9.

Soluble c-Met was measured in urine samples as described in for mouse urine, with the exception that the standard curve was generated in phosphate buffered normal saline, not pooled mouse urine. Urine creatinine was measured using a standard, commercially available spectrophotometric method which is routinely used in medical laboratories. Samples for all assays were performed in triplicate or quadruplicate. Prior to measurements, samples were centrifuged to remove cells and debris, filtered through 0.22 micron filters, and the pH was adjusted to 7.4.

c-Met Shedding in a Group of Normal Patients and Patients with Bladder Cancer.

Soluble c-Met was measured in urine samples as described in for mouse urine, with the exception that the standard curve was generated in phosphate buffered normal saline, not pooled mouse urine. Urine creatinine was measured using a standard, commercially available spectrophotometric method which is routinely used in medical laboratories. Samples for all assays were performed in triplicate or quadruplicate. Prior to measurements, samples were centrifuged to remove cells and debris, filtered through 0.22 micron filters, and the pH was adjusted to 7.4. Values of soluble c-Met were obtained from the raw data using the standard curve (run in every assay); the standard curve was generated as described, final values are expressed as pg/ml. Urine creatinine was also measured against a standard curve and values are expressed as mg/ml. Corrected c-Met/creatinine values are expressed as pg/mg. Statistical analysis was performed using GraphPad Prism software, using assumptions and conventions accepted throughout the research community. t-test at the $p < 0.05$ level is the standard method for comparing the mean value obtained from patient replicates to the normal group mean, i.e. determining if any difference is unlikely to be due to chance. See FIG. 12.

Normals were healthy adult volunteers with no medical history of cancer and asymptomatic for diseases of the kidney, prostate, bladder or urinary tract at the time the sample was obtained, as established by routine medical laboratory tests, physical exam, and interview.

Bladder cancer samples were obtained from adult patients with biopsy proven (graded and staged by a licensed pathologist) bladder cancer or TCC (transitional cell carcinoma, which originates in ureteral epithelium, generally classified with bladder cancer). Patients had active or stable disease and samples were obtained prior to any surgical intervention.

Table 1 below provides the soluble c-Met measurements as pg per mg urine creatinine.

TABLE 1

Statistics	Normals	Bladder Cancer
Number of values	30	12
Minimum	892.3	1498
25% Percentile	3071	4343
Median	4344	6658
75% Percentile	5719	13990
Maximum	7742	33090
Mean	4193	10600
Standard Deviation	1835	9278
Standard Error	335	2678
Lower 95% Confidence Interval	3508	4709
Higher 95% Confidence Interval	4878	16500
KS Distance	0.09099	0.2363
P Value	$P > 0.10$	$P > 0.10$

TABLE 1-continued

Statistics	Normals	Bladder Cancer
Passed Normality Test (* = 0.05)	Yes	Yes
P Value Summary	ns	ns
Coefficient of Variation	43.76%	87.49%
Geometric mean	3689	7644

Adaptation of the Soluble c-Met Immunoassay for Analysis of Human Plasma Samples.

Initial measurements of soluble c-Met in human plasma samples obtained from normal healthy volunteers, using the recombinant 358-MT c-Met-Ig fusion protein (R&D Systems) as standard and the same assay protocol used for cultured cell supernatants, revealed very high soluble c-Met levels were present.

Experiments confirm that the measurements were reliable and not due to a factor present in plasma other than bona fide c-Met protein:

1. Samples of goat plasma and mouse plasma were measured to determine whether a plasma constituent other than c-Met might contribute to signal intensity. Because the assay reagents do not cross-react with c-Met protein from mouse or goat, any reading above background would be grounds to suspect a contributory factor in human plasma other than c-Met. Neither goat nor mouse samples displayed soluble c-Met reading above background.

2. Samples of human plasma were diluted with three different buffers over a four log range and then measured in the assay. In many cases, artifactual readings do not decrease linearly with dilution. Using Tris-HCl, PBS or de-ionized water, the signal present in human plasma directly in proportion to dilution factor.

3. Samples of human plasma were subjected to denaturing conditions (heating to 95° C. for 30 minutes) prior to measurement of soluble c-Met in that assay. Under these conditions, many plasma proteins are precipitated. Samples subjected to this protocol did not show c-Met readings above background.

4. Samples of human plasma were subjected to immunodepletion of c-Met prior to measurement in the assay. Ideally, repeated rounds of immunodepletion should remove soluble c-Met from the sample, reducing the measured value to background levels. The protocol used and results obtained from a representative experiment are shown below. To control for the non-specific removal of c-Met from the sample by the protein G beads used to capture immunocomplexes, a mock immunodepletion was performed on human plasma samples in parallel which using a goat-derived IgG directed against a protein not present in human plasma.

5. Samples of human plasma subjected to immunodepletion were analyzed by SDS-PAGE and immunoblotting with another anti-c-Met antibody not used in the immunoassay. A standard curve of 358-MT samples were run in parallel to estimate the amount of soluble c-Met present. Results were consistent with the immunodepletion experiments shown

below (data not shown). Crude estimates of soluble c-Met in human plasma based on immunoblotting were in the range of 100 ng/ml.

Protocol for Immunodepletion (ID) of Soluble c-Met from Human Plasma:

1. To 5 ml human plasma add anti-c-Met R&D AF276 (1 . mu.g/ml of plasma) and incubate on ice for 1 hour; agitate gently every 15 min. In parallel for MOCK samples, add Goat IgG (Rockland) at the same concentration. Repeat all subsequent steps with both ID and MOCK samples.

2. Add Protein G-Sepharose (Amersham; 30 µl/ml plasma) and rotate at 4 C for 1 hour.

3. Spin at 3000 rpm at 4° C. 1-2 minutes in Eppendorf desktop centrifuge to pellet beads. Remove supernatant. Take sample from supernatant (store on ice) and repeat steps 1-3 three more times for a total of four immunodepletion steps.

In FIG. 13, the upper panel shows rounds 1 through 4 of immunodepletion on the category (x) axis, and signal strength in the assay on the Y-axis. The undepleted plasma samples value is shown as a blue bar next to each round of immunodepletion. Signal values are listed numerically above each bar. Background value is approximately 1500 units. The bottom panel shows the immunodepletion samples alone to better visualize the degree of depletion. Variation between triplicate samples is consistent less than 2%.

The results of all experiments support the conclusion that 100% of the signal generated by assay of soluble c-Met in plasma represent bona fide c-Met protein. Measurement of soluble c-Met protein in human plasma samples obtained from a small set of healthy volunteers suggests that the normal concentration range is approximately 100 ng/ml. Analysis of a larger sample set is underway. Given the sensitivity of the assay and the initial concentration estimate, it should be possible to dilute plasma samples 100-fold prior to measurement, using approximately 100 microliters of final volume per well. This should allow replicates of 4 to 6 wells/sample to be generated from very small (10 microliter) plasma samples.

From the foregoing description, various modifications and changes in the compositions and methods will occur to those skilled in the art. All such modifications coming within the scope of the appended claims are intended to be included therein. Each recited range includes all combinations and sub-combinations of ranges, as well as specific numerals contained therein.

All publications and patent documents cited above are hereby incorporated by reference in their entirety for all purposes to the same extent as if each were so individually denoted.

Although the foregoing invention has been described in detail by way of example for purposes of clarity of understanding, it will be apparent to the artisan that certain changes and modifications are comprehended by the disclosure and can be practiced without undue experimentation within the scope of the appended claims, which are presented by way of illustration not limitation.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 2

<210> SEQ ID NO 1

<211> LENGTH: 980

<212> TYPE: PRT

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 1

Met Lys Ala Pro Ala Val Leu Ala Pro Gly Ile Leu Val Leu Leu Phe
1           5           10           15

Thr Leu Val Gln Arg Ser Asn Gly Glu Cys Lys Glu Ala Leu Ala Lys
20           25           30

Ser Glu Met Asn Val Asn Met Lys Tyr Gln Leu Pro Asn Phe Thr Ala
35           40           45

Glu Thr Pro Ile Gln Asn Val Ile Leu His Glu His His Ile Phe Leu
50           55           60

Gly Ala Thr Asn Tyr Ile Tyr Val Leu Asn Glu Glu Asp Leu Gln Lys
65           70           75           80

Val Ala Glu Tyr Lys Thr Gly Pro Val Leu Glu His Pro Asp Cys Phe
85           90           95

Pro Cys Gln Asp Cys Ser Ser Lys Ala Asn Leu Ser Gly Gly Val Trp
100          105          110

Lys Asp Asn Ile Asn Met Ala Leu Val Val Asp Thr Tyr Tyr Asp Asp
115          120          125

Gln Leu Ile Ser Cys Gly Ser Val Asn Arg Gly Thr Cys Gln Arg His
130          135          140

Val Phe Pro His Asn His Thr Ala Asp Ile Gln Ser Glu Val His Cys
145          150          155          160

Ile Phe Ser Pro Gln Ile Glu Glu Pro Ser Gln Cys Pro Asp Cys Val
165          170          175

Val Ser Ala Leu Gly Ala Lys Val Leu Ser Ser Val Lys Asp Arg Phe
180          185          190

Ile Asn Phe Phe Val Gly Asn Thr Ile Asn Ser Ser Tyr Phe Pro Asp
195          200          205

His Pro Leu His Ser Ile Ser Val Arg Arg Leu Lys Glu Thr Lys Asp
210          215          220

Gly Phe Met Phe Leu Thr Asp Gln Ser Tyr Ile Asp Val Leu Pro Glu
225          230          235          240

Phe Arg Asp Ser Tyr Pro Ile Lys Tyr Val His Ala Phe Glu Ser Asn
245          250          255

Asn Phe Ile Tyr Phe Leu Thr Val Gln Arg Glu Thr Leu Asp Ala Gln
260          265          270

Thr Phe His Thr Arg Ile Ile Arg Phe Cys Ser Ile Asn Ser Gly Leu
275          280          285

His Ser Tyr Met Glu Met Pro Leu Glu Cys Ile Leu Thr Glu Lys Arg
290          295          300

Lys Lys Arg Ser Thr Lys Lys Glu Val Phe Asn Ile Leu Gln Ala Ala
305          310          315          320

Tyr Val Ser Lys Pro Gly Ala Gln Leu Ala Arg Gln Ile Gly Ala Ser
325          330          335

Leu Asn Asp Asp Ile Leu Phe Gly Val Phe Ala Gln Ser Lys Pro Asp
340          345          350

Ser Ala Glu Pro Met Asp Arg Ser Ala Met Cys Ala Phe Pro Ile Lys
355          360          365

Tyr Val Asn Asp Phe Phe Asn Lys Ile Val Asn Lys Asn Asn Val Arg
370          375          380

Cys Leu Gln His Phe Tyr Gly Pro Asn His Glu His Cys Phe Asn Arg
385          390          395          400

```

-continued

Thr Leu Leu Arg Asn Ser Ser Gly Cys Glu Ala Arg Arg Asp Glu Tyr
 405 410 415
 Arg Thr Glu Phe Thr Thr Ala Leu Gln Arg Val Asp Leu Phe Met Gly
 420 425 430
 Gln Phe Ser Glu Val Leu Leu Thr Ser Ile Ser Thr Phe Ile Lys Gly
 435 440 445
 Asp Leu Thr Ile Ala Asn Leu Gly Thr Ser Glu Gly Arg Phe Met Gln
 450 455 460
 Val Val Val Ser Arg Ser Gly Pro Ser Thr Pro His Val Asn Phe Leu
 465 470 475 480
 Leu Asp Ser His Pro Val Ser Pro Glu Val Ile Val Glu His Thr Leu
 485 490 495
 Asn Gln Asn Gly Tyr Thr Leu Val Ile Thr Gly Lys Lys Ile Thr Lys
 500 505 510
 Ile Pro Leu Asn Gly Leu Gly Cys Arg His Phe Gln Ser Cys Ser Gln
 515 520 525
 Cys Leu Ser Ala Pro Pro Phe Val Gln Cys Gly Trp Cys His Asp Lys
 530 535 540
 Cys Val Arg Ser Glu Glu Cys Leu Ser Gly Thr Trp Thr Gln Gln Ile
 545 550 555 560
 Cys Leu Pro Ala Ile Tyr Lys Val Phe Pro Asn Ser Ala Pro Leu Glu
 565 570 575
 Gly Gly Thr Arg Leu Thr Ile Cys Gly Trp Asp Phe Gly Phe Arg Arg
 580 585 590
 Asn Asn Lys Phe Asp Leu Lys Lys Thr Arg Val Leu Leu Gly Asn Glu
 595 600 605
 Ser Cys Thr Leu Thr Leu Ser Glu Ser Thr Met Asn Thr Leu Lys Cys
 610 615 620
 Thr Val Gly Pro Ala Met Asn Lys His Phe Asn Met Ser Ile Ile Ile
 625 630 635 640
 Ser Asn Gly His Gly Thr Thr Gln Tyr Ser Thr Phe Ser Tyr Val Asp
 645 650 655
 Pro Val Ile Thr Ser Ile Ser Pro Lys Tyr Gly Pro Met Ala Gly Gly
 660 665 670
 Thr Leu Leu Thr Leu Thr Gly Asn Tyr Leu Asn Ser Gly Asn Ser Arg
 675 680 685
 His Ile Ser Ile Gly Gly Lys Thr Cys Thr Leu Lys Ser Val Ser Asn
 690 695 700
 Ser Ile Leu Glu Cys Tyr Thr Pro Ala Gln Thr Ile Ser Thr Glu Phe
 705 710 715 720
 Ala Val Lys Leu Lys Ile Asp Leu Ala Asn Arg Glu Thr Ser Ile Phe
 725 730 735
 Ser Tyr Arg Glu Asp Pro Ile Val Tyr Glu Ile His Pro Thr Lys Ser
 740 745 750
 Phe Ile Ser Thr Trp Trp Lys Glu Pro Leu Asn Ile Val Ser Phe Leu
 755 760 765
 Phe Cys Phe Ala Ser Gly Gly Ser Thr Ile Thr Gly Val Gly Lys Asn
 770 775 780
 Leu Asn Ser Val Ser Val Pro Arg Met Val Ile Asn Val His Glu Ala
 785 790 795 800
 Gly Arg Asn Phe Thr Val Ala Cys Gln His Arg Ser Asn Ser Glu Ile
 805 810 815
 Ile Cys Cys Thr Thr Pro Ser Leu Gln Gln Leu Asn Leu Gln Leu Pro

-continued

His Pro Leu His Ser Ile Ser Val Arg Arg Leu Lys Glu Thr Lys Asp
 210 215 220
 Gly Phe Met Phe Leu Thr Asp Gln Ser Tyr Ile Asp Val Leu Pro Glu
 225 230 235 240
 Phe Arg Asp Ser Tyr Pro Ile Lys Tyr Val His Ala Phe Glu Ser Asn
 245 250 255
 Asn Phe Ile Tyr Phe Leu Thr Val Gln Arg Glu Thr Leu Asp Ala Gln
 260 265 270
 Thr Phe His Thr Arg Ile Ile Arg Phe Cys Ser Ile Asn Ser Gly Leu
 275 280 285
 His Ser Tyr Met Glu Met Pro Leu Glu Cys Ile Leu Thr Glu Lys Arg
 290 295 300
 Lys Lys Arg Ser Thr Lys Lys Glu Val Phe Asn Ile Leu Gln Ala Ala
 305 310 315 320
 Tyr Val Ser Lys Pro Gly Ala Gln Leu Ala Arg Gln Ile Gly Ala Ser
 325 330 335
 Leu Asn Asp Asp Ile Leu Phe Gly Val Phe Ala Gln Ser Lys Pro Asp
 340 345 350
 Ser Ala Glu Pro Met Asp Arg Ser Ala Met Cys Ala Phe Pro Ile Lys
 355 360 365
 Tyr Val Asn Asp Phe Phe Asn Lys Ile Val Asn Lys Asn Asn Val Arg
 370 375 380
 Cys Leu Gln His Phe Tyr Gly Pro Asn His Glu His Cys Phe Asn Arg
 385 390 395 400
 Thr Leu Leu Arg Asn Ser Ser Gly Cys Glu Ala Arg Arg Asp Glu Tyr
 405 410 415
 Arg Thr Glu Phe Thr Thr Ala Leu Gln Arg Val Asp Leu Phe Met Gly
 420 425 430
 Gln Phe Ser Glu Val Leu Leu Thr Ser Ile Ser Thr Phe Ile Lys Gly
 435 440 445
 Asp Leu Thr Ile Ala Asn Leu Gly Thr Ser Glu Gly Arg Phe Met Gln
 450 455 460
 Val Val Val Ser Arg Ser Gly Pro Ser Thr Pro His Val Asn Phe Leu
 465 470 475 480
 Leu Asp Ser His Pro Val Ser Pro Glu Val Ile Val Glu His Thr Leu
 485 490 495
 Asn Gln Asn Gly Tyr Thr Leu Val Ile Thr Gly Lys Lys Ile Thr Lys
 500 505 510
 Ile Pro Leu Asn Gly Leu Gly Cys Arg His Phe Gln Ser Cys Ser Gln
 515 520 525
 Cys Leu Ser Ala Pro Pro Phe Val Gln Cys Gly Trp Cys His Asp Lys
 530 535 540
 Cys Val Arg Ser Glu Glu Cys Leu Ser Gly Thr Trp Thr Gln Gln Ile
 545 550 555 560
 Cys Leu Pro Ala Ile Tyr Lys Val Phe Pro Asn Ser Ala Pro Leu Glu
 565 570 575
 Gly Gly Thr Arg Leu Thr Ile Cys Gly Trp Asp Phe Gly Phe Arg Arg
 580 585 590
 Asn Asn Lys Phe Asp Leu Lys Lys Thr Arg Val Leu Leu Gly Asn Glu
 595 600 605
 Ser Cys Thr Leu Thr Leu Ser Glu Ser Thr Met Asn Thr Leu Lys Cys
 610 615 620
 Thr Val Gly Pro Ala Met Asn Lys His Phe Asn Met Ser Ile Ile Ile

-continued

625	630	635	640
Ser Asn Gly His Gly Thr Thr Gln Tyr Ser Thr Phe Ser Tyr Val Asp 645 650 655			
Pro Val Ile Thr Ser Ile Ser Pro Lys Tyr Gly Pro Met Ala Gly Gly 660 665 670			
Thr Leu Leu Thr Leu Thr Gly Asn Tyr Leu Asn Ser Gly Asn Ser Arg 675 680 685			
His Ile Ser Ile Gly Gly Lys Thr Cys Thr Leu Lys Ser Val Ser Asn 690 695 700			
Ser Ile Leu Glu Cys Tyr Thr Pro Ala Gln Thr Ile Ser Thr Glu Phe 705 710 715 720			
Ala Val Lys Leu Lys Ile Asp Leu Ala Asn Arg Glu Thr Ser Ile Phe 725 730 735			
Ser Tyr Arg Glu Asp Pro Ile Val Tyr Glu Ile His Pro Thr Lys Ser 740 745 750			
Phe Ile Ser Thr Trp Trp Lys Glu Pro Leu Asn Ile Val Ser Phe Leu 755 760 765			
Phe Cys Phe Ala Ser Gly Gly Ser Thr Ile Thr Gly Val Gly Lys Asn 770 775 780			
Leu Asn Ser Val Ser Val Pro Arg Met Val Ile Asn Val His Glu Ala 785 790 795 800			
Gly Arg Asn Phe Thr Val Ala Cys Gln His Arg Ser Asn Ser Glu Ile 805 810 815			
Ile Cys Cys Thr Thr Pro Ser Leu Gln Gln Leu Asn Leu Gln Leu Pro 820 825 830			
Leu Lys Thr Lys Ala Phe Phe Met Leu Asp Gly Ile Leu Ser Lys Tyr 835 840 845			
Phe Asp Leu Ile Tyr Val His Asn Pro Val Phe Lys Pro Phe Glu Lys 850 855 860			
Pro Val Met Ile Ser Met Gly Asn Glu Asn Val Leu Glu Ile Lys Gly 865 870 875 880			
Asn Asp Ile Asp Pro Glu Ala Val Lys Gly Glu Val Leu Lys Val Gly 885 890 895			
Asn Lys Ser Cys Glu Asn Ile His Leu His Ser Glu Ala Val Leu Cys 900 905 910			
Thr Val Pro Asn Asp Leu Leu Lys Leu Asn Ser Glu Leu Asn Ile Glu 915 920 925			
Trp Lys Gln Ala Ile Ser Ser Thr Val Leu Gly Lys Val Ile Val Gln 930 935 940			
Pro Asp Gln Asn Phe Thr Gly Leu Ile Ala Gly Val Val Ser Ile Ser 945 950 955 960			
Thr Ala Leu Leu Leu Leu Leu Gly Phe Phe Leu Trp Leu Lys Lys Arg 965 970 975			
Lys Gln Ile Lys Asp Leu Gly Ser Glu Leu Val Arg Tyr Asp Ala Arg 980 985 990			
Val His Thr Pro His Leu Asp Arg Leu Val Ser Ala Arg Ser Val Ser 995 1000 1005			
Pro Thr Thr Glu Met Val Ser Asn Glu Ser Val Asp Tyr Arg Ala 1010 1015 1020			
Thr Phe Pro Glu Asp Gln Phe Pro Asn Ser Ser Gln Asn Gly Ser 1025 1030 1035			
Cys Arg Gln Val Gln Tyr Pro Leu Thr Asp Met Ser Pro Ile Leu 1040 1045 1050			

-continued

Thr Ser Gly Asp Ser Asp Ile Ser Ser Pro Leu Leu Gln Asn Thr 1055 1060 1065
Val His Ile Asp Leu Ser Ala Leu Asn Pro Glu Leu Val Gln Ala 1070 1075 1080
Val Gln His Val Val Ile Gly Pro Ser Ser Leu Ile Val His Phe 1085 1090 1095
Asn Glu Val Ile Gly Arg Gly His Phe Gly Cys Val Tyr His Gly 1100 1105 1110
Thr Leu Leu Asp Asn Asp Gly Lys Lys Ile His Cys Ala Val Lys 1115 1120 1125
Ser Leu Asn Arg Ile Thr Asp Ile Gly Glu Val Ser Gln Phe Leu 1130 1135 1140
Thr Glu Gly Ile Ile Met Lys Asp Phe Ser His Pro Asn Val Leu 1145 1150 1155
Ser Leu Leu Gly Ile Cys Leu Arg Ser Glu Gly Ser Pro Leu Val 1160 1165 1170
Val Leu Pro Tyr Met Lys His Gly Asp Leu Arg Asn Phe Ile Arg 1175 1180 1185
Asn Glu Thr His Asn Pro Thr Val Lys Asp Leu Ile Gly Phe Gly 1190 1195 1200
Leu Gln Val Ala Lys Ala Met Lys Tyr Leu Ala Ser Lys Lys Phe 1205 1210 1215
Val His Arg Asp Leu Ala Ala Arg Asn Cys Met Leu Asp Glu Lys 1220 1225 1230
Phe Thr Val Lys Val Ala Asp Phe Gly Leu Ala Arg Asp Met Tyr 1235 1240 1245
Asp Lys Glu Tyr Tyr Ser Val His Asn Lys Thr Gly Ala Lys Leu 1250 1255 1260
Pro Val Lys Trp Met Ala Leu Glu Ser Leu Gln Thr Gln Lys Phe 1265 1270 1275
Thr Thr Lys Ser Asp Val Trp Ser Phe Gly Val Val Leu Trp Glu 1280 1285 1290
Leu Met Thr Arg Gly Ala Pro Pro Tyr Pro Asp Val Asn Thr Phe 1295 1300 1305
Asp Ile Thr Val Tyr Leu Leu Gln Gly Arg Arg Leu Leu Gln Pro 1310 1315 1320
Glu Tyr Cys Pro Asp Pro Leu Tyr Glu Val Met Leu Lys Cys Trp 1325 1330 1335
His Pro Lys Ala Glu Met Arg Pro Ser Phe Ser Glu Leu Val Ser 1340 1345 1350
Arg Ile Ser Ala Ile Phe Ser Thr Phe Ile Gly Glu His Tyr Val 1355 1360 1365
His Val Asn Ala Thr Tyr Val Asn Val Lys Cys Val Ala Pro Tyr 1370 1375 1380
Pro Ser Leu Leu Ser Ser Glu Asp Asn Ala Asp Asp Glu Val Asp 1385 1390 1395
Thr Arg Pro Ala Ser Phe Trp Glu Thr Ser 1400 1405

What is claimed is:

1. A method of monitoring the status of cancer in a patient comprising:
 - determining the concentration of soluble c-Met ectodomain in a first biological tissue from the patient at a first time point,
 - determining the concentration of soluble c-Met ectodomain in a second biological tissue from the patient at a second time point that is later than the first time point, wherein the first and second biological tissues are the same type of tissues, and
 - determining the status of cancer in the patient by comparing the concentration of soluble c-Met ectodomain in the first and second biological tissues
 - wherein a difference in the concentration of soluble c-Met ectodomain in the first and second biological tissues indicates that the patient's cancer has regressed or worsened; and
 - wherein similar concentrations of soluble c-Met ectodomain in the first and second biological tissues indicates that the patient's cancer has remained stable; and
 - wherein a lower concentration of soluble c-Met ectodomain in the second biological tissue than that in the first biological tissue indicates that the patient's cancer has regressed; and
 - wherein a greater concentration of soluble c-Met ectodomain in the second biological tissue than that in the first biological tissue indicates that the patient's cancer has worsened.
2. The method of claim 1 further comprising determining the concentration of soluble c-Met ectodomain in one or more additional biological tissues of the same type as the first and second biological tissues at time points that are progressively later than the second time point.
3. The method of claim 1 where in the cancer is metastatic cancer.
4. The method of claim 1 wherein the cancer is a cancer of the urinary tract.
5. The method of claim 4 wherein the cancer is kidney cancer, renal cancer, bladder cancer, or renal cell carcinoma.
6. The method of claim 5 wherein the cancer is bladder cancer.
7. The method of claim 1 wherein the cancer is prostate cancer.
8. The method of claim 1 wherein the cancer is breast cancer.

9. A method for measuring a patient's response to cancer therapy comprising:
 - determining the concentration of soluble c-Met ectodomain in a biological tissue from the patient at a first time point,
 - determining the concentration of soluble c-Met ectodomain in a second biological tissue from the patient at a second time point that is later than the first time point, wherein the first and second biological tissues are the same type of tissues, and
 - determining the patient's response to cancer therapy by comparing the concentration of soluble c-Met ectodomain in the first and second biological tissues
 - wherein a difference in the concentration of soluble c-Met ectodomain in the first and second biological tissues indicates that the patient's cancer has regressed or worsened; and
 - wherein similar concentrations of soluble c-Met ectodomain in the first and second biological tissues indicates that the patient has not responded to the cancer therapy; and
 - wherein a lower concentration of soluble c-Met ectodomain in the second biological tissue relative to that in the first biological tissue indicates that the patient has responded positively to the cancer therapy; and
 - wherein a greater concentration of soluble c-Met ectodomain in the second biological tissue than that in the first biological tissue indicates that the patient has not responded to the cancer therapy or has responded negatively to the cancer therapy.
10. The method of claim 9 further comprising determining the concentration of soluble c-Met ectodomain in one or more additional biological tissues of the same type as the first and second biological tissues at time points that are progressively later than the second time point.
11. The method of claim 9 wherein the cancer therapy is an anti-cancer drug.
12. The method of claim 11 wherein the drug is a chemotherapeutic agent.
13. The method of claim 9 wherein the cancer therapy is ionizing radiation therapy or hormone ablation therapy.
14. The method of claim 9 where in the cancer is metastatic cancer.
15. The method of claim 9 wherein the cancer is a cancer of the urinary tract.
16. The method of claim 15 wherein the cancer is kidney cancer, renal cancer, bladder cancer, or renal cell carcinoma.

* * * * *

专利名称(译)	通过测量可溶性c-Met胞外域来诊断和监测癌症进展的方法		
公开(公告)号	US8617831	公开(公告)日	2013-12-31
申请号	US13/627595	申请日	2012-09-26
[标]申请(专利权)人(译)	Bottaro的DONALD P ATHAUDA GAGANI P BURS TERESA LYNN		
申请(专利权)人(译)	Bottaro的, DONALD P. ATHAUDA, GAGANI P. 伯吉斯TERESA LYNN		
当前申请(专利权)人(译)	AMGEN, INC.		
[标]发明人	BOTTARO DONALD P ATHAUDA GAGANI P BURGESS TERESA LYNN		
发明人	BOTTARO, DONALD P. ATHAUDA, GAGANI P. BURGESS, TERESA LYNN		
IPC分类号	G01N33/53		
CPC分类号	G01N2800/52 G01N33/57488		
代理机构(译)	伍德科克WASHBURN LLP		
审查员(译)	沉, BIN		
优先权	13/053560 2012-11-06 US 60/780626 2006-03-09 US 60/734993 2005-11-08 US PCT/US2006/043654 2006-11-08 WO		
其他公开文献	US20130029352A1		
外部链接	Espacenet USPTO		

摘要(译)

提供了测量尿液和血液样品中c-Met水平的方法。还提供了用于癌症的诊断和预后评估的方法。




Figure 1