



US 20030100525A1

(19) **United States**

(12) **Patent Application Publication**  
**Altieri**

(10) **Pub. No.: US 2003/0100525 A1**

(43) **Pub. Date: May 29, 2003**

(54) **SURVIVIN, A PROTEIN THAT INHIBITS  
CELLULAR APOPTOSIS, AND ITS  
MODULATION**

(60) Provisional application No. 60/031,435, filed on Nov.  
20, 1996.

**Publication Classification**

(75) Inventor: **Dario C. Altieri**, Hamden, CT (US)

(51) **Int. Cl.<sup>7</sup>** ..... **A61K 48/00**; A61K 38/17;  
C12N 15/85

Correspondence Address:

**MORGAN LEWIS & BOCKIUS LLP**  
**1111 PENNSYLVANIA AVENUE NW**  
**WASHINGTON, DC 20004 (US)**

(52) **U.S. Cl.** ..... **514/44**; 514/12; 435/455;  
435/320.1

(73) Assignee: **Yale University**

(57) **ABSTRACT**

(21) Appl. No.: **10/138,618**

(22) Filed: **May 6, 2002**

**Related U.S. Application Data**

(62) Division of application No. 09/690,825, filed on Oct.  
18, 2000, which is a division of application No.  
08/975,080, filed on Nov. 20, 1997, now Pat. No.  
6,245,523.

The present invention provides the amino acid of a protein that inhibits cellular apoptosis, herein termed the Survivin protein and nucleic acid molecules that encode Survivin. Based on this disclosure, the present invention provides isolated Survivin protein, isolated Survivin encoding nucleic acid molecules, methods of isolating other members of the Survivin family of proteins, methods for identifying agents that block Survivin mediated inhibition of cellular apoptosis, methods of using agents that block Survivin mediated inhibition or Survivin expression to modulate biological and pathological processes, and methods of assaying Survivin activity.




FIG. 1A




FIG. 1B







FIG. 1C

5' Exon		Intron	3' Exon
(2918)GCGG	GTGAG	-----	(3161)CTGTCCCTTGCAG ATGGC
(3280)CCAT	GTAAG	-----	(5145)TTGATTTTCTAG AGAGG
(5272)AATT	GTATG	-----	(11942)TCTTTATTCCAG GCAAA

FIG. 1D






FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E




FIG. 3F

FIG. 4A


10 20 30 40  
 MGAPTLPPAW CPFLKDHRS TFKNWPFLG CACTPERMAE 40  
 AGFIHCPTEN EPDLAOCFFC FKELEGWEPD DDPIEEHKKH 80  
 SSGCAFLSVK KOFEEITLGE FLKLDREKAK NKIAKETNNK 120  
 KKEFEETAKK VRRALAEOLAA MD 142

FIG. 4B-1


177	EEARLVTFQNWPD-AFL----	TPQELAKAGFYLLGRGDQVQCFA	CGGKLA	Majority	
113	EEARFLTYSMWP-LSFL----	SPAELARAGFYLLGRGDQVQCFA	CGGKLS		L49433.PRO
7	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		L49441.PRO
18	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		P41436.PRO
159	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		P41437.PRO
113	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		U19251.PRO
163	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		U32373.PRO
163	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		U32974.PRO
169	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		U36842.PRO
184	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		U45878.PRO
163	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		U45879.PRO
113	EEARLVTFKDWPN-PNI----	TPQALAKAGFYLLGRGDQVQCFA	CGGKLS		U45880.PRO
15	KDHRISIEKNNWPF-LEG-CACTPERMAEAGFI-----HC				U45881.PRO
					SURVIVIN.PRO

FIG. 4B-2

	360	370	380	390	400	Majority				
YVGIGDKVKCFHCDGGLRDWEPGDDPWEEHAKWFPRCEFLLLAKGQEYVS										
2290	YVDRND	DYKCF	CCDCGLRC	WEPGDD	PWIEHAKWF	PRCEFLIRM	GQEF	VMD	L49433	.PRO
2240	YQKIGD	QVRCF	HCNIGLRS	WQKED	EPWEEHAKWS	PKCQEVLLAK	GPAY	VS	L49441	.PRO
21136	YTGYGD	NTKCF	YCCDGLKD	WEPE	DVPWEQHVR	WEDRCAYVQ	LVKGRDY	M-	P41436	.PRO
21139	YTGCGD	KTRCF	CCDGLKD	WEPE	DAPWOOHARWY	DRCEYVLLV	KGRDEY	-	P41437	.PRO
2306	YTGJTD	LVQCF	SCGGCLEK	WOEGDD	PLDDHT	RCFPCFLLONM	KSSAEV	VT	U19251	.PRO
2240	YQKIGD	QVRCF	HCNIGLRS	WQKED	EPWEEHAKWS	PKCQEVLLAK	GPSY	VS	U32373	.PRO
2291	ALGEGD	KVKCF	HCCGG	LTDW	KPSED	PWEQHAKWY	PGCKYLL	LEQR	U32974	.PRO
2290	ALGEGD	KVKCF	HCCGG	LTDW	KPSED	PWEQHAKWY	PGCKYLL	LEQR	U36842	.PRO
2283	YVGN	SDVKCF	CCDGLRC	WESGDD	PWVQHAKWF	PRCEYL	IRIK	GQEF	U45878	.PRO
2297	YVGRND	DVKCF	CCDGLRC	WESGDD	PWEHAKWF	PRCEFL	IRM	KGQEF	U45879	.PRO
2291	ALGEGD	KVKCF	HCCGG	LTDW	KPSED	PWEQHAKWY	PGCKYLL	LEQR	U45880	.PRO
2240	YQKIGD	QVRCF	HCNIGLRS	WQKED	EPWEEHAKWS	PKCQFVLLAK	GPAY	VS	U45881	.PRO
53	----	DLAQCF	CFKEL	EGWEP	DDDEPI	EEHKKHSSG	CAFLS	----	SURVIVIN	.PRO



**FIG. 4C-1**



**FIG. 4C-2**






FIG. 5A

FIG. 5B

FIG. 5C




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 6E




FIG. 6F



FIG. 6G



FIG. 6H



**FIG. 7A**




FIG. 7B-1




FIG. 7B-2




FIG. 7B-3





FIG. 7B-4

**FIG. 7C**






FIG. 8A

FIG. 8B

FIG. 8C





FIG. 8D

FIG. 9



## FIG. 10A

translation=MGAPTLPPAWQPFLKDHRISTFKNWPFLGCACTPERMAEAGFIHCP  
 TENEPDLAQCFPCFKELEGWEPDDDDPIEEHKKHSSGCAFLSVKKQFEELTLGEFL  
 KLDREKAKNKIAKETNNKKKEFEETAKKVRRAIEQLAAMD

```

1      TCTAGACATG CGGATATATT CAAGCTGGGC ACAGCACAGC AGCCCCACCC
51     CAGGCAGCTT GAAATCAGAG CTGGGGTCCA AAGGGACCAC ACCCCGAGGG
101    ACTGTGTGGG GGTCGGGGCA CACAGGCCAC TGCTTCCCCC CGTCTTTCTC
151    AGCCATTCTT GAAGTCAGCC TCACTCTGCT TCTCAGGGAT TTCAAATGTG
201    CAGAGACTCT GGCACCTTTG TAGAAGCCCC TTCTGGTCCT AACTTACACC
251    TGGATGCTGT GGGGCTGCAG CTGCTGCTCG GGCTCGGGAG GATGCTGGGG
301    GCCCCGTGCC CATGAGCTTT TGAAGCTCCT GGAAGTCGGT TTTGAGGGTG
351    TTCAGGTCCA GGTGGACACC TGGGCTGTCC TTGTCCATGC ATTTGATGAC
401    ATTGTGTGCA GAAGTGAAAA GGAGTTAGGC CGGGCATGCT GGCTTATGCC
451    TGTAATCCCA GCACTTTGGG AGGCTGAGGC GGGTGGATCA CGAGGTCAGG
501    AGTTCAATAC CAGCCTGGCC AAGATGGTGA AACCCGCTCT CTAATAAAAA
551    TACAAAAAAA TTAGCCGGGC ATGGTGGCGG GCGCATGTAA TCCCAGCTAC
601    TGGGGGGGGT GAGGCAGAGA ATTGCTGGAA CCCAGGAGAT GGAGGTTGCA
651    GTGAGCCAAG ATTGTGCCAC TGCACGAC TCCAGCCTGG CGACAGAGCA
701    AGACTCTGTC TCAAAAAAAA AAAAAAAAAG TGAAAAGGAG TTGTTCCCTTT
751    CCTCCCTCCT GAGGGCAGGC AACTGCTGCG GTTGCCAGTG GAGGTGGTGC
801    GTCCTTGGTC TGTGCCTGGG GGCCACCCCA GCAGAGGCCA TGGTGGTGCC
851    AGGGCCCGGT TAGCGAGCCA ATCAGCAGGA CCCAGGGGCG ACCTGCCAAA
901    GTCAACTGGA TTTGATAACT GCAGCGAAGT TAAGTTTCCT GATTTTGATG
951    ATTGTGTTGT GGTGTGTAA GAGAATGAAG TATTTCCGGG TAGTATGGTA
1001   ATGCCTTCAA CTTACAAACG GTTCAGGTAA ACCACCCATA TACATACATA
1051   TACATGCATG TGATATATAC ACATACAGGG ATGTGTGTGT GTTCACATAT
1101   ATGAGGGGAG AGAGACTAGG GGAGAGAAAG TAGGTTGGGG AGAGGGAGAG
1151   AGAAAGGAAA ACAGGAGACA GAGAGAGAGC GGGGAGTAGA GAGAGGGAAG
1201   GGGTAAGAGA GGGAGAGGAG GAGAGAAAGG GAGGAAGAAG CAGAGAGTGA
1251   ATGTTAAAGG AAACAGGCAA AACATAACA GAAAATCTGG GTGAAGGGTA
1301   TATGAGTATT CTTTGTACTA TTCTTGCAAT TATCTTTTAT TTAAATTGAC
1351   ATCGGGCCGG GCGCAGTGGC TCACATCTGT AATCCCAGCA CTTTGGGAGG
1401   CCGAGGCAGG CAGATCACTT GAGGTCAGGA GTTTGAGACC AGCCTGGCAA
1451   ACATGGTGAA ACCCATCTC TACTAAAAAT AAAAAATTA GCCTGGTGTG
1501   GTGGTGCAAT CCTTTAATCT CAGCTACTCG GGAGGCTGAG GCAGGAGAAT
1551   CGCTTGAACC CGTGGCGGGG AGGAGGTTGC AGTGAGCTGA GATCATGCCA
1601   CTGCACTCCA GCCTGGGCGA TAGAGCGAGA CTCAGTTTCA AATAAATAAA
1651   TAAACATCAA AATAAAAAGT TACTGTATTA AAGAATGGGG GCGGGGTGGG
1701   AGGGGTGGGG AGAGGTTGCA AAAATAAATA AATAAATAAA TAAACCCCAA
1751   AATGAAAAAG ACAGTGGAGG CACCAGGCCT GCGTGGGGCT GGAGGGCTAA
1801   TAAGGCCAGG CCTCTTATCT CTGGCCATAG AACCAGAGAA GTGAGTGGAT
1851   GTGATGCCCC GCTCCAGAAG TGACTCCAGA ACACCCTGTT CCAAAGCAGA
1901   GGACACACTG ATTTTTTTTT TAATAGGCTG CAGGACTTAC TGTGCTGGG
1951   ACGCCCTGCT TTGCGAAGGG AAAGGAGGAG TTTGCCCTGA GCACAGGCC
2001   CCACCCCTCA CTGGGCTTTC CCCAGCTCCC TTGTCTTCTT ATCACGGTAG
2051   TGGCCAGTC CCTGGCCCTT GACTCCAGAA GGTGGCCCTC CTGGAAACCC
2101   AGGTCGTGCA GTCAACGATG TACTCGCCGG GACAGCGATG TCTGCTGCAC
2151   TCCATCCCTC CCCTGTTTCAT TTGTCCTTCA TGCCCGTCTG GAGTAGATGC

```



## FIG. 10B

```

2201 TTTTTCAGAG GGTGGCACCC TGTAAGCTC TCCTGTCTGA CTTTTTTTTT
2251 TTTTTCAGAG TGAGTTTTCG TCTTGTGTC TAGGCTGGAG TGCAATGGCA
2301 CAATCTCAGC TCACTGCACC CTCTGCCTCC CGGGTTCAAG CGATTCTCCT
2351 GCCTCAGCCT CCCGAGTAGT TGGGATTACA GGCATGCACC ACCACGCCCC
2401 GCTAATTTTT GTATTTTTAG TAGAGACAAG GTTTCACCGT GATGGCCAGG
2451 CTGGTCTTGA ACTCCAGGAC TCAAGTGATG CTCCTGCCTA GGCCTCTCAA
2501 AGTGTGGGA TTACAGGCGT GAGCCACTGC ACCCGGCCG CACGCGTTCT
2551 TTGAAAGCAG TCGAGGGGGC GCTAGGTGTG GGCAGGGACG AGCTGGCGCG
2601 GCGTCGCTGG GTGCACCGCG ACCACGGGCA GAGCCACGCG GCGGGAGGAC
2651 TACAACCTCC GGCACACCCC GCGCCGCCCC GCCTCTACTC CCAGAAGGCC
2701 GCGGGGGGTG GACCGCCTAA GAGGGCGTGC GCTCCCGACA TGCCCCGCGG
2751 CCGGCCATTA ACCGCCAGAT TTGAATCGCG GGACCCGTTG GCAGAGGTGG
      |----->Start
2801 CCGCGGCGGC ATGGGTGCCC CGACGTTGCC CCCTGCCTGG CAGCCCTTTC
2851 TCAAGGACCA CCGCATCTCT ACATTCAAGA ACTGGCCCTT CTGGAGGGGC
2901 TGCGCCTGCA CCCCGGAGCG GGTGAGACTG CCCGGCCTCC TGGGGTCCCC
2951 CACGCCCCGC TTGCCCTGTC CCTAGCGAGG CCACTGTGAC TGGGCCTCGG
3001 GGGTACAAGC CGCCTCCCC TCCCCGTCT GTCCCCAGCG AGGCCACTGT
3051 GGCTGGGCCC CTGGGTCCA GGCCGGCCTC CCCTCCCTGC TTTGTCCCCA
3101 TCGAGGCTT TGTGGCTGGG CCTCGGGGT CCGGGCTGCC ACGTCCACTC
3151 ACGAGCTGTG CTGTCCCTTG CAGATGGCCG AGGCTGGCTT CATCCACTGC
3201 CCCACTGAGA ACGAGCCAGA CTTGGCCCAG TGTTTCTTCT GCTTCAAGGA
3251 GCTGGAAGGC TGGGAGCCAG ATGACGACCC CATGTAAGTC TTCTCTGGCC
3301 AGCCTCGATG GGCTTTGTTT TGAAGTGAAG TGTCAAAAGA TTTGAGTTGC
3351 AAAGACACTT AGTATGGGAG GGTGCTTTC CACCCTCATT GCTTCTTAAA
3401 CAGCTGTTGT GAACGGATAC CTCTCTATAT GCTGGTGCCT TGGTGATGCT
3451 TACAACCTAA TTAAATCTCA TTTGACCAA ATGCCTTGGG GTGGACGTAA
3501 GATGCCTGAT GCCTTTCATG TTCAACAGAA TACATCAGCA GACCCTGTTG
3551 TTGTGAACTC CCAGGAATGT CCAAGTGCTT TTTTGTAGAT TTTTAAAAA
3601 ACAGTTTAAT TGAAATATAA CCTACACAGC AAAAAATTA CCCTTTGAAA
3651 GTGTGCACTT CACACTTTCG GAGGCTGAGG CGGGCGGATC ACCTGAGGTC
3701 AGGAGTTCAA GACCTGCCTG GCCAACTTGG CGAAACCCCG TCTCTACTAA
3751 AAATACAAAA ATTAGCCGGG CATGGTAGCG CACGCCCGTA ATCCAGCTA
3801 CTCGGGAGGC TAAGGCAGGA GAATCGCTTG AACCTGGGAG GCGGAGGTTG
3851 CAGTGAGCCG AGATTGTGCC AATGCACTCC AGCCTCGGCG ACAGAGCGAG
3901 ACTCCGTCAT AAAAATAAAA AATTGAAAA AAAAAAGAA AGAAAGCATA
3951 TACTTCAGTG TTGTTCTGGA TTTTCTTCT CAAGATGCCT AGTTAATGAC
4001 AATGAAATTC TGTACTCGGA TGGTATCTGT CTTTCCACAC TGTAATGCCA
4051 TATTCTTTTC TCACCTTTTT TTCTGTGCGA TTCAGTTGCT TCCACAGCTT
4101 TAATTTTTTT CCCCTGGAGA TCTTAGTATG TTTGCTATGG TGGTTATACT
4151 GCATCCCCGT AATCACTGGG AAAAGATCAG TGGTATTCTT CTTGAAAATG
4201 AATAAGTGTT ATGATATTTT CAGATTAGAG TTCAACTGG CTGTCTTTTT
4251 GGACTTTGTG TGGCCATGTT TCATTGTAA TGCAGTCTG GTAACGGTGA
4301 TAGTCAGTTA TACAGGGAGA CTCCCCTAGC AGAAAATGAG AGTGTGAGCT
4351 AGGGGGTCCC TTGGGGAACC CGGGGCAATA ATGCCCTTCT CTGCCCTTAA
4401 TCCTTACAGT GGGCCGGGCA CGGTGGCTTA CGCCTGTAAT ACCAGCACTT
4451 TGGGAGGCGG AGGCGGGCGG ATCAGGAGGT CAGGAGATCG AGACCATCTT
4501 GGCTAATACG GTGAAACCCC GTCTCCACTA AAAATACAAA AAATTAGCCG

```

## FIG. 10C

4551	GGCGTGGTGG	TGGGCGCCTG	TAGTCCCAGC	TACTCGGGAG	GCTGAGGCAG
4601	GAGAATGGCG	TGAACCCAGG	AGGCGGAGCT	TGCAGTGAGC	CGAGATTGCA
4651	CCACTGCACT	CCAGCCTGGG	CGACAGAATG	AGACTCCGTC	TCAAAAAAAAA
4701	AAAAAAAAAGA	AAAAAATCTT	TACAGTGGAT	TACATAACAA	TTCCAGTGAA
4751	ATGAAATTAC	TTCAAACAGT	TCCTTGAGAA	TGTTGGAGGG	ATTTGACATG
4801	TAATTCCTTT	GGACATATAC	CATGTAACAC	TTTTCCAAC	AATTGCTAAG
4851	GAAGTCCAGA	TAAAATAGAT	ACATTAGCCA	CACAGATGTG	GGGGGAGATG
4901	TCCACAGGGA	GAGAGAAGGT	GCTAAGAGGT	GCCATATGGG	AATGTGGCTT
4951	GGGCAAAGCA	CTGATGCCAT	CAACTTCAGA	CTTGACGTCT	TACTCCTGAG
5001	GCAGAGCAGG	GTGTGCCCTG	GGAGGGCGTG	GGGAGGTGGC	CCGTGGGGAG
5051	TGGACTGCCG	CTTTAATCCC	TTCAGCTGCC	TTTCCGCTGT	TGTTTTGATT
5101	TTTCTAGAGA	GGAACATAAA	AAGCATTCGT	CCGGTTGCGC	TTTCCTTTCT
5151	GTCAAGAAGC	AGTTTGAAGA	ATTAACCCTT	GGTGAATTTT	TGAAACTGGA
5201	CAGAGAAAGA	GCCAAGAACA	AAATTGTATG	TATTGGGAAT	AAGAACTGCT
5251	CAAAACCCTGT	TCAATGTCTT	TAGCACTAAA	CTACCTAGTC	CCTCAAAGGG
5301	ACTCTGTGTT	TTCCTCAGGA	AGCATTTTTT	TTTTTTTTTCT	GAGATAGAGT
5351	TTCACTCTTG	TTGCCCAGGC	TGGAGTGCAA	TGGTGCAATC	TTGGCTCACT
5401	GCAACCTCTG	CCTCTCGGGT	TCAAGTGATT	CTCCTGCCTC	AGCCTCCCAA
5451	GTAACCTGGGA	TTACAGGGAA	GTGCCACCAC	ACCCAGCTAA	TTTTTGTIATT
5501	TTTAGTAGAG	ATGGGGTTTC	ACCACATTGC	CCAGGCTGGT	CTTGAACCTCC
5551	TGACCTCGTG	ATTCGCCCAC	CTTGGCCTCC	CAAAGTGCTG	GGATTACAGG
5601	CGTGAACCAC	CACGCCCTGG	TTTTTTTTTT	TTGTTCTGAG	ACACAGTTTC
5651	ACTCTGTTAC	CCAGGCTGGA	GTAGGGTGGC	CTGATCTCGG	ATCACTGCAA
5701	CCTCCGCCCTC	CTGGGCTCAA	GTGATTTGCC	TGCTTCAGCC	TCCCAAGTAG
5751	CCGAGATTAC	AGGCATGTGC	CACCACACCC	AGGTAATTTT	TGTATTTTTG
5801	GTAGAGACGA	GGTTTCACCA	TGTTGGCCAG	GCTGGTTTTG	AACTCCTGAC
5851	CTCAGGTGAT	CCACCCGCCT	CAGCCTCCCA	AAGTGCTGAG	ATTATAGGTG
5901	TGAGCCACCA	CACCTGGCCT	CAGGAAGTAT	TTTTATTTTT	AAATTTATTT
5951	ATTTATTTGA	GATGGAGTCT	TGCTCTGTCT	CCCAGGCTAG	AGTGCAGCGA
6001	CGGGATCTCG	GCTCACTGCA	AGCTCCGCCC	CCCAGGTTCA	AGCCATTCTC
6051	CTGCCTCAGC	CTCCCGAGTA	GCTGGGACTA	CAGGCGCCCG	CCACCACACC
6101	CGGCTAATTT	TTTTGTATTT	TTAGTAGAGA	CGGGTTTTCA	CCGTGTTAGC
6151	CAGGAGGGTC	TTGATCTCCT	GACCTCGTGA	TCTGCCTGCC	TCGGCCTCCC
6201	AAAGTGCTGG	GATTACAGGT	GTGAGCCACC	ACACCCGGCT	ATTTTTATTT
6251	TTTTGAGACA	GGGACTCACT	CTGTCACCTG	GGCTGCAGTG	CAGTGGTACA
6301	CCATAGCTCA	CTGCAGCCTC	GAATCCTGA	GCTCAAGTGA	TCCTCCCACC
6351	TCATCCTCAC	AAGTAATTGG	GACTACAGGT	GCACCCACC	ATGCCACCTT
6401	AATTTATTTA	TTTATTTATT	TATTTATTTT	CATAGAGATG	AGGGTTCCCT
6451	GTGTTGTCCA	GGCTGGTCTT	GAATCCTGA	GCTCACGGGA	TCCTTTTGCC
6501	TGGGCCTCCC	AAAGTGCTGA	GATTACAGGC	ATGAGCCACC	GTGCCCAGCT
6551	AGGAATCATT	TTTAAAGCCC	CTAGGATGTC	TGTGTGATTT	TAAAGCTCCT
6601	GGAGTGTTGG	CGGTATAAGT	ATATACCGGT	ATAAGTAAAT	CCCACATTTT
6651	CTGGGCTTTA	TTTATTTATT	TATTTATTTA	TTTATTTTTA	ATTTTTTTTT
6701	TTGAGACGAG	TCTCACTTG	TCACCCAGGC	TGGAGTGCAG	TGGCACGATC
6751	TCGGCTCACT	GCAACCTCTG	CCTCCCGGGG	TCAAGCGATT	CTCCTGCCTT
6801	AGCCTCCCGA	GTAGCTGGGA	CTACAGGCAC	GCACCACCAT	GCCTGGCTAA
6851	TTTTTGATTT	TTTAGTAGAC	GGGGTTTCAC	CATGCTGGCC	AAGCTGGTCT
6901	CAAACCTCTG	ACCTTGTTGAT	CTGCCCCGCT	TAGCCTCCCA	GAGTGCTGGG

## FIG. 10D

6951	ATTACAGGCA	TGAGCCACCA	TGCGTGGTCT	TTTTAAAATT	TTTTGATTTT
7001	TTTTTTTTTT	GAGACAGAGC	CTTGCTCTGT	CGCCCAGGCT	GGAGTGCAGT
7051	GGCACGATCT	CAGCTCACTA	CAAGCTCCGC	CTCCCGGGTT	CACGCCATTC
7101	TTCTGCCTCA	GCCTCCTGAG	TAGCTGGGAC	TACAGGTGCC	CACCACCACG
7151	CCTGGCTAAT	TTTTTTTGGT	ATTTTTATTA	GAGACAAGGT	TTCATCATGT
7201	TGGCCAGGCT	GGTCTCAAAC	TCCTGACCTC	AAGTGATCTG	CCTGCCTCGG
7251	CCTCCCAAAG	CGCTGAGATT	ACAGGTGTGA	TCTACTGCGC	CAGGCCTGGG
7301	CGTCATATAT	TCTTATTTGC	TAAGTCTGGC	AGCCCCACAC	AGAATAAGTA
7351	CTGGGGGATT	CCATATCCTT	GTAGCAAAGC	CCTGGGTGGA	GAGTCAGGAG
7401	ATGTTGTAGT	TCTGTCTCTG	CCACTTGCAG	ACTTTGAGTT	TAAGCCAGTC
7451	GTGCTCATGC	TTTCCTTGCT	AAATAGAGGT	TAGACCCCTT	ATCCCATGGT
7501	TTCTCAGGTT	GCTTTTCAGC	TTGAAAATTG	TATTCCTTTG	TAGAGATCAG
7551	CGTAAAATAA	TTCTGTCCTT	ATATGTGGCT	TTATTTTAAT	TTGAGACAGA
7601	GTGTCACTCA	GTCGCCCAGG	CTGGAGTGTG	GTGGTGCGAT	CTTGGCTCAC
7651	TGCGACCTCC	ACCTCCCAGG	TTCAAGCGAT	TCTCGTGCCT	CAGGCTCCCA
7701	AGTAGCTGAG	ATTATAGGTG	TGTGCCACCA	GGCCCAGCTA	ACTTTTGTAT
7751	TTTTAGTAGA	GACAGGGTTT	TGCCATGTTG	GCTAAGCTGG	TCTCGAACTC
7801	CTGGCCTCAA	GTGATCTGCC	CGCCTTGGCA	TCCCAAAGTG	CTGGGATTAC
7851	AGGTGTGAAC	CACCACACCT	GGCCTCAATA	TAGTGGCTTT	TAAGTGCTAA
7901	GGACTGAGAT	TGTGTTTTGT	CAGGAAGAGG	CCAGTTGTGG	GTGAAGCATG
7951	CTGTGAGAGA	GCTTGTCAAC	TGGTTGAGGT	TGTGGGAGCT	GCAGCGTGGG
8001	AACTGGAAAG	TGGGCTGGGG	ATCATCTTTT	TCCAGGTCAG	GGGTCAGCCA
8051	GCTTTTCTGC	AGCGTGCCAT	AGACCATCTC	TTAGCCCTCG	TGGGTCAGAG
8101	TCTCTGTTGC	ATATTGTCTT	TTGTTGTTTT	TCACAACTTT	TTAGAAACAT
8151	AAAAAGCATT	CTTAGCCCGT	GGGCTGGACA	AAAAAAGGCC	ATGACGGGCT
8201	GTATGGATTT	GGCCCAGCAG	GCCCTTGCTT	GCCAAAGCCCT	GTTTTAGACA
8251	AGGAGCAGCT	TGTGTGCCTG	GAACCATCAT	GGGCACAGGG	GAGGAGCAGA
8301	GTGGATGTGG	AGGTGTGAGC	TGGAAACCAG	GTCCCAGAGC	GCTGAGAAAG
8351	ACAGAGGGTT	TTTGCCCTTG	CAAGTAGAGC	AACTGAAATC	TGACACCATC
8401	CAGTTCCAGA	AAGCCCTGAA	GTGCTGGTGG	ACGCTGCGGG	GTGCTCCGCT
8451	CTAGGGTTAC	AGGGATGAAG	ATGCAGTCTG	GTAGGGGGAG	TCCACTCACC
8501	TGTTGGAAGA	TGTGATTAAG	AAAAGTAGAC	TTTCAGGGCC	GGGCATGGTG
8551	GCTCACGCCT	GTAATCCCAG	CACTTTGGA	GGCCGAGCGG	GGTGGATCAC
8601	GAGGTCAGGA	GATCGAGACC	ATCCTGGCTA	ACATGGTGAA	ACCCCGTCTT
8651	TACTAAAAAT	ACAAAAAATT	AGCTGGGCGT	GCTGGCGGGC	GCCTGTAGTC
8701	CCAGCTACTC	GGGAGGCTGA	GGCAGGAGAA	TGGCGTGAAC	CTGGGAGGTG
8751	GAGCTTGCTG	TGAGCCGAGA	TCGCGCCACT	GCACTCCAGC	CTGGGCGACA
8801	GAGCGAGACT	CCGTCTCAA	AAAAAAAAAA	AAAGTAGGCT	TTCATGATGT
8851	GTGAGCTGAA	GGCGCAGTAG	GCAGAAGTAG	AGGCCTCAGT	CCCTGCAGGA
8901	GACCCCTCGG	TCTCTATCTC	CTGATAGTCA	GACCCAGCCA	CACCTGGAAAG
8951	AGGGGAGACA	TTACAGCCTG	CGAGAAAAGT	AGGGAGATTT	AAAAACTGCT
9001	TGGCTTTTAT	TTTGAACGT	TTTTTTTGTT	TGTTTGTTTT	CCCCAATTCA
9051	GAATACAGAA	TACTTTTATG	GATTTGTTTT	TATTACTTTA	ATTTTGAAAC
9101	AATATAATCT	TTTTTTTGTT	GTTTTTTTGA	GACAGGGTCT	TACTCTGTCA
9151	CCCAGGCTGA	GTGCAGTGGT	GTGATCTTGG	CTCACCTCAG	CCTCGACCCC
9201	CTGGGCTCAA	ATGATTCTCC	CACCTCAGCT	TCCCAAGTAG	CTGGGACCAC
9251	AGGTGCGTGT	GTTGCGCTAT	ACAAATCCTG	AAGACAAGGA	TGCTGTTGCT
9301	GGTGATGCTG	GGGATTCCCA	AGATCCCAGA	TTTGATGGCA	GGATGCCCTT

# FIG. 10E

9351	GTCTGCTGCC	TTGCCAGGGT	GCCAGGAGGG	CGCTGCTGTG	GAAGCTGAGG
9401	CCCGGCCATC	CAGGGCGATG	CATTGGGCGC	TGATTCTTGT	TCCTGCTGCT
9451	GCCTCGGTGC	TTAGCTTTTG	AAACAATGAA	ATAAATTAGA	ACCAGTGTGA
9501	AAATCGATCA	GGGAATAAAT	TTAATGTGGA	AATAAACTGA	ACAACTTAGT
9551	TCTTCATAAG	AGTTTACTTG	GTAAATACTT	GTGATGAGGA	CAAAACGAAG
9601	CACTAGAAGG	AGAGGCGAGT	TGTAGACCTG	GGTGGCAGGA	GTGTTTTGTT
9651	TGTTTTCTTT	GGCAGGGTCT	TGCTCTGTTG	CTCAGGCTGG	AGTACAGTGG
9701	CACAAACACA	GCTCACTATA	GCCTCGACCT	CCTGGACTCA	AGCAATCCTC
9751	CTGCCTCAGC	CTCCCAGTAG	CTGGGACTAC	AGGCGCATGC	CACCATGCCT
9801	GGCTAATTTT	AAATTTTTTT	TTTTCTCTTT	TTTGAGATGG	AATCTCACTC
9851	TGTCGCCCAG	GCTGGAGTGC	AGTGGCGTGA	TCTCGGCTGA	CGGCAAGCTC
9901	CGCCTCCCAG	GTTCACTCCA	TTCGCCTGCC	TCAGCCTCCC	AAGTAGCTGG
9951	GACTACAGGC	GCTGGGATTA	CAAACCCAAA	CCCAAAGTGC	TGGGATTACA
10001	GGCGTGAGCC	ACTGCACCCG	GCCTGTTTTG	TCTTTCAATA	GCAAGAGTTG
10151	TGTTTGCTTC	GCCCCACCT	TTAGTGGAAG	AATGTATAAA	ATGGAGATAT
10201	TGACCTCCAC	ATTGGGGTGG	TTAAATTATA	GCATGTATGC	AAAGGAGCTT
10251	CGCTAATTTA	AGGCTTTTTT	GAAAGAGAAG	AAACTGAATA	ATCCATGTGT
10301	GTATATATAT	TTTAAAAGCC	ATGGTCATCT	TTCCATATCA	GTAAAGCTGA
10351	GGCTCCCCTG	GACTGCAGAG	TTGTCCATCA	CAGTCCATTA	TAAGTGCCTG
10401	GCTGGGGCCG	GTGCAGTGGC	TTGTGCCTGA	ATCCCAGCAC	TTTGGGAGGC
10451	CAAGGCAGGA	GGATTCAATG	AGCCCAGGAG	TTTTGAGGCG	AGCCTGGGCA
10501	ATGTGGCCAG	ACCTCATCTC	TTCAAAAAAT	ACACAAAAAA	TTAGCCAGGC
10551	ATGGTGGCAC	GTGCCTGTAG	TCTCAGCTAC	TCAGGAGGCT	GAGGTGGGAG
10601	GATCACTTTG	AGCCTTGCGG	GTCAAAGCTG	CAGTAAGCCA	TGATCTTGCC
10651	ACTGCATTCC	AGCCTGGATG	ACAGAGCGAG	ACCCTGTCTC	TAAAAAATAA
10701	AAAAACCAAA	CGGTGCACTG	TTTTCTTTTT	TCTTATCAAT	TTATTAATTT
10751	TAAATTAAAT	TTTCTTTTAA	TAATTTATAA	ATTATAAAAT	TATATTAAAA
10801	AATGACAAAT	TTTTATTACT	TATACATGAG	GTAAAACTTA	GGATATATAA
10851	AGTACATATT	GAAAAGTAAT	TTTTTGGCTG	GCACAGTGGC	TCACACCTGT
10901	AATCCCAGCA	CTTTGGGAGG	CCGTGGCGGG	CAGATCACAT	GAGATCATGA
10951	GTTCGAGACC	AACCTGACCA	ACATGGAGAG	ACCCCATCTC	TACTAAAAAT
11001	ACAAAATTAG	CCGGGGTGGT	GGCGCATGCC	TGTAATCCCA	GCTACTCGGG
11051	AGGCTGAGGC	AGGAGAATCT	CTTGAACCCG	CGAGGCAGAG	GTTGCGGTGA
11101	GCCAAGATCG	TGCCTTTGCA	CACCAGCCTA	GGCAACAAGA	GCGAAAGTCC
11151	GTCTCAAAAA	AAAAGTAATT	TTTTTTAAGT	TAACCTCTGT	CAGCAAACAA
11201	ATTTAACCCT	ATAAAGGTCT	TTGTTTTTTA	ATGTAGTAGA	GGAGTTAGGG
11251	TTTATAAAAA	ATATGGTAGG	GAAGGGGGTC	CCTGGATTTG	CTAATGTGAT
11301	TGTCATTTGC	CCCTTAGGAG	AGAGCTCTGT	TAGCAGAATG	AAAAAATTGG
11351	AAGCCAGATT	CAGGGAGGGA	CTGGAAGCAA	AAGAAATTTCT	GTTCGAGGAA
11401	GAGCCTGATG	TTTGCCAGGG	TCTGTTTAAC	TGGACATGAA	GAGGAAGGCT
11451	CTGGACTTTC	CTCCAGGAGT	TTCAGGAGAA	AGGTAGGGCA	GTGGTTAAGA
11501	GCAGAGCTCT	GCCTAGACTA	GCTGGGGTGC	CTAGACTAGC	TGGGGTGCCC
11551	AGACTAGCTG	GGGTGCCTAG	ACTAGCTGGG	TACTTTGAGT	GGCTCCTTCA
11601	GCCTGGACCT	CGGTTTCCTC	ACCTGTATAG	TAGAGATATG	GGAGCACCCA
11651	GCGCAGGATC	ACTGTGAACA	TAAATCAGTT	AATGGAGGAA	GCAGGTAGAG
11701	TGGTGCTGGG	TGCATACCAA	GCACTCCGTC	AGTGTTCCTT	GTTATTTCGAT
11751	GATTAGGAGG	CAGCTTAAAC	TAGAGGGAGT	TGAGCTGAAT	CAGGATGTTT
11801	GTCCCAGGTA	GCTGGGAATC	TGCCTAGCCC	AGTGCCAGT	TTATTTAGGT

## FIG. 10F

9351	GTCTGCTGCC	TTGCCAGGGT	GCCAGGAGGG	CGCTGCTGTG	GAAGCTGAGG
9401	CCCGGCCATC	CAGGGCGATG	CATTGGGCGC	TGATTCTTGT	TCCTGCTGCT
9451	GCCTCGGTGC	TTAGCTTTTG	AAACAATGAA	ATAAAATTAGA	ACCACTGTGA
9501	AAATCGATCA	GGGAATAAAT	TTAATGTGGA	AATAAACTGA	ACAACTTAGT
9551	TCTTCATAAG	AGTTTACTTG	GTAATACTT	GTGATGAGGA	CAAAACGAAG
9601	CACTAGAAAG	AGAGGCGACT	TGTAGACCTG	GGTGGCAGGA	GTGTTTTGTT
9651	TGTTTTCTTT	GGCAGGGTCT	TGCTCTGTTG	CTCAGGCTGG	AGTACAGTGG
9701	CACAATCACA	GCTCACTATA	GCCTCGACCT	CCTGGACTCA	AGCAATCCTC
9751	CTGCCTCAGC	CTCCCAGTAG	CTGGGACTAC	AGGCGCATGC	CACCATGCCT
9801	GGCTAATTTT	AAATTTTTTT	TTTTCTCTTT	TTTGAGATGG	AATCTCACTC
9851	TGTCGCCCAG	GCTGGAGTGC	AGTGGCGTGA	TCTCGGCTGA	CGGCAAGCTC
9901	CGCCTCCCAG	GTTCACTCCA	TTGCGCTGCC	TCAGCCTCCC	AAGTAGCTGG
9951	GACTACAGGC	GCTGGGATTA	CAAACCCAAA	CCCAAAGTGC	TGGGATTACA
10001	GGCGTGAGCC	ACTGCACCCG	GCCTGTTTTG	TCTTTCAATA	GCAAGAGTTG
10151	TGTTTGCTTC	GCCCCTACCT	TTAGTGGAAG	AATGTATAAA	ATGGAGATAT
10201	TGACCTCCAC	ATTGGGGTGG	TTAAATTATA	GCATGTATGC	AAAGGAGCTT
10251	CGCTAATTTA	AGGCTTTTTT	GAAAGAGAAG	AAACTGAATA	ATCCATGTGT
10301	GTATATATAT	TTTAAAAGCC	ATGGTCATCT	TTCCATATCA	GTAAAGCTGA
10351	GGCTCCCTGG	GACTGCAGAG	TTGTCCATCA	CAGTCCATTA	TAAGTGCGCT
10401	GCTGGGCCAG	GTGCAGTGGC	TTGTGCCTGA	ATCCCAGCAC	TTTGGGAGGC
10451	CAAGGCAGGA	GGATTCATTG	AGCCCAGGAG	TTTTGAGGCG	AGCCTGGGCA
10501	ATGTGGCCAG	ACCTCATCTC	TTCAAAAAAT	ACACAAAAAA	TTAGCCAGGC
10551	ATGGTGGCAC	GTGCCTGTAG	TCTCAGCTAC	TCAGGAGGCT	GAGGTGGGAG
10601	GATCACTTTG	AGCCTTGCGAG	GTCAAAGCTG	CAGTAAGCCA	TGATCTTGCC
10651	ACTGCATTCC	AGCCTGGATG	ACAGAGCCAG	ACCCTGTCTC	TAAAAAATAA
10701	AAAAACCAAA	CGGTGCACTG	TTTTCTTTTT	TCTTATCAAT	TTATTATTTT
10751	TAAATTAAAT	TTTCTTTTAA	TAATTTATAA	ATTATAAAAT	TATATTAAAA
10801	AATGACAAAT	TTTTATTACT	TATACATGAG	GTAAAACTTA	GGATATATAA
10851	AGTACATATT	GAAAAGTAAT	TTTTTGGCTG	GCACAGTGGC	TCACACCTGT
10901	AATCCCAGCA	CTTTGGGAGG	CCGTGGCGGG	CAGATCACAT	GAGATCATGA
10951	GTTTCGAGAC	AACCTGACCA	ACATGGAGAG	ACCCCATCTC	TACTAAAAAT
11001	ACAAAATTAG	CCGGGGTGGT	GGCGCATGCC	TGTAATCCCA	GCTACTCGGG
11051	AGGCTGAGGC	AGGAGAATCT	CTTGAACCCG	GGAGGCAGAG	GTTCGGGTGA
11101	GCCAAGATCG	TGCCTTTGCA	CACCAGCCTA	GGCAACAAGA	GCGAAAGTCC
11151	GTCTCAAAAA	AAAAGTAATT	TTTTTTAAGT	TAACCTCTGT	CAGCAAACAA
11201	ATTTAACCCA	ATAAAGGTCT	TTGTTTTTTA	ATGTAGTAGA	GGAGTTAGGG
11251	TTTATAAAAA	ATATGGTAGG	GAAGGGGGTC	CCTGGATTTG	CTAATGTGAT
11301	TGTCATTTGC	CCCTTAGGAG	AGAGCTCTGT	TAGCAGAATG	AAAAAATTGG
11351	AAGCCAGATT	CAGGGAGGGA	CTGGAAGCAA	AAGAATTTCT	GTTCGAGGAA
11401	GAGCCTGATG	TTTGCCAGGG	TCTGTTTAAC	TGGACATGAA	GAGGAAGGCT
11451	CTGGACTTTC	CTCCAGGAGT	TTCAGGAGAA	AGGTAGGGCA	GTGGTTAAGA
11501	GCAGAGCTCT	GCCTAGACTA	GCTGGGGTGC	CTAGACTAGC	TGGGGTGCCC
11551	AGACTAGCTG	GGGTGCCTAG	ACTAGCTGGG	TACTTTGAGT	GGCTCCTTCA
11601	GCCTGGACCT	CGGTTTCCTC	ACCTGTATAG	TAGAGATATG	GGAGCACCCA
11651	GCGCAGGATC	ACTGTGAACA	TAAATCAGTT	AATGGAGGAA	GCAGGTAGAG
11701	TGGTGCTGGG	TGCATACCAA	GCACTCCGTC	AGTGTTTCCT	GTTATTTCGAT
11751	GATTAGGAGG	CAGCTTAAAC	TAGAGGGAGT	TGAGCTGAAT	CAGGATGTTT
11801	GTCCCAGGTA	GCTGGGAATC	TGCCTAGCCC	AGTGCCAGT	TTATTTAGGT

## FIG. 10G

11851	GCTCTCTCAG	TGTTCCCTGA	TTGTTTTTTC	CTTTGTCATC	TTATCTACAG
11901	GATGTGACTG	GGAAGCTCTG	GTTTCAGTGT	CATGTGTCTA	TTCTTTATTT
11951	CCAGGCAAAG	GAAACCAACA	ATAAGAAGAA	AGAATTTGAG	GAAACTGCCA
12001	AGAAAGTGCG	CCGTGCCATC	GAGCAGCTGG	CTGCCATGGA	TTGAGGCCCTC
12051	TGGCCGGAGC	TGCCTGGTCC	CAGAGTGGCT	GCACCACTTC	CAGGGTTTAT
12101	TCCCTGGTGC	CACCAGCCTT	CCTGTGGGCC	CCTTAGCAAT	GTCTTAGGAA
12151	AGGAGATCAA	CATTTTCAAA	TTAGATGTTT	CAACTGTGCT	CCTGTTTTGT
12201	CTTGAAAGTG	GCACCAGAGG	TGCTTCTGCC	TGTGCAGCGG	GTGCTGCTGG
12251	TAAACAGTGGC	TGCTTCTCTC	TCTCTCTCTC	TTTTTTGGGG	GCTCATTTTT
12301	GCTGTTTTGA	TTCCCGGGCT	TACCAGGTGA	GAAAGTAGGG	AGGAAGAAGG
12351	CAGTGTCCCT	TTTGCTAGAG	CTGACAGCTT	TGTCGCGTGT	GGCAGAGCCT
12401	TCCACAGTGA	ATGTGTCTGG	ACCTCATGTT	GTTGAGGCTG	TCACAGTCCT
12451	GAGTGTGGAC	TTGGCAGGTG	CCTGTTGAAT	CTGAGCTGCA	GGTTCCTTAT
12501	CTGTCACACC	TGTGCCTCCT	CAGAGGACAG	TTTTTTTGTT	GTTGTGTTTT
12551	TTTGTTTTTT	TTTTTTGGTA	GATGCATGAC	TTGTGTGTGA	TGAGAGAATG
12601	GAGACAGAGT	CCCTGGCTCC	TCTACTGTTT	AACAACATGG	CTTTCITTAT
12651	TTGTTTGAAT	TGTTAATTCA	CAGAATAGCA	CAAACATAAA	TTAAACTAA
12701	GCACAAAGCC	ATTCTAAGTC	ATTGGGGAAA	CGGGGTGAAC	TTCAGGTGGA
12751	TGAGGAGACA	GAATAGAGTG	ATAGGAAGCG	TCTGGCAGAT	ACTCCTTTTG
12801	CCACTGCTGT	GTGATTAGAC	AGGCCAGTG	AGCCGCGGGG	CACATGCTGG
12851	CCGCTCCTCC	CTCAGAAAAA	GGCAGTGGCC	TAAATCCTTT	TTAAATGACT
12901	TGGCTCGATG	CTGTGGGGGA	CTGGCTGGGC	TGCTGCAGGC	CGTGTGCTG
12951	TCAGCCCAAC	CTTCACATCT	GTCACGTTCT	CCACACGGGG	GAGAGACGCA
13001	GTCCGCCCAG	GTCCCCGCTT	TCTTTGGAGG	CAGCAGCTCC	CGCAGGGCTG
13051	AAGTCTGGCG	TAAGATGATG	GATTTGATTC	GCCCTCCTCC	CTGTCATAGA
13101	GCTGCAGGGT	GGATTGTTAC	AGCTTCGCTG	GAAACCTCTG	GAGGTCATCT
13151	CGGCTGTTCC	TGAGAAATAA	AAAGCCTGTC	ATTTCAAACA	CTGCTGTGGA
13201	CCCTACTGGG	TTTTTAAAAT	ATTGTCAGTT	TTTCATCGTC	GTCCCTAGCC
13251	TGCCAACAGC	CATCTGCCCA	GACAGCCGCA	GTGAGGATGA	GCGTCTCGGC
13301	AGAGACGCAG	TTGTCTCTGG	GCGCTTGCCA	GAGCCACGAA	CCCCAGACCT
13351	GTTTGTATCA	TCCGGGCTCC	TTCCGGGCAG	AAACAACTGA	AAATGCACTT
13401	CAGACCCACT	TATTTATGCC	ACATCTGAGT	CGGCCTGAGA	TAGACTTTTC
13451	CCTCTAAACT	GGGAGAATAT	CACAGTGGTT	TTTGTTAGCA	GAAAATGCAC
13501	TCCAGCCTCT	GTAATCATCT	AAGCTGCTTA	TTTTTGATAT	TTGTGTCAGT
13551	CTGTAAATGG	ATACTTCACT	TTAATAACTG	TTGCTTAGTA	ATTGGCTTTG
13601	TAGAGAAGCT	GGAAAAAAT	GGTTTTGTCT	TCAACTCCTT	TGCATGCCAG
13651	GCGGTGATGT	GGATCTCGGC	TTCTGTGAGC	CTGTGCTGTG	GGCAGGGCTG
13701	AGCTGGAGCC	GCCCCCTCTCA	GCCCGCCTGC	CACGGCCTTT	CCTTAAAGGC
13751	CATCCTTAAA	ACCAGACCCT	CATGGCTGCC	AGCACCTGAA	AGCTTCCTCG
13801	ACATCTGTTA	ATAAAGCCGT	AGGCCCTTGT	CTAAGCGCAA	CCGCCTAGAC
13851	TTTCTTTCAG	ATACATGTCC	ACATGTCCAT	TTTTCAGGTT	CTCTAAGTTG
13901	GAGTGGAGTC	TGGGAAGGGT	TGTGAATGAG	GCTTCTGGGC	TATGGGTGAG
13951	GTTCCAATGG	CAGGTTAGAG	CCCCTCGGGC	CAACTGCCAT	CCTGGAAAGT
14001	AGAGACAGCA	GTGCCCCTG	CCCAGAAGAG	ACCAGCAAGC	CAAACTGGAG
14051	CCCCCATTCG	AGGCTGTGCG	CATGTGGAAA	GAGTAACTCA	CAATTGCCAA
14101	TAAAGTCTCA	TGTGGTTTTA	TCTACTTTTT	TTTTCTTTTT	CTTTTTTTTT
14151	GAGACAAGGC	CTTGCCCTCC	CAGGCTGGAG	TGCAGTGGAA	TGACCACAGC
14201	TCACCGCAAC	CTCAAATTCT	TGCGTTCAAG	TGAACCTCCC	ACTTTAGCCT

## FIG. 10H

```
14251   CCCAAGTAGC TGGGACTACA GGCGCACGCC ATCACACCCG GCTAATTGAA
14301   AAATTTTTTT TTTTGTTTAG ATGGAATCTC ACTTTGTTGC CCAGGCTGGT
14351   CTCAAACTCC TGGGCTCAAG TGATCATCCT GCTTCAGCGT CCGACTTGTT
14401   GGTATTATAG GCGTGAGCCA CTGGGCCTGA CCTAGCTACC ATTTTTTAAT
14451   GCAGAAATGA AGACTTGTAG AAATGAAATA ACTTGTCCAG GATAGTCGAA
14501   TAAGTAACTT TTAGAGCTGG GATTGAACC CAGGCAATCT GGCTCCAGAG
14551   CTGGGCCCTC ACTGCTGAAG GACACTGTCA GCTTGGGAGG GTGGCTATGG
14601   TCGGCTGTCT GATTCTAGGG AGTGAGGGCT GTCTTTAAAG CACCCCATTC
14651   CATTTTCAGA CAGCTTTGTC AGAAAGGCTG TCATATGGAG CTGACACCTG
14701   CCTCCCAAG GCTTCCATAG ATCCTCTCTG TACATTGTAA CCTTTTATTT
14751   TGAAATGAAA ATTCACAGGA AGTTGTAAGG CTAGTACAGG GGATCC
```




FIG. 11A




FIG. 11B






FIG. 11C




**FIG. 12**




**FIG. 13**




**FIG. 14A**




**FIG. 14B**




**FIG. 15A**




**FIG. 15B**



**FIG. 16**



**FIG. 17**



## **SURVIVIN, A PROTEIN THAT INHIBITS CELLULAR APOPTOSIS, AND ITS MODULATION**

### **STATEMENT OF RELATED APPLICATIONS**

[0001] This application is based upon provisional U.S. Application Serial No. 60/031,435, filed Nov. 20, 1996, the disclosure of which is hereby incorporated by reference.

### **FIELD OF THE INVENTION**

[0002] The present invention relates to the field of modulating cell apoptosis, particularly agents useful to inhibit apoptosis, as well as to diagnostic and prognostic assays involving conditions mediated by the expression of inhibitors of apoptosis. The invention specifically relates to the identification of a novel human gene, tentatively named Survivin. Survivin encodes a protein, Survivin, that inhibits cellular apoptosis, particularly in cancer cells and embryonic cells.

### **BACKGROUND OF THE INVENTION**

[0003] Regulation of cell proliferation by programmed cell death (apoptosis) maintains tissue homeostasis during development and differentiation (Raff, M. D., *Nature* (1992) 356:397-400; Vaux, D. L. et al., *Cell* (1994) 76:777-779). This process involves an evolutionarily conserved multi-step cascade (Oltvai, Z. et al., *Cell* (1994) 79:189-192), and is controlled by proteins that promote or counteract apoptotic cell death. Apoptosis also involves cell surface receptors (Smith, A. et al., *Cell* (1994) 76, 959-962), and associated signal transducers (Tartaglia, L. A. et al., *Immunol Today* (1992) 13:151-153), protease gene families (Martin, S. J. et al., *Cell* (1995) 82:349-352), intracellular second messengers (Kroemer, G. et al., *FASEB J* (1995) 9:1277-1287), tumor suppressor genes (Haffner, R. et al., *Curr Op Gen Dev* (1995) 5:84-90), and negative regulatory proteins that counteract apoptotic cell death (Hockenbery, D. et al., *Nature* (1990) 348:334-336). Aberrantly increased apoptosis or abnormally prolonged cell survival (Oltvai, Z. N. et al., *Cell* (1994) 79:189-192) may both contribute to the pathogenesis of human diseases, including autoimmune disorders, neurodegenerative processes, and cancer (Steller, H., *Science* (1995) 267:1445-1449; Thompson, C. B., *Science* (1995) 267:1456-1462).

[0004] Specifically, for example, inhibitors of apoptosis, most notably of the bcl-2 family (Reed, J., *J Cell Biol* (1994) 124:1-6, and Yang, E. et al., *Blood* (1996) 88:386-401), maintain lymphoid homeostasis and morphogenesis in adult (Hockenbery, D. et al., *Proc Natl Acad Sci USA* (1991) 88:6961-6965) and fetal (LeBrun, D. et al. (1993) 142:743-753) tissues. Deregulated expression of bcl-2 has also been implicated in cancer, by aberrantly prolonging cell survival and facilitating the insurgence of transforming mutations.

[0005] In addition to bcl-2, several members of a new gene family of inhibitors of apoptosis related to the baculovirus IAP gene (Bimbaum, M. J. et al., *J Virology* (1994) 68:2521-2528; Clem, R. J. et al., *Mol Cell Biol* (1994) 14:5212-5222) have been identified in *Drosophila* and mammalian cells (Duckett, C. S. et al., *EMBO J* (1996) 15:2685-2694; Hay, B. A. et al., *Cell* (1995) 83:1253-1262; Liston, P. et al., *Nature* (1996) 379:349-353; Rothe, M. et al., *Cell* (1995) 83:1243-1252; Roy, N. et al., *Cell* (1995) 80:167-178). These molecules are highly conserved evolutionarily; they

share a similar architecture organized in two or three approximately 70 amino acid amino terminus Cys/His baculovirus IAP repeats (BIR) and by a carboxy terminus zinc-binding domain, designated RING finger (Duckett, C. S. et al., *EMBO J* (1996) 15:2685-2694; Hay, B. A. et al., *Cell* (1995) 83:1253-1262; Liston, P. et al., *Nature* (1996) 379:349-353; Rothe, M. et al., *Cell* (1995) 83:1243-1252; Roy, N. et al., *Cell* (1995) 80:167-178). Recombinant expression of IAP proteins blocks apoptosis induced by various stimuli in vitro (Duckett, C. S. et al., *EMBO J* (1996) 15:2685-2694; Liston, P. et al., *Nature* (1996) 379:349-353), and promotes abnormally prolonged cell survival in the developmentally-regulated model of the *Drosophila* eye, in vivo (Hay, B. A. et al., *Cell* (1995) 83:1253-1262). Finally, deletions in a IAP neuronal inhibitor of apoptosis, NAIIP, were reported in 75% of patients with spinal muscular atrophy, thus suggesting a potential role of this gene family in human diseases (Roy, N. et al., *Cell* (1995) 80:167-178).

[0006] Therapeutic and diagnostic uses of nucleic acids that encode various inhibitors of apoptosis relating to a member of the IAP family have been described in the patent literature. See, for example, International Patent Applications No. WO 97/06255, WO 97/26331, and WO 97/32601. In particular, the uses of such genes and gene products are contemplated for the novel protein and its encoding nucleic acid discussed below.

[0007] Recently, a novel gene encoding a structurally unique IAP apoptosis inhibitor, designated Survivin has been identified. Survivin is a ~16.5 kD cytoplasmic protein containing a single BIR, and a highly charged carboxyl-terminus coiled-coil region instead of a RING finger, which inhibits apoptosis induced by growth factor (IL-3) withdrawal when transferred in B cell precursors (Ambrosini, G. et al., *Nature Med.* (1997) 3:917-921). At variance with bcl-2 or other IAP proteins, Survivin is undetectable in adult tissues, but becomes prominently expressed in all the most common human cancers of lung, colon, breast, pancreas, and prostate, and in ~50% of high-grade non-Hodgkin's lymphomas, in vivo. Intriguingly, the coding strand of the Survivin gene was highly homologous to the sequence of Effector cell Protease Receptor-1 (EPR-1) (Altieri, D. C., *FASEB J* (1995) 9:860-865), but oriented in the opposite direction, thus suggesting the existence of two separate genes duplicated in a head-to-head configuration.

[0008] The present invention is based on the identification of a novel human gene which is nearly identical to EPR-1, but oriented in the opposite direction. The antisense EPR-1 gene product, designated Survivin, is a distantly related member of the IAP family of inhibitors of apoptosis (Duckett, C. S. et al., *EMBO J* (1996) 15:2685-2694; Hay, B. A. et al., *Cell* (1995) 83:1253-1262; Liston, P. et al., *Nature* (1996) 379:349-353; Rothe, M. et al., *Cell* (1995) 83:1243-1252; Roy, N. et al., *Cell* (1995) 80:167-178), and is prominently expressed in actively proliferating transformed cells and in common human cancers, in vivo, but not in adjacent normal cells. Functionally, inhibition of Survivin expression by up-regulating its natural antisense EPR-1 transcript resulted in massive apoptosis and decreased cell growth.

### **SUMMARY OF THE INVENTION**

[0009] The present invention is based, in part, on the isolation and identification of a protein that is expressed in

most cancer cells and inhibits cellular apoptosis, hereinafter Survivin or the Survivin protein. Based on this observation, the present invention provides purified Survivin protein.

**[0010]** The present invention further provides nucleic acid molecules that encode the Survivin protein. Such nucleic acid molecules can be in an isolated form, or can be operably linked to expression control elements or vector sequences.

**[0011]** The present invention further provides methods of identifying other members of the Survivin family of proteins. Specifically, the nucleic acid sequence of Survivin can be used as a probe, or to generate PCR primers, in methods to identify nucleic acid molecules that encode other members of the Survivin family of proteins.

**[0012]** The present invention further provides antibodies that bind to Survivin. Such antibodies can be either polyclonal or monoclonal. Anti-Survivin antibodies can be used in a variety of diagnostic formats and for a variety of therapeutic methods.

**[0013]** The present invention further provides methods for isolating Survivin binding partners. Survivin binding partners are isolated using the Survivin protein as a capture probe. Alternatively, Survivin can be used as bait in the yeast two-hybrid system to screen an expression library and identify genes that encode proteins that bind to the Survivin protein. Binding partners isolated by these methods are useful in preparing antibodies and also serve as targets for drug development.

**[0014]** The present invention further provides methods to identify agents that can block or modulate the association of Survivin with a binding partner. Specifically, an agent can be tested for the ability to block, reduce or otherwise modulate the association of Survivin with a binding partner by contacting Survivin, or a fragment thereof, and a binding partner with a test agent and determining whether the test agent blocks or reduces the binding of the Survivin protein to the binding partner.

**[0015]** The present invention further provides methods for reducing or blocking the association of Survivin with one or more of its binding partners. Specifically, the association of Survivin with a binding partner can be blocked or reduced by contacting Survivin, or the binding partner, with an agent that blocks the binding of Survivin to the binding partner. The method can utilize an agent that binds to Survivin or to the binding partner.

**[0016]** The present invention further provides methods of regulating the expression of Survivin within a cell. Expression of Survivin within a cell can be regulated so as to produce or inhibit the production of Survivin.

**[0017]** Blocking Survivin/binding partner associations or Survivin expression can be used to modulate biological and pathological processes that require Survivin. For example, methods that reduce Survivin production induce apoptosis of tumor cells. Stimulation of Survivin production can be used as a means of extending the culturability of cells or tissues.

**[0018]** The biological and pathological processes that require Survivin or Survivin/binding partner interactions can further be modulated using gene therapy methods. Additional genetic manipulation within an organism can be used to alter the expression of a Survivin gene or the production of a Survivin protein in an animal model. For example, a

Survivin gene can be altered to correct a genetic deficiency; peptide modulators of Survivin activity can be produced within a target cell using genetic transformation methods to introduce a modulator encoding nucleic acid molecules into a target cell; etc. The use of nucleic acids for antisense and triple helix therapies and interventions are expressly contemplated.

**[0019]** The present invention further provides methods of reducing the severity of pathological processes that require Survivin. Since expression of Survivin or association of Survivin with a binding partner is required for Survivin-mediated biological processes, agents that block Survivin expression, Survivin activity or the association of Survivin with a binding partner, can be used in therapeutic methods.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0020]** **FIG. 1** shows the identification of a complementary EPR-1 gene. A, B. Chromosomal location. A digoxigenin-labeled human P1 genomic clone selected by hybridization with the EPR-1 cDNA, was incubated with metaphase chromosomes isolated from phytohemagglutinin-stimulated PBMC in 50% formamide, 10% dextran sulfate and 2×SSC. The EPR-1-hybridizing gene was mapped in single-color labeling to the long arm of a group E chromosome (A, green staining), and in two-color staining with probe D17Z1, specific for the centromere of chromosome 17 (B, red staining), to the long arm of chromosome 17 (B, green staining), to band 17q25. C. Map of the antisense EPR-1 gene. A contig spanning 14796 bp was derived from two EPR-1-hybridizing P1 clones, subcloned in pBSKS<sup>+</sup>, and completely sequenced on both strands. Orientation of the map is 5'→3' with respect to the position of intron-exon boundaries (see below). Exons are solid boxes, a putative CpG island upstream exon 1 is an open box. The translational initiation codon (ATG) is indicated. Restriction sites are: B, BamHI, H, HindIII; P, PstI; S, SmaI; X, XbaI. D. Intron-exon boundaries of the antisense EPR-1 gene. Positions of the intron-exon boundaries in bp are indicated in parenthesis.

**[0021]** **FIG. 2** shows the complexity and evolutionary conservation of EPR-1-related sequences. A. Southern blot of human genomic DNA. Samples were digested with the indicated restriction enzymes, transferred to GeneScreen nylon membranes and hybridized with the EPR-1 cDNA, in 5×SSC, 0.5% SDS, 5×Denhardt's and 0.1% sodium pyrophosphate at 65° C. Radioactive bands indicated by an arrow (7.6 kb BamHI, 7.5 kb XbaI and HindIII fragments of 15, 7.5, 6.4, and 3.7 kb) do not derive from the antisense EPR-1 gene in **FIG. 1C**. B. Southern blot of pulsed field gel electrophoresis. High molecular weight human genomic DNA was digested with the indicated restriction enzymes, separated by pulsed field gel electrophoresis for 20 h at 200 V with a pulse time of 75 sec, transferred to nylon membrane, and hybridized with the EPR-1 cDNA, as described in A. C. Multiple species Southern blot. EcoRI-digested genomic DNA from the indicated species was hybridized with a 3' 548 bp fragment of the EPR-1 cDNA, as described in A. For all panels, molecular weight markers in kb are shown on the left.

**[0022]** **FIG. 3** shows the discordant tissue distribution of sense/antisense EPR-1 transcripts. Northern hybridization was carried out on a multiple tissue adult or fetal mRNA blot

with single strand-specific probes in 5×SSPE, 10×Denhardt's solution, 2% SDS, 100 mg/ml denatured salmon sperm DNA at 60° C. for 14 h. After washes in 2×SSC at 60° C. and in 0.2×SSC at 22° C., radioactive bands were visualized by autoradiography. A. EPR-1-specific single-strand probe. B. Antisense EPR-1-specific single-strand probe. C. Control actin probe. Molecular weight markers in kb are shown on the left.

**[0023] FIG. 4** shows the sequence analysis of Survivin and expression in cell lines. A. Predicted translation of the antisense EPR-1 gene product (Survivin). B. Sequence alignment of the BIR in Survivin and in other IAP proteins by the Clustal method. IAP proteins are identified by accession number, L49433, TNFR2-TRAF signaling complex-associated IAP; L49441, apoptosis 2 inhibitor (*Drosophila*); P41436, IAP gene from *Cydia pomonella* granulosus virus; P41437, IAP gene from *Orgyia pseudotsugata* nuclear polyhedrosis virus; U19251, NAIP, neuronal inhibitor of apoptosis; U32373, IAP-like protein ILP from *Drosophila melanogaster*; U32974, human IAP-like protein ILP; U36842, mouse inhibitor of apoptosis; U45878, human inhibitor of apoptosis 1; U45879, human inhibitor of apoptosis 2; U45880, X-linked inhibitor of apoptosis; U45881, *Drosophila* inhibitor of apoptosis. Conserved residues are boxed, identities between Survivin and NAIP (U19251) are boxed and shaded. C. Immunoblotting with anti-Survivin antibody JC700. Protein-normalized aliquots of SDS-extracts of cell lines HEL (erythroleukemia), Daudi and JY (B lymphoma), THP-1 (monocytic), Jurkat and MOLT13 (T leukemia), or non transformed human lung Lu18 fibroblasts, HUVEC or PBMC were separated by electrophoresis on a 5-20% SDS gradient gel, transferred to Immobilon and immunoblotted with control non-immune rabbit IgG (RbIgG), or anti-Survivin antibody JC700 (Survivin). Protein bands were visualized by alkaline phosphatase-conjugated goat anti-rabbit IgG and tetrazolium salts. Molecular weight markers in kDa are shown on the left.

**[0024] FIG. 5** shows the regulation of Survivin expression by cell growth/differentiation. HL-60 cells were terminally differentiated to a mature monocytic phenotype by a 72 h culture with 0.1 mM vitamin D<sub>3</sub> plus 17.8 mg/ml indomethacin. Survivin expression before or after vitamin D<sub>3</sub> differentiation was detected by immunoblotting with JC700 antibody, or by Northern hybridization with a Survivin-specific single strand probe. RbIgG, control non-immune rabbit IgG. Protein molecular weight markers in kDa and position of ribosomal bands are shown on the left of each blot.

**[0025] FIG. 6** shows the over-expression of Survivin in human cancer, in vivo. A. Immunohistochemical staining of human lung adenocarcinoma with affinity-purified anti-Survivin antibody JC700 (20 µg/ml). B. Inhibition of JC700 staining of lung adenocarcinoma by pre-absorption with the immunizing Survivin 3-19 peptide. C. Immunohistochemical expression of Survivin in squamous lung cell carcinoma, but not in the adjacent normal gland epithelium of the lung (C, arrow). D. In-situ hybridization of Survivin mRNA in squamous lung cell carcinoma with a Survivin-specific riboprobe. E. Expression of Survivin in pancreatic adenocarcinoma by immunohistochemistry with JC700. F. Normal pancreas, negative for Survivin expression by immunohistochemistry. G. In situ hybridization of Survivin mRNA expression in colon adenocarcinoma, but H, not in the

adjacent non neoplastic colon gland epithelium (H, arrow). Magnifications are ×200, except G, ×400.

**[0026] FIG. 7** shows the effect of Survivin on apoptosis/proliferation. A. EPR-1-regulation of Survivin expression. HeLa cells were transfected with control vector pML1 or the EPR-1 cDNA (which is antisense to Survivin) by electroporation, and selected in hygromycin (0.4 mg/ml). Aliquots of vector control HeLa cells (Vector) or Survivin antisense transfectants (Antisense) were induced with 200 mM ZnSO<sub>4</sub> detergent-solubilized, and immunoblotted with the anti-Survivin JC700 antibody. Molecular weight markers are shown on the left. B. Effect of Survivin on apoptosis. Survivin antisense transfectants (1, 2), or vector control HeLa cells (3, 4) were induced with Zn<sup>2+</sup> ions in 0% FBS for 24 h and stained by the AptoTag method with TdT-catalyzed dUTP labeling of 3'-OH DNA ends and immunoperoxidase (1, 3), or by hematoxylin-eosin (HE) (2, 4). 1. Prominent nuclear DNA fragmentation detected by AptoTag staining in serum-starved Survivin antisense transfectants; 2. HE staining of antisense transfectants reveals the presence of numerous apoptotic bodies (arrows); 3. AptoTag staining of vector control HeLa cells detects a few sparse apoptotic cell (arrow); 4. HE staining of vector control HeLa cells. The arrow indicates a single apoptotic body. Magnification ×400. C. Effect of Survivin on cell growth. Twenty thousands vector control HeLa cells (Vector) or Survivin antisense transfectants (Antisense) were seeded in 24-well plates, induced with ZnSO<sub>4</sub>, harvested at the indicated time points, and cell proliferation was determined microscopically by direct cell count. Data are the mean±SEM of replicates of a representative experiment out of seven independent determinations.

**[0027] FIG. 8** shows the expression of Survivin in HL-60 cells. HL-60 cells were examined via Western and Northern blots for Survivin expression.

**[0028] FIG. 9** presents a structural analysis of Survivin. The Survivin protein was analyzed using the Chou-Fasman, Garnier-Robson, Kyle-Doolittle, Eisenberg, Karplus-Schultz, Jameson-Wolf and Emini analysis methods.

**[0029] FIG. 10** shows the nucleotide Sequence of Survivin.

**[0030] FIG. 11** shows the expression of Survivin and the generation and characterization of anti-Survivin mAb 8E2 by ELISA and immunoblotting.

**[0031] FIG. 12** shows the site-directed mutagenesis of Survivin and identification of key functional residues involved in apoptosis inhibition.

**[0032] FIG. 13** shows the cytoprotective effect of Survivin addition on endothelial cell apoptosis.

**[0033] FIG. 14** shows that the presence of Survivin is a negative predictive-prognostic factor in neuroblastoma

**[0034] FIG. 15** shows that the presence of Survivin is a negative predictive prognostic factor in high-grade non-Hodgkin's lymphoma.

**[0035] FIG. 16** shows the down regulation of Survivin induced by inflammatory and cytostatic cytokines.

**[0036] FIG. 17** shows the effects of Survivin constructs or XIAP on apoptosis induced in NIH3T3 cells by hydrogen peroxide.



## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

### [0037] I. General Description

[0038] The present invention is based in part on identifying a novel protein that is expressed in tumor cells and inhibits cellular apoptosis, hereinafter the Survivin protein or Survivin. Survivin is also found to be expressed in embryonic tissues.

[0039] The Survivin protein can be used as an agent, or serve as a target for agents, that can be used to inhibit or stimulate Survivin mediated inhibition of cellular apoptosis, for example to block abnormal cell growth or to extend cell growth in culture.

[0040] As used herein, modulation of apoptosis means increasing or decreasing the number of cells that would otherwise undergo apoptosis in a given cell population. This can be effected by increasing or decreasing the amount of Survivin present in a cell or by increasing or decreasing the activity of the Survivin. Preferably, the given cell population in which apoptosis is to be modulated is found in a tumor or other tissue or group of cells in which beneficial effect results from the modulation. Also, preferably, the increase or decrease in number of cells that would otherwise undergo apoptosis in a given cell population is at least about 10%, 20%, 40% or more preferably at least about 50% of the cells in that population.

[0041] The present invention is further based on the development of methods for isolating proteins that bind to Survivin. Probes based on the Survivin protein or fragments of Survivin as discussed below are used as capture probes to isolate Survivin binding proteins. Dominant negative proteins, DNAs encoding these proteins, antibodies to these binding proteins, peptide fragments of these proteins or mimics of these proteins may be introduced into cells to affect Survivin function. Additionally, these proteins provide novel targets for screening of synthetic small molecules and combinatorial or naturally occurring compound libraries to discover novel therapeutics to regulate Survivin function.

### [0042] II. Identification, General Characterization and Tissue Distribution of Survivin

[0043] The present invention is based on the identification on chromosome 17q25 of a novel member of the IAP family of inhibitors of apoptosis, designated Survivin, which may confer a selective advantage for cancer cell growth. Relevant features of the Survivin gene include its developmentally- and differentiation-regulated expression, its nearly identical and complementary DNA sequence with the factor Xa receptor EPR-1, and its abundant in vivo expression in common human malignancies, but not in the adjacent non-neoplastic population. As described below, targeting Survivin expression by metallothionein-induction of EPR-1 mRNA resulted in apoptosis and inhibition of proliferation of HeLa cell transfectants.

[0044] In addition to their contribution to hemostasis, cellular receptors for blood proteases have recently emerged as pleiotropic signaling molecules, playing a crucial role in embryologic development (Connolly, A. J. et al., *Nature* (1996) 381:516-519), and vasculogenesis (Carmeliet, P. et al., *Nature* (1996) 383:73-75). In this context, the Survivin gene was isolated by hybridization with the cDNA for

EPR-1, a receptor for factor Xa contributing to procoagulant activity (Altieri, D. C., *FASEB J* (1995) 9:860-865), and T cell activation (Duchosal, M. A. et al., *Nature* (1996) 380:352-356). Although the Survivin coding sequence was found to be nearly identical to the EPR-1 cDNA, its orientation was unambiguously assigned to the antisense EPR-1 strand for the position of the consensus splice sites at intron-exon boundaries (Padgett, R. A. et al., *Ann Rev Biochem* (1986) 55:1119-1150). On the other hand, the authenticity of the EPR-1 "sense" strand was demonstrated in previous studies, when mammalian cells transfected with the EPR-1 cDNA or with chimeric EPR-1 constructs (Ambrosini, G. et al., *J Biol Chem* (1996) 271:1243-1248 and Altieri, D. C., *FASEB J* (1995) 9:860-865), were recognized by anti-EPR-1 mAbs and bound factor Xa in a specific and saturable reaction.

[0045] These findings could be reconciled by the existence of multiple, highly homologous, EPR-1 transcripts oriented in opposite directions. The heterogeneity of EPR-1 mRNA and the complex pattern of Southern hybridization support this hypothesis. Previously, double strand EPR-1 probes detected three strongly hybridizing bands of 1.9, 3.4 and ~1.5 kb in mRNA of EPR-1<sup>+</sup> cells (Altieri, D. C., *FASEB J* (1995) 9:860-865). Here, single strand-specific probes confirmed the presence of multiple mature and polyadenylated EPR-1-related messages, and revealed that the 1.9 and 3.4 kb bands corresponded to two highly regulated, antisense EPR-1 transcripts, while the 1.5 kb band, more accurately defined as 1.2 kb, coincided with a genuine EPR-1-encoding message. While the 1.9 kb antisense transcript clearly originated from the Survivin gene described here, a gene encoding the 1.2 kb "sense" EPR-1 message has not yet been identified.

[0046] However, (i) the presence of several genomic EPR-1-hybridizing bands unrelated to the Survivin gene, (ii) the different restriction pattern of EPR-1 sequences in various species, and (iii) the numerous expressed sequence tag database entries matching ( $P=0.018-7 \times 10^{-11}$ ) the positive (accession n. W46267), or the negative (accession n. W34764, W83810, T29149) EPR-1 strand, altogether suggest the existence of at least a second, highly-related, EPR-1 gene oriented in the opposite direction to that described here, and encoding the previously characterized factor Xa receptor (Altieri, D. C., *FASEB J* (1995) 9:860-865).

[0047] A similar situation could arise from gene duplication event(s) involving EPR-1 sequences. Interestingly, the single hybridization signal detected on chromosome 17q25, and the single hybridizing bands identified in a Southern blot of high molecular weight genomic DNA, suggest that EPR-1-related sequences potentially oriented in opposite directions may be adjacent in close proximity, within a physical interval of 75-130 kb.

[0048] The presence of multiple EPR-1 transcripts oriented in opposite directions implies a reciprocal regulatory mechanism by naturally occurring antisense. This is consistent with the predominantly discordant and mutually exclusive distribution of sense and antisense EPR-1 messages in developing or adult tissues in vivo, and during HL-60 cell terminal differentiation. While antisense regulation is common in prokaryotes (Green, P. J. et al., *Annu Rev Biochem* (1986) 55:569-597), a growing number of eukaryotic gene products have been recently characterized for the occurrence

of functional antisense transcripts potentially participating in gene regulation, including basic fibroblast growth factor (Kimmelman, D. et al., *Cell* (1989) 59:687-696; Murphy, P. R. et al., *Molecular Endocrinology* (1994) 8:852-859),  $\alpha 1(I)$  collagen (Farrell, C. M. et al., *J Biol Chem* (1995) 270:3400-3408 and Lukens, 1995), n-myc (Krystal, G. W. et al., *Mol Cell Biol* (1990) 10:4180-4191), c-myc (Celano, P. et al., *J Biol Chem* (1992) 267:15092-15096), p53 (Khochbin, S. et al., *EMBO J* (1989) 8:4107-4114), c-erbAa (Lazar, M. A. et al., *Mol Cell Biol* (1989) 9:1128-1136), and CD3  $\zeta/\eta/\theta$  locus (Lerner, A. et al., *J Immunol* (1993) 151:3152-3162).

[0049] As described below, the existence of a EPR-1/Survivin gene balance regulated by functional antisense was demonstrated in HeLa cell transfectants, when metallothionein-induced transcription of the EPR-1 "sense" strand suppressed the expression of Survivin and profoundly influenced apoptosis/cell proliferation (see below). This regulatory mechanism was not due to a potential protein association between EPR-1 and Survivin, since the EPR-1 construct used for these experiments lacked a translational initiation codon. Additional experiments have evaluated the ability of a Survivin antisense to inhibit cell growth. This was done by transiently co-transfecting the Survivin antisense with a lacZ reported plasmid and making a determination of cell viability after a 48-h transfection in  $\beta$ -galactosidase expressing cells. The results indicated that the viability of Survivin antisense transfectants was <20% of control cells transfected with the empty vector. A control antisense of ICAM-1 (intercellular adhesion molecule-1) similarly co-transfected in HeLa cells was ineffective.

[0050] Survivin was found to be a small protein of 142 amino acids (~16.5 kDa) with no amino acid sequence homology to EPR-1, and designated Survivin for the presence of a BIR-homologous domain (Birnbaum, M. J. et al., *J Virology* (1994) 68:2521-2528; Clem, R. J. et al., *Mol Cell Biol* (1994) 14:5212-5222) found in IAP inhibitors of apoptosis (Duckett, C. S. et al., *EMBO J* (1996) 15:2685-2694; Hay, B. A. et al., *Cell* (1995) 83:1253-1262; Liston, P. et al., *Nature* (1996) 379:349-353; Rothe, M. et al., *Cell* (1995) 83:1243-1252; Roy, N. et al., *Cell* (1995) 80:167-178). Based on overall sequence conservation, the absence of a carboxy terminus RING finger and the presence of a single, partially conserved, BIR domain, Survivin is the most distantly related member of the IAP family, sharing the highest degree of similarity with NAIP (Roy, N. et al., *Cell* (1995) 80:167-178). Thus, unlike bcl-2 or other IAP proteins, Survivin is undetectable in adult tissues, but becomes prominently expressed in all the most common human cancers of lung, colon, breast, pancreas, and prostate, and in ~50% of high-grade non-Hodgkin's lymphomas, in vivo. Additionally, unlike other IAP proteins (Deveraux, Q. et al., *Nature* (1997) 388:300-304), Survivin does not bind caspases in a cell-free system (Roy, N. et al., *Blood* (1997) 595:2645).

[0051] Consistent with the anti-apoptosis properties of IAP proteins in vitro (Duckett, C. S. et al., *EMBO J* (1996) 15:2685-2694; Liston, P. et al., *Nature* (1996) 379:349-353), and in vivo (Hay, B. A. et al., *Cell* (1995) 83:1253-1262), inhibition of Survivin expression by the EPR-1 transcript (which naturally is antisense to Survivin) resulted in increased apoptosis, as determined by in situ internucleosomal DNA fragmentation in HeLa cell transfectants. The ability of a RING finger-less IAP protein to counteract

apoptosis is not without a precedent, as demonstrated by the suppression of apoptosis mediated by NAIP (Liston, P. et al., *Nature* (1996) 379:349-353), and by the in vivo gain-of-function of a Drosophila IAP protein following deletion of the RING finger (Hay, B. A. et al., *Cell* (1995) 83:1253-1262). Although anti-apoptosis genes are thought to play an indirect role in cell growth, by favoring the accumulation of oncogenic mutations(s) in aberrantly long-living cells (Reed, J. C., *J Cell Biol* (1994) 124:1-6), down-regulation of Survivin resulted in a profound inhibition of HeLa cell proliferation. While this may derive from rapid disappearance of HeLa cells expressing the highest levels of antisense transcripts by apoptosis, a similar decrease in tumor cell proliferation has been reported in vivo after antisense inhibition of bcl-2 (Reed, J. C. et al., *Proc Natl Acad Sci USA* (1990) 87:3660-3664).

[0052] The possibility that IAP proteins may play a more general role in cell proliferation, not exclusively restricted to apoptosis inhibition, has been proposed earlier. Rothe et al., have recently demonstrated that the amino terminus BIR in two IAP proteins (cIAPs) physically interacts with the signal transducers associated with the 75 kDa TNF receptor (Rothe, M. et al., *Cell* (1995) 83:1243-1252), a molecule primarily implicated in cell proliferation and survival rather than apoptotic signaling (Tartaglia, L. A. et al., *Immunol Today* (1992) 13:151-153). While it is not known if Survivin is physically linked to signaling molecules (Rothe, M. et al., *Cell* (1995) 83:1243-1252), the structural divergence of its BIR as compared with other IAP proteins (Duckett, C. S. et al., *EMBO J* (1996) 15:2685-2694; Hay, B. A. et al., *Cell* (1995) 83:1253-1262; Liston, P. et al., *Nature* (1996) 379:349-353; Rothe, M. et al., *Cell* (1995) 83:1243-1252; Roy, N. et al., *Cell* (1995) 80:167-178), may confer specificity for supramolecular interaction(s) potentially relevant to its particular mechanism of apoptosis inhibition/cell growth.

[0053] Dysregulation of programmed cell death (apoptosis) has recently emerged as a primary mechanism contributing to the pathogenesis of various human diseases, including cancer (Steller, H., *Science* (1995) 267:1445-1449; Thompson, C. B., *Science* (1995) 267:1456-1462). While the impact of anti-apoptosis gene(s) in neoplasia is highlighted by the role of bcl-2 in follicular lymphoma (Korsmeyer, S. J., *Blood* (1992) 80:879-886), a potential distribution of IAP proteins in cancer had not been previously investigated. In this context, one of the most striking characteristics of Survivin was its abundant expression in actively proliferating transformed cell lines, and in all the most common human malignancies of lung, colon, pancreas, and breast, in vivo, but not in the non-neoplastic adjacent cell population. This distribution in multiple human cancers may signal a fundamental role of this molecule in apoptosis/cell proliferation mechanisms in neoplasia. By analogy with the paradigm of bcl-2, over-expression of Survivin in cancer may lead to aberrantly prolonged cell viability (Veis, D. J. et al., *Cell* (1993) 75:229-240), increased resistance to chemotherapy-induced apoptosis (Miyashita, T. et al., *Blood* (1993) 81:151-157), and, as suggested by the in vitro studies reported above, in a direct advantage for transformed cell proliferation.

[0054] On the other hand, for its presence in normal PBMC and benign breast adenomas, in vivo (unpublished observations), Survivin expression cannot be interpreted per

se as a marker of malignant transformation but may reflect a more general, developmental- or cell type-specific response to certain stimuli. This is consistent with the presence of Survivin during normal embryonic (our unpublished observations) and fetal development, and its rapid disappearance in growth-arrested cell types (i.e. vitamin D<sub>3</sub>-treated HL-60), and terminally-differentiated tissues, in vivo. At variance with other IAP proteins which are constitutively found in adult mature tissues (Duckett, C. S. et al., *EMBO J* (1996) 15:2685-2694; Liston, P. et al., *Nature* (1996) 379:349-353; Rothe, M. et al., *Cell* (1995) 83:1243-1252), this pattern of expression is reminiscent of the distribution of bcl-2 in fetal tissues (LeBrun, D. P. et al., *Am J Pathol* (1993) 142:743-753), and its more restricted presence in differentiated cells, correlating with susceptibility to apoptosis (Hockenbery, D. M. et al., *Proc Natl Acad Sci USA* (1991) 88:6961-6965).

[0055] In summary, these findings identify Survivin as a novel link between IAP proteins and cancer, in vivo. A key implication of the data presented below is the possibility to balance the effect of this potent anti-apoptosis gene by manipulating a normal cell regulatory mechanism, centered on the expression of EPR-1 (Altieri, D. C., *FASEB J* (1995) 9:860-865). Targeting Survivin may then remove a selective advantage for transformed cell growth and be therapeutically beneficial to increase the susceptibility of cancer cells to chemotherapy-induced apoptosis. Along the same line, identification of polymorphic markers and construction of extended aptotypes within and around the EPR-1/Survivin locus may provide new insights on the population genetics of susceptibility to chemotherapy.

### [0056] III. Specific Embodiments

#### [0057] A. Survivin Protein

[0058] The present invention provides isolated Survivin protein, as well as allelic variants of the Survivin protein, and conservative amino acid substitutions of the Survivin protein. As used herein, the Survivin protein (or Survivin) refers to a protein that has the amino acid sequence of human Survivin depicted in FIG. 4. The term "Survivin protein" also includes naturally occurring allelic variants of Survivin, naturally occurring proteins that have a slightly different amino acid sequence than that specifically recited above. Allelic variants, though possessing a slightly different amino acid sequence than those recited above, will still have the requisite ability to inhibit cellular apoptosis.

[0059] As used herein, the Survivin family of proteins refers to Survivin proteins that have been isolated from organisms in addition to humans. The methods used to identify and isolate other members of the Survivin family of proteins are described below.

[0060] Survivin is a member of the IAP (inhibitory apoptosis proteins) family of protein. However, Survivin is the first member of a unique subfamily of IAP proteins that differ from other IAP proteins in significant ways. Despite homology and sequence conservation in the BIR module between Survivin and other members of this gene family, there are important structural differences that are unique to members of the Survivin family of proteins. First unlike any other IAP protein, Survivin has only one BIR module (most of the other molecules have 2-3). Further, Survivin does not contain a carboxy-terminal RING finger but has a predicted

coiled-coil instead. Only the Neuronal Apoptosis Inhibitory Protein (NAIP) in the IAP family lacks a RING finger, but does not contain a carboxy-terminus coiled coil. Finally there is no DNA sequence similarity between Survivin and other IAP proteins (PCR primers designed on Survivin are unlikely to detect other IAP proteins and vice-versa).

[0061] The Survivin proteins of the present invention are preferably in isolated form. As used herein, a protein is said to be isolated when physical, mechanical or chemical methods are employed to remove the Survivin protein from cellular constituents that are normally associated with the Survivin protein. A skilled artisan can readily employ standard purification methods to obtain an isolated Survivin protein.

[0062] The Survivin proteins of the present invention further include conservative variants of the Survivin proteins herein described. As used herein, a conservative variant refers to alterations in the amino acid sequence that do not adversely affect the ability of the Survivin protein to bind to a Survivin binding partner and/or to inhibit cellular apoptosis. A substitution, insertion or deletion is said to adversely affect the Survivin protein when the altered sequence prevents the Survivin protein from associating with a Survivin binding partner and/or prevents the Survivin protein from inhibiting cellular apoptosis. For example, the overall charge, structure or hydrophobic/hydrophilic properties of Survivin can be altered without adversely affecting the activity of Survivin. Accordingly, the amino acid sequence of Survivin can be altered, for example to render the peptide more hydrophobic or hydrophilic, without adversely affecting the activity of Survivin.

[0063] The allelic variants, the conservative substitution variants and the members of the Survivin family of proteins, will have the ability to inhibit cellular apoptosis. Such proteins will ordinarily have an amino acid sequence having at least about 75% amino acid sequence identity with the human Survivin sequence, more preferably at least about 80%, even more preferably at least about 90%, and most preferably at least about 95%. Identity or homology with respect to such sequences is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the known peptides, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent homology, and including any conservative substitutions as being homologous. N-terminal, C-terminal or internal extensions, deletions, or insertions into the peptide sequence shall not be construed as affecting homology.

[0064] Thus, the Survivin proteins of the present invention include molecules having the amino acid sequences disclosed in FIG. 1; fragments thereof having a consecutive sequence of at least about 3, 5, 10 or 15 amino acid residues of the Survivin protein; amino acid sequence variants of such sequence wherein an amino acid residue has been inserted N- or C-terminal to, or within, the disclosed Survivin sequence; amino acid sequence variants of the disclosed Survivin sequence, or their fragments as defined above, that have been substituted by another residue. Contemplated variants further include those containing predetermined mutations by, e.g., homologous recombination, site-directed or PCR mutagenesis, and the corresponding Survivin proteins of other animal species, including but not

limited to rabbit, rat, murine, porcine, bovine, ovine, equine and non-human primate species, and the alleles or other naturally occurring variants of the Survivin family of proteins; and derivatives wherein the Survivin protein has been covalently modified by substitution, chemical, enzymatic, or other appropriate means with a moiety other than a naturally occurring amino acid (for example a detectable moiety such as an enzyme or radioisotope). The recombinant Survivin protein also can be used to solve the molecular structure of Survivin by 2D-NMR, circular dichroism and X-ray crystallography, thus integrating the site-directed mutagenesis approach and the rational design of specific small molecule inhibitors.

[0065] As described below, members of the Survivin family of proteins can be used: 1) as a target to block Survivin mediated inhibition of cellular apoptosis, 2) to identify and isolate binding partners that bind Survivin, 3) in methods to identify agents that block the association of Survivin with a Survivin binding partner, 4) as a target to assay for Survivin mediated inhibition of cellular apoptosis, 5) as an agent to block cellular apoptosis, administered alone or as part of a combination therapy, 6) as a binding partner in an assay to quantitate circulating levels of anti-Survivin antibodies, 7) as an antigen to elicit production of anti-Survivin antibodies that in turn can be used in an assay to quantitate circulating levels of Survivin and or can be used for immunohistochemical purposes, and 8) as a therapeutic anticancer vaccine, or component of a polyvalent vaccine.

#### [0066] B. Anti-Survivin Antibodies

[0067] The present invention further provides antibodies that selectively bind to a Survivin protein. The anti-Survivin antibodies particularly contemplated include monoclonal and polyclonal antibodies as well as fragments containing the antigen binding domain and/or one or more complement determining regions.

[0068] Antibodies are generally prepared by immunizing a suitable mammalian host using a Survivin protein, or fragment, in isolated or immunoconjugated form (Harlow, Antibodies, Cold Spring Harbor Press, NY (1989)). FIG. 9 provides a Jameson-Wolf plot of the antigenic index of various regions of Survivin. Such regions, in combination with the other structural analysis provided in FIG. 9, provide suitable fragments for use in generating Survivin specific antibodies. Methods for preparing immunogenic conjugates of a protein with a carrier such as BSA, KLH, or other carrier proteins are well known in the art. In some circumstances, direct conjugation using, for example, carbodiimide reagents may be used; in other instances linking reagents such as those supplied by Pierce Chemical Co., Rockford, Ill., may be effective.

[0069] Administration of the Survivin immunogen is conducted generally by injection over a suitable time period and with use of a suitable adjuvant, as is generally understood in the art. During the immunization schedule, titers of antibodies can be taken to determine adequacy of antibody formation.

[0070] While the polyclonal antisera produced in this way may be satisfactory for some applications, for pharmaceutical compositions, monoclonal antibody preparations are preferred. Immortalized cell lines which secrete a desired monoclonal antibody may be prepared using the standard

method of Kohler and Milstein or modifications which effect immortalization of lymphocytes or spleen cells, as is generally known. The immortalized cell lines secreting the desired antibodies are screened by immunoassay in which the antigen is the Survivin peptide. When the appropriate immortalized cell culture secreting the desired antibody is identified, the cells can be cultured either in vitro or by production in ascites fluid.

[0071] The desired monoclonal antibodies are then recovered from the culture supernatant or from the ascites supernatant. Fragments of the monoclonals or the polyclonal antisera which contain the immunologically significant portion can be used as antagonists, as well as the intact antibodies. Use of immunologically reactive fragments, such as the Fab, Fab', of F(ab')<sub>2</sub> fragments is often preferable, especially in a therapeutic context, as these fragments are generally less immunogenic than the whole immunoglobulin.

[0072] The antibodies or fragments may also be produced, using current technology, by recombinant means. Regions that bind specifically to the desired regions of receptor can also be produced in the context of chimeras or CDR grafted antibodies of multiple species origin.

[0073] The antibodies thus produced are useful not only as modulators of the association of Survivin with a Survivin binding partner, but are also useful in immunoassays for detecting Survivin expression/activity and for the purification of Survivin and associated binding partners.

#### [0074] C. Survivin Encoding Nucleic Acid Molecules

[0075] The present invention further provides nucleic acid molecules that encode Survivin, and the related Survivin proteins herein described, preferably in isolated form. For convenience, all Survivin encoding nucleic acid molecules will be referred to as the Survivin encoding nucleic acid molecule, the Survivin gene, or Survivin. As used herein, "nucleic acid" is defined as RNA or DNA that encodes a peptide as defined above, or is complementary to a nucleic acid sequence encoding such peptides, or hybridizes to such a nucleic acid and remains stably bound to it under stringent conditions, or encodes a polypeptide sharing at least 75% sequence identity, preferably at least 80%, and more preferably at least 85%, with the peptide sequences. Specifically contemplated are genomic DNA, cDNA, mRNA and antisense molecules, as well as nucleic acids based on an alternative backbone or including alternative bases whether derived from natural sources or synthesized. Such hybridizing or complementary nucleic acid, however, is defined further as being novel and unobvious over any prior art nucleic acid including that which encodes, hybridizes under appropriate stringency conditions, or is complementary to a nucleic acid encoding a Survivin protein according to the present invention.

[0076] As used herein, "stringent conditions" are conditions in which hybridization yields a clear and detectable sequence. Stringent conditions are those that (1) employ low ionic strength and high temperature for washing, for example, 0.015M NaCl/0.0015M sodium titrate/0.1% SDS at 50° C., or (2) employ during hybridization a denaturing agent such as formamide, for example, 50% (vol/vol) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH

6.5 with 750 mM NaCl, 75 mM sodium citrate at 42° C. Another example is use of 50% formamide, 5×SSC (0.75M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5×Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42° C., with washes at 42° C. in 0.2×SSC and 0.1% SDS. A skilled artisan can readily determine and vary the stringency conditions appropriately to obtain a clear and detectable hybridization signal.

**[0077]** As used herein, a nucleic acid molecule is said to be "isolated" when the nucleic acid molecule is substantially separated from contaminant nucleic acid encoding other polypeptides from the source of nucleic acid.

**[0078]** The present invention further provides fragments of the Survivin encoding nucleic acid molecule. As used herein, a fragment of a Survivin encoding nucleic acid molecule refers to a small portion of the entire protein encoding sequence. The size of the fragment will be determined by the intended use. For example, if the fragment is chosen so as to encode an active portion of the Survivin protein, such as the C-terminal  $\beta$  coils or the IAP motif, the fragment will need to be large enough to encode the functional region(s) of the Survivin protein. If the fragment is to be used as a nucleic acid probe or PCR primer, then the fragment length is chosen so as to obtain a relatively small number of false positives during probing/priming. **FIG. 1** identifies fragments of the Survivin gene that are particularly useful as selective hybridization probes or PCR primers.

**[0079]** Fragments of the Survivin encoding nucleic acid molecules of the present invention (i.e., synthetic oligonucleotides) that are used as probes or specific primers for the polymerase chain reaction (PCR), or to synthesize gene sequences encoding Survivin proteins can easily be synthesized by chemical techniques, for example, the phosphotriester method of Matteucci, et al., *J Am Chem Soc* (1981) 103:3185-3191 or using automated synthesis methods. In addition, larger DNA segments can readily be prepared by well known methods, such as synthesis of a group of oligonucleotides that define various modular segments of the Survivin gene, followed by ligation of oligonucleotides to build the complete modified Survivin gene.

**[0080]** The Survivin encoding nucleic acid molecules of the present invention may further be modified so as to contain a detectable label for diagnostic and probe purposes. As described above such probes can be used to identify other members of the Survivin family of proteins and as described below, such probes can be used to detect Survivin expression and tumor growth potential. A variety of such labels are known in the art and can readily be employed with the Survivin encoding molecules herein described. Suitable labels include, but are not limited to, biotin, radiolabeled nucleotides and the like. A skilled artisan can employ any of the art known labels to obtain a labeled Survivin encoding nucleic acid molecule.

**[0081]** Since the Survivin gene is an antisense or reverse orientation of the EPR-1 gene, particularly preferred are single-stranded probes for use in diagnostic purposes. Specifically, single-stranded diagnostic probes can be used to selectively hybridize to mRNA that encodes Survivin. Single-stranded probes can be generated using known methods in which one strand of a double-stranded probe is isolated or in which a single stranded RNA probe is generated.

**[0082]** Modifications to the primary structure itself by deletion, addition, or alteration of the amino acids incorporated into the protein sequence during translation can be made without destroying the activity of the protein. Such substitutions or other alterations result in proteins having an amino acid sequence encoded by DNA falling within the contemplated scope of the present invention.

#### **[0083]** D. Isolation of Other Survivin Encoding Nucleic Acid Molecules

**[0084]** As described above, the identification of the human Survivin encoding nucleic acid molecule allows a skilled artisan to isolate nucleic acid molecules that encode other members of the Survivin family of proteins in addition to the human sequence herein described.

**[0085]** Essentially, a skilled artisan can readily use the amino acid sequence of Survivin to generate antibody probes to screen expression libraries prepared from cells. Typically, polyclonal antiserum from mammals such as rabbits immunized with the purified Survivin protein (as described below) or monoclonal antibodies can be used to probe a mammalian cDNA or genomic expression library, such as lambda gtl1 library, to obtain the appropriate coding sequence for Survivin, or other members of the Survivin family of proteins. The cloned cDNA sequence can be expressed as a fusion protein, expressed directly using its own control sequences, or expressed by constructions using control sequences appropriate to the particular host used for expression of the enzyme. **FIG. 1** identifies important antigenic and/or putative operative domains found in the Survivin protein sequence. Such regions are preferred sources of antigenic portions of the Survivin protein for the production of probe, diagnostic, and therapeutic antibodies.

**[0086]** Alternatively, a portion of the Survivin encoding sequence herein described can be synthesized and used as a probe to retrieve DNA encoding a member of the Survivin family of proteins from any mammalian organisms that contains such a protein. Oligomers containing approximately 18-20 nucleotides (encoding about a 6-7 amino acid stretch) are prepared and used to screen genomic DNA or cDNA libraries to obtain hybridization under stringent conditions or conditions of sufficient stringency to eliminate an undue level of false positives.

**[0087]** Additionally, pairs of oligonucleotide primers can be prepared for use in a polymerase chain reaction (PCR) to selectively clone a Survivin-encoding nucleic acid molecule. A PCR denature/anneal/extend cycle for using such PCR primers is well known in the art and can readily be adapted for use in isolating other Survivin encoding nucleic acid molecules. **FIG. 1** identifies regions of the human Survivin gene that are particularly well suited for use as a probe or as primers.

#### **[0088]** E. rDNA Molecules Containing a Survivin Encoding Nucleic Acid Molecule

**[0089]** The present invention further provides recombinant DNA molecules (rDNAs) that contain a Survivin encoding sequence. As used herein, a rDNA molecule is a DNA molecule that has been subjected to molecular manipulation in vitro. Methods for generating rDNA molecules are well known in the art, for example, see Sambrook et al., *Molecular Cloning* (1989). In the preferred rDNA mol-

ecules, a Survivin encoding DNA sequence is operably linked to expression control sequences and/or vector sequences.

**[0090]** The choice of vector and/or expression control sequences to which one of the Survivin encoding sequences of the present invention is operably linked depends directly, as is well known in the art, on the functional properties desired, e.g., protein expression, and the host cell to be transformed. A vector contemplated by the present invention is at least capable of directing the replication or insertion into the host chromosome, and preferably also expression, of the Survivin gene included in the rDNA molecule.

**[0091]** Expression control elements that are used for regulating the expression of an operably linked protein encoding sequence are known in the art and include, but are not limited to, inducible promoters, constitutive promoters, secretion signals, and other regulatory elements. Preferably, the inducible promoter is readily controlled, such as being responsive to a nutrient in the host cell's medium.

**[0092]** In one embodiment, the vector containing a Survivin encoding nucleic acid molecule will include a prokaryotic replicon, i.e., a DNA sequence having the ability to direct autonomous replication and maintenance of the recombinant DNA molecule extrachromosomally in a prokaryotic host cell, such as a bacterial host cell, transformed therewith. Such replicons are well known in the art. In addition, vectors that include a prokaryotic replicon may also include a gene whose expression confers a detectable marker such as a drug resistance. Typical bacterial drug resistance genes are those that confer resistance to ampicillin or tetracycline.

**[0093]** Vectors that include a prokaryotic replicon can further include a prokaryotic or viral promoter capable of directing the expression (transcription and translation) of the Survivin encoding gene sequences in a bacterial host cell, such as *E. coli*. A promoter is an expression control element formed by a DNA sequence that permits binding of RNA polymerase and transcription to occur. Promoter sequences compatible with bacterial hosts are typically provided in plasmid vectors containing convenient restriction sites for insertion of a DNA segment of the present invention. Typical of such vector plasmids are pUC8, pUC9, pBR322 and pBR329 available from Biorad Laboratories, (Richmond, Calif.), pPL and pKK223 available from Pharmacia, Piscataway, N.J.

**[0094]** Expression vectors compatible with eukaryotic cells, preferably those compatible with vertebrate cells, can also be used to form rDNA molecules that contain a Survivin encoding sequence. Eukaryotic cell expression vectors are well known in the art and are available from several commercial sources. Typically, such vectors are provided containing convenient restriction sites for insertion of the desired DNA segment. Typical of such vectors are PSVL and pKSV-10 (Pharmacia), pBPV-1/pML2d (International Biotechnologies, Inc.), pTDT1 (ATCC, #31255), the vector pCDM8 described herein, and the like eukaryotic expression vectors.

**[0095]** Eukaryotic cell expression vectors used to construct the rDNA molecules of the present invention may further include a selectable marker that is effective in an eukaryotic cell, preferably a drug resistance selection

marker. A preferred drug resistance marker is the gene whose expression results in neomycin resistance, i.e., the neomycin phosphotransferase (neo) gene. Southern et al., *J Mol Anal Genet* (1982) 1:327-341. Alternatively, the selectable marker can be present on a separate plasmid, and the two vectors are introduced by co-transfection of the host cell, and selected by culturing in the appropriate drug for the selectable marker.

**[0096]** F. Host Cells Containing an Exogenously Supplied Survivin Encoding Nucleic Acid Molecule

**[0097]** The present invention further provides host cells transformed with a nucleic acid molecule that encodes a Survivin protein of the present invention. The host cell can be either prokaryotic or eukaryotic. Eukaryotic cells useful for expression of a Survivin protein are not limited, so long as the cell line is compatible with cell culture methods and compatible with the propagation of the expression vector and expression of the Survivin gene product. Preferred eukaryotic host cells include, but are not limited to, yeast, insect and mammalian cells, preferably vertebrate cells such as those from a mouse, rat, monkey or human fibroblastic cell line, the most preferred being cells that do not naturally express a Survivin protein. Preferred eukaryotic host cells include the murine IL-3 dependent cell line BaF3, and the like eukaryotic tissue culture cell lines.

**[0098]** Any prokaryotic host can be used to express a Survivin-encoding rDNA molecule. The preferred prokaryotic host is *E. coli*.

**[0099]** Transformation of appropriate cell hosts with a rDNA molecule of the present invention is accomplished by well known methods that typically depend on the type of vector used and host system employed. With regard to transformation of prokaryotic host cells, electroporation and salt treatment methods are typically employed, see, for example, Cohen et al., *Proc Natl Acad Sci USA* (1972) 69:2110; and Maniatis et al., *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1982). With regard to transformation of vertebrate cells with vectors containing rDNAs, electroporation, cationic lipid or salt treatment methods are typically employed, see, for example, Graham et al., *Virology* (1973) 52:456; Wigler et al., *Proc Natl Acad Sci USA* (1979) 76:1373-76.

**[0100]** Successfully transformed cells, i.e., cells that contain a rDNA molecule of the present invention, can be identified by well known techniques. For example, cells resulting from the introduction of an rDNA of the present invention can be cloned to produce single colonies. Cells from those colonies can be harvested, lysed and their DNA content examined for the presence of the rDNA using a method such as that described by Southern, *J Mol Biol* (1975) 98:503, or Berent et al., *Biotech* (1985) 3:208 or the proteins produced from the cell assayed via an immunological method.

**[0101]** G. Production of Survivin Using a rDNA Molecule Encoding a Survivin Protein

**[0102]** The present invention further provides methods for producing a Survivin protein that uses one of the Survivin encoding nucleic acid molecules herein described. In general terms, the production of a recombinant form of a Survivin protein typically involves the following steps.

[0103] First, a nucleic acid molecule is obtained that encodes a Survivin protein, such as the nucleic acid molecule depicted in **FIG. 1**. If the Survivin encoding sequence is uninterrupted by introns, it is directly suitable for expression in any host. If not, then a spliced form of the Survivin encoding nucleic acid molecule can be generated and used or the intron containing nucleic acid molecule can be used in a compatible eukaryotic expression system.

[0104] The Survivin encoding nucleic acid molecule is then preferably placed in operable linkage with suitable control sequences, as described above, to form an expression unit containing the Survivin encoding sequences. The expression unit is used to transform a suitable host and the transformed host is cultured under conditions that allow the production of the Survivin protein. Optionally the Survivin protein is isolated from the medium or from the cells; recovery and purification of the protein may not be necessary in some instances where some impurities may be tolerated.

[0105] Each of the foregoing steps can be done in a variety of ways. For example, the desired coding sequences may be obtained from genomic fragments and used directly in appropriate hosts. The construction of expression vectors that are operable in a variety of hosts is accomplished using appropriate replicons and control sequences, as set forth above. The control sequences, expression vectors, and transformation methods are dependent on the type of host cell used to express the gene and were discussed in detail earlier. Suitable restriction sites can, if not normally available, be added to the ends of the coding sequence so as to provide an excisable gene to insert into these vectors. A skilled artisan can readily adapt any host/expression system known in the art for use with Survivin encoding sequences to produce a Survivin protein.

#### [0106] H. Inhibition of Cell Death Using Survivin

[0107] As provided above, Survivin has been shown to inhibit cellular apoptosis. Accordingly, Survivin can be used in methods to extend the life of cells. In general, cellular apoptosis can be inhibited by contacting a cell with Survivin.

[0108] There are a number of situations in which it is desirable to inhibit cellular apoptosis. For example, the death of cells in tissues and organs being prepared for transport and transplant can be inhibited using the Survivin protein. Alternatively, cell lines can be established for long term culture using Survivin encoding nucleic acid molecules expressed in the cell line.

[0109] Hence, Survivin protein or Survivin gene expression can be used as a means to inhibit cellular apoptosis. In cell culture systems, the Survivin protein can be introduced into a cell, for example via liposomal, Penetrin-1 delivery, or inclusion in the cell growth media, to inhibit apoptosis. Alternatively, the Survivin gene can be introduced and expressed in cells to increase the longevity of cells in culture. These provide means and methods for increasing the ability of cultured cells to produce desired compounds as well as provide methods of establishing long-term culture of primary explants of cells and tissues.

[0110] In tissue transplant, typically tissues and organs are stored and transported prior to transplant. Cell death, by mechanisms similar to apoptosis, can lead to the loss of viability of the tissues or organs. In this setting, infusion

with Survivin protein can be used as a method to inhibit cell death in such tissues and organs.

[0111] There are pathological conditions characterized by premature and unwanted cellular apoptosis, for example in accelerated aging disorders. It is already known that inactivating mutations in a IAP protein may cause human diseases. The example is for the NAIP (see above). Studies of patients with SMA (Spinal muscular atrophy, a neurodegenerative disease that is thought to be caused by aberrantly increased apoptosis) has demonstrated that the NAIP gene is inactivated and deleted in 75% of these patients (Roy et al. 1995, *Cell* 80:167). By extension, inactivating mutations in Survivin can result in degenerative diseases characterized by aberrantly increased cell death. Haplotypic markers within and around the Survivin locus on chromosome 17q25 can be used in studies of population genetics to determine if that locus has already been implicated in diseases with increased apoptosis. In such cases, the Survivin gene or the Survivin protein can be used to treat the conditions. Accordingly, the Survivin protein, or a Survivin encoding nucleic acid molecule is administered to an individual as a means of treating abnormal apoptosis.

#### [0112] I. Methods to Identify Survivin Binding Partners

[0113] Another embodiment of the present invention provides methods for use in isolating and identifying binding partners of Survivin. Specifically, the Survivin protein can be used as a capture probe to identify Survivin binding partners. As used herein, a Survivin binding partner is a biomolecule (such as a protein, DNA or other cofactor) that binds to Survivin and mediates Survivin inhibition of cellular apoptosis.

[0114] In detail, a Survivin protein is mixed with an extract or fraction of a cell that expresses Survivin under conditions that allow the association of a binding partner with Survivin. After mixing, peptides that have become associated with Survivin are separated from the mixture. The binding partner that bound Survivin can then be removed and further analyzed.

[0115] To identify and isolate a binding partner, the entire Survivin protein can be used. Alternatively, a fragment of a Survivin protein can be used.

[0116] As used herein, a cellular extract refers to a preparation or fraction that is made from a lysed or disrupted cell. The preferred source of cellular extracts will be cells that naturally express Survivin. Examples of such cells include, but are not limited to tumor cells and embryonic tissues.

[0117] A variety of methods can be used to obtain an extract of a cell. Cells can be disrupted using either physical or chemical disruption methods. Examples of physical disruption methods include, but are not limited to, sonication and mechanical shearing. Examples of chemical lysis methods include, but are not limited to, detergent lysis and the enzyme lysis. In addition, the cellular extract can be prepared from cells that have been freshly isolated from a subject or from cells or cell lines which have been cultured. A skilled artisan can readily adapt methods for preparing cellular extracts in order to obtain extracts for use in the present methods.

[0118] Once an extract of a cell is prepared, the extract is mixed with the Survivin protein under conditions in which

association of Survivin with the binding partner can occur. A variety of conditions can be used, the most preferred being conditions that closely resemble conditions found in the cytoplasm of a Survivin-expressing cell. Features such as osmolarity, pH, temperature, and the concentration of cellular extract used, can be varied to optimize the association of the Survivin with the binding partner.

[0119] After mixing under appropriate conditions, Survivin is separated from the mixture. A variety of techniques can be utilized to separate the mixture. For example, antibodies specific to Survivin can be used to immunoprecipitate the Survivin and associated binding partner. Alternatively, standard chemical separation techniques such as chromatography and density/sediment centrifugation can be used.

[0120] After removal of nonassociated cellular constituents found in the extract, the binding partner can be dissociated from the Survivin protein using conventional methods. For example, dissociation can be accomplished by altering the salt concentration or pH of the mixture.

[0121] To aid in separating associated Survivin/binding partner pairs from the mixed extract, the Survivin protein can be immobilized on a solid support. For example, Survivin can be attached to a nitrocellulose matrix or acrylic beads. Attachment of Survivin to a solid support further aids in separating peptide/binding partner pair from other constituents found in the extract.

[0122] Alternatively, the Survivin-encoding nucleic acid molecule can be used in a yeast two-hybrid system. The yeast two-hybrid system has been used to identify other protein partner pairs and can readily be adapted to employ the Survivin encoding molecules herein described.

#### [0123] J. Use of Survivin Binding Partners

[0124] Once isolated, the Survivin binding partners obtained using the above described methods can be used for a variety of purposes. The binding partners can be used to generate antibodies that bind to the Survivin binding partner using techniques known in the art. Antibodies that bind a Survivin binding partner can be used to assay Survivin activity, as a therapeutic agent to modulate a biological or pathological process mediated by Survivin, or to purify the binding partner. These uses are described in detail below.

#### [0125] K. Methods to Identify Agents that Block Survivin/Binding Partner Interactions

[0126] Another embodiment of the present invention provides methods for identifying agents that reduce or block the association of Survivin with a Survivin binding partner. Specifically, Survivin is mixed with a Survivin binding partner in the presence and absence of an agent to be tested. After mixing under conditions that allow association of Survivin with the Survivin binding partner, the two mixtures are analyzed and compared to determine if the agent reduced or blocked the association of Survivin with the Survivin binding partner. Agents that block or reduce the association of Survivin with the Survivin binding partner will be identified as decreasing the amount of association present in the sample containing the tested agent.

[0127] As used herein, an agent is said to reduce or block Survivin/Survivin binding partner association when the presence of the agent decreases the extent to which or prevents the Survivin binding partner from becoming asso-

ciated with Survivin. One class of agents will reduce or block the association by binding to the Survivin binding partner while another class of agents will reduce or block the association by binding to Survivin.

[0128] The Survivin binding partner used in the above assay can either be an isolated and fully characterized protein or can be a partially characterized protein that binds to Survivin or a Survivin binding partner that has been identified as being present in a cellular extract. It will be apparent to one of ordinary skill in the art that so long as the Survivin binding partner has been characterized by an identifiable property, e.g., molecular weight, the present assay can be used.

[0129] Agents that are assayed in the above method can be randomly selected or rationally selected or designed. As used herein, an agent is said to be randomly selected when the agent is chosen randomly without considering the specific sequences involved in the association of the Survivin with the Survivin binding partner. An example of randomly selected agents is the use a chemical library or a peptide combinatorial library, or a growth broth of an organism.

[0130] As used herein, an agent is said to be rationally selected or designed when the agent is chosen on a nonrandom basis which takes into account the sequence of the target site and/or its conformation in connection with the agent's action. As described above, there are two sites of action for agents that block Survivin/Survivin binding partner interaction: the binding partner contact site on Survivin and the Survivin contact site on the Survivin binding partner. Agents can be rationally selected or rationally designed by utilizing the peptide sequences that make up the contact sites of the Survivin/Survivin binding partner pair. For example, a rationally selected peptide agent can be a peptide whose amino acid sequence is identical to the Survivin contact site on the Survivin binding partner. Such an agent will reduce or block the association of Survivin with the binding partner by binding to the Survivin binding partner.

[0131] The agents of the present invention can be, as examples, peptides, small molecules, vitamin derivatives, as well as carbohydrates. A skilled artisan can readily recognize that there is no limit as to the structural nature of the agents of the present invention. One class of agents of the present invention are peptide agents whose amino acid sequences are chosen based on the amino acid sequence of the Survivin protein.

[0132] The peptide agents of the invention can be prepared using standard solid phase (or solution phase) peptide synthesis methods, as is known in the art. In addition, the DNA encoding these peptides may be synthesized using commercially available oligonucleotide synthesis instrumentation and produced recombinantly using standard recombinant production systems. The production using solid phase peptide synthesis is necessitated if non-gene-encoded amino acids are to be included.

[0133] Another class of agents of the present invention are antibodies immunoreactive with critical positions of the Survivin or Survivin binding partner. As described above, antibodies are obtained by immunization of suitable mammalian subjects with peptides, containing as antigenic regions, those portions of the Survivin or binding partner, intended to be targeted by the antibodies. Critical regions



include the contact sites involved in the association of the Survivin with the Survivin binding partner.

[0134] As discussed below, the important minimal sequence of residues involved in Survivin activity define a functional linear domain that can be effectively used as a bait for two-hybrid screening and identification of potential Survivin-associated molecules. Use of such Survivin fragments will significantly increase the specificity of the screening as opposed to using the full length molecule or the entire BIR domain and is therefore preferred. Similarly, this linear sequence can be also used as an affinity matrix also to isolate Survivin binding proteins using a biochemical affinity purification strategy.

[0135] L. Uses for Agents that Block the Association of Survivin with a Survivin Binding Partner

[0136] As provided in the Background section, Survivin inhibits cellular apoptosis. Agents that reduce or block the interactions of Survivin with a Survivin binding partner can be used to modulate biological and pathologic processes associated with Survivin function and activity.

[0137] In detail, a biological or pathological process mediated by Survivin can be modulated by administering to a subject an agent that blocks the interaction of Survivin with a Survivin binding partner.

[0138] As used herein, a subject can be any mammal, so long as the mammal is in need of modulation of a pathological or biological process mediated by Survivin. The term "mammal" is meant an individual belonging to the class Mammalia. The invention is particularly useful in the treatment of human subjects.

[0139] As used herein, a biological or pathological process mediated by Survivin or Survivin binding to a Survivin binding partner refers to the wide variety of cellular events mediated by Survivin. Pathological processes refer to a category of biological processes which produce a deleterious effect. For example, a pathological process mediated by Survivin is the inhibition of cellular apoptosis in tumor cells. This pathological process can be modulated using agents that reduce or block Survivin/Survivin binding partner association or block Survivin expression.

[0140] As used herein, an agent is said to modulate a pathological process when the agent reduces the degree or severity of the process. For example, an agent is said to modulate tumor cell proliferation when the agent decrease the rate or extent of cell division.

[0141] M. Administration of Survivin or Agents that Affect Survivin Activity

[0142] The agents of the present invention, whether they be agents that block Survivin/binding partner association or the Survivin protein, can be administered via parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, or buccal routes. Alternatively, or concurrently, administration may be by the oral route. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired. For example, to treat tumor cells as a means of blocking Survivin inhibition of apoptosis, an agent that blocks Survivin expression or the interaction of Survivin with a binding partner, is administered systemically or locally to the indi-

vidual being treated. As described below, there are many methods that can readily be adapted to administer such agents.

[0143] The present invention further provides compositions containing Survivin or one or more agents that block Survivin/binding partner association. While individual needs vary, a determination of optimal ranges of effective amounts of each component is within the skill of the art. Typical dosages comprise 0.1 to 100  $\mu\text{g/kg}$  body wt. The preferred dosages comprise 0.1 to 10  $\mu\text{g/kg}$  body wt. The most preferred dosages comprise 0.1 to 1  $\mu\text{g}$  body wt.

[0144] In addition to the pharmacologically active agent, the compositions of the present invention may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically for delivery to the site of action. Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water-soluble salts. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers. Liposomes can also be used to encapsulate the agent for delivery into the cell.

[0145] The pharmaceutical formulation for systemic administration according to the invention may be formulated for enteral, parenteral or topical administration. Indeed, all three types of formulations may be used simultaneously to achieve systemic administration of the active ingredient.

[0146] Suitable formulations for oral administration include hard or soft gelatin capsules, pills, tablets, including coated tablets, elixirs, suspensions, syrups or inhalations and controlled release forms thereof.

[0147] In practicing the methods of this invention, the compounds of this invention may be used alone or in combination, or in combination with other therapeutic or diagnostic agents. In certain preferred embodiments, the compounds of this invention may be coadministered along with other compounds typically prescribed for these conditions according to generally accepted medical practice, such as chemotherapeutic agents.

[0148] N. Combination Therapy

[0149] Survivin, as well as agents of the present invention that modulate Survivin activity, can be provided alone, or in combination with another agents that modulate a particular biological or pathological process. For example, an agent of the present invention that reduces Survivin inhibited apoptosis can be administered in combination with other anti-cancer agents in methods to control cancer cell growth. Alternatively, Survivin can be administered with other protective agents as a means for reducing cellular apoptosis. As used herein, two agents are said to be administered in combination when the two agents are administered simultaneously or are administered independently in a fashion such that the compounds will act at the same time.

**[0150]** Inhibition of Survivin activity/expression can be used in combination with conventional chemotherapies. The timing for using a chemotherapeutic agent in combination with inhibiting Survivin activity/expression depends upon chemotherapeutic agent used and the tumor cell type treated. Examples of chemotherapeutic agents that can be used in combination with agents the effect Survivin activity/expression, includes, but is not limited to alkylating agents, such as cyclophosphamide (CTX; cytoxan), chlorambucil (CHL; leukan), cisplatin (CisP; platinol) busulfan (myleran), melphalan, carmustine (BCNU), streptozotocin, triethylenemelamine (TEM), mitomycin C., and the like alkylating agents; anti-metabolites, such as methotrexate (MTX), etoposide (VP16; vepesid) 6-mercaptopurine (6MP), 6-thioguanine (6TG), cytarabine (Ara-C), 5-fluorouracil (5FU), dacarbazine DTIC, and the like anti-metabolites; antibiotics, such as actinomycin D, doxorubicin (DXR; adriamycin), daunorubicin (daunomycin), bleomycin, mitramycin and the like antibiotics; alkaloids, such as vinca alkaloids such as vincristine (VCR), vinblastine, and the like; and other antitumor agents, such as taxol and taxol derivatives, the cytostatic agents glucocorticoids such as dexamethasone (DEX; decadron) and corticosteroids such as prednisone, nucleoside enzyme inhibitors such as hydroxyurea, amino acid depleting enzymes such as asparaginase, and the like diverse antitumor agents.

**[0151]** The use of the cytotoxic agents described above in chemotherapeutic regimens is generally well characterized in the cancer therapy arts, and their use herein falls under the same considerations for monitoring tolerance and effectiveness and for controlling administration routes and dosages, with some adjustments. For example, the actual dosages of the cytotoxic agents may vary depending upon the patient's cultured cell response determined by using the present histoculture methods. Generally, the dosage will be reduced compared to the amount used in the absence of agents the effect Survivin activity/expression.

**[0152]** Typical dosages of an effective cytotoxic agent can be in the ranges recommended by the manufacturer, and where indicated by in vitro responses or responses in animal models, can be reduced by up to about one order of magnitude concentration or amount. Thus, the actual dosage will depend upon the judgment of the physician, the condition of the patient, and the effectiveness of the therapeutic method based on the in vitro responsiveness of the primary cultured malignant cells or histocultured tissue sample, or the responses observed in the appropriate animal models.

**[0153]** O. Methods for Identifying Survivin Expression and Survivin-Mediated Inhibition of Apoptosis

**[0154]** The present invention further provides methods for identifying cells involved in Survivin-mediated inhibition of apoptosis as well as techniques that can be applied to diagnose biological and pathological processes associated with Survivin activity, the progression of such conditions, the susceptibility of such conditions to treatment and the effectiveness of treatment for such conditions. Specifically, Survivin-mediated inhibition of apoptosis can be identified by determining whether the Survivin protein is expressed in a cell. Cells expressing Survivin are considered to be inhibited from natural cellular apoptosis.

**[0155]** A variety of immunological and nucleic acid techniques can be used to determine if the Survivin protein, or

a Survivin encoding mRNA, is produced in a particular cell. In one example, an extract of cells is prepared. The extract is then assayed to determine whether Survivin is expressed in the cell. The degree of expression provides a measurement of the degree of inhibition of apoptosis. An increase in expression is a measurement of an increased inhibition of apoptosis.

**[0156]** The measurement of Survivin expression can be used as a marker for a variety of purposes. In tumors, the present of Survivin expression correlates with the proliferative potential of the tumor. In the Examples, it is shown that lymphomas display varying levels of Survivin expression; lymphomas showing little to no Survivin expression are low grade lymphomas that can be effectively treated while lymphomas showing high levels of Survivin expression are high grade aggressive lymphomas that typically cannot be effectively treated. Accordingly, the level of Survivin expression in a lymphoma, or other tumor, can be used as a predictive measurement of the aggressiveness and treatability of the tumor: the higher the level of Survivin expression, the higher the aggressiveness of the tumor and the more difficult the treatment will be.

**[0157]** For example, to determine a tumor's proliferative potential or easy/prognosis of treatment, an extract is made of the tumor cells and the extract is then analyzed, for example, by gel electrophoresis, to determine whether a Survivin protein is present. The presence and level of Survivin correlates with the proliferative potential of the cancer and the ease of treatment. Alternatively, as described above, single-strand probes can be used to identify Survivin-encoding mRNA in the cellular extracts.

**[0158]** In addition to being a marker of tumor aggressiveness and treatment potential, Survivin expression can be used as a measurement of the effectiveness of anti-tumor therapy. In the Examples, it is shown that HL-60, a promyelocytic cell line, had high levels of Survivin expression. Treatment of HL-60 cells with retenoic acid, and anti-cancer agent that acts by causing the differentiation of tumor cells, resulted in a reduction and elimination of Survivin expression. The reduction in expression correlated with the degree of differentiation, the greater the differentiation, the lower the level of Survivin expression. Accordingly, Survivin expression can be used to measure the effectiveness of anti-tumor treatment: if Survivin expression decreases during treatment, the treatment protocol is effective and can be continued, whereas if Survivin expression remains unaltered, a different therapeutic regime or protocol needs to be performed.

**[0159]** P. Other Methods to Control Survivin Expression

**[0160]** The present invention further provides additional methods that can be used to control Survivin expression in a cell. As discussed above and below, the Survivin promoter has a CPG island upstream from its promoter. CPG islands are known targets for DNA methylation. The DNA methylation sites in the CPG island serves as a means for regulating Survivin expression: methylation of CPG islands results in the suppression of transcription of the gene found downstream from the promoter. Accordingly, agents that methylate DNA, such as DNA methylase, and agents that stimulate the production of endogenous methylases, can be used to control Survivin expression. Specifically, Survivin expression in a cell can be reduced or eliminated by causing

the cell to increase the level of DNA methylation, particularly at the CPG island found upstream from the Survivin gene.

[0161] In another method, Survivin expression can be reduced by increasing the level of EPR-1 expression. As shown in the Examples, Survivin expression and EPR-1 expression are generally mutually exclusive, expression of EPR-1 results in a decrease or elimination of Survivin expression and visa-a-versa. Accordingly, Survivin expression can be reduced by causing a cell to increase EPR-1 expression.

#### [0162] Q. Animal Models

[0163] We have isolated almost the complete structure of the mouse Survivin gene. The gene is very conserved with its human counterpart including sizes of introns, exons and intron-exon boundaries. The coding regions of the mouse Survivin gene are 88%, to the extent sequenced, identical to the human protein, thereby demonstrating strong evolutionary conservation. We have also determined the differential and developmentally-regulated distribution of Survivin during both human and mouse development. The availability of the complete structure of the mouse Survivin gene and protein will allow the preparation of targeting vectors for gene knockout experiments and a more rational approach for the generation of transgenic mice expressing Survivin under the control of tissue-specific promoters.

[0164] The Survivin gene and the Survivin protein can serve as a target for gene therapy in a variety of contexts. For example, in one application, Survivin-deficient non-human animals can be generated using standard knock-out procedures to inactivate a Survivin gene or, if such animals are non-viable, inducible Survivin antisense molecules can be used to regulate Survivin activity/expression. Alternatively, an animal can be altered so as to contain a Survivin or antisense-Survivin expression unit that directs the expression of Survivin or the antisense molecule in a tissue specific fashion. In such a uses, a non-human mammal, for example a mouse or a rat, is generated in which the expression of the Survivin gene is altered by inactivated or activation. This can be accomplished using a variety of art-known procedures such as targeted recombination. Once generated, the Survivin-deficient animal, the animal that expresses Survivin in a tissue specific manner, or an animal that expresses an antisense molecule can be used to 1) identify biological and pathological processes mediated by Survivin, 2) identify proteins and other genes that interact with Survivin, 3) identify agents that can be exogenously supplied to overcome Survivin deficiency and 4) serve as an appropriate screen for identifying mutations within Survivin that increase or decrease activity.

[0165] For example, it is possible to generate transgenic mice expressing the human minigene for Survivin in a tissue specific-fashion and test the effect of over-expression of the protein in district that normally do not contain Survivin. This strategy has been successfully used for another family of apoptosis inhibitors, namely bcl-2 (Veis et al., *Cell* (1993) 75:229). Such an approach can readily be applied to the Survivin protein and can be used to address the issue of a potential beneficial effect of Survivin in a specific tissue area to protect cells from apoptosis (transplant).

#### [0166] R. Survivin Gene Therapy

[0167] In another embodiment, genetic therapy can be used as a means for modulating a Survivin-mediated biological or pathological processes. For example, in tumor therapy, it may be desirable to introduce into the subject being treated a genetic expression unit that encodes a modulator of Survivin expression, such as an antisense encoding nucleic acid molecule. Such a modulator can either be constitutively produced or inducible within a cell or specific target cell. This allows a continual or inducible supply of a modulator of Survivin expression within the subject. Blocking Survivin expression allows for the control of tumor cell growth. Similarly, cells may be genetically engineered to express Survivin, e.g., in allograft pancreatic  $\beta$  cells for transplantation.

[0168] The level of Survivin gene expression may correlate with the level of resistance to apoptosis. Thus, Survivin genes also find use in anti-apoptosis gene therapy. In particular, a functional Survivin gene may be used to sustain neuronal cells that undergo apoptosis in the course of a neurodegenerative disease, lymphocytes (i.e., T cells and B cells), or cells that have been injured by ischemia.

[0169] Retroviral vectors, adenoviral vectors, adeno-associated viral vectors, or other viral vectors with the appropriate tropism for cells likely to be involved in apoptosis (for example, epithelial cells) may be used as a gene transfer delivery system for a therapeutic Survivin gene construct. Numerous vectors useful for this purpose are generally known (Miller, *Human Gene Therapy* 15-14, 1990; Friedman, *Science* 244:1275-1281, 1989; Eglitis and Anderson, *Bio Techniques* 6:608-614, 1988; Tolstoshev and Anderson, current opinion in biotechnology 1:55-61, 1990; Sharp, *The Lancet* 337:1277-1278, 1991; Cornetta et al., *Nucleic Acid Research and Molecular Biology* 36:311-322, 1987; Anderson, *Science* 226:401-409, 1984; Moen, *blood Cells* 17:407-416, 1991; Miller et al., *Biotechniques* 7:980-990, 1989; Le Gal La Salle et al., *Science* 259:988-990, 1993; and Johnson, *Chest* 107:77S-83S, 1995). Retroviral vectors are particularly well developed and have been used in clinical settings (Rosenberg et al., *N. Engl. J. Med* 323:370, 1990; Anderson et al., U.S. Pat. No. 5,399,346).

[0170] Non-viral approaches may also be employed for the introduction of therapeutic DNA into cells otherwise predicted to undergo apoptosis. For example, Survivin may be introduced into a neuron or a T cell by lipofection (Feigner et al., *Proc. Natl. Acad. Sci. USA* 84:7413, 1987; Ono et al., *Neurosci. Lett.* 117:259, 1990; Brigham et al., *Meth. Enz.* 101:512, 1983), asialorosonucoid-polylysine conjugation (Wu et al., *J. Biol. Chem.* 263:14621, 1988; Wu et al., *J. Biol. Chem.* 264:16985, 1989); or, less preferably, microinjection under surgical conditions (Wolff et al., *Science* 247:1465, 1990).

[0171] For any of the methods of application described above, the therapeutic Survivin nucleic acid construct is preferably applied to the site of the predicted apoptosis event (for example, by injection). However, it may also be applied to tissue in the vicinity of the predicted apoptosis event or to a blood vessel supplying the cells predicted to undergo apoptosis.

[0172] In the constructs described, Survivin cDNA expression can be directed from any suitable promoter (e.g., the

human cytomegalovirus (CMV), simian virus 40 (SV40), or metallothionein promoters), and regulated by any appropriate mammalian regulatory element. For example, if desired, enhancers known to preferentially direct gene expression in neural cells, T cells, or B cells may be used to direct Survivin expression. The enhancers used could include, without limitation, those that are characterized as tissue- or cell-specific in their expression. Alternatively, if a Survivin genomic clone is used as a therapeutic construct (for example, following its isolation by hybridization with the Survivin cDNA described above), regulation may be mediated by the cognate regulatory sequences or, if desired, by regulatory sequences derived from a heterologous source, including any of the promoters or regulatory elements described above.

**[0173] S. Use of the Survivin Promoter to Direct Gene Expression**

**[0174]** The present invention further provides the promoter of the Survivin gene in a form that can be used in generating expression vectors. Specifically, the Survivin promoter, identified as being 5' from the ATG start codon in of Survivin, can be used to direct the expression of an operably linked protein encoding DNA sequence. Since the Survivin promoter does not have a TATA box, a skilled artisan would use a 5' fragment, such as nucleotides 2560-2920 (including exon 1). The Survivin promoter is expressed in fetal tissues and can therefore be used to target protein expression in specific cell types during specific stages of development. As discussed below, transfection of 3T3 cells with the c-myc oncogene results in the up-regulation of Survivin mRNA as detected by Northern blots. Accordingly, DNA encoding anti-tumor polypeptides under the control of the Survivin promoter could be used to transfect tumor cell where they would be expressed. A skilled artisan can readily use the Survivin promoter in expression vectors using methods known in the art.

**[0175] T. Preventative Anti-Apoptotic Therapy**

**[0176]** In a patient diagnosed to be heterozygous for a Survivin mutation or to be susceptible to Survivin mutations (even if those mutations do not yet result in alteration or loss of Survivin biological activity), or a patient diagnosed with a degenerative disease (e.g., motor neuron degenerative diseases such as SMA or ALS diseases), or diagnosed as HIV positive, any of the disclosed therapies may be administered before the occurrence of the disease phenotype. For example, the therapies may be provided to a patient who is HIV positive but does not yet show a diminished T cell count or other overt signs of AIDS. In particular, compounds shown to increase Survivin expression or Survivin biological activity may be administered by any standard dosage and route of administration. Alternatively, gene therapy using a Survivin expression construct may be undertaken to reverse or prevent the cell defect prior to the development of the degenerative disease.

**[0177]** The methods of the instant invention may be used to reduce or diagnose the disorders described herein in any mammal, for example, humans, domestic pets, or livestock. Where a non-human mammal is treated or diagnosed, the Survivin polypeptide, nucleic acid, or antibody employed is preferably specific for that species.

**[0178] U. Examples of Additional Apoptosis Assays**

**[0179]** In addition to the foregoing discussion, specific examples of apoptosis assays are also provided in the following references. Assays for apoptosis in lymphocytes are disclosed by: Li et al., "Induction of apoptosis in uninfected lymphocytes by HIV-1 Tat protein", *Science* 268:429-431, 1995; Gibellini et al., "Tat-expressing Jurkat cells show an increased resistance to different apoptotic stimuli, including acute human immunodeficiency virus-type 1 (HIV-1) infection", *Br. J. Haematol.* 89:24-33, 1995; Martin et al., "HIV-1 infection of human CD4<sup>+</sup> T cells in vitro. Differential induction of apoptosis in these cells." *J. Immunol.* 152:330-42, 1994; Terai et al., "Apoptosis as a mechanism of cell death in cultured T lymphoblasts acutely infected with HIV-1", *J. Clin. Invest.* 87:1710-5, 1991; Dhein et al., "Autocrine T-cell suicide mediated by APO-1/(Fas/CD95)11, *Nature* 373:438-441, 1995; Katsikis et al., "Fas antigen stimulation induces marked apoptosis of T lymphocytes in human immunodeficiency virus-infected individuals", *J. Exp. Med.* 181:2029-2036, 1995; Westendorp et al., "Sensitization of T cells to CD95-mediated apoptosis by HIV-1 Tat and gp120", *Nature* 375:497, 1995; DeRossi et al., *Virology* 198:234-44, 1994.

**[0180]** Assays for apoptosis in fibroblasts are disclosed by: Vossbeck et al., "Direct transforming activity of TGF-beta on rat fibroblasts", *Int. J. Cancer* 61:92-97, 1995; Goruppi et al., "Dissection of c-myc domains involved in S phase induction of HIH3T3 fibroblasts", *Oncogene* 9:1537-44, 1994; Fernandez et al., "Differential sensitivity of normal and Ha-ras transformed C3H mouse embryo fibroblasts tumor necrosis factor; induction of bcl-2, c-myc, and manganese superoxide dismutase in resistant cells", *Oncogene* 9:2009-17, 1994; Harrington et al., "c Myc-induced apoptosis in fibroblasts is inhibited by specific cytokines", *EMBO J.*, 13:3286-3295, 1994; Itoh et al., "A novel protein domain required for apoptosis. Mutational analysis of human Fas antigen", *J. Biol. Chem.* 268:10932-7, 1993.

**[0181]** Assays for apoptosis in neuronal cells are disclosed by: Melino et al., "Tissue transglutaminase and apoptosis: sense and antisense transfection studies with human neuroblastoma cells", *Mol. Cell Biol.* 14:6584-6596, 1994; Rosenblum et al., "Evidence for hypoxia-induced, programmed cell death of cultured neurons", *Ann. Neurol.* 36:864-870, 1994; Sato et al., "Neuronal differentiation of PC12 cells as a result of prevention of cell death by bcl-2", *J. Neurobiol.* 25:1227-1234, 1994; Ferrari et al., "N-acetylcysteine D- and L-stereoisomers prevents apoptotic death of neuronal cells", *J. Neurosci.* 15:2857-2866, 1995; Talley et al., "Tumor necrosis factor alpha-induced apoptosis in human neuronal cells: protection by the antioxidant N-acetylcysteine and the genes bcl-2 and crma", *Mol. Cell Biol.* 15:2359-2366, 1995; Talley et al., "Tumor Necrosis Factor Alpha-Induced Apoptosis in Human Neuronal Cells: Protection by the Antioxidant N-Acetylcysteine and the Genes bcl-2 and crma", *Mol. Cell. Biol.* 15:2359-2366, 1995; Walkinshaw et al., "Induction of apoptosis in catecholaminergic PC12 cells by L-DOPA. Implications for the treatment of Parkinson's disease," *J. Clin. Invest.* 95:2458-2464, 1995.

**[0182]** Assays for apoptosis in insect cells are disclosed by: Clem et al., "Prevention of apoptosis by a baculovirus gene during infection on insect cells", *Science* 254:1388-90, 1991; Crook et al., "An apoptosis-inhibiting baculovirus gene with a zinc finger-like motif", *J. Virol.* 67:2168-74,

1993; Rabizadeh et al., "Expression of the baculovirus p35 gene inhibits mammalian neural cell death", *J. Neurochem.* 61:2318-21, 1993; Birnbaum et al., "An apoptosis inhibiting gene from a nuclear polyhedrosis virus encoding a polypeptide with Cys/His sequence motifs", *J. Virol.* 68:2521-8, 1994; Clem et al., *Mol. Cell. Biol.* 14:5212-5222, 1994.

**[0183]** V. Use of Survivin in Tissue and Organ Transplantation The present invention includes methods of inhibiting or preventing tissue or organ transplant rejection in a subject, comprising the local administration of a Survivin polypeptide, Survivin polypeptide fragment, an apoptosis-, inhibiting peptidomimetic thereof, a transgene encoding a Survivin polypeptide or a transgene encoding a Survivin polypeptide fragment to the tissue, organ or to a site proximal to the transplant. Local delivery of the polypeptides, peptidomimetics to the tissue, organ or to a site proximal to the transplant is accomplished by any means commonly available, including but not limited to direct local perfusion, injection, microsponges, microcapsules, liposomes or time-released delivery vehicles.

**[0184]** Local delivery of a transgene encoding a Survivin polypeptide or a transgene encoding a Survivin polypeptide fragment to the tissue, organ or to a site proximal to the transplant may be accomplished with any available vector, via lipofection or via direct plasmid DNA injection. See Qin et al. (1995) *Transplantation* 59(6): 809-816; Le Coultré et al. (1997) *Eur. J. Pediatr. Surg.* 7(4):221-226; Wang et al. (1992) *Transplantation* 53(3):703-705; Wang et al. (1996) *Transplantation* 61(12):1726-1729; Schmid et al., (1997) *Eur. J. Cardiothorac. Surg.* 11(6):1023-28; and Boasquevisque, C. et al. (1997) *Ann. Thorac. Surg.* 63(6):1556-1561. Vectors encoding the transgene include both replicable and replication-, defective vectors, such as retroviral vectors, adenovirus vectors or other vectors with the appropriate tropism for the cells likely to be involved in apoptosis or cells proximal to the site of apoptosis. In the transgene constructs, expression can be directed from any suitable promoter, including tissue specific promoters which direct gene expression in specific cell types, such as the human insulin promoter. Local delivery of the transgene to the tissue, organ or to a site proximal to the transplant is accomplished by any means commonly available, including but not limited to direct local perfusion, injection, microsponges, microcapsules, liposomes or time-released delivery vehicles.

**[0185]** Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. The following working examples therefore, specifically point out preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure. Other generic configurations will be apparent to one skilled in the art. All journal articles and other published documents such as patents and patent applications are hereby incorporated by reference in their entireties.

## EXAMPLES

### Example 1

#### Experimental procedures and cloning

**[0186]** Cells and cell culture. The following cell lines were obtained from American Type Culture Collection (ATCC,

Rockville, Md.), erythroleukemia HEL, B-lymphoma Daudi and JY, monocytic THP-1, T leukemia Jurkat, epithelial carcinoma HeLa, promyelocytic HL-60, and non-transformed human lung fibroblast Lu18. The T leukemia cell line MOLT13 was characterized previously (Altieri, D. C., *FASEB J* (1995) 9:860-865). Cells were maintained in culture in complete medium RPMI 1640 or DMEM (HeLa, Lu18) (BioWhittaker, Walkersville, Md.), supplemented with 10% heat-inactivated fetal bovine serum (FBS, Whitaker), 2 mM L-glutamine, and 10 mM HEPES. Human umbilical vein endothelial cells (HUVEC) were isolated by collagenase treatment and maintained in culture in DMEM medium supplemented with 20% FBS, 2 mM L-glutamine and endothelial cell growth factor (Biomedical Technologies, Stoughton, Mass.).

**[0187]** Peripheral blood mononuclear cells (PBMC) were isolated from heparinized blood collected from normal informed volunteers by differential centrifugation on Ficoll-Hypaque (Pharmacia, Piscataway, N.J.) at 400 g for 22° C., and washed in phosphate buffered saline (PBS), pH 7.4. In some experiments, HL-60 cells were terminally differentiated to a mature monocytic phenotype by a 72 h culture in the presence of 0.1  $\mu$ M 1, 25-dihydroxy-vitamin D<sub>3</sub> and 17.8  $\mu$ g/ml indomethacin (Sigma Chemical Co., St. Louis, Mo.). De novo induction of differentiation-dependent markers on vitamin D<sub>3</sub>-treated HL-60 cells, including CD11b/CD18 integrin (Hickstein, D. D. et al., *J Immunol* (1987) 138:513-519) was determined by flow cytometry with anti-CD11b mAb LM2/1.

**[0188]** Genomic and cDNA cloning, chromosomal localization and Southern blots. A human P1 genomic library (Genome Systems, St. Louis, Mo.) was screened by hybridization with a 1.6 kb fragment containing the complete human EPR-1 cDNA (Altieri, D. C., *FASEB J* (1995) 9:860-865). Three overlapping clones were isolated, purified and confirmed by Southern hybridization with the EPR-1 cDNA. Hybridizing fragments generated by restriction digest with BamHI, HindIII and XbaI (Boehringer Mannheim, Indianapolis, Ind.) were cloned in pBluescript (pBSKS<sup>+</sup>, Stratagene, San Diego, Calif.) for further analysis. An overlapping contig spanning 14796 bp from two EPR-1-hybridizing P1 clones was arrayed, characterized by restriction analysis, and completely sequenced on both strands by Taq FS polymerase-based automated sequencing using an Applied Biosystem Prism 377 automated sequencer (Foster City, Calif.). In some experiments, 10 mg of total RNA extracted from HeLa cells by the guanidinium isothiocyanate method was primed with EPR-1 forward "sense" oligonucleotide C3/27 (bp 80-102) and reverse transcribed in the presence of 200 U of Superscript II (Life Science, Grand Island, N.Y.) for 50 min at 42° C.

**[0189]** The resulting cDNA was amplified by PCR in the presence of 0.5 mg of EPR-1-derived primers T5/27 (bp 161-184) and G11/16 (1124-1098, numbering from the EPR-1 coding sequence), 200 mM dNTPs (New England Biolabs, Beverly, Mass.) and 2 U Vent DNA polymerase (New England Biolabs) in a total volume of 50  $\mu$ l. After 35 cycles of amplification with annealing at 58° C. for 1 min, denaturation at 94° C. for 1 min and extension at 72° C. for 1 min, the product was analyzed by agarose gel electrophoresis, subcloned in pCRII (Invitrogen Corp., San Diego, Calif.), and completely sequenced on both strands. Contig assembly, and DNA and protein sequence analyses were

performed using Lasergene (DNASTAR, Madison, Wis.) and MacVector (Eastman Kodak, Rochester, N.Y.) software packages. Chromosomal location of the EPR-1-hybridizing gene was carried out by fluorescence in situ hybridization. Purified DNA from a EPR-1-hybridizing P1 clone was labeled with digoxigenin dUTP (Amersham Corp., Arlington Heights, Ill.) by nick translation.

**[0190]** The labeled probe was combined with sheared human DNA and hybridized to normal metaphase chromosomes derived from phytohemagglutinin-stimulated PBMC in a solution containing 50% formamide, 10% dextran sulfate and 2×SSC. For two-color staining, biotin-conjugated probe D17Z1, specific for the centromere of chromosome 17, was co-hybridized with the digoxigenin-labeled P1 clone. Specific staining was detected by incubating the hybridized slides with fluoresceinated anti-digoxigenin antibodies and Texas red avidin. Slides were counterstained with propidium iodide for one color labeling, or with DAPI for two color labeling. A total of 80 metaphase cells were analyzed with 69 cells exhibiting specific labeling. For Southern hybridization, human genomic DNA was extracted from HeLa cells according to published protocols, digested with EcoRI, BamHI, XbaI or HindIII, separated on a 0.8% agarose gel and transferred to GeneScreen nylon membranes (New England Nuclear, Boston, Mass.).

**[0191]** After UV cross-linking (Stratalinker, Stratagene, San Diego, Calif.), the membrane was prehybridized with 100 mg/ml of denatured salmon sperm DNA (Promega Corp. Madison, Wis.) in 5×SSC, 0.5% SDS, 5×Denhardt's solution and 0.1% sodium pyrophosphate at 65° C. in a roller hybridization oven (Hoefer Scientific, San Francisco, Calif.). Hybridization was carried out with gel-purified (GeneClean Bio101, Vista, Calif.), <sup>32</sup>P-dCTP (Amersham) random-primed labeled (Boehringer-Mannheim, Indianapolis, Ind.) 1.6 kb EPR-1 cDNA for 16 h at 65° C.

**[0192]** After two washes in 2×SSC, 1% SDS for 30 min at 65° C., and 0.2×SSC at 22° C., radioactive bands were visualized by autoradiography using a Kodak X-Omat AR X-ray film and intensifying screens (DuPont de Nemours, Wilmington, Del.). In other experiments, cultured lymphoblastoid cells were embedded in LMP agarose (Bio Rad, Richmond, Calif.) at the concentration of 2×10<sup>6</sup>/220 μl block and DNA was extracted by standard procedures. After block digestion with MluI or NotI, samples were separated by pulsed field gel electrophoresis on a 1% agarose gel for 20 h at 200 V with a pulse time of 75 sec using a Bio-Rad CHEF DRII apparatus (Hercules, Calif.). After transfer to nylon membranes, and UV cross-linking, hybridization with the EPR-1 cDNA and washes were carried out as described above.

**[0193]** In another series of experiments, a blot containing aliquots of genomic DNA isolated from several species (Clontech, San Francisco, Calif.) was hybridized with a 3' 548 bp fragment of the EPR-1 cDNA, as described above.

**[0194]** Northern blots. Single strand probes specific for sense or antisense EPR-1 sequences were generated by asymmetric PCR amplification of a 301 bp fragment of the EPR-1 cDNA. The template, comprising the first 5' 226 bp of the EPR-1 coding sequence plus 75 bp of the retained regulatory intron (Altieri, D. C., *FASEB J* (1995) 9:860-865), was generated by restriction digest of the EPR-1 cDNA with EcoRI (cloning site) and SacII, gel-purified, and

mixed in a total volume of 10 ml with 15 pmol dNTP (New England Biolabs), 7.5 pmol dCTP, and 25 mCi <sup>32</sup>P-dCTP (Amersham), in the presence of 20 mM Tris HCl, 50 mM KCl, pH 8.4, 1.5 mM MgCl<sub>2</sub>, and 2.5 U of Taq DNA polymerase (Life Science).

**[0195]** Generation of a EPR-1-specific antisense probe was carried out by addition of 0.2 mg/ml of a "SacII" reverse oligonucleotide 5'TGCTGGCCGCTCCTCCCTC3', while extension of the EPR-1 positive strand and generation of a Survivin-specific probe was performed using 0.2 mg/ml of forward F11 oligonucleotide 5'ATGACCTCCAGAG-GTTTC3'. Twenty-five cycles of amplification were carried with denaturation at 94° C. for 1 min, annealing at 52° C. for 1 min, and extension at 72° C. for 1 min. The EPR-1 sense or antisense probes were centrifuged through a Sephadex G-50 spin column (Worthington Biochemical Corp., Freehold, N.J.) at 14,000 g for 5 min to separate free from incorporated radioactivity, heated at 100° C. for 2 min, and immediately added to the hybridization reaction.

**[0196]** Identical strand-specific probes were used for hybridization of multiple tissue blots of adult or fetal human mRNA (Clontech), in 5×SSPE, 10×Denhardt's solution, 2% SDS, 100 mg/ml denatured salmon sperm DNA at 60° C. for 14 h, and washes at 60° C., as described above. Aliquots of total RNA extracted from undifferentiated or vitamin D<sub>3</sub> terminally differentiated HL-60 cells, were processed for Northern hybridization with Survivin-specific single strand probe, as described above.

#### Example 2

##### Production of Anti-Survivin Antibodies

**[0197]** A Survivin sequence-specific antibody, called JC700, was produced and characterized as follows. A seventeenmer peptide corresponding to the Survivin sequence A<sup>3</sup>PTLPPAWQPFLKDHRI<sup>19</sup> was synthesized and characterized by mass spectrometry. One hundred mg of the Survivin peptide were coupled in a 1:1 ratio to Keyhole Limpet Hemocyanin and injected s. c. into a rabbit in complete Freund's adjuvant. After a 4-week interval, animals were boosted with s. c. injection of 100 mg of peptide in incomplete Freund's adjuvant and sequentially boosted and bled at alternate weeks.

**[0198]** Purification of the anti-Survivin antibody was carried out by affinity chromatography on a peptide-Sepharose matrix (5 mg/ml of peptide) with elution of the specific IgG fraction in 1 mM glycine, pH 2.5. Specificity of the affinity-purified anti-Survivin antibody, designated JC700, was determined by ELISA against the immobilized Survivin peptide or a control EPR-1 peptide by absorbance at OD<sub>405</sub>.

#### Example 3

##### Production of a Monoclonal Antibody Against a Survivin Fusion Protein

**[0199]** The Survivin cDNA was expressed as a GST-fusion protein in *E. Coli* BL21 strain and purified to homogeneity with removal of the GST frame. The purified protein was used to inject mice and generate monoclonal antibodies using standard hybridoma technology. Three independent mAbs were isolated, cloned twice by limiting dilution and further characterized. One of the new anti-Survivin mAbs,

designated 8E2, recognized the immobilized, purified recombinant Survivin by ELISA and bound to Survivin in immunoblots, as shown in **FIG. 11**.

#### Example 4

##### Immunoblotting and In Situ Hybridization

**[0200]** For immunoblotting, aliquots of SDS-solubilized extracts of various transformed cell lines, non-transformed HUVEC, PBMC or Lu18, or undifferentiated or vitamin D<sub>3</sub>-differentiated HL-60 cells, were normalized for protein content by absorbance at OD<sub>280</sub>, separated by electrophoresis on a 5-20% SDS polyacrylamide gradient gel under non-reducing conditions, and electroblotted to Immobilon membranes (Millipore Corp., New Bedford, Mass.) at 1.1 A for 30 min at 22° C. The membrane was blocked in TBS, pH 7.4, plus 5% milk, and incubated with 20 mg/ml of control non-immune rabbit IgG or anti-Survivin antibody JC700 for 1 h at 22° C., followed by washes in TBS, pH 7.4, and addition of a 1:7500 dilution of alkaline phosphatase-conjugated goat anti-rabbit IgG (Promega) for 30 min at 22° C. Binding of the primary antibody was revealed by addition of 75 mg/ml nitro blue tetrazolium in 70% dimethylformamide (Sigma Chemical Co., St. Louis, Mo.) plus 50 mg/ml 5-bromo-4-chloro-3-indolyl phosphate (Sigma) in 100% dimethylformamide.

**[0201]** Tissue samples, immunohistochemistry and in situ hybridization. Tissue samples from colon adenocarcinoma (6 cases), lung squamous cell carcinoma (6 cases), lung adenocarcinoma (9 cases), pancreas adenocarcinoma (2 cases), invasive breast adenocarcinoma (7 cases), were obtained from the archives of Yale-New Haven Hospital and used in the present study. Samples of 44 high grade lymphoma tissues and 7 low grade lymphoma tissue was also obtained. Tissue samples were fixed in formalin, embedded in paraffin, cut in 5  $\mu$ m sections, deparaffinized in xylene, and rehydrated in graded alcohol followed by quenching of endogenous peroxidase activity by treatment with 2% H<sub>2</sub>O<sub>2</sub> in methanol.

**[0202]** For immunostaining, the slides were boiled for 5 min in a standard pressure cooker, blocked in 10% normal goat serum, and incubated with affinity-purified anti-Survivin antibody JC700 (20  $\mu$ g/ml) for 14 h at 4° C. After washes in PBS, pH 7.4, slides were further incubated with biotin-conjugated goat anti-rabbit IgG (Vector Laboratories, Burlingame, Calif.) for 30 min at 22° C., and washed in PBS, pH 7.4. After addition of streptavidin-biotin conjugated peroxidase (Boehringer Mannheim) for 30 min at 22° C., slides were washed, and binding of the primary mAbs was revealed by addition of 3'-3'-diamino-benzidine (DAB) and counterstaining with hematoxylin.

**[0203]** Negative controls were carried out by replacing the primary antibody with normal goat serum, under the same experimental conditions. In some experiments, aliquots of JC700 antibody were pre-adsorbed with 25 mg/ml of the Survivin 3-19 peptide before tissue staining. For in situ hybridization, 1  $\mu$ g of the Survivin cDNA containing the entire coding sequence plus 271 bp of 3' untranslated region in pcDNA3 (Invitrogen), was completely digested with EcoRI and transcribed in the antisense orientation using T7 RNA polymerase in the presence of digoxigenin 11-uridine-5' triphosphate (Boehringer Mannheim). Tissue slides were

coated with 1% gelatin, 0.1% chrome-alum, baked at 120° C. for 2 h, and stored dust-free at 22° C. Sections were deparaffinized and rehydrated through graded alcohol, digested with proteinase K (1  $\mu$ g/ml in 100 mM Tris HCl pH 8.7, 50 mM EDTA) for 30 min at 37° C., and acetylated in 0.25% acetic anhydride acid and 100 mM triethanolamine pH 8.0 for 10 min at 22° C.

**[0204]** Detection of Survivin mRNA in human tissues was carried out by in situ hybridization of the Survivin antisense riboprobe in a buffer containing 4 $\times$ SSC, 1 $\times$ Denhardt's solution, 50% deionized formamide, 250  $\mu$ g/ml yeast tRNA, 500  $\mu$ g/ml salmon sperm DNA and 5% dextran for 16 h at 50° C. After washes in 2 $\times$ SSC for 90 min at 48° C., immobilized digoxigenin was detected using an anti-digoxigenin mAb (Boehringer Mannheim) at a 1:3000 dilution, and revealed by alkaline phosphatase staining with NBT/BCIP cytochemical stain.

#### Example 5

##### Expression of Survivin in Human Cancers

**[0205]** Survivin is prominently expressed in human cancer. For its abundant distribution in transformed cell types, a potential expression of Survivin in neoplasia was investigated, in vivo. Immunohistochemical analysis of formalin-fixed, paraffin embedded tissue sections with the affinity-purified anti-Survivin JC700 antibody demonstrated abundant expression of Survivin in all cases examined of human lung cancer, including adenocarcinoma (**FIG. 6A**), and squamous cell carcinoma (**FIG. 6C**). Consistent with the topography of other IAP proteins (Duckett, C. S. et al., *EMBO J* (1996) 15:2685-2694), expression of the protein was exclusively localized to the cytoplasm of tumor cells, while the adjacent normal gland epithelium of the lung did not express Survivin (**FIG. 6C**, arrow). No staining was observed when the anti-Survivin antibody was substituted with control goat serum (not shown), or after pre-adsorption with the immunizing Survivin 3-19 peptide (**FIG. 6B**), thus confirming the specificity of the observed recognition.

**[0206]** Prominent accumulation of Survivin mRNA in squamous lung cell carcinoma was independently demonstrated by in situ hybridization with a Survivin-specific single strand riboprobe (**FIG. 6D**). Survivin was also abundantly detected in all cases examined of adenocarcinoma of pancreas (**FIG. 6E**), and breast (not shown) by immunohistochemistry, and colon (**FIG. 6G**) by in situ hybridization. However, consistent with its absence in non-transformed cell types HUVEC and Lu18 (**FIG. 4C**), in mature tissues (**FIG. 3**), and in terminally-differentiated HL-60 cells (**FIG. 5**), no reactivity of the anti-Survivin JC700 antibody was observed with normal exocrine pancreatic epithelial cells by immunohistochemistry (**FIG. 6F**), and no Survivin mRNA was found in the adjacent non-neoplastic colon gland epithelium by in situ hybridization (**FIG. 6H**).

**[0207]** Expression of Survivin in Lymphoma Tissue. Tissue samples were obtained from 44 patients with aggressive, high grade lymphoma and 7 samples were obtained from 7 patients with non-aggressive, low grade lymphoma. The sample were treated as described above and examined for Survivin expression. None of the low grade lymphoma samples displayed Survivin expression whereas 27 samples (61%) from patients with high grade lymphoma expressed Survivin.

## Example 6

## Expression of Survivin in other Cancers

[0208] In addition to the malignant forms of cancer discussed above, the expression of Survivin in other types of cancers was investigated in the inventors' laboratory or collaboratively with other academic investigators. Survivin was found prominently expressed in the most aggressive and metastatic forms of malignant thymoma (~100 cases tested), in head and-neck squamous cell carcinoma (~140 cases) and in all forms of prostate cancer (15 cases), including the transition lesion of benign prostate hyperplasia. The most aggressive forms of neuroblastoma are also positive for Survivin as discussed below.

## Example 7

## Tissue Specific Expression of Survivin

[0209] Survivin, was recently found in all the most common human cancers but not in normal, terminally differentiated adult tissues. The expression of Survivin in embryonic and fetal development was investigated. Immunohistochemistry and in situ hybridization studies demonstrated strong expression of Survivin in several apoptosis-regulated fetal tissues, including the stem cell layer of stratified epithelia, endocrine pancreas and thymic medulla, with a pattern non-overlapping with that of another apoptosis inhibitor, i.e. bcl-2. A sequence-specific antibody to Survivin immunoblotted a single ~16.5 kD Survivin band in human fetal lung, liver, heart, kidney and gastrointestinal tract. In mouse embryo, prominent and nearly ubiquitous distribution of Survivin was found at embryonic date (E) 11.5, whereas at E15-21, Survivin expression was restricted to the distal bronchiolar epithelium of the lung and neural crest-derived cells, including dorsal root ganglion neurons, hypophysis and the chorioid plexus. These data suggest that expression of Survivin in embryonic and fetal development may contribute to tissue homeostasis and differentiation independently of bcl-2.

## Example 8

## Preparation of Survivin Transfectants

[0210] Inducible Survivin antisense transfectants and apoptosis/proliferation experiments. A 708 bp SmaI-EcoRI fragment comprising nucleotides 379-1087 of the EPR-1 cDNA, was directionally cloned in the sense orientation in the mammalian cell expression vector pML1 (generously provided by Dr. R. Pytela, Cardiovascular Research Institute, University of California, San Francisco). The vector is derived from the episomal mammalian cell expression vector pCEP4 by replacing the cytomegalovirus promoter cassette with the mMT1 promoter, directing  $Zn^{2+}$ -dependent expression of recombinant proteins in mammalian cells (Lukashev, M. E. et al., *J Biol Chem* (1994) 269:18311-18314).

[0211] Ten million HeLa cells were incubated with 10 mg of pML1 DNA containing the Survivin antisense construct plus 50 mg of salmon sperm DNA for 15 min on ice, followed by a single electric pulse delivered by a Gene Pulser apparatus (Bio-Rad) at 350 V at 960  $\mu$ F. Forty-eight h after transfection, cells were diluted fifteen fold, plated onto 100 mm diameter tissue culture dishes and selected for

4 weeks in complete growth medium containing 0.4 mg/ml hygromycin. Apoptosis in control cultures or Survivin antisense HeLa cell transfectants was evaluated by in situ detection of internucleosomal DNA degradation after  $Zn^{2+}$ -dependent induction of EPR-1 transcription under serum-starving conditions.

[0212] Briefly, control or antisense Survivin transfectants were treated with 200 mM  $ZnSO_4$  in 0% FBS for 24 h at 37° C. Cells were harvested, centrifuged at 800 g for 10 min at 4° C., and the pellet was fixed in 10% formalin overnight, dehydrated, embedded in paraffin blocks, and sections of 3-5 mm were put on high adhesive slides. Samples were treated with 20 mg/ml proteinase K for 15 min at 22° C., washed in distilled water, quenched of endogenous peroxidase in 2%  $H_2O_2$  in PBS, and subsequently mixed with digoxigenin-labeled dUTP in the presence of terminal deoxynucleotidyl transferase (TdT) followed by peroxidase conjugated anti-digoxigenin antibody.

[0213] Nuclear staining in apoptotic cells was detected by DAB, according to the manufacturer's instructions (AptoTag, Oncor, Gaithersburg, Md.). Control experiments were performed by omitting the enzyme incubation step. Morphologic features of apoptotic cells (apoptotic bodies) under the various conditions tested were detected by hematoxylin/eosin staining of the same slides.

[0214] For proliferation experiments, vector control HeLa cells or Survivin antisense transfectants were plated at  $20 \times 10^4$ /well onto 24-well tissue culture plates (Costar), induced with 200 mM  $ZnSO_4$  for 16 h at 37° C., harvested at 24 h intervals, and cell proliferation under the various conditions tested was determined microscopically by direct cell count. Down-regulation of Survivin expression under these experimental conditions was assessed by immunoblotting with JC700 antibody.

## Example 9

## Identification of EPR-1 Complementary Gene

[0215] Three overlapping clones were isolated by hybridization screening of a human P1 plasmid genomic library with the EPR-1 cDNA and confirmed by Southern blot. This gene was located to the long arm of chromosome 17, to band 17q25, by fluorescence in situ hybridization (FIG. 1A, B).

[0216] A contig of P1 fragments spanning 14796 bp was cloned in pBSKS<sup>+</sup> and completely sequenced on both strands (FIG. 1C). Three putative splice sites, matching perfectly the consensus sequences for eukaryotic intron-exon boundaries (Padgett, R. A. et al., *Ann Rev Biochem* (1986) 55:1119-1150), were identified at position 2922, 3284, and 5276 (donor), and 3173, 5157, and 11954 (acceptor), thus defining a gene organization in four exons and three introns of 252, 1874, and 6678 bp, respectively (FIG. 1D).

[0217] Sequence analysis of the putative coding regions demonstrated a nearly complete identity with the EPR-1 cDNA (Altieri, D. C., *FASEB J* (1995) 9:860-865), except for 5 nucleotide changes and 6 nucleotide insertions. However, the three splice sites were found on the complementary, antisense strand of the EPR-1 coding sequence. Consistent with this unexpected orientation, the EPR-1 complementary gene revealed a 5' GC rich region, comprising nucleotides



2560-2920 and including exon 1 (see below), which fulfilled the base composition criteria of a CpG island (Gardiner-Garde, M. et al., *J Mol Biol* (1987) 196:261-282 and Frommer, 1987). Sequencing the 2.5 kb upstream the CpG island revealed a TATA-less promoter with numerous Sp1 sites (not shown).

[0218] Complex hybridization pattern and evolutionary conservation of EPR-1 sequences. Probing human genomic DNA with the EPR-1 cDNA revealed several hybridizing fragments (**FIG. 2A**). Of these, a ~7.5 kb XbaI, a 7.6 kb BamHI, and 4 HindIII fragments of ~15, 7.5, 6.4, and 3.7 kb, respectively (**FIG. 2A**, arrows), could not be recapitulated by the restriction map of the antisense EPR-1 gene (**FIG. 1C**). In contrast, other bands of comparable intensity, including a 5.15 kb XbaI and a 7.1 kb BamHI fragment, genuinely originated from the antisense EPR-1 gene and comprised the first two, or three exons, respectively (**FIG. 2A**).

[0219] At variance with this complex hybridization pattern, Southern blot of high molecular weight human genomic DNA digested with MluI or NotI and separated by pulsed field gel electrophoresis, revealed single EPR-1-hybridizing bands of ~75 kb and 130 kb, respectively (**FIG. 2B**). Finally, Southern blots of multiple species genomic DNA revealed significant evolutionary conservation of EPR-1-related sequences (**FIG. 2C**), with numerous strongly hybridizing bands in mammalian species and fainter signals in rabbit or chicken genomic DNA, under high stringency hybridization conditions (**FIG. 2C**).

[0220] Discordant tissue distribution of sense/antisense EPR-1 transcripts. The potential expression of distinct sense or antisense EPR-1 transcripts was investigated in Northern blots with single strand-specific probes. Consistent with the size of the spliced EPR-1 message (Altieri, D. C., *FASEB J* (1995) 9:860-865), an EPR-1 strand-specific probe detected a prominent ~1.2 kb band in mRNA extracted from all adult and terminally-differentiated human tissues examined (**FIG. 3A**). In contrast, no specific bands hybridized with a EPR-1 antisense-specific single strand probe in adult tissues, under the same experimental conditions (**FIG. 3B**). A similar ~1.2 kb band was detected by the single strand EPR-1-specific probe in fetal kidney, and, to a lesser extent, in fetal liver, lung and brain (**FIG. 3A**). At variance with the absence of hybridization in adult tissues, the EPR-1 antisense-specific probe recognized a prominent ~1.9 kb band, and a larger 3.2 kb species corresponding to the size of an incompletely processed transcript, in fetal liver, while fainter hybridization bands were also seen in fetal kidney, lung and brain (**FIG. 3B**). A control hybridization with an actin probe confirmed comparable loading of mRNA in adult or fetal samples (**FIG. 3C**).

[0221] Characterization of the antisense EPR-1 gene product. Inspection of the 5' CpG island in the antisense EPR-1 gene revealed a putative ATG initiation codon at position 2811, surrounded by a sequence (CGGCATGG) that conformed well to the consensus for eukaryotic initiation of translation (Kozak, M., *Nucleic Acids Res* (1984) 12:857-872). Analysis of the antisense EPR-1 sequence in the 5'→3' direction dictated by the position of intron-exon boundaries revealed an open reading frame of 426 bp, spanning all four exons, and terminating with a TGA codon at position 12042 in exon 4. A canonical polyadenylation signal (AATAAA) was found at position 13166. PCR products amplified from

reverse-transcribed HeLa cell RNA primed with EPR-1 "sense" oligonucleotides matched perfectly the genomic sequence and confirmed the open reading frame and the predicted intron-exon boundaries (not shown).

[0222] Two  $\lambda$ gt11 cDNA clones isolated by hybridization of a HEL library with the EPR-1 CDNA, also matched the consensus genomic sequence and revealed a homopolymeric A tail on the antisense EPR-1 strand at position 13186, 14 bp downstream the polyadenylation signal, generating a 3' untranslated region of 1144 bp. In these clones, the 5' untranslated region upstream from the initiating ATG was of 49 bp, beginning at position 2762 in the genomic sequence, and contained an in-frame termination codon. Translation of the antisense EPR-1 open reading frame predicted a new protein of 142 amino acids, with an estimated molecular weight of 16,389 and an acidic pI of 5.74, lacking an amino-terminus signal peptide or a carboxy terminus hydrophobic stretch for membrane insertion (**FIG. 4A**).

[0223] A coiled coil was predicted for the last 40 carboxy terminus residues (Lupas, A. et al., *Science* (1991) 252:1162-1164). BLAST database searches revealed a significant degree of similarity between residues 18-88 of the antisense EPR-1 gene product and the BIR module in the IAP family of inhibitors of apoptosis (Birnbaum, M. J. et al., *J Virology* (1994) 68:2521-2528; Clem, R. J. et al., *Mol Cell Biol* (1994) 14:5212-5222). For this analogy, the antisense EPR-1 gene product was designated Survivin. At variance with other IAP proteins, Survivin contained only one BIR, encoded by the first three exons of the gene, and lacked a carboxy terminus RING finger, without additional/alternative exon(s) potentially encoding this domain (**FIG. 1C**).

[0224] An alignment by the Clustal method between the Survivin BIR and that of other known IAP proteins is shown in **FIG. 4B**. Despite the overall match of the consensus and several conservative substitutions, phylogenetic analysis suggested that Survivin is a distantly related member of the IAP family, most closely related to NAIP, which also lacked a RING finger (**FIG. 4B**, shaded boxes) (Roy, N. et al., *Cell* (1995) 80:167-178).

[0225] A rabbit polyclonal antiserum designated JC700, was raised against residues A<sup>3</sup>PTLPPAWQPFLKDHR<sup>19</sup> of Survivin, purified by affinity chromatography on a peptide-Sepharose column, and used in Western blots. Consistent with the predicted molecular weight of Survivin, JC700 antibody immunoblotted a single band of ~16.5 kDa from detergent-solubilized extracts of all transformed cell lines examined, including B lymphoma Daudi and JY, T leukemia Jurkat and MOLT13, monocytic THP-1, and erythroleukemia HEL (**FIG. 4C**).

[0226] Survivin was also found in isolated peripheral blood mononuclear cells (PBMC). In contrast, no expression of Survivin was detected in non-transformed Lu-18 human lung fibroblasts or human umbilical vein endothelial cells (HUVEC) (**FIG. 4C**). No specific bands were immunoblotted by control non-immune rabbit IgG, under the same experimental conditions (**FIG. 4C**).

[0227] Identification of agents that modulate transcription of the EPR-1 gene. Agents that increase the transcription of the EPR-1 gene may be identified by conventional techniques. Preferably, a candidate agent is brought into contact with a cell that expresses the EPR-1 gene product and the

level of expression of this product or the level of transcription are determined and agents that increase or decrease EPR-1 gene transcripts may readily be identified. Alternatively, the EPR-1 transcriptional regulatory elements may be placed upstream of a reporter gene such as CAT or  $\beta$ -galactosidase.

#### Example 10

##### Regulation of Survivin Expression by Cell Growth/Differentiation

[0228] Consistent with the expression of Survivin in transformed cell lines (FIG. 4C), undifferentiated and actively proliferating promyelocytic HL-60 cells constitutively expressed high levels of Survivin, as demonstrated by immunoblotting of a single ~16.5 kDa band with JC700 antibody, and Northern hybridization of a ~1.9 kb transcript with a single strand-specific probe (FIG. 5). In contrast, no specific bands were recognized by control non-immune rabbit IgG under the same experimental conditions (FIG. 5).

[0229] Vitamin D<sub>3</sub>-induced terminal differentiation of HL-60 cells to a mature monocytic phenotype resulted in growth arrest of these cells and de novo induction of differentiation-specific markers, including a ~200-fold increased expression of leukocyte CD11b/CD18 integrin detected by flow cytometry (not shown), and in agreement with previous observations (Hickstein, D. D. et al., *J Immunol* (1987) 138:513-519). Under these experimental conditions, the anti-Survivin JC700 antibody failed to immunoblot specific bands from vitamin D<sub>3</sub>-treated HL-60 extracts, and no Survivin transcript(s) were detected by Northern hybridization with a single strand-specific probe (FIG. 5).

[0230] In contrast, an anti-EPR-1 polyclonal antibody immunoblotted a single ~62 kDa band corresponding to EPR-1 in vitamin D<sub>3</sub>-differentiated HL-60 extracts under the same experimental conditions (not shown). Moreover, down-regulation of Survivin in vitamin D<sub>3</sub>-differentiated HL-60 cells was accompanied by a 5- to 10-fold increased surface expression of EPR-1 in these cells, as detected by flow cytometry with anti-EPR-1 monoclonal antibodies B6 or 12H1 (FIG. 8).

[0231] As shown in FIG. 16, Survivin is down regulated by the combination of cytokines  $\gamma$  interferon and tumor necrosis factor  $\alpha$ , but not by either cytokine alone. Similarly, the transfection of 3T3 cells with the c-myc oncogene results in the up-regulation of Survivin mRNA as detected by Northern blots.

#### Example 11

##### Promoting Apoptosis with Survivin

[0232] Targeting Survivin promotes apoptosis and inhibits cell proliferation. Transfection of the Survivin cDNA in mouse or hamster cell lines (NIH 3T3, CHO) was not suitable for the presence of immunochemically indistinguishable endogenous homologues in these cells (not shown). Similarly, initial attempts to target the Survivin gene in stable antisense transfectants were unsuccessful for slow cell growth and rapid loss of viability (not shown). Therefore, Survivin<sup>+</sup> HeLa cells were transfected with the 3' end of the EPR-1 cDNA (Survivin antisense) under the control of a metallothionein-inducible promoter (Lukashev,

M. E. et al., *J Biol Chem* (1994) 269:18311-18314), selected in hygromycin, and analyzed for apoptosis and cell proliferation after Zn<sup>2+</sup>-dependent activation of transcription.

[0233] Consistent with the expression of Survivin in transformed cell lines (FIG. 4C), the JC700 antibody immunoblotted a single molecular species of ~16.5 kDa in extracts of control HeLa cells transfected with the vector alone (FIG. 7A). In contrast, no specific bands were recognized by JC700 antibody in metallothionein-induced HeLa cells transfected with the EPR-1 cDNA (Survivin antisense) (FIG. 7A). Under these experimental conditions, in situ analysis of internucleosomal DNA fragmentation by AptoTag staining revealed only a few apoptotic cells in serum-starved, Zn<sup>2+</sup>-induced, vector control HeLa cells (FIG. 7B).

[0234] In contrast, as discussed above, inhibition of Survivin expression in Zn<sup>2+</sup>-induced antisense HeLa cell transfectants was associated with prominent nuclear staining in the vast majority of cells examined (FIG. 7B). No nuclear staining was detected in the absence of TdT tagging of the digoxigenin-labeled dUTP (not shown). Typical morphologic features of apoptosis, including numerous apoptotic bodies, were also demonstrated in induced antisense HeLa cell transfectants by hematoxylin/eosin staining, while only occasional apoptotic bodies were observed in vector control HeLa cultures, under the same experimental conditions (FIG. 7B).

[0235] A potential effect of Survivin on cell growth was also investigated. In these experiments, metallothionein-controlled, EPR-1-dependent, inhibition of Survivin expression caused a profound reduction of serum-dependent HeLa cell proliferation (FIG. 7C). Three days after Zn<sup>2+</sup> induction, the cell count in vector control HeLa cultures increased by 288%, as opposed to only a 20% increase in Survivin antisense transfectants, under the same experimental conditions (FIG. 7C).

#### Example 12

##### Structure—Function Relationship of Survivin

[0236] The minimal structural requirements involved in Survivin-mediated inhibition of apoptosis have been identified through a mutagenesis strategy of Ala substitutions of the most evolutionarily conserved residues in the single Survivin BIR (baculovirus IAP repeat) module. These residues included in the amino-terminal half of the Survivin BIR, Arg<sup>18</sup>, Phe<sup>22</sup>, Trp<sup>25</sup>, Pro<sup>26</sup>, Pro<sup>35</sup>, Ala<sup>39</sup>, Ala<sup>41</sup>, Gly<sup>42</sup>, and Cys<sup>46</sup>. In the carboxyl-terminal half of the Survivin BIR, Ala mutants were first targeted at the Cys<sup>37</sup>X<sub>2</sub>Cys<sup>60</sup>X<sub>16</sub>His<sup>77</sup>X<sub>6</sub>Cys<sup>84</sup> putative zinc binding motif. Additional conserved residues targeted by mutagenesis include Asp<sup>53</sup>, Leu<sup>64</sup>, Trp<sup>67</sup>, Pro<sup>69</sup>, Asp<sup>71</sup>, Asp<sup>72</sup> and Pro<sup>73</sup>.

[0237] The Survivin mutants are characterized in stable and transiently transfected cells, IL-3-dependent BaF3 cells and NIH3T3, respectively. In addition to these point mutants, a Survivin chimeric molecule containing a carboxyl-terminal RING finger has also been generated and screened for apoptosis inhibition (the RING finger is a domain found in most other IAP proteins, but not in Survivin). Secondly, a truncated form of Survivin has also been generated, in which the last 40 carboxylterminus residues, containing a predicted coiled-coil structure, have been

deleted. As shown in **FIG. 12**, Ala mutagenesis of key conserved residues in Survivin Trp<sup>67</sup>-Pro<sup>73</sup>-Cys<sup>84</sup> produced a recombinant molecule which lacked the ability to protect BaF3 cells from apoptosis induced by IL-3 withdrawal.

#### Example 13

##### Cytoprotective Effects of Survivin

[0238] Classical examples of cell damage to stable cell populations mediated by apoptosis include allograft rejection by infiltrating lymphocytes, Alzheimer's disease and reperfusion injury following myocardial infarction. In addition to being expressed in cancer, thereby functioning as a growth-advantage factor for cancer cells, the targeted expression of Survivin is useful to protect stable cell populations from apoptosis and other cellular insults. This application of Survivin was tested by adding increasing concentrations of purified recombinant Survivin to monolayers of human endothelial cells injured with hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), a classical apoptosis-inducing stimulus. The results are summarized in **FIG. 13**. Increasing concentrations of added Survivin resulted in a significant increased viability of the treated cells as opposed to control cultures treated with control protein myoglobin. Similarly, Survivin protected NIH3T3 cells from apoptosis induced by hydrogen peroxide after transient co-transfection with a lacZ reporter gene as shown in **FIG. 17**.

#### Example 14

##### Survivin as a Predictive-Prognostic Factor

[0239] The presence of Survivin can be utilized as predictive-prognostic negative factor in neuroblastoma and non-Hodgkin's lymphoma, and in other cancers.

[0240] Neuroblastoma. A large series of neuroblastoma cases (72) was screened for Survivin expression in a multicentric study. As shown in **FIG. 14**, Survivin expression increased dramatically when patients contained at least one negative prognostic factor for aggressive and rapidly progressing disease. Secondly, expression of Survivin strongly correlated with a more aggressive disease and unfavorable histology. Importantly, expression of Survivin was a more sensitive prognostic index than simple histology. Survivin-positive cases with early diagnosis of favorable histology were found to contain at least one negative prognostic factor for disease progression and dissemination.

[0241] Hodgkin's Lymphoma. A similar multicentric study has been recently completed on analysis of Survivin expression in high grade non-Hodgkin's lymphoma (n=8). The results are similar to those observed for neuroblastoma. As shown in **FIG. 15**, expression of Survivin strongly correlated with a more widespread disease predominantly in stage IV. Clinically, Survivin-expressing patients had fewer episodes of complete remission and more episodes of incomplete remission, no remission or relapses as compared with Survivin-negative patients.

[0242] Potential implications. The demonstrated role of Survivin as a negative predictive prognostic factor in these two embryologically different types of cancer iterates the potential use of this molecules a diagnostic tool to monitor disease progression and response to the therapy. It can also be used for staging purposes and to identify populations of

patients potentially susceptible to multi-drug resistance (groups with no remissions or incomplete remissions). Also, Survivin derived primers easily designed from the complete sequence of the Survivin gene can be used as a screening tool to identify potential cases of cancer in which the Survivin gene has been deleted or mutated. These cases will be very important to identify because targeted inactivation of the Survivin gene would confer a favorable prognostic factor to cancer patients, removing a potential drug-resistance gene. Inactivating mutations in the Survivin gene can target the same key residues identified in our initial screening of Ala-based mutagenesis or result in an abortive or truncated protein for premature termination of translation.

#### Example 15

##### Survivin Cancer Vaccine

[0243] Vaccines directed against Survivin, as found in various types of cancer, may be developed as with other disease-related intracellular protein targets. These techniques are commonly available and representative approaches are described by the references cited below. Vaccines may also include the systemic administration of peptide fragments of Survivin and the use of vectors to deliver mini-genes encoding Survivin peptides to tumor cell targets are contemplated. As mentioned above, Survivin is not expressed in normal cells, even in proliferating stem cells in the bone marrow. This ensures that the immune response mounted against Survivin will be highly selective and specific and will not involve normal cells.

[0244] Development and Administration of Polypeptide-based vaccines

[0245] Methods of the use of peptide components in a monovalent or a polyvalent cancer immunotherapy-vaccine product are described by Nardi, N. et al., *Mol. Med.* (1995) 1(5):563-567. Additional references that discuss the different cancer vaccine and cancer immunotherapies currently being used include: N. P. Restifo and M. Sznol "Cancer Vaccines," in DeVita's Cancer: Principles & Practice of Oncology 3023-3043 (Lippincott-Raven, Philadelphia; 1997); J. Galea-Lauri et al., *Cancer Gene Ther.* (1996) 3(3): 202-214; D. C. Linehan et al., *Ann. Surg. Oncol.* (1996) 3(2): 219-228; and J. Vieweg et al., *Cancer Invest.* (1995) 13(2): 193-201.

[0246] Consistent with the foregoing approach, Survivin polypeptides or full length Survivin are synthesized either chemically by known techniques or recombinantly by expressing appropriate cDNAs in prokaryotic or eukaryotic cells. Survivin proteins so produced are then purified as necessary to remove contaminating proteins, such as serum or bacterial proteins. Survivin can be further purified using columns containing antibodies that bind Survivin, such as the monoclonal antibody JC700 or the antibody 8E2 (both described above) which recognize and bind to Survivin. In purifying an antibody-based vaccine, the recombinantly produced Survivin would bind to the antibodies while other proteins and cellular debris would be washed out. Survivin polypeptides are then be isolated and concentrated to a desired strength.

[0247] Alternatively Survivin polypeptides are created by cleaving the native Survivin with one or more proteases (e.g., trypsin). Proteolytic fragments are then be separated

and recovered using SDS-PAGE, high-resolution/high-pressure separation techniques, or reverse-phase HPLC. See R. J. BEYNON AND J. S. BOND, *PROTEOLYTIC ENZYMES: A PRACTICAL APPROACH* (Oxford University Press, New York 1989). These isolated peptides are then be concentrated to a desired final concentration.

**[0248]** Once purified, Survivin polypeptides or full length Survivin molecules may then placed in an emulsion containing an adjuvant. Adjuvants contemplated for use with Survivin include aluminum adjuvants, Freund's adjuvant, oil-in-water emulsions containing tubercle bacilli, and interleukin-2 (IL-2). Additional preparations include combining the Survivin polypeptides with other appropriate tumor-associated antigens and, optionally, other immunomodulatory agents such as cytokines. Other suitable carriers or excipients can be used including bovine serum albumin, coupling the Survivin polypeptide with haptens, keyhole limpet hemocyanin, ovalbumin, and purified protein derivative of tuberculin. Peptides may be coupled to carriers using techniques such as those described in ED HARLOW AND DAVID LANE, *ANTIBODIES: A LABORATORY MANUAL* (Cold Spring Harbor Laboratory, 1988).

**[0249]** Vaccines in human subjects may be administered in the form of an emulsion injected subcutaneously, intradermally or intramuscularly (IM); vaccines appropriately formulated can be taken orally. With vaccines containing adjuvants, the vaccine is generally preferably be given IM, e.g., in the deltoid.

**[0250]** The amount of Survivin vaccine or Survivin peptide vaccine to be administered to a patient will correspond to values typically used in for other cancer vaccines. Dosage concentrations will range from about 0.25 g to about 1000 g per day. More preferred ranges will be from about 10  $\mu$ g to about 500  $\mu$ g per day.

#### Example 16

##### Diagnostic use of Anti-Survivin Antibodies

**[0251]** Frequently, tumor associated antigens (TAA) are shed from tumor cells into the surrounding plasma or into the blood. As a result, TAA often are found in the blood, and blood samples obtained from patients may be used in detecting the presence of cancer, as well as used as a factor is staging cancers (e.g., stage I, II, III or IV). Survivin is one such TAA, and healthy, normal individuals do not express Survivin. Results from studies of several cancers have indicated that the presence of Survivin (or Survivin fragments) correlates with and is predictive that the disease may be aggressive or may have metastasized. A similar strategy of detecting and quantifying the levels of Survivin or Survivin fragments can be used to determine residual tumor burden in patients undergoing chemotherapy or radiation therapy for cancer treatment. Elevated or increasing levels of Survivin may reflect late stage neoplastic disease.

**[0252]** For diagnostic uses, blood is drawn from patients, by well known techniques, who have known cancer loads or from patients suspected of having cancer. The blood sample is prepared by known techniques and is tested for binding with antibodies to Survivin that are prepared and, optionally, labeled, as discussed above. Such general antibody detection protocols and associated reagents are well established in the art. Other biological fluid samples such as semen, urine, or

saliva can also be monitored for the presence of Survivin. This diagnostic technique also can be used to monitor disease progression and response to individualized therapies. This method offers a relatively non-invasive means of tracking cancer progression or regression.

#### Example 17

##### Detection of Survivin by Immunobioassay

**[0253]** An illustrative example of an immunobioassay to test for the presence of Survivin in the blood of patient relies on the ability of the monoclonal antibodies to Survivin to bind Survivin and remove the detectable Survivin from solution by immunoprecipitation. Such an immunobioassay is used to detect Survivin in suspected cancer patients and in fractions eluted from fractionation columns. An aliquot of each patient sample is incubated for 2 hours at 4° C. with a monoclonal antibody that specifically recognizes and binds Survivin, such as the Mab 8E2, described above. The monoclonal antibody is insolubilized on anti-mouse IgG agarose beads, which can be acquired from Sigma Chemical Co., St. Louis, Mo.

**[0254]** The agarose bead anti-mouse (IgG(H+L))-Survivin complex is prepared by first washing the agarose beads with binding buffer containing 0.01 M phosphate buffer, (pH 7.2), and 0.25 M NaCl and then incubating the beads with the Survivin monoclonal antibody for 18 hours at 4° C. in the same buffer. The agarose beads may then be sedimented by centrifugation for 30 seconds at 16,000 $\times$ g in a microcentrifuge and non-specific sites may be blocked by incubation with 2% non-fat dry milk in 0.5 M NaCl-TMK for 30 minutes at 4° C. After blocking, the beads may be washed 3 times with 0.5 M NaCl-TMK and resuspended in an equal volume of the same buffer. 20:1 of the agarose bead-monoclonal antibody complex may then incubated with each 250:1 of the patient test sample for 2 hours at 4° C. Any Survivin present in the patient test sample will be found by the Survivin monoclonal antibody on the beads. The bead complex, now with Survivin bound, may be removed by centrifugation for 30 seconds at 16,000 $\times$ g. The supernatant is then assayed for Survivin activity in the bioassay as described below. Control samples are treated with blocked beads that lacked the Survivin monoclonal antibody and tested for Survivin activity in the bioassay.

#### Example 16

##### Detecting Survivin using a Direct Elisa Test

**[0255]** Samples of normal plasma (control) and cancer patient-plasma are diluted 1:1 with phosphate buffered saline (PBS). One volume of each mixture is added to centricon-10 filter having a 10 kD molecular weight limit and centrifuged at 5000 $\times$ g (7000 rpm) for 1 hour. One volume of PBS is added to the retentate and centrifuged for 30 min. The final dilution is about 1:3. The ELISA plate wells are then coated with retentate at 1:6, 1:12, 1:24, 1:48 and 1:96 final dilution in bicarbonate coating buffer, having a pH 9.6 overnight at 4° C. C. The plates are then washed 2 times with wash buffer containing 5% Tween 20 in phosphate buffered saline. Residual binding sites are blocked with 4% bovine serum albumin (BSA), 300  $\mu$ /well for 2 hours. Plates are then washed 2 times with wash buffer. Next, 100  $\mu$ l of a monoclonal antibody that specifically

recognizes and binds to Survivin, such as Mab 8E2, is used at 1:200 dilution in 1% BSA is added to the wells and incubated for 1 hour with agitation. Plates are washed 5 times with wash buffer. Next, 100  $\mu$ l horseradish peroxidase conjugated secondary antibody is added, typically at a 1:2,000 dilution to each well, and incubated for 1 hour. Plates are again washed 5 times with wash buffer. Next, 100  $\mu$ l/well of substrate containing 5  $\mu$ g of Survivin and 5  $\mu$ l  $H_2O_2$ /10 ml citrate-phosphate buffer is added to each well and incubated for 5 minutes. The enzyme reaction is stopped by adding 50  $\mu$ l/well 2 M  $H_2SO_4$ . The absorbance of light is measured at 492 nm in an EIA reader. Patient samples that contain Survivin will produce a positive reading, whereas those samples that do not contain Survivin will be negative.

#### Example 18

##### Survivin Fragments, Peptides and Small Molecule Antagonists

[0256] As described above, key functional residues in Survivin required for apoptosis have been identified. These data provide a template upon which to produce synthetic peptides and small molecule antagonists and competitive inhibitors of Survivin function. Preferably, the peptides are produced from native Survivin or include substitutions from the native Survivin peptide backbone that include the functionally relevant residues Trp<sup>67</sup>-Pro<sup>73</sup>-Cys<sup>84</sup>. Peptide fragments of native Survivin can be generated by standard techniques, including protein digests. A determination of which fragments compete with Survivin can readily be made by using the apoptosis measurement systems and apoptosis assay systems described above. These results provide a unique opportunity to identify a discrete linear sequence in Survivin, that is essential for inhibition of apoptosis.

[0257] Consistent with the general paradigm of IAP proteins-dependent inhibition of apoptosis, it also was predicted that a structural region in the molecule required for the anti-apoptotic function is the primary candidate for being a site of interaction with other molecules (such as binding partners). The functionally relevant peptide sequence in Survivin, based on the mutagenesis data, is: EGWEPD-DDPIEEHKKHSSGC. Ala substitutions of the underlined residues results in a complete loss of function of Survivin in transfected cells. This linear sequence can be synthesized and used as a much more stringent and specific reagent to isolate associated molecules using standard biochemical procedures of affinity chromatography or as a bait for the yeast two-hybrid system.

[0258] Also, preferably, the  $\beta$ COOH coiled-coil region of Survivin is included in Survivin fragments and peptides. Recent data indicates that this Survivin domain is important for the anti-apoptosis function of Survivin. We have generated a recombinant truncated form of Survivin lacking the last 40  $\beta$ COOH terminus amino acids comprising the coiled-coil domain. This truncated form was co-transfected with a lacZ plasmid in NIH3T3 cells side-by-side with wild type Survivin and XIAP, another member of the IAP gene family. The results, shown in FIG. 17, indicate that the truncated Survivin had lost most (~80%) of the cytoprotective effect at preventing apoptosis in transfected cells induced by hydrogen peroxide. Incidentally, in this system, Survivin was more potent than NAIP at preventing apoptosis.

[0259] Agonists and antagonists of Survivin also can readily be identified through conventional techniques.

Designed, synthetic peptides based on the native linear sequence also function as competitive inhibitors of Survivin's interaction with as yet unidentified partner molecules. However, this inhibition should be sufficient to block the anti-apoptosis function of Survivin.

[0260] A similar peptide-based strategy has been successful to block caspase activation in vitro and in vivo, protecting cells from apoptosis. See, e.g., Milligan, C. E. et al., (1995) *Neuron* 15:385-393.

#### Example 19

##### Therapeutic uses of Antisense Survivin DNA

[0261] As described above, the transcription of a Survivin antisense sequence altered the EPR-1/Survivin gene balance. This was demonstrated in HeLa cell transfectants, in which metallothionein-induced transcription of the EPR-1 "sense" strand suppressed the expression of Survivin and profoundly influenced apoptosis/cell proliferation. Additionally, transiently co-transfecting a Survivin antisense construct with a lacZ reported plasmid decreased the viability of Survivin antisense transfectants after a 48-h transfection in  $\beta$ -galactosidase expressing cells. Accordingly, the level of expression of Survivin in a Survivin expressing cell or tissue, such as a tumor, is decreased by transfecting the cell or tissue with the EPR-1 sense strand of DNA. Alternatively, a Survivin antisense-encoding DNA is used to transfect a target cell or tissue. Such therapy effectively decreases the translation of Survivin-encoding mRNA into Survivin protein.

#### Example 20

##### Use of Survivin as a Protective Agent Against Apoptosis

[0262] Survivin has been shown to protect cells from apoptosis when administered to cells that have been exposed to hydrogen peroxide or other agents that typically induce apoptosis. It is contemplated that cellular permeability may need to be increased, preferably in a transient manner in order to facilitate delivery of Survivin, or fragments thereof effective to reduce apoptosis. Certain conditions involving transient metabolic inhibition or transient hypoxia are likely to increase cellular permeability without the need for additional, external agents. Agents that may be appropriate include, metabolic inhibitors like 2-deoxyglucose and sodium azide. However, the ability of Survivin to mediate cytoprotection during a transient increase in cellular permeability offers the possibility of using therapeutic infusion of recombinant Survivin to decrease reperfusion injury and cellular damage during myocardial infarction and stroke. It is contemplated that such processes are mediated by increased tissue damage due to apoptosis. Treatment with Survivin could reduce the extent and magnitude of the injured tissue.

[0263] The use of Survivin or allelic variants of Survivin in subjects to modulate or prevent apoptosis related cell death would be beneficial in treating or ameliorating the effects of a variety of apoptosis-related indications. These indications include, but are not limited to, dermatological effects of aging (e.g., baldness that is caused by apoptosis of cells of hair follicle cells), disorders and diseases such as immunosuppression, gastrointestinal perturbations (e.g.,

damage of lining of the gut, ulcers, and radiation or chemotherapy induced damage), cardiovascular disorders, apoptosis related to reperfusion damage (e.g., coronary artery obstruction, cerebral infarction, spinal/head trauma and concomitant severe paralysis, damage due to insults such as frostbite or burns, and any indication previously thought to be treatable by superoxide dismutase), rejection of tissue transplantation (e.g., graft versus host disease), and Alzheimer's disease. The administration of Survivin also may be cytoprotective against chemotherapy or radiation-induced apoptosis.

[0264] Survivin protein for administration can be produced as described above, e.g., using the cDNA described herein. The protein may require purification for purposes of pharmaceutical administration and such purification steps preferably utilize monoclonal antibody separation and purification techniques as also described above.

[0265] In a clinical setting, Survivin is administered to patients in pharmaceutically effective dosages, i.e., in dosages effective to reduce the level or extent of apoptosis otherwise present, via several routes. For example, to treat dermatological ailments that involve apoptosis, Survivin can be administered in a salve, cream, ointment or powder form. Topical formulations may contain additional pharmaceutical or cosmetic compositions such as moisturizers, humectants, odor modifiers, buffer, pigment, preservatives, vitamins (such as A, C or E), emulsifiers, dispersing agents, wetting

agents, stabilizers, propellants, antimicrobial agents, sunscreen, enzymes and the like. Typical dosages of Survivin that may be administered to patients will be 0.01% to 1.0% by weight. Additional topical pharmaceutical compositions are described in S. Nakai et al., U.S. Pat. No. 5,672,603.

[0266] Survivin may also be administered, as may be appropriate for the condition being treated, in the form of pills, solutions, suspensions, emulsions, granules or capsules. Survivin can be administered orally; injected in solutions administered intravenously either alone or in admixture with conventional fluids for parenteral infusion (e.g., fluids containing glucose, amino acids etc.); injected intramuscularly, intradermally, subcutaneously or intraperitoneally; using suppositories; and in the form of ophthalmic solutions such as eye drops. Survivin can also be administered using delayed release carriers, such as liposomes, microsponges, microspheres or microcapsules that are deposited in close proximity to the tissue being treated for prevention of apoptosis related cell death.

[0267] Concentrations of Survivin or functional allelic variants of Survivin administered via routes other than topical administration typically would range in dose from about 10 µg per day to about 25 mg per day depending on the route of administration. Of course, it would be expected that skilled artisans, such as physicians, may alter these values on a case by case basis as required for the particular patient.

---

#### SEQUENCE LISTING

##### (1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 35

##### (2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 19 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
 (A) DESCRIPTION: /desc = "oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TGCTGGCCGC TCCTCCCTC

19

##### (2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 18 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
 (A) DESCRIPTION: /desc = "oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

ATGACCTCCA GAGGTTTC

18

---

-continued

---

## (2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 17 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Ala Pro Thr Leu Pro Pro Ala Trp Gln Pro Phe Leu Lys Asp His Ar  
1                   5                   10                   15  
Ile

## (2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Glu Gly Trp Glu Pro Asp Asp Asp Pro Ile Glu Glu His Lys Lys Hi  
1                   5                   10                   15  
Ser Ser Gly Cys  
                 20

## (2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GCGGGTGAGC TGTCCCTTGC AGATGGC 27

## (2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CCATGTAAGT TGATTTTCT AGAGAGG 27

## (2) INFORMATION FOR SEQ ID NO: 7:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

AATTGTATGT CTTTATTTC AGGCAAA

27

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: <Unknown>
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Glu Glu Ala Arg Leu Val Thr Phe Gln Asn Trp Pro Asp Ala Phe Leu  
 1 5 10 15  
 Thr Pro Gln Glu Leu Ala Lys Ala Gly Phe Tyr Tyr Leu Gly Arg Glu  
 20 25 30  
 Asp Gln Val Gln Cys Phe Ala Cys Gly Gly Lys Leu Ala  
 35 40 45

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: <Unknown>
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Glu Glu Ala Arg Phe Leu Thr Tyr Ser Met Trp Pro Leu Ser Phe Le  
1 5 10 15  
Ser Pro Ala Glu Leu Ala Arg Ala Gly Phe Tyr Tyr Ile Gly Pro Gl  
20 25 30  
Asp Arg Val Ala Cys Phe Ala Cys Gly Gly Lys Leu Ser  
35 40 45

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: <Unknown>
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Glu Ala Asn Arg Leu Val Thr Phe Lys Asp Trp Pro Asn Pro Asn Ile  
1 5 10 15

Thr Pro Gln Ala Leu Ala Lys Ala Gly Phe Tyr Tyr Leu Asn Arg Leu  
20 25 30

Asp His Val Lys Cys Val Trp Cys Asn Gly Val Ile Ala  
35 40 45

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 45 amino acids



-continued

(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

```
Glu Glu Val Arg Leu Asn Thr Phe Glu Lys Trp Pro Val Ser Phe Le
1           5           10           15
Ser Pro Glu Thr Met Ala Lys Asn Gly Phe Tyr Tyr Leu Gly Arg Se
          20           25           30
Asp Glu Val Arg Cys Ala Phe Cys Lys Val Glu Ile Met
          35           40           45
```

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 45 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

```
Lys Ala Ala Arg Leu Gly Thr Tyr Thr Asn Trp Pro Val Gln Phe Le
1           5           10           15
Glu Pro Ser Arg Met Ala Ala Ser Gly Phe Tyr Tyr Leu Gly Arg Gl
          20           25           30
Asp Glu Val Arg Cys Ala Phe Cys Lys Val Glu Ile Thr
          35           40           45
```

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 47 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

```
Glu Glu Ala Arg Leu Ala Ser Phe Arg Asn Trp Pro Phe Tyr Val Gl
1           5           10           15
Gly Ile Ser Pro Cys Val Leu Ser Glu Ala Gly Phe Val Phe Thr Gl
          20           25           30
Lys Gln Asp Thr Val Gln Cys Phe Ser Cys Gly Gly Cys Leu Gly
          35           40           45
```

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 45 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

```
Glu Ala Asn Arg Leu Val Thr Phe Lys Asp Trp Pro Asn Pro Asn Il
1           5           10           15
```

-continued

Thr Pro Gln Ala Leu Ala Lys Ala Gly Phe Tyr Tyr Leu Asn Arg Le  
                   20                  25                  30

Asp His Val Lys Cys Val Trp Cys Asn Gly Val Ile Ala  
           35                  40                  45

## (2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 46 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: <Unknown>
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

Glu Glu Ala Arg Leu Lys Ser Phe Gln Asn Trp Pro Asp Tyr Ala Hi  
   1                  5                  10                  15

Leu Thr Pro Arg Glu Leu Ala Ser Ala Gly Leu Tyr Tyr Thr Gly Il  
           20                  25                  30

Gly Asp Gln Val Gln Cys Phe Cys Cys Gly Gly Lys Leu Lys  
           35                  40                  45

## (2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 46 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: <Unknown>
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Glu Glu Ala Arg Leu Lys Ser Phe Gln Asn Trp Pro Asp Tyr Ala Hi  
   1                  5                  10                  15

Leu Thr Pro Arg Glu Leu Ala Ser Ala Gly Leu Tyr Tyr Thr Gly Al  
           20                  25                  30

Asp Asp Gln Val Gln Cys Phe Cys Cys Gly Gly Lys Leu Glu  
           35                  40                  45

## (2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 45 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: <Unknown>
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

Glu Asn Ala Arg Leu Leu Thr Phe Gln Thr Trp Pro Leu Thr Phe Le  
   1                  5                  10                  15

Ser Pro Thr Asp Leu Ala Arg Ala Gly Phe Tyr Tyr Thr Gly Pro Gl  
           20                  25                  30

Asp Arg Val Ala Cys Phe Ala Cys Gly Gly Lys Leu Ser  
           35                  40                  45

## (2) INFORMATION FOR SEQ ID NO: 18:

---

-continued

---

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 45 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Glu Glu Ala Arg Phe Leu Thr Tyr His Met Trp Pro Leu Thr Phe Le  
1                   5                   10                   15  
Ser Pro Ser Glu Leu Ala Arg Ala Gly Phe Tyr Tyr Ile Gly Pro Gl  
                  20                   25                   30  
Asp Arg Val Ala Cys Phe Ala Cys Gly Gly Lys Leu Ser  
          35                   40                   45

(2) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 46 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Glu Glu Ala Arg Leu Lys Ser Phe Gln Asn Trp Pro Asp Tyr Ala Hi  
1                   5                   10                   15  
Leu Thr Pro Arg Glu Leu Ala Ser Ala Gly Leu Tyr Tyr Thr Gly Il  
                  20                   25                   30  
Gly Asp Gln Val Gln Cys Phe Cys Cys Gly Gly Lys Leu Lys  
          35                   40                   45

(2) INFORMATION FOR SEQ ID NO: 20:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 45 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Glu Ala Asn Arg Leu Val Thr Phe Lys Asp Trp Pro Asn Pro Asn Il  
1                   5                   10                   15  
Thr Pro Gln Ala Leu Ala Lys Ala Gly Phe Tyr Tyr Leu Asn Arg Le  
                  20                   25                   30  
Asp His Val Lys Cys Val Trp Cys Asn Gly Val Ile Ala  
          35                   40                   45

(2) INFORMATION FOR SEQ ID NO: 21:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 50 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: <Unknown>  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

## -continued

---

Tyr Val Gly Ile Gly Asp Lys Val Lys Cys Phe His Cys Asp Gly Gl  
 1                    5                    10                    15  
 Leu Arg Asp Trp Glu Pro Gly Asp Asp Pro Trp Glu Glu His Ala Ly  
                   20                    25                    30  
 Trp Phe Pro Arg Cys Glu Phe Leu Leu Leu Ala Lys Gly Gln Glu Ty  
                   35                    40                    45  
 Val Ser  
       50

## (2) INFORMATION FOR SEQ ID NO: 22:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 50 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: <Unknown>  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Tyr Val Asp Arg Asn Asp Asp Val Lys Cys Phe Cys Cys Asp Gly Gl  
 1                    5                    10                    15  
 Leu Arg Cys Trp Glu Pro Gly Asp Asp Pro Trp Ile Glu His Ala Ly  
                   20                    25                    30  
 Trp Phe Pro Arg Cys Glu Phe Leu Ile Arg Met Lys Gly Gln Glu Ph  
                   35                    40                    45  
 Val Asp  
       50

## (2) INFORMATION FOR SEQ ID NO: 23:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 50 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: <Unknown>  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

Tyr Gln Lys Ile Gly Asp Gln Val Arg Cys Phe His Cys Asn Ile Gl  
 1                    5                    10                    15  
 Leu Arg Ser Trp Gln Lys Glu Asp Glu Pro Trp Phe Glu His Ala Ly  
                   20                    25                    30  
 Trp Ser Pro Lys Cys Gln Phe Val Leu Leu Ala Lys Gly Pro Ala Ty  
                   35                    40                    45  
 Val Ser  
       50

## (2) INFORMATION FOR SEQ ID NO: 24:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 49 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: <Unknown>  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

Tyr Thr Gly Tyr Gly Asp Asn Thr Lys Cys Phe Tyr Cys Asp Gly Gl

-continued

---

1	5	10	15
Leu Lys Asp Trp Glu Pro Glu Asp Val Pro Trp Glu Gln His Val Ar	20	25	30
Trp Phe Asp Arg Cys Ala Tyr Val Gln Leu Val Lys Gly Arg Asp Ty	35	40	45
Val			

(2) INFORMATION FOR SEQ ID NO: 25:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 49 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: <Unknown>
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

Tyr Thr Gly Gln Gly Asp Lys Thr Arg Cys Phe Cys Cys Asp Gly Gl	1	5	10	15
Leu Lys Asp Trp Glu Pro Asp Asp Ala Pro Trp Gln Gln His Ala Ar	20	25	30	
Trp Tyr Asp Arg Cys Glu Tyr Val Leu Leu Val Lys Gly Arg Asp Ph	35	40	45	
Val				

(2) INFORMATION FOR SEQ ID NO: 26:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 50 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: <Unknown>
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Tyr Thr Gly Ile Lys Asp Ile Val Gln Cys Phe Ser Cys Gly Gly Cy	1	5	10	15
Leu Glu Lys Trp Gln Glu Gly Asp Asp Pro Leu Asp Asp His Thr Ar	20	25	30	
Cys Phe Pro Asn Cys Pro Phe Leu Gln Asn Met Lys Ser Ser Ala Gl	35	40	45	
Val Thr	50			

(2) INFORMATION FOR SEQ ID NO: 27:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 50 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: <Unknown>
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

Tyr Gln Lys Ile Gly Asp Gln Val Arg Cys Phe His Cys Asn Ile Gl	1	5	10	15
Leu Arg Ser Trp Gln Lys Glu Asp Glu Pro Trp Phe Glu His Ala Ly				

(2) INFORMATION FOR SEO ID NO: 28:

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

(2) INFORMATION FOR SEQ ID NO: 29:

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

(2) INFORMATION FOR SEQ ID NO: 30:

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

Tyr Val Gly Asn Ser Asp Asp Val Lys Cys Phe Cys Cys Asp Gly Gl  
1 5 10 15  
Leu Arg Cys Trp Glu Ser Gly Asp Asp Pro Trp Val Gln His Ala Ly  
20 25 30

-continued

Trp Phe Pro Arg Cys Glu Tyr Leu Ile Arg Ile Lys Gly Gln Glu Ph  
 35 40 45

Ile Arg  
 50

(2) INFORMATION FOR SEQ ID NO: 31:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 50 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: <Unknown>  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

Tyr Val Gly Arg Asn Asp Asp Val Lys Cys Phe Gly Cys Asp Gly Gl  
 1 5 10 15

Leu Arg Cys Trp Glu Ser Gly Asp Asp Pro Trp Val Glu His Ala Ly  
 20 25 30

Trp Phe Pro Arg Cys Glu Phe Leu Ile Arg Met Lys Gly Gln Glu Ph  
 35 40 45

Val Asp  
 50

(2) INFORMATION FOR SEQ ID NO: 32:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 50 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: <Unknown>  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

Ala Leu Gly Glu Gly Asp Lys Val Lys Cys Phe His Cys Gly Gly Gl  
 1 5 10 15

Leu Thr Asp Trp Lys Pro Ser Glu Asp Pro Trp Glu Gln His Ala Ly  
 20 25 30

Trp Tyr Pro Gly Cys Lys Tyr Leu Leu Glu Gln Lys Gly Gln Glu Ty  
 35 40 45

Ile Asn  
 50

(2) INFORMATION FOR SEQ ID NO: 33:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 50 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: <Unknown>  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

Tyr Gln Lys Ile Gly Asp Gln Val Arg Cys Phe His Cys Asn Ile Gl  
 1 5 10 15

Leu Arg Ser Trp Gln Lys Glu Asp Glu Pro Trp Phe Glu His Ala Ly  
 20 25 30

-continued

Trp Ser Pro Lys Cys Gln Phe Val Leu Leu Ala Lys Gly Pro Ala Ty  
 35 40 45

Val Ser  
 50

## (2) INFORMATION FOR SEQ ID NO: 34:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 142 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: <Unknown>  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

Met Gly Ala Pro Thr Leu Pro Pro Ala Trp Gln Pro Phe Leu Lys As  
 1 5 10 15  
 His Arg Ile Ser Thr Phe Lys Asn Trp Pro Phe Leu Glu Gly Cys Al  
 20 25 30  
 Cys Thr Pro Glu Arg Met Ala Glu Ala Gly Phe Ile His Cys Pro Th  
 35 40 45  
 Glu Asn Glu Pro Asp Leu Ala Gln Cys Phe Phe Cys Phe Lys Glu Le  
 50 55 60  
 Glu Gly Trp Glu Pro Asp Asp Asp Pro Ile Glu Glu His Lys Lys Hi  
 65 70 75 80  
 Ser Ser Gly Cys Ala Phe Leu Ser Val Lys Lys Gln Phe Glu Glu Le  
 85 90 95  
 Thr Leu Gly Glu Phe Leu Lys Leu Asp Arg Glu Arg Ala Lys Asn Ly  
 100 105 110  
 Ile Ala Lys Glu Thr Asn Asn Lys Lys Lys Glu Phe Glu Glu Thr Al  
 115 120 125  
 Lys Lys Val Arg Arg Ala Ile Glu Gln Leu Ala Ala Met Asp  
 130 135 140

## (2) INFORMATION FOR SEQ ID NO: 35:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 14796 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

TCTAGACATG CGGATATATT CAAGCTGGGC ACAGCACAGC AGCCCCACCC CAGGCAGCTT 60  
 GAAATCAGAG CTGGGGTCCA AAGGGACCAC ACCCGAGGG ACTGTGTGGG GGTCTGGGGC 120  
 CACAGGCCAC TGCTTCCCC CGTCTTTCTC AGCCATTCCT GAAGTCAGCC TCACTCTGC 180  
 TCTCAGGGAT TTCAAATGTG CAGAGACTCT GGCACCTTTG TAGAAGCCCC TTCTGGTCC 240  
 AACTTACACC TGGATGCTGT GGGGCTGCAG CTGCTGCTCG GGCTCGGGAG GATGCTGGG 300  
 GCCCGGTGCC CATGAGCTTT TGAAGCTCCT GGAAGTCGGT TTTGAGGGTG TTCAGGTCC 360  
 GGTGGACACC TGGGCTGTCC TTGTCCATGC ATTTGATGAC ATTGTGTGCA GAAGTGAAA 420  
 GGAGTTAGGC CGGGCATGCT GGCTTATGCC TGTAATCCCA GCACTTTGGG AGGCTGAGG 480  
 GGTGGATCA CGAGGTCAGG AGTTCAATAC CAGCCTGGCC AAGATGGTGA AACCCCGTC 540



-continued

CTACTAAAAA TACAAAAAAA TTAGCCGGGC ATGGTGGCGG GCGCATGTAA TCCCAGCTA	600
TGGGGGGGCT GAGGCAGAGA ATTGCTGGAA CCCAGGAGAT GGAGGTGCA GTGAGCCAA	660
ATTGTGCCAC TGCACCTGCAC TCCAGCCTGG CGACAGAGCA AGACTCTGTC TCAAAAAAA	720
AAAAAAAAG TGAAAAGGAG TTGTTCCCTT CCTCCCTCCT GAGGGCAGGC AACTGCTGC	780
GTTGCCAGTG GAGGTGGTGC GTCCTTGGTC TGTGCCTGGG GGCACCCCA GCAGAGGCC	840
TGGTGGTGCC AGGGCCCGGT TAGCGAGCCA ATCAGCAGGA CCCAGGGGCG ACCTGCCAA	900
GTCAACTGGA TTTGATAACT GCAGCGAAGT TAAGTTTCCT GATTTTGTATG ATTGTGTTG	960
GTTTGTGTAA GAGAATGAAG TATTTGCGGG TAGTATGGTA ATGCCTTCAA CTTACAAA	1020
GTTCAGGTAA ACCACCCATA TACATACATA TACATGCATG TGATATATAC ACATACAG	1080
ATGTGTGTGT GTTCACATAT ATGAGGGGAG AGAGACTAGG GGAGAGAAAG TAGGTTGG	1140
AGAGGGAGAG AGAAAGGAAA ACAGGAGACA GAGAGAGAGC GGGGAGTAGA GAGAGGGA	1200
GGGTAAGAGA GGGAGAGGAG GAGAGAAAG GAGGAAGAAG CAGAGAGTGA ATGTTAAA	1260
AAACAGGCAA AACATAACA GAAAATCTGG GTGAAGGGTA TATGAGTATT CTTTGTAC	1320
TTCTTGCAAT TATCTTTTAT TTAAATTGAC ATCGGGCCGG GCGCAGTGGC TCACATCT	1380
AATCCCAGCA CTTTGGGAGG CCGAGGCAGG CAGATCACTT GAGGTCAGGA GTTTGAGA	1440
AGCCTGGCAA ACATGGTGAA ACCCCATCTC TACTAAAAAT AAAAAATTA GCCTGGTG	1500
GTGGTGCATG CCTTTAATCT CAGCTACTCG GGAGGCTGAG GCAGGAGAAT CGCTTGAA	1560
CGTGCGGGG AGGAGGTTGC AGTGAGCTGA GATCATGCCA CTGCACTCCA GCCTGGGC	1620
TAGAGCGAGA CTCAGTTTCA AATAAATAA TAAACATCAA AATAAAAAGT TACTGTAT	1680
AAGAATGGGG GCGGGGTGGG AGGGGTGGGG AGAGGTTGCA AAAATAAATA AATAAATA	1740
TAAACCCCAA AATGAAAAAG ACAGTGGAGG CACCAGGCCT GCGTGGGGCT GGAGGGCT	1800
TAAGGCCAGG CCTCTTATCT CTGGCCATAG AACCAGAGAA GTGAGTGGAT GTGATGCC	1860
GCTCCAGAAG TGAATCCAGA ACACCCTGTT CCAAAGCAGA GGACACACTG ATTTTTTTT	1920
TAATAGGCTG CAGGACTTAC TGTGTGGTGG ACGCCCTGCT TTGCGAAGGG AAAGGAGG	1980
TTTGCCCTGA GCACAGGCCC CCACCCTCCA CTGGGCTTTC CCCAGCTCCC TTGTCTTC	2040
ATCACGGTAG TGGCCCAGTC CCTGGCCCCT GACTCCAGAA GGTGGCCCTC CTGGAAAC	2100
AGGTCGTGCA GTCAACGATG TACTCGCCGG GACAGCGATG TCTGCTGCAC TCCATCCC	2160
CCCTGTTCAT TTGTCCCTCA TGCCCGTCTG GAGTAGATGC TTTTTCGAGA GGTGGCAC	2220
TGTAAAGCTC TCCTGTCTGA CTTTTTTTTT TTTTITAGAC TGAGTTTTCG TCTTGTTC	2280
TAGGCTGGAG TGCAATGGCA CAATCTCAGC TCACTGCACC CTCTGCCTCC CGGGTTCA	2340
CGATTCTCCT GCCTCAGCCT CCCGAGTAGT TGGGATTACA GGCATGCACC ACCACGCC	2400
GCTAATTTTT GTATTTTTAG TAGAGACAAG GTTTCACCGT GATGGCCAGG CTGGTCTT	2460
ACTCCAGGAC TCAAGTGATG CTCCTGCCTA GGCCTCTCAA AGTGTTGGGA TTACAGGC	2520
GAGCCACTGC ACCCGGCCCTG CACGCGTTCT TTGAAAGCAG TCGAGGGGGC GCTAGGTG	2580
GGCAGGGACG AGCTGGCGCG GCGTCGCTGG GTGCACCGCG ACCACGGGCA GAGCCACG	2640
GCGGGAGGAC TACAATCCC GGCACACCCC GCGCCGCCCC GCCTCTACTC CCAGAAGG	2700
GCGGGGGGTG GACCGCCTAA GAGGGCGTGC GCTCCCGACA TGCCCCGCGG CGCGCCAT	2760
ACCGCCAGAT TTGAATCGCG GGACCCGTTG GCAGAGGTGG CGGCGGCGGC ATGGGTGC	2820

-continued

---

CGACGTTGCC	CCCTGCCTGG	CAGCCCTTTC	TCAAGGACCA	CCGCATCTCT	ACATTCAA	2880
ACTGGCCCTT	CTTGAGAGGC	TGCGCCTGCA	CCCCGGAGCG	GGTGAGACTG	CCCGGCCT	2940
TGGGGTCCCC	CACGCCCGCC	TTGCCCTGTC	CCTAGCGAGG	CCACTGTGAC	TGGGCCTC	3000
GGGTACAAGC	CGCCCTCCCC	TCCCCGTCCT	GTCCCCAGCG	AGGCCACTGT	GGCTGGGC	3060
CTTGGGTCCA	GGCCGGCCTC	CCCTCCCTGC	TTTGTCCCCA	TCGAGGCCTT	TGTGGCTG	3120
CCTCGGGGTT	CCGGGCTGCC	ACGTCCACTC	ACGAGCTGTG	CTGTCCCTTG	CAGATGGC	3180
AGGCTGGCTT	CATCCACTGC	CCCACTGAGA	ACGAGCCAGA	CTTGGCCCAG	TGTTTCTT	3240
GCTTCAAGGA	GCTGGAAGGC	TGGGAGCCAG	ATGACGACCC	CATGTAAGTC	TTCTCTGG	3300
AGCCTCGATG	GGCTTTGTTT	TGAAC TGAGT	TGTCAAAAGA	TTTGAGTTGC	AAAGACAC	3360
AGTATGGGAG	GGTTGCTTTC	CACCCCTATT	GCTTCTTAAA	CAGCTGTTGT	GAACGGAT	3420
CTCTCTATAT	GCTGGTGCCT	TGGTGATGCT	TACAACCTAA	TTAAATCTCA	TTTGACCA	3480
ATGCCTTGGG	GTGGACGTAA	GATGCCTGAT	GCCTTTCATG	TTCAACAGAA	TACATCAG	3540
GACCCGTGTT	TGTGAACTC	CCAGGAATGT	CCAAGTGCTT	TTTTTGAGAT	TTTTTAAA	3600
ACAGTTTAAT	TGAAATATAA	CCTACACAGC	ACAAAAATTA	CCCTTTGAAA	GTGTGCAC	3660
CACACTTTCG	GAGGCTGAGG	CGGGCGGATC	ACCTGAGGTC	AGGAGTTCAA	GACCTGCC	3720
GCCAACTTGG	CGAAACCCCG	TCTCTACTAA	AAATACAAAA	ATTAGCCGGG	CATGGTAG	3780
CACGCCCGTA	ATCCCAGCTA	CTCGGGAGGC	TAAGGCAGGA	GAATCGCTTG	AACCTGGG	3840
GCGGAGGTTG	CAGTGAGCCG	AGATTGTGCC	AATGCACTCC	AGCCTCGGCG	ACAGAGCG	3900
ACTCCGTCAT	AAAAATAAAA	AATTGAAAAA	AAAAAAGAA	AGAAAGCATA	TACTTCAG	3960
TTGTTCTGGA	TTTTTTTCTT	CAAGATGCCT	AGTTAATGAC	AATGAAATTC	TGTACTCG	4020
TGGTATCTGT	CTTCCACAC	TGTAATGCCA	TATTCTTTTC	TCACCTTTTT	TTCTGTCG	4080
TTCACTTGCT	TCCACAGCTT	TAATTTTTTT	CCCCTGGAGA	ATCACCCAG	TTGTTTTT	4140
TTTTGGCCAG	AAGAGAGTAG	CTGTTTTTTT	TCTTAGTATG	TTTGCTATGG	TGTTTATA	4200
GCATCCCCGT	AATCACTGGG	AAAAGATCAG	TGGTATTCTT	CTTGAAAATG	AATAAGTG	4260
ATGATATTTT	CAGATTAGAG	TTACAAC TGG	CTGTCTTTTT	GGACTTTGTG	TGGCCATG	4320
TTCATTGTAA	TGCAGTTCTG	GTAACGGTGA	TAGTCAGTTA	TACAGGGAGA	CTCCCCTA	4380
AGAAAATGAG	AGTGTGAGCT	AGGGGGTCCC	TTGGGGAACC	CGGGGCAATA	ATGCCCTT	4440
CTGCCCTTAA	TCCTTACAGT	GGCCCGGGCA	CGGTGGCTTA	CGCCTGTAAT	ACCAGCAC	4500
TGGGAGGCCG	AGGCGGGCGG	ATCACGAGGT	CAGGAGATCG	AGACCATCTT	GGCTAATA	4560
GTGAAACCCC	GTCTCCACTA	AAAATACAAA	AAATTAGCCG	GGCGTGGTGG	TGGGCGCC	4620
TAGTCCCAGC	TACTCGGGAG	GCTGAGGCAG	GAGAATGGCG	TGAACCCAGG	AGGCGGAG	4680
TGCAGTGAGC	CGAGATTGCA	CCACTGCACT	CCAGCCTGGG	CGACAGAATG	AGACTCCG	4740
TCAAAAAAAA	AAAAAAAAGA	AAAAAATCTT	TACAGTGGAT	TACATAACAA	TTCCAGTG	4800
ATGAAATTAC	TTCAACAGT	TCCTTGAGAA	TGTTGGAGGG	ATTTGACATG	TAATTCCT	4860
GGACATATAC	CATGTAACAC	TTTTCCAAC T	AATTGCTAAG	GAAGTCCAGA	TAAAATAG	4920
ACATTAGCCA	CACAGATGTG	GGGGGAGATG	TCCACAGGGA	GAGAGAAGGT	GCTAAGAG	4980
GCCATATGGG	AATGTGGCTT	GGGCAAGACA	CTGATGCCAT	CAACTTCAGA	CTTGACGT	5040
TACTCCTGAG	GCAGAGCAGG	GTGTGCCTGT	GGAGGGCGTG	GGGAGGTGGC	CCGTGGGG	5100

-continued

---

TGGACTGCCG	CTTTAATCCC	TTCAGCTGCC	TTTCCGCTGT	TGTTTGTATT	TTTCTAGA	5160
GGAACATAAA	AAGCATTCGT	CCGGTTGCGC	TTTCCTTTCT	GTCAAGAAGC	AGTTTGAA	5220
ATTAACCCCTT	GGTGAATTTT	TGAAACTGGA	CAGAGAAAGA	GCCAAGAACA	AAATTGTA	5280
TATTGGGAAT	AAGAACTGCT	CAAACCTGT	TCAATGTCTT	TAGCACTAAA	CTACCTAG	5340
CCTCAAAGGG	ACTCTGTGTT	TTCTCAGGA	AGCATTTTTT	TTTTTTTTCT	GAGATAGA	5400
TTCACTCTTG	TTGCCCAGGC	TGGAGTGCAA	TGGTGCAATC	TTGGCTCACT	GCAACCTC	5460
CCTCTCGGGT	TCAAGTGATT	CTCCTGCCCT	AGCCTCCCAA	GTAAC TGGA	TTACAGGG	5520
GTGCCACCAC	ACCCAGCTAA	TTTTTGATTT	TTTAGTAGAG	ATGGGGTTTC	ACCACATT	5580
CCAGGCTGGT	CTTGAATCC	TGACCTCGTG	ATTGCGCCAC	CTTGGCCTCC	CAAAGTGC	5640
GGATTACAGG	CGTGAACCAC	CACGCCTGGC	TTTTTTTTTT	TTGTTCTGAG	ACACAGTT	5700
ACTCTGTTAC	CCAGGCTGGA	GTAGGGTGGC	CTGATCTCGG	ATCACTGCAA	CCTCCGCC	5760
CTGGGCTCAA	GTGATTTGCC	TGCTTCAGCC	TCCAAGTAG	CCGAGATTAC	AGGCATGT	5820
CACCACACCC	AGGTAATTTT	TGTATTTTGT	GTAGAGACGA	GGTTTCACCA	TGTTGGCC	5880
GCTGGTTTTG	AACTCCTGAC	CTCAGGTGAT	CCACCCGCCT	CAGCCTCCCA	AAGTGCTG	5940
ATTATAGGTG	TGAGCCACCA	CACCTGGCCT	CAGGAAGTAT	TTTTATTTTT	AAATTTAT	6000
ATTTATTTGA	GATGGAGTCT	TGCTCTGTCT	CCCAGGCTAG	AGTGCAGCGA	CGGGATCT	6060
GCTCACTGCA	AGCTCCGCCC	CCAGGTTC	AGCCATTCTC	CTGCCTCAGC	CTCCCGAG	6120
GCTGGGACTA	CAGGCGCCCC	CCACCACACC	CGGCTAATTT	TTTTGTATTT	TTAGTAGA	6180
CGGGTTTTCA	CCGTGTTAGC	CAGGAGGGTC	TTGATCTCCT	GACCTCGTGA	TCTGCCTG	6240
TCGGCCTCCC	AAAGTGCTGG	GATTACAGGT	GTGAGCCACC	ACACCCGGCT	ATTTTTAT	6300
TTTTGAGACA	GGGACTCACT	CTGTCACCTG	GGCTGCAGTG	CAGTGGTACA	CCATAGCT	6360
CTGCAGCCTC	GAACTCCTGA	GCTCAAGTGA	TCCTCCCACC	TCATCCTCAC	AAGTAATT	6420
GACTACAGGT	GCACCCACCC	ATGCCACCT	AATTTATTTA	TTTATTTATT	TATTTATT	6480
CATAGAGATG	AGGGTTCCTT	GTGTTGTCCA	GGCTGGTCTT	GAACTCCTGA	GCTCACGG	6540
TCCTTTTGCC	TGGGCCTCCC	AAAGTGCTGA	GATTACAGGC	ATGAGCCACC	GTGCCCAG	6600
AGGAATCATT	TTTAAAGCCC	CTAGGATGTC	TGTGTGATTT	TAAAGCTCCT	GGAGTGTG	6660
CGGTATAAGT	ATATACCGGT	ATAAGTAAAT	CCCACATTTT	GTGTCAGTAT	TTACTAGA	6720
CTTAGTCATT	TATCTGAAGT	TGAAATGTAA	CTGGGCTTTA	TTTATTTATT	TATTTATT	6780
TTTATTTTTA	ATTTTTTTTT	TTGAGACGAG	TCTCACTTTG	TCACCCAGGC	TGGAGTGC	6840
TGGCAGCATC	TCGGCTCACT	GCAACCTCTG	CCTCCCGGGG	TCAAGCGATT	CTCCTGCC	6900
AGCCTCCCGA	GTAGCTGGGA	CTACAGGCAC	GCACCACCAT	GCCTGGCTAA	TTTTTGTA	6960
TTTAGTAGAC	GGGGTTTAC	CATGCTGGCC	AAGCTGGTCT	CAAACCTCTG	ACCTTGTG	7020
CTGCCCCTTT	TAGCCTCCCA	GAGTGCTGGG	ATTACAGGCA	TGAGCCACCA	TGCGTGGT	7080
TTTTAAATTT	TTTGTATTTT	TTTTTTTTTT	GAGACAGAGC	CTTGCTCTGT	CGCCCAGG	7140
GGAGTGCAGT	GGCAGCATCT	CAGCTCACTA	CAAGCTCCGC	CTCCCGGGT	CACGCCAT	7200
TTCTGCCTCA	GCCTCCTGAG	TAGCTGGGAC	TACAGGTGCC	CACCACCACG	CCTGGCTA	7260
TTTTTTTGGT	ATTTTTTATTA	GAGACAAGGT	TTCATCATGT	TGGCCAGGCT	GGTCTCAA	7320
TCCTGACCTC	AAGTGATCTG	CCTGCCTCGG	CCTCCCAAAG	CGCTGAGATT	ACAGGTGT	7380

-continued

TCTACTGCGC	CAGGCCTGGG	CGTCATATAT	TCTTATTTGC	TAAGTCTGGC	AGCCCCAC	7440
AGAATAAGTA	CTGGGGGATT	CCATATCCTT	GTAGCAAAGC	CCTGGGTGGA	GAGTCAGG	7500
ATGTTGTAGT	TCTGTCTCTG	CCACTTGCAG	ACTTTGAGTT	TAAGCCAGTC	GTGCTCAT	7560
TTTCCTTGCT	AAATAGAGGT	TAGACCCCT	ATCCCATGGT	TTCTCAGGTT	GCTTTTCA	7620
TTGAAAATTG	TATTCCTTTG	TAGAGATCAG	CGTAAAAATA	TTCTGTCCTT	ATATGTGG	7680
TTATTTTAAT	TTGAGACAGA	GTGTCACTCA	GTGCCCCAGG	CTGGAGTGTG	GTGGTGCG	7740
CTTGCGTCAC	TGCGACCTCC	ACCTCCCAGG	TTCAAGCGAT	TCTCGTGCCT	CAGGCTCC	7800
AGTAGCTGAG	ATTATAGGTG	TGTGCCACCA	GGCCCAGCTA	ACTTTTGTAT	TTTTAGTA	7860
GACAGGGTTT	TGCCATGTTG	GCTAAGCTGG	TCTCGAACTC	CTGGCCTCAA	GTGATCTG	7920
CGCCTTGCCA	TCCCAAAGTG	CTGGGATTAC	AGGTGTGAAC	CACCACACCT	GGCCTCAA	7980
TAGTGGCTTT	TAAGTGCTAA	GGACTGAGAT	TGTGTTTGTG	CAGGAAGAGG	CCAGTTGT	8040
GTGAAGCATG	CTGTGAGAGA	GCTTGTCAAC	TGGTTGAGGT	TGTGGGAGCT	GCAGCGTG	8100
AACTGGAAAG	TGGGCTGGGG	ATCATCTTTT	TCCAGGTCAG	GGGTCAGCCA	GCTTTTCT	8160
AGCGTGCCAT	AGACCATCTC	TTAGCCCTCG	TGGGTCAGAG	TCTCTGTTGC	ATATTGTC	8220
TTGTTGTTTT	TCACAACCTT	TTAGAAACAT	AAAAAGCATT	CTTAGCCCGT	GGGCTGGA	8280
AAAAAAGGCC	ATGACGGGCT	GTATGGATTT	GGCCCAGCAG	GCCCTTGCTT	GCCAAGCC	8340
GTTTTAGACA	AGGAGCAGCT	TGTGTGCCTG	GAACCATCAT	GGGCACAGGG	GAGGAGCA	8400
GTGGATGTGG	AGGTGTGAGC	TGGAACCAG	GTCCCAGAGC	GCTGAGAAAG	ACAGAGGG	8460
TTTGCCCTTG	CAAGTAGAGC	AACTGAAATC	TGACACCATC	CAGTTCAGAG	AAGCCCTG	8520
GTGCTGGTGG	ACGCTGCGGG	GTGCTCCGCT	CTAGGGTTAC	AGGGATGAAG	ATGCAGTC	8580
GTAGGGGGAG	TCCACTCACC	TGTTGAAGA	TGTGATTAAG	AAAAGTAGAC	TTTCAGGG	8640
GGGCATGGTG	GCTCAGCCT	GTAATCCAG	CACCTTGGA	GGCCGAGGCG	GGTGGATC	8700
GAGGTCAGGA	GATCGAGACC	ATCCTGGCTA	ACATGGTGAA	ACCCCGTCTT	TACTAAAA	8760
ACAAAAAATT	AGCTGGGCGT	GGTGGCGGGC	GCCTGTAGTC	CCAGCTACTC	GGGAGGCT	8820
GGCAGGAGAA	TGGCGTGAAC	CTGGGAGGTG	GAGCTTGCTG	TGAGCCGAGA	TCGCGCCA	8880
GCACTCCAGC	CTGGGCGACA	GAGCGAGACT	CCGTCTCAAA	AAAAAAAAAA	AAAGTAGG	8940
TTCATGATGT	GTGAGCTGAA	GGCGCAGTAG	GCAGAAGTAG	AGGCCTCAGT	CCCTGCAG	9000
GACCCCTCGG	TCTCTATCTC	CTGATAGTCA	GACCCAGCCA	CACTGGAAAG	AGGGGAGA	9060
TTACAGCCTG	CGAGAAAAGT	AGGGAGATTT	AAAAACTGCT	TGGCTTTTAT	TTTGAACT	9120
TTTTTTTGTT	TGTTTGTTTT	CCCCAATTCA	GAATACAGAA	TACTTTTATG	GATTTGTT	9180
TATTACTTTA	ATTTTGAAAC	AATATAATCT	TTTTTTTGTT	GTTTTTTTGA	GACAGGGT	9240
TACTCTGTCA	CCCAGGCTGA	GTGCAGTGGT	GTGATCTTGG	CTCACCTCAG	CCTCGACC	9300
CTGGGCTCAA	ATGATTCTCC	CACCTCAGCT	TCCCAAGTAG	CTGGGACCAC	AGGTGCGT	9360
GTTGCGCTAT	ACAAATCCTG	AAGACAAGGA	TGCTGTTGCT	GGTGATGCTG	GGGATTCC	9420
AGATCCCAGA	TTTGATGGCA	GGATGCCCTT	GTCTGCTGCC	TTGCCAGGGT	GCCAGGAG	9480
CGCTGCTGTG	GAGCTGAGG	CCCGGCCATC	CAGGGCGATG	CATTGGGCGC	TGATTCTT	9540
TCCTGCTGCT	GCCTCGGTGC	TTAGCTTTTG	AAACAATGAA	ATAAATTAGA	ACCAGTGT	9600
AAATCGATCA	GGGAATAAAT	TTAATGTGGA	AATAAACTGA	ACAACTTAGT	TCTTCATA	9660

-continued

---

AGTTTACTTG	GTAAATACTT	GTGATGAGGA	CAAAACGAAG	CACTAGAAGG	AGAGGCGA	9720
TGTAGACCTG	GGTGGCAGGA	GTGTTTGTGT	TGTTTCTTTT	GGCAGGGTCT	TGCTCTGT	9780
CTCAGGCTGG	AGTACAGTGG	CACAATCACA	GCTCACTATA	GCCTCGACCT	CCTGGACT	9840
AGCAATCCTC	CTGCCTCAGC	CTCCCAGTAG	CTGGGACTAC	AGGCGCATGC	CACCATGC	9900
GGCTAATTTT	AAATTTTTTT	TTTTCTCTTT	TTTGAGATGG	AATCTCACTC	TGTCGCCC	9960
GCTGGAGTGC	AGTGGCGTGA	TCTCGGCTGA	CGGCAAGCTC	CGCCTCCCAG	GTTCACT	10020
TTGCCTGCC	TCAGCCTCCC	AAGTAGCTGG	GACTACAGGC	GCTGGGATTA	CAAACCC	10080
CCCAAAGTGC	TGGGATTACA	GGCGTGAGCC	ACTGCACCCG	GCCTGTTTTG	TCTTTCA	10140
GCAAGAGTTG	TGTTTGCTTC	GCCCCACCT	TTAGTGGA	AAATGTATAA	ATGGAGA	10200
TGACCTCCAC	ATTGGGGTGG	TTAAATTATA	GCATGTATGC	AAAGGAGCTT	CGCTAAT	10260
AGGCTTTTTT	GAAAGAGAAG	AACTGAATA	ATCCATGTGT	GTATATATAT	TTTAAAA	10320
ATGGTCATCT	TTCCATATCA	GTAAAGCTGA	GGCTCCCTGG	GACTGCAGAG	TTGTCCA	10380
CAGTCCATTA	TAAGTGCGCT	GCTGGGCCAG	GTGCAGTGGC	TTGTGCCTGA	ATCCCAG	10440
TTTGGGAGGC	CAAGGCAGGA	GGATTCAATTG	AGCCCAGGAG	TTTGGAGCG	AGCCTGG	10500
ATGTGGCCAG	ACCTCATCTC	TTCAAAAAAT	ACACAAAAAA	TTAGCCAGGC	ATGGTGG	10560
GTGCCTGTAG	TCTCAGCTAC	TCAGGAGGCT	GAGGTGGGAG	GATCACTTTG	AGCCTTG	10620
GTCAAAGCTG	CAGTAAGCCA	TGATCTTGCC	ACTGCATTCC	AGCCTGGATG	ACAGAGC	10680
ACCCTGTCTC	TAAAAA	AAAAACCAA	CGGTGCACTG	TTTTCTTTTT	TCTTATC	10740
TTATTATTTT	TAAATTAAAT	TTTCTTTTAA	TAATTTATAA	ATTATAAAT	TATATTA	10800
AATGACAAAT	TTTTATTACT	TATACATGAG	GTAAACTTA	GGATATATAA	AGTACAT	10860
GAAAAGTAAT	TTTTTGCTG	GCACAGTGGC	TCACACCTGT	AATCCCAGCA	CTTTGGG	10920
CCGTGGCGGG	CAGATCACAT	GAGATCATGA	GTTGAGAGCC	AACCTGACCA	ACATGGA	10980
ACCCCATCTC	TACTAAAAAT	ACAAAATTAG	CCGGGTGGT	GGCGCATGCC	TGTAATC	11040
GCTACTCGGG	AGGCTGAGGC	AGGAGAATCT	CTTGAACCCG	GGAGGCAGAG	GTTGCGG	11100
GCCAAGATCG	TGCCTTTGCA	CACCAGCCTA	GGCAACAAGA	GCGAAAGTCC	GCTCTAA	11160
AAAAGTAATT	TTTTTTAAGT	TAACCTCTGT	CAGCAACAA	ATTTAACCCA	ATAAAGG	11220
TTGTTTTTTA	ATGTAGTAGA	GGAGTTAGGG	TTTATAAAAA	ATATGGTAGG	GAAGGGG	11280
CCTGGATTGG	CTAATGTGAT	TGTCATTGTC	CCCTTAGGAG	AGAGCTCTGT	TAGCAGA	11340
AAAAAATTGG	AAGCCAGATT	CAGGGAGGGA	CTGGAAGCAA	AAGAATTTCT	GTTGAGG	11400
GAGCCTGATG	TTTGCCAGGG	TCTGTTTAAC	TGGACATGAA	GAGGAAGGCT	CTGGACT	11460
CTCCAGGAGT	TTCAGGAGAA	AGGTAGGGCA	GTGGTTAAGA	GCAGAGCTCT	GCCTAGA	11520
GCTGGGGTGC	CTAGACTAGC	TGGGGTGCCC	AGACTAGCTG	GGGTGCCTAG	ACTAGCT	11580
TACTTTGAGT	GGCTCCTTCA	GCCTGGACCT	CGGTTTCCTC	ACCTGTATAG	TAGAGAT	11640
GGAGCACCCA	GCGCAGGATC	ACTGTGAACA	TAAATCAGTT	AATGGAGGAA	GCAGGTA	11700
TGGTGCTGGG	TGCATACCAA	GCACTCCGTC	AGTGTTTCCT	GTTATTCGAT	GATTAGG	11760
CAGCTTAAAC	TAGAGGGAGT	TGAGCTGAAT	CAGGATGTTT	GTCCCAGGTA	GCTGGGA	11820
TGCCTAGCCC	AGTGCCAGT	TTATTTAGGT	GCTCTCTCAG	TGTTCCCTGA	TTGTTTT	11880
CTTTGTCATC	TTATCTACAG	GATGTGACTG	GGAAGCTCTG	GTTTCAGTGT	CATGTGT	11940

-continued

TTCTTTATTT	CCAGGCAAAG	GAAACCAACA	ATAAGAAGAA	AGAATTTGAG	GAAACTG	12000
AGAAAGTGCG	CCGTGCCATC	GAGCAGCTGG	CTGCCATGGA	TTGAGGCCTC	TGGCCGG	12060
TGCCTGGTCC	CAGAGTGGCT	GCACCACTTC	CAGGGTTTAT	TCCCTGGTGC	CACCAGC	12120
CCTGTGGGCC	CCTTAGCAAT	GTCTTAGGAA	AGGAGATCAA	CATTTTCAAA	TTAGATG	12180
CAACTGTGCT	CCTGTTTTGT	CTTGAAAGTG	GCACCAGAGG	TGCTTCTGCC	TGTGCAG	12240
GTGCTGCTGG	TAACAGTGGC	TGCTTCTCTC	TCTCTCTCTC	TTTTTTGGGG	GCTCATT	12300
GCTGTTTTGA	TTCCCGGGCT	TACCAGGTGA	GAAGTGAGGG	AGGAAGAAGG	CAGTGTC	12360
TTTGCTAGAG	CTGACAGCTT	TGTTGCGCTG	GGCAGAGCCT	TCCACAGTGA	ATGTGTC	12420
ACCTCATGTT	GTTGAGGCTG	TCACAGTCCT	GAGTGTGGAC	TTGGCAGGTG	CCTGTTG	12480
CTGAGCTGCA	GGTTCCTTAT	CTGTACACCC	TGTGCCTCCT	CAGAGGACAG	TTTTTTTT	12540
GTTGTGTTTT	TTTGTTTTTT	TTTTTTGGTA	GATGCATGAC	TTGTGTGTGA	TGAGAGA	12600
GAGACAGAGT	CCCTGGCTCC	TCTACTGTTT	AACAACATGG	CTTCTTTATT	TTGTTTG	12660
TGTTAATTCA	CAGAATAGCA	CAAACTACAA	TTAAACTTAA	GCACAAAGCC	ATTCTAA	12720
ATTGGGGAAA	CGGGGTGAAC	TTCAGGTGGA	TGAGGAGACA	GAATAGAGTG	ATAGGAA	12780
TCTGGCAGAT	ACTCCTTTTG	CCACTGCTGT	GTGATTAGAC	AGGCCCAGTG	AGCCGCG	12840
CACATGCTGG	CCGCTCCTCC	CTCAGAAAAA	GGCAGTGGCC	TAAATCCTTT	TTAAATG	12900
TGGCTCGATG	CTGTGGGGGA	CTGGCTGGGC	TGCTGCAGGC	CGTGTGCTCG	TCAGCCC	12960
CTTCACATCT	GTCACGTTCT	CCACACGGGG	GAGAGACGCA	GTCCGCCCAG	GTCCCCG	13020
TCTTTGGAGG	CAGCAGCTCC	CGCAGGGCTG	AAGTCTGGCG	TAAGATGATG	GATTTGA	13080
GCCCTCCTCC	CTGTCATAGA	GCTGCAGGGT	GGATTGTTAC	AGCTTCGCTG	GAAACCT	13140
GAGGTCATCT	CGGCTGTTCC	TGAGAAATAA	AAAGCCTGTC	ATTTCAAACA	CTGCTGT	13200
CCCTACTGGG	TTTTTAAAAA	ATTGTCAGTT	TTTCATCGTC	GTCCCTAGCC	TGCCAAC	13260
CATCTGCCCA	GACAGCCGCA	GTGAGGATGA	GCGTCCTGGC	AGAGACGCAG	TTGTCTC	13320
GCGCTTGCCA	GAGCCACGAA	CCCCAGACCT	GTTTGATCA	TCCGGGCTCC	TTCCGGG	13380
AAACAACTGA	AAATGCACTT	CAGACCCACT	TATTTATGCC	ACATCTGAGT	CGGCCTG	13440
TAGACTTTTC	CCTCTAAACT	GGGAGAATAT	CACAGTGGTT	TTTGTAGCA	GAAAATG	13500
TCCAGCCTCT	GTACTCATCT	AAGCTGCTTA	TTTTTGATAT	TTGTGTCAGT	CTGTAAA	13560
ATACTTCACT	TTAATAACTG	TTGCTTAGTA	ATTGGCTTTG	TAGAGAAGCT	GGAAAAA	13620
GGTTTTGTCT	TCAACTCCTT	TGCATGCCAG	GCGGTGATGT	GGATCTCGGC	TTCTGTG	13680
CTGTGCTGTG	GGCAGGGCTG	AGCTGGAGCC	GCCCCTCTCA	GCCCGCCTGC	CACGGCC	13740
CCTTAAAGGC	CATCCTTAAA	ACCAGACCCT	CATGGCTGCC	AGCACCTGAA	AGCTTCC	13800
ACATCTGTTA	ATAAAGCCGT	AGGCCCTTGT	CTAAGCGCAA	CCGCCTAGAC	TTTCTTT	13860
ATACATGTCC	ACATGTCCAT	TTTTCAGGTT	CTCTAAGTTG	GAGTGGAGTC	TGGGAAG	13920
TGTGAATGAG	GCTTCTGGGC	TATGGGTGAG	GTTCCAATGG	CAGTTAGAG	CCCCTCG	13980
CAACTGCCAT	CCTGAAAAGT	AGAGACAGCA	GTGCCCCTG	CCCAGAAGAG	ACCAGCA	14040
CAAACTGGAG	CCCCCATTGC	AGGCTGTCGC	CATGTGGAAA	GAGTAACTCA	CAATTGC	14100
TAAAGTCTCA	TGTGGTTTTA	TCTACTTTTT	TTTTCTTTTT	CTTTTTTTTT	GAGACAA	14160
CTTGCCCTCC	CAGGCTGGAG	TGCAGTGAA	TGACCACAGC	TCACCGCAAC	CTCAAAT	14220

-continued

---

TGCGTTCAAG TGAACCTCCC ACTTTAGCCT CCCAAGTAGC TGGGACTACA GCGGCAC	14280
ATCACACCCG GCTAATTGAA AAATTTTTTT TTTTGTTTAG ATGGAATCTC ACTTTGT	14340
CCAGGCTGGT CTCAAACTCC TGGGCTCAAG TGATCATCCT GCTTCAGCGT CCGACTT	14400
GGTATTATAG GCGTGAGCCA CTGGGCTGA CCTAGCTACC ATTTTTTAAT GCAGAAA	14460
AGACTTGTAG AAATGAAATA ACTTGTCCAG GATAGTCGAA TAAGTAACTT TTAGAGC	14520
GATTTGAACC CAGGCAATCT GGCTCCAGAG CTGGGCCCTC ACTGCTGAAG GACTCTG	14580
GCTTGGGAGG GTGGCTATGG TCGGCTGTCT GATTCTAGGG AGTGAGGGCT GTCTTTA	14640
CACCCCATTC CATTTTCAGA CAGCTTTGTC AGAAAGGCTG TCATATGGAG CTGACAC	14700
CCTCCCCAAG GCTTCCATAG ATCCTCTCTG TACATTGTAA CCTTTTATTT TGAAATG	14760
ATTACACAGGA AGTTGTAAGG CTAGTACAGG GGATCC	14796

---

What is claimed:

1. A method of modulating apoptosis in a cell, comprising the step of administering to the cell an agent that modulates the amount or activity of Survivin in the cell.

2. The method of claim 1, wherein the level of apoptosis is increased by decreasing the amount or activity of Survivin in the cell.

3. The method of claim 1, wherein the level of apoptosis is decreased by increasing the amount or activity of Survivin in the cell.

4. A method of inhibiting apoptosis in a cell, comprising the step of administering to the cell a Survivin polypeptide, Survivin polypeptide fragment or an apoptosis-inhibiting peptidomimetic thereof in an amount effective to inhibit apoptosis in the cell.

5. A method of inhibiting apoptosis in a cell, comprising the step of administering to the cell a transgene encoding a Survivin polypeptide or Survivin polypeptide fragment thereof, the transgene being effective to cause expression of the Survivin polypeptide or fragment thereof in an amount effective to inhibit apoptosis in the cell.

6. A method of increasing apoptosis in a cell, comprising the step of administering to the cell an agent that decreases the apoptosis-inhibiting activity of the Survivin polypeptide present in the cell, in an amount that is effective to increase the level of apoptosis in the cell.

7. The method of claim 6, wherein said agent is a polypeptide comprising the sequence EGWEPDDDDPIEEH-KKHSSGC, its conservatively substituted homologs or small molecule peptidomimetics thereof.

8. A method of increasing the level of apoptosis in a cell, comprising the step of administering to the cell an agent that increases the transcription of the sense strand of EPR-1, in an amount effect to inhibit the translation of mRNA encoding a Survivin polypeptide, thereby decreasing the transcription of the Survivin polypeptide and increasing the level of apoptosis in the cell.

9. An isolated nucleic acid molecule that encodes the amino acid sequence depicted in **FIG. 10**, allelic variants of the amino acid sequence of **FIG. 10**, and fragments thereof that are effective to inhibit apoptosis.

10. The isolated nucleic acid molecule of claim 9, wherein said nucleic acid molecule is operably linked to one or more expression control elements.

11. The isolated nucleic acid molecule of claim 9, wherein said nucleic acid molecule is included in a vector.

12. An isolated nucleic acid molecule that encodes a member of the Survivin family of proteins, wherein said nucleic acid molecule hybridizes to a nucleic acid molecule of claim 9 under conditions of sufficient stringency to produce a clear signal.

13. A host cell transformed to contain the nucleic acid molecule of claim 9.

14. The host cell of claim 13, wherein said host is selected from the group consisting of prokaryotic hosts and eukaryotic hosts.

15. A method for producing a Survivin protein comprising the step of culturing a host transformed with the nucleic acid molecule of claim 11 under conditions in which the Survivin protein is expressed.

16. The method of claim 15, wherein said host is selected from the group consisting of prokaryotic hosts and eukaryotic hosts.

17. An isolated polypeptide comprising the amino acid sequence depicted in **FIG. 10**, allelic variants thereof and fragments thereof that retain the ability to inhibit cellular apoptosis.

18. A polypeptide comprising the sequence EGWEPD-DDPIEEHKKHSSGC, its conservatively substituted homologs and small molecule peptidomimetics thereof.

19. A monoclonal antibody that binds to the polypeptide, allelic variants thereof and fragments thereof that retain the ability to inhibit cellular apoptosis of claim 17.

20. The monoclonal antibody of claim 19 which has been humanized.

21. A method for reducing the severity of a pathological state mediated by Survivin comprising the step of reducing Survivin expression or activity.

22. The method of claim 21 wherein said pathological state is caused by abnormal cell growth.

23. The method of claim 21 wherein said Survivin expression is reduced by contacting affected cells with an RNA molecule that is complementary to a Survivin encoding mRNA molecule.

**24.** The method of claim 21, wherein said Survivin activity is reduced by blocking the intracellular interaction of Survivin with a Survivin binding partner comprising the step of contacting said Survivin with an agent that blocks the binding of Survivin to said binding partner.

**25.** The method of claim 24 wherein said agent blocks the binding of said Survivin to said binding partner by selectively binding to Survivin.

**26.** The method of claim 25 wherein said agent blocks the binding of said Survivin to said binding partner by selectively binding to the binding partner.

**27.** The method of claim 26 wherein said agent is a polypeptide fragment of Survivin.

**28.** The method of claim 26 wherein said agent comprises the polypeptide sequence EGWEPDDDDPIEEHKKHSSGC, its conservatively substituted homologs and small molecule peptidomimetics thereof.

**29.** A method for identifying agents that block the interaction of Survivin with a Survivin binding partner comprising the steps of:

- a) incubating Survivin, or a fragment thereof, or a polypeptide comprising the sequence EGWEPDDDDPIEEHKKHSSGC, and a binding partner, or a fragment thereof, with an agent to be tested, and
- b) determining whether said agent blocks the binding of Survivin to said Survivin binding partner.

**30.** A method to assay for the presence of Survivin comprising the step of determining whether a Survivin protein is expressed by a sample.

**31.** The method of claim 30, wherein said sample is selected from the group consisting of a tissue biopsy, stool, blood, urine and saliva.

**32.** The method of claim 30 further comprising the steps of:

- a) preparing an extract of the cells in said sample, and
- b) examining the proteins of said cell extract to determine the presence of a Survivin protein.

**33.** The method of claim 30 further comprising the steps of:

- a) preparing an extract of the cells in said sample, and
- b) examining the mRNA of said cell extract to determine the presence of a Survivin encoding mRNA.

**34.** The method of claim 30 wherein said method is used to determine the growth potential of a tumor cell by correlating the level of Survivin expression with control samples to indicate tumor growth potential.

**35.** A method of detecting or monitoring the stage or progression of a cancer comprising the steps of:

- obtaining a biological fluid sample from a subject;
- contacting said sample with a monoclonal antibody that recognizes and binds to Survivin; and
- determining whether the monoclonal antibody recognizes and binds to Survivin in said sample, the presence of Survivin thereby predicting the presence of cancer.

**36.** The method of claim 35, wherein the presence of Survivin predicts late stage neoplastic disease.

**37.** A method for preserving the growth of cells in culture, comprising the step of contacting the cells with an amount of Survivin that is effective to reduce apoptosis.

**38.** A kit for detecting the presence of Survivin in a sample, comprising an antibody that binds specifically to Survivin and reagents to detect the antibody-Survivin binding pair.

**39.** A therapeutic vaccine that modulates the level of Survivin-mediated apoptosis in a host.

**40.** The vaccine of claim 39, comprising a component that decreases the activity of Survivin in the host, selected from the group of components consisting of: (1) Survivin protein or fragments thereof sufficient to evoke a cellular mediated response, (2) a DNA molecule that is antisense to Survivin mRNA or portions thereof sufficient to inhibit translation of Survivin, (3) the sense DNA strand of EPR-1 acid or portions thereof sufficient to inhibit translation of Survivin.

**41.** A method of prophylactic or preventative anti-apoptotic therapy, comprising the administration of a Survivin polypeptide, Survivin polypeptide fragment, an apoptosis-inhibiting peptidomimetic thereof, a transgene encoding a Survivin polypeptide, or a transgene encoding a Survivin polypeptide fragment to a subject before the occurrence of a disease phenotype associated with cell apoptosis.

**42.** The method of claim 41, wherein the disease phenotype is a diminished T cell count in a subject infected with HIV.

**43.** The method of claim 41, wherein the disease phenotype is a degenerative disease.

**44.** The method of any of claim 6 and **20** to **22**, further comprising the administration of one or more additional agents to control cell growth.

**45.** A method to treat cancer in a patient comprising the step of increasing the level of apoptosis in one or more cells of the cancer by decreasing the amount or activity of Survivin in said one or more cancer cells.

**46.** The method of claim 45, further comprising the step of administering to the patient an agent that decreases the translation of Survivin mRNA in said one or more cancer cells, in an amount that is effective to increase the level of apoptosis in said one or more cancer cells.

**47.** The method of claim 45, further comprising the step of administering to the patient an agent that decreases the apoptosis-inhibiting activity of the Survivin polypeptide present in said one or more cancer cells, in an amount that is effective to increase the level of apoptosis in the cell.

**48.** The method of claim 47, wherein said decreased apoptosis-inhibiting activity of Survivin is effected by the administration an agent that blocks the intracellular interaction of Survivin with a Survivin binding partner.

**49.** The method of claim 48 wherein said agent blocks the binding of said Survivin to said binding partner by selectively binding to Survivin.

**50.** The method of claim 48 wherein said agent blocks the binding of said Survivin to said binding partner by selectively binding to said binding partner.

**51.** The method of claim 50 wherein said agent is a fragment of Survivin.

**52.** The method of claim 50 wherein said agent comprises the polypeptide sequence EGWEPDDDDPIEEHKKHSSGC, its conservatively substituted homologs and small molecule peptidomimetics thereof.

**53.** A method of inhibiting or reversing reperfusion injury in a subject comprising the step of administering to subject in need thereof a Survivin polypeptide, an apoptosis-inhibiting Survivin polypeptide fragment or an apoptosis-inhibiting peptidomimetic thereof.



**54.** The method of claim 53 wherein said step of administering comprises local administration at the site of injury.

**55.** A method for inhibiting or preventing tissue or organ transplant rejection, comprising a local administration of a Survivin polypeptide, an apoptosis-inhibiting Survivin polypeptide fragment, an apoptosis-inhibiting peptidomimetic thereof, a transgene encoding a Survivin polypeptide or a transgene encoding an apoptosis-inhibiting Survivin polypeptide fragment to the tissue or organ or to a site proximal to the transplant.

**56.** A method for enhancing viability of organs and tissues prior to their transplantation into a subject, comprising the

perfusion of the organs or tissues with a a Survivin polypeptide, an apoptosis-inhibiting Survivin polypeptide fragment, or an apoptosis-inhibiting peptidomimetic thereof.

**57.** The method of claim 56, wherein said transgene is delivered via a viral vector.

**58.** The method of claim 57 wherein said vector is replication defective.

**59.** The method of claim 56 wherein said transgene is delivered as a naked nucleic acid.

\* \* \* \* \*

专利名称(译)	Survivin , 一种抑制细胞凋亡的蛋白质及其调节作用		
公开(公告)号	<a href="#">US20030100525A1</a>	公开(公告)日	2003-05-29
申请号	US10/138618	申请日	2002-05-06
[标]申请(专利权)人(译)	耶鲁大学		
申请(专利权)人(译)	耶鲁大学		
当前申请(专利权)人(译)	耶鲁大学		
[标]发明人	ALTIERI DARIO C		
发明人	ALTIERI, DARIO C.		
IPC分类号	C12N15/09 A61K38/00 A61K38/17 A61K39/00 A61K39/395 A61K48/00 A61P35/00 A61P43/00 C07K14/47 C07K16/18 C12N1/15 C12N1/19 C12N1/21 C12N5/10 C12N15/12 C12P21/02 C12P21/08 C12Q1/68 G01N33/53 G01N33/68 C12N15/85		
CPC分类号	A61K38/00 G01N2510/00 A61K48/00 A61K2039/505 C07K14/4747 C07K16/3015 C07K16/3023 C07K16/303 C07K16/3046 C07K16/3061 C07K16/3069 C07K2317/73 G01N33/5008 G01N33/5011 G01N33/5017 G01N33/5044 G01N33/57407 G01N33/68 G01N33/6893 A61K39/00		
优先权	60/031435 1996-11-20 US		
其他公开文献	US6800737		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

# 摘要(译)

本发明提供抑制细胞凋亡的蛋白质的氨基酸, 本文称为存活蛋白和编码存活蛋白的核酸分子。基于本公开内容, 本发明提供分离的存活蛋白, 分离的存活蛋白编码核酸分子, 分离存活蛋白家族的其他成员的方法, 鉴定阻断存活蛋白介导的细胞凋亡抑制的药剂的方法, 使用药剂的方法阻断存活蛋白介导的抑制或存活蛋白表达以调节生物学和病理学过程, 以及测定存活蛋白活性的方法。




FIG. 1A

FIG. 1B



FIG. 1C

5' Exon	Intron	3' Exon
(2918)GCGG GTGAG	(3161)CTGTCCCTTGACAG ATGGC	
(3280)CCAT GTAAG	(5145)ITGATTTTCTAG AGAGG	
(5272)AATT GTATG	(11942)TCITTATTCCAG GCAAA	

FIG. 1D