



US 20100166769A1

(19) **United States**

(12) **Patent Application Publication**
Hsiao et al.

(10) **Pub. No.: US 2010/0166769 A1**
(43) **Pub. Date: Jul. 1, 2010**

(54) **HIGH-YIELD TRANSGENIC MAMMALIAN
EXPRESSION SYSTEM FOR GENERATING
VIRUS-LIKE PARTICLES**

(75) Inventors: **Pei-Wen Hsiao**, Taipei (TW); **Chia-
Ying Wu**, Kaohsiung City (TW);
Yi-Chun Yeh, Taipei (TW)

Correspondence Address:
**PANITCH SCHWARZE BELISARIO & NADEL
LLP**
**ONE COMMERCE SQUARE, 2005 MARKET
STREET, SUITE 2200**
PHILADELPHIA, PA 19103 (US)

(73) Assignee: **Academia Sinica**, Taipei (TW)

(21) Appl. No.: **12/704,928**

(22) Filed: **Feb. 12, 2010**

Related U.S. Application Data

(63) Continuation-in-part of application No. 11/515,843,
filed on Sep. 5, 2006.

Publication Classification

(51) Int. Cl.	
<i>A61K 39/42</i>	(2006.01)
<i>C12P 21/00</i>	(2006.01)
<i>C12N 9/26</i>	(2006.01)
<i>A61K 39/145</i>	(2006.01)
<i>G01N 33/53</i>	(2006.01)
<i>C07K 16/08</i>	(2006.01)
<i>C12N 5/07</i>	(2010.01)
<i>C12N 15/00</i>	(2006.01)
<i>A61P 31/16</i>	(2006.01)
(52) U.S. Cl.	424/159.1; 435/69.1; 435/201;
	424/209.1; 435/7.1; 530/389.4; 435/364;
	435/455

(57) **ABSTRACT**

Virus-like particles (VLPs) of mammalian-hosted viruses, such as SARS-CoV and influenza viruses, have been recombinantly produced from Vero cells. The VLPs closely emulate the exterior of authentic virus particles and are highly immunogenic. They can elicit not only humoral but also cellular immune responses in a mammal. Compositions and methods related to the VLPs are also described.

Fig. 1A





Fig. 1B

Fig. 2B




Fig. 2A





Fig. 2C

Fig. 2D

Fig. 3A




Fig. 3B







Fig. 3C

Fig. 3D

Fig. 3E



Fig. 4B




Fig. 4A

Figure 5




Figure 7A

H3N2-VLPs
Average diam. 108.2 ± 17.9 nm




Figure 7B

H3N2 Virus
Average diam. 133.5 ± 15.4 nm




Figure 7C

H5N1-VLPs
Average diam. 125.6 ± 10.5 nm




Figure 7D

H5N1 Virus
Average diam. 104.1 ± 12.4 nm






Figure 8


A SDS-PAGE




B Anti-H3& M1& M2



C SDS-PAGE



D Anti-H5& N1& M1& M2



E






Figure 9



B




100 nm


Anti-clathrin

Figure 10


A SDS-PAGE



B Anti-H5



C Anti-N1



D Anti-H3




Figure 11A




Figure 11B




Figure 11C




Figure 11D




Figure 11E




Figure 11F





Figure 12



HIGH-YIELD TRANSGENIC MAMMALIAN EXPRESSION SYSTEM FOR GENERATING VIRUS-LIKE PARTICLES

REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation-in-part patent application of U.S. patent application Ser. No. 11/515,843, filed on Sep. 5, 2006, which was published as US 20080063664 on Mar. 13, 2008, the disclosure of which is incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The present invention relates to a mammalian expression system for generating virus-like particles (VLPs), and uses of VLPs generated by the mammalian expression system.

BACKGROUND OF THE INVENTION

[0003] The spread of a newly evolved coronavirus (CoV) caused a global threat of severe acute respiratory syndrome (SARS) pandemics in 2003 (Kuiken, T. et al., 2003, *Lancet* 362: 263-270). Coronaviruses are taxonomically classified in the order Nidovirales, based on features of their genome organization and replication strategy. As with other coronaviruses, SARS-CoV has the morphology of enveloped particles with typical peripheral projections, termed "corona" or "spikes," surrounding the surface of a viral core (Ksiazek, T. G. et al., 2003, *N Engl J Med* 348: 1953-1966; Lin, Y. et al., 2004, *Antivir Ther* 9: 287-289). Outside the coronavirus particle core is a layer of lipid envelope containing mainly three membrane proteins, the most abundant M (membrane) protein, the small E (envelope) protein, and the S (spike) protein. The homo-trimers of the S protein collectively form the aforementioned corona, which is involved in viral binding to host receptors, membrane fusion for viral entry, cell-to-cell spread and tissue tropism of coronaviruses. The viral core inside the envelope, termed nucleocapsid, harbors a positive-strand viral genome RNA of approximately 30 kb packaged by the N (nucleocapsid) protein.

[0004] Unlike other human coronaviruses, such as HCoV-229E and HCoV-OC43, that can cause only symptoms like the common cold, SARS-CoV causes a highly transmittable, severe and virulent disease that can often be lethal in adults and especially the elderly. Research and clinical interest on SARS-CoV has grown rapidly owing to the high infectivity and mortality. There is especially an urgent need for an effective and safe vaccine against SARS-CoV to deal with possible future reemergence of the SARS epidemics.

[0005] Most antiviral vaccines currently in use contain whole viruses, either inactivated or live-attenuated. Inactivated, or killed, viruses are treated chemically or by irradiation to disable their replication and are generally safe and easy to make. While eliciting neutralizing antibodies, they are unlikely to deliver viral antigens to cytosol for cytotoxic CD8+ T lymphocytes (CTLs) activation, which is critical to defend animals from infection. Live-attenuated vaccines are significantly more potent than killed vaccines. However, live-attenuated viruses pose the risk of reversion or recombination with circulating wild type into a virulent strain. Moreover, the manufacture of vaccines based on whole viruses also carries the risk of viral escape.

[0006] To avoid the danger of using the whole virus (such as killed or attenuated viruses) as a vaccine, recombinant viral

proteins have been pursued not only as research tools but also as potential advanced subunit vaccines. However, subunit vaccines are known to suffer often from poor immunogenicity, owing to incorrect folding, poor antigen presentation, or difference in carbohydrate and lipid composition. Virus-like particles (VLPs) are self-assembled microscopic antigenic structures that resemble a virus in size and shape but lack genetic materials. VLPs can concurrently present viral proteins, carbohydrates and lipids in a similar and authentic conformation and thus have been viewed as an ideal vaccine against viruses (McGuigan, L. C. et al., 1993, *Vaccine* 11: 675-678). VLPs display intact viral antigens on the surface in a repeated arrangement, with which they afford the natural binding of a large viral entity to membrane receptors of antigen-presenting cells (APCs), such as dendritic cells (DCs). DC-targeted uptake of VLPs enables potent stimulation of CD4+ T cells against VLP-associated antigens. Besides stimulating humoral immunity, VLPs are permissive for cross-presentation in DCs that allows priming of CTL response with VLP-associated antigens (Moron, G. et al., 2002, *J Exp Med* 195: 1233-45).

[0007] VLPs for over thirty different viruses have been generated in insect and mammalian systems for vaccine purpose (Noad, R. and Roy, P., 2003, *Trends Microbiol* 11: 438-44). It has been shown that cellular expression of the M protein accompanied by the E protein of coronaviruses was a minimal requirement and sufficient for the assembly of VLPs (Vennema, H. et al., 1996, *EMBO J.* 15: 2020-2028). While being dispensable in forming VLPs, the S protein can be integrated into the VLPs whenever available (Godeke, G. J. et al., 2000, *J Virol* 74: 1566-1571).

[0008] Researchers have used baculovirus expression systems to produce SARS VLPs (Ho, Y. et al., 2004, *Biochem Biophys Res Commun* 318: 833-838; Mortola, E. and Roy, P., 2004, *FEBS Lett* 576: 174-178). However, due to the intrinsic differences between insect cells and mammalian cells, the VLPs assembled in the insect (SF9) cells exhibited a size of 110 nm in diameter, which is much larger than the 78 nm of the authentic SARS-CoV virions (Lin, Y. et al., 2004, supra, and Ho, Y. et al., 2004, supra). Moreover, immunogenicity of the insect cell-based SARS-VLP remains uninvestigated. Other researchers also tried to use mammalian expression systems to produce SARS VLPs (Huang, Y. et al., 2004, *J Virol* 78: 12557-65). However, the extracellular release of VLPs is not efficient, and the yield of VLPs is not satisfying.

[0009] Therefore, there is still a need for an efficient method for the large-scale production of SARS VLPs in order to provide an effective and safe vaccine against SARS.

[0010] Influenza infection is a major threat to human health and results in significant morbidity and mortality worldwide. According to World Health Organization estimates, seasonal influenza epidemics influence 5-15% of the global populations annually and are responsible for more than 3-5 million hospitalizations and about 250,000 to 500,000 deaths per year (www.who.int/mediacentre/factsheets/fs211/en/index.html). Recently, in addition to the yearly circulating seasonal influenza variants caused by antigenic drift, other influenza virus strains with pandemic potential such as the highly pathogenic avian H5N1 or emerged novel A/H1N1 pose greater threats than in the past (www.who.int/csr/disease/avian_influenza/country/en/, and www.who.int/csr/don/2009_08_19/en/index.html) since they have become better adapted to humans by reassortment. The most efficient way of reducing the transmission of and the subsequent huge economic loss caused by

seasonal or pandemic outbreaks of influenza is preventive vaccination. The manufacture of the current licensed influenza vaccines, either in the form of a split subvirion (disrupted, highly purified virus) or a subunit vaccine (purified hemagglutinin, HA, and neuraminidase, NA), is absolutely dependent on fertilized chicken eggs as a production bioreactor. This method has substantial limitations since the manufacturing capacity is restricted by the availability of eggs, which may be insufficient to meet the urgent requirements for vaccine during a pandemic [1,2,3]. In addition, these vaccines induce antibodies primarily to the viral HA and are efficacious in healthy adults, but display lower protective rates in high-risk groups (e.g., the elderly) and may be poorly immunogenic in young children. These problems are compounded once the wild population of virus undergoes significant antigenic drift in the HA component [1,2,4,5,6]. Consequently, the protective immunity elicited by inactivated vaccines is of too short a duration to protect from newly developed influenza variants. Therefore, the development of vaccines with cross-protective efficacy to allow a rapid response to influenza evolution and/or to prolong the efficacy of vaccination needs to be addressed.

[0011] Alternatively, an improvement in the preparation of seasonal influenza vaccines licensed in Europe uses reverse genetics in mammalian cell-based culture systems rather than in eggs [2]. Using mammalian cell culture systems such as Vero or MDCK cells as adaptive hosts for vaccine viruses has several advantages, not only increasing the flexibility and consistency of the manufacture process but also recovering the host-dependent specific glycosylation of viral antigens which may not be glycosylated properly in egg- or baculovirus-dependent systems. In eukaryotic cells, protein glycosylation is involved in correct folding or directing the cellular localization of newly translated proteins and plays important roles in protein function. Different glycosylation patterns underlie some of the differences between various strains of the influenza virus.

[0012] Recently, the use of noninfectious virus-like particles (VLPs) that self-assemble by spontaneous interactions of viral structural proteins has been considered to offer good potential for advanced vaccines for a wide range of viruses that cause disease in humans [7]. The VLP-based vaccine approach is an attractive alternative to replace or complement the conventional inactivated virus vaccines or subunit vaccines with improved safety and efficacy, especially for children and the elderly. It is worth noting that a VLP-based human papillomavirus (HPV) vaccine produced in yeast system which is capable of inducing protective immune response against the HPV responsible for cervical cancer was approved for the market in 2006 [8, 27, 28]. Influenza VLPs expressed by recombinant baculovirus systems that present multi-component antigens, including HA and matrix 1 (M1), with or without NA, and that are capable of inducing cognate or innate immune responses against homologous or heterologous strains of influenza virus, have been described [3,9,10, 11,12,13,14, 29]. Clinical studies for baculovirus-expressed influenza VLPs are currently being undertaken.

[0013] In light of the great threats posed by seasonal and pandemic influenza infection, there is a need for further improved means for the development of flexible, effective, and safe vaccine for influenza infection.

BRIEF SUMMARY OF THE INVENTION

[0014] The present invention provides an efficient method for generating VLPs, wherein the resulting VLPs are highly

immunogenic and can serve as a useful vaccine; particularly SARS VLPs for use as a vaccine against SARS, and influenza VLPs for use as a vaccine or to stimulate immune response against influenza infection.

[0015] A flexible platform based on the production of influenza VLPs from recombinant Vero cells presents a practical new approach to safe and effective vaccine production, which does not have the drawbacks of the egg-based or the baculovirus culture-based methodology, and is an alternative to the conventional reverse genetics approaches used in influenza vaccine manufacture.

[0016] In some embodiments of the present invention, there is provided a method for generating virus-like particles (VLPs) of a mammalian-hosted virus, the method comprising:

[0017] constructing a plasmid comprising a nucleotide sequence encoding a combination of at least two structural proteins of the virus;

[0018] transfecting Vero cells with the plasmid; and

[0019] expressing the viral structural proteins in the transfected cells to generate VLPs of the virus.

[0020] In other embodiments of the present invention, there is provided a method for generating antibodies against SARS-CoV, comprising immunizing a mammal or bird with SARS-VLPs generated according to the present invention, and harvesting antibodies against the VLPs from the blood of the mammal or bird.

[0021] In further embodiments of the present invention, there is provided a method for detecting an infection of SARS-CoV in a subject, comprising contacting a serum sample from the subject with SARS-VLPs generated according to the present invention, and determining the presence in the sample of an antibody/antigen complex, whereby the presence of the complex indicates a positive result.

[0022] In further embodiments of the present invention, there is provided a method for detecting an infection of SARS-CoV in a subject, comprising contacting a tissue sample from the subject with antibodies against the SARS-VLPs generated according to the present invention, and determining the presence in the sample of an antibody/antigen complex, whereby the presence of the complex indicates a positive result.

[0023] In still other embodiments of the present invention, there is provided a method for preventing an infection of SARS-CoV in a subject, comprising immunizing the subject with SARS-VLPs generated according to the present invention.

[0024] In still other embodiments of the present invention, there is provided an immunogenic composition comprising SARS-VLPs generated according to the present invention.

[0025] A further aspect of the present invention relates to a method of preparing an influenza virus-like particle (VLP), the method comprising:

[0026] obtaining a founder Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2;

[0027] constructing at least one recombinant DNA molecule comprising a sequence encoding an influenza hemagglutinin (HA) and a sequence encoding an influenza neuraminidase (NA);

[0028] introducing the at least one recombinant DNA molecule into the founder Vero cell to obtain a co-expression Vero cell stably transfected with the sequences encoding the influenza M1 and the influenza M2, and further transfected with

the sequences encoding the influenza HA and the influenza NA, wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems;

[0029] culturing the co-expression Vero cell under conditions to allow expressions of the influenza M1, the influenza M2, the influenza HA and the influenza NA, and assembly of the VLP comprising the influenza M1, the influenza M2, the influenza HA and the influenza NA; and

[0030] isolating the VLP from the culture supernatant of the co-expression Vero cell.

[0031] Another aspect of the present invention relates to a method of preparing an influenza virus-like particle (VLP), the method comprising:

[0032] obtaining a co-expression Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2, and further transfected with a sequence encoding an influenza hemagglutinin (HA) and a sequence encoding an influenza neuraminidase (NA), wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems; and

[0033] culturing the co-expression Vero cell under conditions to allow expressions of the influenza M1, the influenza M2, the influenza HA and the influenza NA, and assembly of the VLP comprising the influenza M1, the influenza M2, the influenza HA and the influenza NA; and

[0034] isolating the VLP from the culture supernatant of the co-expression Vero cell.

[0035] Another aspect of the present invention relates to an influenza virus-like particle (VLP), comprising:

[0036] an influenza M1, an influenza M2, an influenza hemagglutinin (HA) and an influenza neuraminidase (NA), wherein the influenza proteins are recombinantly expressed from a Vero cell; and

[0037] at least one cellular protein of the Vero cell.

[0038] Another aspect of the invention relates to a founder Vero cell that is a Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2, wherein the expressions of the influenza M1 and M2 proteins in the founder Vero cell are controlled by an inducible expression system. In one embodiment, the founder Vero cell is recombinantly made from Vero E6 cell.

[0039] Another aspect of the invention relates to a method of obtaining a founder Vero cell. The method comprises:

[0040] introducing into a Vero cell a sequence encoding an influenza M1 and a sequence encoding an influenza M2; and

[0041] obtaining the founder Vero cell stably transfected with the sequence encoding the influenza M1 and the sequence encoding the influenza M2,

[0042] wherein the expressions of the influenza M1 and M2 proteins in the founder Vero cell are controlled by an inducible expression system.

[0043] An embodiment of the invention relates to a co-expression Vero cell that is a Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2, and further transfected with a sequence encoding an influenza HA and a sequence encoding an influenza NA, wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems. In one embodiment, the co-expression Vero cell is recombinantly made from Vero E6 cell.

[0044] Another embodiment of the invention relates to a method of obtaining a co-expression Vero cell. The method comprises:

[0045] obtaining a founder Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2;

[0046] introducing into the founder Vero cell a sequence encoding an influenza HA and a sequence encoding an influenza NA; and

[0047] obtaining a co-expression Vero cell stably transfected with the sequence encoding the influenza M1 and the sequence encoding the influenza M2, and further transfected with the sequence encoding the influenza HA and the sequence encoding the influenza NA,

[0048] wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems.

[0049] Embodiments of the invention further relate to immunogenic compositions comprising the influenza VLPs according to embodiments of the present invention, and antibodies against the influenza VLPs according to embodiments of the present invention.

[0050] Methods related to the influenza VLPs, the immunogenic compositions and antibodies are also included in the present invention.

[0051] Other aspects, features and advantages of the invention will be apparent from the following disclosure, including the detailed description of the invention and its preferred embodiments and the appended claims.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

[0052] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawings will be provided by the Office upon request and payment of the necessary fee.

[0053] The foregoing summary, as well as the following detailed description of the invention, will be better understood when read in conjunction with the appended drawings. For the purpose of illustrating the invention, there are shown in the drawings embodiments which are presently preferred. It should be understood, however, that the invention is not limited to the precise arrangements and instrumentalities shown.

[0054] In the drawings:

[0055] FIG. 1 comprises FIGS. 1A and 1B. FIG. 1A comprises an illustration of the construction of the fluorescent SARS VLP-expressing plasmid. FIG. 1B comprises fluorescent images showing the locations of the expressed VLPs. Regarding FIG. 1A, two tet operator-regulated, CMV promoter-driven expression cassettes were constructed into the same plasmid for inducible expression of M-GFP fusion protein (i.e., the M protein fused with a green fluorescent protein (GFP)) and E protein from one cassette, and S protein from the other. FIG. 1B shows the results of the expression and assembly of fluorescent SARS VLPs in the VeroE6/S-MG-E-55 producer cell line, wherein cells were induced by adding 1 µg/ml doxycycline (Dox) to culture medium for 1 day, fixed, and then stained indirectly with antibodies specifically against M, GFP, S and E proteins as marked. The green fluorescence from GFP in the stained cells was scanned and merged for co-localization with different proteins contained in the VLP inside the producer cells.

[0056] FIG. 2 comprises FIGS. 2A-2D, and shows the results of the purification and characterization of Vero E6-secreted SARS-VLPs. Regarding FIG. 2A, secreted VLPs were purified by sucrose gradient ultra-centrifugation. Protein concentration (measured by Bradford Assay) and GFP fluorescence level in each fraction were plotted as marked. Regarding FIG. 2B, proteins contained in each fraction were analyzed by SDS-PAGE and Coomassie blue staining. Regarding FIG. 2C, identities of the protein bands marked in FIG. 2B were verified by western blot analysis using antibodies against S, M, E, or GFP proteins. FIG. 2D is an electron microscopic image of negatively stained SARS-VLPs (fractions 9 to 15 of FIG. 2B) purified by sucrose gradient from cell culture medium (the bar indicates a scale of 50 nm).

[0057] FIG. 3 comprises FIGS. 3A-3E and shows the results of immunization with SARS-VLPs induced humoral immune responses in mice. Regarding FIG. 3A, a diagram of immunization protocol, groups of four mice were subcutaneously injected with different dosage of SARS-VLPs at two time points as marked. Serum samples were examined for VLP-specific antibody responses in tested mice by ELISA after serial dilution. FIG. 3B shows graphs relating to ELISA titers of VLP-specific IgG, IgG1, and IgG2a using SARS-VLP as the capture antigen. Serum samples were collected on the 28th day after primary immunization. Dilution of test samples is marked on the X-axis. The background-subtracted absorbance (450 nm) was plotted as means \pm standard deviations (error bar). Presented data summarize the results of three different experiments. FIG. 3C is a graph that relates to cross-reaction of VLP-specific IgG antibodies with real SARS-CoV. Anti-sera as shown in FIG. 3B were diluted (1:250) in PBS. The SARS-specific antibody titer elicited by SARS-VLP vaccination was detected by a commercial SARS ELISA test kit (Euroimmun) according to the manufacturer's protocol, except for a modification by replacing the anti-human IgG secondary antibody with anti-mouse IgG. Mean titer and standard deviation in each group of immunized mice was summarized and plotted as means \pm standard deviations. FIG. 3D is a graph relating to a time course of VLP-elicited antibody responses. Serum samples were collected from immunized mice at the indicated time points. Anti-sera were diluted (1:250) in PBS and titers of VLP-specific IgG were measured by ELISA analysis as in FIG. 3B. FIG. 3E relates to antigen determinants of VLP-elicited antibodies. Three doses (100, 10, 1 ng) of purified VLP were loaded as Western blot antigens. Anti-sera as shown in FIG. 3B were diluted (1:1000) in PBS and subjected to western blot analysis.

[0058] FIG. 4 comprises FIGS. 4A and 4B and relates to immunization with SARS-VLPs induced cellular immune responses in mice. Primary culture of splenocytes obtained from tested mice 28 days after priming as shown in FIG. 3B were re-stimulated with SARS-VLP for 40 hours. Responsive cells that secrete Interferon- γ (FIG. 4A) and interleukin-4 (FIG. 4B) were determined by ELISPOT assays. Presented data summarize the results of three different experiments as means \pm standard deviations (error bar).

[0059] FIG. 5 illustrates the construction and cellular expression of influenza virus-like particles (VLPs) in Vero cells:

[0060] FIG. 5A are diagrams of mammalian expression vectors of influenza VLPs, which indicate the arrangement of various genetic components including the CMV constitutive promoter (CMV), tetracycline repressor gene (TetR), polyadenylation signals (PA), CMV doxycycline (Dox)-inducible

promoter (CMV-TO), chimeric intron (In), encephalomyocarditis virus internal ribosomal entry site (IRES), and coding sequences of influenza proteins (M1 and M2, matrix 1 and 2; HA, hemagglutinin; NA, neuraminidase);

[0061] FIGS. 5B and 5C are pictures showing expression of viral genes in selected quadruple VLP producer cells, where expressions of M1, M2, H3, H5, and N1 were detected by Western blot assays: the total cell lysates were extracted from VLP producing Vero cells without (-) or with (+) Dox induction, proteins in the cell lysates were separated by SDS-PAGE, then probed with respective specific antibodies against HA, NA, M1 and M2 in Western blot assays, molecular masses of expressed HA, NA, M1 and M2 are indicated on the right; expression of N2 (shown in FIG. 5B) without (-) or with (+) Dox induction was detected by RT-PCR; and

[0062] FIGS. 5D and 5E show *in vivo* immunofluorescence staining of expressed viral proteins in VLP producing Vero cells with antibodies against HA and NA (red) as marked and counterstained with DAPI (blue).

[0063] FIG. 6 shows the morphology and antigen presence determinations of purified influenza VLPs produced by Vero cells:

[0064] FIGS. 6A and 6B are the pictures of purified secreted influenza VLPs under TEM: the secreted VLPs were purified by sucrose density gradient ultracentrifugation, negatively stained with 2% uranyl acetate, and observed by TEM at 100,000 \times magnification; pictures of the respective virus strains are also shown below the VLPs; and

[0065] FIGS. 6C and 6D are pictures of purified secreted influenza VLPs from immunogold electron microscopy analysis: the primary antibodies used are labeled under individual panels; secondary antibody was goat anti-rabbit conjugated to 12 nm gold beads, bars represent 100 nm.

[0066] FIGS. 7A-D illustrate the dynamic light scattering (DLS) measurements of average particle sizes and distributions of H3N2 VLPs, H3N2 virus, H5N1 VLPs and H5N1 virus in solution, respectively: the average particle diameters of H3N2 VLPs and H5N1 VLPs in phosphate buffer (pH 7.4), at 25 $^{\circ}$ C. were 108.2 \pm 17.9 nm (A) and 125.6 \pm 10.5 nm (C), respectively; the average particle diameters of H3N2 virus and H5N1 virus in phosphate buffer (pH 7.4), at 25 $^{\circ}$ C. were 133.5 \pm 15.4 nm (B) and 104.1 \pm 12.4 nm (D), respectively; two representative determinations of different batches of VLPs or virus are shown as red and green lines; the size distribution of VLP populations ranged from 70 to 200 nm with a 95% confidence interval (CI).

[0067] FIG. 8 shows the characterization of purified influenza VLPs produced by Vero cells:

[0068] FIGS. 8A and 8C show pictures of SDS-PAGE analysis of H3N2 VLPs and H5N1 VLPs as compared with H3N2 virus and H5N1 virus, respectively: the total proteins of the VLPs or virus were resolved by SDS-PAGE in a 7.5-17.5% gradient gel and stained with Coomassie blue; each relevant band on the gel, as marked with an arrow and a number, was subjected to LC/MS/MS analysis to identify its composition; the identified viral protein bands of the VLPs are marked with parentheses in panels; and molecular masses of protein markers are labeled on the right;

[0069] FIGS. 8B and 8D show pictures of Western blot analyses of viral proteins in H3N2 VLPs and H5N1 VLPs as compared with H3N2 virus and H5N1 virus, respectively: after the SDS-PAGE described in FIGS. 8A and 8C, Western blot was performed with specific antibodies as labeled: the relative abundances of HA and NA, quantified by Chemige-

nus 2 (SYNGENE, Frederick, Md.) and GeneTools (version 3.07) software, were 4:1 for H3N2 VLPs and 3:2 for H5N1-VLPs; the HA protein attributes to 12.8% or 18% of total proteins in H3N2 VLPs or H5N1 VLPs, respectively; and

[0070] FIG. 8E demonstrates the assessment of HA function by hemagglutination assay: the amounts of VLPs or virus used are indicated, in a twofold serial dilution; THE (buffer of VLPs) and PBS were included as the negative controls in the hemagglutination assays.

[0071] FIG. 9 shows specific integration of Vero cellular proteins in H5N1 VLPs:

[0072] FIG. 9A shows pictures of Western blot analyses of viral proteins (left) and Vero cell proteins (right) in H5N1 VLPs: purified VLPs were either mock-treated (Untr., lane 1) or digested overnight with trypsin (Prot., lane 2) followed by purification with a 20% sucrose cushion; 2.5 µg each of the purified VLPs were further analyzed by SDS-PAGE followed by Western blotting with antibodies against the indicated proteins including HA, NA, M1, M2, β-actin, tubulin, annexin A2, and clathrin; molecular weight markers are marked on the left; and

[0073] FIG. 9B shows immunogold labeling of clathrin on the surface of purified H5N1 VLPs: secreted VLPs purified from conditioned medium were immunogold labeled with antibody against clathrin, negatively stained with 2% uranyl acetate, and observed by electron microscopy (100,000× magnification); bar represents 100 nm.

[0074] FIG. 10 shows glycosylation profiling of HA and NA in influenza VLPs produced by Vero cells: mock-treated and deglycosylase-treated proteins of H5N1-VLPs were separated by SDS-PAGE, stained with Coomassie blue (A), and further analyzed by Western blotting using antibodies against H5 (B) and N1 (C); proteins of H5N1 virus were also analyzed by Western blotting using antibodies against H5 (B); mock-treated and deglycosylase-treated proteins of H3N2-VLPs or H3N2 virus were also analyzed by Western blotting using the antibody against H3 (D); the molecular markers are labeled on the left; the positions of PNGase F and Endo-H are indicated; the glycosylated HA1, HA2, NA, and H3 are labeled as HA1**, HA2**, NA** and, H3**, respectively; # represents glycosylated proteins harboring a residue moiety of complex-type glycans sensitive to PNGase F, but not Endo-H; π represents unknown posttranslational modifications on the NA of H5N1-VLPs.

[0075] FIG. 11 illustrates humoral immune response of influenza VLPs made from Vero cells:

[0076] FIG. 11A shows the regimen of prime and boost vaccination followed by viral challenge: IM0, IM1, and IM2 represented the mouse serum collected at pre-immune, and 14 days after priming, and boosting, respectively;

[0077] FIG. 11B shows antigen-specific IgG antibodies from the serum of each mouse group taken at IM2 assayed against distinct antigens of H3N2-VLP, H5 protein (recombinant baculovirus expressed), and H5N1-VLP (same as the immunization antigen) by ELISA after vaccination of H5N1-VLP or inactivated whole virus (IWV): groups of mice (n=8-12) were either intramuscularly immunized with 2.5 µg and 10 µg dose of VLPs, or IWV, as marked; the dilution of used serum samples in the ELISA assays are labeled at the top; H3N2-VLPs contain all the host cell proteins integrated in the VLP antigen except with different subtype of HA and NA as negative control, whereas the baculovirus-produced H5 protein was positive control; the used amounts of coating antigens were equivalent as labeled;

[0078] FIG. 11C shows results of ELISA of serum IgG antibodies induced by H5N1-VLPs or IWV vaccines against H5N1 virus as ELISA antigen;

[0079] FIG. 11D shows results of Western blot analysis of mice serum IgG antibodies elicited by either H5N1-VLP (left panel) or IWV (right panel) vaccines: the used antigens and individual amounts are labeled at the top of panels;

[0080] FIG. 11E shows the specific IgG isotype and IgA elicited by VLPs and IWV vaccines assayed using H5N1-VLP as ELISA antigen; and

[0081] FIG. 11F shows HI titer of each vaccinated mouse and plot of the mean values of the same group: HI titer of 40 was set as threshold of seroprotection.

[0082] FIGS. 12A-D show vaccine protection against lethal-dose challenge of H5N1 virus: at day 42, vaccinated mice were challenged intranasally with a lethal dose (100 LD₅₀) of recombinant H5N1 (NIBRG-14) virus and monitored daily for weight loss and mortality; the percentages of survival rate and changes of body weight and temperature were recorded: (A) survival for H5N1-VLP groups; (B) survival for IWV groups, mice that lost greater than 30% body weight were euthanized; (C) body weight and temperature for H5N1-VLP groups; for the groups receiving 0.3 µg and 1.5 µg antigen dose, only data of surviving mice are shown; and (D) body weight and temperature for IWV groups.

DETAILED DESCRIPTION OF THE INVENTION

[0083] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention pertains. Otherwise, certain terms used herein have the meanings as set in the specification. All patents, published patent applications and publications cited herein are incorporated by reference as if set forth fully herein. It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise.

[0084] To generate VLPs as a SARS vaccine, technical challenges include mammalian post-translational modifications and correct folding of viral proteins, their delicate organization into a lipid envelope, and sustainable yield for practical usage. The SARS-S protein is deduced as a huge glycoprotein containing 1255 aa residues with 23 putative N-linked glycosylation sites, and at least 12 N-glycans have been identified (Krokhin, O. et al., 2003, Mol Cell Proteomics 2: 346-56). In SARS-CoV infected cells and purified virion, protein M contains one high-mannose type N-glycan (Voss, D. et al., 2006, FEBS Lett 580: 968-73). Thus, mammalian expression and cell culture-based approaches are of interest to the inventors to attain massive production of SARS-VLPs.

[0085] In one aspect, the present invention provides a method for generating virus-like particles (VLPs) of a mammalian-hosted virus, such as SARS-CoV, the method comprising:

[0086] constructing a plasmid comprising a nucleotide sequence encoding a combination of at least two structural proteins of the virus;

[0087] transfecting Vero cells with the plasmid; and

[0088] expressing the viral structural proteins in the transfected cells to generate VLPs of the virus.

[0089] The method of the present invention is suitable for generating various mammalian-hosted viruses, including but not limited to arenaviruses, coronaviruses, hepadnaviruses, herpes viruses, orthomyxoviruses, paramyxoviruses,

papovaviruses, parvoviruses, and retroviruses. In a preferred embodiment of the present invention, the mammalian-hosted virus is a coronavirus. More preferably, the mammalian-hosted virus is SARS-CoV.

[0090] The term “viral structural protein” or “structural protein of a virus” and equivalent terms as used herein refers to viral genome-encoded proteins that form the structure of a virus, including membrane glycoproteins and capsid proteins. The genome of a virus also encodes non-structural regulatory proteins involved in virus replication. For example, the structural proteins of a coronavirus comprise the M (membrane), E (envelope), S (spike) and N (nucleocapsid) proteins.

[0091] In an embodiment of the method used to generate SARS-VLPs according to the invention, the structural proteins to be expressed in transfected cells can be any combinations derived with the E, M, N and S proteins of SARS-CoV, such as, for example, M+E, M+E+S, M+S, N+M+E, N+M+E+S, and N+M+S. In a preferred embodiment, the combination of the structural proteins is M+E. Most preferably, the combination of the structural proteins is M+E+S.

[0092] The plasmid used in the present invention can be any plasmid or vector suitable for expressing heterologous proteins in mammalian cells. Many commercially available mammalian expression vectors can be readily used in the present invention, for example, the pcDNA™ series by Invitrogen Corporation (Carlsbad, Calif., USA).

[0093] To construct the recombinant plasmid used in the present invention, nucleotide sequences encoding a combination of the viral structural proteins can be grouped into one or more “expression cassettes” for controlled expression. As used herein, the term “expression cassette” refers to a nucleic acid construct, generated recombinantly or synthetically, with a series of specified nucleic acid elements which permit transcription of a nucleotide sequence of interest in a host cell. The expression cassette can be incorporated into a plasmid or chromosome. Typically, the expression cassette portion of an expression vector includes, among other sequences, a nucleotide sequence to be transcribed, a promoter, and a poly-adenylation signal. In the present invention, the term “expression cassette” is used interchangeably with the term “transgene.”

[0094] For optimal expression of the viral proteins of the present invention, the expression cassette may include an inducible system that allows high-level expression upon induction. In a preferred embodiment of the present invention, a tetracycline-inducible expression system is utilized for high-level expression of the viral proteins, wherein the induction is achieved by the addition of doxycycline into the culture medium. Examples of commercially available inducible expression systems include but not limited to the T-REx™ System and GeneSwitch™ System by Invitrogen Corporation, and the BD Tet-On™ and BD Tet-Off™ Gene Expression Systems by Clontech Laboratories, Inc. (Mountain View, Calif., USA).

[0095] According to an embodiment of the present invention, the cells used in the generation of VLPs are Vero cells. The Vero cell line, i.e. the cell line of ATCC No. CCL-81™, was initiated from the kidney of a normal adult African green monkey on Mar. 27, 1962, by Y. Yasumura and Y. Kawakita at the Chiba University in Chiba, Japan. The cell line was brought to the Laboratory of Tropical Virology, National Institute of Allergy and Infectious Diseases, National Institutes of Health in the 93rd passage from Chiba University by

B. Simizu on Jun. 15, 1964. In addition to its use as a vaccine cell substrate, this cell line has been used extensively for virus replication studies and plaque assays. In the present invention, the term “Vero cell” includes not only cells from the original Vero cell line, but also those derived from Vero-derived cell lines such as Vero 76 (ATCC No. CRL-1587™) and Vero E6 (ATCC No. CRL-1586™).

[0096] Transfection can be performed by any known method and can result in either transient or stable transfection. Stable transfection is preferred to establish a cell line producing VLPs of interest. Methods for obtaining stable transfection are well known and include, for example, selection for spontaneously stable transfectants, transfection with immortalizing genes, and selection for genes providing resistance to antibiotics such as neomycin, puromycin, zeocin, hygromycin B, and blasticidin S.

[0097] As demonstrated in the following examples, SARS-VLPs generated by the method of the present invention can induce high titers of SARS-CoV-specific antibodies in mice. Therefore, the present invention also provides a method for generating antibodies against SARS-CoV, comprising immunizing a mammal or bird with SARS-VLPs generated according to the present invention, and harvesting antibodies against the VLPs from the blood of the mammal or bird.

[0098] According to the following examples, in addition to eliciting humoral immune responses, SARS-VLPs generated by the method of the present invention also stimulates systemic activation of T helper (T_H) cells. Therefore, the present invention also provides a method for preventing an infection of SARS-CoV in a subject, comprising immunizing the subject with SARS-VLPs generated according to the present invention. Preferably, the subject is a mammal, such as a dog, a cat, a rabbit, a rat, a mouse, a pig, a sheep; a goat, and a cow, and more preferably, a human.

[0099] Immunization can be performed traditionally. Suitable regimes for initial administration and booster doses are variable, but may include an initial administration followed by subsequent booster administrations. The quantity of SARS-VLPs to be administered depends on the subject to be immunized, including, for example, the capacity of the individual's immune system to synthesize antibodies, and if needed, to produce a cell-mediated immune response. Precise amounts of VLPs required to be administered depend on the judgment of the practitioner. However, suitable dosage ranges are readily determinable by one skilled in the art without undue experimentation in view of the present disclosure. The dosage may also depend on the route of administration and will vary according to the size of the host. Non-limiting exemplary dosages include, for instance, a preferred dosage of about 0.01 mg/kg to about 10 mg/kg body weight, and a more preferred dosage of about 0.1 mg/kg to about 1 mg/kg body weight.

[0100] In another aspect, the present invention provides a method for detecting an infection of SARS-CoV in a subject, comprising contacting a serum sample from the subject with SARS-VLPs generated according to the present invention, and determining the presence in the sample of an antibody/antigen complex, whereby the presence of the complex indicates a positive result.

[0101] Preferably, the method involves an immunoassay. In a particularly preferred embodiment of the present invention, the method involves an enzyme-linked immunosorbent assay (ELISA). In ELISA assays, the VLPs are immobilized onto a selected surface, for example, a surface capable of binding

proteins, such as the wells of a polystyrene microtiter plate. After washing to remove incompletely adsorbed material, a nonspecific protein, such as a solution of bovine serum albumin (BSA) that is known to be antigenically neutral with regard to the test sample may be bound to the selected surface. This allows for blocking of nonspecific adsorption sites on the immobilizing surface and thus reduces the background caused by nonspecific binding of proteins in the antisera onto the surface.

[0102] The immobilizing surface is then contacted with a sample, such as a serum sample from a subject suspected of a SARS-CoV infection, in a manner conducive to immune complex (antigen/antibody) formation. This may include diluting the sample with diluents, such as solutions of BSA, bovine gamma globulin (BGG) and/or phosphate buffered saline (PBS)/Tween. The sample is then allowed to incubate for about 2 to about 4 hours, at suitable incubation temperatures, such as of the order of about 25° C. to about 37° C. Following incubation, the sample-contacted surface is washed to remove non-immunocomplexed material. The washing procedure may include washing with a solution, such as PBS/Tween™ or a borate buffer. Following formation of specific immunocomplexes between the test sample and the bound protein, and subsequent washing, the occurrence, and even the amount, of immunocomplex formation may be determined by subjecting the immunocomplex to a second antibody having specificity for the first antibody. If the test sample is of human origin, the second antibody is an antibody having specificity for human immunoglobulins and in general IgG. To provide for the detection, the second antibody may have an associated activity such as an enzymatic activity that will generate, for example, a color development upon incubating with an appropriate chromogenic substrate. Quantification may then be achieved by measuring the degree of color generation using, for example, a spectrophotometer.

[0103] The present invention also provides another method for detecting an infection of SARS-CoV in a subject, comprising contacting a tissue sample from the subject with antibodies against the SARS-VLPs generated according to the present invention, and determining the presence in the sample of an antibody/antigen complex, whereby the presence of the complex indicates a positive result.

[0104] Preferably, the method involves an immunoassay. In a particularly preferred embodiment of the present invention, the method involves indirect immunofluorescence staining. Indirect immunofluorescence staining involves intracellular staining of specific proteins with antibodies and tracking of the signals via respective fluorescence-labeled second antibodies. For example, target cells were first fixed, permeated, and washed, and the cells were blocked with 1% gelatin/PBST for 1 hour and then reacted with the first antibody (such as anti-S, M, E and GFP) in appropriate dilution with 1% gelatin/PBST at 4° C. for overnight. Subsequent to another three washes in PBST, the cells were incubated with the fluorescence-conjugated secondary antibody, washed and scanned under a confocal microscope.

[0105] In a further aspect, the present invention provides an immunogenic composition comprising SARS-VLPs generated according to the present invention. An immunogenic composition preferably generates immunological responses, such as antibody or T-cell responses, in a subject to whom it is administered.

[0106] SARS-VLPs generated according to the present invention can be purified after being harvested from a culture

medium or cell suspension and before being used in an immunogenic composition. Any method can be used that is known to separate VLPs or viruses from surrounding proteins, lipids, nucleic acids, membranes, intact cells, and the like. Especially preferred are affinity chromatography methods; for example, an immobilized monoclonal antibody specific for SARS-VLPs can be used. Additional suitable methods are gel filtration chromatography, ion exchange chromatography, and density gradient sedimentation.

[0107] The immunogenicity of SARS-VLPs generated according to the present invention may be further improved when co-administered with adjuvants. Adjuvants enhance the immunogenicity of an antigen but are not necessarily immunogenic themselves. Adjuvants may act by retaining the antigen locally near the site of administration to produce a depot effect facilitating a slow, sustained release of antigen to cells of the immune system. Adjuvants can also attract cells of the immune system to an antigen depot and stimulate such cells to elicit immune responses.

[0108] For example, preferred adjuvants to enhance effectiveness of an immunogenic composition include, but are not limited to: (1) aluminum salts (alum), such as aluminum hydroxide, aluminum phosphate, aluminum sulfate, etc; (2) oil-in-water emulsion formulations (with or without other specific immunostimulating agents such as muramyl peptides (see below) or bacterial cell wall components), such as for example (a) MF59™, containing 5% Squalene™, 0.5% Tween™ 80, and 0.5% Span™ 85 (optionally containing various amounts of MTP-PE (see below), although not required) formulated into submicron particles using a microfluidizer such as Model 1 10Y microfluidizer (Microfluidics, Newton, Mass., U.S.A.), (b) SAF™, containing 10% Squalene™, 0.4% Tween™ 80, 5% pluronic-blocked polymer L121, and thr-MDP (see below) either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion, and (c) Ribi™ adjuvant system (RAS), (Ribi Immunochem, Hamilton, Mont., U.S.A.) containing 2% Squalene™, 0.2% Tween™ 80, and one or more bacterial cell wall components from the group consisting of monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL+CWS (Detox™); (3) saponin adjuvants, such as Stimulon™ (Cambridge Bioscience, Worcester, Mass., U.S.A.) may be used or particles generated therefrom such as ISCOMs (immunostimulating complexes); (4) Complete Freund's Adjuvant (CFA) and Incomplete Freund's Adjuvant (IFA); (5) cytokines, such as interleukins (e.g., IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12, etc.), interferons (e.g., gamma interferon), macrophage colony stimulating factor (M-CSF), tumor necrosis factor (TNF), etc; (6) detoxified mutants of a bacterial ADP-ribosylating toxin such as a cholera toxin (CT), a pertussis toxin (PT), or an *E. coli* heat-labile toxin (LT), particularly LT-K63, LT-R72, CT-Si09, PT-K9/G129; and (7) other substances that act as immunostimulating agents to enhance the effectiveness of the composition.

[0109] As mentioned above, muramyl peptides include, but are not limited to, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-s-n-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), etc.

[0110] Pharmaceutically acceptable salts can also be used in immunogenic compositions of the present invention. For example, mineral salts such as hydrochlorides, hydrobro-

mides, phosphates, or sulfates, as well as salts of organic acids such as acetates, propionates, malonates, or benzoates.

[0111] Immunogenic compositions of the present invention generally contain pharmaceutically acceptable excipients, such as water, saline, glycerol, and ethanol, and may contain substances such as wetting agents, emulsifying agents, or pH buffering agents.

[0112] Immunogenic compositions of the present invention may be prepared as indictable, as liquid solutions, suspensions or emulsions, and administered parenterally, by injection subcutaneous, intradermal or intramuscularly injection. Alternatively, the immunogenic compositions of the present invention may be formulated and delivered in a manner to evoke an immune response at mucosal surfaces. Thus, the immunogenic composition may be administered to mucosal surfaces by, for example, the nasal or oral (intra-gastric) routes. Alternatively, other modes of administration including suppositories and oral formulations may be desirable. Oral formulations can take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders.

[0113] The immunogenic composition of the present invention may further comprise antigens from other pathogens to be a multivalent immunogenic composition.

[0114] In another aspect of the present invention, the mammalian-hosted virus is an orthomyxovirus, i.e., a family of RNA viruses that includes five genera: influenza virus A (such as H1N1, H2N2, H3N2, H5N1, H7N7, H1N2, H9N2, H7N2, H7N3, H10N7), influenza virus B, influenza virus C, isavirus and thogotovirus. Preferably, the mammalian-hosted virus is an influenza virus.

[0115] The structural proteins of an influenza virus, include, but are not limited to, hemagglutinin (HA), neuraminidase (NA), matrix (M1) and proton ion-channel protein (M2).

[0116] M1 is the most abundant protein in influenza particles. It forms a layer inside the viral envelope, which is derived from portions of the host cell membranes (phospholipids and proteins), but include some viral glycoproteins. M1 is able to direct viral assembly and budding. It is reported that the expression of M1 alone induces the release of VLPs from insect cells (see US20050186621A1 and references therein).

[0117] M2 is a proton-selective ion channel protein, integral in the viral envelope. It has an important role in the life cycle of the influenza virus, enables hydrogen ions to enter the viral particle (virion) from the endosome, thus lowering the pH inside of the virus and causing dissociation of the viral matrix protein M1 from the ribonucleoprotein RNP. This is a crucial step in uncoating of the virus and exposing its content to the cytoplasm of the host cell.

[0118] HA is an antigenic glycoprotein found on the surface of the influenza viruses. It is a type of hemagglutinin, i.e., a protein has the ability to cause red blood cells (erythrocytes) to clump together (agglutinate) in vitro. HA mediates binding of the influenza virus to target cells and entry of the viral genome into the target cell, e.g., by binding to the sialic acid-containing receptors on the surface of its target cells and causing the fusion of host endosomal membrane with the viral membrane. At least 16 different influenza HA subtypes have been discovered so far, H1 to H16.

[0119] NA is another antigenic glycoprotein found on the surface of the influenza viruses. It is involved in the release of progeny virus from infected cells, e.g., by catalyze the hydrolysis of terminal sialic acid residues from the newly

formed virions and from the host cell receptors. Its activities also include assistance in the mobility of virus particles through the respiratory tract mucus and in the elution of virion progeny from the infected cell. At least 9 different influenza NA subtypes have been discovered so far, N1 to N9.

[0120] HA and NA are the sources of the major immunodominant epitopes for virus neutralization and protective immunity. They are considered the most important components for prophylactic influenza vaccines. Influenza A viruses are classified into subtypes based on antibody responses to HA and NA. Changes in HA and/or NA can significantly alter the host specificity of an influenza virus, e.g., making the virus infectious to a new host species or more infectious to an old host. When a new strain of influenza virus having a new subtype of HA and/or NA emerges, antibodies developed after vaccination against older influenza strains may not provide effective protection against the new strain. New influenza vaccines must be developed to take into account of the antigenic drift in flu viruses.

[0121] Currently, there are two types of seasonal influenza vaccines: the flu shot and the nasal-spray flu vaccine. The flu shot contains killed influenza virus that is given with a needle, usually in the arm. The flu shot is approved for use in people older than 6 months, including healthy people and people with chronic medical conditions. The nasal-spray flu vaccine contains live, weakened flu viruses that do not cause the flu (sometimes called LAIV for "live attenuated influenza vaccine" or FluMist®). LAIV (FluMist®) is approved for use in healthy people 2-49 years of age who are not pregnant.

[0122] Each seasonal influenza vaccine are formulated as a blend of three strains of influenza viruses, e.g., two strains of influenza A and one strain of influenza B. The viruses in the seasonal vaccine change each year based on international surveillance and scientists' estimations about which types and strains of viruses will circulate in a given year. About 2 weeks after vaccination, antibodies that provide protection against influenza virus infection develop in the body. However, the seasonal influenza vaccine may not be able to provide protection against all influenza in circulation among the human population for a given year. For example, the seasonal flu vaccine containing one influenza A (H3N2) virus, one regular seasonal A (H1N1) virus, and one influenza B virus does not provide protection against 2009 H1N1 flu, which has an unusual mix of swine, avian and human influenza genetic sequences. An influenza vaccine separate from the seasonal flu vaccine, such as that against 2009 H1N1, may be required to effectively prevent the outbreak or pandemic of influenza infection.

[0123] Yearly flu vaccination begins in September or as soon as vaccine is available and continues throughout the influenza season, into December, January, and beyond. This is because the timing and duration of influenza seasons vary. While influenza outbreaks can happen as early as October, most of the time influenza activity peaks in January or later.

[0124] According to embodiments of the present invention, a platform to generate influenza VLPs from Vero cells, such as Vero E6 cell, presents a practical new approach to safe and effective vaccine production. This platform does not have the drawbacks of the egg-based or the baculovirus culture-based methodology, and is an alternative to the conventional reverse genetics approaches used in influenza vaccine manufacture. The Vero cell system according to embodiments of the present invention produces influenza VLPs including not only HA and NA but also the matrix proteins M1 and M2, because

both M1 and M2 have equally critical roles in influenza virus assembly and budding processes, suggesting their similar importance for mammalian VLP budding efficiency [31,32, 33,34,35,36]. The incorporation of M1 and M2 into influenza VLPs not only increased the VLP production yield from Vero cells, but also supplemented interior viral antigens, which can provide highly conserved T-cell and B-cell epitopes to fight homologous and heterologous viruses [37,38].

[0125] The flexibility of this approach has been demonstrated by exchanging the surface antigens of HA and NA to generate VLPs mimicking two subtypes of influenza (FIGS. 6 and 8). This can shorten the lead time for adjusting the match of vaccine specificity against the circulating strains of viruses. The preclinical-scale production of influenza VLPs from recombinant Vero cells has been achieved. This Vero cell VLP system alleviates safety restrictions and bottlenecks associated with dependence on live viruses. It also allows rapid and scaleable production, independent of a reliance on egg availability for manufacturing vaccines.

[0126] Generation of influenza VLPs from a mammalian cell has been previously reported by transient coexpression of HA and NA proteins in human 293T cells [30]. However, the present invention is the first report to reveal the proteome of VLPs made from a mammalian cell with sufficient quantities of HA and NA proteins as confirmed by SDS-PAGE analysis, including their further characterization by deglycosylation and hemagglutination assays. It is now discovered, for the first time, that VLPs expressed from Vero cells include multiple proteins from Vero cells, e.g. beta-actin, tubulin, etc., which are missing from VLPs made from other expression systems.

[0127] It is also discovered, for the first time, that VLPs expressed from Vero cells induce predominantly IgG1 antibodies, which is the same as that induced by the split or subunit vaccine and is indicative of Th2 type immune response. Vaccination of animals with VLPs expressed from Vero cells elicited HA-specific IgG1 antibodies and resulted in full protection against lethal infection of homologous virus. Whereas VLPs expressed from baculovirus induce IgG2a dominant antibody, which is the same as that induced by inactivated whole virus vaccine and is indicative of Th1 dominant immune response. The Th1 type response is prone to have adverse side effects.

[0128] It is further discovered, for the first time, that HA in Vero cell expressed VLPs contains both complex type and high-mannose type of glycans. Such glycosylated HA is not possible to make from the baculovirus system. By modifying the host insect cell, the baculovirus system can only choose one kind of glycosylation.

[0129] According to an embodiment of the present invention, an influenza virus-like particle (VLP) is prepared by a method comprising:

[0130] obtaining a founder Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2;

[0131] constructing at least one recombinant DNA molecule comprising a sequence encoding an influenza hemagglutinin (HA) and a sequence encoding an influenza neuraminidase (NA);

[0132] introducing the at least one recombinant DNA molecule into the founder Vero cell to obtain a co-expression Vero cell stably transfected with the sequences encoding the influ-

enza M1 and the influenza M2, and transfected further with the sequences encoding the influenza HA and the influenza NA;

[0133] culturing the co-expression Vero cell under conditions to allow expressions of the influenza M1, the influenza M2, the influenza HA and the influenza NA, and assembly of the VLP comprising the influenza M1, the influenza M2, the influenza HA and the influenza NA; and

[0134] isolating the VLP from the culture supernatant of the co-expression Vero cell.

[0135] The founder Vero cell can be obtained by stably transfecting a Vero cell with a DNA sequence encoding an influenza M1 and a DNA sequence encoding an influenza M2. The coding sequences of the influenza M1 and M2 can be derived from any influenza virus, such as H1N1, H5N1, H3N2, etc. The sequences encoding influenza M1 and M2 can be derived from the same influenza virus, or different influenza viruses. The sequences can be the same as those naturally occurring in the viruses. The sequences can also contain one or more modifications, preferably those do not alter the functions of the encoded influenza M1 and M2. For example, the sequences can be modified based on the degeneracy of genetic codon to optimize gene expression in Vero cells, without introducing any changes in the amino acid sequences of the encoded influenza M1 and M2.

[0136] The coding sequences of an influenza M1 and M2 are cloned into one or more expression vectors so that they are operably linked to expression control elements that allow expression of the influenza M1 and M2 in Vero cells. The expression control elements comprise a promoter for Vero cell expression, translation initiation codon, transcription and translation termination sequences. The expression control elements can also contain a regulatory sequence that allows regulation of gene expression, e.g., that for inducible expression of the influenza proteins.

[0137] The expression vector is introduced into a Vero cell using methods known in the art in view of the present disclosure. Sequences encoding the influenza M1 and M2, which include the coding sequences of the influenza M1 and M2 and the operably linked expression control elements, are integrated into the genome of the Vero cell. The resulting Vero cell, i.e., the founder Vero cell, is stably transfected with the sequences encoding the influenza M1 and M2. Such stably transfected cells are selected and verified using methods known in the art in view of the present disclosure. The sequences encoding the influenza M1 and M2 can be stably transfected into the Vero cell using a single expression vector, or two separate expression vectors.

[0138] In a particular embodiment of the present invention, the founder Vero cell is stably transfected with a sequence comprising SEQ ID NO:12, which encodes the influenza M1 and M2 of influenza A/Taiwan/083/2006. In SEQ ID NO:12, a CMV/TO promoter controls the transcription of the M1 and M2 coding sequences, which are linked with an internal ribosome entry site (IRES). A sequence (SEQ ID NO:13) that constitutively expresses a tet repressor is also stably transfected into the Vero cell to regulate the expression of all genes controlled by the CMV/TO promoter. In SEQ ID NO:13, a CMV promoter controls the transcription of the tet repressor coding sequence.

[0139] Expression vectors, cell lines, and methods similar to that describe above for the construction of recombinant Vero cells for SARS-CoV VLPs can be used for the construc-

tion of the founder Vero cells stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2.

[0140] Live cultures of the founder Vero cells can be stored, e.g., by cryogenic storage at very low temperature, such as in liquid nitrogen. The cells can be conveniently retrieved and used for subsequent manipulation in a method according to embodiments of the present invention.

[0141] The coding sequences of HA and NA of one or more influenza viruses can be obtained using standard molecular biology techniques, such as RT-PCR, followed by DNA cloning and sequencing. The influenza viruses of interest can be potentially pandemic or seasonal influenza virus strains, i.e., the types and strains of flu viruses that will likely circulate among the human population for a given year based on international surveillance and scientists' estimations. The coding sequences of HA and NA can be derived from the same influenza virus, or different influenza viruses. The sequences can be the same as those naturally occurring in the viruses. The sequences can also contain one or more modifications, preferably those do not alter the antigenic specificity of the encoded influenza HA and NA. For example, the sequences can be modified based on the degeneracy of genetic codon to optimize gene expression in Vero cells, without introducing any changes in the amino acid sequences of the encoded influenza HA and NA. Or, selective residuals of the amino acid sequences can be mutated for specific purposes, such as enabling a protease cleavage site, disabling glycosylation sites or enabling fusion of designed epitopes.

[0142] The coding sequences of an influenza HA and NA are cloned into one or more expression vectors so that they are operably linked to expression control elements that allow expression of the influenza HA and NA in Vero cells. The expression control elements comprise a promoter for Vero cell expression, translation initiation codon, transcription and translation termination sequences. The expression control elements can also contain a regulatory sequence that allows regulation of gene expression, e.g., that for inducible expression of the influenza proteins.

[0143] The expression vector is introduced into the founder Vero cell using methods known in the art in view of the present disclosure. Sequences encoding the influenza HA and NA, which include the coding sequences of the influenza HA and NA and the operably linked expression control elements, can be integrated into the genome of the founder Vero cell, resulting in a co-expression Vero cell stably transfected with sequences encoding influenza proteins M1, M2, HA and NA. Sequences encoding the influenza HA and NA can also remain on the expression vector independent of the genome inside the cell, resulting in a co-expression Vero cell stably transfected with sequences encoding the influenza M1 and M2, but transiently transfected with the influenza HA and NA. Embodiments of the present invention include co-expression Vero cells that are stably or transiently transfected with sequences encoding the influenza HA and NA. The sequences encoding the influenza HA and NA can be transfected into the founder Vero cell using a single expression vector, or two separate expression vectors.

[0144] In particular embodiments of the present invention, the co-expression Vero cell is stably transfected with a sequence comprising SEQ ID NO:12, which encodes the influenza M1 and M2 of influenza A/Taiwan/083/2006, and further transfected, preferably stably transfected, with a sequence comprising SEQ ID NO:14, which encodes H3 and

N2 of influenza A/Taiwan/083/2006. SEQ ID NO:14 also includes the 5'- and 3'-expression control elements for H3 and N2 expressions.

[0145] In another particular embodiments of the present invention, the co-expression Vero cell is stably transfected with a sequence comprising SEQ ID NO:12, which encodes the influenza M1 and M2 of influenza A/Taiwan/083/2006, and further transfected, preferably stably transfected, with a sequence comprising SEQ ID NO:15, which encodes H5 and N1 of influenza A/Hanoi/30408/2005(H5N1). SEQ ID NO:15 also includes the 5'- and 3'-expression control elements for H5 and N1 expressions.

[0146] One or more DNA sequences encoding additional proteins can be further introduced into the founder Vero cell for recombinant production of influenza VLPs further comprising the additional proteins.

[0147] In one embodiment of the present invention, one or more DNA sequences encoding an adjuvant, such as a flagellin of a pathogenic bacterium, can be introduced into the founder cell for recombinant production of VLPs containing the adjuvant.

[0148] In another embodiment of the present invention, sequences encoding HAs and NAs from two or more different influenza virus strains, e.g., two or more potentially pandemic or seasonal influenza virus strains, can be transfected into the founder Vero cell. The resulting co-expression Vero cell is stably transfected with sequences encoding the influenza M1 and M2, and transiently or stably transfected with sequences encoding HAs and NAs from the two or more different influenza virus strains. The co-expression Vero cell produces VLPs comprising a blend of HAs and NAs useful for prophylactic prevention of infection by the two or more different influenza virus strains.

[0149] Expression vectors and methods similar to that described above for the construction of recombinant Vero cells for SARS-CoV VLPs can be used for the construction of the co-expression Vero cells for influenza VLPs.

[0150] In an embodiment of the present invention, expressions of the sequences encoding the viral protein, e.g., influenza M1, M2, HA and NA, are under the control of one or more inducible gene expression systems, so that viral proteins are produced and assembled into influenza VLPs under inducible conditions. Any inducible gene expression system, such as those described above for inducible production of SARS-CoV VLPs, can be used for inducible production of influenza VLPs in view of the present disclosure. The viral proteins can be controlled by the same inducible gene expression system. Each of the viral proteins can also be controlled independently by different inducible gene expression systems.

[0151] Live cultures of the co-expression Vero cells can be stored, e.g., by cryogenic storage at very low temperature, such as in liquid nitrogen. The cells can be conveniently retrieved and used for subsequent production of influenza VLP.

[0152] Accordingly, an embodiment of the present invention relates to a method of preparing an influenza virus-like particle (VLP), the method comprising:

[0153] obtaining a co-expression Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2, and further transfected with a sequence encoding an influenza hemagglutinin (HA) and a sequence encoding an influenza neuraminidase (NA), wherein the expressions of the influenza M1, M2, HA and NA

proteins in the co-expression Vero cell are controlled by one or more inducible expression systems; and

[0154] culturing the co-expression Vero cell under conditions to allow expressions of the influenza M1, the influenza M2, the influenza HA and the influenza NA, and assembly of the VLP comprising the influenza M1, the influenza M2, the influenza HA and the influenza NA; and

[0155] isolating the VLP from the culture supernatant of the co-expression Vero cell.

[0156] Embodiments of the present invention relate to a founder Vero cell that is a Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2, wherein the expressions of the influenza M1 and M2 proteins in the founder Vero cell are controlled by an inducible expression system.

[0157] Methods of obtaining a founder Vero cell are also encompassed by embodiments of the present invention. The method comprises:

[0158] introducing into a Vero cell a sequence encoding an influenza M1 and a sequence encoding an influenza M2; and

[0159] obtaining the founder Vero cell stably transfected with the sequence encoding the influenza M1 and the sequence encoding the influenza M2,

[0160] wherein the expressions of the influenza M1 and M2 proteins in the founder Vero cell are controlled by an inducible expression system.

[0161] The sequence encoding the influenza M1 and the sequence encoding the influenza M2 can be introduced into the Vero cell on a single nucleic acid molecule or on two separate nucleic acid molecules.

[0162] In an embodiment of the present invention, the founder Vero cell is a Vero E6 cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2, wherein the expressions of the influenza M1 and M2 proteins in the founder Vero E6 cell are controlled by an inducible expression system.

[0163] The founder Vero cell can be used as a host cell for construction of recombinant Vero cells comprising one or more transfected sequences encoding one or more influenza proteins that are different from the influenza M1 and M2.

[0164] Embodiments of the present invention relate to a co-expression Vero cell that is a Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2, and further transfected with a sequence encoding an influenza HA and a sequence encoding an influenza NA, wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems. In one embodiment, the co-expression Vero cell is recombinantly made from Vero E6 cell.

[0165] Method of obtaining a co-expression Vero cell are also encompassed by embodiments of the present invention. The method comprises:

[0166] obtaining a founder Vero cell stably transfected with a sequence encoding the influenza M1 and a sequence encoding the influenza M2;

[0167] introducing into the founder Vero cell a sequence encoding an influenza HA and a sequence encoding an influenza NA; and

[0168] obtaining a co-expression Vero cell stably transfected with the sequence encoding the influenza M1 and the sequence encoding the influenza M2, and further transfected with the sequence encoding the influenza HA and the sequence encoding the influenza NA,

[0169] wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems.

[0170] The sequence encoding the influenza HA and the sequence encoding the influenza NA can be introduced into the Vero cell on a single nucleic acid molecule or on two separate nucleic acid molecules. The co-expression cell can be stably or transiently transfected with the sequence encoding the influenza HA and the sequence encoding the influenza NA.

[0171] The expression of each of the influenza M1, M2, HA and NA proteins can be independently controlled by the same or different inducible expression system. In one embodiment, the expression of the influenza M1, M2, HA and NA proteins are controlled by the same inducible expression system.

[0172] Influenza proteins, e.g., M1, M2, HA and NA, are expressed in the co-expression Vero cell under suitable conditions, e.g., when the cell is grown in a medium containing inducer for the inducible expression system. These viral proteins, together with some cellular proteins from Vero cell, self-assemble into noninfectious VLPs with the antigenic HA and NA presented on the surface of the VLPs. The VLPs are secreted into the culture medium of the co-expression Vero cell. They are subsequently separated from the cells and isolated using methods that preserve the integrity of the VLPs, such as by density gradient centrifugation, and the like.

[0173] Methods similar to that described above for the production and isolation of SARS-CoV VLPs can be used for the preparation of influenza VLPs according to embodiments of the present invention.

[0174] It is now discovered for the first time that the cellular constituents specifically present in authentic influenza virus particles were also incorporated into the influenza VLP produced from Vero cells. This indicates that the particular interactions between host proteins and viral proteins involved in the biosynthesis of VLPs reflected similar processes as that during virus assembly and budding of the authentic virus particles (Table 1 and FIG. 9). In addition, several cellular proteins are found to be associated with the influenza VLPs according to embodiments of the present invention, but not with the virus particles, suggesting that these proteins may play a role in Vero cell production of influenza VLPs, but not influenza virus particles.

[0175] Accordingly, another embodiment of the present invention relates to an influenza virus-like particle (VLP), comprising: an influenza M1, an influenza M2, an influenza hemagglutinin (HA) and an influenza neuraminidase (NA), and at least one cellular protein of a Vero cell, wherein the influenza proteins are recombinantly expressed from the Vero cell.

[0176] In a particular embodiment of the present invention, the Vero cell is Vero E6 cell.

[0177] The influenza M1, M2, HA and NA in the influenza VLP can be derived from the same or different influenza virus strains. In one embodiment of the present invention, the influenza M1 and M2 are derived from the same influenza virus, and the HA and NA are derived from one or more different influenza virus strains, such as one or more potentially pandemic or seasonal influenza virus strains. In an embodiment of the present invention, the influenza HA and the influenza NA are derived from the same potentially pandemic or seasonal influenza virus strain.

[0178] In another embodiment of the present invention, the influenza VLP comprise two or more HAs and two or more

NAs derived from two or more different influenza virus strains, e.g., two or more potentially pandemic or seasonal influenza virus strains.

[0179] In particular embodiments of the present invention, the influenza VLP comprises the M1 (SEQ ID NO:6) and the M2 (SEQ ID NO:7) of influenza A/Taiwan/083/2006, and the H3 (SEQ ID NO:8) and N2 (SEQ ID NO:9) of influenza A/Taiwan/083/2006.

[0180] In another particular embodiments of the present invention, the influenza VLP comprises the M1 (SEQ ID NO:6) and the M2 (SEQ ID NO:7) of influenza A/Taiwan/083/2006, and H5 (SEQ ID NO:10) and N1 (SEQ ID NO:11) of influenza A/Hanoi/30408/2005(H5N1).

[0181] In one embodiment of the present invention, the cellular proteins in the influenza VLP according to an embodiment of the present invention include those that are also present in authentic influenza viruses, such as those listed in Table 1, e.g., cytoskeleton protein, extra cellular matrix (ECM) proteins, heat shock proteins, annexins, tetraspanins, and glycolytic enzymes.

[0182] In another embodiment of the present invention, the influenza VLP comprises a cellular protein that is not present in influenza virus particles, such as those listed in Table 2, e.g., one or more selected from the group consisting of clathrin heavy chain 1, spectrin beta, plexin B2, CD109 homolog, prostaglandin F2 receptor negative regulator, Na⁺/K⁺-ATPase alpha 1, tumor rejection antigen (gp96) 1 and flotillin I.

[0183] The glycosylation profiles of HA and NA spike in influenza VLPs according to embodiments of the present invention were examined by N-deglycosylation (FIG. 10). In the case of H3N2-VLPs, the glycosylation profiles of HA were highly similar to that of influenza virus replicating in the Vero cells [25]. When examined by transmission electron microscopy (TEM), the influenza VLPs according to embodiments of the present invention were found to closely resemble influenza virus in size, particle morphology, and fine structure of the surface spikes (FIGS. 6 and 7). It was found that influenza VLPs according to embodiments of the present invention stimulated antibody response in mice administered with the VLPs. It was also demonstrated that vaccination of the influenza VLPs according to embodiments of the present invention provided total protection to mouse against avian influenza infection. All these results indicate that influenza VLPs provide a safe and effective means against influenza infection.

[0184] An embodiment of the present invention relates to an immunogenic composition comprising an immunogenic effective amount of the influenza VLP according to an embodiment of the present invention and a pharmaceutically acceptable excipient. The immunogenic composition can further comprise an adjuvant. Any of the pharmaceutically acceptable excipient or adjuvant, such as those described above for the immunogenic composition comprising SARS-CoV VLPs, can be used in the immunogenic composition comprising the influenza VLP.

[0185] In one embodiment of the present invention, the immunogenic composition according to an embodiment of the present invention is administered to a subject to induce immunity against an influenza virus in the subject. The immunogenic composition comprises an influenza VLP comprising the HA and the NA derived from the target influenza virus. The induction of the immunity in the subject results in the prevention, amelioration, or reduction of at least one symptom related to influenza virus infection in the subject. The

“immunogenic effective amount of the influenza VLP” generally refers to the amount of the influenza VLP sufficient to induce immunity to prevent, ameliorate, or reduce at least one symptom related to infection of influenza virus.

[0186] In another embodiment of the present invention, protection against an influenza virus in a subject is provided by vaccinating the subject with a vaccine comprising an influenza VLP according to an embodiment of the present invention, wherein the influenza VLP comprises the HA and the NA derived from the influenza virus.

[0187] In an embodiment of the present invention, a blend of influenza VLPs comprising HAs and NAs from more than one influenza virus strains is administered to a subject to induce immunity against the more than one influenza virus strains in the subject.

[0188] In another embodiment of the present invention, an influenza VLP according to an embodiment of the present invention is used in a method of diagnosing infection by an influenza virus in a subject. The method comprises:

[0189] obtaining a biological sample from the subject;

[0190] contacting the biological sample with the influenza VLP, wherein the influenza VLP comprises the HA and the NA derived from the influenza virus; and

[0191] measuring in the biological sample the amount of an antibody that forms an antibody-antigen complex with the influenza VLP,

[0192] whereby a higher than a threshold amount of the antibody indicates that the subject has ever been infected by the influenza virus or vaccinated by a similar strain of virus.

[0193] The biological sample can be, for example, a serum sample or a tissue fluid. The amount of the antibody that forms an antibody-antigen complex with the influenza VLP can be measured using any methods, such as ELISA, in view of the present disclosure. The threshold amount of the antibody can be determined from a positive control, e.g., a biological sample taken from a subject known to be infected by a threshold titer of the influenza virus.

[0194] Another embodiment of the present invention relates to an antibody against the influenza VLP according to embodiments of the present invention.

[0195] The antibody can be prepared using methods known in the art in view of the present disclosure. For example, the antibody can be prepared by administering the VLP to a vertebrate; and harvesting the antibody against the VLP from the blood of the vertebrate. The antibody can be polyclonal or monoclonal.

[0196] The antibody can be used for various purposes, such as for the treatment or diagnosis of infection by an influenza virus.

[0197] In one embodiment of the present invention, a method of treating infection by an influenza virus in a subject comprises administering to the subject an antibody against an influenza VLP. The influenza VLP comprises the HA and the NA derived from the target influenza virus.

[0198] The antibody can be administered to the subject together with a pharmaceutically acceptable excipients as those described above. The antibody recognizes and binds to antigenic sites on the HA and NA on the surface of the influenza virus, thus mediates an immune response against the influenza virus in the subject. The HA-specific antibody plays an important role for neutralization of the influenza virus, and the NA-specific antibody lessens the release of virus from infected cells. The antibodies can also block the influenza virus from entering cells in the subject or block

replication of the influenza virus. Early intervention with an antibody therapy or passive immunization with immune plasma involving antibodies against influenza VLP can help to control the outbreak of acute influenza virus infection, particularly in subjects with weak, suppressed, or compromised immune systems.

[0199] In another embodiment, the antibody can be used in a method for detecting infection by an influenza virus in a subject. The method comprises:

[0200] obtaining a biological sample from the subject;

[0201] contacting the biological sample with the antibody of claim 17; and

[0202] detecting in the biological sample an antigen that forms an antibody-antigen complex with the antibody,

[0203] wherein the presence of the antigen in the biological sample indicates that the subject is infected by the influenza virus.

[0204] The biological sample can be, for example, serum, throat swap, tear, or tissue specimen, etc., from the subject.

[0205] Embodiments of the present invention include compositions and methods related to the influenza VLPs produced by Vero cells similar to those described above for SARS-CoV VLPs.

[0206] The present invention is further illustrated by the following examples, which are provided for the purpose of demonstration rather than limitation.

Example 1

Expression and Assembly of SARS-VLPs

[0207] Cell Lines and Plasmids

[0208] Vero E6 cells were obtained from American Type Culture Collection (ATCC No. CRL-1586™) and routinely cultured in MEM medium supplemented with 10% fetal bovine serum. Vero E6-based tetracycline-inducible founder cells, Vero/TR, were derived by a stable transfection with the pcDNA6/TR plasmid (Invitrogen). Inducible M-GFP and E expression cassettes were constructed by PCR linking consecutively a β -globin/IgG chimeric intron (from pCI vector, Promega), M-GFP coding sequence, an internal ribosome entry site (IRES) from the encephalomyocarditis virus (ECMV), and an E coding sequence, and the construct was then inserted into the backbone of the pcDNA4/TO plasmid (Invitrogen). Inducible S expression cassette was constructed by inserting a cDNA of the S protein of TW1 strain into the pcDNA5/TO plasmid (Invitrogen). Subsequently, the entire S expression cassette was inserted into the expression plasmid for M-GFP and E to generate the pcDNA4/TO-S-MG-E vector. The sequence of the entire plasmid was verified by DNA sequencing.

[0209] Plasmid Construction

[0210] As shown in FIG. 1A, transgenes encoding the three SARS-CoV envelope proteins, S, M-GFP (i.e., the M protein fused with a green fluorescent protein (GFP) for tracking the VLPs) and E, were constructed in the same plasmid (pcDNA4/TO-S-MG-E). In one plasmid, the vector harbors two expression cassettes. The CMV/TO-MG-E cassette (SEQ ID NO: 1) transcribes an RNA transcript that holds two open-reading frames encoding the M-GFP and E proteins, which are connected by an internal ribosome entry sequence (IRES), and the CMV/TO-S cassette (SEQ ID NO: 2) expresses only the S protein. Both transcription units are regulated by a tetracycline-inducible promoter.

[0211] VLP Expression

[0212] Stable transfection of the pcDNA4/TO-S-MG-E vector into a planned Vero E6-derived founder cell line was conducted to obtain SARS-VLP expression. The founder cell has been previously stably transfected with a tetracycline repressor gene (pcDNA6/TR); therefore, the recombinant SARS-CoV genes will not express until induction. According to the fluorescence intensity of GFP, two clones were selected for prolific production of SARS-VLP, namely Vero/S-MG-E-55 and Vero/S-MG-E-68. Expression of the viral genes was induced by addition of doxycycline (1 μ g/ml) to the cell culture, as verified by RT-PCR for inducible expression of RNA encoding the S, M and E (data not shown). Expression level of VLP in Vero/S-MG-E-55 is higher than Vero/S-MG-E-68, therefore is primarily used.

[0213] For confocal-microscopy analysis, test cells were grown on 12 mm coverslips and treated with doxycycline (1 μ g/ml) for 1 day. Cells were fixed with 4% paraformaldehyde on ice for 20 mins, permeabilized with 0.2% (v/v) Triton X™-100/PBS, and then washed with PBS three times. After blocking in 1% (v/v) fish gelatin/PBST (PBS with 0.1% Tween™-20) for 1 hr, samples were incubated with a specific antibody at 4° C. for 18 hrs, followed by 3 washes with PBST, and then probed with the respective fluorescence-conjugated secondary antibody for 1 hr at room temperature. Finally, samples were washed with PBST three times and mounted in mounting medium (Vector). The samples were scanned for GFP and antibody-stained signals, thereafter analyzed for co-localization according to the manufacturer's software (Zeiss LSM 510 META).

[0214] Upon induction, GFP dots appear evident inside the producer cells within one day and accruing for longer than five days as shown in microscopic studies (FIG. 1B). The GFP dots of various sizes in the cytoplasm from the peri-nuclear region toward plasma membrane, showing indicative pattern all along the secretory pathway of mammalian cells from endoplasmic reticulum (ER) to the plasma membrane. This intracellular distribution corresponds with the CoV assembly of SARS and others, which is located at ER-Golgi-intermediate-compartment.

[0215] Intracellular expression of each VLP component protein (S, M, E and GFP) and their assembly were next inspected by immuno-fluorescence staining and overlaid with the fluorescence tracks of GFP, as exemplified by VLP producer cells induced for one day (FIG. 1B). Staining with antibodies against either M protein or GFP results in signals which completely overlap with GFP tracks and thus indicates that GFP fusion faithfully labels the M protein (FIG. 1B). In addition to peri-nuclear staining (Golgi complex), the S protein is stained intensely as reticular ER pattern in addition to the profiles of Golgi and secretory vesicles (FIG. 1B). However, the co-localization of S protein with M-GFP principally limits to Golgi and secretory vesicles. More S protein accumulates in ER, suggesting its longer duration for de novo synthesis and glycosylation in ER. While most secretory M-GFP dots co-localize with the staining of E protein, peri-nuclear M-GFP shows two ways, positively and negatively co-localized with E protein. These data collectively suggest that E protein as soon as it is translated initiates VLP assembly with M-GFP and S protein nearby the Golgi and resulting in punctual co-staining of M-GFP, E, and S proteins as secretory vesicles (FIG. 1B). As negative controls, the same immuno-staining with the S, M or E Ab in parental Vero E6 cells detected no signals; neither was seen for fluorescence tracks

of GFP (data not shown). In agreement with previous studies on CoV budding, VLP assembly for SARS-CoV and others in mammalian cells, our data indicated the peri-nuclear assembly of SARS-M, E and S and their co-localization in a secretory vesicle profile. Assembly of the three proteins into SARS-CoV-like particles is further demonstrated by their co-sedimentation in a sucrose gradient and forming spiky spherical particles (FIG. 2D).

Example 2

Purification and Characterization of SARS VLPs

[0216] Purification of VLP was initially performed by concentrating conditioned culture medium of the induced cells on a 45% sucrose cushion by ultracentrifugation at 200,000×g at 4° C. for 2 hrs. The interface was collected and further separated on a step-wise gradient between 25% and 35% sucrose at 200,000×g at 4° C., for 48 hrs. Sedimentation fractions were taken from the bottom of the tube every 0.5 ml volume. Each fraction was analyzed for protein concentration by Coomassie (Bradford) Protein Assay Kit (Pierce) and GFP fluorescence measured by VICTOR²™ fluorometer (PerkinElmer).

[0217] For western blot analysis, polyclonal antibodies against E and M proteins were separately raised in rabbits using *E. coli* expressed M (a.a. 53-221 of SEQ ID NO: 3) and E (a.a. 1-76 of SEQ ID NO: 4) proteins as antigens by intrasplenic injection. Anti-S polyclonal antibodies were raised in ducks using *E. coli* expressed S (a.a. 679-888 of SEQ ID NO: 5) as antigens, and IgY antibodies were purified from egg white (Wu, H. S. et al., 2004, *J Biomed Sci* 11: 117-126).

[0218] As shown in FIG. 2A, distribution of both proteins and GFP exhibited a coherent major peak concentrated in 25% sucrose layer (fractions 9 to 15). Unexpectedly, we also find a minor protein peak concentrated in 35% sucrose layer, which is absent in the Vero/S-MG-E-68 clone (fractions 2 to 6). Protein analysis by SDS-PAGE and Coomassie blue staining reveal that the two distinct peaks are obviously of different protein compositions (FIG. 2B). Each VLP constituent protein of expected size as marked in FIG. 2B is confirmed by western blot analysis using specific antibodies against S, M, E and GFP proteins (FIG. 2C). The SARS-VLP contains multiple forms of S protein, predominantly of mature form with apparent M_r 180 kDa (○), and less with 170 kDa (★), and 140 kDa (+) (FIG. 2B). According to previous studies on individual expression of S and M proteins in mammalian cells, the 180 kDa (○) band represents a complex-type glycosylated form (EndoH-resistant yet PNGaseF-sensitive); the 170 kDa (★) band represents a high-mannose-type glycosylated form (EndoH-sensitive); and the 140 kDa (+) band represents a non-glycosylated form. The purified SARS-CoV contains two forms of M. The more abundant form with apparent M_r 22 kDa is not glycosylated, and the less abundant 27 kDa form contains an EndoH-sensitive, high-mannose-type N-glycan linked to the Asn-4 residue (Voss, D. et al., 2006, *supra*). In agreement, the M-GFP in SARS-VLP is mainly of 65 kDa (#), and less of 70 kDa (*) (FIG. 2B). Since GFP fusion contributes M_r ~27 kDa, both forms of M-GFP in SARS-VLP show an additional 16 kDa increase in apparent M_r due to unknown reasons. The E protein associates with M protein in sucrose gradient sedimentation and perhaps lacks glycosylation as per its 9 kDa size.

[0219] The SARS-VLP resides in the expected major peak; whereas the unexpected minor peak comprises primarily S

protein of the 170 kDa form, less for M-GFP of the 65 kDa form, but no E protein, and not observable by electron microscopy; therefore, it is not further characterized here (data not shown). The SARS-VLP of interest to the inventors (i.e., fractions 9 to 15 in FIGS. 2A-2C) contains primarily M-GFP, with less S protein and the least E protein, which is a ratio similar to CoV of SARS and others. The S protein of the secreted SARS-VLP is the predominant 180 kDa form containing complex-type N-glycans whose maturation was suggested to occur before S protein trimerizes. All forms of S protein contained in the secreted SARS-VLP were found not cleaved as estimated by their mobility in SDS-PAGE.

[0220] Morphology of the SARS-VLP was further examined by transmission electron microscopy (EM). For EM, aliquots of 10 μl of purified SARS-VLPs were loaded onto a carbon-coated grid, and let stand still for 3 mins. Grids were then stained with 2% uranyl acetate for 2 mins, and examined directly under an electron microscope. As can be seen in FIG. 2D, the negatively-stained VLP appeared as spherical particles with a spiky surface resembling SARS-CoV particle and a diameter ranging between 50 nm and 70 nm. The diameter of Vero E6 cell-secreted empty VLP is smaller than the extra-cellular whole SARS-CoV whose diameter is between 60 nm and 100 nm.

[0221] Noteworthy, the protein yield of the SARS-VLP described above is fascinatingly high, which makes the system very attractive for all relevant applications. The result demonstrated in FIG. 2 represents a routine purification of VLP from a pool of 750 ml culture medium collected on day 3 and day 5 after induction. Summation of fractions 9 to 15 (3 ml in each fraction) yields 250 mg protein of purified VLP in total (FIG. 2A). The inventors' routine yield of mammalian cell-based SARS-VLP from Vero/S-MG-E-55 cells is 449.7±69.3 (N=12) mg/L of culture medium (using 1.2×10⁸ producer cells), and is over 1,000-fold higher than the reported level of insect cell-based SARS-VLP (200 μg/L×10⁹ host cells, estimated to be 0.5 to 1 L of cell culture) (Mortola E. and Roy, P., 2004, *supra*). The inventors believe the unprecedentedly high expression level of SARS-VLP in this study may result from the best match of Vero E6 as host cells to express the SARS viral proteins and insertion of the transgenes into a chromatin position which is highly active in gene transcription, because the inventors also isolated many other transgenic Vero E6 clones whose intracellular expression of GFP dots were at apparently lower levels. However, it may also involve with the much stronger expression from the inducible CMV promoter used in our cell line. Production of SARS-VLPs in Vero E6 cells by stable transfection gives the best high yields to the inventors' understanding and the production process is ready to be adapted for large scale manufacture, offering an attractive approach for development of an effective and economical vaccine.

Example 3

Vaccination Experiments with SARS-VLPs

[0222] With the high-yield SARS-VLP available from mammalian expression as described above, the subsequent important question is its immunogenicity and SARS-CoV-neutralizing antibody response. To address this issue, the inventors designed a series of vaccination experiments in mice and examined the systemic immune responses (FIG. 3A). Groups of four female C57BL/6 mice, 6-8 weeks of age, were s.c. injected with 20 μg of SARS-VLP in 100 μl of PBS

without additional adjuvant, and boosted with different dosages (0, 5 μ g, 20 μ g) after 2 weeks. Mock immunization mice were injected with 100 μ l of PBS as controls.

[0223] Immunization with SARS-VLP Elicits an Antigen-Specific IgG1 Response in Mice.

[0224] Two weeks after booster immunization, serum titers of antigen-specific IgG were measured by ELISA using native SARS-VLPs as the absorbent antigen. For ELISA, serum was collected by tail vein bleeding, allowing clotting at 4° C. overnight and cleared up by centrifugation. ELISA plates (Nunc) was coated with 1 μ g native VLP at 4° C. overnight and blocked with 5% dry milk in PBS. ELISA plates were then incubated with serum samples of indicated dilution at 37° C. for 1 hr, traced with HRP-conjugated secondary antibodies, and developed color with TMB substrate (PIERCE). Washes with PBST for 5 times were applied between each step of ELISA. Finally, the ELISA was read out with absorbance of 450 nm wavelength (A_{450}) by a microplate reader (Power Wave XS, Bio-Teck). VLP-specific IgG titer (A_{450}) was calculated by subtracting the background readout of mock samples.

[0225] As shown in FIG. 3B, a single dose of 20 μ g VLP positively induced antibody response up to 50-fold. The specific antibody titers were dose-dependently increased by a booster immunization for over 6250-fold (FIG. 3B). Similar ELISA for various IgG subtypes detected that the antibody response mainly restricted in IgG1 subtype which generally acts on neutralization (FIG. 3B). In contrast, IgG2a subtype of VLP-specific antibody titer was very low in these experiments (FIG. 3B). Together, the response of antibody subtype indicates an induction of T_H2 -type effector functions against the epitopes on the SARS-VLP surface. Most prominently, the IgG antibody stimulated by the SARS-VLP effectively cross-reacted with genuine SARS-CoV virion inactivated by the gamma-radiation and heat, as demonstrated by ELISA using a commercial kit advised in the World Health Organization website (FIG. 3C). The antigen-specific antibody in mice serum retains high titers for longer than 4 weeks following the booster immunization, indicating a long persistence of antibody response caused by SARS-VLP immunization (FIG. 3D). The ELISA results in FIG. 3B-3C are particularly meaningful to SARS-CoV neutralization because they discern the antibody that binds surface of VLP and whole virus. These results endorse the resemblance in surface between the VLP and intact SARS-CoV and indicate a potential neutralizing antibody response induced by SARS-VLP vaccine in mice.

[0226] SARS-VLP-Induced Serum IgG Antibodies Recognize S and M Proteins.

[0227] The antigen determinants with which VLP protein the mouse anti-bodies would react were examined by western blot assay loaded with three different amounts of SARS-VLP. As shown in FIG. 3E, the VLP-specific antibody detects the most intensely against M-GFP, followed by S protein, and minimally to E protein. The VLP-specific antibody efficiently reacted with all forms of S and M proteins. The observations specify that M and S proteins in the context of SARS-VLP are much more immunogenic than E protein, which also agrees with the antibody specificity found in SARS patients.

[0228] Immunization with SARS-VLP Induces Antigen-Specific T Helper (T_H) Responses in Mice.

[0229] The type of T_H response upon SARS-VLP vaccination was investigated by IFN- γ and IL-4 ELISPOT (enzyme-linked immunospot) assays for commitment to secrete T_H1

and T_H2 cytokines by splenocytes. For ELISPOT assays, PVDF-bottom plates (Millipore) were coated with 0.1 ml INF- γ and IL-4 capture antibodies (1:60; R&D systems) at 4° C. overnight. After washing with PBS twice, the plates were and then blocked with 1% BSA in PBS at room temperature for 4 hrs. Splenocytes were isolated from tested mice 14 days after booster administration, and allowing erythrocyte lysis. Splenocytes of single cell were suspended in RPMI containing 10% heat-inactive FBS, 50 μ M β -mercapto-ethanol, and 3×10^5 cells/well were grown in INF- γ or IL-4 ELISPOT plates with 1 μ g VLP for 40 hrs. Washes with PBST for 5 times were applied between each step of ELISPOT. The plates were incubated with 0.1 ml biotinylated INF- γ or IL-4 detection antibodies of 1/60 dilution (R&D systems) at 4° C. overnight, incubated with streptavidin-alkaline phosphatase of 1/60 dilution (R&D systems) at room temperature for 1.5 hrs, washed, and rinsed twice with water. The color of ELISPOT was developed in darkness for 30 mins with BCIP/NBT solution (R&D systems). Development was stopped by washing with water and air-dried. The signals were counted by ImmunoSpot analyzer and analyzed by ImmunoSpot software (CTL).

[0230] When the primary culture of splenocytes isolated from SARS-VLP-immunized mice re-exposed to SARS-VLP ex vivo, both INF- γ - and IL-4-producing populations rise along with the booster dose of SARS-VLP, indicating development of VLP-recognizing T_H1 cells and T_H2 cells in spleen provoked by SARS-VLP vaccination dose-dependently in vivo (FIGS. 4A, 4B). However, a T_H2 -biased Ab response as indicated by induction of IgG1-dominant antibodies in serum further indicates the effector function of T_H1 cells in vaccinated mice was to activate CTL. Further, both T_H1 and CTL can secrete INF- γ when DC presents them against the VLP-antigens (FIGS. 3B, 3C). Together, these data demonstrate that SARS-VLP per se is a potent vaccine that raised humoral and cellular immune responses.

Example 4

Expression and Purification of Influenza VLPs from Vero Cells

[0231] Establishment of Vero Cell System Producing Influenza VLPs

[0232] Vero cells were obtained from the Bioresource Collection and Research Center, (Hsinchu, Taiwan) and maintained in minimal essential medium (HyClone, South Logan, Utah) supplemented with 10% fetal bovine serum (Gibco, San Diego, Calif.) in a humidified incubator at 37° C. with 5% CO₂.

[0233] The cDNAs of M1 and M2 derived from the sequences of influenza A/Taiwan/083/2006 virus were cloned into the backbone modified from pcDNA6/TR (Invitrogen, Carlsbad, Calif.) linked by IRES separately into a single eukaryotic expression vector to give the plasmid of pCI6/TO-M1-M2 (FIG. 5A). The amino acid sequences of the encoded M1 and M2 are SEQ ID NO:6 and SEQ ID NO:7, respectively. The cDNAs of HA and NA were synthesized sequences based on the distinct virus strains of A/Taiwan/083/2006 and A/Hanoi/30408/2005(H5N1) (a kind gift from Dr. Po-Huang Liang at Institute of Biological Chemistry, Academia Sinica) optimized for mammalian codon usage and further cloned into expression vectors as illustrated for pCI4/TO-HA-NA (FIG. 5A). The amino acid sequences of the encoded HA and NA based on A/Taiwan/083/2006 are SEQ

ID NO:8 and SEQ ID NO:9, respectively. The amino acid sequences of the encoded HA and NA based on A/Hanoi/30408/2005(H5N1) are SEQ ID NO:10 and SEQ ID NO:11, respectively.

[0234] For the H3N2- and H5N1-VLPs producing Vero cells, the plasmid pCI6/TO-M1-M2 was stably transfected into Vero cells to derive a founder Vero cell line, which was further transfected with HA-NA expression vectors to obtain the quadruple co-expression Vero cell line with HA, NA, M1, and M2 proteins.

[0235] To confirm the gene expression of N2 in the H3N2-VLP producer cell line, total RNAs were extracted separately from cells with and without doxycycline (Dox) induction and RT-PCR assays performed using a primer pair corresponding to the internal sequence of the N2 gene. These primers were N2-F, 5'-ATTAGCTTTCCGCTGGTGGGGACAT-3' (SEQ ID NO:16) and N2-R, 5'-GCATTCTGACTCCTGGGTCCTGAGGATT-3' (SEQ ID NO:17).

[0236] Expression of the proteins was confirmed by Western blot analyses and immunofluorescence staining as follows. Quadruple VLP-expression cells were induced with Dox for 48 h, or left untreated as a control. The cells were then fixed in 4% paraformaldehyde for 10 min and immersed in 0.05% Triton-X 100 for 1 min. After blocking with 1% gelatin, the cells were incubated with distinct primary specific antibodies, followed with goat anti-mouse or goat anti-rabbit IgG conjugated with Cy3 dye. Fluorescence images were acquired by confocal microscopy (LSM 510 META NLO DuoScan, Carl Zeiss, GmbH). The antibodies used in this study were polyclonal: H3 (ab20084), N1 (ab21305), M1 (ab20734), annexin A2 (ab41803), and clathrin (ab21679) from Abcam (Cambridge, Mass.), β -actin (sc-1616-R) from Santa Cruz Biotechnology (Santa Cruz, Calif.), and monoclonal: M2 (ab5416), and tubulin (ab6160) from Abcam and H5 (MCA2660, used for IFA) from AbD Serotec (Raleigh, N.C.). Rabbit polyclonal antibody against H5 used for Western blotting was provided by Dr. Che Ma (Genomics Research Center, Academia Sinica).

[0237] In this study, mammalian cell culture-based approaches were used to generate influenza VLPs. To stably transfect the viral genes of HA, NA, M1, and M2 critical for VLP production into cultured Vero cells, such as Vero E6 cell, four gene expression cassettes were designed and placed into two vectors as illustrated (FIG. 5A). A tetracycline repressor gene and tet operator-regulated gene expression cassette were also inserted in the plasmid expressing M1 and M2, giving the vector pCI6/TO-M1-M2. By stable transfection of the M1-M2 vector into Vero cells, founder cells that would not express the M1-M2 transgene until doxycycline (Dox) induction were constructed. Another two tet operator-regulated, CMV promoter-driven expression cassettes were inserted separately into the other plasmid carrying HA and NA genes, giving the expression vector pCI4/TO-HA-NA. After stable transfection of pCI4/TO-HA-NA vector into an M1-M2 founder cell line, a mammalian-expressed VLP system in Vero cell was established.

[0238] To verify that co-expression of all four viral genes was indeed driven by the inducible promoter (CMV-TO), total cell lysates of quadruple-transfected Vero cell line of H3N2 were analyzed by Western blot with specific viral antibodies against H3, M1, and M2 (FIG. 5B). As there is no available antibody to N2, RT-PCR was used to confirm the expression of the N2 gene. The cellular localization of H3 was also observed by confocal laser scanning microscope (FIG.

5D). The utility of this system as an alternative platform to reverse genetics for vaccine development was shown by the simple substitution of a separate plasmid carrying the HA and NA genes of H3N2 with those of H5N1 (FIGS. 5C and 5E). The resulting quadruple H5N1-VLP cell line again co-expressed HA and NA, this time of the H5 and N1 varieties. Two subtypes of quadruple Vero cell lines that generated the putative H3N2- and H5N1-VLPs, respectively, have been constructed and verified. Vero cell expression systems for producing other influenza VLPs can also be constructed and verified using similar methods in view of the present disclosure.

[0239] Microcarrier Culture and Purification of Influenza VLPs Made from Vero Cells

[0240] To scale up the cultivation of VLP producer cells, 60 g microcarriers (HyClone) and cells (about 2×10^8) were added to a 3 L spinner flask (BellcoGlass com., Vineland, N.J.), stirred at around 35 rpm with a pendant glass ball, and maintained in minimal essential medium supplemented with 10% fetal bovine serum in a humidified incubator at 37° C. with 5% CO₂. After 7 days cultivation, the cells had attached to the surface of the collagen-coated microcarrier and grown to confluence. For VLP expression and secretion from cells, the culture medium was removed and replaced with serum-free medium (SFM4 MegaVir, HyClone) containing 1 μ g/mL Dox to begin induction.

[0241] After Dox-induction for 72 h, the conditioned medium of VLPs producer cells was harvested, filtered with 0.45 μ m Stericap, concentrated by Vivaflow 50 (Sartorius Stedim Biotech, Göttingen, Germany), and then layered onto a 30% sucrose-TNE (10 mM Tris-HCl, pH 7.4, 100 mM NaCl, 1 mM EDTA) cushion. Following centrifuge at 112,600 \times g for 2 h at 4° C. in a Beckman SW28 rotor (Beckman Coulter, Fullerton, Calif.), the resulting pellet was resuspended in TNE buffer, and further purified over a 20-30-60% sucrose gradient (112,600 \times g, 2 h at 4° C.). Finally, the banded VLPs were collected, dialyzed with TNE buffer overnight, and stored at $\pm 80^\circ$ C. To analyze the protein constituents of purified VLPs, the samples quantified by Quant-iT Protein Assay Kit (Invitrogen) were mixed with Lämmle SDS-PAGE sample buffer, boiled for 5 min, and separated in a 7.5-17.5% gradient gel.

[0242] The mammalian Vero cell system successfully produced influenza VLPs on a preclinical scale by stable co-expression of four viral proteins: M1, M2, HA and NA. Typically, the pilot production of 3 L-scale microcarrier systems yielded an average 1.2 mg/L medium of influenza VLPs after purification. Each inoculation of 3 L-culture attained 10^9 cells and could be induced to continuously express VLPs three times.

Example 5

Characterization of Influenza VLPs Made by Vero Cells

[0243] Morphology and Antigen Presentation of Purified H3N2 VLPs and H5N1 VLPs

[0244] Immunogold electron microscopy was performed on purified influenza VLPs made from Vero cells. Sucrose gradient purified influenza VLPs of 1 μ g were adsorbed onto formvar/carbon-coated nickel grids (Electron Microscopy Sciences, Fort Washington, Pa.). After a 2 min wash with TBS buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl), the sample was blocked with 1% BSA in TBS for 1 h. Primary antibody

(10 $\mu\text{g}/\text{mL}$) was diluted in 1% BSA/TBS and adsorbed onto the grid for 1 h at room temperature. Following three washes with TBS, secondary gold-conjugated antibody was added for 1 h at room temperature. The grids were then washed twice with TBS, fixed with 1% glutaraldehyde, washed with water, and negatively stained with 2% uranyl acetate for 30 sec. The images of stained mammalian VLPs were captured using a Hitachi H-7000 transmission electron microscope.

[0245] The morphologies of mammalian VLPs purified from culture medium of transfected cells were negative stained with 2% uranyl acetate and observed by TEM as compared to their corresponding viruses propagated in Vero cells (FIGS. 6A and 6B).

[0246] The influenza VLPs displayed generally spherical morphologies and densely stained cores. The spike projections on the surface of VLPs were no different in appearance to those on authentic influenza viruses. The HA and NA glycoproteins on the surface of VLPs were immunogold labeled with individual specific antibodies, and counter-stained with gold spheres coupled to secondary antibodies (FIGS. 6C and 6D).

[0247] Dynamic Light Scattering (DLS) Determination of Average Particle Size of Influenza VLPs

[0248] In complement the morphology analysis by TEM, the average sizes of secreted VLPs in native solution were estimated by DLS assays. To be an effective vaccine, it has been proposed that particles ranging from 20 to 200 nm could facilitate the drainage of free antigens to the lymph nodes and induce strong responses in dendritic cells (DC) for long-term protective purposes [15]. Laser-based DLS can monitor changes in Brownian motion of nanoparticles in solution, giving information related to the average size and frequency distribution of particles.

[0249] Stock solutions of influenza VLPs were diluted to 0.1 $\mu\text{g}/\text{mL}$ in 20 mM phosphate buffer at pH 7.4, passed through 0.45 μm filters, and analyzed on a Nano ZS particle-size analyzer (Malvern Zetasizer, Malvern Instruments Ltd, UK). For each sample analyzed by DLS, two consecutive measurements were taken on a single sample and measured with a light-scattering data collection time of 60 sec according to the manipulation instruction. The accompanying software (Nanov510) was used to convert the intensity-based measurement to a size distribution based on the number of particles in each size class, and was presented as a diagram of curves showing the frequency distribution of the sample where the area under the curve was proportional to the numbers of VLPs detected in the relevant size range. The average diameters of VLPs were then calculated as the mean size of VLP population \pm standard deviation (SD) of three independent experiments.

[0250] As shown in FIG. 7, DLS here revealed the average diameters of H3N2- and H5N1-VLPs were 108.2 ± 17.9 nm and 125.6 ± 10.5 nm, respectively, at pH 7.4, 25° C. The sizes of the VLPs are comparable to the sizes of their corresponding viruses, e.g., 133.5 ± 15.4 nm and 104.1 ± 12.4 nm for H3N2 and H5N1 viruses, respectively. The size distributions of both subtypes of VLPs ranged from 70-200 nm (95% CI), suggesting that the influenza VLPs produced by Vero cells were in the preferred size range for DC uptake and promise to stimulate a potent immune response (FIG. 7). DLS will be a useful approach to monitor the batch-to-batch consistency of VLPs by rapidly providing information on the whole population of particles. Together, the DLS and TEM measurements of influ-

enza VLPs were consistent and showed the VLPs made by Vero cells to be of comparable size and morphology to native influenza viruses.

[0251] Identification of Influenza VLP Composition and VLP-Associated Cellular Proteins

[0252] To verify the protein constituents of mammalian expressed VLPs, 10 μg of H3N2- and H5N1-VLPs were separated on a 7.5%-17.5% gradient gel, and stained with Coomassie blue (FIGS. 8A and 8C) or probed with specific antibodies against viral proteins in separate experiments (FIGS. 8B and 8D). Besides the viral proteins of HA, NA, M1, and M2, a wide spectrum of minor bands were also observed in the influenza VLPs similar to those in the authentic viruses (FIGS. 8A and 8C).

[0253] To identify the basic protein profiles of these VLPs, the more obvious protein bands (indicated by arrows in FIGS. 8A and 8C) were excised from the gels, subjected to in-gel trypsin digestions, and analyzed by liquid chromatography tandem mass spectrometry (LC/MS/MS). Briefly, the protein bands from 1-D gel were manually excised from the gel and cut into small pieces (~ 0.5 mm³). The gel pieces were washed in a microcentrifuge tube with a solution containing 50% methanol and 5% acetic acid for 2-3 h, twice with a solution of 25 mM NH_4HCO_3 in 50% acetonitrile for 10 min each, and then dried in a vacuum centrifuge. After DTT reduction and iodoacetamide alkylation, a solution containing 75 ng of sequencing grade modified trypsin (Promega Corporation, Madison, Wis.) in 25 μL of 25 mM NH_4HCO_3 was added and incubated with dried gel pieces at 37° C. for 12-16 h. Following digestion, tryptic peptides were extracted twice with 50% acetonitrile containing 5% formic acid for 15 min each time with vortex. The extracted solutions were pooled and evaporated to dryness under vacuum. The dried pellet was re-dissolved in 10-20 μL of 0.1% formic acid for LC/MS/MS analysis as described below. The NanoLC-nanoESI-MS/MS analysis was performed on a nanoAcquity system (Waters, Milford, Mass.) connected to an LTQ-Orbitrap XL hybrid mass spectrometer (Thermo Electron, Bremen, Germany) equipped with a PicoView nanospray interface (New Objective, Woburn, Mass.). Peptide mixtures were loaded onto a 75 μm ID, 25 cm length C18 BEH column (Waters) packed with 1.7 μm particles with a pore size with of 130 Å and were separated using a segmented gradient in 90 min from 5% to 50% solvent B (acetonitrile with 0.1% formic acid) at a flow rate of 300 mL/min and a column temperature of 35° C. Solvent A was 0.1% formic acid in water. The mass spectrometer was operated in the data-dependant mode. Briefly, survey full-scan MS spectra were acquired in the orbitrap (m/z 350-1600) with the resolution set to 60,000 at m/z 400 and automatic gain control (AGC) target at 106. The 10 most intense ions were sequentially isolated for CID MS/MS fragmentation and detection in the linear ion trap (AGC target at 7000) with previously selected ions dynamically excluded for 90 sec. Ions with single and/or unrecognized charge state were also excluded.

[0254] The MS and MS/MS raw data were processed with Bioworks 3.3.1 and searched against an in-house generated NCBI protein database, using a Mascot Daemon 2.2 server. Search criteria used were trypsin digestion, variable modifications set as carbamidomethyl (C) and oxidation (M), allowing up to 2 missed cleavages, mass accuracy of 10 ppm on the parent ion and 0.60 Da on the fragment ions.

[0255] The major VLP constituents identified by searching against the NCBI database are shown as Tables 1 and 2. It was found that HA and NA proteins of mammalian VLPs were

distributed in multiple gel slices in addition to the expected locations shown in Table 1. This likely reflects the fact that they are the most abundant proteins in VLPs and form HA and NA protein oligomers (FIGS. 8B and 8D).

[0256] In addition, another 22 VLP-associated cellular proteins were identified, which are identical or functionally

analogous to those cellular proteins commonly found in the interior or exterior of influenza virions [16]. Most of them could be classified into functional groups including cytoskeleton protein, extra cellular matrix (ECM) proteins, heat shock proteins, annexins, tetraspanins, and glycolytic enzymes.

TABLE 1

Viral and cellular proteins associated with influenza VLPs identified by mass spectrometry (LC/MS/MS). SDS-PAGE and LC/MS/MS Analysis						
Protein Type	Protein Name	Mass (Da)	Protein band number ^a H5N1/H3N2*	Mascot score ^b	Sequence coverage (%) ^c	Reported in influenza virion ^d
Viral proteins	Hemagglutinin (H5)	64163	5, 6, 10 ^b	227, 812, 1053	26, 18, 40	yes
	Neuraminidase (N1)	51298	6	71	10	
	Hemagglutinin (H3)	63595	/4	/239	/24	
	Neuraminidase (N2)	52018	/5	/230	/40	
	Matrix 1 protein (M1)	27860	10/10	35/45	36/16	
	Matrix 2 protein (M2)	11157	11/11	34/44	50/35	
Cytoskeletal proteins	β -actin	41710	8/8	1519/447	72/73	yes
	β -tubulin	49639	7/7	41/655	15/72	yes
	myosin IA	118204	2/2	175/41	7/2	tropomyosin
	Similar to myosin IC	107102	3/3	898/107	42/15	4 & 1
ECM proteins	integrin alpha 3	118333	2/2	896/69	27/20	Integrin beta 1
	integrin alpha 5	115919	2/2	687/174	37/22	
Heat shock proteins	Heat shock 90 kDa protein	83185	4/4	435/368	40/43	HSP 27 kDa
	Heat shock 70 kDa protein	72288	5/5	615/503	54/50	
	Heat shock 27 kDa protein	22768	6	112	44	yes
Annexin	annexin A11	54443	9/7	284/282	38/38	yes
	annexin A2	53564/ 38576	7, 9/ 7, 9	1087, 361/ 153, 613	43, 41/ 26, 49	yes
Tetraspanin	CD81 molecule	25741	6	117	33	yes
	CD9 molecule	25380	7	27	11	yes
Glycolytic enzymes	enolase 1	47182	9	1011	70	yes
	Similar to phosphoglycerate kinase 1	44558	8	572	71	Phosphoglycerate kinase
	pyruvate kinase	64479	1/6	248/354	45/51	yes
	glyceraldehyde-3-phosphate dehydrogenase	35959	6/6	64/48	23/25	yes
Unclassified proteins	2',3'-cyclic nucleotide 3' phosphodiesterase	47509	8	306	56	yes
	Aldo-keto reductase family 1	35992	7	99	23	Yes
	WD repeat domain 18	48167	8	78	18	WD repeat-containing protein 1

TABLE 1-continued

Viral and cellular proteins associated with influenza VLPs identified by mass spectrometry (LC/MS/MS). SDS-PAGE and LC/MS/MS Analysis						
Protein Type	Protein Name	Mass (Da)	Protein band number ^a H5N1/H3N2*	Mascot score ^b	Sequence coverage (%) ^c	Reported in influenza virion ^d
	Gamma-glutamyl-transferase 1	61261	6/6	76/71	4/10	Yes
	Peroxiredoxin 2	21878	7	28	9	Peroxiredoxin 1

^aExcised protein bands were numbered as indicated by arrows 1-11 of FIG. 4A and C.

^bFor this search a Mascot score ≥ 25 is significant ($p < 0.05$).

^cSequence coverage is based on peptides with unique sequence.

^dViral and cellular proteins have been reported in influenza virion [17].

*The band number, Mascot score, and sequence coverage of cellular proteins both identified in H5N1- and H3N2-VLPs are presented as A/B.

^aThe band number, Mascot score, and sequence coverage of a single protein distributed in multiple locations is presented as A, B.

[0257] Apart from those common proteins identified in virion and VLPs, several unique cellular proteins (as listed in Table 2) were also identified in influenza VLPs with very high Mascot scores in LC/MS/MS analysis. These proteins possibly involved in VLPs biosynthesis.

[0258] For real viruses, recruitment or encapsidation of some cellular proteins into the virion may be a critical behavior supporting the completion of the life cycle by some specific interaction with viral proteins or RNA. However, in this case, the mammalian influenza VLP has the components of viral (by transfection) and cellular proteins (by recruitment) without package of any viral genetic material. The cellular proteins identified in the influenza VLPs might be actively involved in the normal virus life cycle, especially during virus assembly and budding from the host cells.

TABLE 2

Unique cellular proteins identified by LC/MS/MS with high Mascot scores in mammalian VLPs.				
Protein Name	Mass (Da)	Protein band number ^a H5N1/ H3N2*	Mascot score ^b	Sequence coverage (%) ^c
Clathrin heavy chain 1 (Cytoplasmic vesicles formation)	192276	1	3242	63
Spectrin (interacts with actin)	274472	1	1146	35
Beta (non-erythrocytic) Plexin B2 (interacts with cytoskeleton)	203451	1	757	26
Similar to CD109	161515	1	738	23
Prostaglandin F2 receptor negative regulator	116885	2/2	1035/136	44/18
Na ⁺ /K ⁺ -ATPase alpha 1	112838	3/3	1266/452	42/29
Tumor rejection antigen (gp96) 1	92555	3/3	1016/614	52/60
Flotillin 1	47384	9	1048	68

^aExcised protein bands were numbered as indicated by arrows 1-11 of FIG. 4A and C.

^bFor this search a Mascot score ≥ 25 is significant ($p < 0.05$).

^cSequence coverage is based on peptides with unique sequence.

*The band number, Mascot score, and sequence coverage of cellular proteins both identified in H5N1- and H3N2-VLPs are presented as A/B.

[0259] To characterize the functionality of HA spike on mammalian VLPs, hemagglutination assays were performed

(FIG. 8E). VLP preparations reacted with 0.75% guinea pig erythrocytes had significant hemagglutination activities, with titers of 2^7 for H3N2-VLPs and 2^6 for H5N1-VLPs in samples containing 3.5 μg of VLPs. Both H3N2 and H5N1 viruses of same amount as VLPs were at similar titers (2^7) of HA activity. This result suggests the HA spikes compassing around the surface of mammalian VLPs are in a native orientation and function as those of active authentic viruses.

[0260] Generally, the results of two independent LC/MS/MS analyses of H5N1- and H3N2-VLPs and Western blotting in this study resemble the reported proteome of influenza virus [16], which suggests that the assembly of influenza VLPs shares much similarity with real virus assembly and escape. Taken together, these data confirm that the mammalian-expressed influenza VLPs are very similar to the authentic viruses, a considerable advantage to their use in further vaccine development.

[0261] Confirmation of Cellular Proteins Associated and Incorporated into Influenza VLPs

[0262] Following proteomic identification of incorporated cellular proteins, several viral and associated cellular proteins were further characterized by Western blot and immunogold labeling. To rule out the possibility that identified proteins may be due to non-specific contamination such as co-purification of microvesicles or exosomes with VLPs, the mammalian influenza VLP preparations were subjected to a protease protection assay that has been shown to efficiently remove microvesicles from HIV-1 virion preparations [17,18].

[0263] Purified H5N1-VLPs equivalent to 50 μg of proteins were incubated with or without 20 μg of MSG-Trypsin (G-Biosciences, St. Louis, Mo.) in 20 mM Tris-HCl, pH 8.0 and 1 mM CaCl₂ for 18 h at 37° C. After trypsin incubation, the treated and untreated VLPs preparations were separately diluted to 7 mL with THE buffer containing 30 μM PMSF (Sigma, St. Louis, Mo.), and concentrated through a 20% sucrose cushion by ultracentrifugation (200,000 \times g, 2 h at 4° C. in a Beckman SW41Ti rotor) and then subjected to Western blot analyses with antibodies against HA, NA, M1, M2, β -actin, tubulin, annexin A2, and clathrin (FIG. 9A).

[0264] Both HA and NA proteins were lost from the VLPs after trypsin digestion, demonstrating that these proteins were located on the outside of the membrane envelope, integral and attached to the VLP surface. Most co-purified con-

taminants were eliminated by the protease digestion. However, several representative cellular proteins expected to be inside the virion such as actin, tubulin, and annexin A2 were still found to be present in the protease-digested VLPs, which indicates that these cellular proteins were specifically incorporated into the structure of VLPs (FIG. 9A).

[0265] In contrast, a unique cellular protein identified in this study, clathrin heavy chain, an important component of the clathrin-coated pits mediating the endocytosis of many receptors, ion channels, transporters, and other transmembrane proteins as well as various soluble macromolecules and viruses, was lost following protease treatment (FIG. 9A) [19, 20]. This finding raised two possibilities: Firstly, clathrin is associated with contaminants rather than the VLPs, or secondly, that clathrin is indeed incorporated into VLPs but is exposed on the surface like HA and NA. To resolve this issue, immunogold labeling was used to look for the presence of clathrin on the surface of intact, undigested VLPs (FIG. 9B). This assay did show clathrin staining on the surface of intact VLPs, just as for HA and NA (FIGS. 6C and 6D). This new discover, i.e., clathrin, an endocytosis mediator protein, was specifically associated with secretory influenza VLPs, can have implications for the route of VLP biosynthesis and for the late stage of virus assembly and budding.

[0266] The proteomic analysis and protease protection assays of the secreted VLPs revealed that 22 cellular proteins associated with authentic virus of H1N1 were also specifically incorporated into mammalian H3N2 and H5N1 VLPs (Table 1 and FIG. 8). Among these proteins, tubulin, actin, annexins, enolase, GAPDH, gamma-glutamyltransferase, and HSP 27 have been demonstrated to be derived from lipid raft by proteomic analysis in previous studies [39,40,41,42, 43]. Correct assembly and budding of influenza virus requires cooperative action by multiple viral proteins with the lipid bilayers and genomic RNA as well as host proteins [44,45]. In VLP systems, except for the interactions involved in viral RNAs and capsid proteins, the events related to virus particle release are thought to be congruent to real virus assembly and release. A number of the proteins identified in mammalian influenza VLPs shed light on their roles in the influenza virion during assembly/budding stages of the infection process. The similarity between VLP proteomes and the virus cellular protein content also suggests the budding behaviors and constituents of mammalian VLPs are very similar to those of authentic viruses, and therefore that VLPs expressed from a mammalian cell-based system and constituting a non-pathogenic pseudovirion with very similar properties apart from genomic content are most likely to be a promising vaccine candidate.

[0267] Glycosylation Profiling of Influenza VLPs

[0268] Glycosylation of viral surface antigens is critical for immune recognition, receptor binding, inflammation, and pathogenicity, and therefore has a major influence on the efficacy of vaccine antigens [21,22]. For example, the common phenomenon of amino acid substitutions of the viral HA due to egg-adaptation and the consequent altered glycosylations severely affect the antigenicity of influenza virus [23]. As demonstrated by N-glycan footprinting analyses of HA, the use of different cell lines for replication of the same virus results in different N-glycosylation patterns on HA, which can be attributed to host-mediated changes in the amino acid sequence and potential glycosylation sites of HA, further influencing the antigenic properties of manufactured virus [24,25,26]. Therefore, the glycosylation status of HA and NA

antigens in the VLPs were assessed to look for any change that would affect the antigenicity and immune response of a VLP-based vaccine.

[0269] Deglycosylation assay of proteins in influenza VLPs were performed. Purified influenza VLPs equivalent to 10 μ g of proteins were denatured by heating at 100°C. for 10 min in the presence of 0.5% SDS and 40 mM DTT. Next, either PNGase F or Endo-H (New England Biolabs, Ipswich, Mass.) was added and the mixture incubated at 37°C. for 1 h with distinct reaction buffers, before protein gel electrophoresis and subsequent Western blot analyses.

[0270] The N-glycosylation patterns of H3N2- and H5N1-VLPs produced from Vero cells were compared to the glycan profiles of authentic viruses by performing deglycosylation assays with N-endoglycosidases PNGase F and Endo-H. PNGase F can remove all types of N-linked oligosaccharides from glycoproteins such as complex, hybrid, and high-mannose types, whereas Endo-H cleaves the chitobiose core of high-mannose and hybrid oligosaccharides from N-linked glycoproteins. As shown in FIG. 10A, most of the modified HA and NA (labeled as HA 1**+NA** and HA2**) in purified H5N1-VLP was seen as two major bands in the SDS-PAGE gel before deglycosylation (lane 1); their apparent molecular masses were around 56 and 30 kDa, respectively (FIGS. 10B and 10C, lanes 1). After treatment of H5N1-VLPs with PNGase F, HAL HA2, and NA bands increased their mobility to molecular masses of 40, 27, and 52 kDa, respectively (FIGS. 10A, 10B, and 10C, lanes 2), demonstrating that the two predominant viral surface antigens were mainly glycosylated by N-linked oligosaccharides. Of note, one form of NA marked as (π) in FIG. 10C whose mobility was not changed by reaction with both enzymes suggests NA may have other types of post-translational modification. When the glycosylated HA from H5N1-VLP was treated with Endo-H, the deglycosylation reaction was only partial, therefore the original bands and the Endo-H digested residue bands marked as (#) of HA1 and HA2 can be seen simultaneously in FIG. 10B, lane 6. However, the HA of H5N1 virus propagated in Vero cells was resistant to Endo-H digestion, suggesting the glycans linked to the viral HA are complex type (FIG. 10B, lane 8). The partial sensitivity of HA in H5N1-VLPs to Endo-H may be a result of hybrid glycan chains due to the overwhelming expression of viral protein thus incomplete glycosylation. However, as the great majority of NA can be deglycosylated by Endo-H, the NA proteins of H5N1-VLP may possess more high-mannose than complex glycans (FIG. 10C, lane 4). In parallel, the deglycosylation assays were performed on H3N2-VLPs (FIG. 10D). In H3 glycoprotein, Endo-H treatment deduced molecular mass 16 kDa and PNGase F treatment reduced 25 kDa, suggesting a higher content of high-mannose or hybrid types than complex type in the H3 glycan pool (FIG. 10D).

[0271] The results suggest that VLPs generated from Vero cells have similar glycosylation profiles to the authentic viruses that result from infection in the same host species. Collectively, the VLPs made from Vero cells resemble the real viruses in particle size, morphology, protein composition, and glycosylation profiles and therefore offer great potential as safe and effective influenza vaccine antigens.

Example 6

Virus Propagation, Hemagglutination and Serological Tests

[0272] Influenza virus, A/Taiwan/083/2006 and H5N1 (NI-BRG-14) strains (National Institute for Biological Standards

and Control, Potters Bar, U.K.) were propagated in Vero (for VLP comparison) or MDCK cells (for viral challenge). To assess hemagglutination, 3.5 μg of VLPs or virus and their serial 2-fold dilutions were mixed with a 0.75% suspension of guinea pig red blood cells in 96 well plates. Plates were incubated for 1 h and hemagglutination was assessed by eye. The highest dilution of VLPs or virus giving hemagglutination was determined as 1 HA unit.

[0273] To assess hemagglutination inhibition (HI) titers, sera were treated with a receptor-destroying enzyme and heat-inactivated (30 min, 56° C.), tested in 2-fold dilutions starting with an initial dilution of 1:10, then mixed with 8 HA units of H5N1-VLP and incubated at room temperature. After 1 h, a 0.75% suspension of guinea pig red blood cells was added and hemagglutination was assessed 2 h later by eye. HI titer as expressed as the reciprocal of the highest dilution that showed 50% inhibition of hemagglutination. All samples were tested in triplicate.

[0274] ELISA plates (Nunc) were coated with indicated H5 glycoprotein, VLPs, or virus at 4° C. overnight and blocked with 1% casein (Blocker Casein, Pierce, Rockford, Ill.) in PBS. ELISA plates were then incubated with serum samples of indicated dilution at 37° C. for 1 h, traced with HRP-conjugated secondary Ab, and developed color with TMB substrate (Pierce). They were washed with PBST five times between each step of ELISA. Finally, the ELISA was read out with absorbance of 450 nm wavelength (A_{450}) by a microplate reader (Power Wave XS, Bio-Tek) and results were plotted.

Example 7

Vaccination and Viral Challenge

[0275] Female BALB/c mice (6 weeks old) were purchased from National Laboratory Animal Center, randomly assigned to receive two doses of vaccine 21 days apart. Vaccines of 0.3 μg , 1.5 μg , 2.5 μg , or 10 μg H5N1-VLPs comprised of 0.054 μg , 0.27 μg , 0.45 μg , or 1.8 μg of H5 glycoprotein in sequence, and whole virus vaccine were grown in chicken embryo, inactivated by formalin and applied at 2.5 μg or 10 μg doses. Vaccines or PBS (as mock control) were given by intramuscular injection into the quadriceps. Blood was collected from mice via the retro-orbital sinus, transferred to a tube containing a serum separator and clot activator, and allowed to clot at room temperature. Sera were removed after centrifugation and stocked at -80° C. The immunized mice were challenged intranasally with a recombinant H5N1 virus, NIBRG-14, with a lethal dose (100-fold lethal dose to 50% of mice) as performed previously [48]. The mice were monitored daily for 14 days after the challenge for survival and morbidity (i.e. weight loss, inactivity and body temperature). All animal experiments were evaluated and approved by the Institutional Animal Care and Use Committee of Academia Sinica.

[0276] Humoral Immune Response of VLPs

[0277] To investigate the vaccine effect of mammalian expressed VLPs, mice were vaccinated with VLPs without adjuvantation. The vaccinated mice were analyzed for the antibody response and protection against viral infection. Mice (BALB/c; n=12) were vaccinated twice (day 0 and 21) via intramuscular injection with purified H5N1-VLP or inactivated whole virus of H5N1-pseudotyped vaccine strain (recombinant H5N1 engineered by reverse genetics) at two antigen doses (2.5 μg and 10 μg). Blood samples were collected to analyze humoral immune response before primary (day -1)

and after immunizations (day 14 and 35) (FIG. 11A). Sera were tested for influenza virus-specific IgG antibodies by ELISA against baculovirus produced H5 glycoprotein, or mammalian expressed H5N1-VLP and H3N2-VLP (FIG. 11B). Mice vaccinated with H5N1-VLP showed a robust response of IgG antibodies against H5 protein and H5N1-VLP. The ELISA titers of both antigen doses against H5 protein and H5N1-VLP were higher than 1:200,000, in contrast their titers against H3N2-VLP were insignificant (<1:25,000). This suggests that H5N1-VLP was highly immunogenic to stimulate highly specific antibodies against the H5 epitopes. Whole virus vaccine stimulated ELISA titers of approximate 1:100,000, while being highly immunogenic, the H5-specific titer was considerably lower than the VLP vaccine (FIG. 11B). However, ELISA titers for VLP vaccine group reduced to the level of whole virus vaccine group when whole virus was used as ELISA antigen (FIG. 11C). This suggests immunity of H5N1-VLP was more potent and specific to H5 glycoprotein, it may arise from the higher HA content in the VLPs than the viruses. Specificity of the VLP-induced antibody was further detected by Western blotting against all proteins of VLP and the virus. Indeed, the IgG antibodies detected only signals corresponding to the H5 glycoprotein, both HA1 and HA2 fragments in the VLP and virus (FIG. 11D). The lack of signal detecting other host and viral proteins in this experiment indicated the VLP was as "clean" as the inactivated split virus and subunit vaccines, only immunogenic toward the HA glycoprotein.

[0278] The IgG antibody isotypes distribution elicited by vaccination is indicative of the type of T cell immune response, as subsets of antigen-specific helper T cell via secreting different cytokines regulate the production of different IgG isotypes. The IgG1 isotype in mice is believed to signal a Th2 response, whereas the IgG2a isotype indicates more of a Th1 response. ELISA test was further used to measure the class and IgG isotypes of antigen-specific antibodies in response to the VLP and whole virus vaccines. As shown, the antibodies induced by VLP were predominantly IgG1 isotype, much less in IgG2a, and low or undetectable in IgG2b and IgA (FIG. 11E, left). However, the antibodies induced by whole virus vaccine were mainly IgG2a, less in IgG1, and insignificant in IgG2b and IgA. These results suggest that mammalian VLP vaccine at the two antigen doses induced primarily a Th2 response, whereas whole virus vaccine stimulated a mixed Th1/Th2 response with Th1 more dominant at higher antigen-dosage.

[0279] Vaccine Induced Hemagglutination-Inhibition (HI) Activity and Protection Against Viral Infection

[0280] The HI assay is the most widely accepted serological test for influenza immunity and is the gold standard measure of functional HA-specific antibodies after vaccination. The serological criteria currently used for approval of pandemic vaccines in the US are based on seasonal influenza vaccines, with seroconversion (i.e. a minimum 4-fold rise in HI titer) rate >40% and seroprotection (i.e. HI titer >1:40) rate \geq 70% in adults younger than 65. Antibodies elicited by each vaccine candidate were evaluated for ability to inhibit the VLP-induced agglutination of guinea pig red blood cells (FIG. 11E). After the second dose (day 35), the seroprotective HI titers were induced in 83.3% of mice received 2.5 μg and 10 μg VLP vaccine with mean HI titer reached about 1:60. When the antigen dose of VLP vaccine decreased to 0.3 μg and 1.5 μg , the reciprocal seroprotective rate dropped to

12.5% and 25%. Seroprotective rates of corresponding whole virus vaccine were 75% and 87.5% in parallel experiments.

[0281] All mice vaccinated with VLP vaccine, whole virus vaccine or mock control were challenged intranasally with predetermined lethal dose of H5N1-pseudotyped recombinant virus to evaluate the protective efficacy of each vaccine candidate. All mice received 2.5 µg and 10 µg VLP vaccine survived from the viral challenge, in contrast to the mock control mice that all died within 7 days after infection (FIG. 12A). Lower dose (0.3 µg and 1.5 µg) of VLP vaccine indeed compromised survival rate (50% and 25%, respectively), which is consistent with a lower seroprotection rate. Also, whole virus vaccine (2.5 µg and 10 µg) was protective to the viral challenge except one mouse in the 2.5 µg dose group lost >30% body weight (FIG. 12B). Body weight and temperature changes of test mice were indicative of illness and mice vaccinated with high-dose VLP and whole virus vaccine recovered their original weight by day 13 post-challenge, a result consistent with the survival outcome. However, mice vaccinated with low-dose VLP had more prominent weight loss and temperature decrease, despite some of the survivors recovered at later time.

[0282] Efficacy of the VLP created by mammalian expression system to be a new influenza vaccine for human and animal use is demonstrated for the first time in the present study. For example, it was demonstrated that vaccination with mammalian expressed VLPs provided full protection against lethal infection against the homologous strain challenge at doses as low as 2.5 µg VLP (0.45 µg HA) using two dose regimen in BALB/c mice.

[0283] The full protection of mammalian VLP vaccine was well-correlated with functional antibody responses (HI assay), which is the licensure criteria accepted for yearly inter-pandemic vaccines. The presence of numerous cellular proteins integrated in the mammalian expressed VLP may raise concerns of autoimmunity. It was shown that host protein contents of the VLP were akin to the authentic virus in varieties and quantity. Both Vero and Vero E6 cell lines are currently considered as the most widely acceptable cell substrate by regulatory authorities to produce a wide range of viruses for manufacturing human-use vaccine, including influenza, polio virus, rabies virus, smallpox, vesicular stomatitis virus, herpes simplex virus and rotavirus, etc. Furthermore, Vero cells are the only recommended cells to prepare viruses for vaccine production by reverse genetics in the document: "WHO guidance on development of influenza vaccine reference viruses by reverse genetics". For this reason, we chose Vero and Vero E6 cell lines rather than any other human or non-human mammalian cell lines to produce VLPs. In fact, our studies demonstrate that vaccination in mice with VLP void of adjuvant formulation elicited high-titer antibodies against HA only but not other proteins (FIG. 11). Vaccinated mice survived perfectly with two doses of VLP vaccines at 2.5 µg and 10 µg levels via intramuscular immunization, and they all survived the subsequent lethal viral challenge. No adverse effect was found before and after viral challenge throughout the 56-day experiment. Worth of note, the humoral immune response elicited by mammalian expressed VLPs is different from that of baculovirus-derived VLPs, suggesting a distinction between the two forms of VLPs. It may attribute to the glycosylation profile, host protein contents, the overall particle structure that present antigens, or something else.

[0284] It will be appreciated by those skilled in the art that changes could be made to the embodiments described above without departing from the broad inventive concept thereof. It is understood, therefore, that this invention is not limited to the particular embodiments disclosed, but it is intended to cover modifications within the spirit and scope of the present invention as defined by the appended claims.

REFERENCES

- [0285]** 1. Nicholson K G (2009) Influenza and vaccine development: a continued battle. *Expert Rev Vaccines* 8: 373-374.
- [0286]** 2. Sambhara S, Stephenson I (2009) Moving influenza vaccines forward. *Expert Rev Vaccines* 8: 375-377.
- [0287]** 3. Kang S M, Yoo D G, Lipatov A S, Song J M, Davis C T, et al. (2009) Induction of long-term protective immune responses by influenza H5N1 virus-like particles. *PLoS One* 4: e4667.
- [0288]** 4. Demicheli V, Rivetti D, Deeks J J, Jefferson T O (2004) Vaccines for preventing influenza in healthy adults. *Cochrane Database Syst Rev*: CD001269.
- [0289]** 5. Trollfors B (2006) General vaccination of children against influenza? *Acta Paediatr* 95: 774-777.
- [0290]** 6. Johansson B E, Brett I C (2007) Changing perspective on immunization against influenza. *Vaccine* 25: 3062-3065.
- [0291]** 7. Roy P, Noad R (2008) Virus-like particles as a vaccine delivery system: myths and facts. *Hum Vaccin* 4: 5-12.
- [0292]** 8. Garland S M, Hernandez-Avila M, Wheeler C M, Perez G, Harper D M, et al. (2007) Quadrivalent vaccine against human papillomavirus to prevent anogenital diseases. *N Engl J Med* 356: 1928-1943.
- [0293]** 9. Bright R A, Carter D M, Daniluk S, Toapanta F R, Ahmad A, et al. (2007) Influenza virus-like particles elicit broader immune responses than whole virion inactivated influenza virus or recombinant hemagglutinin. *Vaccine* 25: 3871-3878.
- [0294]** 10. Bright R A, Carter D M, Crevar C J, Toapanta F R, Steckbeck J D, et al. (2008) Cross-clade protective immune responses to influenza viruses with H5N1 HA and NA elicited by an influenza virus-like particle. *PLoS One* 3: e1501.
- [0295]** 11. Galarza J M, Latham T, Cupo A (2005) Virus-like particle vaccine conferred complete protection against a lethal influenza virus challenge. *Viral Immunol* 18: 365-372.
- [0296]** 12. Mahmood K, Bright R A, Mytle N, Carter D M, Crevar C J, et al. (2008) H5N1 VLP vaccine induced protection in ferrets against lethal challenge with highly pathogenic H5N1 influenza viruses. *Vaccine* 26: 5393-5399.
- [0297]** 13. Tao P, Luo M, Zhu D, Qu S, Yang Z, et al. (2009) Virus-like particle vaccine comprised of the HA, NA, and M1 proteins of an avian isolated H5N1 influenza virus induces protective immunity against homologous and heterologous strains in mice. *Viral Immunol* 22: 273-281.
- [0298]** 14. Quan F S, Huang C, Compans R W, Kang S M (2007) Virus-like particle vaccine induces protective immunity against homologous and heterologous strains of influenza virus. *J Virol* 81: 3514-3524.

- [0299] 15. Manolova V, Flace A, Bauer M, Schwarz K, Saudan P, et al. (2008) Nanoparticles target distinct dendritic cell populations according to their size. *Eur J Immunol* 38: 1404-1413.
- [0300] 16. Shaw ML, Stone KL, Colangelo CM, Gulcicek E E, Palese P (2008) Cellular proteins in influenza virus particles. *PLoS Pathog* 4: e1000085.
- [0301] 17. Ott D E, Coren L V, Kane B P, Busch L K, Johnson D G, et al. (1996) Cytoskeletal proteins inside human immunodeficiency virus type 1 virions. *J Virol* 70: 7734-7743.
- [0302] 18. Wubbolts R, Leckie R S, Veenhuizen P T, Schwarzmann G, Mobius W, et al. (2003) Proteomic and biochemical analyses of human B cell-derived exosomes. Potential implications for their function and multivesicular body formation. *J Biol Chem* 278: 10963-10972.
- [0303] 19. Kirchhausen T (2000) Three ways to make a vesicle. *Nat Rev Mol Cell Biol* 1: 187-198.
- [0304] 20. Brodsky F M, Chen C Y, Knuehl C, Towler M C, Wakeham D E (2001) Biological basket weaving: formation and function of clathrin-coated vesicles. *Annu Rev Cell Dev Biol* 17: 517-568.
- [0305] 21. Hamby S E, Hirst J D (2008) Prediction of glycosylation sites using random forests. *BMC Bioinformatics* 9: 500.
- [0306] 22. Schwarzer J, Rapp E, Reichl U (2008) N-glycan analysis by CGE-LIF: profiling influenza A virus hemagglutinin N-glycosylation during vaccine production. *Electrophoresis* 29: 4203-4214.
- [0307] 23. Robertson J S (1987) Sequence analysis of the haemagglutinin of A/Taiwan/1/86, a new variant of human influenza A(H1N1) virus. *J Gen Virol* 68 (Pt 4): 1205-1208.
- [0308] 24. Romanova J, Katinger D, Ferko B, Voglauer R, Mochalova L, et al. (2003) Distinct host range of influenza H3N2 virus isolates in Vero and MDCK cells is determined by cell specific glycosylation pattern. *Virology* 307: 90-97.
- [0309] 25. Schwarzer J, Rapp E, Hennig R, Genzel Y, Jordan I, et al. (2009) Glycan analysis in cell culture-based influenza vaccine production: influence of host cell line and virus strain on the glycosylation pattern of viral hemagglutinin. *Vaccine* 27: 4325-4336.
- [0310] 26. Skehel J J, Stevens D J, Daniels R S, Douglas A R, Knossow M, et al. (1984) A carbohydrate side chain on hemagglutinins of Hong Kong influenza viruses inhibits recognition by a monoclonal antibody. *Proc Natl Acad Sci USA* 81: 1779-1783.
- [0311] 27. Evans T G, Bonne W, Rose R C, Koenig S, Demeter L, et al. (2001) A Phase 1 study of a recombinant viruslike particle vaccine against human papillomavirus type 11 in healthy adult volunteers. *J Infect Dis* 183: 1485-1493.
- [0312] 28. Koutsky L A, Ault K A, Wheeler C M, Brown D R, Barr E, et al. (2002) A controlled trial of a human papillomavirus type 16 vaccine. *N Engl J Med* 347: 1645-1651.
- [0313] 29. Galarza J M, Latham T, Cupo A (2005) Virus-like particle (VLP) vaccine conferred complete protection against a lethal influenza virus challenge. *Viral Immunol* 18: 244-251.
- [0314] 30. Chen B J, Leser G P, Morita E, Lamb R A (2007) Influenza virus hemagglutinin and neuraminidase, but not the matrix protein, are required for assembly and budding of plasmid-derived virus-like particles. *J Virol* 81: 7111-7123.
- [0315] 31. Ali A, Avalos R T, Ponimaskin E, Nayak D P (2000) Influenza virus assembly: effect of influenza virus glycoproteins on the membrane association of M1 protein. *J Virol* 74: 8709-8719.
- [0316] 32. Enami M, Enami K (1996) Influenza virus hemagglutinin and neuraminidase glycoproteins stimulate the membrane association of the matrix protein. *J Virol* 70: 6653-6657.
- [0317] 33. Gregoriades A, Frangione B (1981) Insertion of influenza M protein into the viral lipid bilayer and localization of site of insertion. *J Virol* 40: 323-328.
- [0318] 34. Gomez-Puertas P, Albo C, Perez-Pastrana E, Vivo A, Portela A (2000) Influenza virus matrix protein is the major driving force in virus budding. *J Virol* 74: 11538-11547.
- [0319] 35. McCown M F, Pekosz A (2006) Distinct domains of the influenza A virus M2 protein cytoplasmic tail mediate binding to the M1 protein and facilitate infectious virus production. *Virology* 350: 8178-8189.
- [0320] 36. Ruigrok R W, Barge A, Durrer P, Brunner J, Ma K, et al. (2000) Membrane interaction of influenza virus M1 protein. *Virology* 267: 289-298.
- [0321] 37. Johansson B E, Bucher D J, Pokorny B A, Mikhail A, Kilbourne ED (1989) Identification of PR8 M1 protein in influenza virus high-yield reassortants by M1-specific monoclonal antibodies. *Virology* 171: 634-636.
- [0322] 38. McMurry JA, Johansson B E, De Groot A S (2008) A call to cellular & humoral arms: enlisting cognate T cell help to develop broad-spectrum vaccines against influenza A. *Hum Vaccin* 4: 148-157.
- [0323] 39. Blonder J, Terunuma A, Conrads T P, Chan K C, Yee C, et al. (2004) A proteomic characterization of the plasma membrane of human epidermis by high-throughput mass spectrometry. *J Invest Dermatol* 123: 691-699.
- [0324] 40. Foster L J, De Hoog C L, Mann M (2003) Unbiased quantitative proteomics of lipid rafts reveals high specificity for signaling factors. *Proc Natl Acad Sci USA* 100: 5813-5818.
- [0325] 41. Li N, Mak A, Richards D P, Naber C, Keller B O, et al. (2003) Monocyte lipid rafts contain proteins implicated in vesicular trafficking and phagosome formation. *Proteomics* 3: 536-548.
- [0326] 42. Li N, Shaw A R, Zhang N, Mak A, Li L (2004) Lipid raft proteomics: analysis of in-solution digest of sodium dodecyl sulfate-solubilized lipid raft proteins by liquid chromatography-matrix-assisted laser desorption/ionization tandem mass spectrometry. *Proteomics* 4: 3156-3166.
- [0327] 43. von Haller P D, Donohoe S, Goodlett D R, Aebersold R, Watts J D (2001) Mass spectrometric characterization of proteins extracted from Jurkat T cell detergent-resistant membrane domains. *Proteomics* 1: 1010-1021.
- [0328] 44. Scheiffele P, Rietveld A, Wilk T, Simons K (1999) Influenza viruses select ordered lipid domains during budding from the plasma membrane. *J Biol Chem* 274: 2038-2044.
- [0329] 45. Nayak D P, Hui E K, Barman S (2004) Assembly and budding of influenza virus. *Virus Res* 106: 147-165.
- [0330] 46. Jia N, Wang S X, Liu Y X, Zhang P H, Zuo S Q, et al. (2008) Increased sensitivity for detecting avian influenza-specific antibodies by a modified hemagglutination inhibition assay using horse erythrocytes. *J Virol Methods* 153: 43-48.
- [0331] 47. Gillim-Ross L, Subbarao K (2006) Emerging respiratory viruses: challenges and vaccine strategies. *Clin Microbiol Rev* 19: 614-636.
- [0332] 48. Wang C C, Chen J R, Tseng Y C, Hsu C H, Hung Y F, et al. (2009) Glycans on influenza hemagglutinin affect receptor binding and immune response. *Proc Natl Acad Sci USA* 106: 18137-18142.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1

<211> LENGTH: 3484

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: expression cassette

<400> SEQUENCE: 1

```

gttgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata    60
gccatataat ggagttccgc gttacataac ttacggtaaa tggcccgcct ggctgaccgc    120
ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt tcccatagta acgccaatag    180
ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccac ttggcagtac    240
atcaagtgtg tcatatgcca agtacgcccc ctattgacgt caatgacggg aaatggcccc    300
cctggcatta tgcccagtag atgaccttat gggactttcc tacttggcag tacatctacg    360
tattagtcac cgctattacc atggtgatgc ggttttgcca gtacatcaat gggcgtggat    420
agcggtttga ctcacgggga tttccaagtc tccaccccat tgacgtcaat gggagtttgt    480
tttggcacca aaatcaacgg gactttccaa aatgtcgtaa caactccgcc ccattgacgc    540
aaatggcgcg taggcgtgtg cgggtggagg tctatataag cagagctctc cctatcagtg    600
atagagatct ccctatcagt gatagagatc gtcgacgagc tcgtttagtg aaccgtcaga    660
tcgcctggag acgccatcca cgctgttttg acctccatag aagacaccgg gaccgatcca    720
gcctccggac tctagcgttt aaacttaagc ttggtaccga gctcggatcc cttgcagaag    780
ttggtcgtga ggcactgggc aggtaagtat caaggttaca agacaggttt aaggagacca    840
atagaaaactg ggcttgcga gacagagaag actcctgcgt ttctgatagg cacctattgg    900
tcttactgac atccactttg ctttctctc cacaggtgtc cactccagtg tcaattacag    960
ctcttaaggc tagagtactt aatacgactc actataggct agcatggcag acaacggtag    1020
tattaccggt gagagctta aacaactcct ggaacaatgg aacctagtaa taggtttcct    1080
attcctagcc tggattatgt tactacaatt tgccatttct aatcggaaca ggtttttgta    1140
cataataaag cttgttttcc tctggctctt gtggccagta acacttgctt gttttgtgct    1200
tgctgctgtc tacagaatta attgggtgac tggcgggatt gcgattgcaa tggcttgat    1260
tgtaggcttg atgtggctta gctacttcgt tgcttcttc aggctgtttg ctctgacctg    1320
ctcaatgtgg tcattcaacc cagaacaaa cattcttctc aatgtgcctc tccgggggac    1380
aattgtgacc agaccgctca tggaaagtga acttgcattt ggtgctgtga tcattcgtgg    1440
tcacttgcga atggccggac accccctagg gcgctgtgac attaaggacc tgccaaaaga    1500
gatcactgtg gctacatcac gaacgctttc ttattacaaa ttaggagcgt cgcagcgtgt    1560
aggcactgat tcaggttttg ctgcatacaa ccgctaccgt attggaaact ataaattaaa    1620
tacagaccac gccggtagca acgacaatat tgctttgcta gtacaggagc tcgtgagcaa    1680
ggcgaggag ctgttcaacc ggggtgggtcc catcctggtc gagctggacg ggcagctaaa    1740
cggccacaag ttcagcgtgt ccggcgaggg cgaggcgat gccacctacg gcaagctgac    1800
cctgaagttc atctgcacca ccggcaagct gcccggtgcc tggcccgcct tcgtgaccac    1860

```

-continued

```

cctgacctac ggcgtgcagt gcttgagccg ctaccccgac cacatgaagc agcaagcactt 1920
cttcaagtcc gccatgcccg aaggtctact ccaggagcgc accatcttct tcaaggacga 1980
cggcaactac aagaccgcg ccgagggtgaa gttcgagggc gacaccctgg tgaaccgcat 2040
cgagctgaag ggcctcgact tcaaggagga cggcaacatc ctggggcaca agctggagta 2100
caactacaac agccacaacg tctatatcat ggccgacaag cagaagaacg gcatcaaggt 2160
gaacttcaag atccgccaca acatcgagga cggcagcgtg cagctcgccg accactacca 2220
gcagaacacc cccatcgccg acggcccctg gctgctgccc gacaaccact acctgagcac 2280
ccagtcgcc ctgagcaaaag accccaacga gaagcgcgat cacatggtcc tctggagtt 2340
cgtgaccgcc gccgggatca ctctcgcat ggacgagctg tacaagtaag aattccgcc 2400
ctctccctcc ccccccccta acgttactgg ccgaagcgc ttggaataag gccggtgtgt 2460
gtttgtctat atgtgatttt ccaccatatt gccgtctttt ggcaatgtga gggcccggaa 2520
acctggccct gtcttcttga cgagcattcc taggggtctt tcccctctcg ccaaaggaat 2580
gcaaggtctg ttgaatgtcg tgaaggaagc agttctctg gaagctctt gaagacaaaac 2640
aacgtctgta ggcacccttt gcaggcagcg gaacccccca cctggcgaca ggtgcctctg 2700
cggccaaaag ccagctgat aagatacacc tgcaaggcg gcacaacccc agtgccacgt 2760
tgtgagttgg atagttgtgg aaagagtcaa atggctctcc tcaagcgtag tcaacaaggg 2820
gctgaaggat gccagaagg taccccattg tatgggaatc tgatctgggg cctcggtgca 2880
catgctttac atgtgtttag tcgaggttaa aaaaacgtct agggcccccg aaccacgggg 2940
acgtggtttt cctttgaaaa acacgatgat aatatggcca caaccggatc tatgtactca 3000
ttcgtttcgg aagaaacagg tacgttaata gttaatagcg tacttctttt tcttctttc 3060
gtggtattct tgctagtcac actagccatc cttactgcgc ttcgattgtg tgcgtactgc 3120
tgcaatattg ttaacgtgag tttagtaaaa ccaacggttt acgtctactc gcgtgttaaa 3180
aatctgaact cttctgaagg agttctctgat cttctggtct aatctagagg gcccgtttaa 3240
acccgctgat cagcctcgac tgtgcctct agttgccagc catctgtgt ttgccctcc 3300
cccgtgcctt ccttgacctt ggaaggtgcc actcccactg tcccttctta ataaaatgag 3360
gaaattgcat cgcattgtct gagtaggtgt cattctatc tggggggtgg ggtggggcag 3420
gacagcaagg gggaggattg ggaagacaat agcaggcatg ctggggatgc ggtgggctct 3480
atgg 3484

```

```

<210> SEQ ID NO 2
<211> LENGTH: 4837
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: expression cassette

```

```

<400> SEQUENCE: 2

```

```

gttgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata 60
gccatataat ggagttccgc gttacataaac ttacggtaaa tggcccgcct ggetgaccgc 120
ccaacgacct ccgccattg acgtcaataa tgacgtatgt tccatagta acgccaatag 180
ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccac ttggcagtac 240
atcaagtgta tcatatgcca agtacgcccc ctattgacgt caatgacggg aaatggcccc 300

```

-continued

cctggcatta	tgcccagtac	atgaccttat	gggactttcc	tacttggcag	tacatctaeg	360
tattagtcot	cgctattacc	atggatgatgc	ggttttggca	gtacatcaat	gggcgtggat	420
agcggtttga	ctcaecggga	tttccaagtc	tccaccccat	tgacgtcaat	gggagtttgt	480
tttggcacca	aaatcaacgg	gactttccaa	aatgtcgtaa	caactccgcc	ccattgacgc	540
aatggggcgg	taggcgtgta	cggtgggagg	tctatataag	cagagctctc	cctatcagtg	600
atagagatct	cctatcagtg	gatagagatc	gtcgcagcgc	tcgttttagtg	aaccgtcaga	660
tcgcctggag	acgccatcca	cgctgttttg	acctccatag	aagacaccgg	gaccgatcca	720
gcctccggac	tctagcgttt	aaacttaagc	ttggtaccga	gctcggatcc	atgtttattt	780
tcttattatt	tcttactctc	actagtggta	gtgaccttga	cgggtgcacc	acttttgatg	840
atgttcaagc	tctaattac	actcaacata	cttcatctat	gagggggggt	tactatcctg	900
atgaaatfff	tagatcagac	actctttatt	taactcagga	tttatttctt	ccattttatt	960
ctaagtgtac	agggtttcat	actattaatc	atcgttttgg	caaccctgtc	atacctttta	1020
aggatgggat	ttattttgct	gccacagaga	aatcaaatgt	tgtccgtggg	tgggtttttg	1080
gttctaccat	gaacaacaag	tcacagtcgg	tgattattat	taacaattct	actaatgttg	1140
ttatacagac	atgtaacttt	gaattgtgtg	acaacccttt	ctttgctggt	tctaaacceca	1200
tgggtacaca	gacacatact	atgatattcg	ataatgcatt	taattgcact	ttcgagtaca	1260
tatctgatgc	cttttcgctt	gatgtttcag	aaaagtcagg	taattttaaa	cacttacgag	1320
agtttgtgtt	taaaaataaa	gatgggtttc	tctatgttta	taagggetat	caacctatag	1380
atgtagtctg	tgatctacct	tctggtttta	acactttgaa	acctatffff	aagttgcctc	1440
ttggtattaa	cattacaaat	tttagagcca	ttcttacagc	cttttcacct	gctcaagaca	1500
tttggggcac	gtcagctgca	gcctatffff	ttggctatft	aaagccaact	acatttatgc	1560
tcaagtatga	tgaaaatggt	acaatcacag	atgctgttga	ttgttctcaa	aatccacttg	1620
ctgaactcaa	atgctctggt	aagagctttg	agattgacaa	aggaatttac	cagacctcta	1680
atttcagggt	tgttccctca	ggagatgttg	tgagattccc	taatattaca	aacttggtgc	1740
cttttgagga	ggtttttaat	gctactaaat	tcccttctgt	ctatgcatgg	gagagaaaaa	1800
aaatttctaa	ttgtgttgct	gattactctg	tgctctacaa	ctcaacatft	ttttcaacct	1860
ttaagtgcct	tggcgtttct	gccactaagt	tgaatgatct	ttgcttctcc	aatgtctatg	1920
cagattcttt	tgtagtcaag	ggagatgatg	taagacaaat	agcgcacagga	caaactggtg	1980
ttattgtctg	ttataattat	aaattgccag	atgatttcat	gggttgtgtc	cttgcttggga	2040
atactaggaa	cattgatgct	acttcaactg	gtaattataa	ttataaatat	aggtatctta	2100
gacatggcaa	gcttaggccc	tttgagagag	acatatctaa	tgtgcctttc	tcccctgatg	2160
gcaaaccttg	cacccacctc	gctcttaatt	gttattggcc	attaaatgat	tatggttttt	2220
acaccactac	tggcattggc	taccaacctt	acagagtgtg	agtactttct	tttgaacttt	2280
taaatgcacc	ggccacgggt	tgtggacca	aattatccac	tgaccttatt	aagaaccagt	2340
gtgtcaatft	taattttaat	ggactcactg	gtactgggtg	gttaactcct	tcttcaaaga	2400
gatttcaacc	atttcaacaa	tttggccgtg	atgtttctga	tttcaactgat	tccgttcgag	2460
atcctaaaac	atctgaaata	ttagacatft	caccttgctc	ttttgggggt	gtaagtgtaa	2520
ttacacctgg	aacaaatgct	tcactctgaag	ttgctgttct	atatcaagat	gttaactgca	2580

-continued

ctgatgttcc	tacagcaatt	catgcagatc	aactcacacc	agcttggegc	atatattcta	2640
ctggaacaaa	tgtattccag	actcaagcag	gctgtcttat	aggagctgag	catgtcgaca	2700
cttcttatga	gtgcgacatt	cctattggag	ctggcatttg	tgctagttae	catacagttt	2760
ctttattacg	tagtactagc	caaaaatcta	ttgtggctta	tactatgtct	ttaggtgctg	2820
atagttcaat	tgcttactct	aataacacca	ttgctatacc	tactaacttt	tcaattagca	2880
ttactacaga	agtaatgcct	gtttctatgg	ctaaaacctc	cgtagattgt	aatatgtaca	2940
tctgcggaga	ttctactgaa	tgtgctaatt	tgcttctcca	atatggtagc	ttttgcacac	3000
aactaaatcg	tgcaactctca	ggtattgctg	ctgaacagga	tcgcaacaca	cgtgaagtgt	3060
tcgctcaagt	caaacaaatg	tacaaaaacc	caactttgaa	atattttggt	ggttttaatt	3120
tttcacaaat	attacctgac	cctctaaagc	caactaagag	gtcttttatt	gaggacttgc	3180
tctttaataa	ggtgacactc	gctgatgctg	gcttcatgaa	gcaatatggc	gaatgcctag	3240
gtgatattaa	tgctagagat	ctcatttgctg	cgcagaagtt	caatggactt	acagtgttgc	3300
cacctctgct	cactgatgat	atgattgctg	cctacactgc	tgctctagtt	agtggtagctg	3360
ccactgctgg	atggacattt	ggtgctggcg	ctgctcttca	aatacctttt	gctatgcaaa	3420
tggcatatag	gttcaatggc	attggagttta	cccaaaatgt	tctctatgag	aaccaaaaaac	3480
aaatcgccaa	ccaatttaac	aaggcgatta	gtcaaatcca	agaatcactt	acaacaacat	3540
caactgcatt	gggcaagctg	caagacgttg	ttaaccagaa	tgctcaagca	ttaaacacac	3600
ttgttaaaaa	acttagctct	aattttgggtg	caatttcaag	tgtgctaaat	gatatccttt	3660
cgcgacttga	taaagtcgag	gocgaggtac	aaattgacag	gttaattaca	ggcagacttc	3720
aaagccttca	aacctatgta	acacaacaac	taatcagggc	tgctgaaatc	agggtctctg	3780
ctaactctgc	tgctactaaa	atgtctgagt	gtgttcttgg	acaatcaaaa	agagttgact	3840
tttgtggaaa	gggctaccac	cttatgtcct	tcccacaagc	agccccgcac	ggtgttctct	3900
tcctacatgt	cacgtatgtg	ccatcccagg	agaggaactt	caccacagcg	ccagcaattt	3960
gtcatgaagg	caaagcatac	ttccctcgtg	aagggttttt	tgtgtttaat	ggcacttctt	4020
ggtttattac	acagaggaac	ttcttttctc	cacaaataat	tactacagac	aatacatttg	4080
tctcagggaaa	ttgtgatgtc	gttattggca	tcattaacaa	cacagtttat	gatcctctgc	4140
aacctgagct	tgactcattc	aaagaagagc	tggaacaagta	cttcaaaaat	catacatcac	4200
cagatgttga	tcttgccgac	atttcaggca	ttaacgcttc	tgctgcaaac	attcaaaaag	4260
aaattgaccg	cctcaatgag	gtcgtcaaaa	atttaaatga	atcaactcatt	gaccttcaag	4320
aattgggaaa	atatgagcaa	tatattaaat	ggccttggtta	tgtttggctc	ggcttcattg	4380
ctggactaat	tgccatcgtc	atggttacaa	tcttgcttgg	ttgcatgact	agttgttgca	4440
gttgctctca	gggtgcatgc	tcttggtggt	cttgctgcaa	gtttgatgag	gatgactctg	4500
agccagttct	caaggtgtgc	aaattacatt	acacataaaa	gcttgcaatc	actagtgaat	4560
tcgcgccgcg	tcgagtctag	agggcccgtt	taaaccocgt	gatcagcctc	gactgtgctc	4620
tctagttgcc	agccatctgt	tgtttgcctc	tcccctgctc	cttccttgac	cctggaaggt	4680
gccactccca	ctgtccttcc	ctaataaaaat	gaggaaattg	catcgatttg	tctgagttagg	4740
tgctattcta	ttctgggggg	tggggtgggg	caggacagca	agggggagga	ttgggaagac	4800
aatagcaggc	atgctggggg	tgccgtgggg	tctatgg			4837

-continued

```

<210> SEQ ID NO 3
<211> LENGTH: 221
<212> TYPE: PRT
<213> ORGANISM: SARS coronavirus Urbani

<400> SEQUENCE: 3
Met Ala Asp Asn Gly Thr Ile Thr Val Glu Glu Leu Lys Gln Leu Leu
1      5      10      15
Glu Gln Trp Asn Leu Val Ile Gly Phe Leu Phe Leu Ala Trp Ile Met
20     25     30
Leu Leu Gln Phe Ala Tyr Ser Asn Arg Asn Arg Phe Leu Tyr Ile Ile
35     40     45
Lys Leu Val Phe Leu Trp Leu Leu Trp Pro Val Thr Leu Ala Cys Phe
50     55     60
Val Leu Ala Ala Val Tyr Arg Ile Asn Trp Val Thr Gly Gly Ile Ala
65     70     75     80
Ile Ala Met Ala Cys Ile Val Gly Leu Met Trp Leu Ser Tyr Phe Val
85     90     95
Ala Ser Phe Arg Leu Phe Ala Arg Thr Arg Ser Met Trp Ser Phe Asn
100    105    110
Pro Glu Thr Asn Ile Leu Leu Asn Val Pro Leu Arg Gly Thr Ile Val
115    120    125
Thr Arg Pro Leu Met Glu Ser Glu Leu Val Ile Gly Ala Val Ile Ile
130    135    140
Arg Gly His Leu Arg Met Ala Gly His Pro Leu Gly Arg Cys Asp Ile
145    150    155    160
Lys Asp Leu Pro Lys Glu Ile Thr Val Ala Thr Ser Arg Thr Leu Ser
165    170    175
Tyr Tyr Lys Leu Gly Ala Ser Gln Arg Val Gly Thr Asp Ser Gly Phe
180    185    190
Ala Ala Tyr Asn Arg Tyr Arg Ile Gly Asn Tyr Lys Leu Asn Thr Asp
195    200    205
His Ala Gly Ser Asn Asp Asn Ile Ala Leu Leu Val Gln
210    215    220

```

```

<210> SEQ ID NO 4
<211> LENGTH: 76
<212> TYPE: PRT
<213> ORGANISM: SARS coronavirus Urbani

<400> SEQUENCE: 4
Met Tyr Ser Phe Val Ser Glu Glu Thr Gly Thr Leu Ile Val Asn Ser
1      5      10      15
Val Leu Leu Phe Leu Ala Phe Val Val Phe Leu Leu Val Thr Leu Ala
20     25     30
Ile Leu Thr Ala Leu Arg Leu Cys Ala Tyr Cys Cys Asn Ile Val Asn
35     40     45
Val Ser Leu Val Lys Pro Thr Val Tyr Val Tyr Ser Arg Val Lys Asn
50     55     60
Leu Asn Ser Ser Glu Gly Val Pro Asp Leu Leu Val
65     70     75

```

```

<210> SEQ ID NO 5
<211> LENGTH: 1255

```

-continued

<212> TYPE: PRT

<213> ORGANISM: SARS coronavirus TW1

<400> SEQUENCE: 5

```

Met Phe Ile Phe Leu Leu Phe Leu Thr Leu Thr Ser Gly Ser Asp Leu
1      5      10      15
Asp Arg Cys Thr Thr Phe Asp Asp Val Gln Ala Pro Asn Tyr Thr Gln
20      25      30
His Thr Ser Ser Met Arg Gly Val Tyr Tyr Pro Asp Glu Ile Phe Arg
35      40      45
Ser Asp Thr Leu Tyr Leu Thr Gln Asp Leu Phe Leu Pro Phe Tyr Ser
50      55      60
Asn Val Thr Gly Phe His Thr Ile Asn His Thr Phe Gly Asn Pro Val
65      70      75      80
Ile Pro Phe Lys Asp Gly Ile Tyr Phe Ala Ala Thr Glu Lys Ser Asn
85      90      95
Val Val Arg Gly Trp Val Phe Gly Ser Thr Met Asn Asn Lys Ser Gln
100     105     110
Ser Val Ile Ile Ile Asn Asn Ser Thr Asn Val Val Ile Arg Ala Cys
115     120     125
Asn Phe Glu Leu Cys Asp Asn Pro Phe Phe Ala Val Ser Lys Pro Met
130     135     140
Gly Thr Gln Thr His Thr Met Ile Phe Asp Asn Ala Phe Asn Cys Thr
145     150     155     160
Phe Glu Tyr Ile Ser Asp Ala Phe Ser Leu Asp Val Ser Glu Lys Ser
165     170     175
Gly Asn Phe Lys His Leu Arg Glu Phe Val Phe Lys Asn Lys Asp Gly
180     185     190
Phe Leu Tyr Val Tyr Lys Gly Tyr Gln Pro Ile Asp Val Val Arg Asp
195     200     205
Leu Pro Ser Gly Phe Asn Thr Leu Lys Pro Ile Phe Lys Leu Pro Leu
210     215     220
Gly Ile Asn Ile Thr Asn Phe Arg Ala Ile Leu Thr Ala Phe Ser Pro
225     230     235     240
Ala Gln Asp Ile Trp Gly Thr Ser Ala Ala Ala Tyr Phe Val Gly Tyr
245     250     255
Leu Lys Pro Thr Thr Phe Met Leu Lys Tyr Asp Glu Asn Gly Thr Ile
260     265     270
Thr Asp Ala Val Asp Cys Ser Gln Asn Pro Leu Ala Glu Leu Lys Cys
275     280     285
Ser Val Lys Ser Phe Glu Ile Asp Lys Gly Ile Tyr Gln Thr Ser Asn
290     295     300
Phe Arg Val Val Pro Ser Gly Asp Val Val Arg Phe Pro Asn Ile Thr
305     310     315     320
Asn Leu Cys Pro Phe Gly Glu Val Phe Asn Ala Thr Lys Phe Pro Ser
325     330     335
Val Tyr Ala Trp Glu Arg Lys Lys Ile Ser Asn Cys Val Ala Asp Tyr
340     345     350
Ser Val Leu Tyr Asn Ser Thr Phe Phe Ser Thr Phe Lys Cys Tyr Gly
355     360     365
Val Ser Ala Thr Lys Leu Asn Asp Leu Cys Phe Ser Asn Val Tyr Ala
370     375     380

```

-continued

Asp Ser Phe Val Val Lys Gly Asp Asp Val Arg Gln Ile Ala Pro Gly
 385 390 395 400
 Gln Thr Gly Val Ile Ala Asp Tyr Asn Tyr Lys Leu Pro Asp Asp Phe
 405 410 415
 Met Gly Cys Val Leu Ala Trp Asn Thr Arg Asn Ile Asp Ala Thr Ser
 420 425 430
 Thr Gly Asn Tyr Asn Tyr Lys Tyr Arg Tyr Leu Arg His Gly Lys Leu
 435 440 445
 Arg Pro Phe Glu Arg Asp Ile Ser Asn Val Pro Phe Ser Pro Asp Gly
 450 455 460
 Lys Pro Cys Thr Pro Pro Ala Leu Asn Cys Tyr Trp Pro Leu Asn Asp
 465 470 475 480
 Tyr Gly Phe Tyr Thr Thr Thr Gly Ile Gly Tyr Gln Pro Tyr Arg Val
 485 490 495
 Val Val Leu Ser Phe Glu Leu Leu Asn Ala Pro Ala Thr Val Cys Gly
 500 505 510
 Pro Lys Leu Ser Thr Asp Leu Ile Lys Asn Gln Cys Val Asn Phe Asn
 515 520 525
 Phe Asn Gly Leu Thr Gly Thr Gly Val Leu Thr Pro Ser Ser Lys Arg
 530 535 540
 Phe Gln Pro Phe Gln Gln Phe Gly Arg Asp Val Ser Asp Phe Thr Asp
 545 550 555 560
 Ser Val Arg Asp Pro Lys Thr Ser Glu Ile Leu Asp Ile Ser Pro Cys
 565 570 575
 Ser Phe Gly Gly Val Ser Val Ile Thr Pro Gly Thr Asn Ala Ser Ser
 580 585 590
 Glu Val Ala Val Leu Tyr Gln Asp Val Asn Cys Thr Asp Val Ser Thr
 595 600 605
 Ala Ile His Ala Asp Gln Leu Thr Pro Ala Trp Arg Ile Tyr Ser Thr
 610 615 620
 Gly Asn Asn Val Phe Gln Thr Gln Ala Gly Cys Leu Ile Gly Ala Glu
 625 630 635 640
 His Val Asp Thr Ser Tyr Glu Cys Asp Ile Pro Ile Gly Ala Gly Ile
 645 650 655
 Cys Ala Ser Tyr His Thr Val Ser Leu Leu Arg Ser Thr Ser Gln Lys
 660 665 670
 Ser Ile Val Ala Tyr Thr Met Ser Leu Gly Ala Asp Ser Ser Ile Ala
 675 680 685
 Tyr Ser Asn Asn Thr Ile Ala Ile Pro Thr Asn Phe Ser Ile Ser Ile
 690 695 700
 Thr Thr Glu Val Met Pro Val Ser Met Ala Lys Thr Ser Val Asp Cys
 705 710 715 720
 Asn Met Tyr Ile Cys Gly Asp Ser Thr Glu Cys Ala Asn Leu Leu Leu
 725 730 735
 Gln Tyr Gly Ser Phe Cys Thr Gln Leu Asn Arg Ala Leu Ser Gly Ile
 740 745 750
 Ala Ala Glu Gln Asp Arg Asn Thr Arg Glu Val Phe Ala Gln Val Lys
 755 760 765
 Gln Met Tyr Lys Thr Pro Thr Leu Lys Tyr Phe Gly Gly Phe Asn Phe
 770 775 780

-continued

Ser Gln Ile Leu Pro Asp Pro Leu Lys Pro Thr Lys Arg Ser Phe Ile
 785 790 795 800
 Glu Asp Leu Leu Phe Asn Lys Val Thr Leu Ala Asp Ala Gly Phe Met
 805 810 815
 Lys Gln Tyr Gly Glu Cys Leu Gly Asp Ile Asn Ala Arg Asp Leu Ile
 820 825 830
 Cys Ala Gln Lys Phe Asn Gly Leu Thr Val Leu Pro Pro Leu Leu Thr
 835 840 845
 Asp Asp Met Ile Ala Ala Tyr Thr Ala Ala Leu Val Ser Gly Thr Ala
 850 855 860
 Thr Ala Gly Trp Thr Phe Gly Ala Gly Ala Ala Leu Gln Ile Pro Phe
 865 870 875 880
 Ala Met Gln Met Ala Tyr Arg Phe Asn Gly Ile Gly Val Thr Gln Asn
 885 890 895
 Val Leu Tyr Glu Asn Gln Lys Gln Ile Ala Asn Gln Phe Asn Lys Ala
 900 905 910
 Ile Ser Gln Ile Gln Glu Ser Leu Thr Thr Thr Ser Thr Ala Leu Gly
 915 920 925
 Lys Leu Gln Asp Val Val Asn Gln Asn Ala Gln Ala Leu Asn Thr Leu
 930 935 940
 Val Lys Gln Leu Ser Ser Asn Phe Gly Ala Ile Ser Ser Val Leu Asn
 945 950 955 960
 Asp Ile Leu Ser Arg Leu Asp Lys Val Glu Ala Glu Val Gln Ile Asp
 965 970 975
 Arg Leu Ile Thr Gly Arg Leu Gln Ser Leu Gln Thr Tyr Val Thr Gln
 980 985 990
 Gln Leu Ile Arg Ala Ala Glu Ile Arg Ala Ser Ala Asn Leu Ala Ala
 995 1000 1005
 Thr Lys Met Ser Glu Cys Val Leu Gly Gln Ser Lys Arg Val Asp
 1010 1015 1020
 Phe Cys Gly Lys Gly Tyr His Leu Met Ser Phe Pro Gln Ala Ala
 1025 1030 1035
 Pro His Gly Val Val Phe Leu His Val Thr Tyr Val Pro Ser Gln
 1040 1045 1050
 Glu Arg Asn Phe Thr Thr Ala Pro Ala Ile Cys His Glu Gly Lys
 1055 1060 1065
 Ala Tyr Phe Pro Arg Glu Gly Val Phe Val Phe Asn Gly Thr Ser
 1070 1075 1080
 Trp Phe Ile Thr Gln Arg Asn Phe Phe Ser Pro Gln Ile Ile Thr
 1085 1090 1095
 Thr Asp Asn Thr Phe Val Ser Gly Asn Cys Asp Val Val Ile Gly
 1100 1105 1110
 Ile Ile Asn Asn Thr Val Tyr Asp Pro Leu Gln Pro Glu Leu Asp
 1115 1120 1125
 Ser Phe Lys Glu Glu Leu Asp Lys Tyr Phe Lys Asn His Thr Ser
 1130 1135 1140
 Pro Asp Val Asp Leu Gly Asp Ile Ser Gly Ile Asn Ala Ser Val
 1145 1150 1155
 Val Asn Ile Gln Lys Glu Ile Asp Arg Leu Asn Glu Val Ala Lys
 1160 1165 1170
 Asn Leu Asn Glu Ser Leu Ile Asp Leu Gln Glu Leu Gly Lys Tyr

-continued

1175	1180	1185
Glu Gln Tyr Ile Lys Trp	Pro Trp Tyr Val Trp	Leu Gly Phe Ile
1190	1195	1200
Ala Gly Leu Ile Ala Ile	Val Met Val Thr Ile	Leu Leu Cys Cys
1205	1210	1215
Met Thr Ser Cys Cys Ser	Cys Leu Lys Gly Ala	Cys Ser Cys Gly
1220	1225	1230
Ser Cys Cys Lys Phe Asp	Glu Asp Asp Ser Glu	Pro Val Leu Lys
1235	1240	1245
Gly Val Lys Leu His Tyr	Thr	
1250	1255	

<210> SEQ ID NO 6
 <211> LENGTH: 252
 <212> TYPE: PRT
 <213> ORGANISM: influenza A/Taiwan/083/2006

<400> SEQUENCE: 6

Met Ser Leu Leu Thr	Glu Val Glu Thr Tyr Val	Leu Ser Ile Val Pro
1	5	10 15
Ser Gly Pro Leu Lys	Ala Glu Ile Ala Gln Arg	Leu Glu Asp Val Phe
20	25	30
Ala Gly Lys Asn Thr	Asp Leu Glu Ala Leu Met	Glu Trp Leu Lys Thr
35	40	45
Arg Pro Ile Leu Ser	Pro Leu Thr Lys Gly Ile	Leu Gly Phe Val Phe
50	55	60
Thr Leu Thr Val Pro	Ser Glu Arg Gly Leu Gln	Arg Arg Arg Phe Val
65	70	75 80
Gln Asn Ala Leu Asn	Gly Asn Gly Asp Pro Asn	Met Asp Lys Ala
85	90	95
Val Lys Leu Tyr Arg	Lys Leu Lys Arg Glu Ile Thr	Phe His Gly Ala
100	105	110
Lys Glu Ile Ala Leu	Ser Tyr Ser Ala Gly Ala	Leu Ala Ser Cys Met
115	120	125
Gly Leu Ile Tyr Asn	Arg Met Gly Ala Val Thr	Thr Glu Val Ala Phe
130	135	140
Gly Leu Val Cys Ala	Thr Cys Glu Gln Ile Ala	Asp Ser Gln His Arg
145	150	155 160
Ser His Arg Gln Met	Val Ala Thr Thr Asn Pro	Leu Ile Arg His Glu
165	170	175
Asn Arg Met Val Leu	Ala Ser Thr Thr Ala Lys	Ala Met Glu Gln Met
180	185	190
Ala Gly Ser Ser Glu	Gln Ala Ala Glu Ala Met	Glu Ile Ala Ser Gln
195	200	205
Ala Arg Gln Met Val	Gln Ala Met Arg Ala Ile	Gly Thr His Pro Ser
210	215	220
Ser Ser Thr Gly Leu	Arg Asp Asp Leu Leu Glu	Asn Leu Gln Thr Tyr
225	230	235 240
Gln Lys Arg Met Gly	Val Gln Met Gln Arg Phe	Lys
245	250	

<210> SEQ ID NO 7
 <211> LENGTH: 97

-continued

```

<212> TYPE: PRT
<213> ORGANISM: influenza A/Taiwan/083/2006

<400> SEQUENCE: 7

Met Ser Leu Leu Thr Glu Val Glu Thr Pro Ile Arg Asn Glu Trp Gly
1          5          10          15
Cys Arg Cys Asn Asp Ser Ser Asp Pro Leu Val Val Ala Ala Asn Ile
20          25          30
Ile Gly Ile Leu His Leu Ile Leu Trp Ile Leu Asp Arg Leu Phe Phe
35          40          45
Lys Cys Val Tyr Arg Leu Phe Lys His Gly Leu Lys Arg Gly Pro Ser
50          55          60
Thr Glu Gly Val Pro Glu Ser Met Arg Glu Glu Tyr Arg Lys Glu Gln
65          70          75          80
Gln Asn Ala Val Asp Ala Asp Asp Ser His Phe Val Ser Ile Glu Leu
85          90          95

Glu

```

```

<210> SEQ ID NO 8
<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: influenza A/Taiwan/083/2006

<400> SEQUENCE: 8

Met Lys Thr Ile Ile Ala Leu Ser Tyr Ile Leu Cys Leu Val Phe Ala
1          5          10          15
Gln Lys Leu Pro Gly Asn Asp Asn Ser Thr Ala Thr Leu Cys Leu Gly
20          25          30
His His Ala Val Pro Asn Gly Thr Ile Val Lys Thr Ile Thr Asn Asp
35          40          45
Gln Ile Glu Val Thr Asn Ala Thr Glu Leu Val Gln Ser Ser Ser Thr
50          55          60
Gly Gly Ile Cys Asp Ser Pro His Gln Ile Leu Asp Gly Glu Asn Cys
65          70          75          80
Thr Leu Ile Asp Ala Leu Leu Gly Asp Pro Gln Cys Asp Gly Phe Gln
85          90          95
Asn Lys Lys Trp Asp Leu Phe Val Glu Arg Ser Lys Ala Tyr Ser Asn
100         105         110
Cys Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ser Leu Arg Ser Leu Val
115         120         125
Ala Ser Ser Gly Thr Leu Glu Phe Asn Asn Glu Ser Phe Asn Trp Thr
130         135         140
Gly Val Thr Gln Asn Gly Thr Ser Ser Ala Cys Lys Arg Arg Ser Asn
145         150         155         160
Asn Ser Phe Phe Ser Arg Leu Asn Trp Leu Thr His Leu Lys Phe Lys
165         170         175
Tyr Pro Ala Leu Asn Val Thr Met Pro Asn Asn Glu Lys Phe Asp Lys
180         185         190
Leu Tyr Ile Trp Gly Val His His Pro Gly Thr Asp Asn Asp Gln Ile
195         200         205
Phe Leu Tyr Ala Gln Ala Ser Gly Arg Ile Thr Val Ser Thr Lys Arg
210         215         220
Ser Gln Gln Thr Val Ile Pro Asn Ile Gly Ser Arg Pro Arg Val Arg

```

-continued

```

225                230                235                240
Asp Ile Pro Ser Arg Ile Ser Ile Tyr Trp Thr Ile Val Lys Pro Gly
                245                250                255
Asp Ile Leu Leu Ile Asn Ser Thr Gly Asn Leu Ile Ala Pro Arg Gly
                260                265                270
Tyr Phe Lys Ile Arg Ser Gly Lys Ser Ser Ile Met Arg Ser Asp Ala
                275                280                285
Pro Ile Gly Lys Cys Asn Ser Glu Cys Ile Thr Pro Asn Gly Ser Ile
290                295                300
Pro Asn Asp Lys Pro Phe Gln Asn Val Asn Arg Ile Thr Tyr Gly Ala
305                310                315
Cys Pro Arg Tyr Val Lys Gln Asn Thr Leu Lys Leu Ala Thr Gly Met
                325                330                335
Arg Asn Val Pro Glu Lys Gln Thr Arg Gly Ile Phe Gly Ala Ile Ala
                340                345                350
Gly Phe Ile Glu Asn Gly Trp Glu Gly Met Val Asp Gly Trp Tyr Gly
                355                360                365
Phe Arg His Gln Asn Ser Glu Gly Ile Gly Gln Ala Ala Asp Leu Lys
370                375                380
Ser Thr Gln Ala Ala Ile Asn Gln Ile Asn Gly Lys Leu Asn Arg Leu
385                390                395
Ile Gly Lys Thr Asn Glu Lys Phe His Gln Ile Glu Lys Glu Phe Ser
                405                410                415
Glu Val Glu Gly Arg Ile Gln Asp Leu Glu Lys Tyr Val Glu Asp Thr
                420                425                430
Lys Ile Asp Leu Trp Ser Tyr Asn Ala Glu Leu Leu Val Ala Leu Glu
                435                440                445
Asn Gln His Thr Ile Asp Leu Thr Asp Ser Glu Met Asn Lys Leu Phe
450                455                460
Glu Arg Thr Lys Lys Gln Leu Arg Glu Asn Ala Glu Asp Met Gly Asn
465                470                475                480
Gly Cys Phe Lys Ile Tyr His Lys Cys Asp Asn Ala Cys Ile Gly Ser
                485                490                495
Ile Arg Asn Gly Thr Tyr Asp His Asp Val Tyr Arg Asp Glu Ala Leu
500                505                510
Asn Asn Arg Phe Gln Ile Lys Gly Val Glu Leu Lys Ser Gly Tyr Lys
515                520                525
Asp Trp Ile Leu Trp Ile Ser Phe Ala Ile Ser Cys Phe Leu Leu Cys
530                535                540
Val Ala Leu Leu Gly Phe Ile Met Trp Ala Cys Gln Lys Gly Asn Ile
545                550                555                560
Arg Cys Asn Ile Cys Ile
565

```

```

<210> SEQ ID NO 9
<211> LENGTH: 469
<212> TYPE: PRT
<213> ORGANISM: influenza A/Taiwan/083/2006
<400> SEQUENCE: 9

```

```

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Val Ser Leu Thr
1                5                10                15

```

-continued

Ile	Ser	Thr	Ile	Cys	Phe	Phe	Met	Gln	Ile	Ala	Ile	Leu	Ile	Thr	Thr
			20					25				30			
Val	Thr	Leu	His	Phe	Lys	Gln	Tyr	Glu	Phe	Asn	Ser	Pro	Pro	Asn	Asn
		35					40					45			
Gln	Val	Met	Leu	Cys	Glu	Pro	Thr	Ile	Ile	Glu	Arg	Asn	Ile	Thr	Glu
		50				55					60				
Ile	Val	Tyr	Leu	Thr	Asn	Thr	Thr	Ile	Glu	Lys	Glu	Ile	Cys	Pro	Lys
65					70					75					80
Leu	Ala	Glu	Tyr	Arg	Asn	Trp	Ser	Lys	Pro	Gln	Cys	Asn	Ile	Thr	Gly
				85					90					95	
Phe	Ala	Pro	Phe	Ser	Lys	Asp	Asn	Ser	Ile	Arg	Leu	Ser	Ala	Gly	Gly
			100					105					110		
Asp	Ile	Trp	Val	Thr	Arg	Glu	Pro	Tyr	Val	Ser	Cys	Asp	Pro	Asp	Lys
		115					120					125			
Cys	Tyr	Gln	Phe	Ala	Leu	Gly	Gln	Gly	Thr	Thr	Leu	Asn	Asn	Val	His
		130				135						140			
Ser	Asn	Asp	Thr	Val	His	Asp	Arg	Thr	Pro	Tyr	Arg	Thr	Leu	Leu	Met
145					150					155					160
Asn	Glu	Leu	Gly	Val	Pro	Phe	His	Leu	Gly	Thr	Lys	Gln	Val	Cys	Ile
			165					170						175	
Ala	Trp	Ser	Ser	Ser	Ser	Cys	His	Asp	Gly	Lys	Ala	Trp	Leu	His	Val
			180					185					190		
Cys	Val	Thr	Gly	Asp	Asp	Lys	Asn	Ala	Thr	Ala	Ser	Phe	Ile	Tyr	Asn
		195					200					205			
Gly	Arg	Leu	Val	Asp	Ser	Ile	Val	Ser	Trp	Ser	Lys	Glu	Ile	Leu	Arg
	210					215					220				
Thr	Gln	Glu	Ser	Glu	Cys	Val	Cys	Ile	Asn	Gly	Thr	Cys	Thr	Val	Val
225					230					235					240
Met	Thr	Asp	Gly	Ser	Ala	Ser	Gly	Lys	Ala	Asp	Thr	Lys	Ile	Leu	Phe
			245						250					255	
Ile	Glu	Glu	Gly	Lys	Ile	Val	His	Thr	Ser	Thr	Leu	Ser	Gly	Ser	Ala
			260					265						270	
Gln	His	Val	Glu	Glu	Cys	Ser	Cys	Tyr	Pro	Arg	Tyr	Pro	Gly	Val	Arg
		275					280					285			
Cys	Val	Cys	Arg	Asp	Asn	Trp	Lys	Gly	Ser	Asn	Arg	Pro	Ile	Val	Asp
	290					295					300				
Ile	Asn	Ile	Lys	Asp	Tyr	Ser	Ile	Val	Ser	Ser	Tyr	Val	Cys	Ser	Gly
305					310					315					320
Leu	Val	Gly	Asp	Thr	Pro	Arg	Lys	Asn	Asp	Ser	Ser	Ser	Ser	Gly	His
			325						330					335	
Cys	Leu	Asp	Pro	Asn	Asn	Glu	Glu	Gly	Gly	His	Gly	Val	Lys	Gly	Trp
			340					345					350		
Ala	Phe	Asp	Asp	Gly	Asn	Asp	Val	Trp	Met	Gly	Arg	Thr	Ile	Ser	Glu
		355					360					365			
Lys	Leu	Arg	Ser	Gly	Tyr	Glu	Thr	Phe	Lys	Val	Ile	Glu	Gly	Trp	Ser
	370					375					380				
Asn	Pro	Asn	Ser	Lys	Leu	Gln	Ile	Asn	Arg	Gln	Val	Ile	Val	Asp	Arg
385					390					395					400
Gly	Asn	Arg	Ser	Gly	Tyr	Ser	Gly	Ile	Phe	Ser	Val	Glu	Gly	Lys	Ser
			405						410					415	
Cys	Ile	Asn	Arg	Cys	Phe	Tyr	Val	Glu	Leu	Ile	Arg	Gly	Arg	Lys	Glu

-continued

```

          420              425              430
Glu Thr Glu Val Leu Trp Thr Ser Asn Ser Ile Val Val Phe Cys Gly
      435              440              445

Thr Ser Gly Thr Tyr Gly Thr Gly Ser Trp Pro Asp Gly Ala Asp Ile
      450              455              460

Asn Leu Met Pro Ile
465

<210> SEQ ID NO 10
<211> LENGTH: 567
<212> TYPE: PRT
<213> ORGANISM: influenza A/Hanoi/30408/2005(H5N1)

<400> SEQUENCE: 10

Met Glu Lys Ile Val Leu Leu Phe Ala Ile Val Ser Leu Val Lys Ser
1      5      10      15
Asp Gln Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Glu Gln Val
20     25     30
Asp Thr Ile Met Glu Lys Asn Val Thr Val Thr His Ala Gln Asp Ile
35     40     45
Leu Glu Lys Thr His Asn Gly Lys Leu Cys Asp Leu Asp Gly Val Lys
50     55     60
Pro Leu Ile Leu Arg Asp Cys Ser Val Ala Gly Trp Leu Leu Gly Asn
65     70     75     80
Pro Met Cys Asp Glu Phe Ile Asn Val Pro Glu Trp Ser Tyr Ile Val
85     90     95
Glu Lys Ala Asn Pro Val Asn Asp Leu Cys Tyr Pro Gly Asp Phe Asn
100    105    110
Asp Tyr Glu Glu Leu Lys His Leu Leu Ser Arg Ile Asn His Phe Glu
115    120    125
Lys Ile Gln Ile Ile Pro Lys Ser Ser Trp Leu Ser His Glu Ala Ser
130    135    140
Leu Gly Val Ser Ser Ala Cys Pro Tyr Gln Gly Lys Ser Ser Phe Phe
145    150    155    160
Arg Asn Val Val Trp Leu Ile Lys Lys Asn Ser Thr Tyr Pro Thr Ile
165    170    175
Lys Arg Ser Tyr Asn Asn Thr Asn Gln Glu Asp Leu Leu Val Leu Trp
180    185    190
Gly Ile His His Pro Asn Asp Ala Ala Glu Gln Thr Lys Leu Tyr Gln
195    200    205
Asn Pro Thr Thr Tyr Ile Ser Val Gly Thr Ser Thr Leu Asn Gln Arg
210    215    220
Leu Val Pro Arg Ile Ala Thr Arg Ser Lys Val Asn Gly Gln Ser Gly
225    230    235    240
Arg Met Glu Phe Phe Trp Thr Ile Leu Lys Pro Asn Asp Ala Ile Asn
245    250    255
Phe Glu Ser Asn Gly Asn Phe Ile Ala Pro Glu Tyr Ala Tyr Lys Ile
260    265    270
Val Lys Lys Gly Asp Ser Thr Ile Met Lys Ser Glu Leu Glu Tyr Gly
275    280    285
Asn Cys Asn Thr Lys Cys Gln Thr Pro Met Gly Ala Ile Asn Ser Ser
290    295    300

```


-continued

Ile Arg Glu Pro Phe Ile Ser Cys Ser His Leu Glu Cys Arg Thr Phe
100 105 110

Phe Leu Thr Gln Gly Ala Leu Leu Asn Asp Lys His Ser Asn Gly Thr
115 120 125

Val Lys Asp Arg Ser Pro His Arg Thr Leu Met Ser Cys Pro Val Gly
130 135 140

Glu Ala Pro Ser Pro Tyr Asn Ser Arg Phe Glu Ser Val Ala Trp Ser
145 150 155 160

Ala Ser Ala Cys His Asp Gly Thr Ser Trp Leu Thr Ile Gly Ile Ser
165 170 175

Gly Pro Asp Asn Gly Ala Val Ala Val Val Lys Tyr Asn Gly Ile Ile
180 185 190

Thr Asp Thr Ile Lys Ser Trp Arg Asn Asn Ile Leu Arg Thr Gln Glu
195 200 205

Ser Glu Cys Ala Cys Val Asn Gly Ser Cys Phe Thr Val Met Thr Asp
210 215 220

Gly Pro Ser Asn Gly Gln Ala Ser His Lys Ile Phe Lys Met Glu Lys
225 230 235 240

Gly Lys Val Val Lys Ser Val Glu Leu Asp Ala Pro Asn Tyr His Tyr
245 250 255

Glu Glu Cys Ser Cys Tyr Pro Asp Ala Gly Glu Ile Thr Cys Val Cys
260 265 270

Arg Asp Asn Trp His Gly Ser Asn Arg Pro Trp Val Ser Phe Asn Gln
275 280 285

Asn Leu Glu Tyr Gln Ile Gly Tyr Ile Cys Ser Gly Val Phe Gly Asp
290 295 300

Asn Pro Arg Pro Asn Asp Gly Thr Gly Ser Cys Gly Pro Val Ser Ser
305 310 315 320

Asn Gly Ala Tyr Gly Val Lys Gly Phe Ser Phe Lys Tyr Gly Asn Gly
325 330 335

Val Trp Ile Gly Arg Thr Lys Ser Thr Asn Ser Arg Ser Gly Phe Glu
340 345 350

Met Ile Trp Asp Pro Asn Gly Trp Thr Glu Thr Asp Ser Ser Phe Ser
355 360 365

Val Lys Gln Asp Ile Val Ala Ile Thr Asp Trp Ser Gly Tyr Ser Gly
370 375 380

Ser Phe Val Gln His Pro Glu Leu Thr Gly Leu Asp Cys Ile Arg Pro
385 390 395 400

Cys Phe Trp Val Glu Leu Ile Arg Gly Arg Pro Lys Glu Ser Thr Ile
405 410 415

Trp Thr Ser Gly Ser Ser Ile Ser Phe Cys Gly Val Asn Ser Asp Thr
420 425 430

Val Gly Trp Ser Trp Pro Asp Gly Ala Glu Leu Pro Phe Thr Ile Asp
435 440 445

Lys

<210> SEQ ID NO 12
<211> LENGTH: 2920
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CMV-T0-intron-M1 coding sequence-IRES-M2

-continued

coding sequence-polyA

<400> SEQUENCE: 12

```

gttgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata    60
gcccatatat ggagttccgc gttacataac ttacggtaaa tggcccgcct ggctgaccgc    120
ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt toccatagta acgccaatag    180
ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccac ttggcagtac    240
atcaagtgta tcatatgcca agtaacgccc ctattgacgt caatgacggg aaatggcccg    300
cctggcatta tgcccagtac atgaccttat gggactttcc tacttggcag tacatctacg    360
tattagtcat cgtattacc atggtgatgc ggttttgga gtacatcaat gggcgtggat    420
agcggtttga ctcaecggga tttccaagtc tccaccccat tgacgtcaat gggagtttgt    480
tttggcacca aaatcaacgg gactttccaa aatgtcgtaa caactccgcc ccattgacgc    540
aaatgggcgg taggcgtgta cgggtggagg tctatataag cagagctctc cctatcagtg    600
atagagatct cctatcagtg gatagagatc gtcgacgagc tcgttttagtg aaccgtcaga    660
tcgacctggag acgccatcca cgctgttttg acctccatag aagacaccgg gaccgatcca    720
gcctccggac tctagecgtt aaacttaagc ttggtaccga gctcggatcc cttgcagaag    780
ttggtcgtga ggcactgggc aggtaagtat caaggttaca agacaggttt aaggagacca    840
atagaaaactg ggcctgtgca gacagagaag actcctgctt ttctgatagg cacctattgg    900
tcttactgac atccactttg cctttctctc cacaggtgtc cactcccagt tcaattacag    960
ctcttaaggc tagagtactt aatacgactc actataggct agcatgagcc ttctaaccga   1020
ggtcgaaaac gatgttctct ctatogttcc atcaggcccc ctcaaagccg agatcgcgca   1080
gagacttgaa gatgtctttg ctgggaaaaa cacagatctt gaggetctca tggaaatggct   1140
aaagacaaga ccaattctgt cacctctgac taaggggatt ttggggtttg tgttcacgct   1200
caccgtgcca agtgagcggg gactgcagcg tagacgcttt gtccaaaatg ccctcaatgg   1260
gaatggagat ccaataaaca tggcacaagc agttaaactg tataggaaac ttaagaggga   1320
gataacgttc catggggcca aagaaatagc tctcagttat tctgctggtg cacttgccag   1380
ttgcatgggc ctcatataca atagaatggg ggctgtaacc actgaagtgg catttggcct   1440
ggtatgtgca acatgtgagc agattgctga ctcccagcac aggtctcata ggcaaatggt   1500
ggcaacaacc aatccattaa taagacatga gaacagaatg gttttggcca gcactacagc   1560
taaggctatg gagcaaatgg ctggatcaag tgagcaggca gcggaggcca tggagattgc   1620
tagtcaggcc aggcagatgg tgcaggcaat gagagccatt gggactcatc ctagttccag   1680
tactggtcta agagatgac ttcttgaaaa ttgacagacc tatcagaaac gaatgggggt   1740
gcagatgcaa cgattcaagt gagaattccg cccctctccc tcccccccc ctaacgttac   1800
tgccgaagc cgcttggaaat aaggccgggt tgtgtttgtc tatatgtgat tttccaccat   1860
attgocgtct tttggcaatg tgagggcccc gaaacctggc cctgtcttct tgacgagcat   1920
tctaggggt ctttcccctc tcgccaaggg aatgcaaggt ctggtgaatg tcgtgaagga   1980
agcagttcct ctggaagctt cttgaagaca aacaacgtct gtacggaccc tttgcaggca   2040
goggaaaccc ccacctggcg acaggtgctc ctgcgcccaa aagccacgtg tataagatac   2100
acctgcaaag gcggcacaac cccagtgcca cgttgtgagt tggatagttg tggaaagagt   2160

```

-continued

```

caaatggctc tctcaagcg tagtcaacaa ggggctgaag gatgcccaga aggtacccca 2220
ttgtatggga atctgatctg gggcctcggg gcacatgctt tacatgtggt tagtgcgaggt 2280
taaaaaaacg tctaggcccc ccgaaccacg gggacgtggg tttcctttga aaaacacgat 2340
gataaatatg ccacaaccgg atctatgagc cttctaaccg aggtcgaaac acctatcaga 2400
aacgaatggg ggtgcagatg caacgattca agtgaccgcg ttgttgttgc cgcgaatato 2460
attgggatct tgcacttgat attgtggatt cttgatcgtc tttttttcaa atgcgtctat 2520
cgactcttca aacacggcct taaaagaggg ccttctacgg aaggagtacc tgagtctatg 2580
aggaagaat atcgaaagga acagcagaat gctgtggatg ctgacgacag tcattttgtc 2640
agcatagagt tggagtaato tagagggccc gtttaaaccc gctgatcagc ctgcactgtg 2700
cctctagtt gccagccatc tgttgtttgc cctcccccg tgccttctt gaccctggaa 2760
ggtgccactc cactgtcct ttcctaataa aatgaggaaa ttgcatcgca ttgtctgagt 2820
aggtgtcatt ctattctggg ggggtgggtg gggcaggaca gcaaggggga ggattgggaa 2880
gacaatagca ggcctgctgg ggatgcgggt ggctctatgg 2920

```

<210> SEQ ID NO 13

<211> LENGTH: 2246

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: CMV promoter-rabbit beta-globin intron II-Tet R coding sequence-SV40 polyA

<400> SEQUENCE: 13

```

gttgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata 60
gccatataat ggagttccgc gttacataac ttacggtaaa tggcccgcct ggctgaccgc 120
ccaacgaccc ccgccattg acgtcaataa tgacgtatgt tcccatagta acgccaatag 180
ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccac ttggcagtac 240
atcaagtgta tcatatgcca agtacgcccc ctattgacgt caatgacggg aaatggcccc 300
cctggcatta tgcccagtac atgaccttat gggactttcc tacttggcag tacatctacg 360
tattagtcat cgtattacc atggatgatg ggttttgcca gtacatcaat gggcgtggat 420
agcggtttga ctcacgggga tttccaagtc tccaccccat tgacgtcaat gggagtttgt 480
tttggcacca aaatcaacgg gactttccaa aatgtcgtaa caactccgcc ccattgacgc 540
aaatgggccc taggcgtgta cgggtggagg tctatataag cagagctctc tggttaacta 600
gagaaccacc tgcttactgg cttatcgaaa ttaatacgac tcactatagg gagaccaag 660
ctggctagcg tttaaactta agcttggtag ccggggatcc tctagggcct ctgagctatt 720
ccagaagtag tgaagaggct tttttggagg cctaggcttt tgcaaaaagc tccggatcga 780
tcctgagaac ttcagggtga gtttggggac cttgattgt tctttctttt tcgctattgt 840
aaaattcatg ttatatggag ggggcaagt tttcagggtg ttgtttgaa tgggaagatg 900
tcccttgat caccatggac cctcatgata atttgttct tttcacttct tactctgttg 960
acaaccattg tctcctctta ttttctttc attttctgta acttttctgt taaactttag 1020
cttgcatgtg taacgaattt ttaaattcac ttttgtttat ttgtcagatt gtaagtactt 1080
tctctaatca ctttttttcc aaggcaatca gggatatatta tattgtactt cagcagctt 1140
ttagagaaca attgttataa ttaaatgata aggtagaata tttctgcata taaattctgg 1200

```

-continued

```

ctggcgtgga aatattctta ttggtagaaa caactacatc ctggatcatca tctgccttt 1260
ctctttatgg ttacaatgat atacactggt tgagatgagg ataaaatact ctgagtcctaa 1320
accgggcccc tctgctaacc atgttcacgc cttcttcttt ttcctacagc tctgaggcaa 1380
cgtgctgggtt attgtgctgt ctcacatctt tgccaagaa ttgtaatacg actcactata 1440
gggcgaattg atagtctag attagataaa agtaaagtga ttaacagcgc attagagctg 1500
cttaatgagg tcggaatcga aggtttaaca acccgtaaac tcgccagaa gctagggtga 1560
gagcagccta cattgtattg gcacgtaaaa aataagcggg ctttgcctga cgccttagcc 1620
attgagatgt tagataggga ccatactcac ttttgcctt tagaagggga aagctggcaa 1680
gattttttac gtaataacgc taaaagtttt agatgtgctt tactaagtca tcgcatgga 1740
gcaaaagtac atttaggtag ccggcctaca gaaaaacagt atgaaactct cgaatacaa 1800
ttagcctttt tatgccaaca aggtttttca ctagagaatg cattatatgc actcagcgt 1860
gtggggcatt ttactttagg ttgcgtattg gaagatcaag agcatcaagt cgctaagaa 1920
gaaagggaaa cacctactac tgatagtagt ccgccattat tacgacaagc tatcgaatta 1980
tttgatcacc aagggtcaga gccagccttc ttattcggcc tgaattgat catatgcgga 2040
ttagaaaaac aacttaaatg tgaaagtggg tccgcgtaca gcggatccc ggaattcaga 2100
tcttattaaa gcagaacttg tttattgcag cttataatgg ttacaaataa agcaatagca 2160
tcacaaatth cacaaataaa gcattttttt cactgcattc tagttgtggt ttgtccaaa 2220
tcatcaatgt atcttateat gtctgg 2246

```

<210> SEQ ID NO 14

<211> LENGTH: 5646

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: DNA sequence encoding influenza H3 and N2, including the 5' and 3' expression control region

<400> SEQUENCE: 14

```

gttgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata 60
gccatataat ggagttccgc gttacataac ttacggtaaa tggcccgcct ggctgaccgc 120
ccaacgaccc ccgccattg acgtcaataa tgacgtatgt tccatagta acgccaatag 180
ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccac ttggcagtac 240
atcaagtgta tcatatgcca agtacgcccc ctattgacgt caatgacggg aaatggcccg 300
cctggcatta tgcccagtag atgaccttat gggactttcc tacttggcag tacatctacg 360
tattagtcac cgctattacc atggtgatgc ggttttgcca gtacatcaat gggcgtggat 420
agcggtttga ctcacgggga tttccaagtc tccaccccat tgacgtcaat gggagtttgt 480
tttggcacca aaatcaacgg gactttccaa aatgtcgtaa caactccgcc ccattgacgc 540
aaatgggccc taggcgtgta cgggtggagg tctatataag cagagctctc cctatcagtg 600
atagagatct cctatcagt gatagagatc gtcgacgagc tcgtttagtg aaccgtcaga 660
tcgcctggag acgccatcca cgctgttttg acctccatag aagacaccgg gaccgatcca 720
gcctccggac tctagcgttt aaacttaagc ttggtaccga gtcgggatcc cttgcagaag 780
ttggtcgtga ggcactgggc aggttaagat caaggttaca agacaggttt aaggagacca 840

```

-continued

atagaaactg	ggcttgcga	gacagagaag	actcttgcgt	ttctgatagg	cacctattgg	900
tcttactgac	atccactttg	ccttctctc	cacaggtgto	cactcccagt	tcaattacag	960
ctcttaaggc	tagagtactt	aatcagactc	actataggct	agaatgaatc	caaatcaaaa	1020
gataataacg	attggctctg	tttctctc	catttccaca	atatgcttct	tcatgcaaat	1080
tgccatcttg	ataactactg	taacattgca	tttcaagcaa	tatgaattca	actccccccc	1140
aaacaaccaa	gtgatgctgt	gtgaaccaac	aataatagaa	agaaacataa	cagagatagt	1200
gtatctgacc	aacaccacca	tagagaagga	aatatgcccc	aaactagcag	aatacagaaa	1260
ttggtcacaa	ccgcaatgta	acattacagg	atttgcaact	ttttctaagg	acaattcgat	1320
taggctttcc	gctgggtggg	acatctgggt	gacaagagaa	ccttatgtgt	catgcatcc	1380
tgacaagtgt	tatcaatttg	cccttgaca	gggaacaaca	ctaacaacg	tgcatcaaaa	1440
tgacacagta	catgatagga	ccccttatcg	gaccctattg	atgaatgagt	taggtgttcc	1500
attcatctg	gggaccaagc	aagtgtgcat	agcatggctc	agctcaagtt	gtcacgatgg	1560
aaaagcatgg	ctgcatggtt	gtgtaacggg	ggatgataaa	aatgcaactg	ctagcttcat	1620
ttacaatggg	aggctttag	atagtattgt	ttcgtggctc	aaagaaatcc	tcaggacca	1680
ggagtcagaa	tgctttagta	tcaatggaac	ttgtacagta	gtaatgactg	atgggagtg	1740
ttcaggaaaa	gctgatacta	aaatactatt	cattgaggag	gggaaatcg	ttcatactag	1800
cacattgtca	ggaagtgtc	agcatgtcga	ggagtgttcc	tgctatccta	gatatcctgg	1860
tgtcagatgt	gtctgcagag	acaactggaa	agctccaat	aggcccatag	tagatataaa	1920
cataaaggat	tatagcattg	tttccagtta	tgtgtgctca	ggacttgttg	gagacacacc	1980
cagaaaaaac	gacagctcca	gcagtggcca	ttgcttggat	cctaacaatg	aagaaggtgg	2040
tcattggagt	aaaggctggg	cctttgatga	tggaaatgac	gtgtggatgg	gaagaacgat	2100
cagcgagaag	ttacgctcag	gatatgaaac	cttcaaagtc	attgaaggct	ggtccaaccc	2160
taattccaaa	ttgcagataa	ataggcaagt	catagttagc	agaggtaata	ggtccgggta	2220
ttctggtatt	ttctctggtg	aaggcaaaa	ctgcatcaat	cggtgctttt	atgtggagtt	2280
gataagggga	agaaaagagg	aaactgaagt	cttgtggacc	tcaaacagta	ttgttgtgtt	2340
ttgtggcacc	tcaggtagat	atggaacagg	ctcatggcct	gatggggcgg	acatcaatct	2400
catgctata	taatctagag	ggcccgttta	aaccctgta	tcagctcga	ctgtgccttc	2460
tagttgccag	ccatctggtg	tttgcctc	ccccgtgct	tccttgacc	tggaaggtgc	2520
cactcccact	gtcctttcct	aataaaatga	ggaaattgca	tcgcatgtc	tgagttagtg	2580
tcattctatt	ctgggggggtg	gggtggggca	ggacagcaag	ggggaggatt	gggaagacaa	2640
tagcaggcat	gctgggggatg	cggtgggctc	tatggcacgc	gttgacattg	attattgact	2700
agttattaat	agtaatcaat	tacgggggtca	ttagttcata	gccatataat	ggagttccgc	2760
gttacataac	ttacggtaaa	tgccccgct	ggctgaccgc	ccaacgaccc	ccgcccattg	2820
acgtcaataa	tgacgtatgt	tccatagta	acgccaatag	ggactttcca	ttgacgtcaa	2880
tggttgaggt	atttacggta	aactgcccac	ttggcagtac	atcaagtgt	tcatatgcca	2940
agtagcccc	ctattgacgt	caatgacgg	aaatggccc	cctggcatta	tgcccagtac	3000
atgaccttat	gggactttcc	tacttggcag	tacatctacg	tattagtc	cgctattacc	3060
atggtgatgc	ggttttggca	gtacatcaat	ggcgtggat	agcggtttga	ctcacgggga	3120

-continued

tttccaagtc tccaccccat tgacgtcaat gggagtttgt tttggcacca aaatcaacgg	3180
gactttccaa aatgtcgtaa caactccgcc ccattgacgc aaatgggagg taggcgtgta	3240
cggtgggagg tctatataag cagagctctc cctatcagtg atagagatct cctatcagt	3300
gatagagatc gtcgacgagc tcgttttagtg aaccgtcaga tcgcctggag acgccatcca	3360
cgctgttttg acctccatag aagacaccgg gaccgatcca gctccggac tctagcgttt	3420
aaacttaagc ttggtagcga gctcggatcc cttgcagaag ttggtcgtga ggcactgggc	3480
aggtaagtat caaggttaca agacaggttt aaggagacca atagaaactg ggcttgctga	3540
gacagagaag actccttgctt ttctgatagg cacctattgg tcttactgac atccactttg	3600
cctttctctc cacagggtgc cactcccagt tcaattacag ctcttaaggc tagagtactt	3660
aatacgactc actataggct agcatgaaga ctatcattgc tttgagctac attctatgtc	3720
tggttttcgc tcaaaaactt cccggaaatg acaacagcac ggcaacgctg tgccttgggc	3780
atcatgcagt accaaaacgga acgatatgta aaacaatcac gaatgaccaa attgaagtta	3840
ctaagtctac tgagctgggt cagagttcct caacagggtg aatatgcgac agtccctac	3900
agatccttga tggagaaaa tgcacactaa tagatgctct attgggagac cctcagtgtg	3960
atggcttcca aaataagaaa tgggaccttt ttggtgaacg cagcaaagcc tacagcaact	4020
gttaccctta tgatgtgccg gattatgctc cccttaggtc actagttgcc tcatccggca	4080
cactggagtt taacaatgaa agcttcaatt ggactggagt cactcaaat ggaacaagct	4140
ctgcttgcaa aaggagatct aataacagtt tctttagtag attgaattgg ttgaccact	4200
taaaaattcaa ataccagca ttgaacgtga ctatgccaaa caatgaaaaa tttgacaaat	4260
tgtacatttg gggggttcc caccgggta cggacaatga ccaaatcttc ctgtatgctc	4320
aagcatcagg aagaatcaca gtctctacca aaagaagcca acaactgta atcccgaata	4380
tcggatctag acccagagta agggatatcc ccagcagaat aagcatctat tggacaatag	4440
taaaaccggg agacatactt ttgattaaca gcacaggaa tctaattgct cctagggggt	4500
acttcaaat acgaagtggg aaaagctcaa taatgagatc agatgcaccc attggcaaat	4560
gcaattctga atgcatcact ccaaatggaa gcattcccaa tgacaaaacca tttcaaatg	4620
taaacaggat cacatatggg gcctgtccca gatatgtaa gcaaacact ctgaaattgg	4680
caacagggat gcgaaatgta ccagagaaac aaactagagg catatttggc gcaatcggcg	4740
gtttcataga aaatggtttg gagggaatgg tggatggttg gtacggtttc aggcacaaa	4800
attctgaggg aataggacaa gcagcagatc tcaaaagcac tcaagcagca atcaatcaaa	4860
tcaatgggaa gctgaatagg ttgatcggga aaaccaacga gaaattccat cagattgaaa	4920
aagaattctc agaagttaga gggagaatc aggacctga gaaatggtt gaggacacta	4980
aaatagatct ctggtcatac aacgcggagc ttcttgttgc cctggagaac caacatacaa	5040
ttgatctaac tgactcagaa atgaacaaac tgtttgaaag aacaaagaag caactgaggg	5100
aaaatgctga ggatatgggc aatggttgtt tcaaaatata ccacaaatgt gacaatgcct	5160
gcataggatc aatcagaaat ggaacttatg accatgatgt atacagagat gaagcattaa	5220
acaaccggtt ccagatcaaa ggcgttgagc tgaagtcagg atacaaagat tggatectat	5280
ggatttcctt tgccatatac tgttttttgc tttgtgttgc tttgtgggg ttcacatgt	5340
gggcctgcca aaaaggcaac attaggtgca acatttgcatt ttgagctaga gggcccggtt	5400

-continued

```

aaacccgctg atcagcctcg actgtgcctt ctagtgtcca gccatctgtt gtttgeccct 5460
ccccctgccc ttccttgacc ctggaagggt ccactcccac tgtcctttcc taataaaatg 5520
aggaaattgc atcgattgt ctgagtaggt gtcattctat tctggggggt ggggtggggc 5580
aggacagcaa gggggaggat tgggaagaca atagcaggca tgctggggat gcggtgggct 5640
ctatgg 5646

```

```

<210> SEQ ID NO 15
<211> LENGTH: 5598
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence encoding influenza H5 and N1,
including the 5' and 3' expression control region

<400> SEQUENCE: 15

```

```

gttgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata 60
gcccataat ggagttccgc gttacataac ttacggtaaa tggcccgcct ggctgaccgc 120
ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt tcccatagta acgccaatag 180
ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccac ttggcagtac 240
atcaagtgta tcatatgcca agtacgcccc ctattgacgt caatgacggg aaatggcccc 300
cctggcatta tgcccagtac atgaccttat gggactttcc tacttggcag tacatctacg 360
tattagtcac cgetattacc atggtgatgc ggttttgcca gtacatcaat gggcgtggat 420
agcgggttga ctcaagggga tttccaagtc tcaacccat tgacgtcaat gggagtttgt 480
tttggcacca aaatcaacgg gactttccaa aatgtcgtaa caactccgcc ccattgacgc 540
aaatgggccc taggcgtgta cgggtggagg tctatataag cagagctctc cctatcagtg 600
atagagatct ccctatcagt gatagagatc gtcgacgagc tegttagtg aacgcgaga 660
tcgcctggag acgccatcca cgctgttttg acctccatag aagacaccgg gaccgatcca 720
gcctccggac tctagcgttt aaacttaagc ttggtaccga gctcggatcc cttgcagaag 780
ttggtcgtga ggcactgggc aggtaagtat caaggttaca agacaggttt aaggagacca 840
atagaaaactg ggtctgtgca gacagagaag actcctgcgt ttctgatagg cacctattgg 900
tcttactgac atccactttg cctttctctc cacaggtgto cactcccagt tcaattacag 960
ctcttaaggc tagagtactt aatacgactc actataggct agaatgaacc ccaaccagaa 1020
gateatcacc atcggctcca tctgcatggt gaccggtatc gtgtccctga tgctgcagat 1080
cggtaacatg atctccatct ggggtgtcca ctccatccac accggttaacc agcaccagtc 1140
cgagcccac tccaacacca agttctccac cgagaaggct gtggcttccg tgaagctggc 1200
tggtaaactc tcctgtgccc ccatcaacgg ttgggctgtg tactccaagg acaactccat 1260
ccgtatcggg tccaaggggt acgtgttcgt gatccgtgag ccttccatct cctgctccca 1320
cctggagtgc cgtaccttct tcctgaccca ggggtgctctg ctgaacgaca agcactccaa 1380
cggcaccgtg aaggaccgtt cccccaccg taccctgatg tcctgccccg tgggagaggc 1440
tcctccccc tacaactccc gtttcgagtc cgtggcttgg tccgcttccg cttgccacga 1500
cggcacctct tggctgacca tcggatctct cggctcccac aacgggtgctg tggctgtggt 1560
gaagtacaac ggcacatca cgcacacat caagtctcgg cgtaacaaca tctgctgtac 1620
ccaagagtcc gagtgcgctt cgtggaacgg ttctgcttc accgtgatga ccgacggctc 1680

```

-continued

ctccaacggc caggcttccc acaagatctt caagatggag aagggcaagg tggtaagtc	1740
cgtggagctg gacgctccca actaccacta cgaggagtgc tcttgctacc cggacgctgg	1800
cgagatcacc tgcgtgtgcc gtgacaaactg gcacggttcc aacegtecct ggggtgcctt	1860
caaccagaac ctggagtacc agatcgggta catctgctcc ggcgtgttcg gtgacaaacc	1920
ccgtcccaac gacggaaccg gttcttgccg tcccgtgtcc tccaacggtg cttacgggtg	1980
caagggttc tcttcaagt acggtaacgg tgtctggatc ggctgtacca agtccaccaa	2040
ctcccgtcc ggtttcgaga tgatctggga ccccaacggg tggaccgaga ccgactctc	2100
cttctccgtg aagcaggaca tcgtggctat caccgactgg tccggttact ccggttcctt	2160
cgtgcagcac cccgagctga ccggtctgga ctgtatccgt ccctgcttct ggggtggaget	2220
gatcgtggg cgtcccaagg agtccaccat ctggacctcc ggctcctcca tctctttctg	2280
cggtgtgaac tccgacaacc tgggctgggc ctggcccagc ggtgcccagc tgccttccac	2340
catcgacaag taggcggccg cgggcccgtt taaaccgct gatcagcctc gactgtgcct	2400
tctagttgcc agccatctgt gtgttgcctc tccccgtgc ctctcttgac cctggaaggt	2460
gccactccca ctgtccttcc ctaataaaat gaggaattg catcgattg tctgagtagg	2520
tgatattcta tctggggggc tggggtgggc caggacagca agggggagga ttgggaagac	2580
aatagcaggc atgctgggga tgcgggtggc tctatggacg cgttgacatt gattattgac	2640
tagttattaa tagtaataaa ttacgggggc attagttcat agcccatata tggagttccg	2700
cgttacataa cttacggtaa atggcccgc tggctgaccg cccaacgacc cccgcccatt	2760
gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc attgacgtca	2820
atgggtggag tatttaccgt aaactgccc cttggcagta catcaagtgt atcatatgcc	2880
aagtacgccc cctattgacg tcaatgacgg taaatggccc gctggcatt atgcccagta	2940
catgacctta tgggacttcc ctacttgcca gtacatctac gtattagtca tcgctattac	3000
catggtgatg cggttttggc agtacatcaa tgggctgga tagcggtttg actcacgggg	3060
atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc aaaatcaacg	3120
ggactttcca aaatgctgta acaactccgc cccattgacg caaatgggcg gtaggcgtgt	3180
acggtgggag gtctatataa gcagagctct ccctatcagt gatagagatc tccctatcag	3240
tgatagagat cgtcagcagc ctcgtttagt gaaccgtcag atcgccctgga gacgcatcc	3300
acgctgtttt gacctccata gaagacaccg ggaccgatcc agcctccgga ctctagcgtt	3360
taaaactaag cttggtaccg agctcggatc ccttgacgaa gttggctcgtg aggcactggg	3420
caggtaagta tcaaggttac aagacaggtt taaggagacc aatagaaact gggcttgcg	3480
agacagagaa gactcttgcc tttctgatag gcacctattg gtcttactga catccacttt	3540
gcctttctct ccacaggtgt ccaactccag ttcaattaca gctcttaagg ctagagtact	3600
taatcagact cactatagcc tagcatggag aagatcgtgc tgetgttcgc tatcgtgtcc	3660
ctggtgaagt ccgaccagat ctgcatcggg taccacgcta acaactccac cgagcaggtg	3720
gacaccatca tggagaagaa cgtcaccgtg acccacgctc aggacatcct ggaaaagacc	3780
cacaacggca agctgtgcga cctggacggg gtcaagcccc tgatcctgcg tgactgctcc	3840
gtggctgggt ggtcgtggg taaccccatg tgcgacgagt tcatcaacgt gcccgagtgg	3900
tctacatcg tggagaaggc taaccccggt aacgacctgt gctaccccg tgacttcaac	3960

-continued

```

gactacgagg agctgaagca cctgctgtcc cgtatcaacc acttcogagaa gatccagatc 4020
atccccaagt cctcttggct gtcccacgag gcttccctgg gcgtgtcctc cgettgccca 4080
taccagggca agtccagctt ctcccgtaac gtggtgtggc tgatcaagaa gaactctacc 4140
taccocacca tcaagcgttc ctacaacaac accaaccagg aggacctgct ggtcctgtgg 4200
ggatccacc accccaacga cgctgccgag cagaccaago tgtaccagaa ccccaccacc 4260
tacatctcgg tgggcacctc cacccctaac cagcgtctgg tgccccgtat cgtaccctgt 4320
tccaagtgta acggccagtc cggctgtatg gagttcttct ggaccatcct gaagcctaac 4380
gacgctatca acttcgagtc caacggcaac ttcctcgtcc cggagtacgc ttacaagatc 4440
gtgaagaagg gcgactccac catcatgaag tccgagctgg agtacggtaa ctgcaacacc 4500
aagtgccaga cccccatggg tgctataaac tctccatgc ccttcocaaa catccacccc 4560
ctgaccatcg gcgagtgcce caagtacgtg aagtccaacc gctcgtgtgt ggtaccgggt 4620
ctgcgtaact cccccagcg tgagcgtgtg aagaagcgtg gtctgttcgg cgtatcgtct 4680
ggtttcatcg agggcggttg gcagggcatg gtggacggtt ggtacgggta ccaccactct 4740
aacgagcagg gttccgggta cgctcgtgac aaggagtcca cccagaagge tatcgacggc 4800
gtcaccaaca agtggaactc catcatcgac aagatgaaca cccagttcga ggctgtgggt 4860
cgtgagttca acaacctgga gcgctgatac gagaacctga acaagaagat ggaggacgggt 4920
ttcctggacg tgtggaccta caacgccgag ctgctggtgc tgatggagaa cgagcgtacc 4980
ctggacttcc acgactccaa cgtgaagaac ctgtacgaca aggtccgctt gcagctgctg 5040
gacaacgcta aggagctggg taacggttgc ttcgagttct accacaagtg cgacaacgag 5100
tgcatggagt cctgctgtaa cggcacctac gactaccccc agtactccga ggaggccaag 5160
ctgaagcgtg gcgagatctc cggcgtgaag ctggagtcca tcggtatcta ccagatcctg 5220
tccatctact ccaccgtggc ttccctcctg gctctggcta tcatggtggc tggctctgctc 5280
ctgtggatgt gctccaacgg ttccctgcag tgcctatctt gcatctagge ggccgcgcta 5340
gagggcccgt ttaaacccgc tgatcagcct cgactgtgcc ttctagtgc cagccatctg 5400
ttgtttgccc cteccccgtg ccttccctga cctggaagg tgccactccc actgtccttt 5460
cctaataaaa tgaggaaatt gcatcgatt gtctgagtag gtgtcattct attctggggg 5520
gtggggggg gcaggacagc aagggggagg attgggaaga caatagcagg catgctgggg 5580
atgcggggg ctctatgg 5598

```

```

<210> SEQ ID NO 16
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RT-PCR primer

```

```

<400> SEQUENCE: 16

```

```

attaggcttt ccgctggtgg ggacat 26

```

```

<210> SEQ ID NO 17
<211> LENGTH: 28
<212> TYPE: DNA

```

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: RT-PCR primer

<400> SEQUENCE: 17

gcattctgac tctctgggtcc tgaggatt

28

We claim:

1. A method of preparing an influenza virus-like particle (VLP), the method comprising:

obtaining a founder Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2;

constructing at least one recombinant DNA molecule comprising a sequence encoding an influenza hemagglutinin (HA) and a sequence encoding an influenza neuraminidase (NA);

introducing the at least one recombinant DNA molecule into the founder Vero cell to obtain a co-expression Vero cell stably transfected with the sequences encoding the influenza M1 and the influenza M2, and further transfected with the sequences encoding the influenza HA and the influenza NA, wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems;

culturing the co-expression Vero cell under conditions to allow expressions of the influenza M1, the influenza M2, the influenza HA and the influenza NA, and assembly of the VLP comprising the influenza M1, the influenza M2, the influenza HA and the influenza NA; and

isolating the VLP from the culture supernatant of the co-expression Vero cell.

2. The method according to claim 1, wherein the founder Vero cell is derived from Vero E6 cell.

3. The method according to claim 1, wherein the influenza HA and the influenza NA are derived from one or more potentially pandemic or seasonal influenza virus strains.

4. The method according to claim 1, wherein the at least one recombinant DNA molecule further comprises a sequence encoding a second influenza HA and a sequence encoding a second influenza NA, wherein the influenza HA and the second influenza HA are derived from different influenza virus strains, and the influenza NA and the second influenza NA are derived from different influenza virus strains;

the co-expression Vero cell is further transfected with the sequences encoding the second influenza HA and the second influenza NA; and

the culturing step allows expressions of the influenza M1, the influenza M2, the influenza HA, the influenza NA, the second influenza HA and the second influenza NA, and assembly of the VLP comprising the influenza M1, the influenza M2, at least one of the influenza HA and the second influenza HA, and at least one of the influenza NA and the second influenza NA.

5. A method of preparing an influenza virus-like particle (VLP), the method comprising:

obtaining a co-expression Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence

encoding an influenza M2, and further transfected with a sequence encoding an influenza hemagglutinin (HA) and a sequence encoding an influenza neuraminidase (NA), wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems; and

culturing the co-expression Vero cell under conditions to allow expressions of the influenza M1, the influenza M2, the influenza HA and the influenza NA, and assembly of the VLP comprising the influenza M1, the influenza M2, the influenza HA and the influenza NA; and

isolating the VLP from the culture supernatant of the co-expression Vero cell

6. An influenza virus-like particle (VLP), comprising: an influenza M1, an influenza M2, an influenza hemagglutinin (HA) and an influenza neuraminidase (NA), wherein the influenza proteins are recombinantly expressed from a Vero cell; and

at least one cellular protein of the Vero cell.

7. The influenza VLP of claim 6, wherein the Vero cell is Vero E6 cell.

8. The influenza VLP of claim 6, wherein the influenza HA and the influenza NA are derived from one or more potentially pandemic or seasonal influenza virus strains.

9. The influenza VLP of claim 6, further comprising a second influenza HA and a second influenza NA, wherein the influenza HA and the second influenza HA are derived from different influenza virus strains, and the influenza NA and the second influenza NA are derived from different influenza virus strains.

10. The influenza VLP of claim 6, comprising M1 and M2 of influenza A/Taiwan/083/2006, and H3 and N2 of influenza A/Taiwan/083/2006, or H5 and N1 of influenza A/Hanoi/30408/2005(H5N1), respectively.

11. The influenza VLP of claim 6, wherein the at least one cellular protein is selected from the group consisting of clathrin heavy chain 1, spectrin beta, plexin B2, CD109 homolog, prostaglandin F2 receptor negative regulator, Na⁺/K⁺-ATPase alpha 1, tumor rejection antigen (gp96) 1 and flotillin I.

12. An immunogenic composition comprising an immunogenic effective amount of the influenza VLP of claim 6 and a pharmaceutically acceptable excipient.

13. The immunogenic composition of claim 12 further comprising an adjuvant.

14. A method of inducing immunity against an influenza virus in a subject, the method comprising administering to the subject the immunogenic composition of claim 12, wherein the influenza VLP comprises the HA and the NA derived from the influenza virus.

15. A method of providing protection against an influenza virus in a subject, the method comprising vaccinating the subject with a vaccine comprising the influenza VLP of claim

6, wherein the influenza VLP comprises the HA and the NA derived from the influenza virus.

16. A method of diagnosing infection by an influenza virus in a subject, the method comprising:

obtaining a biological sample from the subject;
contacting the biological sample with the influenza VLP of claim **6**, wherein the influenza VLP comprises the HA and the NA derived from the influenza virus; and
measuring in the biological sample the amount of an antibody that forms an antibody-antigen complex with the influenza VLP,

wherein a higher than a threshold amount of the antibody indicates that the subject has ever been infected by the influenza virus or vaccinated by a similar strain of virus.

17. An antibody against the influenza VLP of claim **6**.

18. A method of preparing the antibody of claim **17**, comprising:

administering the VLP to a vertebrate; and
harvesting the antibody against the VLP from the blood of the vertebrate.

19. A method of treating infection by an influenza virus in a subject, the method comprising administering to the subject the antibody of claim **17**, wherein the influenza VLP comprises the HA and the NA derived from the influenza virus.

20. A method for detecting infection by an influenza virus in a subject, the method comprising:

obtaining a biological sample from the subject;
contacting the biological sample with the antibody of claim **17**; and

detecting in the biological sample an antigen that forms an antibody-antigen complex with the antibody,
wherein the presence of the antigen in the biological sample indicates that the subject is infected by the influenza virus.

21. A founder Vero cell being a Vero cell stably transfected with a sequence encoding an influenza M1 and a sequence encoding an influenza M2, wherein the expressions of the influenza M1 and M2 proteins in the founder Vero cell are controlled by an inducible expression system.

22. The founder Vero cell of claim **21**, wherein the Vero cell is Vero E6 cell.

23. The founder Vero cell of claim **21** further transfected with a sequence encoding an influenza protein different from the influenza M1 and M2 proteins.

24. The founder Vero cell of claim **23** being a co-expression Vero cell stably transfected with the sequence encoding the influenza M1 and the sequence encoding the influenza M2, and further transfected with a sequence encoding an influenza hemagglutinin (HA) and a sequence encoding an influenza neuraminidase (NA), wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems.

25. A method of obtaining a founder Vero cell, the method comprising:

introducing into a Vero cell a sequence encoding an influenza M1 and a sequence encoding an influenza M2; and
obtaining the founder Vero cell stably transfected with the sequence encoding the influenza M1 and the sequence encoding the influenza M2,

wherein the expressions of the influenza M1 and M2 proteins in the founder Vero cell are controlled by an inducible expression system.

26. The method of claim **25**, further comprising:

introducing into the founder Vero cell a sequence encoding an influenza hemagglutinin (HA) and a sequence encoding an influenza neuraminidase (NA); and

obtaining a co-expression Vero cell stably transfected with the sequence encoding the influenza M1 and the sequence encoding the influenza M2, and further transfected with the sequence encoding the influenza HA and the sequence encoding the influenza NA,

wherein the expressions of the influenza M1, M2, HA and NA proteins in the co-expression Vero cell are controlled by one or more inducible expression systems.

* * * * *

专利名称(译)	用于产生病毒样颗粒的高产转基因哺乳动物表达系统		
公开(公告)号	US20100166769A1	公开(公告)日	2010-07-01
申请号	US12/704928	申请日	2010-02-12
[标]申请(专利权)人(译)	中央研究院		
申请(专利权)人(译)	中央研究院		
当前申请(专利权)人(译)	中央研究院		
[标]发明人	HSIAO PEI WEN WU CHIA YING YEYI CHUN		
发明人	HSIAO, PEI-WEN WU, CHIA- YING YEYI, YI-CHUN		
IPC分类号	A61K39/42 C12P21/00 C12N9/26 A61K39/145 G01N33/53 C07K16/08 C12N5/07 C12N15/00 A61P31/16		
CPC分类号	A61K39/145 G01N2469/20 A61K2039/5258 C07K14/005 C12N2760/16122 C12N2760/16123 C12N2760/16134 C12N2770/20022 C12N2770/20023 C12N2830/006 C12N2770/20034 A61K39/12 C07K16/1018 C12N2760/16034 G01N33/56983 G01N2333/11 G01N2469/10 A61K39/215		
其他公开文献	US8980281		
外部链接	Espacenet USPTO		

摘要(译)

已经从Vero细胞重组产生哺乳动物宿主病毒(例如SARS-CoV和流感病毒)的病毒样颗粒(VLP)。VLP紧密模拟真实病毒颗粒的外部并具有高度免疫原性。它们不仅可以引发哺乳动物的体液免疫反应,还可以引发细胞免疫反应。还描述了与VLP相关的组合物和方法。

