



US 20100261160A1

(19) **United States**(12) **Patent Application Publication**  
**Wood et al.**(10) **Pub. No.: US 2010/0261160 A1**(43) **Pub. Date: Oct. 14, 2010**(54) **METHODS OF SCREENING FOR  
MODULATORS OF CELL PROLIFERATION  
AND METHODS OF DIAGNOSING CELL  
PROLIFERATION STATES**(75) Inventors: **Kenneth W. Wood**, Foster City, CA  
(US); **Jeffrey T. Finer**, Foster City,  
CA (US); **Christophe Beraud**, San  
Francisco, CA (US); **John Mak**,  
San Bruno, CA (US); **Roman**  
**Sakowicz**, Foster City, CA (US)Correspondence Address:  
**TOWNSEND AND TOWNSEND AND CREW,  
LLP  
TWO EMBARCADERO CENTER, EIGHTH  
FLOOR  
SAN FRANCISCO, CA 94111-3834 (US)**(73) Assignee: **Cytokinetics, Inc.**, South San  
Francisco, CA (US)(21) Appl. No.: **11/503,530**(22) Filed: **Aug. 10, 2006****Related U.S. Application Data**(60) Continuation of application No. 10/601,036, filed on  
Jun. 19, 2003, now abandoned, which is a division of  
application No. 09/428,156, filed on Oct. 27, 1999,  
now Pat. No. 6,617,115.**Publication Classification**(51) **Int. Cl.**  
**C12Q 1/68** (2006.01)  
**C12Q 1/02** (2006.01)  
**C12Q 1/34** (2006.01)  
**G01N 33/53** (2006.01)  
**G01N 27/447** (2006.01)  
(52) **U.S. Cl. .... 435/6; 435/29; 435/18; 435/7.92;**  
204/456(57) **ABSTRACT**Described herein are methods that can be used for diagnosis  
and prognosis of cellular proliferation. Also described herein  
are methods that can be used to screen candidate bioactive  
agents for the ability to modulate cellular proliferation. Addi-  
tionally, methods and molecular targets (genes and their prod-  
ucts) for therapeutic intervention in cancers are described.

gaattccgctc AGGCGTCGC agccaaattc gtctgcgaag agaaagagg agaaggggaa  
gaacatccag gtgggtggtga gatgcagacc atttaatttg gcagagcgga aagcttagcgc  
ccattcaata gtagaatgtg atcctgtacg aaaagaagtt agtgtacgaa ctggaggatt  
ggctgacaag agctcaagga aaacatacac ttttgatatg gtgcttggag catctactaa  
acagattgat gtttacggaa gtgtgtgttg tccaattctg gatgaagtta ttatgggcta  
taattgcact atccttgcgt atggccaaac tggcactgga aaaactttta caatgggaagg  
tgaaagggtca cctaatagaag agtataacctg ggaagaggat cccttggctg gtataattcc  
acgtaccctt catcaaatct ttgagaaact tactgataat ggtactgaat ttctagctaa  
agtgtctctg ttggagatct ataatgaaga gctttctgat ctcttaate catcatctga  
tgtttctgag agactacaga tgtttgatga tccccgtaac aagagaggag tgataactaa  
aggcttagaa gaaattacag tacacaacaa ggatgaagtc tatcaaattt tagaaaaggg  
ggcagcaaaa aggacaactg cagctactct gatgaatgca tactctagtc gttccactc  
agtttctctt gttacaatac atatgaaaga aactacgatt gatggagaag agcttgttaa  
aatcggaag ttgaacttgg ttgatcttgc aggaagtga aacattggcc gttctggagc  
tgttgataag agagctcggg aagctggaaa tataaatcaa tccctgttga ctttgggaag  
ggcattact gcccttctag aaagaacacc tcatgttctt tatcgagaat agcttgttaa  
tagaatcttc caggattctc ttggaggcg tacaagaaca tctataattg caacaattc  
tcctgcatct ctcaatcttg aggaaactct gagtacattg gaatatgtc atagagcaaa  
gaacatatg aataagcctg aagtgaatca gaaactcacc aaaaagctc ttattaggaa  
gtataccgag gagatagaac gtttaaaacg agatcttgc gcagccctg agaaaaattg  
agtggtatct cctgaagaaa attttagagt catgagtggg aatcaactg ttcaagaaga  
gcagattgta gaattgattg aaaaaatttg tgcgttggag gaggagctga atagggttac  
agagttgttt atggataata aaaaatgaact tgaccagtg aaatctgacc tgcaaaataa  
aacacaagaa cttgaaacca ctcaaaaaca ttgcaagaa actaaattac aacttgttaa  
agaagaatc atcacatcag ctttggaaag tactgaggag aaactctatg atgctgccag  
caagctgctt aacacagttg aagaaactac aaaagatgta tctggtctcc attccaaact  
ggatcgtaag aaggcagttg accaacacaa tgcagaagct caggatattt ttggcaaaaa  
cctgaatagt ctgtttaata atatggaaga attaattag gatggcagct caaagcaaaa  
ggccatgcta gaagtacata agaccttact tggtaattct tctcttcca gtgtctctgc  
attagatacc attactacag tagcacttgg atctctcaca tctattccag aaaaatgtgtc  
tactcatgtt tctcagattt ttaatatgat actaaagaa caatcattag cagcagaag  
taaaactgta ctacaggaat tgattaatgt actcaagact garcttctaa gtctcattga  
aargatttta tccccaaact tgggtgtctat actgaaaaac aatagtcaac taagcatal  
tttcaagact tcattgacag tggccgataa gatagaagat caaaaaaaa ggaactcaga  
tggcttctc agtatactgc gtaaccaatct acatgaacta caagaaaaa ccatrtgttc  
cttggttgag tcacaaaaagc aactgtgaaa cctaactgaa gacctgaaga caataaagca  
gacccattcc caggaaactt tacaagaaacc gcaagctaat gaattcttgg acagagagat  
ggaggaaaaa tgtgaaaaata tacagaaacc acttagtagt gtccaggaaa atatacaga  
gaaatcttaag gatcatgtca acaaaaatgac ttttcacagt caaaaaattt ttgctgtctc  
tgatggcttc tcacaggaac tcagaaattt taaccaagaa ggtacaaaat tggtrgaaga  
atctgtgaaa cactctgata aactcaatgg caacctggaa aaaaatatcc aagagactga  
acagagatct gaattctctga acacaagaac agtttatttt tctgaacagt gggatctctc  
cttaaatgaa aggggaacagg aacttcacaa cttatggag gtctgaagcc aatgttgtga  
ggcttcaagt tcagacatca ctgagaaatc agatggagct aaggcagctc atgagaaaca  
gcataacatt tttcttgatc agatgactat tgatgaagat aaattgatag cacaaaactc  
agaacttaat gaaaccataa aaattcggtt gactaagctt aattgctttc tggacagga  
cctgaaactg gatattccaa caggtagcag accacagagg aaaaagtatt tataccatc  
aacactggta agaactgaac cactgtgaaca tctccttgat cagctgaaaa ggaacagcc  
tgagctgtta atgatgctaa actgttcaga tctccttgat gaagagacaa ttccggatgt  
ggatgtagaa gaggcagttc tggggcagta tactgaagaa cctctaagtc aagagccatc  
tgtagatgct ggtgtggatt gttcatcaat tggcggggt ccacttttcc agcataaaaa  
atcacatgga aaagacaaag aaaacagagg cattaacaca ctggagagg cttaagtga  
  
agaaactaca gagcacttgg ttacaaagag cagattacct ctgcgagccc agatcaacct  
tTAAAttcact tgggggttgg caattttatt tttaaaagaa aacttaaaaa taaaacctga  
aaccaccagaa cttgagcctt gtgtatagat tttaaaagaa tatatatatc agccgggcgc  
gtggctctag ctgtaattcc agctaaactt ggaggctgag gcgggtggat tgcctgagcc  
caggagtctg agaccagcct ggccaacgtg cgctaaaacc tctgtctctg ttaaaaatta  
tccggggcgtg gtgggcacac tctgtactc ccagctactg gggaggctga ggcacagaa  
gcacttgaac ccaggaagcg ggttgacgt agccaaaggt acaccactac actccagcct  
gggcaacaga gcaagactcg gtctcaaaa taaaatttaa aaaagatata aggcagtact  
gtaaattcag ttgaattttg atattctacc attttctgt catccctata gttcactttg  
tatcaaatcg ggttccattt gggatttgca atgtaaatc gtatttctag ttttcataa  
aagtagtctt tttaggaatt c

FIGURE 1

MASQPNSSAKKKEEGKNIQVVVRCRPFNLAERKASAHSIVECD  
PVRKEVSVRTGGLADKSSRKTYTFDMVFGASTKQIDVYRSVVCPIIDEVIMGYNCTIF  
AYGQTGTGKTFTMEGERSPNEEYTWEEDPLAGIIPRTLHQIFEKLTONGTEFSVKVSL  
LEIYNEELFDLLNPSSDVSERLQMFDDPRNKRGVLIKGLEEITVHNKDEVYQILEKGA  
AKRTTAATLMNAYSSRSHSVFSVTIHMKETTIDGEELVKIGKLNLDLAGSENIGRSG  
AVDKRAREAGNINQSLTLGRVITALVERTPHVPYRESKLTRILQDSLGGRTRTSIIA  
TISPASLNLEETLSTLEYAHRANKILNKPEVNQKLTKKALIKKEYTEEIERLKRDAAA  
REKNGVYI SEENFRVMSGKLTVQEEQIVELIEKIGAVEEELNRVTLEFMDNKNELDQC  
KSDLQNKTOLETTQKHLQETKLQLVKEEYITSALESTEELHDAASKLLNTVEETTK  
DVSGLSKLDKKAQVQHNAAEQDIFGKNLNSLFNNMEELIKDGSSKQKAMLEVHKT  
FGNLLSSSVSALDTITTVALGSLTSIPENVSTHVSQIFNMILKEQSLAAESKTVLQEL  
INVLKTDLLSSLEMILSPTVVVSILKINSQKHFKTSLTVADKIEDQKKRNSDGFLSI  
LCNNLHELQENTICSLVESQKQCGNLTEDLKTIKQTHSQELCKLMNLWTERFCALEEK  
CENIQKPLSSVQENIQKSKDIVNKMTEHSQKFCADSDGFSQELRNFNQEGTKLVEES  
VKHSDKLNGLNLEKISQETEQRCESLNTRTVYFSEQWVSSLNREQELHNLLEVVSQCC  
EASSSDITEKSDGRKAAHEKQHNIFLDQMTIDEDKLIQNLNELNETIKIGLTKLNCFL  
EQDLKLDIPTGTTPQRKSYLYPSTLVRTEPREHLLDQLKRKQPELLMMLNCSENNKEE  
TIPDQDVVEEAVLGQYTEEPLSQEPSVDAGVDCSSIGGVPPFQHKKSHGDKENRGINT  
LERSKVEETTEHLVTKSRLPLRAQINL

FIGURE 2

FIGURE 4

FIGURE 5

atggcggtggcagccaaattcgtctgcgaagaagaaaggagagaaggggaagaacatccagggtgggtgggtgaga  
tgcagaccattttaatttggcagagcggaaagctagcgcccattcaatagtagaatgtgatccctgtacgaaaa  
gaagttagtgtagaactggaggattggctgacaagagctcaaggaaaacatacacttttgatatgggtgtct  
ggagcatctactaaacagattgatgtttaccgaagtggtgtttgtccaattctggatgaagttattatgggc  
tataattgcactatctttgctgtatggccaaactggcactggaaaaactttacaattggaagggtgaaagggtca  
cctaattgaagagtatacctgggaaggagatcccttggctgggtataaattccacgtacccttcatcaaatTTTT  
gagaaacttactgataaactgtactgaattttcagtc aaagtgtctctgttggagatctataatgaagagcct  
tttgatcttcttaatccatcatctgatgtttctgagagactacagatgtttgatgatccccgtaaacaagaga  
ggagtgataattaaagggtttagaagaaattacagtacacaacaaggatgaagtcctatcaaatTTTtagaaaag  
ggggcagcaaaaaggacaactgcagctactctgatgaatgcatactctagtcgttcccactcagttttctct  
gttacaatacataatgaaagaaactacgattgatggagaagagcctgttataaatcggaaagtgtgaacttgggt  
gatcttgcagggaagtga aaacattggccgttctggagctgttgataagagagctcgggaagctggaaatata  
aatcaatccctgttgactttgggaagggtcattactgcccttgttagaaagaacacctcatgttctcttatcga  
gaatctaaactaaactagaatctctccgggattctcttggagggcgtagaagaacatctataattgcaacaatt  
tctcctgcattctcaatcttggagaaactctgagtagacattggaatatgctcatagagcaaaagaacatttg  
aataagcctgaagtgaatcagaaactcaccaaaaaagctcttattaaaggagtatacggaggagatagaacgt  
tataaacgagatcttctgtcagcccgtagaaaaaatggagtgatatttctgaaagaaaatttttagagtcattg  
agtggaaaattaaactgttcaagaagagcagattgtagaattgattgaaaaaattgggtgctgttgaggaggag  
ctgaatagggttcacagagttgtttatggataataaaaaatgaacttgaccagtgtaaatctgacctgc aaat  
aaaacacaagaacttgaaacactcaaaaacatttgcaagaaactaaattacaacttgttaaaagaagaatat  
atcacatcagcttggaaagtactgaggagaaactcgaagggtaccgagcagaagcttgatcagcgaaggagac  
ctgatcgaagcaccacccaccaccactga

Macqpnsakkkkeekgkniqvvrcripfnlaerkasahsivecdpvrkevsvrtggladkssrktytfamvf  
gastkqidvyrsvvcpildevimgynctifayggtgtgkftmegerspneeytweedplagiiprtlhqif  
ekltdngtefsvkvsilleiyneelfdlpsscvserlqmfdprnkrvgiikgleeitvhnkdevyqilek  
gaakrttaatlmanayssrshsvfsvtihmkettidgeelvkigklnlvdlagsenigrsgavdkrareagni  
ngslltigrvitalivertphvpyreskltrilqdslggrtrtsiiatispaslnleetlstleyahraknil  
nkpevnqkltkkalikeyteeierlkrdlaaarekngvyiseenfrvmgkltvqeequiveliekigaveee  
lnrvtelmdnkneldqcksdlqktgelettqkhiqetklqlvkeeyitsalesteeklegteqkliseed  
liehhhhh

FIGURE 6

FIGURE 3

FIGURE 9

Atggcggtgacagccaaattcgtctgccaagaagaagaggagaaggggaagaacatccaggtgggtgggtgaga  
tgcagaccatttaatttggcagagcggaagctagcgcccatccaatagtagaatgtgatcctgtacgaaaa  
gaagttagtgtagaactggaggattggctgacaagagctcaaggaaacatacacttttgatcgggtgtt  
ggagcactctactaaacagattgatgtttaccgaagtgtgtttgtccaattctggatgaagttattatgggc  
tataattgcactatctttgctgtatggccaaactggcactggaaaaacttttacaatggaaggtgaaaggtca  
cctaatgaagagtatacctgggaagaggatcccttggctgggtataatcccacgtaccctccatcaaattttt  
gagaaacttactgataatggactgaattttcagtcmaaagtgcctctgttggagatctataatgaagagctt  
tttgatcttcttaatccatcatctgatgtttctgagagactacagatgtttgatgatccccgtaacaagaga  
ggagtataaattaaaggttttagaagaaattacagtacacaacaaggatgaagtctatcaaattttagaaaaag  
ggggcagcaaaaaggacaaactgcagctactctgatgaatgcatactctagtcttccactcagctttctct  
gttacaatacatatgaagaagaaactacgattgatggagaagagcttgttaaaatcggaaggttgaacttgggt  
gatcttgcaggaagtgaaaaacattggccgttctggagctgttgataagagagctcgggaagctggaaatata  
aatcaatccctgttgacttgggaaggggtcattactgccttctgtagaagaacacccatgttcccttatcga  
gaatctaaactaactagaatcctccaggattctcttggagggcggtacaagaacatctataattgcaacaatt  
ctctctgcatctctcaatcttgaggaaactcagagtacattggaatatgtccatagagcaagaacatactg  
aataagcctgaagrgaatcagaaatag

maggnssakkkeekgknigvvrzcrpfolaerkasahsivecdpvrkevsvrtggladkssrkytfdmfv  
gastkqidvyrsvvcpldevimgynctifayggtgtgktftmegerspneeytweedplagiiprtlhqif  
ekldngtfevsvkslleiyneelfdlpssdvserlqmfddprnkrgviikgleeitvhnkdevyqilek  
gaakrttaatlmmayssrshsvfsvtihmketridgeelvkigklnlvdlagsenigrsgavdkrareagni  
ngslltgrvitalvertphvpyreskltrilqdslggrtrtsiatispaslnleettstleyahraknil  
nkpevnqk

FIGURE 10




FIGURE 11



## METHODS OF SCREENING FOR MODULATORS OF CELL PROLIFERATION AND METHODS OF DIAGNOSING CELL PROLIFERATION STATES

### FIELD OF THE INVENTION

**[0001]** The invention relates to the use of nucleic acids encoding the kinesin KSP and their gene products to identify modulators of cell proliferation and their use in diagnosis, prognosis and treatment of cell proliferation states and disorders, for example cancer.

### BACKGROUND OF THE INVENTION

**[0002]** Cancer is the second-leading cause of death in industrialized nations. Effective therapeutics include the taxanes and vinca alkyls, agents which act on microtubules. Microtubules are the primary structural element of the mitotic spindle. The mitotic spindle is responsible for distribution of replicate copies of the genome to each of the two daughter cells that result from cell division. It is presumed that it is the disruption of the mitotic spindle by these drugs that results in inhibition of cancer cell division, and also induction of cancer cell death. However, microtubules also form other types of cellular structures, including tracks for intracellular transport in nerve processes. Therefore, the taxanes have side effects that limit their usefulness. Furthermore, taxanes and vinca alkaloids specifically target microtubule polymerization dynamics. There are additional dynamics of the mitotic spindle that these compounds do not target.

**[0003]** Therefore, it is desirable to identify agents and compositions which are specific and therapeutically effective against cancer. It is further desirable to identify agents and compositions which have a novel mechanism of action. It is further desirable to provide methods of diagnosis of hyper or hypo proliferation disorders. Additionally, it is desirable to identify agents and compositions which modulate cell proliferation. Cell proliferation modulation is desirable in a number of cases as discussed below, for example, for treatment of any hyper or hypo proliferation disorder, wound healing, transplantation procedures and for use in the agricultural arena. It is thus desirable to provide such methods of treatment. Moreover, it is desirable to provide assays to quickly identify such agents and compositions.

### SUMMARY OF THE INVENTION

**[0004]** Provided herein are assays for screening for bioactive agents which affect cell proliferation. Also provided herein are methods of diagnosing proliferation states in a cell which are useful for identifying cell proliferation disorders such as cancer. Also provided are methods of prognosis and methods of treatment including treatment for cancer. As is further described below, a number of compositions and methods are provided.

**[0005]** In one aspect, a method of screening drug candidates is provided. In one embodiment, said method comprises providing a cell that expresses recombinant human KSP or a fragment thereof and adding a drug candidate to said cell under conditions where the drug candidate is taken up by the cell. The method further includes determining the effect of said drug candidate on the bioactivity of said recombinant human KSP. The bioactivity of recombinant human KSP, or particularly the changes in the presence of a drug candidate, can be determined by assays such as those for determining

cellular proliferation, cellular viability, and cellular morphology. In a further aspect of the invention, any changes in bioactivity of recombinant human KSP can be determined by assays for determining changes in the mitotic spindle, particularly inhibition of mitosis, and ATP hydrolysis. The methods herein may also determine the bioactivity of recombinant human KSP in the presence and absence of candidate agents by performing assays determining the effect on apoptosis and necrosis.

**[0006]** The methods provided herein can be performed on single individual cells or a population of cells. The cell can be any kind of cell including but not limited to a lymphocyte, cancer cell or an endothelial cell. In one aspect, wherein cancer cells are utilized, cancer growth or inhibition can be determined, and wherein endothelial cells are utilized, angiogenesis or inhibition thereof can be determined.

**[0007]** In another aspect of the invention, a method of screening for a bioactive agent capable of binding to a cellular proliferation protein is provided. Preferably, the cellular proliferation protein is human KSP or a fragment thereof. In one embodiment, said method comprises combining said cellular proliferation protein and a candidate bioactive agent, wherein said candidate bioactive agent is an exogenous agent, and determining the binding of said candidate agent to said cellular proliferation protein.

**[0008]** In a further aspect herein, a method of screening for a candidate protein capable of binding to a cellular proliferation protein, wherein said cellular proliferation protein is KSP or a fragment thereof, is provided. In a preferred method, said method comprises combining a nucleic acid encoding said cellular proliferation protein and a nucleic acid encoding a candidate protein, wherein an identifiable marker is expressed wherein said candidate protein binds to said cellular proliferation protein.

**[0009]** Also provided herein is a method for screening for a bioactive agent capable of interfering with the binding of a cellular proliferation protein, wherein said cellular proliferation protein is KSP or a fragment thereof, and an antibody which binds to said cellular proliferation protein. In one embodiment, the method comprises combining a cellular proliferation protein, wherein said cellular proliferation protein is KSP or fragment thereof, a candidate bioactive agent and an antibody which binds to said cellular proliferation protein and determining the binding of said cellular proliferation protein and said antibody.

**[0010]** In a further aspect of the invention herein, a method for screening for a bioactive agent capable of modulating the activity of a cellular proliferation protein, wherein said cellular proliferation protein is human KSP or a fragment thereof, is provided. In one aspect, said method comprises combining said cellular proliferation protein and a candidate bioactive agent, wherein said candidate bioactive agent is an exogenous agent, and determining the effect of said candidate agent on the activity of said cellular proliferation protein.

**[0011]** Also provided herein is a method of screening drug candidates comprising providing a cell that expresses KSP, adding a drug candidate to said cell, and determining the effect of said drug candidate on the expression of KSP. In a further aspect the method includes comparing the level of expression in the absence of said drug candidate to the level of expression in the presence of said drug candidate, wherein the concentration of said drug candidate can vary when present, and wherein said comparison can occur after addition or removal of the drug candidate. In a preferred embodiment, the

expression of said KSP is decreased as a result of the introduction of the drug candidate. Preferably, the cell utilized is a tumor cell.

**[0012]** In a further aspect, a method of evaluating the effect of a candidate drug on cellular proliferation (a candidate cellular proliferation drug) is provided which comprises administering said drug to a patient, removing a cell sample from said patient, and determining the expression profile of said cell, wherein said expression profile includes a KSP gene. In another aspect, the method includes comparing said expression profile to an expression profile of a healthy individual.

**[0013]** In another aspect herein, a method of diagnosing a hyper-proliferative disorder in an individual is provided herein comprising determining the level of expression a KSP gene in an individual and comparing said level to a standard or control level of expression, wherein an increase indicates that the individual has a hyper-proliferative disorder, such as, but not limited to, cancer.

**[0014]** Also provided herein is a method of evaluating the effect of a candidate cellular proliferation drug comprising administering said drug to a patient wherein said patient has cancer and has been identified as expressing KSP at a level higher than an individual not having cancer, removing a cell sample from said patient, and determining the effect on KSP activity, wherein said KSP activity is mitosis.

**[0015]** In the methods provided herein, the cells can come from a variety of sources. For example, samples can be from, but are not limited to, a blood sample, a urine sample, a buccal sample, a PAP smear, cerebral spinal fluid, and any tissue including, breast tissue, lung tissue and colon tissue. In one embodiment, the patient has cancer.

**[0016]** Also provided herein is a method for inhibiting cellular proliferation, said method comprising administering to a cell a composition comprising an antibody to KSP, wherein said antibody is conjugated to a ligand. In one aspect, the ligand of the antibody is tumor cell specific. In another aspect, the ligand facilitates said antibody entry to said cell. Moreover, the antibody can be a humanized antibody. The methods of inhibition can be performed in vitro on cells or in vivo on an individual. In one embodiment, the cells are cancerous. In a further embodiment, the individual has cancer. Another method of inhibiting cellular proliferation in a cell or individual is provided herein which comprises administering to a cell or individual a composition comprising antisense molecules to KSP.

**[0017]** In yet another embodiment herein, a method for inhibiting cellular proliferation is provided which comprises administering to a cell a composition comprising an inhibitor of KSP. In one embodiment, the inhibitor is of human KSP or a fragment thereof. In one embodiment, the inhibitor is specific to human KSP. In one embodiment, KSP inhibitors are any agent which disrupts or inhibits KSP activity as further described herein. In one aspect of the invention, the inhibitor of KSP is a small molecule as further defined herein. Generally, small molecules have a molecular weight of between 50 kD and 2000 kD, and in some cases, less than 1500 kD, or less than 1000 kD or less than 500 kD. Examples of KSP inhibitors include but are not limited to small molecules, ribozymes, antisense molecules and antibodies. KSP inhibitors are further described herein and in the application filed Oct. 27, 1999, entitled Methods and Compositions Utilizing Quinazolinones (serial number not yet received, named inventor Jeffrey T. Finer), incorporated by reference in its

entirety. The composition which is administered to a cell further comprises an acceptable pharmaceutical carrier in one embodiment. The composition can have a variety of formulations, including, but not limited to those for parental, oral or topical administration.

**[0018]** The methods of inhibiting cellular proliferation can be performed in vitro or in vivo. More particularly, the compositions can be administered to cells in vitro or in an individual. The individual may have a disease or be at risk for disease. Disease states which can be treated by the methods herein are further described below. In one case, the individual has cancer or is at risk for restenosis. The cell can be any cell, preferably a cancer cell. Other preferred cell types include but are not limited to endothelial cells and metastatic cancer cells. In one embodiment, the method of inhibiting by the KSP inhibitor is by disruption of mitosis or induction of apoptosis.

**[0019]** In a further aspect of the invention, a biochip comprising a nucleic acid segment from KSP, wherein said biochip comprises fewer than 1000 nucleic acid probes, is provided. Methods of screening and diagnosing conditions with said biochip are also provided herein.

**[0020]** Other aspects of the invention will become apparent to the skilled artisan by the following description of the invention.

#### DETAILED DESCRIPTION OF THE FIGURES

**[0021]** FIG. 1 shows a cDNA sequence for human KSP, GenBank accession number X85137, wherein the start and stop codons are shown underlined and in bold, beginning at positions 11 and 3182, respectively.

**[0022]** FIG. 2 shows an amino acid sequence encoding human KSP.

**[0023]** FIG. 3 shows a nucleic acid sequence encoding a fragment of KSP, termed KSPL360 herein. Portions differing from the sequences of FIGS. 1 and 2 are indicated in bold typeface and are underlined. Residues at the C-terminus include a myc epitope and a 6-histidine tag.

**[0024]** FIG. 4 shows an amino acid sequence encoding KSPL360.

**[0025]** FIG. 5 shows a nucleic acid sequence encoding a fragment of KSP, termed KSP-K491 herein. Portions differing from the sequences of FIGS. 1 and 2 are indicated in bold typeface and are underlined. Residues at the C-terminus include a myc epitope and a 6-histidine tag.

**[0026]** FIG. 6 shows an amino acid sequence encoding KSP-K491.

**[0027]** FIG. 7 shows a nucleic acid sequence encoding a fragment of KSP, termed KSP-S553 herein. Portions differing from the sequences of FIGS. 1 and 2 are indicated in bold typeface and are underlined. Residues at the C-terminus include a myc epitope and a 6-histidine tag.

**[0028]** FIG. 8 shows an amino acid sequence encoding KSP-S553.

**[0029]** FIG. 9 shows a nucleic acid sequence encoding a fragment of KSP, termed KSP-K368 herein. Portions differing from the sequences of FIGS. 1 and 2 are indicated in bold typeface and are underlined.

**[0030]** FIG. 10 shows an amino acid sequence encoding KSP-K368.

**[0031]** FIG. 11 is a graph showing KSP mRNA levels in matched normal and tumor tissue from breast, lung and colon. mRNA levels were measured by quantitative PCR relative to a standard. The relative magnitudes of overexpression in each tumor sample relative to the matched normal tissue are dis-

played above each pair. All values are normalized to the level of KSP mRNA expression observed in cultured HeLa cells.

#### DETAILED DESCRIPTION OF THE INVENTION

**[0032]** Provided herein are assays for screening for bioactive agents which affect cell proliferation. Also provided herein are methods of diagnosing proliferation states in a cell which are useful for identifying cell proliferation disorders such as cancer. Also provided are methods of prognosis and methods of treatment including treatment for cancer. As is further described below, a number of compositions and methods are provided.

**[0033]** In one aspect, the assays or methods of diagnosis provided herein include the use of a cellular proliferation protein or nucleic acid. The terms “cell proliferation” and “cellular proliferation” are used herein interchangeably. Additionally, the cellular proliferation protein and nucleic acid can be referred to herein as “cellular proliferation sequences” wherein the context will indicate whether the sequence is an amino acid sequence, nucleic acid sequence, or either.

**[0034]** In a preferred embodiment, the cellular proliferation sequence is KSP. KSP belongs to an evolutionarily conserved kinesin subfamily of plus end-directed microtubule motors that assemble into bipolar homotetramers consisting of anti-parallel homodimers. During mitosis KSP associates with microtubules of the mitotic spindle. Microinjection of antibody directed against KSP into cells prevents spindle pole separation during prometaphase, giving rise to monopolar spindles and causing mitotic arrest. KSP and related kinesins bundle antiparallel microtubules and slide them relative to one another, thus forcing the two spindle poles apart. KSP may also mediate in anaphase B spindle elongation and focussing of microtubules at the spindle pole.

**[0035]** Human KSP has been reported on (also termed HsEg5). Galgio, et al., *J. Cell Biol.*, 135(2):399-414 (1996); Kaiser, et al., *JBC*, 274(27):18925-31 (1999); Blangy, et al., *Cell*, 83:1159-69 (1995); Blangy, et al., *J Biol. Chem.*, 272: 19418-24 (1997); Slangy, et al., *Cell Motil Cytoskeleton*, 40:174-82 (1998); Whitehead, et al., *Arthritis Rheum.*, 39:1635-42 (1996); GenBank accession numbers: X85137, NM\_004523 and U37426. Moreover, a fragment of the KSP gene (TRIPS) has been reported on. Lee, et al., *Mol. Endocrinol.*, 9:243-54 (1995); GenBank accession number L40372. Also see, Whitehead and Rattner, *J. Cell Sci.*, 111: 2551-61 (1998).

**[0036]** *Xenopus* KSP homologs (Eg5) have also been reported on. Walczak, et al., *Curr Biol.*, 8(16):903-13 (1998); Le Guellec, et al., *Mol. Cell Biol.*, 11(6):3395-8 (1991); Sawin, et al., *Nature*, 359:540-3 (1992); Sawin and Mitchison, *Mol Biol Cell*, 5:217-26 (1994); Sawin and Mitchison, *PNAS*, 92:4289-93 (1995); Kapoor and Mitchison, *PNAS*, 96:9106-11 (1999); Lockhart and Cross, *Biochemistry*, 35(7):2365-73 (1996); Crevel, et al., *J. Mol. Biol.*, 273:160-170 (1997). Additionally, *Drosophila* KLP61F/KRP130 has been reported on. Heck, et al., *J Cell Biol*, 123:665-79 (1993); Cole, et al., *J. Biol. Chem.*, 269(37):22913-6 (1994); Barton, et al., *Mol. Biol. Cell*, 6:1563-74 (1995).

**[0037]** In the preferred embodiment herein, a sequence as shown in the figures is utilized. As indicated herein, in some embodiments a fragment of KSP is utilized. Preferred protein fragments are shown in FIGS. 2, 4, 6, and 8. In one embodiment, the cellular proliferation fragment shown in FIG. 4 is preferred. Preferred fragments of KSP have kinesin activity

as further described below. Moreover, in one embodiment, KSP peptides or fragments have at least one, and preferably at least two epitope tags. In a preferred embodiment, a KSP fragment comprises a myc epitope and a histidine tag.

**[0038]** In another preferred embodiment herein, the cellular proliferation protein is non-glycosylated. For example, in one embodiment the protein is, for example, human, expressed in bacteria, for example, *E. Coli*. Moreover, phosphorylation and/or methylation of KSP as used herein may differ from KSP as found in its native form within a cell.

**[0039]** Thus, while it is preferred that the cellular proliferation sequences are from humans, sequences from other organisms may be useful in animal models of disease and drug evaluation; thus, in alternative embodiments, other sequences are provided such as from vertebrates, including mammals, including rodents (rats, mice, hamsters, guinea pigs, etc.), primates, farm animals (including sheep, goats, pigs, cows, horses, etc), *Xenopus*, and *Drosophila*.

**[0040]** In another embodiment, the sequences are naturally-occurring allelic variants of the sequences set forth in the figures. In another embodiment, the sequences are sequence variants as further described herein.

**[0041]** In one embodiment, a cellular proliferation sequence can be initially identified by substantial nucleic acid and/or amino acid sequence homology to the cellular proliferation sequences outlined herein. Such homology can be based upon the overall nucleic acid or amino acid sequence, and is generally determined as outlined below, using either homology programs or hybridization conditions.

**[0042]** Thus, in one embodiment, a nucleic acid is a “cellular proliferation nucleic acid” if the overall homology of the nucleic acid sequence to the nucleic acid sequences of FIG. 1, 3, 5, 7 or 9 (the nucleic acid figures) is preferably greater than about 75%, more preferably greater than about 80%, even more preferably greater than about 85% and most preferably greater than 90%. In some embodiments the homology will be as high as about 93 to 95 or 98%. Homology as used herein is in reference to sequence similarity or identity, with identity being preferred. This homology will be determined using standard techniques known in the art, including, but not limited to, the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *PNAS USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux et al., *Nucl. Acid Res.* 12:387-395 (1984), preferably using the default settings, or by inspection.

**[0043]** One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. It can also plot a tree showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J. Mol. Evol.* 35:351-360 (1987); the method is similar to that described by Higgins & Sharp *CABIOS* 5:151-153 (1989). Useful PILEUP parameters including a default gap weight of 3.00, a default gap length weight of 0.10, and weighted end gaps.

**[0044]** Another example of a useful algorithm is the BLAST algorithm, described in Altschul et al., *J. Mol. Biol.*

215, 403-410, (1990) and Karlin et al., PNAS USA 90:5873-5787 (1993). A particularly useful BLAST program is the WU-BLAST-2 program which was obtained from Altschul et al., *Methods in Enzymology*, 266: 460-480 (1996); <http://blast.wustl.edu/blast/REACRCE.html>). WU-BLAST-2 uses several search parameters, most of which are set to the default values. The adjustable parameters are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched; however, the values may be adjusted to increase sensitivity. A % amino acid sequence identity value is determined by the number of matching identical residues divided by the total number of residues of the "longer" sequence in the aligned region. The "longer" sequence is the one having the most actual residues in the aligned region (gaps introduced by WU-Blast-2 to maximize the alignment score are ignored).

**[0045]** Thus, "percent (%) nucleic acid sequence identity" is defined as the percentage of nucleotide residues in a candidate sequence that are identical with the nucleotide residues of the sequence shown in the nucleic acid figures. A preferred method utilizes the BLASTN module of WU-BLAST-2 set to the default parameters, with overlap span and overlap fraction set to 1 and 0.125, respectively.

**[0046]** The alignment may include the introduction of gaps in the sequences to be aligned. In addition, for sequences which contain either more or fewer nucleosides than those of the nucleic acid figures, it is understood that the percentage of homology will be determined based on the number of homologous nucleosides in relation to the total number of nucleosides. Thus, for example, homology of sequences shorter than those of the sequences identified herein and as discussed below, will be determined using the number of nucleosides in the shorter sequence.

**[0047]** In one embodiment, the cellular proliferation nucleic acid is determined through hybridization studies. Thus, for example, nucleic acids which hybridize under high stringency to the nucleic acid sequences identified in the figures, or a complement, are considered a cellular proliferation sequence in one embodiment herein. High stringency conditions are known in the art; see for example Maniatis et al., *Molecular Cloning: A Laboratory Manual*, 2d Edition, 1989, and *Short Protocols in Molecular Biology*, ed. Ausubel, et al., both of which are hereby incorporated by reference. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, stringent conditions are selected to be about 5-10° C. lower than the thermal melting point (T<sub>m</sub>) for the specific sequence at a defined ionic strength pH. The T<sub>m</sub> is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at T<sub>m</sub>, 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion

concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g. 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g. greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

**[0048]** In another embodiment, less stringent hybridization conditions are used; for example, moderate or low stringency conditions may be used, as are known in the art; see Maniatis and Ausubel, *supra*, and Tijssen, *supra*.

**[0049]** In addition, in one embodiment the cellular proliferation nucleic acid sequences of the invention are fragments of larger genes, i.e. they are nucleic acid segments. "Genes" in this context includes coding regions, non-coding regions, and mixtures of coding and non-coding regions. Accordingly, as will be appreciated by those in the art, using the sequences provided herein, additional sequences of the cellular proliferation genes can be obtained, using techniques well known in the art for cloning either longer sequences or the full length sequences; see Maniatis et al., and Ausubel, et al., *supra*, hereby expressly incorporated by reference.

**[0050]** Once the cellular proliferation nucleic acid is identified, it can be cloned and, if necessary, its constituent parts recombined to form the entire cellular proliferation nucleic acid. Once isolated from its natural source, e.g., contained within a plasmid or other vector or excised therefrom as a linear nucleic acid segment, the recombinant cellular proliferation nucleic acid can be further-used as a probe to identify and isolate other cellular proliferation nucleic acids, for example additional coding regions. It can also be used as a "precursor" nucleic acid to make modified or variant cellular proliferation nucleic acids and proteins. "Recombinant" as used herein refers to a nucleic acid or protein which is not in its native state. For example, the nucleic acid can be genetically engineered, isolated, inserted into a man-made vector or be in a cell wherein it is not natively expressed in order to be considered recombinant.

**[0051]** In another aspect, the cellular proliferation nucleic acid and protein sequences are differentially expressed in cells having varying states of cellular proliferation, including cancer cells which over proliferate compared to non cancerous cells. As outlined below, cellular proliferation sequences include those that are up-regulated (i.e. expressed at a higher level) during cellular proliferation, as well as those that are down-regulated (i.e. expressed at a lower level) in cellular proliferation. In a preferred embodiment, the cellular proliferation sequences are upregulated during cellular proliferation in their native state, i.e., without the administration of modulators or therapeutics.

**[0052]** The term "nucleic acid" refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. Unless specifically limited, the term encompasses nucleic acids containing known analogues of natural nucleotides which have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions) and complementary sequences and as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer et al.,

*Nucleic Acid Res.* 19:5081 (1991); Ohtsuka et al., *J. Biol. Chem.* 260:2605-2608 (1985); Cassol et al., 1992; Rossolini et al., *Mol. Cell. Probes* 8:91-98 (1994)). The term nucleic acid is used interchangeably with gene, cDNA, and mRNA encoded by a gene.

**[0053]** The cellular proliferation nucleic acids of the present invention are used in several ways. In a preferred embodiment, cellular proliferation nucleic acids encoding cellular proliferation proteins are used to make a variety of expression vectors to express cellular proliferation proteins which can then be used in screening assays, as described below. The expression vectors may be either self-replicating extrachromosomal vectors or vectors which integrate into a host genome. Generally, these expression vectors include transcriptional and translational regulatory nucleic acid operably linked to the nucleic acid encoding the cellular proliferation protein. The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

**[0054]** Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice. The transcriptional and translational regulatory nucleic acid will generally be appropriate to the host cell used to express the cellular proliferation protein; for example, transcriptional and translational regulatory nucleic acid sequences from *Bacillus* are preferably used to express the cellular proliferation protein in *Bacillus*. Numerous types of appropriate expression vectors, and suitable regulatory sequences are known in the art for a variety of host cells.

**[0055]** In general, the transcriptional and translational regulatory sequences may include, but are not limited to, promoter sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, and enhancer or activator sequences. In a preferred embodiment, the regulatory sequences include a promoter and transcriptional start and stop sequences.

**[0056]** Promoter sequences encode either constitutive or inducible promoters. The promoters may be either naturally occurring promoters or hybrid promoters. Hybrid promoters, which combine elements of more than one promoter, are also known in the art, and are useful in the present invention.

**[0057]** In addition, the expression vector may comprise additional elements. For example, the expression vector may have two replication systems, thus allowing it to be maintained in two organisms, for example in mammalian or insect cells for expression and in a procaryotic host for cloning and

amplification. Furthermore, for integrating expression vectors, the expression vector contains at least one sequence homologous to the host cell genome, and preferably two homologous sequences which flank the expression construct. The integrating vector may be directed to a specific locus in the host cell by selecting the appropriate homologous sequence for inclusion in the vector. Constructs for integrating vectors are well known in the art.

**[0058]** In addition, in a preferred embodiment, the expression vector contains a selectable marker gene to allow the selection of transformed host cells. Selection genes are well known in the art and will vary with the host cell used.

**[0059]** The cellular proliferation proteins of the present invention can be produced by culturing a host cell transformed with an expression vector containing nucleic acid encoding a cellular proliferation protein, under the appropriate conditions to induce or cause expression of the cellular proliferation protein. The conditions appropriate for cellular proliferation protein expression will vary with the choice of the expression vector and the host cell, and will be easily ascertained by one skilled in the art through routine experimentation. For example, the use of constitutive promoters in the expression vector will require optimizing the growth and proliferation of the host cell, while the use of an inducible promoter requires the appropriate growth conditions for induction. In addition, in some embodiments, the timing of the harvest is important. For example, the baculoviral systems used in insect cell expression are lytic viruses, and thus harvest time selection can be crucial for product yield.

**[0060]** Appropriate host cells include yeast, bacteria, archaeobacteria, fungi, and insect and animal cells, including mammalian cells. Of particular interest are *Drosophila melanogaster* cells, *Saccharomyces cerevisiae* and other yeasts, *E. coli*, *Bacillus subtilis*, Sf9 cells, C129 cells, 293 cells, *Neurospora*, BHK, CHO, COS, HeLa cells, THP1 cell line (a macrophage cell line) and human cells and cell lines.

**[0061]** In one embodiment, the cellular proliferation proteins are expressed in mammalian cells. Mammalian expression systems are also known in the art, and include retroviral systems. A preferred expression vector system is a retroviral vector system such as is generally described in PCT/US97/01019 and PCT/US97/01048, both of which are hereby expressly incorporated by reference. Of particular use as mammalian promoters are the promoters from mammalian viral genes, since the viral genes are often highly expressed and have a broad host range. Examples include the SV40 early promoter, mouse mammary tumor virus LTR promoter, adenovirus major late promoter, herpes simplex virus promoter, and the CMV promoter. Typically, transcription termination and polyadenylation sequences recognized by mammalian cells are regulatory regions located 3' to the translation stop codon and thus, together with the promoter elements, flank the coding sequence. Examples of transcription terminator and polyadenylation signals include those derived from SV40.

**[0062]** The methods of introducing exogenous nucleic acid into mammalian hosts, as well as other hosts, is well known in the art, and will vary with the host cell used. Techniques include dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, viral infection, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

**[0063]** In a preferred embodiment, cellular proliferation proteins are expressed in bacterial systems. Bacterial expression systems are well known in the art. Promoters from bacteriophage may also be used and are known in the art. In addition, synthetic promoters and hybrid promoters are also useful; for example, the tac promoter is a hybrid of the trp and lac promoter sequences. Furthermore, a bacterial promoter can include naturally occurring promoters of non-bacterial origin that have the ability to bind bacterial RNA polymerase and initiate transcription. In addition to a functioning promoter sequence, an efficient ribosome binding site is desirable. The expression vector may also include a signal peptide sequence that provides for secretion of the cellular proliferation protein in bacteria. The protein is either secreted into the growth media (gram-positive bacteria) or into the periplasmic space, located between the inner and outer membrane of the cell (gram-negative bacteria). The expression vector may also include an epitope tag providing for affinity purification of the cellular proliferation protein. The bacterial expression vector may also include a selectable marker gene to allow for the selection of bacterial strains that have been transformed. Suitable selection genes include genes which render the bacteria resistant to drugs such as ampicillin, chloramphenicol, erythromycin, kanamycin, neomycin and tetracycline. Selectable markers also include biosynthetic genes, such as those in the histidine, tryptophan and leucine biosynthetic pathways. These components are assembled into expression vectors. Expression vectors for bacteria are well known in the art, and include vectors for *Bacillus subtilis*, *E. coli*, *Streptococcus cremoris*, and *Streptococcus lividans*, among others. The bacterial expression vectors are transformed into bacterial host cells using techniques well known in the art, such as calcium chloride treatment, electroporation, and others.

**[0064]** In one embodiment, cellular proliferation proteins are produced in insect cells. Expression vectors for the transformation of insect cells, and in particular, baculovirus-based expression vectors, are well known in the art.

**[0065]** In another embodiment, cellular proliferation protein is produced in yeast cells. Yeast expression systems are well known in the art, and include expression vectors for *Saccharomyces cerevisiae*, *Candida albicans* and *C. maltose*, *Hansenula polymorpha*, *Kluyveromyces fragilis* and *K. lactis*, *Pichia guilliermondii* and *P. pastoris*, *Schizosaccharomyces pombe*, and *Yarrowia lipolytica*.

**[0066]** The cellular proliferation protein may also be made as a fusion protein, using techniques well known in the art. Thus, for example, for the creation of monoclonal antibodies, if the desired epitope is small, the cellular proliferation protein may be fused to a carrier protein to form an immunogen. Alternatively, the cellular proliferation protein may be made as a fusion protein to increase expression, or for other reasons. For example, when the cellular proliferation protein is a cellular proliferation peptide, the nucleic acid encoding the peptide may be linked to other nucleic acid for expression purposes.

**[0067]** In one embodiment, the cellular proliferation nucleic acids, proteins and antibodies of the invention are labeled. By "labeled" herein is meant that a compound has at least one element, isotope or chemical compound attached to enable the detection of the compound. In general, labels fall into three classes: a) isotopic labels, which may be radioactive or heavy isotopes; b) immune labels, which may be antibodies or antigens; and c) colored or fluorescent dyes. The labels may be incorporated into the cellular proliferation nucleic

acids, proteins and antibodies at any position. For example, the label should be capable of producing, either directly or indirectly, a detectable signal. The detectable moiety may be a radioisotope, such as  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ , or  $^{125}\text{I}$ , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the label may be employed, including those methods described by Hunter et al., *Nature*, 144:945 (1962); David et al., *Biochemistry*, 13:1014 (1974); Pain et al., *J. Immunol. Meth.*, 40:219 (1981); and Nygren, *J. Histochem. and Cytochem.*, 30:407 (1982).

**[0068]** Accordingly, the present invention also provides cellular proliferation protein sequences. A cellular proliferation protein of the present invention may be identified in several ways. "Protein" in this sense includes proteins, polypeptides, and peptides. As will be appreciated by those in the art, the nucleic acid sequences of the invention can be used to generate protein sequences.

**[0069]** Also included within one embodiment of cellular proliferation proteins are amino acid variants of the naturally occurring sequences, as determined herein. Preferably, the variants are preferably greater than about 75% homologous to the wild-type sequence, more preferably greater than about 80%, even more preferably greater than about 85% and most preferably greater than 90%. In some embodiments the homology will be as high as about 93 to 95 or 98%. As for nucleic acids, homology in this context means sequence similarity or identity, with identity being preferred. This homology will be determined using standard techniques known in the art as are outlined above for the nucleic acid homologies. The proteins of the present invention may be shorter or longer than the wild type amino acid sequences. Thus, in a preferred embodiment, included within the definition of cellular proliferation proteins are portions or fragments of the wild type sequences. Preferred fragments have a binding domain to a modulating agent or antibody as discussed below. In addition, as outlined above, the cellular proliferation nucleic acids of the invention may be used to obtain additional coding regions, and thus additional protein sequence, using techniques known in the art.

**[0070]** In one embodiment, the cellular proliferation proteins are derivative or variant cellular proliferation proteins as compared to the wild-type sequence. That is, as outlined more fully below, the derivative cellular proliferation peptide will contain at least one amino acid substitution, deletion or insertion, with amino acid substitutions being particularly preferred. The amino acid substitution, insertion or deletion or combination thereof may occur at any residue within the cellular proliferation peptide. These variants ordinarily are prepared by site specific mutagenesis of nucleotides in the DNA encoding the cellular proliferation protein, using cassette or PCR mutagenesis or other techniques well known in the art, to produce DNA encoding the variant, and thereafter expressing the DNA in recombinant cell culture as outlined above. However, variant cellular proliferation protein fragments having up to about 100-150 residues may be prepared by in vitro synthesis using established techniques. Amino acid sequence variants are characterized by the predetermined nature of the variation, a feature that sets them apart from naturally occurring allelic or interspecies variation of the cellular proliferation protein amino acid sequence. The variants typically exhibit the same qualitative biological

activity as the naturally occurring analogue, although variants can also be selected which have modified characteristics as will be more fully outlined below.

**[0071]** While the site or region for introducing an amino acid sequence variation is predetermined, the mutation per se need not be predetermined. For example, in order to optimize the performance of a mutation at a given site, random mutagenesis may be conducted at the target codon or region and the expressed cellular proliferation variants screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, for example, M13 primer mutagenesis and PCR mutagenesis. Screening of the mutants is done using assays of cellular proliferation protein activities.

**[0072]** Amino acid substitutions are typically of single residues; insertions usually will be on the order of from about 1 to 20 amino acids, although considerably larger insertions may be tolerated. Deletions range from about 1 to about 20 residues, although in some cases deletions may be much larger.

**[0073]** Substitutions, deletions, insertions or any combination thereof may be used to arrive at a final derivative. Generally these changes are done on a few amino acids to minimize the alteration of the molecule. However, larger changes may be tolerated in certain circumstances. When small alterations in the characteristics of the cellular proliferation protein are desired, substitutions are generally made in accordance with the following chart:

CHART I

Original Residue	Exemplary Substitutions
Ala	Ser
Arg	Lys
Asn	Gln, His
Asp	Glu
Cys	Ser
Gln	Asn
Glu	Asp
Gly	Pro
His	Asn, Gln
Ile	Leu, Val
Leu	Ile, Val
Lys	Arg, Gln, Glu
Met	Leu, Ile
Phe	Met, Leu, Tyr
Ser	Thr
Thr	Ser
Trp	Tyr
Tyr	Trp, Phe
Val	Ile, Leu

**[0074]** Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative than those shown in Chart I. For example, substitutions may be made which more significantly affect: the structure of the polypeptide backbone in the area of the alteration, for example the alpha-helical or beta-sheet structure; the charge or hydrophobicity of the molecule at the target site; or the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in the polypeptide's properties are those in which (a) a hydrophilic residue, e.g. seryl or threonyl is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g. lysyl, arginyl, or histidyl, is substituted for (or by) an elec-

tronegative residue, e.g. glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g. phenylalanine, is substituted for (or by) one not having a side chain, e.g. glycine.

**[0075]** The variants typically exhibit the same qualitative biological activity and will elicit the same immune response as the naturally-occurring analogue, although variants also are selected to modify the characteristics of the cellular proliferation proteins as needed. Alternatively, the variant may be designed such that the biological activity of the cellular proliferation protein is altered.

**[0076]** Covalent modifications of cellular proliferation polypeptides are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a cellular proliferation polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues of a cellular proliferation polypeptide. Derivatization with bifunctional agents is useful, for instance, for crosslinking cellular proliferation protein to a water-insoluble support matrix or surface for use in the method for purifying anti-KSP antibodies or screening assays, as is more fully described below. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio] propioimide.

**[0077]** Other modifications include deamidation of glutamyl and asparaginyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl, threonyl or tyrosyl residues, methylation of the  $\alpha$ -amino groups of lysine, arginine, and histidine side chains [T. E. Creighton, *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

**[0078]** Another type of covalent modification of the cellular proliferation polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence cellular proliferation polypeptide, and/or adding one or more glycosylation sites that are not present in the native sequence cellular proliferation polypeptide.

**[0079]** Addition of glycosylation sites to cellular proliferation polypeptides may be accomplished by altering the amino acid sequence thereof. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence cellular proliferation polypeptide (for O-linked glycosylation sites). The cellular proliferation amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the cellular proliferation polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

**[0080]** Another means of increasing the number of carbohydrate moieties on the cellular proliferation polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in

WO 87/05330 published 11 Sep. 1987, and in Aplin and Wriston, *cellular proliferation* Crit. Rev. Biochem., pp. 259-306 (1981).

**[0081]** Removal of carbohydrate moieties present on the cellular proliferation polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., *Arch. Biochem. Biophys.*, 259:52 (1987) and by Edge et al., *Anal. Biochem.*, 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., *Meth. Enzymol.*, 138:350 (1987).

**[0082]** Another type of covalent modification of cellular proliferation comprises linking the cellular proliferation polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Pat. No. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

**[0083]** The cellular proliferation polypeptides of the present invention may also be modified in one embodiment in a way to form chimeric molecules comprising a cellular proliferation polypeptide fused to another, heterologous polypeptide or amino acid sequence. In one embodiment, such a chimeric molecule comprises a fusion of a cellular proliferation polypeptide with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. Preferred tags include the myc epitope and 6-histidine. The epitope tag is generally placed at the amino- or carboxyl-terminus of the cellular proliferation polypeptide. The presence of such epitope-tagged forms of a cellular proliferation polypeptide can be detected using an antibody against the tag polypeptide as further discussed below. Also, provision of the epitope tag enables the cellular proliferation polypeptide to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. In an alternative embodiment, the chimeric molecule may comprise a fusion of a cellular proliferation polypeptide with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule, such a fusion could be to the Fc region of an IgG molecule.

**[0084]** Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., *Mol. Cell. Biol.*, 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., *Molecular and Cellular Biology*, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., *Protein Engineering*, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., *BioTechnology*, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., *Science*, 255:192-194 (1992)]; tubulin epitope peptide [Skinner et al., *J. Biol. Chem.*, 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., *Proc. Natl. Acad. Sci. USA*, 87:6393-6397 (1990)].

**[0085]** Also included with the definition of cellular proliferation protein in one embodiment are other cellular proliferation proteins of the cellular proliferation family, and cel-

lular proliferation proteins from other organisms, which are cloned and expressed as outlined below. Thus, probe or degenerate polymerase chain reaction (PCR) primer sequences may be used to find other related cellular proliferation proteins from humans or other organisms. As will be appreciated by those in the art, particularly useful probe and/or PCR primer sequences include the unique areas of the cellular proliferation nucleic acid sequence. As is generally known in the art, preferred PCR primers are from about 15 to about 35 nucleotides in length, with from about 20 to about 30 being preferred, and may contain inosine as needed. The conditions for the PCR reaction are well known in the art.

**[0086]** In addition, as is outlined herein, cellular proliferation proteins can be made that are longer than those depicted in the figures, for example, by the elucidation of additional sequences, the addition of epitope or purification tags, the addition of other fusion sequences, etc.

**[0087]** Cellular proliferation proteins may also be identified as being encoded by cellular proliferation nucleic acids. Thus, in one embodiment, cellular proliferation proteins are encoded by nucleic acids that will hybridize to the sequences of the nucleic acid figures, or their complements, as outlined herein.

**[0088]** In a preferred embodiment, the cellular proliferation protein is purified or isolated after expression. Cellular proliferation proteins may be isolated or purified in a variety of ways known to those skilled in the art depending on what other components are present in the sample. Standard purification methods include electrophoretic, molecular, immunological and chromatographic techniques, including ion exchange, hydrophobic, affinity, and reverse-phase HPLC chromatography, and chromatofocusing. For example, the cellular proliferation protein may be purified using a standard anti-KSP antibody column. Ultrafiltration and diafiltration techniques, in conjunction with protein concentration, are also useful. For general guidance in suitable purification techniques, see Scopes, R., *Protein Purification*, Springer-Verlag, NY (1982). The degree of purification necessary will vary depending on the use of the cellular proliferation protein. In some instances no purification will be necessary.

**[0089]** The terms "isolated" "purified" or "biologically pure" refer to material that is substantially or essentially free from components which normally accompany it as found in its native state. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. A protein that is the predominant species present in a preparation is substantially purified. The term "purified" denotes that a nucleic acid or protein gives rise to essentially one band in an electrophoretic gel. Particularly, it means that the nucleic acid or protein is at least 85% pure, more preferably at least 95% pure, and most preferably at least 99% pure. In a preferred embodiment, a protein is considered pure wherein it is determined that there is no contaminating activity.

**[0090]** Once expressed and purified if necessary, the cellular proliferation proteins and nucleic acids are useful in a number of applications. In a number of methods provided herein, wherein either the nucleic acid or a protein is used, a candidate bioactive agent is used to determine the effect on the cellular proliferation sequence, cellular proliferation, cancer, etc., as further discussed below.

**[0091]** In preferred embodiments, the bioactive agents modulate the cellular proliferation sequences or expression



profiles provided herein. In a particularly preferred embodiment, the candidate agent suppresses a cellular proliferation phenotype, for example to inhibit proliferation, inhibit tumor growth, or to a normal tissue fingerprint as further discussed below. Similarly, the candidate agent preferably suppresses a severe cellular proliferation phenotype. Suppression might take the form of cell or tumor growth arrest, with continued viability. Alternatively, suppression may take the form of inducing cell death of cells, thereby eliminating proliferation. As further discussed below, preferred bioactive agents are identified which cause cell death selectively of tumor cells or proliferating cells. Generally a plurality of assay mixtures are run in parallel with different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, i.e., at zero concentration or below the level of detection.

**[0092]** The term “candidate bioactive agent” or “drug candidate” or grammatical equivalents as used herein describes any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, purine analog, etc., to be tested for bioactive agents that are capable of directly or indirectly altering either the cellular proliferation phenotype or the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases, alteration of cellular proliferation protein binding and/or activity is screened. In the case where protein binding or activity is screened, preferred embodiments exclude molecules already known to bind to that particular protein, for example, polymer structures such as microtubules, and energy sources such as ATP. Preferred embodiments of assays herein include candidate agents which do not bind the cellular proliferation protein in its endogenous native state termed herein as “exogenous” agents. In another preferred embodiment, exogenous agents further exclude antibodies to KSP.

**[0093]** Candidate agents can encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Small molecules are further defined herein as having a molecular weight of between 50 kD and 2000 kD. In another embodiment, small molecules have a molecular weight of less than 1500, or less than 1200, or less than 1000, or less than 750, or less than 500 kD. In one embodiment, a small molecule as used herein has a molecular weight of about 100 to 200 kD. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Particularly preferred are peptides.

**[0094]** Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Addition-

ally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification to produce structural analogs.

**[0095]** In a preferred embodiment, the candidate bioactive agents are proteins. By “protein” herein is meant at least two covalently attached amino acids, which includes proteins, polypeptides, oligopeptides and peptides. The protein may be made up of naturally occurring amino acids and peptide bonds, or synthetic peptidomimetic structures. Thus “amino acid”, or “peptide residue”, as used herein means both naturally occurring and synthetic amino acids. For example, homo-phenylalanine, citrulline and noreucine are considered amino acids for the purposes of the invention. “Amino acid” also includes imino acid residues such as proline and hydroxyproline. The side chains may be in either the (R) or the (S) configuration. In the preferred embodiment, the amino acids are in the (S) or L-configuration. If non-naturally occurring side chains are used, non-amino acid substituents may be used, for example to prevent or retard in vivo degradations.

**[0096]** In a preferred embodiment, the candidate bioactive agents are naturally occurring proteins or fragments of naturally occurring proteins. Thus, for example, cellular extracts containing proteins, or random or directed digests of proteinaceous cellular extracts, may be used. In this way libraries of procaryotic and eucaryotic proteins may be made for screening in the methods of the invention. Particularly preferred in this embodiment are libraries of bacterial, fungal, viral, and mammalian proteins, with the latter being preferred, and human proteins being especially preferred.

**[0097]** In a preferred embodiment, the candidate bioactive agents are peptides of from about 5 to about 30 amino acids, with from about 5 to about 20 amino acids being preferred, and from about 7 to about 15 being particularly preferred. The peptides may be digests of naturally occurring proteins as is outlined above, random peptides, or “biased” random peptides. By “randomized” or grammatical equivalents herein is meant that each nucleic acid and peptide consists of essentially random nucleotides and amino acids, respectively. Since generally these random peptides (or nucleic acids, discussed below) are chemically synthesized, they may incorporate any nucleotide or amino acid at any position. The synthetic process can be designed to generate randomized proteins or nucleic acids, to allow the formation of all or most of the possible combinations over the length of the sequence, thus forming a library of randomized candidate bioactive proteinaceous agents.

**[0098]** In one embodiment, the library is fully randomized, with no sequence preferences or constants at any position. In a preferred embodiment, the library is biased. That is, some positions within the sequence are either held constant, or are selected from a limited number of possibilities. For example, in a preferred embodiment, the nucleotides or amino acid residues are randomized within a defined class, for example, of hydrophobic amino acids, hydrophilic residues, sterically biased (either small or large) residues, towards the creation of nucleic acid binding domains, the creation of cysteines, for cross-linking, prolines for SH-3 domains, serines, threonines, tyrosines or histidines for phosphorylation sites, etc., or to purines, etc.

**[0099]** In a preferred embodiment, the candidate bioactive agents are nucleic acids. By “nucleic acid” or “oligonucle-

otide" or grammatical equivalents herein means at least two nucleotides covalently linked together. A nucleic acid of the present invention will generally contain phosphodiester bonds, although in some cases, as outlined below, nucleic acid analogs are included that may have alternate backbones, comprising, for example, phosphoramidate (Beaucage et al., *Tetrahedron* 49(10):1925 (1993) and references therein; Letsinger, *J. Org. Chem.* 35:3800 (1970); Sprinzl et al., *Eur. J. Biochem.* 81:579 (1977); Letsinger et al., *Nucl. Acids Res.* 14:3487 (1986); Sawal et al., *Chem. Lett.* 805 (1984), Letsinger et al., *J. Am. Chem. Soc.* 110:4470 (1988); and Pauwels et al., *Chemica Scripta* 26:141 (1986)), phosphorothioate (Mag et al., *Nucleic Acids Res.* 19:1437 (1991); and U.S. Pat. No. 5,644,048), phosphorodithioate (Briu et al., *J. Am. Chem. Soc.* 111:2321 (1989), O-methylphosphoramidite linkages (see Eckstein, *Oligonucleotides and Analogues: A Practical Approach*, Oxford University Press), and peptide nucleic acid backbones and linkages (see Egholm, *J. Am. Chem. Soc.* 114:1895 (1992); Meier et al., *Chem. Int. Ed. Engl.* 31:1008 (1992); Nielsen, *Nature*, 365:566 (1993); Carlsson et al., *Nature* 380:207 (1996), all of which are incorporated by reference). Other analog nucleic acids include those with positive backbones (Denpcy et al., *Proc. Natl. Acad. Sci. USA* 92:6097 (1995); non-ionic backbones (U.S. Pat. Nos. 5,386,023, 5,637,684, 5,602,240, 5,216,141 and 4,469,863; Kiedrowski et al., *Angew. Chem. Intl. Ed. English* 30:423 (1991); Letsinger et al., *J. Am. Chem. Soc.* 110:4470 (1988); Letsinger et al., *Nucleoside & Nucleotide* 13:1597 (1994); Chapters 2 and 3, *ASC Symposium Series 580, "Carbohydrate Modifications in Antisense Research"*, Ed. Y. S. Sanghui and P. Dan Cook; Mesmaeker et al., *Bioorganic & Medicinal Chem. Lett.* 4:395 (1994); Jeffs et al., *J. Biomolecular NMR* 34:17 (1994); *Tetrahedron Lett.* 37:743 (1996)) and non-ribose backbones, including those described in U.S. Pat. Nos. 5,235,033 and 5,034,506, and Chapters 6 and 7, *ASC Symposium Series 580, "Carbohydrate Modifications in Antisense Research"*, Ed. Y. S. Sanghui and P. Dan Cook. Nucleic acids containing one or more carbocyclic sugars are also included within the definition of nucleic acids (see Jenkins et al., *Chem. Soc. Rev.* (1995) pp 169-176). Several nucleic acid analogs are described in Rawls, *C & E News* Jun. 2, 1997 page 35. All of these references are hereby expressly incorporated by reference. These modifications of the ribose-phosphate backbone may be done to facilitate the addition of additional moieties such as labels, or to increase the stability and half-life of such molecules in physiological environments. In addition, mixtures of naturally occurring nucleic acids and analogs can be made. Alternatively, mixtures of different nucleic acid analogs, and mixtures of naturally occurring nucleic acids and analogs may be made. The nucleic acids may be single stranded or double stranded, as specified, or contain portions of both double stranded or single stranded sequence. The nucleic acid may be DNA, both genomic and cDNA, RNA or a hybrid, where the nucleic acid contains any combination of deoxyribo- and ribo-nucleotides, and any combination of bases, including uracil, adenine, thymine, cytosine, guanine, inosine, xanthine hypoxanthine, isocytosine, isoguanine, etc.

**[0100]** As described above generally for proteins, nucleic acid candidate bioactive agents may be naturally occurring nucleic acids, random nucleic acids, or "biased" random nucleic acids. For example, digests of procaryotic or eucaryotic genomes may be used as is outlined above for proteins.

**[0101]** In a preferred embodiment, the candidate bioactive agents are organic chemical moieties, a wide variety of which are available in the literature.

**[0102]** In a preferred embodiment, as outlined above, screens may be done on individual genes and gene products (proteins). In a preferred embodiment, the gene or protein has been identified as described below as a differentially expressed gene important in a particular state. Thus, in one embodiment, screens are designed to first find candidate agents that can bind to differentially expressed proteins, and then these agents may be used in assays that evaluate the ability of the candidate agent to modulate differentially expressed activity. Thus, as will be appreciated by those in the art, there are a number of different assays which may be run; binding assays and activity assays.

**[0103]** In a preferred embodiment, binding assays are provided. In one embodiment, the methods comprise combining a cellular proliferation protein and a candidate bioactive agent in the presence or absence of microtubules, and determining the binding of the candidate agent to the cellular proliferation protein. Preferred embodiments utilize the human cellular proliferation protein, although other mammalian proteins may also be used as discussed above, for example for the development of animal models of human disease. In some embodiments, as outlined herein, variant or derivative cellular proliferation proteins may be used.

**[0104]** Generally, in a preferred embodiment of the methods herein, the cellular proliferation protein or the candidate agent is non-diffusably bound to an insoluble support having isolated sample receiving areas (e.g. a microtiter plate, an array, etc.). The insoluble supports may be made of any composition to which the compositions can be bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports may be solid or porous and of any convenient shape. Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (e.g., polystyrene), polysaccharides, nylon or nitrocellulose, Teflon™, etc. Microtiter plates and arrays are especially convenient because a large number of assays can be carried out simultaneously, using small amounts of reagents and samples. The particular manner of binding of the composition is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the composition and is nondiffusable. Preferred methods of binding include the use of antibodies (which do not sterically block either the ligand binding site or activation sequence when the protein is bound to the support), direct binding to "sticky" or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the protein or agent, excess unbound material is removed by washing. The sample receiving areas may then be blocked through incubation with bovine serum albumin (BSA), casein or other innocuous protein or other moiety.

**[0105]** In a preferred embodiment, the cellular proliferation protein is bound to the support, and, in the presence or absence of microtubules, a candidate bioactive agent is added to the assay. Alternatively, the candidate agent is bound to the support and the cellular proliferation protein is added. Novel binding agents include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. A wide variety of assays may be used for this purpose, including labeled in vitro protein-protein binding

assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like. Moreover, in another aspect, screening assays are performed herein where neither the drug candidate nor cellular proliferation protein are bound to a solid support. Soluble assays are known in the art. In one embodiment, binding of a cellular proliferation protein, or fragment thereof, to a drug candidate can be determined by changes in fluorescence of either the cellular proliferation protein or the drug candidate, or both. Fluorescence may be intrinsic or conferred by labeling either component with a fluorophor. As an example that is not meant to be limiting, binding could be detected by fluorescence polarization.

**[0106]** The determination of the binding of the candidate bioactive agent to the cellular proliferation protein may be done in a number of ways. In a preferred embodiment, the candidate bioactive agent is labeled, and binding determined directly. For example, this may be done by attaching all or a portion of the cellular proliferation protein to a solid support, adding a labeled candidate agent (for example a fluorescent label), washing off excess reagent, and determining whether the label is present on the solid support. Various blocking and washing steps may be utilized as is known in the art.

**[0107]** By "labeled" herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g. radioisotope, fluorophores including organo-metallic fluorescent compounds, enzyme, antibodies, particles such as magnetic particles, chemiluminescers, or specific binding molecules, etc. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

**[0108]** In some embodiments, only one of the components is labeled. For example, the proteins (or proteinaceous candidate agents) may be labeled at tyrosine positions using  $^{125}\text{I}$ , or with fluorophores. Alternatively, more than one component may be labeled with different labels; using  $^{125}\text{I}$  for the proteins, for example, and a fluorophor for the candidate agents.

**[0109]** In a preferred embodiment, the binding of the candidate bioactive agent is determined through the use of competitive binding assays. In this embodiment, the competitor is a binding moiety known to bind to the target molecule (i.e. cellular proliferation protein), such as ATP, microtubules, an antibody, peptide; binding partner, ligand, etc. Under certain circumstances, there may be competitive binding as between the bioactive agent and the binding moiety, with the binding moiety displacing the bioactive agent.

**[0110]** In one embodiment, the candidate bioactive agent is labeled. Either the candidate bioactive agent, or the competitor, or both, is added first to the protein for a time sufficient to allow binding, if present. Incubations may be performed at any temperature which facilitates optimal activity, typically between 4 and 40° C. Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

**[0111]** In a preferred embodiment, the competitor is added first, followed by the candidate bioactive agent. Displacement of the competitor is an indication that the candidate bioactive agent is binding to the cellular proliferation protein and thus is capable of binding to, and potentially modulating, the activity of the cellular proliferation protein. In this embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate bioactive agent is labeled, the presence of the label on the support indicates displacement.

**[0112]** In an alternative embodiment, the candidate bioactive agent is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor may indicate that the bioactive agent is bound to the cellular proliferation protein with a higher affinity. Thus, if the candidate bioactive agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate that the candidate agent is capable of binding to the cellular proliferation protein.

**[0113]** In another aspect herein, proteins which bind to KSP or a fragment thereof are identified. Genetic systems have been described to detect protein-protein interactions. The first work was done in yeast systems, namely the "yeast two-hybrid" system. The basic system requires a protein-protein interaction in order to turn on transcription of a reporter gene. Subsequent work was done in mammalian cells. See Fields et al., *Nature* 340:245 (1989); Vasavada et al., *PNAS USA* 88:10686 (1991); Fearon et al., *PNAS USA* 89:7958 (1992); Dang et al., *Mol. Cell. Biol.* 11:954 (1991); Chien et al., *PNAS USA* 88:9578 (1991); and U.S. Pat. Nos. 5,283,173, 5,667,973, 5,468,614, 5,525,490, and 5,637,463.

**[0114]** In a preferred embodiment, the binding site of the cellular proliferation protein is identified and provided herein. This can be done in a variety of ways. For example, once the cellular proliferation protein has been identified as binding to a bioactive agent, the protein is fragmented or modified and the assays repeated to identify the necessary components for binding.

**[0115]** In a preferred embodiment, the methods comprise differential screening to identify bioactive agents that are capable of modulating the activity of the cellular proliferation proteins. In this embodiment, the methods comprise combining a cellular proliferation protein and a competitor in a first sample. A second sample comprises a candidate bioactive agent, a cellular proliferation protein and a competitor. The binding of the competitor is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of an agent capable of binding to the cellular proliferation protein and, in one embodiment, modulating its activity. Methods of determining modulation of activity are further described below. That is, if the binding of the competitor is different in the second sample relative to the first sample, the agent is capable of binding to the cellular proliferation protein.

**[0116]** Alternatively, a preferred embodiment utilizes differential screening to identify drug candidates that, in the presence or absence of microtubules, bind to the native cellular proliferation protein, but cannot bind to modified cellular proliferation proteins. The structure of the cellular proliferation protein may be modeled, and used in rational drug design to synthesize agents that interact with that site. Drug candidates that affect cellular proliferation bioactivity are

also identified by screening drugs for the ability to either enhance or reduce the activity of the protein in the presence or absence of microtubules.

**[0117]** Positive controls and negative controls may be used in the assays. Preferably all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples is for a time sufficient for the binding of the agent to the protein. Following incubation, all samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

**[0118]** A variety of other reagents may be included in the screening assays. These include reagents like salts, neutral proteins, e.g. albumin, detergents, etc which may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

**[0119]** Screening for agents that modulate the activity of cellular proliferation proteins may also be done. In a preferred embodiment, methods for screening for a bioactive agent capable of modulating the activity of cellular proliferation proteins comprise the steps of adding a candidate bioactive agent to a sample of cellular proliferation proteins in the presence or absence of microtubules, as above, and determining an alteration in the biological activity of cellular proliferation proteins. "Modulating the activity of cellular proliferation" includes an increase in activity, a decrease in activity, or a change in the type or kind of activity present. Thus, in this embodiment, the candidate agent should both bind to cellular proliferation proteins (although this may not be necessary), and alter its biological or biochemical activity as defined herein. The methods include both in vitro screening methods, as are generally outlined above, and in vivo screening of cells for alterations in the presence, distribution, activity or amount of cellular proliferation proteins.

**[0120]** Thus, in this embodiment, the methods comprise combining a cellular proliferation sample and a candidate bioactive agent, and evaluating the effect on cellular proliferation activity. By "cellular proliferation protein activity" or grammatical equivalents herein is meant at least one of the cellular proliferation protein's biological activities, including, but not limited to, kinesin activity, regulation of spindle pole separation, mitosis, mitotic spindle assembly, satisfaction of the mitotic cell cycle checkpoint, cell cycle progression, apoptosis, cell proliferation, mitotic and involvement in tumor growth. An inhibitor of cellular proliferation activity is the inhibition of any one or more cellular proliferation protein activities.

**[0121]** Kinesin activity is known in the art and includes one or more kinesin activities. Kinesin activities include the ability to affect ATP hydrolysis, microtubule binding, gliding and polymerization/depolymerization (effects on microtubule dynamics), binding to other proteins of the spindle, binding to proteins involved in cell-cycle control, or serving as a substrate to other enzymes, such as kinases or proteases and specific kinesin cellular activities such as spindle separation.

**[0122]** Methods of performing motility assays are well known to those of skill in the art (see, e.g., Hall, et al. (1996), *Biophys. J.*, 71: 3467-3476, Turner et al., 1996, *Anal. Bio-*

*chem.* 242 (1):20-5; Gittes et al., 1996, *Biophys. J.* 70(1): 418-29; Shirakawa et al., 1995, *J. Exp. Biol.* 198: 1809-15; Winkelmann et al., 1995, *Biophys. J.* 68: 2444-53; Winkelmann et al., 1995, *Biophys. J.* 68: 72S, and the like).

**[0123]** In addition to the assays described above, methods known in the art for determining ATPase activity can be used. Preferably, solution based assays are utilized. Alternatively, conventional methods are used. For example, P<sub>i</sub> release from kinesin can be quantified. In one preferred embodiment, the ATPase activity assay utilizes 0.3 M PCA (perchloric acid) and malachite green reagent (8.27 mM sodium molybdate II, 0.33 mM malachite green oxalate, and 0.8 mM Triton X-100). To perform the assay, 10  $\mu$ L of reaction is quenched in 90  $\mu$ L of cold 0.3 M PCA. Phosphate standards are used so data can be converted to mM inorganic phosphate released. When all reactions and standards have been quenched in PCA, 100  $\mu$ L of malachite green reagent is added to the relevant wells in e.g., a microtiter plate. The mixture is developed for 10-15 minutes and the plate is read at an absorbance of 650 nm. If phosphate standards were used, absorbance readings can be converted to mM P<sub>i</sub> and plotted over time. Additionally, ATPase assays known in the art include the luciferase assay.

**[0124]** In another preferred method, kinesin activity is measured by the methods disclosed in Ser. No. 09/314,464, filed May 18, 1999, entitled, Compositions and Assay Utilizing ADP or Phosphate for Detecting Protein Modulators.

**[0125]** In a preferred embodiment, the activity of the cellular proliferation protein is increased; in another preferred embodiment, the activity of the cellular proliferation protein is decreased. Thus, bioactive agents that are antagonists are preferred in some embodiments, and bioactive agents that are agonists may be preferred in other embodiments.

**[0126]** In one aspect of the invention, cells containing cellular proliferation sequences are used in drug screening assays by evaluating the effect of drug candidates on cellular proliferation. Cell type include normal cells, and more preferably cells with abnormal proliferative rates including tumor cells, most preferably human tumor cells. Methods of assessing cellular proliferation are known in the art and include growth and viability assays using cultured cells. In such assays, cell populations are monitored for growth and or viability, often over time and comparing samples incubated with various concentrations of the bioactive agent or without the bioactive agent. Cell number can be quantified using agents that such as 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) [U.S. Pat. No. 5,185,450] and Alamar Blue which are converted to colored or fluorescent compounds in the presence of metabolically active cells. Alternatively, dyes that bind to cellular protein such as sulforhodamine B (SRB) or crystal violet can be used to quantify cell number. Alternatively, cells can be directly counted using a particle counter, such as a Coulter Counter® manufactured by Beckman Coulter, or counted using a microscope to observe cells on a hemocytometer. Preferably, cells counted using the hemocytometer are observed in a solution of trypan blue to distinguish viable from dead cells. Other methods of quantifying cell number are known to those skilled in the art. These assays can be performed on any of the cells, including those in a state of necrosis.

**[0127]** Moreover, apoptosis can be determined by methods known in the art. For example, markers for apoptosis are known, and TUNEL (TdT-mediated dUTP-fluorescein nick

end labeling) kits can be bought commercially, for example, Boehringer Mannheim kit, catalog no. 168795.

**[0128]** In a preferred embodiment, the methods comprise adding a candidate bioactive agent, as defined above, to a cell comprising cellular proliferation proteins. Preferred cell types include almost any cell. The cells contain a nucleic acid, preferably recombinant, that encodes a cellular proliferation protein. In a preferred embodiment, a library of candidate agents are tested on a plurality of cells.

**[0129]** In one aspect, the assays are evaluated in the presence or absence or previous or subsequent exposure to physiological signals, for example hormones, antibodies, peptides, antigens, cytokines, growth factors, action potentials, pharmacological agents including chemotherapeutics, radiation, carcinogenics, or other cells (i.e. cell-cell contacts). In another example, the determinations are determined at different stages of the cell cycle process.

**[0130]** In one aspect of the invention, the cellular proliferation sequences and cells containing cellular proliferation sequences are used in drug screening assays by evaluating the effect of drug candidates on a "gene expression profile" or expression profile genes. In a preferred embodiment, the expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes after treatment with a candidate agent. See, Zlokarnik, et al., *Science* 279, 84-8 (1998).

**[0131]** In one aspect, the expression levels of genes are determined for different cellular states in the cellular proliferation phenotype; that is, the expression levels of genes in normal tissue in proliferating and non-proliferating states, and in abnormal cellular proliferation tissue (and in some cases, for varying severities of cellular proliferation that relate to prognosis, as outlined below) are evaluated to provide expression profiles. Abnormal states include cancer states and other hyper or hypo proliferation states as further defined below.

**[0132]** An expression profile of a particular cell state or point of development is essentially a "fingerprint" of the state; while two states may have any particular gene similarly expressed, the evaluation of a number of genes simultaneously allows the generation of a gene expression profile that is unique to the state of the cell. By comparing expression profiles of cells in different states, information regarding which genes are important (including both up- and down-regulation of genes) in each of these states is obtained. Then, diagnosis may be done or confirmed: does tissue from a particular patient have the gene expression profile of normal or abnormal cellular proliferation tissue.

**[0133]** "Differential expression," or grammatical equivalents as used herein, refers to both qualitative as well as quantitative differences in the genes' temporal and/or cellular expression patterns within and among the cells. Thus, a differentially expressed gene can qualitatively have its expression altered, including an activation or inactivation, in, for example, normal versus abnormal cellular proliferation tissue. That is, genes may be turned on or turned off in a particular state, relative to another state or have a different timing pattern, for example, cancerous cells may have genes which stay on. As is apparent to the skilled artisan, any comparison of two or more states can be made and repeated at various time points. Such a qualitatively regulated gene will exhibit an expression pattern within a state or cell type which is detectable by standard techniques in one such state or cell type, but is not detectable in both. Alternatively, the determination is

quantitative in that expression is increased or decreased: that is, the expression of the gene is either upregulated, resulting in an increased amount of transcript, or downregulated, resulting in a decreased amount of transcript. The degree to which expression differs need only be large enough to quantify via standard characterization techniques as outlined below, such as by use of Affymetrix GeneChip™ expression arrays, Lockhart, *Nature Biotechnology*, 14:1675-1680 (1996), hereby expressly incorporated by reference. Other techniques include, but are not limited to, quantitative reverse transcriptase PCR, Northern analysis and RNase protection. As outlined above, preferably the change in expression (i.e. upregulation or downregulation) is at least about 50%, more preferably at least about 100%, more preferably at least about 150%, more preferably, at least about 200%, with from 300 to at least 1000% being especially preferred.

**[0134]** As will be appreciated by those in the art, this may be done by evaluation at either the gene transcript, or the protein level; that is, the amount of gene expression may be monitored using nucleic acid probes to the DNA or RNA equivalent of the gene transcript, and the quantification of gene expression levels, or, alternatively, the final gene product itself (protein) can be monitored, for example through the use of antibodies to the cellular proliferation protein and standard immunoassays (ELISAs, etc.) or other techniques, including mass spectroscopy assays, 2D gel electrophoresis assays, etc. Thus, the proteins corresponding to cellular proliferation genes, i.e. those identified as being important in a cellular proliferation phenotype, can be evaluated in a cellular proliferation diagnostic test.

**[0135]** In a preferred embodiment nucleic acids encoding the cellular proliferation protein are detected. Although DNA or RNA encoding the cellular proliferation protein may be detected, of particular interest are methods wherein the mRNA encoding a cellular proliferation protein is detected. The presence of mRNA in a sample is an indication that the cellular proliferation gene has been transcribed to form the mRNA, and suggests that the protein is expressed. Probes to detect the mRNA can be any nucleotide/deoxynucleotide probe that is complementary to and base pairs with the mRNA and includes but is not limited to oligonucleotides, cDNA or RNA. Probes also should contain a detectable label, as defined herein. In one method the mRNA is detected after immobilizing the nucleic acid to be examined on a solid support such as nylon membranes and hybridizing the probe with the sample. Following washing to remove the non-specifically bound probe, the label is detected. In another method detection of the mRNA is performed in situ. In this method permeabilized cells or tissue samples are contacted with a detectably labeled nucleic acid probe for sufficient time to allow the probe to hybridize with the target mRNA. Following washing to remove the non-specifically bound probe, the label is detected. For example a digoxigenin labeled ribo-probe (RNA probe) that is complementary to the mRNA encoding a cellular proliferation protein is detected by binding the digoxigenin with an anti-digoxigenin secondary antibody and developed with nitro blue tetrazolium and 5-bromo-4-chloro-3-indoyl phosphate.

**[0136]** In one case, having identified a particular gene as up regulated in cellular proliferation, candidate bioactive agents may be screened to modulate this gene's response; preferably to down regulate the gene, although in some circumstances to up regulate the gene. "Modulation" thus includes both an increase and a decrease in gene expression or a change in

temporal pattern. The preferred amount of modulation will depend on the original change of the gene expression in normal versus tumor tissue, with changes of at least 10%, preferably 50%, more preferably 100-300%, and in some embodiments 300-1000% or greater. Thus, if a gene exhibits a 4 fold increase in tumor compared to normal tissue, a decrease of about four fold is desired; a 10 fold decrease in tumor compared to normal tissue gives a 10 fold increase in expression for a candidate agent is desired.

**[0137]** In a preferred embodiment, gene expression monitoring is done and a number of genes, i.e. an expression profile, is monitored simultaneously, although multiple protein expression monitoring can be done as well.

**[0138]** In one embodiment, the cellular proliferation nucleic acid probes are attached to biochips as outlined below for the detection and quantification of cellular proliferation sequences in a particular cell.

**[0139]** Generally, in a preferred embodiment, a candidate bioactive agent is added to the cells prior to analysis. Any cell can be used, including normal and abnormal cells, including tumor and non-tumor mammalian, preferably human cells. In some cases, plant cells are used. After the candidate agent has been added and the cells allowed to incubate for some period of time, the sample containing the target sequences to be analyzed is added to the biochip. If required, the target sequence is prepared using known techniques. For example, the sample may be treated to lyse the cells, using known lysis buffers, electroporation, etc., with purification and/or amplification such as PCR occurring as needed, as will be appreciated by those in the art. For example, an in vitro transcription with labels covalently attached to the nucleosides is done. Generally, the nucleic acids are labeled with biotin-FITC or PE, or with cy3 or cy5.

**[0140]** In a preferred embodiment, the target sequence is labeled with, for example, a fluorescent, a chemiluminescent, a chemical, or a radioactive signal, to provide a means of detecting the target sequence's specific binding to a probe. The label also can be an enzyme, such as, alkaline phosphatase or horseradish peroxidase, which when provided with an appropriate substrate produces a product that can be detected. Alternatively, the label can be a labeled compound or small molecule, such as an enzyme inhibitor, that binds but is not catalyzed or altered by the enzyme. The label also can be a moiety or compound, such as, an epitope tag or biotin which specifically binds to streptavidin. For the example of biotin, the streptavidin is labeled as described above, thereby, providing a detectable signal for the bound target sequence. As known in the art, unbound labeled streptavidin is removed prior to analysis.

**[0141]** As will be appreciated by those in the art, these assays can be direct hybridization assays or can comprise "sandwich assays", which include the use of multiple probes, as is generally outlined in U.S. Pat. Nos. 5,681,702, 5,597,909, 5,545,730, 5,594,117, 5,591,584, 5,571,670, 5,580,731, 5,571,670, 5,591,584, 5,624,802, 5,635,352, 5,594,118, 5,359,100, 5,124,246 and 5,681,697, all of which are hereby incorporated by reference. In this embodiment, in general, the target nucleic acid is prepared as outlined above, and then added to the biochip comprising a plurality of nucleic acid probes, under conditions that allow the formation of a hybridization complex.

**[0142]** A variety of hybridization conditions may be used in the present invention, including high, moderate and low stringency conditions as outlined above. The assays are generally

run under stringency conditions which allows formation of the label probe hybridization complex only in the presence of target. Stringency can be controlled by altering a step parameter that is a thermodynamic variable, including, but not limited to, temperature, formamide concentration, salt concentration, chaotropic salt concentration pH, organic solvent concentration, etc.

**[0143]** These parameters may also be used to control non-specific binding, as is generally outlined in U.S. Pat. No. 5,681,697. Thus it may be desirable to perform certain steps at higher stringency conditions to reduce non-specific binding.

**[0144]** The reactions outlined herein may be accomplished in a variety of ways, as will be appreciated by those in the art. Components of the reaction may be added simultaneously, or sequentially, in any order, with preferred embodiments outlined below. In addition, the reaction may include a variety of other reagents may be included in the assays. These include reagents like salts, buffers, neutral proteins, e.g. albumin, detergents, etc which may be used to facilitate optimal hybridization and detection, and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used, depending on the sample preparation methods and purity of the target.

**[0145]** Once the assay is run, the data is analyzed to determine the expression levels, and changes in expression levels as between states, of individual genes, forming a gene expression profile.

**[0146]** In one aspect, the screens are done to identify drugs or bioactive agents that modulate the cellular proliferation phenotype. Specifically, there are several types of screens that can be run. A preferred embodiment is in the screening of candidate agents that can induce or suppress a particular expression profile, thus preferably generating the associated phenotype. That is, candidate agents that can mimic or produce an expression profile in cellular proliferation similar to the expression profile of normal non-cancerous tissue is expected to result in a suppression of the cellular proliferation phenotype. Thus, in this embodiment, mimicking an expression profile, or changing one profile to another, is the goal.

**[0147]** In a preferred embodiment, as for the diagnosis and prognosis applications discussed below, having identified the differentially expressed genes important in any one state as further described below, screens can be run to alter the expression of the genes individually. That is, screening for modulation of regulation of expression of a single gene can be done that is, rather than try to mimic all or part of an expression profile, screening for regulation of individual genes can be done. Thus, for example, particularly in the case of target genes whose presence, absence or temporal pattern is unique between two states, screening is done for modulators of the target gene expression. In a preferred embodiment, the target gene encodes the cellular proliferation protein described herein. Thus, screening of candidate agents that modulate the cellular proliferation phenotype either at the gene expression level or the protein level can be done.

**[0148]** In addition screens can be done for novel genes that are induced in response to a candidate agent. After identifying a candidate agent based upon its ability to suppress a cellular proliferation expression pattern leading to a normal expression pattern, or modulate a single cellular proliferation gene expression profile so as to mimic the expression of the gene

from normal tissue, a screen as described above can be performed to identify genes that are specifically modulated in response to the agent. Comparing expression profiles between normal tissue and agent treated cellular proliferation tissue reveals genes that are not expressed in normal tissue or cellular proliferation tissue, but are expressed in agent treated tissue. These agent specific sequences can be identified and used by any of the methods described herein for cellular proliferation genes or proteins. In particular these sequences and the proteins they encode find use in marking or identifying agent treated cells. In addition, antibodies can be raised against the agent induced proteins and used to target novel therapeutics to the treated cellular proliferation tissue sample.

**[0149]** In one embodiment, a candidate agent is administered to a population of cellular proliferation cells, that thus has an associated Cellular proliferation expression profile. By “administration” or “contacting” herein is meant that the candidate agent is added to the cells in such a manner as to allow the agent to act upon the cell, whether by uptake and intracellular action, or by action at the cell surface. In some embodiments, nucleic acid encoding a proteinaceous candidate agent (i.e. a peptide) may be put into a viral construct such as a retroviral construct and added to the cell, such that expression of the peptide agent is accomplished; see PCT US97/01019, hereby expressly incorporated by reference. The phrase “under conditions which allow the cell to uptake the candidate agent” means that the cell is biologically involved in the uptake and intracellular action, or by action at the cell surface in that the agent is not injected into the cell. It is understood that targeting ligands and biochemically agents can be used to facilitate the uptake, however, this differs from mechanical injection. Mechanical injection is explicitly excluded from the definition of “taken up by the cell” as used herein, and is excluded from conditions inductive to high throughput assays as used herein:

**[0150]** Once the candidate agent has been administered to the cells, the cells can be washed if desired and are allowed to incubate under preferably physiological conditions for some period of time. The cells are then harvested and a new gene expression profile is generated, as outlined herein.

**[0151]** Thus, for example, cellular proliferation tissue may be screened for agents that reduce or suppress the cellular proliferation phenotype. A change in at least one gene of the expression profile indicates that the agent has an effect on cellular proliferation activity. By defining such a signature for the cellular proliferation phenotype, screens for new drugs that alter the phenotype can be devised. With this approach, the drug target need not be known and need not be represented in the original expression screening platform, nor does the level of transcript for the target protein need to change.

**[0152]** In all the methods provided herein, bioactive agents are identified. Similarly, compounds which interfere with binding or interaction between the cellular proliferation protein and an identified binding or modulating agent can be identified. Moreover, transgenic models as discussed below may be used to identify bioactive agents. Compounds with pharmacological activity are able to enhance or interfere with the activity of the cellular proliferation protein. The compounds can be used in further assays so as to confirm activity wherein necessary or optimize conditions including varying the identified molecules. In a preferred embodiment, the agents are used as therapeutics as discussed below.

**[0153]** In a further aspect of the present invention, methods of modulating cellular proliferation in cells or organisms are

provided. In one embodiment, the methods comprise administering to a cell an anti-cellular proliferation antibody as further discussed below that reduces or eliminates the biological activity of an endogenous cellular proliferation protein. In a preferred embodiment, a nucleic acid encoding said antibody is administered. Agents identified to modulate cellular proliferation can also be used. Alternatively, the methods comprise administering to a cell or organism a composition comprising a cellular proliferation sequence.

**[0154]** In a preferred embodiment, for example when the cellular proliferation sequence is down-regulated in cellular proliferation, the activity of the cellular proliferation gene is increased by increasing the amount of cellular proliferation in the cell, for example by overexpressing the endogenous cellular proliferation or by administering a gene encoding the cellular proliferation sequence, using known gene-therapy techniques, for example. In a preferred embodiment, the gene therapy techniques include the incorporation of the exogenous gene using enhanced homologous recombination (EHR), for example as described in PCT/US93/03868, hereby incorporated by reference in its entirety. Alternatively, for example when the cellular proliferation sequence is up-regulated in cellular proliferation, the activity of the endogenous cellular proliferation gene is decreased, for example by the administration of a cellular proliferation antisense nucleic acid. Preferably, as discussed below, cellular proliferation is inhibited.

**[0155]** Thus, In one embodiment, a method of inhibiting cell division is provided. In a preferred embodiment, a method of inhibiting tumor growth is provided. In a further embodiment, methods of treating cells or individuals with cancer are provided. The method comprises administration of a cellular proliferation inhibitor.

**[0156]** In one embodiment, a cellular proliferation inhibitor is an antibody as discussed above and further described below. In another embodiment, the cellular proliferation inhibitor is an antisense molecule as discussed above. Antisense molecules as used herein include antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target mRNA (sense) or DNA (antisense) sequences for cellular proliferation molecules. A preferred antisense molecule is for KSP or for a ligand or activator thereof. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment generally at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, for example, Stein and Cohen (*Cancer Res.* 48:2659, 1988) and van der Krol et al. (*BioTechniques* 6:958, 1988).

**[0157]** Antisense molecules may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell. Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO



90/10448. It is understood that the use of antisense molecules or knock out and knock in models may also be used in screening assays as discussed above, in addition to methods of treatment. Moreover, knock out models can include knocking out expression, rather than the genome, such as by the use of ribozymes. In one case, ribozymes are a preferred KSP inhibitor.

**[0158]** As discussed above, the methods and compositions herein are not limited to cancer. Disease states which can be treated by the methods and compositions provided herein include, but are not limited to, cancer (further discussed below), restenosis, autoimmune disease, arthritis, graft rejection, inflammatory bowel disease, proliferation induced after medical procedures, including, but not limited to, surgery, angioplasty, and the like. It is appreciated that in some cases the cells may not be in a hyper or hypo proliferation state (abnormal state) and still require treatment. For example, during wound healing, the cells may be proliferating “normally”, but proliferation enhancement may be desired. Similarly, as discussed above, in the agriculture arena, cells may be in a “normal” state, but proliferation modulation may be desired to enhance a crop by directly enhancing growth of a crop, or by inhibiting the growth of a plant or organism which adversely affects the crop. Thus, in one embodiment, the invention herein includes application to cells or individuals afflicted or impending affliction with any one of these disorders or states.

**[0159]** The compositions and methods provided herein are particularly deemed useful for the treatment of cancer including solid tumors such as skin, breast, brain, cervical carcinomas, testicular carcinomas, etc. More particularly, cancers that may be treated by the compositions and methods of the invention include, but are not limited to: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Log: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, calm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondroblastoma), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull

(osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis); brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma [embryonal rhabdomyosarcoma], fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]; Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. The cancer can be solid tumors or metastatic. Thus, the term “cancerous cell” as provided herein, includes a cell afflicted by any one of the above identified conditions.

**[0160]** In another aspect herein, diagnostic assays are provided herein. In one embodiment, the cellular proliferation sequences are used in the diagnostic assays. This can be done on an individual gene or corresponding polypeptide level. In a preferred embodiment, the expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes and/or corresponding polypeptides. In a preferred embodiment in situ hybridization of labeled cellular proliferation nucleic acid probes to tissue arrays is done. For example, arrays of tissue samples, including cellular proliferation tissue in various states and or time points and/or normal tissue, are made. In situ hybridization as is known in the art can then be done. It is understood that conventional antibody and protein localization methods can also be used in diagnostic assays herein.

**[0161]** It is understood that when comparing the fingerprints between an individual and a standard, the skilled artisan can make a diagnosis as well as a prognosis. It is further understood that the genes which indicate the diagnosis may differ from those which indicate the prognosis.

**[0162]** In a preferred embodiment, the cellular proliferation sequences are used in prognosis assays. As above, gene expression profiles can be generated that correlate to cellular proliferation severity, in terms of long term prognosis. Again, this may be done on either a protein or gene level, with the use of genes being preferred. In both the diagnostic and prognostic assays, the cellular proliferation probes can be attached to biochips as described below for the detection and quantification of cellular proliferation sequences in a tissue or patient.

**[0163]** Accordingly, disorders based on mutant or variant cellular proliferation genes may also be determined. In one embodiment, the invention provides methods for identifying cells containing variant cellular proliferation genes comprising determining all or part of the sequence of at least one endogenous cellular proliferation genes in a cell. As will be



appreciated by those in the art, this may be done using any number of sequencing techniques. In a preferred embodiment, the invention provides methods of identifying the cellular proliferation genotype of an individual comprising determining all or part of the sequence of at least one cellular proliferation gene of the individual. This is generally done in at least one tissue of the individual, and may include the evaluation of a number of tissues or different samples of the same tissue. The method may include comparing the sequence of the sequenced cellular proliferation gene to a known cellular proliferation gene, i.e. a wild-type gene.

**[0164]** The sequence of all or part of the cellular proliferation gene can then be compared to the sequence of a known cellular proliferation gene to determine if any differences exist. This can be done using any number of known homology programs, such as Bestfit, etc. In a preferred embodiment, the presence of a difference in the sequence between the cellular proliferation gene of the patient and the known cellular proliferation gene is indicative of a disease state or a propensity for a disease state, as outlined herein.

**[0165]** In a preferred embodiment, the cellular proliferation genes are used as probes to determine the number of copies of the cellular proliferation gene in the genome.

**[0166]** In another preferred embodiment cellular proliferation genes are used as probes to determine the chromosomal localization of the cellular proliferation genes. Information such as chromosomal localization finds use in providing a diagnosis or prognosis in particular when chromosomal abnormalities such as translocations, and the like are identified in cellular proliferation gene loci.

**[0167]** Once a determination has been made regarding the proliferation state of a cell, if desired, the compositions or agents described herein can be administered. The compounds having the desired pharmacological activity may be administered in a physiologically acceptable carrier (also called a pharmaceutically acceptable carrier) to a host. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways as discussed below. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt. %. The agents may be administered alone or in combination with other treatments, e.g., radiation.

**[0168]** Thus, in a preferred embodiment, cellular proliferation proteins and modulators are administered as therapeutic agents. Similarly, cellular proliferation genes (including both the full-length sequence, partial sequences, or regulatory sequences of the cellular proliferation coding regions) can be administered in gene therapy applications, as is known in the art. These cellular proliferation genes can include antisense applications, either as gene therapy (i.e. for incorporation into the genome) or as antisense compositions, as will be appreciated by those in the art.

**[0169]** In the preferred embodiment, the pharmaceutical compositions are in a water soluble form, such as being present as pharmaceutically acceptable salts, which is meant to include both acid and base addition salts. "Pharmaceutically acceptable acid addition salt" refers to those salts that retain the biological effectiveness of the free bases and that are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic

acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like. "Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium, and magnesium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine.

**[0170]** The pharmaceutical compositions can be prepared in various forms, such as granules, tablets, pills, suppositories, capsules, suspensions, salves, lotions and the like. Pharmaceutical grade organic or inorganic carriers and/or diluents suitable for oral and topical use can be used to make up compositions containing the therapeutically-active compounds. Diluents known to the art include aqueous media, vegetable and animal oils and fats. Stabilizing agents, wetting and emulsifying agents, salts for varying the osmotic pressure or buffers for securing an adequate pH value, and skin penetration enhancers can be used as auxiliary agents. The pharmaceutical compositions may also include one or more of the following: carrier proteins such as serum albumin; buffers; fillers such as microcrystalline cellulose, lactose, corn and other starches; binding agents; sweeteners and other flavoring agents; coloring agents; and polyethylene glycol. Additives are well known in the art, and are used in a variety of formulations.

**[0171]** The administration of the cellular proliferation proteins and modulators of the present invention can be done in a variety of ways as discussed above, including, but not limited to, orally, subcutaneously, intravenously, intranasally, transdermally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly. In some instances, for example, in the treatment of wounds and inflammation, the cellular proliferation proteins and modulators may be directly applied as a solution or spray.

**[0172]** In one embodiment, a therapeutically effective dose of a cellular proliferation protein or modulator thereof is administered to a patient. By "therapeutically effective dose" herein is meant a dose that produces the effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. As is known in the art, adjustments for cellular proliferation degradation, systemic versus localized delivery, and rate of new protease synthesis, as well as the age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art.

**[0173]** a "patient" for the purposes of the present invention includes both humans and other animals, particularly mammals, and organisms. Thus the methods are applicable to both human therapy and veterinary applications. In the preferred embodiment the patient is a mammal, and in the most preferred embodiment the patient is human.

**[0174]** In a preferred embodiment, cellular proliferation genes are administered as DNA vaccines, either single genes or combinations of cellular proliferation genes. Naked DNA

vaccines are generally known in the art. Brower, *Nature Biotechnology*, 16:1304-1305 (1998).

**[0175]** In one embodiment, cellular proliferation genes of the present invention are used as DNA vaccines. Methods for the use of genes as DNA vaccines are well known to one of ordinary skill in the art, and include placing a cellular proliferation gene or portion of a cellular proliferation gene under the control of a promoter for expression in a cellular proliferation patient. The cellular proliferation gene used for DNA vaccines can encode full-length cellular proliferation proteins, but more preferably encodes portions of the cellular proliferation proteins including peptides derived from the cellular proliferation protein. In a preferred embodiment a patient is immunized with a DNA vaccine comprising a plurality of nucleotide sequences derived from a cellular proliferation gene. Similarly, it is possible to immunize a patient with a plurality of cellular proliferation genes or portions thereof as defined herein. Without being bound by theory, expression of the polypeptide encoded by the DNA vaccine, cytotoxic T-cells, helper T-cells and antibodies are induced which recognize and destroy or eliminate cells expressing cellular proliferation proteins.

**[0176]** In a preferred embodiment, the DNA vaccines include a gene encoding an adjuvant molecule with the DNA vaccine. Such adjuvant molecules include cytokines that increase the immunogenic response to the cellular proliferation polypeptide encoded by the DNA vaccine. Additional or alternative adjuvants are known to those of ordinary skill in the art and find use in the invention.

**[0177]** In another preferred embodiment cellular proliferation genes find use in generating animal models of cellular proliferation. As is appreciated by one of ordinary skill in the art, when the cellular proliferation gene identified is repressed or diminished in cellular proliferation tissue, gene therapy technology wherein antisense RNA directed to the cellular proliferation gene will also diminish or repress expression of the gene. An animal generated as such serves as an animal model of cellular proliferation that finds use in screening bioactive drug candidates. Similarly, gene knockout technology, for example as a result of homologous recombination with an appropriate gene targeting vector, will result in the absence of the cellular proliferation protein. When desired, tissue-specific expression or knockout of the cellular proliferation protein may be necessary.

**[0178]** It is also possible that the cellular proliferation protein is overexpressed in cellular proliferation. As such, transgenic animals can be generated that overexpress the cellular proliferation protein. Depending on the desired expression level, promoters of various strengths can be employed to express the transgene. Also, the number of copies of the integrated transgene can be determined and compared for a determination of the expression level of the transgene. Animals generated by such methods find use as animal models of cellular proliferation and are additionally useful in screening for bioactive molecules to treat cellular proliferation.

**[0179]** In a preferred embodiment, biochips are provided herein. Nucleic acid probes to cellular proliferation nucleic acids (both the nucleic acid sequences outlined in the figures and/or the complements thereof) are made. The nucleic acid probes attached to the biochip are designed to be substantially complementary to the cellular proliferation nucleic acids, i.e. the target sequence (either the target sequence of the sample or to other probe sequences, for example in sandwich assays), such that hybridization of the target sequence and the probes

of the present invention occurs. As outlined below, this complementarity need not be perfect; there may be any number of base pair mismatches which will interfere with hybridization between the target sequence and the single stranded nucleic acids of the present invention. However, if the number of mutations is so great that no hybridization can occur under even the least stringent of hybridization conditions, the sequence is not a complementary target sequence. Thus, by "substantially complementary" herein is meant that the probes are sufficiently complementary to the target sequences to hybridize under normal reaction conditions, particularly high stringency conditions, as outlined herein.

**[0180]** A nucleic acid probe is generally single stranded but can be partially single and partially double stranded. The strandedness of the probe is dictated by the structure, composition, and properties of the target sequence. In general, the nucleic acid probes range from about 8 to about 100 bases long, with from about 10 to about 80 bases being preferred, and from about 30 to about 50 bases being particularly preferred. That is, generally whole genes are not used. In some embodiments, much longer nucleic acids can be used, up to hundreds of bases.

**[0181]** In a preferred embodiment, more than one probe per sequence is used, with either overlapping probes or probes to different sections of the target being used. That is, two, three, four or more probes, with three being preferred, are used to build in a redundancy for a particular target. The probes can be overlapping (i.e. have some sequence in common), or separate.

**[0182]** As will be appreciated by those in the art, nucleic acids can be attached or immobilized to a solid support in a wide variety of ways. By "immobilized" and grammatical equivalents herein is meant the association or binding between the nucleic acid probe and the solid support is sufficient to be stable under the conditions of binding, washing, analysis, and removal as outlined below. The binding can be covalent or non-covalent. By "non-covalent binding" and grammatical equivalents herein is meant one or more of either electrostatic, hydrophilic, and hydrophobic interactions. Included in non-covalent binding is the covalent attachment of a molecule, such as, streptavidin to the support and the non-covalent binding of the biotinylated probe to the streptavidin. By "covalent binding" and grammatical equivalents herein is meant that the two moieties, the solid support and the probe, are attached by at least one bond, including sigma bonds, pi bonds and coordination bonds. Covalent bonds can be formed directly between the probe and the solid support or can be formed by a cross linker or by inclusion of a specific reactive group on either the solid support or the probe or both molecules. Immobilization may also involve a combination of covalent and non-covalent interactions.

**[0183]** In general, the probes are attached to the biochip in a wide variety of ways, as will be appreciated by those in the art. As described herein, the nucleic acids can either be synthesized first, with subsequent attachment to the biochip, or can be directly synthesized on the biochip.

**[0184]** The biochip comprises a suitable solid substrate. By "substrate" or "solid support" or other grammatical equivalents herein is meant any material that can be modified to contain discrete individual sites appropriate for the attachment or association of the nucleic acid probes and is amenable to at least one detection method. As will be appreciated by those in the art, the number of possible substrates are very large, and include, but are not limited to, glass and modified or

functionalized glass, plastics (including acrylics, polystyrene and copolymers of styrene and other materials, polypropylene, polyethylene, polybutylene, polyurethanes, Teflon, etc.), polysaccharides, nylon or nitrocellulose, resins, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses, plastics, etc. In general, the substrates allow optical detection and do not appreciably fluoresce. a preferred substrate is described in copending application entitled Reusable Low Fluorescent Plastic Biochip filed Mar. 15, 1999, herein incorporated by reference in its entirety.

**[0185]** Generally the substrate is planar, although as will be appreciated by those in the art, other configurations of substrates may be used as well. For example, the probes may be placed on the inside surface of a tube, for flow-through sample analysis to minimize sample volume. Similarly, the substrate may be flexible, such as a flexible foam, including closed cell foams made of particular plastics.

**[0186]** In a preferred embodiment, the surface of the biochip and the probe may be derivatized with chemical functional groups for subsequent attachment of the two. Thus, for example, the biochip is derivatized with a chemical functional group including, but not limited to, amino groups, carboxy groups, oxo groups and thiol groups, with amino groups being particularly preferred. Using these functional groups, the probes can be attached using functional groups on the probes. For example, nucleic acids containing amino groups can be attached to surfaces comprising amino groups, for example using linkers as are known in the art; for example, homo- or hetero-bifunctional linkers as are well known (see 1994 Pierce Chemical Company catalog, technical section on cross-linkers, pages 155-200, incorporated herein by reference). In addition, in some cases, additional linkers, such as alkyl groups (including substituted and heteroalkyl groups) may be used.

**[0187]** In this embodiment, the oligonucleotides are synthesized as is known in the art, and then attached to the surface of the solid support. As will be appreciated by those skilled in the art, either the 5' or 3' terminus may be attached to the solid support, or attachment may be via an internal nucleoside.

**[0188]** In an additional embodiment, the immobilization to the solid support may be very strong, yet non-covalent. For example, biotinylated oligonucleotides can be made, which bind to surfaces covalently coated with streptavidin, resulting in attachment.

**[0189]** Alternatively, the oligonucleotides may be synthesized on the surface, as is known in the art. For example, photoactivation techniques utilizing photopolymerization compounds and techniques are used. In a preferred embodiment, the nucleic acids can be synthesized in situ, using well known photolithographic techniques, such as those described in WO 95/25116; WO 95/35505; U.S. Pat. Nos. 5,700,637 and 5,445,934; and references cited within, all of which are expressly incorporated by reference; these methods of attachment form the basis of the Affimetrix GeneChip™ technology.

**[0190]** In another preferred embodiment anti-cellular proliferation antibodies are provided. In one case, the cellular proliferation protein is to be used to generate antibodies, for example for immunotherapy. Wherein a fragment of the cellular proliferation protein is used, the cellular proliferation protein should share at least one epitope or determinant with the full length protein. By "epitope" or "determinant" herein is meant a portion of a protein which will generate and/or bind

an antibody or T-cell receptor in the context of MHC. Thus, in most instances, antibodies made to a smaller cellular proliferation protein will be able to bind to the full length protein. In a preferred embodiment, the epitope is unique; that is, antibodies generated to a unique epitope show little or no cross-reactivity.

**[0191]** In one embodiment, the term "antibody" includes antibody fragments, as are known in the art, including Fab, Fab<sub>2</sub>, single chain antibodies (Fv for example), chimeric antibodies, etc., either produced by the modification of whole antibodies or those synthesized de novo using recombinant DNA technologies.

**[0192]** Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the KSP or fragment thereof or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid a, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

**[0193]** The antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro. The immunizing agent will typically include the KSP polypeptide or fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding *Monoclonal Antibodies: Principles and practice*, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

**[0194]** In one embodiment, the antibodies are bispecific antibodies. Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one

of the binding specificities is for the KSP or a fragment thereof, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit, preferably one that is tumor specific.

**[0195]** In a preferred embodiment, the antibodies to cellular proliferation are capable of reducing or eliminating the biological function of cellular proliferation, as is described below. That is, the addition of anti-KSP antibodies (either polyclonal or preferably monoclonal) to cellular proliferation (or cells containing cellular proliferation) may reduce or eliminate the cellular proliferation activity. Generally, at least a 25% decrease in activity is preferred, with at least about 50% being particularly preferred and about a 95-100% decrease being especially preferred.

**[0196]** In a preferred embodiment the antibodies to the cellular proliferation proteins are humanized antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric molecules of immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues form a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., *Nature*, 321: 522-525 (1986); Riechmann et al., *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992)].

**[0197]** Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as import residues, which are typically taken from an import variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., *Nature* 321:522-525 (1986); Riechmann et al., *Nature* 332:323-327 (1988); Verhoeyen et al. *Science*, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such humanized antibodies are chimeric antibodies (U.S. Pat. No. 4,816, 567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

**[0198]** Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.* 227:381

(1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)). Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al., *Bio/Technology* 10, 779-783 (1992); Lonberg et al., *Nature* 368 856-859 (1994); Morrison *Nature* 368, 812-13 (1994); Fishwild et al., *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

**[0199]** By immunotherapy is meant treatment of cellular proliferation with an antibody raised against cellular proliferation proteins. As used herein, immunotherapy can be passive or active. Passive immunotherapy as defined herein is the passive transfer of antibody to a recipient (patient). Active immunization is the induction of antibody and/or T-cell responses in a recipient (patient). Induction of an immune response is the result of providing the recipient with an antigen to which antibodies are raised. As appreciated by one of ordinary skill in the art, the antigen may be provided by injecting a polypeptide against which antibodies are desired to be raised into a recipient, or contacting the recipient with a nucleic acid capable of expressing the antigen and under conditions for expression of the antigen.

**[0200]** As will be appreciated by one of ordinary skill in the art, the antibody may be a competitive, non-competitive or uncompetitive inhibitor of protein binding to the cellular proliferation protein. Preferably, the antibody is also an antagonist of the cellular proliferation protein. In one aspect, when the antibody prevents the binding of other molecules to the cellular proliferation protein, the antibody prevents growth of the cell. The antibody also sensitizes the cell to cytotoxic agents, including, but not limited to TNF- $\alpha$ , TNF- $\beta$ , IL-1, INF- $\gamma$  and IL-2, or chemotherapeutic agents including 5FU, vinblastine, actinomycin D, cisplatin, methotrexate, and the like.

**[0201]** In another preferred embodiment, the antibody is conjugated to a therapeutic moiety. In one aspect the therapeutic moiety is a small molecule that modulates the activity of the cellular proliferation protein. In another aspect the therapeutic moiety modulates the activity of molecules associated with or in close proximity to the cellular proliferation protein.

**[0202]** In a preferred embodiment, the therapeutic moiety may also be a cytotoxic agent. In this method, targeting the cytotoxic agent to tumor tissue or cells, results in a reduction in the number of afflicted cells, thereby reducing symptoms associated with cellular proliferation. Cytotoxic agents are numerous and varied and include, but are not limited to, cytotoxic drugs or toxins or active fragments of such toxins. Suitable toxins and their corresponding fragments include diphtheria a chain, exotoxin a chain, ricin a chain, abrin a chain, curcin, croton, phenomycin, enomycin and the like. Cytotoxic agents also include radiochemicals made by con-

jugating radioisotopes to antibodies raised against cellular proliferation proteins, or binding of a radionuclide to a chelating agent that has been covalently attached to the antibody. Targeting the therapeutic moiety to cellular proliferation proteins not only serves to increase the local concentration of therapeutic moiety in the cellular proliferation afflicted area, but also serves to reduce deleterious side effects that may be associated with the therapeutic moiety.

**[0203]** Preferably, the antibody is conjugated to a protein which facilitates entry into the cell. In one case, the antibody enters the cell by endocytosis. In another embodiment, a nucleic acid encoding the antibody is administered to the individual or cell. The nucleic acid is identified based on the sequence of the antibody, determined by standard recombinant techniques. Moreover, wherein the cellular proliferation

protein can be targeted within a cell, i.e., the nucleus, an antibody thereto contains a signal for that target localization, i.e., a nuclear localization signal.

**[0204]** In a preferred embodiment, the cellular proliferation antibodies of the invention specifically bind to cellular proliferation proteins. By “specifically bind” herein is meant that the antibodies bind to the protein with a binding constant in the range of at least  $10^{-4}$ - $10^{-8}$   $M^{-1}$ , with a preferred range being  $10^{-7}$ - $10^{-9}$   $M^{-1}$ .

**[0205]** It is understood that the examples described above in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes. All references cited herein are incorporated by reference in their entirety as well as the sequences cited therein or having a GenBank accession number.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1

<211> LENGTH: 3741

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<223> OTHER INFORMATION: human KSP (kinesin-related protein) cDNA

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (11)..(3184)

<223> OTHER INFORMATION: KSP

<400> SEQUENCE: 1

```

gaattccgtc atggcgctgc agccaaattc gtctgcgaag aagaaagagg agaaggggaa      60
gaacatccag tgggtggtga gatgcagacc atttaatttg gcagagcgga aagctagcgc      120
ccattcaata gtagaatgtg atcctgtacg aaaagaagtt agtgtacgaa ctggaggatt      180
ggctgacaag agctcaagga aaacatacac ttttgatatg gtgtttggag catctactaa      240
acagattgat gtttaccgaa gtgtttgttg tccaattctg gatgaagtta ttatgggcta      300
taattgcact atctttgcgt atggccaaac tggcactgga aaaactttta caatggaagg      360
tgaaaggtea cctaatagaag agtatacctg ggaagaggat cccttggtcg gtataattcc      420
acgtaccctt catcaaatTT ttgagaaact tactgataat ggtactgaat ttctagtc aa      480
agtgtctctg ttggagatct ataatagaag gctttttgat cttcttaatc catcatctga      540
tgtttctgag agactacaga tgtttgatga tccccgtaac aagagaggag tgataattaa      600
aggtttagaa gaaattacag tacacaacaa ggatgaagtc tatcaaatTT tagaaaaggg      660
ggcagcaaaa aggacaactg cagctactct gatgaatgca tactctagtc gttcccactc      720
agttttctct gttacaatac atatgaaaga aactacgatt gatggagaag agcttggtta      780
aatcggaagg ttgaacttgg ttgatcttgc aggaagtga aacattggcc gttctggagc      840
tgttgataag agagctcggg aagctggaaa tataaatcaa tcctgtttga ctttgggaag      900
ggtcattact gcccttgtag aaagaacacc tcatgttctt tatcgagaat ctaaaactaa      960
tagaatcttc caggattctc ttggagggcg tacaagaaca tctataattg caacaatttc     1020
tcctgcatct ctcaatcttg aggaaactct gactacattg gaatatgctc atagagcaaa     1080
gaacatattg aataagcctg aagtgaatca gaaactcacc aaaaaagctc ttattaagga     1140

```

-continued

gtatacggag gagatagaac gtttaaaacg agatcttgct gcagcccgtg agaaaaatgg	1200
agtgatatatt tctgaagaaa attttagagt catgagtgga aaattaactg ttcaagaaga	1260
gcagattgta gaattgattg aaaaaattgg tgctgttgag gaggagctga atagggttac	1320
agagttgttt atggataata aaaatgaact tgaccagtgt aaatctgacc tgcaaaataa	1380
aacacaagaa cttgaaacca ctcaaaaaca ttgcaagaa actaaattac aacttggtta	1440
agaagaatat atcacatcag ctttggaag tactgaggag aaacttcattg atgctgccag	1500
caagctgctt aacacagttg aagaaactac aaaagatgta tctggtctcc attccaaact	1560
ggatcgtaag aaggcagttg accaacacaa tgcagaagct caggatatatt ttggcaaaaa	1620
cctgaatagt ctgtttaata atatggaaga attaattaag gatggcagct caaagcaaaa	1680
ggccatgcta gaagtacata agaccttatt tggtaatctg ctgtcttcca gtgtctctgc	1740
attagatacc attactacag tagcacttgg atctctcaca tctattccag aaaatgtgtc	1800
tactcatggt tctcagattt ttaatatgat actaaaagaa caatcattag cagcagaaaag	1860
taaaactgta ctacaggaat tgattaatgt actcaagact gatcttctaa gttcactgga	1920
aatgatttta tcccaactg tgggtgtctat actgaaaac aatagtcaac taaagcatat	1980
tttcaagact tcattgacag tggccgataa gatagaagat caaaaaaaaa ggaactcaga	2040
tggctttctc agtatactgt gtaacaatct acatgaacta caagaaaata ccatttggtc	2100
cttggttgag tcacaaaagc aatgtggaaa cctaactgaa gacctgaaga caataaagca	2160
gacccattcc caggaacttt gcaagttaat gaatctttgg acagagagat tctgtgcttt	2220
ggaggaaaaag tgtgaaaata tacagaaacc acttagtagt gtccaggaaa atatacagca	2280
gaaatctaag gatatagtca acaaaatgac ttttcacagt caaaaatttt gtgctgattc	2340
tgatggcttc tcacaggaac tcagaaattt taaccaagaa ggtacaaaat tggttgaaga	2400
atctgtgaaa cactctgata aactcaatgg caacctggaa aaaatatctc aagagactga	2460
acagagatgt gaatctctga acacaagaac agtttatctt tctgaacagt gggatatctc	2520
cttaaatgaa agggaacagg aacttcacaa cttattggag gttgtaagcc aatgttggtg	2580
ggcttcaagt tcagacatca ctgagaaac agatggacgt aaggcagctc atgagaaaca	2640
gcataacatt tttcttgatc agatgactat tgatgaagat aaattgatag caaaaatct	2700
agaacttaat gaaaccataa aaattggttt gactaagctt aattgctttc tggaacagga	2760
tctgaaactg gatatcccaa caggtaacg accacagagg aaaagttatt tatacccatc	2820
aacactggta agaactgaac cagtgaaaca tctccttgat cagctgaaaa ggaacagcc	2880
tgagctgtta atgatgctaa actgttcaga aaacaacaaa gaagagacaa ttccgatgt	2940
ggatgtagaa gaggcagttc tggggcagta tactgaagaa cctctaagtc aagagccatc	3000
tgtagatgct ggtgtggatt gttcatcaat tggcgggggt ccatttttcc agcataaaaa	3060
atcacatgga aaagacaaaag aaaacagagg cattaacaca ctggagaggt ctaaagtgga	3120
agaaactaca gagcacttgg ttacaaagag cagattacct ctgcgagccc agatcaacct	3180
ttaattcact tgggggttgg caattttatt tttaaagaaa aacttaaaaa taaaacctga	3240
aacccagaa cttgagcctt gtgtatagat tttaaagaa tatatatatc agccgggcgc	3300
gtggctctag ctgtaatccc agctaaactt ggaggctgag ggggtggat tgcttgagcc	3360
caggagtttg agaccagcct ggccaacgtg cgctaaaacc ttcgtctctg ttaaaaatta	3420

-continued

---

```

gccgggcggtg gtgggcacac tcctgtaatc ccagctactg gggaggctga ggcacgagaa 3480
tcacttgaac ccagaagcgg gggtgcagtg agccaaaggt acaccactac actccagcct 3540
gggcaacaga gcaagactcg gtctcaaaaa taaaatttaa aaaagatata aggcagtact 3600
gtaaattcag ttgaattttg atatctaccc atttttctgt catccctata gttcactttg 3660
tattaaattg gggtttcattt gggatttgca atgtaaatac gtatttctag ttttcatata 3720
aagtagttct ttttaggaatt c 3741

```

```

<210> SEQ ID NO 2
<211> LENGTH: 1057
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<223> OTHER INFORMATION: human KSP (kinesin-related protein)

```

```

<400> SEQUENCE: 2

```

```

Met Ala Ser Gln Pro Asn Ser Ser Ala Lys Lys Lys Glu Glu Lys Gly
 1             5             10             15
Lys Asn Ile Gln Val Val Val Arg Cys Arg Pro Phe Asn Leu Ala Glu
 20             25             30
Arg Lys Ala Ser Ala His Ser Ile Val Glu Cys Asp Pro Val Arg Lys
 35             40             45
Glu Val Ser Val Arg Thr Gly Gly Leu Ala Asp Lys Ser Ser Arg Lys
 50             55             60
Thr Tyr Thr Phe Asp Met Val Phe Gly Ala Ser Thr Lys Gln Ile Asp
 65             70             75             80
Val Tyr Arg Ser Val Val Cys Pro Ile Leu Asp Glu Val Ile Met Gly
 85             90             95
Tyr Asn Cys Thr Ile Phe Ala Tyr Gly Gln Thr Gly Thr Gly Lys Thr
 100            105            110
Phe Thr Met Glu Gly Glu Arg Ser Pro Asn Glu Glu Tyr Thr Trp Glu
 115            120            125
Glu Asp Pro Leu Ala Gly Ile Ile Pro Arg Thr Leu His Gln Ile Phe
 130            135            140
Glu Lys Leu Thr Asp Asn Gly Thr Glu Phe Ser Val Lys Val Ser Leu
 145            150            155            160
Leu Glu Ile Tyr Asn Glu Glu Leu Phe Asp Leu Leu Asn Pro Ser Ser
 165            170            175
Asp Val Ser Glu Arg Leu Gln Met Phe Asp Asp Pro Arg Asn Lys Arg
 180            185            190
Gly Val Ile Ile Lys Gly Leu Glu Glu Ile Thr Val His Asn Lys Asp
 195            200            205
Glu Val Tyr Gln Ile Leu Glu Lys Gly Ala Ala Lys Arg Thr Thr Ala
 210            215            220
Ala Thr Leu Met Asn Ala Tyr Ser Ser Arg Ser His Ser Val Phe Ser
 225            230            235            240
Val Thr Ile His Met Lys Glu Thr Thr Ile Asp Gly Glu Glu Leu Val
 245            250            255
Lys Ile Gly Lys Leu Asn Leu Val Asp Leu Ala Gly Ser Glu Asn Ile
 260            265            270
Gly Arg Ser Gly Ala Val Asp Lys Arg Ala Arg Glu Ala Gly Asn Ile
 275            280            285

```

Asn	Gln	Ser	Leu	Leu	Thr	Leu	Gly	Arg	Val	Ile	Thr	Ala	Leu	Val	Glu
290						295				300					
Arg	Thr	Pro	His	Val	Pro	Tyr	Arg	Glu	Ser	Lys	Leu	Thr	Arg	Ile	Leu
305				310					315						320
Gln	Asp	Ser	Leu	Gly	Gly	Arg	Thr	Arg	Thr	Ser	Ile	Ile	Ala	Thr	Ile
				325					330					335	
Ser	Pro	Ala	Ser	Leu	Asn	Leu	Glu	Glu	Thr	Leu	Ser	Thr	Leu	Glu	Tyr
			340					345					350		
Ala	His	Arg	Ala	Lys	Asn	Ile	Leu	Asn	Lys	Pro	Glu	Val	Asn	Gln	Lys
		355					360					365			
Leu	Thr	Lys	Lys	Ala	Leu	Ile	Lys	Glu	Tyr	Thr	Glu	Glu	Ile	Glu	Arg
						375					380				
Leu	Lys	Arg	Asp	Leu	Ala	Ala	Arg	Glu	Lys	Asn	Gly	Val	Tyr	Ile	
385				390					395					400	
Ser	Glu	Glu	Asn	Phe	Arg	Val	Met	Ser	Gly	Lys	Leu	Thr	Val	Gln	Glu
				405					410					415	
Glu	Gln	Ile	Val	Glu	Leu	Ile	Glu	Lys	Ile	Gly	Ala	Val	Glu	Glu	Glu
			420					425				430			
Leu	Asn	Arg	Val	Thr	Glu	Leu	Phe	Met	Asp	Asn	Lys	Asn	Glu	Leu	Asp
		435					440					445			
Gln	Cys	Lys	Ser	Asp	Leu	Gln	Asn	Lys	Thr	Gln	Glu	Leu	Glu	Thr	Thr
450						455					460				
Gln	Lys	His	Leu	Gln	Glu	Thr	Lys	Leu	Gln	Leu	Val	Lys	Glu	Glu	Tyr
465					470					475					480
Ile	Thr	Ser	Ala	Leu	Glu	Ser	Thr	Glu	Glu	Lys	Leu	His	Asp	Ala	Ala
				485					490					495	
Ser	Lys	Leu	Leu	Asn	Thr	Val	Glu	Glu	Thr	Thr	Lys	Asp	Val	Ser	Gly
				500				505					510		
Leu	His	Ser	Lys	Leu	Asp	Arg	Lys	Lys	Ala	Val	Asp	Gln	His	Asn	Ala
		515					520					525			
Glu	Ala	Gln	Asp	Ile	Phe	Gly	Lys	Asn	Leu	Asn	Ser	Leu	Phe	Asn	Asn
						535					540				
Met	Glu	Glu	Leu	Ile	Lys	Asp	Gly	Ser	Ser	Lys	Gln	Lys	Ala	Met	Leu
545				550						555					560
Glu	Val	His	Lys	Thr	Leu	Phe	Gly	Asn	Leu	Leu	Ser	Ser	Ser	Val	Ser
				565					570					575	
Ala	Leu	Asp	Thr	Ile	Thr	Thr	Val	Ala	Leu	Gly	Ser	Leu	Thr	Ser	Ile
			580					585					590		
Pro	Glu	Asn	Val	Ser	Thr	His	Val	Ser	Gln	Ile	Phe	Asn	Met	Ile	Leu
		595					600					605			
Lys	Glu	Gln	Ser	Leu	Ala	Ala	Glu	Ser	Lys	Thr	Val	Leu	Gln	Glu	Leu
					615						620				
Ile	Asn	Val	Leu	Lys	Thr	Asp	Leu	Leu	Ser	Ser	Leu	Glu	Met	Ile	Leu
625				630						635					640
Ser	Pro	Thr	Val												



-continued

690					695					700					
Cys	Gly	Asn	Leu	Thr	Glu	Asp	Leu	Lys	Thr	Ile	Lys	Gln	Thr	His	Ser
705					710					715					720
Gln	Glu	Leu	Cys	Lys	Leu	Met	Asn	Leu	Trp	Thr	Glu	Arg	Phe	Cys	Ala
				725					730					735	
Leu	Glu	Glu	Lys	Cys	Glu	Asn	Ile	Gln	Lys	Pro	Leu	Ser	Ser	Val	Gln
			740					745					750		
Glu	Asn	Ile	Gln	Gln	Lys	Ser	Lys	Asp	Ile	Val	Asn	Lys	Met	Thr	Phe
			755				760					765			
His	Ser	Gln	Lys	Phe	Cys	Ala	Asp	Ser	Asp	Gly	Phe	Ser	Gln	Glu	Leu
			770				775					780			
Arg	Asn	Phe	Asn	Gln	Glu	Gly	Thr	Lys	Leu	Val	Glu	Glu	Ser	Val	Lys
				790						795					800
His	Ser	Asp	Lys	Leu	Asn	Gly	Asn	Leu	Glu	Lys	Ile	Ser	Gln	Glu	Thr
				805					810					815	
Glu	Gln	Arg	Cys	Glu	Ser	Leu	Asn	Thr	Arg	Thr	Val	Tyr	Phe	Ser	Glu
			820					825					830		
Gln	Trp	Val	Ser	Ser	Leu	Asn	Glu	Arg	Glu	Gln	Glu	Leu	His	Asn	Leu
			835				840					845			
Leu	Glu	Val	Val	Ser	Gln	Cys	Cys	Glu	Ala	Ser	Ser	Ser	Asp	Ile	Thr
			850				855					860			
Glu	Lys	Ser	Asp	Gly	Arg	Lys	Ala	Ala	His	Glu	Lys	Gln	His	Asn	Ile
				870						875					880
Phe	Leu	Asp	Gln	Met	Thr	Ile	Asp	Glu	Asp	Lys	Leu	Ile	Ala	Gln	Asn
				885					890					895	
Leu	Glu	Leu	Asn	Glu	Thr	Ile	Lys	Ile	Gly	Leu	Thr	Lys	Leu	Asn	Cys
			900					905					910		
Phe	Leu	Glu	Gln	Asp	Leu	Lys	Leu	Asp	Ile	Pro	Thr	Gly	Thr	Thr	Pro
			915				920					925			
Gln	Arg	Lys	Ser	Tyr	Leu	Tyr	Pro	Ser	Thr	Leu	Val	Arg	Thr	Glu	Pro
			930				935					940			
Arg	Glu	His	Leu	Leu	Asp	Gln	Leu	Lys	Arg	Lys	Gln	Pro	Glu	Leu	Leu
				950						955					960
Met	Met	Leu	Asn	Cys	Ser	Glu	Asn	Asn	Lys	Glu	Glu	Thr	Ile	Pro	Asp
				965					970					975	
Val	Asp	Val	Glu	Glu	Ala	Val	Leu	Gly	Gln	Tyr	Thr	Glu	Glu	Pro	Leu
			980					985					990		
Ser	Gln	Glu	Pro	Ser	Val	Asp	Ala	Gly	Val	Asp	Cys	Ser	Ser	Ile	Gly
			995				1000					1005			
Gly	Val	Pro	Phe	Phe	Gln	His	Lys	Lys	Ser	His	Gly	Lys	Asp	Lys	Glu
			1010				1015					1020			
Asn	Arg	Gly	Ile	Asn	Thr	Leu	Glu	Arg	Ser	Lys	Val	Glu	Glu	Thr	Thr
				1030						1035					1040
Glu	His	Leu	Val	Thr	Lys	Ser	Arg	Leu	Pro	Leu	Arg	Ala	Gln	Ile	Asn
				1045					1050					1055	

Leu

<210> SEQ ID NO 3  
 <211> LENGTH: 1149  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence:KSP fragment  
KSP-L360 plus C-terminal myc epitope and  
6-histidine tag

<400> SEQUENCE: 3

```

atggcgtgcc agccaaattc gtctgcgaag aagaaagagg agaaggggaa gaacatccag      60
gtggttggtga gatgcagacc atttaatttg gcagagcgga aagctagcgc ccattcaata      120
gtagaatgtg atcctgtacg aaaagaagtt agtgtacgaa ctggaggatt ggctgacaag      180
agctcaagga aaacatacac ttttgatatg gtgtttggag catctactaa acagattgat      240
gtttaccgaa gtgtgttttg tccaattctg gatgaagtta ttatgggcta taattgcact      300
atctttgcgt atggccaaac tggcactgga aaaactttta caatggaagg tgaagggtca      360
cctaatagaag agtatacctg ggaagaggat cccttggctg gtataattcc acgtaccctt      420
catcaaattt ttgagaaact tactgataat ggtactgaat tttcagtcaa agtgtctctg      480
ttggagatct ataataaga gctttttgat cttcttaatc catcatctga tgtttctgag      540
agactacaga tgtttgatga tccccgtaac aagagaggag tgataattaa aggttttagaa      600
gaaattacag tacacaacaa ggatgaagtc tatcaaattt tagaaaaggg ggcagcaaaa      660
aggacaactg cagctactct gatgaatgca tactctagtc gttccactc agttttctct      720
gttacaatac atatgaaga aactacgatt gatggagaag agcttgtaa aatcggaag      780
ttgaacttgg ttgatcttgc aggaagtga aacattggcc gttctggagc tgttgataag      840
agagctcggg aagctggaaa tataaatcaa tccctgttga ctttgggaag ggtcattact      900
gccctttag aaagaacacc tcatgttctc tatcgagaat ctaactaac tagaatcctc      960
caggattctc ttggaggggc tacaagaaca tctataattg caacaatttc tctgcactc      1020
ctcaatcttg aggaaactct gactacattg gaatatgctc atagagcaaa gaacatattg      1080
ctcgagggtg ccgagcagaa gctgatcagc gaggaggacc tgatcgagca ccaccaccac      1140
caccactga                                         1149

```

<210> SEQ ID NO 4

<211> LENGTH: 382

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:KSP fragment  
KSP-L360 plus C-terminal myc epitope and  
6-histidine tag

<400> SEQUENCE: 4

```

Met Ala Cys Gln Pro Asn Ser Ser Ala Lys Lys Lys Glu Glu Lys Gly
 1             5             10             15

Lys Asn Ile Gln Val Val Val Arg Cys Arg Pro Phe Asn Leu Ala Glu
 20             25             30

Arg Lys Ala Ser Ala His Ser Ile Val Glu Cys Asp Pro Val Arg Lys
 35             40             45

Glu Val Ser Val Arg Thr Gly Gly Leu Ala Asp Lys Ser Ser Arg Lys
 50             55             60

Thr Tyr Thr Phe Asp Met Val Phe Gly Ala Ser Thr Lys Gln Ile Asp
 65             70             75             80

Val Tyr Arg Ser Val Val Cys Pro Ile Leu Asp Glu Val Ile Met Gly
 85             90             95

```

-continued

Tyr	Asn	Cys	Thr	Ile	Phe	Ala	Tyr	Gly	Gln	Thr	Gly	Thr	Gly	Lys	Thr
				100				105					110		
Phe	Thr	Met	Glu	Gly	Glu	Arg	Ser	Pro	Asn	Glu	Glu	Tyr	Thr	Trp	Glu
		115					120					125			
Glu	Asp	Pro	Leu	Ala	Gly	Ile	Ile	Pro	Arg	Thr	Leu	His	Gln	Ile	Phe
	130					135					140				
Glu	Lys	Leu	Thr	Asp	Asn	Gly	Thr	Glu	Phe	Ser	Val	Lys	Val	Ser	Leu
145					150					155					160
Leu	Glu	Ile	Tyr	Asn	Glu	Glu	Leu	Phe	Asp	Leu	Leu	Asn	Pro	Ser	Ser
				165					170					175	
Asp	Val	Ser	Glu	Arg	Leu	Gln	Met	Phe	Asp	Asp	Pro	Arg	Asn	Lys	Arg
			180						185				190		
Gly	Val	Ile	Ile	Lys	Gly	Leu	Glu	Glu	Ile	Thr	Val	His	Asn	Lys	Asp
		195					200					205			
Glu	Val	Tyr	Gln	Ile	Leu	Glu	Lys	Gly	Ala	Ala	Lys	Arg	Thr	Thr	Ala
	210					215					220				
Ala	Thr	Leu	Met	Asn	Ala	Tyr	Ser	Ser	Arg	Ser	His	Ser	Val	Phe	Ser
225					230					235					240
Val	Thr	Ile	His	Met	Lys	Glu	Thr	Thr	Ile	Asp	Gly	Glu	Glu	Leu	Val
				245					250					255	
Lys	Ile	Gly	Lys	Leu	Asn	Leu	Val	Asp	Leu	Ala	Gly	Ser	Glu	Asn	Ile
			260					265					270		
Gly	Arg	Ser	Gly	Ala	Val	Asp	Lys	Arg	Ala	Arg	Glu	Ala	Gly	Asn	Ile
	275						280					285			
Asn	Gln	Ser	Leu	Leu	Thr	Leu	Gly	Arg	Val	Ile	Thr	Ala	Leu	Val	Glu
	290					295					300				
Arg	Thr	Pro	His	Val	Pro	Tyr	Arg	Glu	Ser	Lys	Leu	Thr	Arg	Ile	Leu
305					310					315					320
Gln	Asp	Ser	Leu	Gly	Arg	Thr	Arg	Thr	Ser	Ile	Ile	Ala	Thr	Ile	
			325					330					335		
Ser	Pro	Ala	Ser	Leu	Asn	Leu	Glu	Glu	Thr	Leu	Ser	Thr	Leu	Glu	Tyr
			340					345					350		
Ala	His	Arg	Ala	Lys	Asn	Ile	Leu	Leu	Glu	Gly	Thr	Glu	Gln	Lys	Leu
		355					360					365			
Ile	Ser	Glu	Glu	Asp	Leu	Ile	Glu	His	His	His	His	His	His		
	370					375					380				

&lt;210&gt; SEQ ID NO 5

&lt;211&gt; LENGTH: 1542

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:KSP fragment  
KSP-K491 plus C-terminal myc epitope and  
6-histidine tag

&lt;400&gt; SEQUENCE: 5

atggcgtgcc agccaaattc gtctgcgaag aagaaagagg agaaggggaa gaacatccag	60
gtggtggtga gatgcagacc atttaatttg gcagagcgga aagctagcgc ccattcaata	120
gtagaatgtg atcctgtacg aaaagaagtt agtgtacgaa ctggaggatt ggctgacaag	180
agctcaagga aaacatacac ttttgatatg gtgtttggag catctactaa acagattgat	240
gtttaccgaa gtgtgtgttg tccaattctg gatgaagtta ttatgggcta taattgcact	300

-continued

```

atctttgcgt atggccaaac tggcactgga aaaactttta caatggaagg tgaagggtca 360
cctaataaag agtataacctg ggaagaggat cccttggtcg gtataattcc acgtaccctt 420
catcaaattt ttgagaaact tactgataat ggtactgaat ttccagtc aa agtgtctctg 480
ttggagatct ataatagaaga gctttttgat cttcttaatc catcatctga tgtttctgag 540
agactacaga tgtttgatga tccccgtaac aagagaggag tgataattaa aggttttagaa 600
gaaattacag tacacaacaa ggatgaagtc tatcaaattt tagaaaaggg ggcagcaaaa 660
aggacaactg cagctactct gatgaatgca tactctagtc gttccactc agttttctct 720
gttacaatac atatgaaaga aactacgatt gatggagaag agcttggtta aatcggaag 780
ttgaacttgg ttgatcttgc aggaagtga aacattggcc gttctggagc tgttgataag 840
agagctcggg aagctgga aa tataaatcaa tccctgttga ctttggaag ggtcattact 900
gcccttgtag aaagaacacc tcatgttct tctcgagaat ctaactaac tagaatctc 960
caggattctc ttggagggcg tacaagaaca tctataattg caacaatttc tctgcatct 1020
ctcaatcttg aggaactct gagtacattg gaatatgctc atagagcaaa gaacatattg 1080
aataagcctg aagtgaatca gaaactcacc aaaaaagctc ttattaagga gtatacggag 1140
gagatagaac gtttaaaacg agatcttgct gcagcccggtg agaaaaatgg agtgtatatt 1200
tctgaagaaa attttagagt catgagtga aaattaactg ttcaagaaga gcagattgta 1260
gaattgattg aaaaaattgg tgctgttgag gaggagctga atagggttac agagttgttt 1320
atggataata aaaatgaact tgaccagtgt aaatctgacc tgcaaaataa aacacaagaa 1380
cttgaaacca ctcaaaaaca ttgcaagaa actaaattac aacttggtta agaagaatat 1440
atcacatcag ctttggaag tactgaggag aaactcgagg gtaccgagca gaagctgac 1500
agcgaggagg acctgatcga gcaccaccac caccaccact ga 1542

```

```

<210> SEQ ID NO 6
<211> LENGTH: 513
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:KSP fragment
KSP-K491 plus C-terminal myc epitope and
6-histidine tag

```

```

<400> SEQUENCE: 6

```

```

Met Ala Cys Gln Pro Asn Ser Ser Ala Lys Lys Lys Glu Glu Lys Gly
 1             5             10            15
Lys Asn Ile Gln Val Val Val Arg Cys Arg Pro Phe Asn Leu Ala Glu
 20            25            30
Arg Lys Ala Ser Ala His Ser Ile Val Glu Cys Asp Pro Val Arg Lys
 35            40            45
Glu Val Ser Val Arg Thr Gly Gly Leu Ala Asp Lys Ser Ser Arg Lys
 50            55            60
Thr Tyr Thr Phe Asp Met Val Phe Gly Ala Ser Thr Lys Gln Ile Asp
 65            70            75            80
Val Tyr Arg Ser Val Val Cys Pro Ile Leu Asp Glu Val Ile Met Gly
 85            90            95
Tyr Asn Cys Thr Ile Phe Ala Tyr Gly Gln Thr Gly Thr Gly Lys Thr
100           105           110
Phe Thr Met Glu Gly Glu Arg Ser Pro Asn Glu Glu Tyr Thr Trp Glu

```

115					120					125						
Glu	Asp	Pro	Leu	Ala	Gly	Ile	Ile	Pro	Arg	Thr	Leu	His	Gln	Ile	Phe	
130					135					140						
Glu	Lys	Leu	Thr	Asp	Asn	Gly	Thr	Glu	Phe	Ser	Val	Lys	Val	Ser	Leu	
145					150					155					160	
Leu	Glu	Ile	Tyr	Asn	Glu	Glu	Leu	Phe	Asp	Leu	Leu	Asn	Pro	Ser	Ser	
165					170					175						
Asp	Val	Ser	Glu	Arg	Leu	Gln	Met	Phe	Asp	Asp	Pro	Arg	Asn	Lys	Arg	
180					185					190						
Gly	Val	Ile	Ile	Lys	Gly	Leu	Glu	Glu	Ile	Thr	Val	His	Asn	Lys	Asp	
195					200					205						
Glu	Val	Tyr	Gln	Ile	Leu	Glu	Lys	Gly	Ala	Ala	Lys	Arg	Thr	Thr	Ala	
210					215					220						
Ala	Thr	Leu	Met	Asn	Ala	Tyr	Ser	Ser	Arg	Ser	His	Ser	Val	Phe	Ser	
225					230					235					240	
Val	Thr	Ile	His	Met	Lys	Glu	Thr	Thr	Ile	Asp	Gly	Glu	Glu	Leu	Val	
245					250					255						
Lys	Ile	Gly	Lys	Leu	Asn	Leu	Val	Asp	Leu	Ala	Gly	Ser	Glu	Asn	Ile	
260					265					270						
Gly	Arg	Ser	Gly	Ala	Val	Asp	Lys	Arg	Ala	Arg	Glu	Ala	Gly	Asn	Ile	
275					280					285						
Asn	Gln	Ser	Leu	Leu	Thr	Leu	Gly	Arg	Val	Ile	Thr	Ala	Leu	Val	Glu	
290					295					300						
Arg	Thr	Pro	His	Val	Pro	Tyr	Arg	Glu	Ser	Lys	Leu	Thr	Arg	Ile	Leu	
305					310					315					320	
Gln	Asp	Ser	Leu	Gly	Gly	Arg	Thr	Arg	Thr	Ser	Ile	Ile	Ala	Thr	Ile	
325					330					335						
Ser	Pro	Ala	Ser	Leu	Asn	Leu	Glu	Glu	Thr	Leu	Ser	Thr	Leu	Glu	Tyr	
340					345					350						
Ala	His	Arg	Ala	Lys	Asn	Ile	Leu	Asn	Lys	Pro	Glu	Val	Asn	Gln	Lys	
355					360					365						
Leu	Thr	Lys	Lys	Ala	Leu	Ile	Lys	Glu	Tyr	Thr	Glu	Glu	Ile	Glu	Arg	
370					375					380						
Leu	Lys	Arg	Asp	Leu	Ala	Ala	Ala	Arg	Glu	Lys	Asn	Gly	Val	Tyr	Ile	
385					390					395					400	
Ser	Glu	Glu	Asn	Phe	Arg	Val	Met	Ser	Gly	Lys	Leu	Thr	Val	Gln	Glu	
405					410					415						
Glu	Gln	Ile	Val	Glu	Leu	Ile	Glu	Lys	Ile	Gly	Ala	Val	Glu	Glu	Glu	
420					425					430						
Leu	Asn	Arg	Val	Thr	Glu	Leu	Phe	Met	Asp	Asn	Lys	Asn	Glu	Leu	Asp	
435					440					445						
Gln	Cys	Lys	Ser	Asp	Leu	Gln	Asn	Lys	Thr	Gln	Glu	Leu	Glu	Thr	Thr	
450					455					460						
Gln	Lys	His	Leu	Gln	Glu	Thr	Lys	Leu	Gln	Leu	Val	Lys	Glu	Glu	Tyr	
465					470					475					480	
Ile	Thr	Ser	Ala	Leu	Glu	Ser	Thr	Glu	Glu	Lys	Leu	Glu	Gly	Thr	Glu	
485					490					495						
Gln	Lys	Leu	Ile	Ser	Glu	Glu	Asp	Leu	Ile	Glu	His	His	His	His	His	
500					505					510						
His																

---

-continued

---

<210> SEQ ID NO 7  
<211> LENGTH: 1728  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence:KSP fragment  
KSP-S553 plus C-terminal myc epitope and  
6-histidine tag

<400> SEQUENCE: 7

```
atggcgtgcc agccaaatcc gtctgcgaag aagaaagagg agaaggggaa gaacatccag    60
gtggtgtgtga gatgcagacc atttaatttg gcagagcgga aagctagcgc ccattcaata    120
gtagaatgtg atcctgtacg aaaagaagtt agtgtacgaa ctggaggatt ggctgacaag    180
agctcaagga aaacatacac ttttgatatg gtgtttggag catctactaa acagattgat    240
gtttaccgaa gtgttggttg tccaattctg gatgaagtta ttatgggcta taattgcact    300
atctttgcgt atggccaaac tggcactgga aaaactttta caatggaagg tgaagggtca    360
cctaatagaag agtatacctg ggaagaggat cccttggtcg gtataattcc acgtaccctt    420
catcaaattt ttgagaaact tactgataat ggtactgaat tttcagtcaa agtgtctctg    480
ttggagatct ataataaga gctttttgat cttcttaatc catcatctga tgtttctgag    540
agactacaga tgtttgatga tcccgtaac aagagaggag tgataattaa aggttttagaa    600
gaaattacag tacacaacaa ggatgaagtc tatcaaattt tagaaaaggg ggcagcaaaa    660
aggacaactg cagctactct gatgaatgca tactctagtc gttccactc agttttctct    720
gttacaatac atatgaaga aactacgatt gatggagaag agcttgtaa aatcggaag    780
ttgaacttgg ttgatcttgc aggaagtga aacattggcc gttctggagc tgttgataag    840
agagctcggg aagctggaat tataaatcaa tccctgttga ctttgggaag ggtcattact    900
gccctttag aaagaacacc tcatgttctc tatcgagaat ctaaaactaac tagaatcctc    960
caggattctc ttggagggcg tacaagaaca tctataattg caacaatttc tctgcatct    1020
ctcaatcttg aggaactct gagtacattg gaatatgctc atagagcaaa gaacatattg    1080
aataagcctg aagtgaatca gaaactcacc aaaaaagctc ttattaagga gtatacggag    1140
gagatagaac gtttaaaacg agatcttgct gcagcccgtg agaaaaatgg agtgtatatt    1200
tctgaagaaa attttagagt catgagtgga aaattaactg ttcaagaaga gcagattgta    1260
gaattgattg aaaaaattgg tgctgttgag gaggagctga ataggggtac agagttgttt    1320
atggataata aaaaatgaac tgaccagtgt aaatctgacc tgcaaaataa aacacaagaa    1380
cttgaaacca ctcaaaaaca tttgcaagaa actaaattac aacttgtaa agaagaatat    1440
atcacatcag ctttggaag tactgaggag aaacttcag atgctgccag caagctgctt    1500
aacacagttg aagaaactac aaaagatgta tctggtctcc attccaaact ggatcgtaag    1560
aaggcagttg accaacacaa tgcagaagct caggatattt ttggcaaaaa cctgaatagt    1620
ctgtttaata atatggaaga attaatgaag gatggcagcc tcgagggtac cgagcagaag    1680
ctgatcagcg aggaggacct gatcgagcac caccaccacc accactga    1728
```

<210> SEQ ID NO 8  
<211> LENGTH: 575  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Description of Artificial Sequence:KSP fragment  
KSP-S553 plus C-terminal myc epitope and  
6-histidine tag

<400> SEQUENCE: 8

```

Met Ala Cys Gln Pro Asn Ser Ser Ala Lys Lys Lys Glu Glu Lys Gly
 1           5           10           15

Lys Asn Ile Gln Val Val Val Arg Cys Arg Pro Phe Asn Leu Ala Glu
 20           25           30

Arg Lys Ala Ser Ala His Ser Ile Val Glu Cys Asp Pro Val Arg Lys
 35           40           45

Glu Val Ser Val Arg Thr Gly Gly Leu Ala Asp Lys Ser Ser Arg Lys
 50           55           60

Thr Tyr Thr Phe Asp Met Val Phe Gly Ala Ser Thr Lys Gln Ile Asp
 65           70           75           80

Val Tyr Arg Ser Val Val Cys Pro Ile Leu Asp Glu Val Ile Met Gly
 85           90           95

Tyr Asn Cys Thr Ile Phe Ala Tyr Gly Gln Thr Gly Thr Gly Lys Thr
100           105           110

Phe Thr Met Glu Gly Glu Arg Ser Pro Asn Glu Glu Tyr Thr Trp Glu
115           120           125

Glu Asp Pro Leu Ala Gly Ile Ile Pro Arg Thr Leu His Gln Ile Phe
130           135           140

Glu Lys Leu Thr Asp Asn Gly Thr Glu Phe Ser Val Lys Val Ser Leu
145           150           155           160

Leu Glu Ile Tyr Asn Glu Glu Leu Phe Asp Leu Leu Asn Pro Ser Ser
165           170           175

Asp Val Ser Glu Arg Leu Gln Met Phe Asp Asp Pro Arg Asn Lys Arg
180           185           190

Gly Val Ile Ile Lys Gly Leu Glu Glu Ile Thr Val His Asn Lys Asp
195           200           205

Glu Val Tyr Gln Ile Leu Glu Lys Gly Ala Ala Lys Arg Thr Thr Ala
210           215           220

Ala Thr Leu Met Asn Ala Tyr Ser Ser Arg Ser His Ser Val Phe Ser
225           230           235           240

Val Thr Ile His Met Lys Glu Thr Thr Ile Asp Gly Glu Glu Leu Val
245           250           255

Lys Ile Gly Lys Leu Asn Leu Val Asp Leu Ala Gly Ser Glu Asn Ile
260           265           270

Gly Arg Ser Gly Ala Val Asp Lys Arg Ala Arg Glu Ala Gly Asn Ile
275           280           285

Asn Gln Ser Leu Leu Thr Leu Gly Arg Val Ile Thr Ala Leu Val Glu
290           295           300

Arg Thr Pro His Val Pro Tyr Arg Glu Ser Lys Leu Thr Arg Ile Leu
305           310           315           320

Gln Asp Ser Leu Gly Gly Arg Thr Arg Thr Ser Ile Ile Ala Thr Ile
325           330           335

Ser Pro Ala Ser Leu Asn Leu Glu Glu Thr Leu Ser Thr Leu Glu Tyr
340           345           350

Ala His Arg Ala Lys Asn Ile Leu Asn Lys Pro Glu Val Asn Gln Lys
355           360           365

Leu Thr Lys Lys Ala Leu Ile Lys Glu Tyr Thr Glu Glu Ile Glu Arg

```

-continued

370	375	380	
Leu Lys Arg Asp	Leu Ala Ala Ala Arg	Glu Lys Asn Gly Val Tyr Ile	
385	390	395	400
Ser Glu Glu Asn Phe	Arg Val Met Ser Gly Lys Leu Thr Val Gln Glu		
	405	410	415
Glu Gln Ile Val Glu	Leu Ile Glu Lys Ile Gly Ala Val Glu Glu Glu		
	420	425	430
Leu Asn Arg Val Thr	Glu Leu Phe Met Asp Asn Lys Asn Glu Leu Asp		
	435	440	445
Gln Cys Lys Ser Asp	Leu Gln Asn Lys Thr Gln Glu Leu Glu Thr Thr		
	450	455	460
Gln Lys His Leu Gln	Glu Thr Lys Leu Gln Leu Val Lys Glu Glu Tyr		
	465	470	475
Ile Thr Ser Ala Leu	Glu Ser Thr Glu Glu Lys Leu His Asp Ala Ala		
	485	490	495
Ser Lys Leu Leu Asn	Thr Val Glu Glu Thr Thr Lys Asp Val Ser Gly		
	500	505	510
Leu His Ser Lys Leu	Asp Arg Lys Lys Ala Val Asp Gln His Asn Ala		
	515	520	525
Glu Ala Gln Asp Ile	Phe Gly Lys Asn Leu Asn Ser Leu Phe Asn Asn		
	530	535	540
Met Glu Glu Leu Ile	Lys Asp Gly Ser Leu Glu Gly Thr Glu Gln Lys		
	545	550	555
Leu Ile Ser Glu Glu	Asp Leu Ile Glu His His His His His His		
	565	570	575
<210> SEQ ID NO 9			
<211> LENGTH: 1107			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Description of Artificial Sequence:KSP fragment			
KSP-K368			
<400> SEQUENCE: 9			
atggcgtgcc agccaaatc	gtctgcgaag aagaaagagg	agaaggggaa gaacatccag	60
gtggtgtgtga gatgcagacc	atttaatttg gcagagcgga	aagctagcgc ccattcaata	120
gtagaatgtg atcctgtacg	aaaagaagtt agtgtacgaa	ctggaggatt ggctgacaag	180
agctcaagga aaacatacac	ttttgatatg gtgtttggag	catctactaa acagattgat	240
gtttaccgaa gtgtttgttg	tccaattctg gatgaagtta	ttatgggcta taattgcact	300
atctttgcgt atggccaaac	tggcactgga aaaactttta	caatggaagg tgaaaggtca	360
cctaataaag agtataacctg	ggaagaggat cccttggtcg	gtataattcc acgtaccctt	420
catcaaattt ttgagaaact	tactgataat ggtactgaat	tttcagtcaa agtgtctctg	480
ttggagatct ataataaaga	gctttttgat cttcttaatc	catcatctga tgtttctgag	540
agactacaga tgtttgatga	tccccgtaac aagagaggag	tgataattaa aggtttagaa	600
gaaattacag tacacaacaa	ggatgaagtc tatcaaat	tagaaaaggg ggcagcaaaa	660
aggacaactg cagctactct	gatgaatgca tactctagtc	gttcccactc agttttctct	720
gttacaatac atatgaaaga	aactacgatt gatggagaag	agcttggttaa aatcggaag	780
ttgaacttgg ttgatcttgc	aggaagtga aacattggcc	gttctggagc tgttgataag	840



-continued

```

agagctcggg aagctggaaa tataaatcaa tccctgttga ctttgggaag ggtcattact    900
gcccttgtag aaagaacacc tcatgttctt tatcgagaat ctaaaactaac tagaatcctc    960
caggattctc ttggagggcg tacaagaaca tctataattg caacaatttc tctgcatct    1020
ctcaatcttg aggaaactct gagtacattg gaatatgctc atagagcaaa gaacatattg    1080
aataagcctg aagtgaatca gaaatag                                          1107

```

```

<210> SEQ ID NO 10
<211> LENGTH: 368
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:KSP fragment
KSP-K368

```

```

<400> SEQUENCE: 10

```

```

Met Ala Cys Gln Pro Asn Ser Ser Ala Lys Lys Lys Glu Glu Lys Gly
 1           5           10          15
Lys Asn Ile Gln Val Val Val Arg Cys Arg Pro Phe Asn Leu Ala Glu
 20          25          30
Arg Lys Ala Ser Ala His Ser Ile Val Glu Cys Asp Pro Val Arg Lys
 35          40          45
Glu Val Ser Val Arg Thr Gly Gly Leu Ala Asp Lys Ser Ser Arg Lys
 50          55          60
Thr Tyr Thr Phe Asp Met Val Phe Gly Ala Ser Thr Lys Gln Ile Asp
 65          70          75          80
Val Tyr Arg Ser Val Val Cys Pro Ile Leu Asp Glu Val Ile Met Gly
 85          90          95
Tyr Asn Cys Thr Ile Phe Ala Tyr Gly Gln Thr Gly Thr Gly Lys Thr
 100         105         110
Phe Thr Met Glu Gly Glu Arg Ser Pro Asn Glu Glu Tyr Thr Trp Glu
 115         120         125
Glu Asp Pro Leu Ala Gly Ile Ile Pro Arg Thr Leu His Gln Ile Phe
 130         135         140
Glu Lys Leu Thr Asp Asn Gly Thr Glu Phe Ser Val Lys Val Ser Leu
 145         150         155         160
Leu Glu Ile Tyr Asn Glu Glu Leu Phe Asp Leu Leu Asn Pro Ser Ser
 165         170         175
Asp Val Ser Glu Arg Leu Gln Met Phe Asp Asp Pro Arg Asn Lys Arg
 180         185         190
Gly Val Ile Ile Lys Gly Leu Glu Glu Ile Thr Val His Asn Lys Asp
 195         200         205
Glu Val Tyr Gln Ile Leu Glu Lys Gly Ala Ala Lys Arg Thr Thr Ala
 210         215         220
Ala Thr Leu Met Asn Ala Tyr Ser Ser Arg Ser His Ser Val Phe Ser
 225         230         235         240
Val Thr Ile His Met Lys Glu Thr Thr Ile Asp Gly Glu Glu Leu Val
 245         250         255
Lys Ile Gly Lys Leu Asn Leu Val Asp Leu Ala Gly Ser Glu Asn Ile
 260         265         270
Gly Arg Ser Gly Ala Val Asp Lys Arg Ala Arg Glu Ala Gly Asn Ile
 275         280         285

```

-continued

---

```

Asn Gln Ser Leu Leu Thr Leu Gly Arg Val Ile Thr Ala Leu Val Glu
 290                295                300

Arg Thr Pro His Val Pro Tyr Arg Glu Ser Lys Leu Thr Arg Ile Leu
305                310                315                320

Gln Asp Ser Leu Gly Gly Arg Thr Arg Thr Ser Ile Ile Ala Thr Ile
                325                330                335

Ser Pro Ala Ser Leu Asn Leu Glu Glu Thr Leu Ser Thr Leu Glu Tyr
                340                345                350

Ala His Arg Ala Lys Asn Ile Leu Asn Lys Pro Glu Val Asn Gln Lys
 355                360                365

```

---

```

<210> SEQ ID NO 11
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:6-histidine
        epitope tag

<400> SEQUENCE: 11

His His His His His His
1                5

```

---

1-33. (canceled)

**34.** A method of evaluating a hyper-proliferative disorder comprising:

- (a) determining the level of expression of kindle-like spindle protein (KSP) in a sample from a tissue having the hyper-proliferative disorder; and
- (b) comparing said KSP expression level to a standard or control level of expression, wherein an increase indicates that the hyper-proliferative disorder is KSP-mediated.

**35.** The method of claim **34** wherein said hyper-proliferative disorder is cancer.

36-59. (canceled)

**60.** The method of claim **34**, wherein the sample is selected from the group consisting of a blood sample, a urine sample, a buccal sample, a PAP smear, a cerebral spinal fluid sample, a breast tissue sample, a lung tissue sample, and a colon tissue sample.

**61.** The method of claim **34**, wherein the standard or control level is representative of normal cells not in a hyper-proliferative state.

**62.** The method of claim **61**, wherein the hyper-proliferative disorder is cancer, and step (b) is performed by nucleic acid hybridization.

**63.** The method of claim **34**, wherein step (a) further comprises determining the expression level of at least one additional cell proliferation protein; and step (b) comprises the expression levels of each cell proliferation protein with the expression level of the same protein in the standard or control level.

**64.** The method of claim **63**, wherein the additional cell proliferation protein is a kinesin.

**65.** The method of claim **63**, wherein the standard or control level for each cell proliferation protein is representative of normal cells not in a hyper-proliferative state.

**66.** The method of claim **65**, wherein the hyper-proliferative disorder is cancer.

**67.** The method of claim **34**, wherein step (a) comprises determining the amount of nucleic acid encoding KSP in the sample.

**68.** The method of claim **67**, wherein the amount of nucleic acid encoding KSP is determined by nucleic acid hybridization.

**69.** The method of claim **67**, wherein the nucleic acid hybridization is in situ hybridization.

**70.** The method of claim **67**, wherein the nucleic acid is DNA.

**71.** The method of claim **67**, wherein the nucleic acid is RNA.

**72.** The method of claim **34**, wherein step (a) comprises determining the amount of KSP protein in the sample.

**73.** The method of claim **72**, wherein the amount of KSP protein is determined by mass spectroscopy.

**74.** The method of claim **72**, wherein the amount of KSP protein is determined by an immunological method.

**75.** The method of claim **74**, wherein the immunological method is an enzyme-linked immunoassay (ELISA).

**76.** The method of claim **72**, wherein the amount of KSP protein is determined by two-dimensional gel electrophoresis.

**77.** The method of claim **34**, further comprising the step of administering a therapeutically effective amount of a KSP inhibitor, when an increase is detected in step (b).

**78.** The method of claim **77**, wherein the hyper-proliferative disorder is cancer.

**79.** The method of claim **77**, wherein the KSP inhibitor is a small molecule having a molecular weight of between 50 kD and 2000 kD.

\* \* \* \* \*

