



US007368255B2

(12) **United States Patent**
Bae et al.

(10) **Patent No.:** **US 7,368,255 B2**
(45) **Date of Patent:** **May 6, 2008**

(54) **RUNX3 GENE SHOWING ANTI-TUMOR ACTIVITY AND USE THEREOF**

(76) Inventors: **Suk-Chul Bae**, 107-302 Samik Apt., Gaesin-dong, Heungduck-ku, Chungcheongbuk-do (KR) 361-240; **Yoshiaki Ito**, 15-14 Higashidacho, Kamitakano, Sakyo-ku, Kyoto (JP) 606-0078

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 418 days.

(21) Appl. No.: **10/466,938**

(22) PCT Filed: **Jan. 30, 2001**

(86) PCT No.: **PCT/KR01/00121**

§ 371 (c)(1),
(2), (4) Date: **Jul. 18, 2003**

(87) PCT Pub. No.: **WO02/061069**

PCT Pub. Date: **Aug. 8, 2002**

(65) **Prior Publication Data**

US 2004/0146986 A1 Jul. 29, 2004

(30) **Foreign Application Priority Data**

Jan. 29, 2001 (KR) 2001-4018

(51) **Int. Cl.**
G01N 33/574 (2006.01)

(52) **U.S. Cl.** **435/7.23**; 435/6

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

OTHER PUBLICATIONS

Levanon et al, *Oncogene* 23:4211-4219, 2004.*
Kyeong-Sook Lee, et al., Runx2 Is a Common Target of Transforming Growth Factor β 1 and Bone Morphogenetic Protein 2, and Cooperation between Runx2 and Smad5 Induces Osteoblast-Specific Gene Expression in the Pluripotent Mesenchymal Precursor Cell Line C2C12, *Molecular and Cellular Biology*, Dec. 2000, p. 8783-8792.

Scott C. Kogen, et al., The *PEBP2 β MYH11* Fusion Created Inv(16)(p13;q22) in Myeloid Leukemia Impairs Neutrophil Maturation and Contributes to Granulocytic Dysplasia, *Proc. Natl. Acad. Sci. USA* vol. 95, pp. 11863-11868, Sep. 1998.

S-C. Bae, et al., Regulation Mechanisms for the Heterodimeric Transcription Factor, PEBP2/CBF, *Histol Histopathol* (1999) 14: 1213-1221.

Suk Chul Bae, et al., Isolation of *PEBP2 alpha β* cDNA Representing the Mouse Homolog of Human Acute Myeloid Leukemia Gene, *AML1*, *Oncogene* (1993), 8, 809-814.

Mee-Young, Ahn, et al., Comparison of the Human Genomic Structure of the Runt Domain-Encoding PEBP2/CBF α Gene Family, *Gene*, 168 (1996) 279-280.

Amjad Javed, et al., Groucho/TLE/R-esp Proteins Associate with the Nuclear Matrix and Repress RUNX (CBF α /AML/PEBP2 α) Dependent Activation of Tissue-Specific Gene Transcription, *Journal of Cell Science* 113, 2221-2231 (2000).

Yen-Yee Tang, et al., Energetic and Functional Contribution of Residues in the Core Binding Factor β (CBF β) Subunit to Heterodimerization with CBF α , *The Journal of Biological Chemistry*, vol. 275, No. 50, p. 39579-39588, 2000.

* cited by examiner

Primary Examiner—Misook Yu
Assistant Examiner—Mark Halvorson
(74) *Attorney, Agent, or Firm*—Lucas & Mercanti, LLP

(57) **ABSTRACT**

The present invention relates to a RUNX3 gene showing anti-tumor activity which is essentially involved in TGF- β dependent-programmed cell death (apoptosis) and use thereof. In addition, the present invention finds that the RUNX3 gene expression is suppressed in the various gastric cancer and lung cancer cell lines. The suppression of the RUNX3 gene expression is due to hyper-methylation of CpG island located around RUNX3 exon (1). The RUNX3 gene and its gene product of the present invention can be used effectively for the development of anti-cancer agents. CpG island around RUNX3 exon (1) could also be used not only for the development of anti-cancer agents which regulate the abnormal DNA methylation and there by induce RUNX3 expression but also for the development of methods for cancer diagnosis by measuring the abnormal DNA methylation.

7 Claims, 17 Drawing Sheets

FIG. 1a

CCGCCACTTGATTCCTGGAGGATTTGTTCTGGGGCTGCGGCCGCGGAGTCG
 Ps-NA
 GGGCGGCCGCGGGCGAGCTTCGGGGCGGGAGGCGGGCGCAGCGGCACAGC 100
 CCCGCGCGGGCCCCCGCCGCGGCCAGGCAGCCGGGACAGCCACGAGGGGC
 GGCAGCAGCGGGCCGCGCGCCGAGGATGCGGGACTAGCCGGGCAGGCT 200
 GCGGGCGGCCGTGCGGCCAGCGAGGCCCTCGCAGCGGGCGGGCCCTGGCGA
 GTATTGGCCGGGCGCGGCCCTCGGCCCTGATGCCCGGGCCCGCCGCT 300
 TCTGCTTCCCGTTCTCGCGGCAGCGGGCGCCGAGGAGCGCCCGCGCC
 GGCGGCCCGGGGGAAGCCGCGCCGTCTCCGCCTGCCCGGCCTGAC 400
 GGCCGCTGTATGCGTATTCCCGTAGACCCAAGCACCAGCCGCGCTTCA
 CACCTCCCTCCCGGCCCTTCCCTGCGGCGGGCGGGCGGCAAGATGGGC 500
 GAGAACAGCGGCCCGCTGAGCGCGCAGCGGGCCGTGGGGCCCGGAGGGCG
 CGCCCGGCCGAGGTGCGCTCGATGGTGGACGTGCTGCGGACCACGCAG 600
 CGGACTCGTGCGCACCGACAGCCCCAACTTCCTCGCTCCGTGCTGCCC
 TCGCACTGGCGTGAACAAGACGCTGCCCGTCCCTCAAGGTGGTGGC 700
 AATTGGGGACGTGCCGGATGGTACGGTGGTACTGTGATGGCAGGCAATG
 ACGAGAATACTCCGCTGAGCTGCGCAATGCCTCGGCCGTGATGAAGAAC 800
 CAGGTGGCCAGGTTCAACGACCTTCGCTTCGTGGGCCGAGTGGGCGAGG
 GAAGAGTTTACCCTGACCATCACTGTGTTACCAACCCCAACCAAGTGG 900
 Ps-CA
 CGACCTACCACCGAGCCATCAAGGTGACCGTGGACGGACCCCGGGAGCCC
 AGACGGCACCGCAGAAGCTGGAGGACCAGACCAAGCCGTTCCCTGACCG 1000
 CTTTGGGGACCTGGAACGGCTGCGCATGCGGGTGACACCGAGCACACCCA
 GCCCCGAGGCCTCACTCAGCACACAAGCCACTTCAGCAGCCAGCCCCAG 1100
 Ps-NB
 ACCCCAATCCAAGGCACCTCGGAACGAACCCATTCTCCGACCCCGCCA
 GTTTGACCGCTCCTTCCCCACGCTGCCAACCCCTCACGGAGAGCCGCTTCC 1200
 CAGACCCAGGATGCATTATCCCGGGGCCATGTCAGCTGCCTTCCCTTAC
 AGCGCCACGCCCTCGGGCAGGAGCATCAGCAGCCTCAGCGTGGCGGGCAT 1300
 GCCGGCCACCAGCCGCTTCCACCATACCTACCTCCCGCCACCCCTACCCGG
 GGGCCCCGAGAACCAGAGCGGGCCCTTCCAGGCCAACCCGTCCCCCTAC 1400
 CACCTCTACTACGGGACATCCTCTGGCTCCTACCAGTTCTCCATGGTGGC
 CGGCAGCAGCAGTGGGGGCGACCGCTCACCTACCCGCATGCTGGCTCTT 1500
 GCACCAGCAGCGCTGCCTCTGTGCGCCGCGGCAACCTCATGAACCCAGC
 CTGGGGCGCCAGAGTGATGGCGTGGAGGCCGACGGCAGCCACAGCAACTC 1600
 ACCACGGCCCTGAGCACGCCAGGCCGCATGGATGAGGCCGTGTGGCGGC
 CCTACTGACCGCCCTGGTGGACTCCTCCCGCTGGAGGCGGGGACCCTAAC 1700
 AACCTTCAAGACCAGTGATGGGCCGGCTCCGAGGCTCCGGGCGGGAATGG
 Ps-CB
 GACCTGCGCTCCAGGTTGGTCTCGGTCCAGGGTGGTCCCAGCTGGTGGG 1800
 AGCCTCTGGCTGCATCTGTGCAGCCACATCCTTGTACAGAGGCATAGGTT
 ACCACCCCAACCCCGGCCCGGGATACTGCCCCCGGCCAGATCCTGGCCG 1900
 TCTCATCCCATACTTCTGTGGGAATCAGCCTCCTGCCACCCCCCGGAA
 GGACCTCACTGTCTCCAGCTATGCCAGTGTGCATGGGACCCATGTCTC 2000
 CTGGGACAGAGGCCATCTCTTCCAGAGAGAGGCAGCATTTGGCCACAG
 GATAAGCCTCAGGCCCTGGGAAACCTCCCGACCCCTGCACCTCGTTGGA 2100
 GCCCTGCATCCCTGGGTCCAGCCCCCTCTGCATTTACACAGATTTGAG
 TCAGAACTGGAAGTGTCCCCCACCCCAACCCCTCGAGCGGGGTCCC 2200
 CTCATTGTACAGATGGGGCAGGACCCAGCACGCTGCTGGCAGAGATGGTT
 TGAGAACACATCCAAGCCAGTCCCCCAGCCAGCTTCCCTCCGTTCCCT 2300
 AACTGTTGGCTTCCCCAGCCGACGGGTCCCAGGCCCAAGAGAAGATG
 AGTCTATGGCATCAGGTTCTTAAACCCAGGAAAGCACCTACAGACCGGCT 2400
 CCTCATGCACTTTACCAGCTCAACGCATCCACTCTCTGTTCTTTGGCA
 GGGCGGGGAGGGGGGATAGGAGGTCCTTCCCTAGGTGGTCTCATA 2500
 ATTCCATTTGTGGAGAGAACAGGAGGGCCAGATAGATAGGTCTTAGCAGA
 AGGCATTGAGGTGAGGGATCATTTGGGTGAGACATCAATGTCCCTGTCC 2600
 CCCCTGGGTCCAGCCAAGCTGTGCCCATCCCCAAGCCTCCTGGGAGGA

FIG. 1b

TCCAGCCAAATCTTGCGACTCCTGGCACACACCTGTCTGTAACCTGTTTT 2700
GTGCTCTGAAAGCAAATAGTCCTGAGCAAAAAAAAAAAAAAAAAACAAAAA
ACAAAAAAAAAACAAAACAGTTTTTAAAACCTGATTTTAGAAAAAGAAGCT 2800
TAATCTAACGTTTTCAACACAAGGTCTCTTACAGGTATAGTTCCGTGAT
TATGATAGCTCTGTGATTATAAGCAACATCCCCGCCCTCTCCCCCG 2900
CGGACCCCGAGCTGCCTCCTGAGGGTGTGGGGTTATTAGGGTCTCAATAC
TTTCTCAAGGGGCTACACTCCCATCAGGCAGCATCCCACCAGCCTGCAC 3000
CACAGGCTCCCCTGGGAGGACGAGGGAAACGCTGATGAGACGCTGGGCAT
CTCTCCTCTGTGGCTCTAGGACATCTGTCCAGGAGGCTGGGCGGAGGTGG 3100
GCAGGATGTGAGAGGTGGGGAGTACTGGCTGTGCGTGGCAGGACAGAAGC
ACTGTAAGGGCTCTCCAGCCGCAGCTCAGCTGCACTGCGTCCGAGGTG 3200
AAGTCTTGCCCTGAATTTTGCAAATGGGAAAGTGGGCGCTTGCCCAAG
GGCCAGGCTGCATGGATTCTCAGCATCAGAGTTCTCTGGCCCTAGAAAGGC 3300
TTAGAAAAGGCGTAAGGGAACCTATAAAGGCTAGCAGCATGCGGTATTTT
AACTTTCTGCCTCGGCCTCTGTGGATGCAGAAATCTGCCCTACAAAATGC 3400
TCTTCATTGGTTGTCTCTGTGAGAGCACTGTCCCCACCCAACCTGTCACA
ACGGCCAGAACCATACACCAGAGACACACTGGCAGGTTAGGCAGTCCTTC 3500
TGGTGATCCTATTCCATTCCCTCCTGCTGCGGTTTCTCTTGGCCTGTCT
CACTGGAAAAACAGTCTCCATCTCCTCAAATAGTTGCTGACTCCCTGCA 3600
CCCAAGGGGCCTCTCCATGCCTTCTTAGGAAGCAGCTATGAATCCATTGT
CCTTGTAGTTTCTTCCCTCCTGTTCTCTGGTTATAGCTGGTCCCAGGTCA 3700
GCGTGGGAGGCACCTTTGGGTTCCAGTGCCAGCACTTTGTAGTCTCAT
CCCAGATTACTAACCCTTCTGATCCTGGAGAGGCAGGGATAGTAAATAA 3800
ATTGCTCTTCCATACCCATCCCCATCCCCTGACAAAAGTGACGGCAGC
CGTACTGAGTCTGTAAGGCCCAAAGTGGGTACAGACAGCCTGGGCTGGTA 3900
AAAGTAGGTCCTTATTTACAAGGCTGCGTTAAAGTTGTAAGGCAAACA
CACTGATGTAGGAAGCACGAGGAAAGGAAGACGTTTTGATATAGTGTTAC 4000
TGTGAGCCTGTCAGTAGTGGGTACCAATCTTTTGTGACATATTGTCATGC
TGAGGTGTGACACCTGCTGCACTCATCTGATGTAACCATCCCAGAGCT 4100
GGCGAGAGGATGGAGCTGGGTGGAACTGCTTTGCACTATCGTTTGCTTG
GTGTTTGTTTTTAACGCACAACCTTGCTTGTACAGTAAACTGTCTTCTGTA 4200
CTATTTAACTGTA 3'

FIG. 1c

MRI PVDPSTSRRFTPPSPAFFPCGGGGGKMGENSGALSAQAAVGPGRARP
EVRSMVDVLADHAGELVRTDSPNFLCSVLPSHWRCNKTLPVAFKVVALGD
VPDGTVVTVMAGNDENYSAELRNASAVMKNQVARFNDLRFVGRSGRGKSF
TLTITVFTNPTQVATYHRAIKVTVDGPREPRRHRQKLEDQTKPFPDRFGD
LERLRMRVTPSTPSPRGSLSTTSHFSSQPQTPIQGTSELNPFSDPRQFDR
SFPTLPTLTESRFPDPRMHYPGAMSAAFPYSATPSGTSISSLSVAGMPAT
SRFHHTYLPPPYPGAPQNQSGPFQANPSPYHLYYGTSSGSYQFSMVAGSS
SGGDRSPTRMLASCTSSAASVAAGNLMNPSLGGQSDGVEADGSHSNSPTA
LSTPGRMDEAVWRPY

FIG. 2a

CCTTCTGCTTCCTAGCCCTGCTGTGGACAACTTAGGGTGCTCTTAGGTGG
GGGCCACTGGGGAGAACTGGCCTGTTTGTCCATCGATCTGATGGAAGAG
GGAGAAAAGACGACGGTCCATGCCAACTGGGGAAGGGCGAGGGTGTCTGC
ATGCCCCAGGTGGGGGAGTCGGAGTTCTCCCTCCCATCAAACAGACGACA
ATFTTTGTCGGTCCGGGATGGGGAGGAAGCAGGTGGAAATTGGGAACAAG
CTAGCCTGTTCTGTGGGTGTTCCCGGGTGCCTGAAAACGCGAAGACAGA
AGGCGTCCATCTTATCAGAGGTGAGGAAGGCGGCCTTGGTGTGACACAAA
GCCATCGGTTTGTCTGAGACCTGGCCGCATGGATGGCAAGGAAAGGGCAG
CTCCTCGGGGGCGGTGCGCGCGCCTGCACCCGGGGCCGCAGCTGGCGCGC
ATCTGTAGCCCGGCCGGGCCCGCACCTCCGCGGCTGGCAGGGCGCGGGCG
CCAAGTAGCGGGCGGCTCCCGCCACTGTGCCTGCCAGGGCGGCCCCCGCACC
TTGGTCTGGGGACCCAAGGGCCTGCACCCGCCCCCTCCCCCGCCGCGC
CTGGTCCCTTGATCTGGTGCCACGGGGAGCCAGCGCCCTACGGAGAGC
CGGAGCCGTGCCGGGCCCTGCGCCAGGTGCCAGGGCCACAAGAGTCCCT
CATTCTCTGGAACTTGTCTGTGAACCCATCGTAAGCGGAGAGGGAGAAA
TCAGGCGGAGGAACAAGCAAAGGGCACGGTGCAAACCGAAACCATTTCGAC
AAATGGAATTTACCACCACCTGAAACAGCGGGTCATGGTCCCGCAACCTG
CTCGAGGGCAGCACGTGTTGCCCGCCCCGGCCAGGGCCCTACCTGGCCA
CGACGCGCTGCGCCTTCTCGGAGACGTTCCCGGAGGTGGGAGCGCCAGG
CTGGATCACTCGCTTTCCTCTAGTTCTGCTGCTCGTGCCAGCGCGTCCGA
GGGCGCGCGGGCCTGGGTCCGCAATCGACAACCTGCCAGGCGCAGGCTCTC
TTAAAAGGTTTCAAGTAAGGGACCTTTGCCGTCTTCCTTTCGACACGGCCT
GAGGGCGTGCTGTGAGGTCCCGAGGTGGGTAGGGGCCAGCTCTCCCGGTG
GTCGGGGTGAGTCCAGAGTCCCTTCGCCCCGGAGCGCACGCGGGGCTTGA
TTTCGTTTGGCAACGACGAAATGGCGCGCGCTGAGCAGGGGTGAGATCCA
TGATGAGATCTTGCGGCCACCGTCCGATCCTAAGCTTCTTTGCTTCCGAG
GCTTGGAAATTCATTGTTTTGCACCTGTCGAGAGCCGGAGGCAACGAAAT
CTAGCCCCGTCTCAAAGCGGCGGGGAGGCTCAGCACGCGTTCGTTCCCC
AGAGTCTAGGGAGGTGTCTGGGGCGATAATTTCGGAATGATTGTGGCTTGA
TCTTTCCCGTTGCCCTCCCAACTGTAGCCGGCCCTAGGTCTGCTCGACA
GACTTAGGAGGCGGGAGAGGAAGGGGTGATTTGCAGTGAAGCCCAGGAGA
GGTTGGGCCACGCGGCTGGGAGTGGGAGCGGGGACCCGGAGCCGGGCGGG
CAGGCAGTGCCTTGGCGAAGCTGTCCGCGGTCCCTGCGGCGCAGCCGGAG
CGCACGGGCCCAAGAAGAAGTGGGGTTGGACCCGCAGAGGCCACTTCCA
CCCGCATGGAGAAAGAAAATTCTCTCTCTGAAAGCGAGGGCCCTTAGCT
TTGCAGCCACTGCTGTTTTTCTTTTCCACCGACGCGCGTACCGTTTCAC
GATGCAGGACCGTGGTTACATGCGTAAAGGAAAAAAGAAAAACGCATTT
TGCAGGCCTCGTCGTGTTTTTCAAAGAGCCACAGGCCGCTACAACGAAGA

FIG. 2b

ACGACGCCGCGAGGCCTGCAAGATCCTGAACTTGTTTTGAGGGGAGAGC
AGAGAGGAAAGGGGTTGTTGGCCCAGGCTACTTAGGGTCCCTAGGAGAC
TCCCTCCGCTGTCCCCGGTTTGGCACAGGGGCCACCGAGGCTGGGACC
AAAGCCGCGCAGGGCTGGGAGCAGCAAAGGCCGCCGGCCGGGCGTGGACG
ACGCGCAAATCCCCTGTGGGGTGGAGGCTCTTGGGTCAGAATAATGTGC
GGGACGAGGGAGGTGAGTAACCTCTTGGGGCGGCTCCAGTGCGGCGTC
ACCGGCCCTGAGACCCCGCGGCCCCAGCCCGGGGTTGCAGAAGTCACAG
GCCCGAAGCAGCAAGAGCTGGGGAAGCCCGGCCGCGGCCAGCGGGGAGGA
GGAGCGAAGGGGTTGCGCCCCAGCGTCAGGGAGCTACGACCCGAGAGAGG
GCGGCAAGGGCGCCTTCCGTGGGACCCGGACGTTCTAAGCAAATTTCTAG
CATTTGCCCCGGGCTCCAGAGCTCTCGGGGGCCCTGGGCTGTGGCACTG
GGGCCTCCTCCGCGGGGTGGCGCCTTCCGCCCTCCCCGTTGGGCGGCCT
CCGGCAGGCCCCGTTCCCTCCCCGCGAACGCCACCGAGGTGCCCGCGATGG
GGGCTCCGCCGATTGGCTGTGCGACGCGTCGCTCCGCCAGCCCCGCCCCG
CGGGCCCCGGGGTACTAACCCCGCGCGGGCGGCCGCGGCCCCGCCACTT
GATTCTGGAGGATTTGTTCTGGGGCTGCGGCCGCGGAGTCGGGGCGGCCG
CGGGCGAGCTTCGGGGCGGGAGGCGGCGGCAGCGGCACAGCCCCGCGCGG
GCCCCGCCGCGGCCAGGCAGCCGGGACAGCCACGAGGGGCGGCCGCACG
CGGGGCCGCGCGCCGAGGATGCGGGACTAGCCGGGCAGGCTGCGGGCGGC
CGTCGGGCCAGCGAGGCCTCGCAGCGGGCGGGCCCTGGCGAGTAGTGGCC
GGGCGCCGCCCTGCGCCCTGAGGCCCGGGCCCCGCCGCTTCTGCTTTC
CCGCTTCTCGCGGCAGCGGCGGCCGAGGAGGCGCCCGCGCCGGCCGCCCC
CGGGGGAAGCCGCGCCGTCTCCGCCTGCCCGGCGCCCTGACGGCCGCTGT
TATGCGTATTCCTGATAGACCCAAGCACCAGCCGCCGCTTACACCTCCCT
CCCCGGCCTTCCCTGCGGCGGCGGGCGGCGGCAAGATGGGCGAGAACAGC
GCGGCTCTGAGCGCCAGGCGGCCGTTGGGGCCCGGAGGGCGCGCCCGGCC
CGAGGTGCGCTCGATGGTGGACGTGCTGGCGGACCACGCAGGCGAGCTCG
TGCGCACCGACAGCCCCAACTTCTCTGCTCCGTGCTGCCCTCGCACTGG

FIG. 3




FIG. 4a




FIG. 4b




FIG. 5a




FIG. 5b




FIG. 5c




FIG. 6a




FIG. 6b

RT-PCR




FIG. 6c




FIG. 7a

Genomic Southern






FIG. 7b



○: SmaI ●: HaeII ▲: BamHI □: MluI ◆: NheI

FIG. 8



RUNX3 GENE SHOWING ANTI-TUMOR ACTIVITY AND USE THEREOF

This patent application claims a benefit of priority from Korean Patent Application No. 2001-4018 filed Jan. 29, 2001 through PCT Application Ser. No. PCT/KR01/00121 filed Jan. 30, 2001, the contents of each of which are incorporated herein by reference.

FIELD OF THE INVENTION

The present invention relates to the RUNX3 gene and uses thereof as an anticancer agent. More specifically, the present invention relates to characterization of the RUNX3 gene, its expression and tumor suppressor activity, and its use in the development of cancer diagnosis methods and anticancer agents.

BACKGROUND OF THE INVENTION

Tumors are assemblies of cells that show excessive autonomous proliferation, and are classified into malignant tumors, which may result in death, and benign tumors; however, it can be difficult to discriminate between these types. Cancers are genetic diseases attributed to the mutation of genes such as oncogenes and tumor suppressor genes, and their causes can be found at a cellular level. Oncogene products constitute a signaling network within cells, and are involved in regulating signal transduction systems for cell division and differentiation. When their regulation becomes abnormal, cells do not differentiate further, but rather divide infinitely; that is, they become tumorigenic. If a way could be found to enable such an abnormal signal transduction pathway to be converted into a normal one, anticancer agents could be developed that would obtain excellent therapeutic effects without side effects.

At present, cancers are for the most part treated in three ways: with surgical therapy, chemical therapy, and radiation therapy. In practice, combinations of these therapies or further combination with laser therapy are prevalent. However, chemical therapy is preferred to other therapies when the pain accompanying therapy and metastatic conditions are taken into account. Numerous anticancer agents have been developed, most of which are based on the selective killing of cells that are actively dividing. However, these anticancer agents suffer from the disadvantage of also killing some normal cells, such as immune cells and hair root cells, with concomitant significant side effects; they are thus not able to be used for long periods of time. Therefore, there remains a need for novel anticancer agents. A therapeutic agent based on the causes of cancer might be expected to be highly effective and to have few or no accompanying side effects.

Abnormal activation of oncogenes induces cell proliferation, and is one cause of cancer. In contrast, tumor suppressor genes function to prevent abnormal cell proliferation or to trigger programmed cell death (apoptosis). Often, tumor suppressor genes trigger apoptosis to kill the cells with abnormally activated oncogenes thus preventing the formation of cancerous cells. Where tumor suppressor genes show normal activity, cells with abnormally activated oncogenes cannot progress toward cancer, but are annihilated. Therefore, to become cancerous cells, cells must have inactivated tumor suppressor genes as well as activated oncogenes.

One of the mechanisms by which tumor suppressor genes are inactivated is by hyper-methylation of CpG islands (Jones and Laird, *Nature Genet.* Vol. 21, 163-167, 1999).

Methylation of CpG islands is performed by DNA methyltransferase. After significant methylation, DNA binding proteins such as methyl cytosine binding protein 2 (MECP2) bind to the methylated cytosine of the DNA, which recruits histone deacetylase (HDAC) to repress gene expression. In detail, HDAC removes the acetyl groups associated with histones, and the chromosomal DNA in the vicinity of the deacetylated histones becomes dense, which leads to repression of gene transcription. If gene expression is repressed by DNA methylation, DNA methyltransferase or HDAC inhibitors may be useful for inducing gene expression. Tumor suppressor genes whose expression is repressed by DNA methylation are exemplified by RB1, TP53, VHL, CDKN2A, CDKN2B, MLH1, and APC (Jones and Laird, *Nature Genet.* Vol. 21, 163-167, 1999).

As mentioned previously, histone acetylation and deacetylation are known to play important roles in regulating DNA transcription in eukaryotic cells (Grunstein M., *Nature*, 389, 349-352, 1997). Some naturally occurring compounds have been found to prevent cells from progressing toward cancer by inhibiting HDAC. Exemplified by trapoxin, trichostatin A, and depudecin, HDAC inhibitors have been studied for their ability to reverse the transformation of cancerous cells. Of the HDAC inhibitors, depudecin is the best characterized as to its anti-angiogenic activity *in vivo* and *in vitro*. In addition, the HDAC inhibitors have been studied with regard to cellular responses, including cell cycle interruption, alteration of gene expression patterns, and induction of apoptosis.

The TGF- β signal transduction system is well known for its tumor suppressor activity. The binding of TGF- β to TGF- β receptors causes the activation of the receptors, which in turn activate Smad proteins by phosphorylation. Once activated, Smad proteins move into the nucleus and regulate gene expression in cooperation with other transcription factors, thereby suppressing cell division or inducing apoptosis (Massague et al., *Cell*, 103 (2):295-309, 2000). Runx3 is one of the transcription factors that physically interact with Smad proteins (Hanai et al., *J. Biol. Chem.* 274: 31577-31582, 1999). Deletion or mutation of TGF- β receptors or Smad genes is observed in cells of various types of cancer. The tumor suppressor activity of TGF- β receptors was also demonstrated by an experiment in a cell strain lacking the TGF- β receptor. When the cell was transfected to express the TGF- β receptor, cell proliferation was reduced and tumorigenesis was decreased in an assay in nude mice (Chang et al., *Cancer Res.*, 57 (14):2856-2859, 1997). The TGF- β signal transduction system is well characterized as to the repression of cell proliferation, which is achieved by promoting the expression of the CDK inhibitor protein p21. However, the mechanism by which TGF- β induces apoptosis remains to be clearly elucidated.

PEBP2 (polyoma virus enhancer binding protein 2) is composed of two subunits, α and β . There are three genes which encode the α subunit: RUNX1/PEBP2 α B/CBFA2/AML1, RUNX2/PEBP2 α A/CBFA2/AML2, and RUNX3/PEBP2 α C/CBFA3/AML2 (Bae and Ito, *Histol. Histo-pathol.* 14(4):1213-1221, 1999). The RUNX1, RUNX2, and RUNX3 genes show homology of about 60-70% in amino acid sequence among them. They are highly conserved evolutionarily, with homology of about 95% between mouse and human.

Regarded as an important causative gene in leukemia, the RUNX1 gene becomes associated with other genes by chromosome translocation to cause acute myeloid leukemia or acute lymphoid leukemia in humans (Miyoshi et al., *EMBO J.*, 12:2715-2721, 1993; Romana et al., *Blood*,

85:3662-3670, 1995; Okuda et al., Blood, 91:3134-3143, 1998; Okuda et al., Cell, 84:321-330, 1996). The RUNX2 gene plays a crucial role in osteogenesis, and its disruption has been implicated in causing cleidocranial dysplasia (Komori et al., Cell, 89:755-764, 1997; Lee et al., Nat. Genet., 15:307-310, 1997; Mundlos et al., Cell, 89:773-779, 1997; Otto et al., Cell, 89:756-771, 1997). Also, it has been reported that the RUNX2 gene shows oncogenic activity in the formation of T-cell lymphoma (Stewart et al., Proc. Natl. Acad. Sci. U.S.A., 94(16):8646-8651, 1997).

The RUNX3 gene was identified by the present inventors several years ago as a member of the PEBP2 family (Bae et al., Gene, 159(2):245-248, 1995; Levanon et al., Genomics, 23(2):425-532, 1994). However, diseases associated with the activation or inactivation of the RUNX3 gene, other than what we describe in this invention, hasn't been reported.

SUMMARY OF THE INVENTION

Leading to the present invention, the thorough and intensive research on diseases associated with the activation of the RUNX3 gene, conducted by the present inventors, resulted in the finding that RUNX3 gene products have tumor suppressor activity. RUNX3 is indispensable for TGF- β -induced apoptosis, and inactivation of the RUNX3 gene by DNA methylation at loci near RUNX3 exon 1 is closely correlated with cancer development. In the present invention, the RUNX3 gene is characterized as to its expression and tumor suppressor activity, offering the possibility of developing cancer diagnosis methods and anticancer agents from it.

The objectives of the present invention are to elucidate the tumor suppressor activity of the RUNX3 gene and its mechanism, and to provide a diagnostic method and a therapeutic agent for cancer based on the tumor suppressor activity of RUNX3.

To achieve the above objectives, the present invention provides a cell strain having a sense RUNX3 cDNA, in which the RUNX3 cDNA is over-expressed, and a cell strain having an antisense RUNX3 cDNA, in which the expression of the RUNX3 gene is inhibited are provided.

Also, the present invention provides a RUNX3 cDNA with tumor suppressor activity and its corresponding protein.

In the present invention, the RUNX3 gene is shown to play a crucial role in TGF- β -dependent apoptosis, using cell strains that either overexpress RUNX3 cDNA or selectively inhibit the expression of the RUNX3 gene.

Additionally, it is shown that the suppression of RUNX3 gene expression in various cell lines can be attributed to DNA methylation in the vicinity of exon 1 of the RUNX3 gene.

Further, the present invention provides a pharmaceutical composition including the RUNX3 gene or its protein.

Finally, the present invention provides uses of the RUNX3 gene and its proteins in the development of anti-cancer agents and diagnostic methods.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1a is a cDNA nucleotide sequence of the human RUNX3 gene with arrows denoting the primers used for RT-PCR analysis and bold letters denoting the translation initiation codon (ATG) and translation stop codon (TGA).

FIG. 1b continues the cDNA nucleotide sequence of the human RUNX3 gene from FIG. 1a.

FIG. 1c is an amino acid sequence deduced from the sequence of the cDNA of the human RUNX3 gene shown in FIG. 1a.

FIG. 2a is a nucleotide sequence showing CpG islands present in exon 1 of the human RUNX3 gene with a site used as a probe in genomic DNA Southern blot analysis denoted by underlining and a translation initiation codon (ATG) denoted by bold letters.

FIG. 2b continues the nucleotide sequence of the CpG islands present in exon 1 of the human RUNX3 gene from FIG. 2a.

FIG. 3 is an autoradiogram resulting from Northern blotting analysis for determining whether SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12, both having an antisense RUNX3 gene, and MKN28-Rx3, having a sense RUNX3 gene, express the antisense and sense RUNX3 genes.

FIG. 4a is a graph in which counts of viable SNU16-Rx3-AS-c11 cells are plotted with regard to time, along with those of the control SNU16 cells, after treatment with 1 ng/ml of TGF- β to analyze TGF- β -induced cell death.

FIG. 4b is a photograph showing the results of the electrophoresis of the genomic DNAs obtained from SNU16, SNU16-Rx3-AS-c11, and SNU16-Rx3-AS-c12 after treatment with 1 ng/ml of TGF- β for 6 hours so as to determine whether RUNX3 is involved in the TGF- β -induced programmed cell death pathway.

FIG. 5a is a graph showing that the size of the tumors formed in nude mice were measured periodically for over 98 days after the subcutaneous injection of SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12, both having restricted RUNX3 expression due to the over-expression of the antisense RUNX3 gene therein; and control-SNU16 which expresses the RUNX3 gene normally.

FIG. 5b is a graph in which tumors formed in nude mice were measured for size periodically for over 37 days after the subcutaneous injection of control MKN28, which does not express the RUNX3 gene, and MKN28-Rx3, in which a RUNX3 cDNA is expressed.

FIG. 5c shows a comparison of cancerous masses excised from the mice sacrificed on the final day of the experiment.

FIG. 6a is a schematic view showing the structure of the RUNX3 cDNA with arrows denoting the RT-PCR primer positions, and a striped box denoting the Runt domain, which plays an important role in binding of the RUNX3 protein to DNA.

FIG. 6b shows the results of electrophoresis of RT-PCR products obtained from total RNAs of 15 gastric cancer cell lines by use of pairs of PS-N primers (PS-N), PS-C primers (PS-C), human PEBP2 β /CBFB cDNA primers (PEBP2 β), and human β -actin cDNA primers (β -actin), and the results of Southern blotting (PS-C ST). For the Southern blot analysis, RT-PCR products obtained by use of PS-C primers were hybridized with RUNX3 cDNA as a probe.

FIG. 6c shows the results of electrophoresis of RT-PCR products obtained from total RNAs of 17 lung cancer cell lines by use of pairs of PS-N primers (PS-N), PS-C primers (PS-C), and human β -actin cDNA primers (β -actin), to analyze the production of RUNX3 mRNA.

FIG. 7a is a schematic illustration showing restriction enzyme sites in the vicinity of RUNX3 exon 1 and the CpG frequency within typical CpG islands, with tick bars denoting the position of a DNA fragment used as a probe in genomic Southern blot analysis.

FIG. 7a is an autoradiogram resulting from Southern blotting analysis for DNA methylation in the vicinity of RUNX3 exon 1 in various gastric cancer cell lines by use of DNA probes which can detect parts of the CpG islands. The

genomic DNAs were isolated from the cancer cell lines and treated with *Sma*I (upper), which can not digest methylated DNA, and *Bam*HI (lower), which can digest DNA irrespective of DNA methylation.

FIG. 8 shows the results of electrophoresis of RT-PCR products obtained from total RNAs of NUGC3, MKN28, and MKN74 after treatment with 5-aza-2-deoxycytidine (AZA, 300 nM) or trichostatin A (TSA, 1 mM), separately or in combination. The results indicate that the treatment with the DNA methyltransferase inhibitors or HDAC inhibitors allows the RUNX3 gene to be expressed.

DETAILED DESCRIPTION OF THE INVENTION

In the present invention, the role of RUNX3 in cells is elucidated. For this purpose, a sense RUNX3 cDNA or an antisense RUNX3 cDNA is introduced into cells to establish strains over-expressing sense or antisense RUNX3 RNA, respectively.

Therefore, in one aspect of the present invention, cell lines are provided that harbor a sense RUNX3 cDNA or an antisense RUNX3 cDNA to excessively express the RUNX3 protein and to selectively inhibit the expression of the RUNX3 gene, respectively.

The human RUNX3 gene cloned by the present inventors in 1995 (Bae et al., *Gene*, 159 (2):245-248, 1995; cDNA nucleotide sequence GenBank Accession No. Z35278) was used to construct a plasmid vector capable of RUNX3 expression. The cDNA of the RUNX3 gene has the nucleotide sequence SEQ. ID. NO: 1. This cDNA has an open reading frame ranging from the translation start codon to the translation stop codon, and encodes a polypeptide consisting of 415 amino acids whose putative amino acid sequence is SEQ. ID. NO: 2 (see FIGS. 1a to 1c). A DNA fragment comprising CpG islands in the vicinity of exon 1 of the RUNX3 gene has the nucleotide sequence SEQ. ID. NO: 3 (see FIGS. 2a and 2b).

A 2,244 bp DNA fragment comprising a coding region of the RUNX3 gene cDNA SEQ. ID. NO: 1 (see FIG. 1a) was inserted in the sense direction into a pEF-BOS vector (Mizushima et al., *Nucleic Acids Res.*, 18(17):5322, 1990) to construct a recombinant plasmid vector, named pEF-BOS-Rx3, which expresses a sense RUNX3 gene. Separately, the RUNX3 cDNA was inserted in the antisense orientation into a pEF-BOS vector to construct a recombinant vector, named pEF-BOS-Rx3-AS.

After the plasmid vector pEF-BOS-Rx3 was co-transfected with another vector bearing a neo resistance selection marker into various cancer cell strains, stable transfectants over-expressing the sense RUNX3 RNA were selected. In a preferred embodiment of the present invention, the pEF-BOS-Rx3 plasmid and a plasmid with a neo selection marker were co-transfected into the MKN28 strain (which does not express a RUNX3 gene) using Lipofectamine (Gibco-BRL), and the selection of transfectants was achieved by means of G418 resistance. The MKN28 strain that shows over-expression of the sense RUNX3 gene was named MKN28-Rx3 and deposited with the Korean Collection for Type Culture of Korea Research Institute of Biotechnology and Biotechnology (KRIBB) under the deposition No. KCTC 0933BP on Jan. 9, 2001.

The strains that express the antisense RUNX3 RNA were established by co-transfecting the plasmid vector pEF-BOS-Rx3-AS and a plasmid with a selection marker into cells of the SNU16 gastric cancer cell line (Kim et al., *J. Natl. Cancer Inst.*, 8(13):938-943, 1991) which has an inactivated

p53 gene and a normal TGF- β gene system (Park et al., *Proc. Natl. Acad. Sci. U.S.A.*, 91 (10):8772-8776, 1994). Stable transfectants were obtained in the same manner as above. Two transfectants which expressed the antisense RUNX3 RNA in excess were named SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12, and deposited with the Korean Collection for Type Culture of KRIBB under the deposition Nos. KCTC 0934BP and KCTC 0935BP on Jan. 9, 2001.

To determine whether the transfectants MKN28-Rx3, SNU16-Rx3-AS-c11, and SNU16-Rx3-AS-c12 actually express the sense RUNX3 gene or the antisense RUNX3 gene, the RUNX3 cDNA of SEQ. ID. NO: 1 was used as a DNA probe for Northern blotting analysis. A RUNX3 transcript from the MKN28-Rx3 strain was detected as a single band at a position almost identical to that of 18S RNA, verifying the expression of RUNX3 mRNA in the cell. In contrast, antisense RUNX3 RNAs of various sizes were detected from SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 cells. In the control SNU16 strain, the expression of the endogenous RUNX3 gene was hardly detected by Northern blotting analysis (see FIG. 3).

In another aspect of the present invention, a RUNX3 gene and its product, which show tumor suppressor activity, are provided.

In the present invention, the function of the RUNX3 gene is elucidated by using the MKN28-Rx3 strain, which expresses the RUNX3 gene, and the SNU16-Rx-AS-11c and the SNU16-Rx-AS-12c strains, whose RUNX3 gene expression is selectively inhibited.

To analyze the role of RUNX3 in the TGF- β -induced programmed cell death (apoptosis), the SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 strains in which RUNX3 expression is selectively inactivated were treated with TGF- β 1; normal SNU16 cells were used as a control. Counts of viable cells revealed that TGF- β 1 induces apoptosis in the control strain, but not in the mutant strains. In detail, treatment with TGF- β 1 killed all SNU16 cells within 2 days. However, treatment with TGF- β 1 did not kill the SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 cells (see FIG. 4a). Accordingly, these results show that the RUNX3 gene is involved in the TGF- β 1-induced cell death in SNU16 cells.

In the present invention, inventors also determine whether the mechanism of the RUNX3-dependent TGF- β 1-induced cell death in SNU16 cells treated with TGF- β 1 is via the apoptosis pathway or a necrosis pathway. Electrophoresis of DNA from cells killed through apoptosis yields specific apoptotic DNA bands. These bands are apparent in the DNA from SNU16 cells treated with TGF- β 1 (see FIG. 4b Lanes 1 and 2). However, no apoptotic bands were observed in the DNA of the SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 cells (see FIG. 4b, Lanes 3 to 6). Therefore, TGF- β 1 stimuli kill SNU16 cells through an apoptotic pathway, and RUNX3 is indispensable for this process.

It is known that TGF- β 1-induced apoptosis is a very important mechanism for suppressing the progression of normal cells toward cancer, and that most of the factors involved in the TGF- β pathway have tumor suppressor activity (Chang et al., *Cancer Res.*, 57(14):2856-2859, 1997). To examine the anticancer activity of RUNX3, a tumorigenesis assay was performed using nude mice. For the tumor suppressor activity assay, the cell lines are introduced into nude mice by subcutaneous injection, after which the cancerous masses formed are periodically measured for their external sizes. After the mice are sacrificed on the final day, the cancerous masses are excised for comparison.

The tumors formed in the nude mice grew at a higher rate in the mice injected with the cells expressing SNU16-Rx3-

AS-c11 or the SNU16-Rx3-AS-c12 than those in the mice injected with the control SNU16 cells (see FIG. 5a). In detail, the cancerous masses were measured on the final day. Those of the experimental group into which SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 had been injected were 1,600 mm³ in volume, while those of the control group were about 800 mm³. The cancerous masses were excised from the mice 98 days after the injection and compared to one another. The cancerous masses from the experimental group were remarkably larger than those from the control group (see FIG. 5c).

The tumorigenicity of control MKN28 cells and MKN28-Rx3 cells in which the sense RUNX3 RNA is expressed was also assayed, using nude mice in the manner described above for the SNU16 cells. Significantly reduced tumorigenesis was observed in the mice injected with MKN28-Rx3 cells compared to that of the mice injected with control MKN28 cells (see FIG. 5b). In detail, the cancerous masses of the control group were 600 mm³ in volume on the final day, in contrast to those of the cancerous masses of the MKN28-Rx3-injected experimental group, which were found to be as small as 250 mm³. This represents a reduction of tumor growth of 58%. The cancerous masses were excised from the mice 30 days after the injection and compared to one another (FIG. 5c).

These results demonstrate that the RUNX3 gene has tumor suppressor activity.

In the present invention, the RUNX3 gene was analyzed for its expression pattern in various cancer cell lines. Total RNAs were isolated from 15 gastric cancer cell lines and 17 lung gastric cancer cell lines and used to determine whether these cell lines produce RUNX3 mRNAs, by use of RT-PCR. The gastric cancer cell lines used in this RT-PCR were SNU1, SNU5, SNU719, NUGC3, MKN1, MKN7, MKN28, MKN45, MKN74, AGS, KatoIII, RF1, RF48, and AZ521. The lung cancer cell lines were NCI-H522, 86-2, A549, LK79, LK87, LCSC#1, LCSC#2, NCI-H23, NCI-H226, NCI-460, NCI-H322, Sq-19, NCI-H1915, NCI-H630, Hs 888Lu, Lu65, and LX-1. These cell lines were obtained from the Korean Research Institute of Bioscience and Biotechnology (KRIBB).

RT-PCR was conducted using two primer pairs, named PS-N and PS-C (FIG. 1a). Among the gastric cancer cell lines, SNU5, SNU16, SNU719, MKN1, MKN45, RF1, RF48, MKN7, and AZ521 were found to produce RT-PCR products with expected sizes (1,080 bp with the PS-N primer pair and 870 bp with the PS-C primer pair), although MKN7 produced RT-PCR products in very small quantities (FIG. 6b). In contrast, no RT-PCR products were obtained from the SNU1, NUGC3, MKN28, MKN74, AGS, and KatoIII cell lines when using either the PS-N primer pair or the PS-C primer pair.

In the case of the lung cancer cell lines, the expression of the RUNX3 gene was found to be suppressed significantly in the HS888Lu cell line and completely in NCI-H226, NCI-460, NCI-H630, and Lu65 cell lines (see FIG. 6c).

Southern blotting analysis was used to verify that the RT-PCR products were amplified from the cDNA of the RUNX3 gene (see FIG. 6b, PS-3 ST). Additionally, the amplification of PEBP2 β (see FIG. 6b PEBP2 β) and β -actin cDNAs (see FIG. 6b, β -actin) from identical first cDNA strands was observed, indicating that the lack of amplification of RUNX3 cDNAs in the gastric cancer cell lines SNU1, NUGC3, MKN28, MKN74, AGS, KatoIII, NCI-H226, NCI-460, NCI-H630, and Lu65 can be attributed to a lack of expression of the RUNX3 gene in those cell lines.

Likewise, the low levels of the RT-PCR products in the MKN7 and Hs888Lu cell lines result from the low expression levels of the RUNX3 gene in the cell lines.

The above results show that the RUNX3 gene is remarkably or completely suppressed in about 47% of the gastric cancer cell lines and about 30% of the lung cancer cell lines. Therefore, the suppression of the RUNX3 gene expression can be an important diagnostic index in about 47% of gastric cancer cells and about 30% of lung cancer cells (in about 37% of the total number of gastric and lung cancer cell lines we examined).

Mutation analysis of the RUNX3 gene in cancer cell lines was performed using all exons of the RUNX3 gene, amplified from various cancer cell lines. No point mutation affecting the amino acid sequence was found. These results led to the conclusion that cancer cell lines in which the expression of the RUNX3 gene was not detected contain a normal RUNX3 gene, but do not express it. To determine whether the suppression of the RUNX3 gene expression can be attributed to DNA methylation, the correlation between the expression of the RUNX3 gene and the hypermethylation of CpG islands at loci near RUNX3 exon 1 was analyzed through genomic DNA Southern blotting using restriction enzymes sensitive to DNA methylation. Genomic DNAs isolated from cancer cell lines were separately treated with restriction enzymes which are unable to cut methylated DNA and enzymes that are able to cut DNA irrespective of methylation.

In one preferred embodiment of the present invention, SmaI, which cannot digest methylated DNA, and BamHI, which can digest DNA irrespective of its methylation, were used. The DNA was specifically protected from SmaI digestion in the gastric cancer cell lines whose RUNX3 gene expression was not detected, that is, the SNU1, NUGC3, MKN28, MKN74, AGS, and KatoIII cell lines. The MKN7 cell line in which the expression level was very low showed a partially methylated pattern (see FIG. 7a). These results suggest that the expression of the RUNX3 gene is closely correlated with DNA methylation at loci near RUNX3 exon 1 in cancer cell lines.

To examine further whether DNA methylation causes RUNX3 gene silencing, the influence of a DNA methyltransferase inhibitor and a histone deacetylase inhibitor on the reactivation of RUNX3 gene expression was analyzed. 5-aza-3-deoxycytidine (AZA, 300 nM) and trichostatin A (TSA, 1 mM) were used as DNA methylase and histone deacetylase inhibitors, respectively. Upon treatment with either the DNA methyltransferase inhibitor or the histone deacetylase inhibitor, the expression of the RUNX3 gene was induced (see FIG. 8). These results demonstrate that the hyper-methylation of CpG islands at loci near exon 1 of the RUNX3 gene is responsible for the suppression of RUNX3 gene expression in about 37% of the cancer cells. This result also indicates that agents capable of promoting the reactivation of the RUNX3 gene can be developed from DNA methyltransferase inhibitors or histone deacetylase inhibitors and used as anticancer agents.

According to the present invention, therefore, the product of the RUNX3 gene is involved in TGF- β 1-dependent apoptosis and shows tumor suppressor activity. The expression of the RUNX3 gene is suppressed in a substantial portion of the cancer cells by the hyper-methylation of CpG islands at loci near exon 1 of the gene.

In a further aspect of the present invention, expression vectors are provided that are capable of expressing RUNX3

cDNA, and pharmaceutical compositions are provided that comprise proteins of the genes as pharmaceutically effective ingredients.

The pharmaceutical compositions of the present invention can be used for the diagnosis, prophylaxis, and treatment of the diseases attributed to abnormal TGF- β -dependent apoptosis, including gastric and lung cancers.

As the pharmaceutically effective ingredients, RUNX3 cDNA, RUNX3 RNA, or proteins produced therefrom may be used. Expression vectors capable of expressing the RUNX3 gene are preferable because they can produce RUNX3 proteins continuously if they are injected. Where a pharmaceutical composition comprising such an expression vector is injected into a bio-tissue via a parenteral route, the bio-tissue can produce RUNX3. That is, the RUNX3 gene of the present invention or an expression vector anchoring the RUNX3 gene can be used as a pharmaceutically effective ingredient of a pharmaceutical composition for use in gene therapy.

Administration of the RUNX3 cDNA of the present invention and/or the RUNX3 RNA or the RUNX3 protein produced therefrom may take an oral or a parenteral route, with preference for parenteral injection.

The expression vector comprising the RUNX3 gene or the RUNX3 protein produced therefrom may be administered in various oral or parenteral dosage forms. For the formulation of the pharmaceutically useful ingredient, diluents, expedients, and/or carriers, such as fillers, thickeners, binders, wetting agents, disintegrating agents, surfactants, etc., may be used. Solid formulations for oral administration may be formed into pills, tablets, powders, granules, capsules, etc. These solid formulations may comprise at least one expedient, such as starch, calcium carbonate, sucrose, lactose, gelatin, etc. In addition to simple expedients, lubricants such as magnesium stearate and talc may be used. Liquid formulations for oral administration may be exemplified by suspensions, internal solutions, emulsions, syrups, etc. These liquid formulations may comprise various expedients, for example, wetting agents, sweeteners, odorants, and preservatives, as well as simple diluents such as water, liquid paraffin, etc. Examples of formulations for parenteral administration include sterile aqueous solutions, non-aqueous solutions, suspensions, emulsions, freeze-dried agents, and suppositories. For the formation of non-aqueous solutions or suspensions, vegetable oils such as polypropylene glycol, polyethylene glycol, and olive oil, and an injectable ester such as ethyloleate may be used. As bases of suppositories, witepsol, microgol, Tween 61, cacao oil, laurine, and glycerogelatin may be used. In order to help the absorption of the pharmaceutically useful ingredient into cells, liposomal dosage forms may be used.

The therapeutically effective dose of the pharmaceutically effective ingredients is within the range of 0.002 to 1 mg/kg of body weight and preferably within the range of 0.02 to 0.2 mg/kg of body weight for RUNX3 cDNA, within the range of 0.001 to 0.5 mg/kg of body weight and preferably within the range of 0.01 to 0.1 mg/kg of body weight for RUNX3 RNA, and within the range of 0.04 to 4 mg/kg and preferably within the range of 0.4 to 1 mg/kg of body weight for RUNX3 protein.

In accordance with a further aspect of the present invention, uses of the RUNX3 gene and its proteins in the development of anticancer agents and the diagnosis of cancers are provided.

As described above, the RUNX3 gene of the present invention shows excellent tumor suppressor activity and is indispensable for TGF- β -dependent apoptosis. The expres-

sion of RUNX3 is suppressed in about 47% of gastric cancer cell lines and about 30% of lung cancer cell lines (in about 37% of the total number of gastric and lung cancer cell lines). In addition, the suppression of RUNX3 gene expression is attributed to abnormal DNA methylation of CpG islands located at loci near exon 1 of the RUNX3 gene. Further, treatment with a DNA methylase inhibitor or a histone deacetylase inhibitor can induce the expression of the RUNX3 gene. Therefore, the RUNX3 gene of the present invention and proteins produced therefrom can be utilized in developing not only anticancer agents, but also diagnostic methods for detecting cancers, in which abnormal DNA methylation of CpG islands at the loci near exon 1 of the RUNX3 gene is measured.

Based on the facts presented in the present invention, pharmaceutical compositions comprising materials capable of regulating the methylation of a nucleotide sequence that includes CpG islands near RUNX3 exon 1 can be provided for the prophylaxis and treatment of cancers as pharmaceutically effective ingredients.

A method for diagnosing cancers characterized by detecting the methylation of the nucleotide sequence containing CpG islands present near exon 1 of the RUNX3 gene comprises the steps of:

- isolating genomic DNA from a tumor tissue of a subject;
- digesting the genomic DNA with a restriction enzyme sensitive to DNA methylation;
- separating the genomic DNA digests on an agarose gel by electrophoresis and transferring them onto a membrane;
- hybridizing the genomic DNA digests with a radiolabeled DNA probe for the RUNX3 gene; and
- exposing the membrane to a film to detect the expression of the RUNX3 gene.

In accordance with another embodiment of the present invention, a method for diagnosing cancers comprises the steps of:

- Isolating genomic DNA from blood or a tumor tissue of a subject;
- digesting the genomic DNA with a restriction enzyme sensitive to DNA methylation;
- performing a polymerase chain reaction (PCR) with the genomic DNA digests serving as templates and with parts of the nucleotide sequence of SEQ. ID. NO: 3 serving as primers;
- separating the PCR products on an agarose gel by electrophoresis to detect the amplification of a DNA sequence of interest.

The diagnostic method using PCR is based on the principle that PCR products are obtained if the DNA region of interest is methylated; otherwise, the products are not obtained.

In accordance with a still further aspect of the present invention, a biological microchip for use in cancer diagnosis is provided, which comprises a part of the RUNX3 cDNA SEQ. ID. NO: 1 or proteins produced therefrom.

Generally, a biological microchip for the diagnosis of diseases, which is based on principles of molecular biology and electronic engineering, has tens to tens of thousands of DNAs or proteins integrated on a small chip. Thanks to the high level of integration, the biological microchip can be used to search for at least hundreds of genes or proteins within a short period of time. Therefore, the biological chip can be used to perform a search for novel genes or to diagnose diseases very rapidly and simply, compared to conventional bioengineering techniques. Biological chips can be classified into oligonucleotide chips and cDNA chips according to the genetic materials integrated thereon, and

they can be fabricated according to use, for example, genes to be searched, diseases to be diagnosed, etc. Microchips using parts of the RUNX3 cDNA nucleotide sequence or antibodies against proteins produced from the RUNX3 cDNA fragments can be used for the diagnosis of various cancers, including gastric and lung cancers.

EXAMPLES

A better understanding of the present invention may be obtained in light of the following examples which are set forth to illustrate, but are not to be construed to limit, the present invention.

Example 1

Establishment of the SNU16-Rx3-AS and MKN28-Rx3 Cell Strains

To elucidate the function of the RUNX3 gene in vivo, cell strains capable of over-expressing a sense RUNX3 gene and an antisense RUNX3 gene were established as follows.

1-1 Construction of Expression Plasmid Vectors Containing Sense and Antisense RUNX3 Genes

The human RUNX3 gene cloned by the present inventors in 1995 (Bae et al., *Gene*, 159 (2):245-248, 1995; cDNA nucleotide sequence GenBank Accession No. Z35278) was used to construct plasmid vectors capable of expressing RUNX3. The cDNA of the RUNX3 gene has the nucleotide sequence SEQ. ID. NO: 1 with an open reading frame ranging from the translation start codon to the translation stop codon, which encodes a polypeptide consisting of 415 amino acids whose putative amino acid sequence is SEQ. ID. NO: 2 (FIGS. 1a to 1c). A DNA fragment comprising CpG islands present in the vicinity of exon 1 of the RUNX3 gene has the nucleotide sequence SEQ. ID. NO: 3 (FIGS. 2a and 2b).

A 2,244 bp BamHI DNA fragment (FIG. 1a) comprising a coding region out of the cDNA of the RUNX3 gene SEQ. ID. NO: 1 was inserted in the sense direction into a pEF-BOS vector (Mizushima et al., *Nucleic Acids Res.*, 18(17): 5322, 1990) at an XbaI site to construct a recombinant plasmid vector, named pEF-BOS-Rx3, which expresses a sense RUNX3 gene.

Separately, the RUNX3 cDNA was inserted in the antisense direction into a pEF-BOS vector at an XbaI site to construct a recombinant vector, named pEF-BOS-Rx3-AS.

1-2 Stable Transfection

The plasmid vector pEF-BOS-Rx3, established in Example 1-1, and pcDNA3.1, which contains a neo selection marker, were co-transfected into the MKN28 strain by the Lipofectamine (Gibco BRL) method as specified by the manufacturer. MKN28 does not endogenously express RUNX3. Stable transfectants that harbored pEF-BOS-Rx3 (sense RUNX3 RNA) were selected in the presence of G418. The MKN28 strain overexpressing sense RUNX3 RNA was named MKN28-Rx3 and deposited with the Korean Collection for Type Culture of Korea Research Institute of Bioscience and Biotechnology (KRIBB) under the deposition No. KCTC 0933BP on Jan. 9, 2001.

The strains capable of expressing the antisense RUNX3 gene were established by co-transfecting the plasmid vector pEF-BOS-Rx3-AS along with pcDNA3.1, which contains a selection marker, into the SNU16 strain, a gastric cancer cell line (Kim et al., *J. Natl. Cancer Inst.*, 8(13):938-943, 1991) that has an inactivated p53 gene and a normal TGF- β

signaling system (Park et al., *Proc. Natl. Acad. Sci. U.S.A.*, 91 (10):8772-8776, 1994). Stable transfectants were obtained in the same manner as above. Two transfectants that excessively express the antisense RUNX3 gene to selectively inhibit the expression of the RUNX3 gene were named SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12, and deposited with the Korean Collection for Type Culture of KRIBB under the deposition Nos. KCTC 0934BP and KCTC 0935BP on Jan. 9, 2001.

For comparison, pcDNA3.1 was transfected alone into SNU16 and MKN28, which were cultured in the presence of G418. The clones obtained were used as controls, named control-SNU16 and control-MKN28.

1-3 Northern Blot Analysis

To determine whether the transfectants MKN28-Rx3, SNU16-Rx3-AS-c11, and SNU16-Rx3-AS-c12, established in Example 1-2, express the sense RUNX3 gene or the antisense RUNX3 gene in large quantities, Northern blotting analysis was conducted.

Total cellular RNA was prepared from each cell strain according to a standard method (Sambrook et al., *Molecular Cloning*, a laboratory manual, 2nd Ed., 1988, Cold Spring Harbor Laboratory). Five micrograms of each RNA preparation was separated on a 1.2% formaldehyde/agarose gel by electrophoresis and transferred onto a Hybond-N⁺ nylon membrane (Amersham). The RUNX3 cDNA of SEQ. ID. NO: 1 (Bae et al., *Gene*, 159 (2):245-248, 1995) was labeled with [α -³²P]dCTP for use as a probe for blotting analysis. Hybridization was conducted by incubating the Hybond-N⁺ nylon membrane and the radiolabeled RUNX3 DNA probe at 65° C. for 16 hours in 5 \times Denhardt's solution comprising 5 \times SSPE, 0.5% SDS and 100 mg/ml of salmon sperm. Afterwards, the nylon membrane was washed twice with 2 \times SSPE/0.1% SDS solution at room temperature for 15 min and twice with 0.1 \times SSPE/0.1% SDS solution at 65° C. for 15 min. The membrane was exposed to Kodak XAR-5 film for 16 hours at -70° C. to yield an autoradiogram.

On the autoradiogram obtained from the MKN28-Rx3 strain, a single band was observed at a position close to that of 18S RNA, verifying the expression of the RUNX3 cDNA in the strain. In contrast, SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 cells, which harbor the antisense RUNX3 cDNA, were found to give antisense RUNX3 gene transcripts of various sizes as detected by multiple bands on the autoradiogram. In the control-SUN16 strain, the expression of the endogenous RUNX3 gene was not detected by Northern blotting analysis (see FIG. 3).

Example 2

Cell Death Effect of TGF- β and Role of the RUNX3 Gene in TGF- β -Induced Cell Death

2-1 Cell Death Effect of TGF- β

SNU16 cells, which normally express the RUNX3 gene, and SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 cells in which an antisense RUNX3 cDNA is over-expressed to selectively inhibit the expression of the normal RUNX3 gene, were individually treated with TGF- β 1 (1 ng/ml), after which viable cells were counted over a period of time. The results are shown in FIG. 4a.

As seen in FIG. 4a, the TGF- β 1 treatment induced cell death in SNU16 cells. Almost all SNU16 cells were killed within 2 days after treatment with TGF- β 1 at a concentration of 1 ng/ml. However, the SNU16-Rx3-AS-c11 cells, in which the antisense RUNX3 gene was excessively

expressed to selectively inhibit the expression of the RUNX3 gene, were not killed after treatment with TGF- β 1. These results suggest that the RUNX3 gene is involved in TGF- β 1-induced cell death in SNU16 cells.

2-2 Role of RUNX3 Gene in TGF- β Induced Cell Death.

Cells dying through the programmed cell death pathway reveal a distinct pattern of DNA bands when electrophoresed in an agarose gel. After being treated with 1 ng/ml TGF- β 1 for 6 hours, the control-SNU16, SNU16-Rx3-AS-c11, and SNU16-Rx3-AS-c12 strains were lysed to isolate genomic DNAs according to a standard method (Sambrook et al., 1988, *Molecular Cloning: a laboratory manual*, 2nd Ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., USA). 0.2 μ g of each of the genomic DNA preparations was electrophoresed on a 1.5% agarose gel and visualized with ethidium bromide under UV light. The results are given in FIG. 4b.

As seen in this figure, apparent programmed cell death-specific apoptotic DNA bands were formed from the SNU16 cells treated with TGF- β 1 (FIG. 4b, lanes 1 and 2). No apoptotic bands were observed in the DNA from SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 cells, both of which over-express the antisense RNX3 gene, and therefore have a selective reduction in RUNX3 expression (FIG. 4b, lanes 3 to 6).

We concluded from these results that SNU16 cells are killed by TGF- β 1 stimuli through an apoptotic pathway and that RUNX3 is required for the process.

Example 3

Suppressive Activity of RUNX3 Against Tumorigenicity

It is known that TGF- β 1-induced apoptosis is a very important mechanism for suppressing the progression of normal cells toward cancer and that most of the factors in the TGF- β 1 pathway have tumor suppressor activity. The RUNX3 gene was assayed for tumor suppressor activity using nude mice, with the expectation that an anticancer effect would be seen.

To this end, the SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 strains, in which RUNX3 gene expression is selectively inhibited and the control-SUN16 strain were suspended in 0.85% phosphate buffered saline (PBS) at a density of 3×10^7 cells/ml, and 0.3 ml of each suspension was subcutaneously injected into groups of nude mice. Separately, the MKN28-Rx3 strain that shows over-expression of the sense RUNX3 RNA and the control-MKN28 strain were suspended in 0.85% PBS at a density of 5×10^7 cells/ml, and 0.3 ml of each suspension was subcutaneously injected into other groups of nude mice. Each strain clone was administered to 9 nude mice, which constituted one group. The tumorigenicity of MKN28-Rx3 and control MKN-28 strains was observed for 37 days after the injection. The tumorigenicity of SNU16-Rx3-AS-c11, SNU16-Rx3-AS-c12, and control-SNU16 cells was monitored for 98 days. During these time periods, the cancerous masses that formed were periodically measured for their external sizes. On the final days, the mice were sacrificed to excise the cancerous masses for comparison.

It was observed that tumors developed at higher rates in the mice into which the SNU16-Rx3-AS-c11 or the SNU16-Rx3-AS-c12 cells were injected than the mice into which the control SNU16 cells were injected, as shown in FIG. 5a. In detail, cancerous masses formed in the control SNU16

cell-injected group were measured to be 800 mm³ in volume at the final day. In contrast, cancerous masses of the groups into which SNU16-Rx3-AS-c11 and SNU16-Rx3-AS-c12 were injected were measured at 1,600 mm³ in volume, twice as large as those of the control group. Ninety-eight days after the injection, the cancerous masses were excised from the mice and compared to one another. The cancerous masses from the experimental group were remarkably larger than those from the control group, as seen in FIG. 5c.

MKN28-Rx3 cells, in which the sense RUNX3 gene is expressed, and control MKN28 cells were also assayed for tumor suppressor activity using nude mice in the same manner as was used for the SNU16 cells. Mice injected with MKN28-Rx3 cells exhibited significantly reduced tumorigenesis compared to the mice injected with the control MKN28 cells, as shown in FIG. 5b. In detail, the cancerous masses of the control group were measured at 600 mm³ in volume on the final day. In contrast, the cancerous masses of the MKN28-Rx3-injected experimental group were found to be as small as 250 mm³, demonstrating that the tumor growth was reduced by 58%. Thirty days after the injection, the cancerous masses were excised from the mice and compared to one another. The cancerous masses excised from the MKN28-Rx3-injected experimental group were macroscopically smaller than those excised from the control group (FIG. 5c).

Taken together, these results demonstrate that the RUNX3 gene has tumor suppressor activity.

Example 4

Analysis of the Expression Pattern of the RUNX3 Gene in Cancer Cells

4-1 RT-PCR Analysis

The expression pattern of the RUNX3 gene was analyzed in various cancer cell lines by RT-PCR. Total RNAs were isolated from 15 gastric cancer cell lines and 17 lung cancer cell lines and used to determine whether these cell lines produce RUNX3 mRNAs. The gastric cancer cells used in this analysis were SNU1, SNU5, SNU719, NUGC3, MKN1, MKN7, MKN28, MKN45, MKN74, AGS, KatoIII, RF1, RF48, and AZ521. The lung cancer cell lines were NCI-H522, 86-2, A549, LK79, LK87, LCSC#1, LCSC#2, NCI-H23, NCI-H226, NCI-460, NCI-H322, Sq-19, NCI-H1915, NCI-H630, Hs 888Lu, Lu65, and LX-1. All were obtained from the Korean Collection for Type Culture of Korea Research Institute of Bioscience and Biotechnology (KRIBB).

Total RNA was isolated from each cancer cell line, according to a standardized single-step guanidium method (Sambrook et al., *Molecular Cloning: a laboratory manual*, 2nd Ed., 1988, Cold Spring Harbor Laboratory). Using a Superscript kit (Gibco BRL) according to the manufacturer's protocol, reverse transcription was performed to prepare cDNA from the isolated total RNA with oligo-dTs serving as primers. Based on the cDNA, the RUNX3 gene was amplified by RT-PCR to determine the production of RUNX3 mRNA. For this RT-PCR, two pairs of PCR primers, PS-N and PS-C, were synthesized, which were composed of a set of the sense primer PS-NA SEQ. ID. NO: 4 and the antisense primer PS-NB SEQ. ID. NO: 5 and a set of the sense primer PS-CA SEQ. ID. NO: 6 and the antisense primer PS-CB SEQ. ID. NO: 7. The nucleotide sequences of these two pairs of PCR primers were designed on the basis of the cDNA nucleotide sequence of the RUNX3 gene SEQ. ID. NO: 1 (FIG. 5a).

For comparison, PEBP2 β /CBFB cDNA and β -actin cDNA were used as controls in the RT-PCR. To amplify the controls, a primer pair for PEBP2 β /CBFB, consisting of the sense primer h- β 5 SEQ. ID. NO: 8 and the antisense primer h- β 3 SEQ. ID. NO:9, and a primer pair for β -actin, consisting of the sense primer h-actin 5 SEQ. ID. NO: 10 and the antisense primer h-actin 3 SEQ. ID. NO: 11, were synthesized. Nucleotide sequences of these primer pairs were designed on the basis of human PEBP2 β cDNA and β -actin cDNA nucleotide sequences.

In a thermal cycler, such as "Model 9600" manufactured by Perkin-Elmer Cetus, RT-PCR was started with a 95° C. pre-denaturation of 50 μ l of each PCR solution for 2 min and carried out with 30 cycles of denaturing at 95° C. for 15 sec, annealing at 50° C. for 1 min, and extending at 72° C. for 1 min. After completion of the RT-PCR, 5 μ l of the RT-PCR product was electrophoresed on a 1.2% agarose gel and dyed with ethidium bromide for observation under UV light.

Among the gastric cancer cell lines, SNU5, SNU16, SNU719, MKN1, MKN45, RF1, RF48, MKN7, and AZ521 were found to produce RT-PCR products with the expected sizes (1,080 bp with the PS-N primer pair and 870 bp with the PS-C primer pair). However, MKN7 produced RT-PCR products in very small quantities, as shown in FIG. 6b. No RT-PCR products were obtained from the SNU1, NUGC3, MKN28, MKN74, AGS, and KatoIII cell lines when using either the PS-N primer pair or the PS-C primer pair.

Among the 17 lung cancer cell lines, expression of the RUNX3 gene was suppressed significantly in the HS888Lu cell line and completely in the NCI-H226, NCI-460, NCI-H630, and Lu65 cell lines, as shown in FIG. 6c.

4-2 DNA Southern Blot Analysis

Southern blotting analysis was conducted to verify the amplification of the RT-PCR products from the cDNA of the RUNX3 gene. The RT-PCR products were run on a 1.5% agarose gel under an electric field and the separated DNA fragments were transferred onto a Hybond-N⁺ nylon membrane.

Using the 2,244 bp BamHI fragment of the RUNX3 cDNA as a probe, hybridization was conducted in the same manner as in the Northern blotting analysis of Example 1-3, followed by autoradiography on XAR-5 films. The results are given in FIG. 6b.

As apparent from the autoradiogram, the RT-PCR products were amplified from the RUNX3 cDNA (PS-3 ST). In addition, the amplification of PEBP2 β and β -actin cDNAs from identical first cDNA strands was observed, indicating that the lack of amplification of RUNX3 cDNAs in the gastric cancer cell lines SNU1, NUGC3, MKN28, MKN74, AGS, KatoIII, NCI-H226, and lung cancer cell lines, NCI-460, NCI-H630, and Lu65 can be attributed to a lack of expression of the RUNX3 gene in those cell lines. Likewise, the low levels of the RT-PCR products in the MKN7 and Hs888Lu cell lines result from the low expression levels of the RUNX3 gene in those cell lines.

From the above results, it is apparent that the RUNX3 gene is remarkably or completely suppressed in about 47% of the gastric cancer cell lines and about 30% of the lung cancer cell lines. Because TGF- β -induced apoptosis does not occur and the tumorigenicity of cancer cells is increased in the substantial absence of RUNX3, the suppression of RUNX3 gene expression can be an important diagnostic index in about 47% of gastric cancer cells and about 30% of lung cancer cells (in about 37% of the total cancer cells examined).

Analysis of DNA Methylation Around Exon 1 of the RUNX3 Gene in Cancer Cell Lines

5-1 Analysis of DNA Methylation by Southern Blotting Analysis of Genomic DNA

To determine whether the suppression of RUNX3 gene expression was caused by DNA methylation, correlation between the expression of the RUNX3 gene and the hypermethylation of CpG islands at loci near RUNX3 exon 1 was analyzed through genomic DNA Southern blotting using restriction enzymes sensitive to DNA methylation.

Genomic DNAs were isolated from cancer cell lines according to a standardized SDS/protease K method (Sambrook et al., 1989). The isolated genomic DNAs were treated with a restriction enzyme that is unable to cut methylated DNA (SmaI), and separately, with one that is able to cut DNA irrespective of methylation (BamHI). The restriction enzyme digests were electrophoresed on a 1.5% agarose gel and then transferred onto a Hybond-N⁺ membrane. A genomic DNA NheI/MluI fragment comprising CpG islands of the RUNX3 gene was used as a DNA probe (FIG. 7b), and hybridization was carried out in the same manner as in the Northern blotting analysis of Example 1-3, followed by autoradiography on XAR-5 films. The genomic DNA was isolated from human genome DNA libraries with the RUNX3 cDNA of SEQ. ID. NO: 1 serving as a probe.

The SmaI digestion was observed to be greatly reduced specifically in those gastric cancer cell lines whose RUNX3 gene expression was not detected, that is, in the SNU1, NUGC3, MKN28, MKN74, AGS, and KatoIII cell lines, as shown in FIG. 7a. The MKN7 cell line in which the expression level was significantly low showed a partially methylated pattern (FIG. 7a). These results indicate that the expression of the RUNX3 gene is closely correlated with the level of DNA methylation at loci near RUNX3 exon 1 in cancer cell lines. Since RUNX3 gene-specific bands are detected in all of the cell lines despite differences in methylation, it can be concluded that some cancer cell lines in which the expression of the RUNX3 gene is not detected under normal conditions have the normal RUNX3 gene, but do not express it due to abnormal methylation.

5-2 Reactivation of the RUNX3 Gene

To determine whether the DNA methylation observed in Example 5-1 is the direct cause of the inactivation of the RUNX3 gene, the influence of a DNA methyltransferase inhibitor and a histone deacetylase inhibitor on the reactivation of the RUNX3 gene expression was analyzed.

The NUGC3, MKN28, and MKN74 cell lines (selected randomly from the cell lines showing no RUNX3 gene expression by RT-PCR analysis) were treated with 5-aza-3-deoxycytidine (AZA, 300 nM), which inhibits DNA methyltransferase, and trichostatin A (TSA, 1 mM), which inhibits histone deacetylase, alone or in combination. After 3 days of the treatment, total RNA was isolated from each cell line in accordance with the same standardized single-step guanidium method used in Example 4-1, and then analyzed for the expression of the RUNX3 gene by RT-PCR using the PS-N primer pair of SEQ. ID. NOS: 4 and 5.

Upon treatment with the DNA methyltransferase inhibitor or the histone deacetylase inhibitor, the expression of the RUNX3 gene was observed, as seen in FIG. 8. These results reveal that hyper-methylation suppresses the expression of the RUNX3 gene in about 37% of the gastric and lung

cancer cells owing to the hyper-methylation of CpG islands at loci near the exon 1 of the RUNX3 gene.

The expression plasmid capable of expressing the sense RUNX3 gene, constructed in Example 1-1, was assayed for acute toxicity as follows.

Example 6

Acute Toxicity Test upon Parenteral Administration of pEF-BOS-Rx3 to Rat

An acute toxicity test was conducted using rats of a specific pathogen-free SD lineage at 6 weeks of age. pEF-BOS-Rx3 (prepared as described in Example 1-1) was suspended in 1 ml of a physiological saline solution and administered to the anterior tibialis of two rats at a dose of 1 mg/kg by intramuscular injection. After the injection, the rats were observed for death, clinical traits, and body weight changes, and serological and serochemical analyses were conducted. In addition, the rats were sacrificed in order to observe the abdominal and thoracic organs. In neither animal were specifically noteworthy clinical traits found. No animals died from the administration of the test material, and no changes were observed in the body weight. The sero-

logical assay, serochemical assay, and autopsy test were all normal. Accordingly, pEF-BOS-Rx3 was identified as safe, with an LD₅₀ of at least 1 mg/kg upon parenteral injection.

5

INDUSTRIAL APPLICABILITY

As described previously herein, the RUNX3 gene of the present invention shows excellent tumor suppressor activity and is indispensably involved in TGF- β -dependent programmed cell death (apoptosis). Additionally, the suppression of RUNX3 gene expression is attributed to abnormal DNA methylation of CpG islands located near exon 1 of the RUNX3 gene. Accordingly, the RUNX3 gene and its products (RNA and proteins) can be utilized for the development of gene therapy for cancers. Measurement of the abnormal DNA methylation in the vicinity of exon 1 of the RUNX3 gene is useful for diagnosis of cancers. In addition, chemicals capable of regulating the abnormal DNA methylation and its consequences (for examples DNA methyltransferase inhibitors and HDAC inhibitors) can be used as anticancer agents because of their ability to promote the expression of the RUNX3 gene.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1

<211> LENGTH: 4213

<212> TYPE: DNA

<213> ORGANISM: human SNUX3 cDNA gene

<400> SEQUENCE: 1

```

cggccacttg attctggagg atttgttctg ggggtgcggc cgcggagtgc gggcggcgcg      60
gggcgagctt cggggcgggg ggcggcgcca gcggcacagc ccgcgcgggg ccccgccgcg      120
gcccaggcag ccgggacagc cacgaggggc ggccgcacgc ggggcccgcg gccgaggatg      180
cgggactagc cgggcaggct gcgggcgccc gtcgggccag cagggcctcg cagcgggccc      240
gcctggcgga gtattggccg gcgcgcgccc cctgogccct gatgcccggg ccccgccgct      300
tctgcttccc cgcttctcgc gcgagcggcg gccgaggagg cccccgcgcc ggccgcccccc      360
gggggaagcc gcgcgctctc gcgctccccg gcgcctgac ggccgctggt atgcgtattc      420
ccgtagacct aagcaccagc cgcgcttca cacctccctc cccggccttc cctgcccggc      480
gcggcgccgc caagatgggc gagaacagcg gcgcgctgag cgcgcaggcg gccgtggggc      540
ccggagggcg cgcggcgccc gaggtgcgct cgatggtgga cgtgctggcg gaccacgcag      600
gagagctcgt gcgcaccgac agccccaaact tcctctgctc cgtgctgccc tcgactggc      660
gtgcaacaa gaecgtgccc gtcgccttca aggtggtggc attgggggac gtgcccgatg      720
gtacggtggt gactgtgatg gcaggcaatg acgagaacta ctccgctgag ctgcccgaatg      780
cctcgccgct catgaagaac caggtggcca ggttcaacga ccttcgcttc gtgggcccga      840
gtgggcgagg gaagagtctc accctgacca tcaactgtgtt caccaacccc acccaagtgg      900
cgacctacca ccgaccatc aaggtgaccg tggacggacc ccgggagccc agacggcacc      960
ggcagaagct ggaggaccag accaagccgt tcctgaccg ctttggggac ctggaacggc     1020
tgcgcatgcg ggtgacaccg agcacacca gcccccaggg ctcaactcagc accacaagcc     1080

```

-continued

acttcagcag	ccagccccag	accccaatcc	aaggcaccto	ggaactgaac	ccattctccg	1140
acccccgcca	gtttgaccgc	tccttcccca	cgctgccaac	cctcacggag	agccgcttcc	1200
cagaccccag	gatgcattat	cccggggcca	tgctagctgc	cttcccctac	agegccaagc	1260
cctcggggcac	gagcatcagc	agcctcagcg	tgccggggcat	gccggccacc	agccgcttcc	1320
accataccta	cctcccccca	ccctaccogg	gggccccgca	gaaccagagc	gggcccttcc	1380
aggccaaccc	gtccccctac	cacctctact	acgggacatc	ctctggctcc	taccagttct	1440
ccatggtggc	cggcagcagc	agtgggggcg	accgctcacc	taccgcgatg	ctggcctctt	1500
gcaccagcag	cgctgcctct	gtgcgcgcgc	gcaacctcat	gaacccccagc	ctggggcgcc	1560
agagtgatgg	cgtggaggcc	gacggcagcc	acagcaactc	acccacggcc	ctgagcacgc	1620
caggccgeat	ggatgaggcc	gtgtggcgcc	cctactgacc	gccctggtgg	actcctcccg	1680
ctggaggcgg	ggaccctaac	aaccttcaag	accagtgatg	ggccggctcc	gaggctccgg	1740
gcgggaatgg	gacctgcgct	ccagggtggt	ctcgggccca	gggtggtccc	agctggtggg	1800
agcctctggc	tgcatctgtg	cagccacatc	cttgtacaga	ggcatagggt	accaccccc	1860
ccccggcccg	ggatactgcc	cccggcccag	atcctggccg	tctcatcccc	tacttctgtg	1920
gggaatcagc	ctcctgccac	ccccccggaa	ggacctcact	gtctccagct	atgcccagtg	1980
ctgcatggga	cccattgtct	ctgggacaga	ggccatctct	cttccagaga	gaggcagcat	2040
tggcccacag	gataagcctc	aggccctggg	aaacctcccc	acccctgcac	cttcgttggg	2100
gccctctcat	cccctgggtc	cagccccctc	tgcatttaca	cagatttgag	tcagaactgg	2160
aaagtgtccc	ccacccccac	caccctcgag	cggggttccc	ctcattgtac	agatggggca	2220
ggacccagca	cgctgctggc	agagatgggt	tgagaacaca	tccaagccag	tccccccagc	2280
ccagcttccc	ctcctgtcct	aactgttggc	tttccccag	cgcacgggt	cccaggcccc	2340
agagaagatg	agtctatggc	atcaggttct	taaaccaggg	aaagcaccta	cagaccggct	2400
cctccatgca	ctttaccagc	tcaacgcatc	cactctctgt	tctcttggca	ggcgggggga	2460
ggggggatag	gaggtcccct	ttcccctagg	tggtctcata	attccatttg	tggagagaac	2520
aggagggcca	gatagatagg	tcctagcaga	aggcattgag	gtgagggatc	attttgggtc	2580
agacatcaat	gtccctgtcc	cccctgggtc	cagccaagct	gtgccccatc	ccccaaagct	2640
cctgggagga	tccagccaaa	tcttgcgact	cctggcacac	acctgtctgt	aacctgtttt	2700
gtgctctgaa	agcaaatagt	cctgagcaaa	aaaaaaaaaa	aaacaaaaaa	acaaaaaaa	2760
aaacaaaacag	tttttaaaac	tgattttaga	aaaagaagct	taatctaacg	ttttcaaaac	2820
caaggtctct	tacaggtata	gttccgtgat	tatgatagct	ctgtgattat	aagcaacatc	2880
ccccccccct	ctcccccccg	cggaccccc	gctgcctcct	gaggggtgtg	ggttattagg	2940
gtctcaatac	tttctcaagg	ggctacactc	cccatcaggc	agcatcccac	cagcctgcac	3000
cacaggctcc	cctgggagga	cgagggaaac	gctgatgaga	cgctgggcat	ctctcctctg	3060
tggctctagg	acatctgtcc	aggaggctgg	gcggagggtg	gcaggatgtg	agaggtgggg	3120
agtactggct	gtcgtgggca	ggacagaagc	actgtaaagg	gctctccagc	cgcagctcag	3180
ctgcaactgc	ttccgaggtg	aagtcttgcc	cctgaatctt	gcaaaatggg	aaagtggggc	3240
cttgcccag	ggccaggctg	catggattct	cacatcagag	ttctctggcc	ctagaaaggc	3300
ttagaaaagg	cgtaagggaa	ctcataaagg	ctagcagcat	gcggtatctt	aactttctgc	3360
ctcggcctct	gtggatgcag	aaatctgccc	tacaaaaatg	tcttcattgg	ttgtctctgt	3420
gagagcactg	tccccacca	acctgtcaca	acggccagaa	ccatacacca	gagacacact	3480

-continued

```

ggcagggttag gcagtccttc tgggtgacct attccattcc ctctgctgc ggtttctctt 3540
ggcctgtcct cactggaaaa acagtctcca tctcctcaaa atagtgtctg actccctgca 3600
cccaaggggc ctctccatgc cttcttagga agcagctatg aatccattgt cctttagtct 3660
tcttccctcc tgttctctgg ttatagctgg tcccagggtca gcgtgggagg cacctttggg 3720
ttcccagtgc ccagcacttt gtagtctcat cccagattac taacccttcc tgatcctgga 3780
gaggcagggga tagtaataa attgctcttc ctaccccatc ccccatcccc tgacaaaaag 3840
tgacggcagc cgtactgagt ctgtaaggcc caaagtgggt acagacagcc tgggctggta 3900
aaagtaggtc cttatttaca aggtgctggt aaagtgtac taggcaaaca cactgatgta 3960
ggaagcacga gaaaggaag acgttttgat atagtgttac tgtgagcctg tcagtagtgg 4020
gtaccaatct tttgtgacat attgtcatgc tgagggtgta cacctgctgc actcatctga 4080
tgtaaaacca tcccagagct ggcgagagga tggagctggg tggaaactgc tttgcactat 4140
cgtttgcttg gtgtttggtt ttaacgcaca acttgcttgt acagtaaact gtcttctgta 4200
ctatttaact gta 4213

```

<210> SEQ ID NO 2

<211> LENGTH: 415

<212> TYPE: PRT

<213> ORGANISM: human SNUX3 protein

<400> SEQUENCE: 2

```

Met Arg Ile Pro Val Asp Pro Ser Thr Ser Arg Arg Phe Thr Pro Pro
 1           5           10           15
Ser Pro Ala Phe Pro Cys Gly Gly Gly Gly Lys Met Gly Glu Asn
 20           25           30
Ser Gly Ala Leu Ser Ala Gln Ala Val Gly Pro Gly Gly Arg Ala
 35           40           45
Arg Pro Glu Val Arg Ser Met Val Asp Val Leu Ala Asp His Ala Gly
 50           55           60
Glu Leu Val Arg Thr Asp Ser Pro Asn Phe Leu Cys Ser Val Leu Pro
 65           70           75           80
Ser His Trp Arg Cys Asn Lys Thr Leu Pro Val Ala Phe Lys Val Val
 85           90           95
Ala Leu Gly Asp Val Pro Asp Gly Thr Val Val Thr Val Met Ala Gly
100           105           110
Asn Asp Glu Asn Tyr Ser Ala Glu Leu Arg Asn Ala Ser Ala Val Met
115           120           125
Lys Asn Gln Val Ala Arg Phe Asn Asp Leu Arg Phe Val Gly Arg Ser
130           135           140
Gly Arg Gly Lys Ser Phe Thr Leu Thr Ile Thr Val Phe Thr Asn Pro
145           150           155           160
Thr Gln Val Ala Thr Tyr His Arg Ala Ile Lys Val Thr Val Asp Gly
165           170           175
Pro Arg Glu Pro Arg Arg His Arg Gln Lys Leu Glu Asp Gln Thr Lys
180           185           190
Pro Phe Pro Asp Arg Phe Gly Asp Leu Glu Arg Leu Arg Met Arg Val
195           200           205
Thr Pro Ser Thr Pro Ser Pro Arg Gly Ser Leu Ser Thr Thr Ser His
210           215           220
Phe Ser Ser Gln Pro Gln Thr Pro Ile Gln Gly Thr Ser Glu Leu Asn
225           230           235           240

```

-continued

Pro Phe Ser Asp Pro Arg Gln Phe Asp Arg Ser Phe Pro Thr Leu Pro
 245 250 255
 Thr Leu Thr Glu Ser Arg Phe Pro Asp Pro Arg Met His Tyr Pro Gly
 260 265 270
 Ala Met Ser Ala Ala Phe Pro Tyr Ser Ala Thr Pro Ser Gly Thr Ser
 275 280 285
 Ile Ser Ser Leu Ser Val Ala Gly Met Pro Ala Thr Ser Arg Phe His
 290 295 300
 His Thr Tyr Leu Pro Pro Pro Tyr Pro Gly Ala Pro Gln Asn Gln Ser
 305 310 315 320
 Gly Pro Phe Gln Ala Asn Pro Ser Pro Tyr His Leu Tyr Tyr Gly Thr
 325 330 335
 Ser Ser Gly Ser Tyr Gln Phe Ser Met Val Ala Gly Ser Ser Ser Gly
 340 345 350
 Gly Asp Arg Ser Pro Thr Arg Met Leu Ala Ser Cys Thr Ser Ser Ala
 355 360 365
 Ala Ser Val Ala Ala Gly Asn Leu Met Asn Pro Ser Leu Gly Gly Gln
 370 375 380
 Ser Asp Gly Val Glu Ala Asp Gly Ser His Ser Asn Ser Pro Thr Ala
 385 390 395 400
 Leu Ser Thr Pro Gly Arg Met Asp Glu Ala Val Trp Arg Pro Tyr
 405 410 415

<210> SEQ ID NO 3
 <211> LENGTH: 3300
 <212> TYPE: DNA
 <213> ORGANISM: CpG island in EXON 1 of human RUNX3 gene
 <400> SEQUENCE: 3

ccttctgctt cctagccctg ctgtggacaa cttaggggtgc tcttaggtgg gggccactgg 60
 ggagaaactg gcctgtttgt ccatcgatct gatggaagag ggagaaaaga cgacggtcca 120
 tgccaactgg ggaaggcgca ggggtgtctgc atgccccagg tgggggagtc ggagttctcc 180
 ctcccatcaa acagacgaca atttttgtcg gtccgggatg gggaggaagc aggtggaaat 240
 tgggaacaag ctagcctgtt ctgtgggtct tccccgggtg cctgaaaacg cgaagacaga 300
 aggcgtccat cttatcagag gtgaggaagg cggccttggg ttgacacaaa gccatcggtt 360
 tgtctgagac ctggccgcat ggatggcaag gaaagggcag ctcctcgggg gcggtcgccc 420
 cgctgcacc cggggccgca gctggcgcg atctgtagcc eggccgggccc cgcacctccc 480
 cggttgccag ggcgcggggc ccaactagcg gcggtcccgc ccactgtgcc tgccaggcgg 540
 ccccgcgacc ttggtctggg gacccaaggg cctgcaccgc cccccctcc cccgcggccc 600
 ctggtccctt ggatctggtg cccacgggga gccagcgccc tacggagagc cggagccgtg 660
 ccgggcccctg cgccaggtgc cagggccacc aagagtcctt cattctctgg aaacttgtct 720
 gtgaaccat cgtaagcgga gaggagaaa tcaggcggag gaacaagcaa agggcacggt 780
 gcaaacgaa accattcgac aaatggaatt taccaccacc tgaacagcg ggtcatggtc 840
 ccgcaactg ctcgagggca gcacgtgttg cccgccccgc gccagggccc tacctggcca 900
 cgacgcgctg cgcttctcg gagacgttcc cggaggtggg agcggccagg ctggatcaact 960
 cgctttcttc tagttctgct gctcgtgcca gcgcgtccga gggcgcgcg gcctgggtcc 1020
 gcaatcgaca actgccaggc gcaggctctc ttaaaaggtt cagtaaggga cctttgccgt 1080
 ccttccttc gacacggcct gagggcgtgc tgtgaggtcc cgaggtgggt aggggccagc 1140

-continued

```

tctcccgggtg gtcgggggtga gtcacagatc ctctgcccct ggagcgcacg cgggggcttga 1200
tttcgtttgg caacgacgaa atggcgcgcg ctgagcaggg gtcagatcca tgatgagatc 1260
ttgcggccac cgtcggatcc taagcttctt tgcttccgag gcttgggaatt cattgttttg 1320
cacctgtcga gagccggagg caacgaaaat ctagccccct ctccaaaagc gcggggaggc 1380
tcagcacgcg ttogttcccc agagtctagg gaggtgtctg gggcgataat tcggaatgat 1440
tgtggcttga tctttcccgt tgccctccca actgtagccg gcccttaggt ctgctcgaca 1500
gacttaggag gcgggagagc aaggggtgat ttgcagtcaa gccacagaga ggttggggca 1560
cgcggctggg agtgggagcg gggaccggga gccggcggg caggcagtgc cttggcgaag 1620
ctgtccgctg tccctcgccg gcagccggag cgcacgggcc caagaagaag tggggttggg 1680
cccgcagagg ccactttcca cccgcctgga gaaagaaaat tctctcctct gaaagcggag 1740
gcccttagct ttgcagccac tgctgttttt cttttgccac cgacgcgctg accgtttcac 1800
gatgcaggac cgtggttaca tgcgtaaagg aaaaaagaa aaacgcattt tgcaggcttc 1860
gtcgtgtttt tcaagagacc acagcccgcc acaacgaaga acgacgccgc gaggcctgca 1920
agatcctgaa acttgttttg aggggagagc agagaggaaa ggggttgttg gcccaggt 1980
acttagggtc ctaggagac tccctccgct ctgtcccggg tttggcacag gggccaccga 2040
ggctgggacc aaagccgccc agggctggga gcagcaaagg ccgcccggcg ggcgtggagc 2100
acgcgcaaaa tccctgtgtg ggtggaggct cttgggtcag aataatgtgc gggacgaggg 2160
aggtgagtaa cctctttggg gcggtcccca gtgcggcgtc accggccctg agaccgccg 2220
gccccagcc cggggttgca gaagtacag gccccaagca gcaagagctg ggaagcccc 2280
gccgcccga gcggggagga ggagcgaagg ggttgcgccc cagcgtcagg gagctacgac 2340
ccgagagagg gcggcaaggc gccttccgt gggaccggga cgttctaagc aaatttctag 2400
catttcccc gggctcccag agctctcggg ggccctgggc tgtggcactg gggcctctc 2460
cgcggggttg gccttccgc cctccccgt tgggcggcct ccggcaggcc cgttctctc 2520
ccgcgaacgc caccgagtg cccgcgatgg gggctccgcc gattggctgt gcgacgcgtc 2580
gctccgccag ccccgcctcg cgggcccggg ggtactaac cccgcgcggg cggcccgcgc 2640
cccgccactt gattctggag gatttgttct ggggctgegg ccgcggagtc ggggcccgcg 2700
cgggagagct tcggggcggg agggcggcgc agcggcacag ccccgccggg gcccccgcgc 2760
ggcccaggca gccgggacag ccacgagggg cggccgcacg cggggccgcg cgcagaggat 2820
gcgggactag ccgggcaggc tgcgggcggc cgtcgggcca gcgagcctc gcagcgggcg 2880
ggccctggcg agtagtgccc gggcgcggcc ccctgcgccc tgaggcccgg gcccccgcgc 2940
ttctgcttcc cgccttctcg cggcagcggc ggccgagggg gcgcccgcgc cggccgcccc 3000
cgggggaagc cgcgcgctct ccgctgccc ggccgctga cggccgctgt tatgctatt 3060
cccgtagacc caagcaccag ccgcccgttc acacctccct ccccggcctt cccctcgggc 3120
ggcggcggcg gcaagatggc cgagaacagc ggcgcgctga gcgcgcaggc ggcctggggg 3180
cccggagggc gcgcccggcc cgaggtgcgc tcgatggtgg acgtgctggc ggaccacgca 3240
ggcgagctcg tgcgcaccga cagcccaaac ttctctgct cgtgctgccc ctgcactgg 3300
3300

```

```

<210> SEQ ID NO 4
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: sense
primer of Ps-N primer pair

<400> SEQUENCE: 4

cgccacttga ttctggagga tttgt 25

<210> SEQ ID NO 5
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: antisense
primer of Ps-N primer pair

<400> SEQUENCE: 5

tgaagtggct tgtggtgctg agtg 25

<210> SEQ ID NO 6
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: sense
primer of Ps-C primer pair

<400> SEQUENCE: 6

gagtttcacc ctgaccatca ctgtg 25

<210> SEQ ID NO 7
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: antisense
primer of Ps-C primer pair

<400> SEQUENCE: 7

gcccatcact ggtcttgaag gttgt 25

<210> SEQ ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: sense
primer of PEBP2 beta/CBFB cDNA primer pair

<400> SEQUENCE: 8

cacgcttcca gaacgctgc 20

<210> SEQ ID NO 9
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: antisense
primer of PEBP2 beta/CBFB cDNA primer pair

<400> SEQUENCE: 9

cgagcctctt caaaggcctg t 21

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: sense

-continued

primer of beta-actin primer pair
 <400> SEQUENCE: 10
 ggacttcgag caagagatgg 20

<210> SEQ ID NO 11
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: antisense
 primer beta-actin cDNA primer pair

<400> SEQUENCE: 11
 gtcaccttca ccgttccagt 20

What is claimed is:

1. A method for diagnosing cancers caused by a loss of RUNX3 expression due to the methylation of the nucleotide sequence of SEQ. ID. NO: 3, characterized by detecting the methylation of a nucleotide sequence SEQ. ID. NO: 3, containing CpG islands present near exon 1 of the RUNX3 gene, comprising the steps of:

isolating genomic DNA from a tumor tissue of a subject; digesting the genomic DNA with a restriction enzyme sensitive to DNA methylation;

separating the genomic DNA digests on an agarose gel by electrophoresis and transferring them onto a membrane;

hybridizing the genomic DNA digests with a radiolabeled DNA probe for a RUNX3 gene; and

exposing the membrane to a film and detecting the methylation of the nucleotide sequence of SEQ. ID.

No: 3, whereby the detection of methylation indicates a cancer caused by a loss of RUNX3 expression.

2. A method for diagnosing cancers caused by a loss of RUNX3 expression due to the methylation of the nucleotide sequence of SEQ. ID. NO: 3, characterized by detecting the methylation of a nucleotide sequence SEQ. ID. NO: 3, containing CpG islands present near exon 1 of the RUNX3 gene, comprising the steps of:

isolating genomic DNA from the blood or tumor tissue of a subject;

digesting the genomic DNA with a restriction enzyme sensitive to DNA methylation;

performing a polymerase chain reaction (PCR) with the genomic DNA digests serving as templates and with parts of the nucleotide sequence of SEQ. ID. NO: 3 serving as primers;

separating the PCR products on an agarose gel by electrophoresis and detecting the amplification of a DNA sequence of interest, whereby the detection of amplification indicates a cancer caused by a loss of RUNX3 expression.

3. A method for diagnosing cancer caused by a loss of RUNX3 expression due to the methylation of the nucleotide sequence of SEQ. ID. NO: 3 comprising:

detecting methylation of a nucleotide sequence containing CpG islands present near exon 1 of the RUNX3 gene, which is isolated from a subject.

4. The method of claim 3, wherein the nucleotide sequence containing CpG islands comprises SEQ. ID. NO: 3.

5. The method of claim 1, wherein the cancer is gastric cancer or lung cancer.

6. The method of claim 2, wherein the cancer is gastric cancer or lung cancer.

7. The method of claim 3, wherein the cancer is gastric cancer or lung cancer.

* * * * *

专利名称(译)	RUNX3基因显示出抗肿瘤活性及其用途		
公开(公告)号	US7368255	公开(公告)日	2008-05-06
申请号	US10/466938	申请日	2001-01-30
[标]申请(专利权)人(译)	BAE SUK CHUL ITO昭		
申请(专利权)人(译)	BAE SUK 哲 ITO昭		
当前申请(专利权)人(译)	BAE SUK 哲 ITO昭		
[标]发明人	BAE SUK CHUL ITO YOSHIAKI		
发明人	BAE, SUK-CHUL ITO, YOSHIAKI		
IPC分类号	G01N33/574 G01N33/53 A61K38/00 A61K38/55 A61K48/00 A61P35/00 C07K14/47 C12N5/10 C12N15/00 C12N15/12 C12Q1/68 G01N33/566		
CPC分类号	C07K14/4747 A61P11/00		
优先权	1020010004018 2001-01-29 KR		
其他公开文献	US20040146986A1		
外部链接	Espacenet USPTO		

摘要(译)

本发明涉及显示抗肿瘤活性的RUNX3基因，其基本上参与TGF-β依赖性程序性细胞死亡(细胞凋亡)及其用途。此外，本发明发现RUNX3基因表达在各种胃癌和肺癌细胞系中受到抑制。RUNX3基因表达的抑制是由于位于RUNX3外显子(1)周围的CpG岛的超甲基化。本发明的RUNX3基因及其基因产物可有效地用于开发抗癌剂。RUNX3外显子(1)周围的CpG岛不仅可以用于开发调节异常DNA甲基化的抗癌剂，并且通过诱导RUNX3表达而且还可以用于通过测量异常DNA甲基化来开发癌症诊断方法。

FIG. 1a

```

CCGCCATTGATTCGAGGATTCCTGCGGGGCTGCGGGCCGGAGTCCG
          F1-HA
CCCCCGCCGGCCCGCCGAGCTTCGGCCGGAGCCGGGACAGCCAGAGGGG 100
CCCCCGCCGGCCCGCCGAGCTTCGGCCGGAGCCGGGACAGCCAGAGGGG 200
GGCCACAGCCGGCCCGCCGAGATTCGGCCGGAGCCGGGACAGCCAGAG 300
GGGGCCGGCCCGCCGAGGAGCCCTCCGAGGGGGCCGGCCCGCCGGGGA
GTATTGGCCGGCCCGCCGAGGAGCCCTCCGAGGGGGCCGGCCCGCCGG 400
TCTGCTTCCCGCTTCTCCGGCAGGGGGGGCCGGAGGGGGCCCGCCGGC
GGCCGCTTATPCCGTATTCCTTAGACCCAGAGCCAGCCAGCCCGCCCTCA
CACTCTCTCCGGGAGCCGGCTTCCCTGAGCCGGCCGGCCGGCCAGAGTGG 500
GAGAACAGCCGGCCGCTGAGCCGAGCCAGCCGGCCGGCCGGCCGGCCGG 600
CGGAGCTCCCTCCGAGCCAGAGCCCAACTTCCCTTCCCTGCTGCTGGCC
TCGGACTGGCCGGAGAGAGCCGGCCCGCCCTTCCAGAGGGTGGCC 700
ATTGGGGAGCTTCCCGGATGCTGAGGCTGGTGGTGGTGGTGGTGGTGGT
ACGAGACTACTCCGCTGAGCTGGGAGTGGCCGGCCGGCCGGCCGGAGAG 800
CAGGTGGCCAGCTTCAACGACTTCCCTTCCCTTCCCTTCCCTTCCCTTCC
GAGAGCTTCCAGCTTCCAGACTTCTTCCCTTCCCTTCCCTTCCCTTCCAG 900
          F2-7A
CGACTACCACCCGAGCCATCAAGGAGCCGTTGACGGAGCCCGGGAGCCG 1000
AGAGGGCAGCCGGCAGAGGCTGGAGGACAGAGCCAGCCAGCCCTTCCCTAC 1100
CTTTGGGGAGCTTGGAGCCGCTCCGATCCGGGCGAGACCGGAGCAGCCCA
GGCCCCGAGGCCTGACTCAGGACCAAGAGCCGATTCAGGACCCAGCCCCAG
          F2-NB
ACCCCAATCCAGGACCTTCGGAACTCAACCCATTCTCCGAGCCCGCCCA 1200
GTTTGACCCCTCCTTCCGACCTTGGCAACCTCGAGGGAGCCGCTTCC 1300
CAGACCCAGGATGCATATCTCCGGGCCATGTCAGCTGCCCTTCCCTTAC
AGCCACAGCCCTGCGCAGAGATTCAGCAGCCCTCAGCCCTGGGGGCT 1400
GCCGGCCAGCAGCCCTTCCACCATACCTACTCCCGCCAGCCCTTACC 1500
CACCTCTACTAGGGGACATCCCTCGGCTCCTACAGGTTCTCCCTATGGTGG
CGGAGCCAGAGTGGGGGAGCCCTTCCCTTCCCTTCCCTTCCCTTCCCT 1600
GCACAGCAGCCCTTCCCTTCTTCCGGCCGGCCAGCTCATGAAGCCCGAG
CTGGCCGGCCAGGATGATGGCTGGAGCCGAGCCAGCCAGCCAGCT 1700
ACCCAGSCTTGAAGCCGAGCCGATGGATGGATGGAGCCCTTGGCCGG
CCTACGAGCCGCTGAGCTTCCCTTCCCTTCCCTTCCCTTCCCTTCCAG
          F2-CB
AACCTTCAAGACCACTGATGGCCGGCTCCGAGGCTCCGGGGGGGAATGG 1800
GACCTGCGCTCCAGGGTGGTCTCGGCTCCAGGTTGCTCCAGCTGGTGGG
AGCCCTTGGCTGCACTCTGTGGAGCCATCCCTTGTACAGAGGATAGGTT 1900
ACCAGCCCGCCGAGATTCCTGGGAACTTGGCCCGCCGGCCAGATGGCCG
TCTCATCCCATACTCTGTGGGGAATCAGCCCTTCCGACCCCGCCGAA 2000
GGAGCCAGCTGCTCAGCTATGGCCAGAGAGAGCCAGGATTTGGCCAG
GATAGCCCTCAGCCCTTGGAACTTCCGAGCCGGCCAGCCAGCCAGCT 2100
GCCCTGCTGATCTTGGTCCAGCCCTTCCCTTCCCTTCCCTTCCCTTCC
TCAGAACTGAAAGTGTCCCGCCAGCCAGCCCTTCCGAGCCGATTTCC 2200
CTCATTTAGAGATGGCCAGAGCCAGCCAGCCCTTCCCTTCCCTTCCCT 2300
TGAGAGCAGATCCAGGACAGTCCCGCCAGCCAGCCCTTCCCTTCCCT 2400
AGCTATGGCATCAGGTTCTTAAACCCAGGAAAGCCCTTACAGAGCCGTT
CTCCATGACTTACCCCTGAGCCAGTACACTCTTCTTCTTCTTCTTCT 2500
CGCCGGGAGGGGGATAGAGGTTCCCTTCCCTTCCCTTCCCTTCCCTT 2600
ATTCCATTTGTGGAGAACCTGAGGGCCAGATAGATAGGCTCCAGCA 2700
AGGCATTGAGGTGAGGATCATTTTGGGTCAGACATCAATCTCCCTGCC
CCCTGGGTCAGCCAGCTGTGCCCATCCCCAGCCCTCTGGAGGA

```