



US 20030138791A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0138791 A1**

**Haalck et al.**

(43) **Pub. Date: Jul. 24, 2003**

(54) **3-SPIRO-CYANIN FLUOROCHROMES AND THEIR USE IN BIOASSAYS**

(52) **U.S. Cl.** ..... **435/6; 546/277.4; 546/15; 548/159; 548/217; 548/300.7; 548/408**

(76) **Inventors: Lutz Haalck, Muenster (DE); Erk Gedig, Muenster (DE); Hans-Juergen Hoelpert, Muenster (DE); Viola Podsadlowski, Muenster (DE)**

(57) **ABSTRACT**

Correspondence Address:  
**Felix J D'Ambrosio  
Jones Tullar & Cooper  
P O Box 2266 Eads Station  
Arlington, VA 22202 (US)**

The invention concerns sterically shielded, stabilized fluorescent dyes based on symmetrical or unsymmetrical 3-spirocyanines and their utilization as marker-molecules in analytical procedures. The fluorescent dyes according to the invention are characterized by a strongly reduced proneness towards aggregation due to the insertion of bulky spiro-substituents and thus enable to reach a high degree of coupling to the target molecule without a significant loss of fluorescence due to self quenching effects. They therefor exhibit a high quantum yield and are thus excellently suitable as fluorescent dyes, especially as NIR-fluorescent dyes, for the coupling or binding to biological and non-biological target molecules used in analytical methods. The analytical methods comprise all methods in which fluorescence-optical methods are used to detect biological and non-biological molecules.

(21) **Appl. No.: 10/130,547**

(22) **PCT Filed: Sep. 3, 2001**

(86) **PCT No.: PCT/EP01/10130**

(30) **Foreign Application Priority Data**

Sep. 19, 2000 (DE)..... 100 46 215.4

**Publication Classification**

(51) **Int. Cl.<sup>7</sup>** ..... **C12Q 1/68; C07D 417/02; C07D 43/02; C07D 413/02**




Fig. 1




Fig. 2




Fig. 3




Fig. 4






Fig. 6




Fig. 7





Fig. 8




5

10

15

Fig. 9



5

10

15 Fig. 10




Fig. 11




Fig. 12




Fig. 13

### 3-SPIRO-CYANIN FLUOROCHROMES AND THEIR USE IN BIOASSAYS

[0001] The invention concerns sterically shielded, stabilized fluorescent dyes on the basis of symmetrical or unsymmetrical 3-spirocyanines and their application as marker-molecules in analytical methods.

[0002] Fluorescent dyes are increasingly used to label enzymes, antibodies and nucleic acids and employed in immunoassays, fluorescence microscopy and for sequencing. Traditionally, fluorescent dyes are used which are excited using light from the UV or visible region of the electromagnetic spectrum. Their excitation and emission spectra overlap with the given natural serum fluorescence, making them only partially suitable for analysis of natural samples (e.g. blood and serum samples, cells). Fluorescent dyes with excitation wavelength in the near infrared region of the light spectrum are thus used since a couple of years, allowing the measurement of biological samples with low background fluorescence and concomitantly increased sensitivity.

[0003] The demand of biochips (protein-, DNA-arrays) that enable the specific detection of proteins as well as DNA or fragments thereof, grows exponentially due to scientific developments in the proteomics and genomics sector. Most of these applications also involve the use of fluorescent dyes.

[0004] The use of reactive and water soluble NIR-fluorophores for coupling with biomolecules was first described in U.S. Pat. No. 5,268,486. Chemical groups are inserted, preferably as N-substituents, that are suitable for coupling with reactive amino acid side chains (e.g. amino-, thio-, carbonyl-, or hydroxy groups), some of these chemical groups being isothiocyanates, thiocyanates, hydrazines, hydroxy-succinimidylesters, disulfides etc. This principle was extended in recent time beyond symmetrical cyanomethine dyes (2,2'-indo-cyanines) to encompass structurally related merocyanines and styrylcyanine dyes.

[0005] All dyes of the cyanine type as well as other commonly used fluorescent dyes, e.g., rhodamines, show a characteristic behavior in aqueous solution. Due to the planar chemical structure (arrangement) of the fluorophores, aggregation or dimer-formation occurs especially at higher coupling ratios (molar ratio dye/protein; D/P>5). This leads to drastically reduced fluorescence due to radiation free intermolecular transition processes. This phenomenon arises in solutions, e.g. in presence of high concentrations of salts, as well as with protein bound fluorophores. Dyes without additional hydrophilic groups and amino-substituted cyanine dyes are especially sensitive to this problem.

[0006] The dyes described in U.S. Pat. No. 5,268,486 and DE 39 120 46 could only reduce but not eliminate this problem through insertion of arylsulfonates to raise their water solubility. In general, the quantum yield in aqueous solutions, however, is reduced dramatically compared to organic solvents.

[0007] Many different interactions are utilized for the labelling of oligonucleotides or DNA/RNA. Besides the classical intercalation dyes like ethidium bromide, the specific ionic binding to phosphate groups of nucleotides (U.S. Pat. No. 5,410,030) by positively charged dyes and various covalent couplings of dyes to modified purin/pyrimidinbases (U.S. Pat. No. 6,027,709) as well as to phosphate groups of

nucleotides are feasible. This development reflects the high increase in demand of fluorescent dyes for DNA labelling and sequencing.


[0008] It is the task of the presented invention to suggest fluorescent dyes that exhibit a high quantum yield and simultaneously a low tendency towards aggregation. Furthermore, these fluorophores shall cover a wide variety of different ways to bind to a vast number of target molecules and thus allow for a wide variety of application fields.

[0009] The task is solved by the characteristic features described in claim 1. The other dependant claims show further advantageous refinements (embodiments).


[0010] The application of fluorophores according to the invention is described by claims 37 to 40. Fluorescent dyes are suggested according to this invention, which are based on a basic structure of symmetrical or unsymmetrical cyanine dyes with at least one sterically demanding spiro-substituent in 3-position of the indole headgroup. These fluorescent dyes will be referred to as spironines. The invention based fluorescent dyes have the following basic structure:

[0011] Fluorescent dyes of the general structure A-Z-A' with


[0012] A a headgroup of general formula I



[0013] wherein X is




[0014] O, S, SO, SO<sub>2</sub> oder




[0015] W<sup>-</sup>=counter ion, preferably selected from the group of halogen, perchlorate or tosylate,


[0016] A' is selected from a headgroup having the general formulas II to V,




or




or




or




[0017] wherein Y is O, S, NR<sub>17</sub>,




oder



[0018] and T is



[0019] O, S, SO, SO<sub>2</sub> or




III

[0020] and

[0021] Z represents a group with the general formula VI or VII


VI



or

VII

IV



[0022] with n is an integer selected from the group consisting of 0, 1, 2, or 3 or

[0023] wherein at least one of the residues R<sub>1</sub> to R<sub>26</sub> is selected from a group consisting of

[0024] a) a chemical reactive group for covalent coupling to a target molecule or

V

[0025] b) an ionic group for coupling to a target molecule by ionic interaction forces or

[0026] c) a lipophilic group for coupling to a target molecule by adsorption.

[0027] By means of the selective introduction of the groups a) to c) into the fluorescent dye, it is made possible to bind the dye to different target molecules and thus allow for different fields of applications.

[0028] The other groups are chosen independently from each other from the groups H, alkyl (C<sub>1</sub>-C<sub>10</sub>), alkoxy (C<sub>1</sub>-C<sub>10</sub>), trifluoromethyl, halogen, sulfonic acid, sulfonate, phosphoric acid and phosphonate.

[0029] Through use of cyanine dyes with bulky substituents in 3-position of the indole headgroup, especially cyclic substituents, so called spiroes, e.g. spiro-1'-cyclohexan, spiro-4'-1'-piperidine, 4'-1'-tetrahydropyran, spiro-4'-1'-tetrahydrothiopyran, spiro-4'-1'-oxotetrahydrothiopyran, the central polymethine chain can be extensively shielded.

[0030] Surprisingly, it could be shown according to the invention that fluorescent dyes substituted like this exhibit a significant reduced tendency towards aggregation. Furthermore, a higher photostability compared to commercially-


available cyanine dyes was observed. Particularly interesting in this matter is the utilization of cyclic substituents, e.g. substituted spiro-cyclohexanes, spiro-piperidines, spiro-tetrahydrooxopyranes or spiro-tetrahydrothiopyranes. Based on these spiro compounds, a new elegant way is created to control the water solubility by choosing a polar or non-polar substituent as well as to positively influence the binding behavior of the dye by insertion of reactive groups  $R_x$  or ionic groups. The shielding and thus stabilizing effect of the spirane groups can be amplified through further insertion of bulky substituents at the spiro-residues.

[0031] The invention based fluorophores can be excited by light of a wavelength of 600-1000 nm. Typical for the fluorescent dyes are high molar extinction coefficients and very high quantum yields. The Stokes shift normally is at least 15 nm. A utilization of these dyes in a broad field of applications is made possible by cost effective laser diodes emitting light in the range of 670-830 nm.



[0032] The groups  $R_1$ ,  $R_4$ ,  $R_6$  to  $R_{10}$ ,  $R_{13}$  and  $R_{17}$  to  $R_{21}$  in the general formulas I to V are of higher significance. These positions are especially suitable for insertion of substituents to increase the water solubility, the insertion of non-polar substituents to increase the lipophily and/or at least one reactive group  $R_x$  for the coupling to biomolecules.

[0033] Thus, the fluorescent dyes contain at least one reactive group  $R_x$  for the coupling to target molecules in the general formulas I to VII advantageously as groups  $R_1$ ,  $R_4$ ,  $R_6$  to  $R_{10}$ ,  $R_{13}$  and  $R_{17}$  to  $R_{21}$ , particular advantageously as groups  $R_1$ ,  $R_6$  to  $R_{10}$  and  $R_{18}$  to  $R_{21}$  and particular advantageously as groups  $R_6$  to  $R_9$  and  $R_{18}$  to  $R_{21}$ . The reactive group  $R_x$  can be covalently bound directly to the dye or through a bridge made by several atoms and exhibits a suitable chemistry. The reactive group  $R_x$  is preferentially chosen from the group of carbonic acids, activated esters, acylazides, acylhalogenides, acylnitriles, aldehydes, anhydrides, arylamides, alkyl halides, anilines, alkylsulfonates, arylhalogenides, thiols, azides, aziridines, borates, carbodiimides, diazoalkanes, epoxides, glycerols, haloacetamines, halotriazines, hydrazines, hydroxylamines, imidoesters, isocyanates, isothiocyanates, maleimides, phosphoramidites, silyl halides, sulfonates and sulfonylchlorides. Subsequently some exemplary groups  $R_x$  and their structures are shown.

[0034] activated carbonic acid




-continued




[0035] with n, m and p independently from each other equal to 1 to 8

[0036] carbodiimides




[0037] with n equal to 1 to 8

[0038] anhydrides



[0039] with n equal to 1 to 8


[0040] carbonic acid azides



[0041] With n equal to 1 to 8



[0042] Coupling of nucleophiles through

[0043] epoxides





[0044] with n equal to 1 to 8

[0045] isothiocyanates




[0046] with n equal to 1 to 8

[0047] isocyanates




[0048] with n equal to 1 to 8


[0049] aziridines



[0050] with n equal to 1 to 8




[0051] maleimides




[0052] with n equal to 1 to 8

[0053] pyridyl-disulfide activated groups




[0054] with n and m independently from each other equal to 1 to 8

[0055] halodi- und triazines




[0056] with W is equal to chlorine or bromine and n equal to 1 to 8

[0057] vinylsulfones




[0058] with n is equal to 1 to 8

[0059] acylimidazoles



[0060] with n is equal to 1 to 8

[0061] phosphoramidite



[0062] with n is equal to 1 to 8

[0063] Tab. 1 shows an overview over possible ways of creating dye-conjugates by covalent binding.

[0064] Additionally, the fluorescent dye according to the invention may contain at least one ionic group. Some of the preferred ionic groups for example are selected from the group of carbonic acids, sulfonic acids, sulfonates, phosphoric acids, phosphonates, phosphordiesteres, phosphortriesteres and primary to quaternary amines. Advantageously at least one of the groups  $R_1$ ,  $R_4$ ,  $R_6$  to  $R_{10}$ ,  $R_{13}$  and  $R_{17}$  to  $R_{21}$ , particular advantageously at least one of the groups  $R_1$ ,  $R_6$  to  $R_{10}$  and  $R_{18}$  to  $R_{21}$  and advantageously at least one of the groups  $R_6$  to  $R_9$  and  $R_{18}$  to  $R_{21}$ .

[0065] Further, one group of the group A, chosen from the group  $R_1$ ,  $R_6$  to  $R_9$ , and one group of the group A', chosen from the group  $R_{10}$ ,  $R_{17}$  to  $R_{21}$ , can represent ionic groups with opposite charges. This has the special advantage that an additional stabilization of the fluorescent dye is made possible by this intramolecular ionic bridge. It is also possible for the ionic groups in the groups A and A' to have the same charge and to form a complex with an ion of opposite charge. It is preferred hereby to have two anionic groups that form a complex with a metal-ion from the 1. to 3. main- or subgroup.

[0066] To raise the solubility in water, for example groups like carbon acids, carbohydrates, sulfonic acids, sulfonates, phosphates, phosphonates, amines, halogens, polyoles or polyethers could be chosen and inserted at any position of the fluorescent dye, advantageously as at least one of the groups  $R_1$ ,  $R_4$ ,  $R_6$  to  $R_{10}$ ,  $R_{13}$  and  $R_{17}$  to  $R_{21}$ , particular advantageously as at least one of the groups  $R_1$ ,  $R_6$  to  $R_{10}$  and  $R_{18}$  to  $R_{21}$  and most particular advantageously as at least one of the groups  $R_6$  to  $R_9$  and  $R_{18}$  to  $R_{21}$ .

[0067] To raise the lipophily, long-chain saturated or unsaturated alkyl groups (preferably  $C_6$  to  $C_{18}$ ) or fatty

acids/fatty alcohols could be chosen and inserted at any position of the fluorophor, advantageously at least one of the groups  $R_1$ ,  $R_4$ ,  $R_6$  to  $R_{10}$ ,  $R_{13}$  and  $R_{17}$  to  $R_{21}$ , particularly advantageously at least one of the groups  $R_1$ ,  $R_6$  to  $R_{10}$  and  $R_{18}$  to  $R_{21}$  and especially advantageously at least one of the groups  $R_6$  to  $R_9$  and  $R_{18}$  to  $R_{21}$ .

[0068] In another advantageous embodiment of the invention two ortho residues at the aromatic ring may be combined to form at least one additional aromatic, carbo- or heterocyclic ring.

[0069] The dye can form complexes or conjugates with a biological or non-biological target molecule by creation of one or more covalent bonds using one or more groups  $R_x$ , the target molecule preferably being from the group of antibodies, proteins, peptides, enzyme substrates, hormones, lymphokines, lipids, phospholipids, metabolites, receptors, antigens, haptens, lectines, toxins, carbon hydrates, oligosaccharides, polysaccharides, nucleic acids, desoxyribonucleic acids, derivatized desoxyribonucleic acid, derivatized nucleic acids, DNA-fragments, RNA-fragments, drugs, virus particles, virus components, yeast, yeast components, bacteria, bacteria components, blood cells, blood cell components, biologic cells, non-cellular blood components, poisons, polymers, polymer particles, glass particles, glass surfaces, plastic surfaces, plastic particles, polymer membranes, metals, conductors or semiconductors.


[0070] In another advantageous refinement fluorescent dyes are bound to the surface of or incorporated into nano- or microparticles, mostly on the basis of polymer materials. These particles can then be used advantageously in analytical methods.

[0071] The fluorescent dyes according to the invention especially distinguish themselves from the state of the art by a strongly reduced proneness towards aggregation due to the insertion of bulky spiro-substituents and thus a high degree of coupling to the target molecule without self-quenching of the fluorescence. They therefore exhibit a high quantum yield and are thus excellently suitable as fluorescent dyes, especially as NIR-fluorescent dyes, for the coupling or binding to biomolecules used in bioassays. The analytical methods comprise all methods in which fluorescence-optical methods are used to detect biomolecules. A preferred way of realization are fluorescence-immuno-tests which are based on known biochemical assays of general receptor-ligand systems, for example antibody-antigen, lectin-carbohydrate, DNA or RNA-complementary nucleic acids, DNA or RNA-proteins, hormone receptors, enzyme-enzyme co-factors, protein G or protein A-immunoglobulin or avidin-biotin.


[0072] If the fluorescent dye according to the invention contains adequate nucleophilic groups, preferably amino-, thio-, or hydroxy-groups, a coupling of the dye, for example after activation as phosphoramidite, to a nucleotide becomes possible. This process is especially important for the preparation of dye-labelled (desoxy-)nucleotides, which for example can be used in sequencing machines.

[0073] Some representative fluorescent dyes according to the invention are summarized in the following.

Formula VIII




Formula XI




[0074] with  $M^+$ =metal cation and with X, T, and n defined according to claim 1;

[0077] with  $M^+$ =metal cation, X, Y, T,  $W^-$ , m and n are defined according to claim 1.

Formula IX



Formula XII




[0075] with  $M^+$ =metal cation, NHS=N-Hydroxy-succinimide and with X, T,  $W^-$  and n defined according to claim 1.


[0078] with X, Y, T,  $W^-$ , m and n defined according to claim 1, p is equal to 1 to 8.

[0079] Formula XIII

Formula X





Formula XIII




[0076] with X, T,  $W^-$  and n defined according to claim 1, m is equal to 0, 1-17.

[0080] with X and n defined according to claim 1, q is equal to 1 to 8.




[0083] with  $W^-$  and  $n$  defined according to claim 1.


[0081] with  $M^+$ =metal cation and  $n$  defined according to claim 1, NHS=N-hydroxy-succinimid,  $r$  and  $s$  independently from each other equal to 1 to 8.



[0082] with  $W^-$  and  $n$  defined according to claim 1.




[0084] with  $W^-$  and  $n$  defined according to claim 1, NHS=N-hydroxy-succinimide, PEG=poly ethylene glycole and with  $t$  and  $u$  independently from each other equal to 1 to 8.




[0085] with W and n defined according to claim 1, R'=trityl, R''=phosphoramidite, v and w independently equal to 0 to 6.

Formula XIX



[0086] with M<sup>+</sup>=metal cation, X, Y, T, W<sup>-</sup> and n defined according to claim 1.

Formula XX



[0087] with M<sup>+</sup>=metal cation, X, Y, T, W<sup>-</sup> and n defined according to claim 1, z equal to 0, 1 to 6.

[0088] The invention will be described in more detail by way of figures and examples below, without restricting it to them.

[0089] FIG. 1 Synthesis of compounds S1 to S3. (Part 1)

[0090] FIG. 2 Synthesis of compounds S1 to S3. (Part 2)

[0091] FIG. 3 Synthesis of compounds S4 to S7. (Part 1)

[0092] FIG. 4 Synthesis of compounds S4 to S7. (Part 2)

[0093] FIG. 5 Synthesis of compound S8.

[0094] FIG. 6 Synthesis of compound S9. (Part 1)

[0095] FIG. 7 Synthesis of compound S9. (Part 2)

[0096] FIG. 8 Absorption spectrum of the fluorescent dyes (compounds S1 to S3) according to the invention.

[0097] FIG. 9 Absorption- and emission-spectrum of compound S2.

[0098] FIG. 10 Absorption- and emission spectrum of the protein conjugate according to example 10.

[0099] FIG. 11 Photostability of compound S3 (according to example 3).

[0100] FIG. 12 Fluorescence of the protein conjugate (compound) according to example 10 depending of the incubation time.

[0101] FIG. 13 Fluorescence of compound in the Fluorescence of compound in presence of high protein concentrations presence of high protein concentrations.

[0102] Preparation of dyes of the symmetrical 3-spiro-1'-cyclohexan indole type

#### EXAMPLE 1:

[0103] Synthesis of dye S1

[0104] Preparation of 2-methyl-3-spiro-1'-cyclohexan-5-sulfo-3H-indole; potassium salt (FIG. 1, 1a)

[0105] 30 mmol (5.646 g) phenylhydrazine sulfonic acid is heated with 60 mmol (7.5729 g) cyclohexylmethylketone and 60 mmol (5.889 g) potassium acetate in 30 ml glacial acetic acid for 6 h under reflux. Subsequently the volatile components were evaporated under reduced pressure. The brown residue is transferred into an extraction thimble and extracted with 200 ml 2-propanole using a Soxhlet apparatus. The product starts crystallising in the flask even in the heat, after cooling it is filtered with suction and dried in vacuum.

[0106] Yield: 6.044 g (66%)

[0107] Preparation of 1-(5-carboxypentyl)-2-methyl-3-spiro-1'-cyclohexan-5-sulfo-3H-indolium bromide; potassium salt (1b)

[0108] 4.5 mmol (1.418 g) 2-methyl-3-spiro-1'-cyclohexan-5-sulfo-3H-indole is mixed with 5.5 mmol (1.073 g) 6-bromohexanoic acid and reacted in the melt at 110° C. for 2.5 h. After cooling the mixture is triturated with acetone, filtered, repeatedly washed with acetone and finally dried in vacuum.

[0109] Yield: 1.505 g (66%)

[0110] The quaternary spiro-indole (2c) is the key component for the further steps in the synthesis.

[0111] Dye S1

[0112] 1-(5-carboxypentyl)-2-[5-(1-(5-carboxypentyl)-1,3-dihydro-3-spiro-1'-cyclohexan-5-sulfo-2H-indole-2-ylidene)-propenyl]-3-spiro-1'-cyclohexan-5-sulfo-3H-indolium hydroxide, inner salt, potassium salt: (FIG. 2, 1c)

[0113] 0.4 mmol (205 mg) 1-(5-carboxypentyl)-2-methyl-3-spiro-1'-cyclohexan-5-sulfo-3H-indolium bromide (potassium salt) together with 1 mmol (148 mg) triethoxymethane in 2 ml pyridine are heated under reflux for 30 minutes. After cooling the crude dye is precipitated with ether and purified by preparative HPLC using RP-18 and ethanol:water gradient.

[0114] According to this procedure cyanine dyes with longer polymethine chain are also accessible.

#### EXAMPLE 2:

[0115] Synthesis of dye S2

[0116] 1-(5-carboxypentyl)-2-[5-(1-(5-carboxypentyl)-1,3-dihydro-3-spiro-1'-cyclohexan-5-sulfo-2H-indole-2-ylidene)-penta-1,3-dienyl]-3-spiro-1'-cyclohexan-5-sulfo-3H-indolium hydroxide, inner salt, potassium salt: (FIG. 2, 1d)

[0117] 0.22 mmol (111.6 mg) 1-(5-carboxypentyl)-2-methyl-3-spiro-1'-cyclohexan-5-sulfoindolium bromide (potassium salt) were dissolved in 1 ml of dry pyridine and refluxed. Then 0,5 ml tetraethoxypropane is added in por-

tions over a period of 2.5 h. The precipitated dye is repeatedly washed with pyridine, finally washed with diethylether and purified by preparative HPLC using methanol water gradient on silicagel RP-18.

#### EXAMPLE 3:

[0118] Synthesis of dye S3

[0119] 1-(5-carboxypentyl)-2-[5-(1-(5-carboxypentyl)-1,3-dihydro-3-spiro-1'-cyclohexan-5-sulfo-2H-indole-2-ylidene)-hepta-1,3,5-trienyl]-3-spiro-1'-cyclohexan-5-sulfo-3H-indolium hydroxide, inner salt, potassium salt (FIG. 2, 1e)

[0120] 0.32 mmol (125 mg) 1-(5-carboxypentyl)-2-methyl-3-spiro-1'-cyclohexan-5-sulfoindolium bromide (potassium salt) together with 0.16 mmol (45 mg) glutaconedianil hydrochloride and 0.65 mmol (63 mg) potassium acetate in a mixture of 2 ml acetic anhydride and 0.5 ml glacial acetic acid are heated to reflux for 30 minutes. The crude dye is precipitated with acetone, filtered off and purified by preparative HPLC using a ethanol-water gradient on silicagel RP-18.

[0121] Preparation of dyes of the asymmetric 3-spiro-1-cyclohexan indole type

#### EXAMPLE 4:

[0122] Synthesis of dye S4

[0123] 2-methyl-3-spiro-1'-cyclohexan-3H-indole (FIG. 3, 2a)

[0124] 1.25 mol (135.2 g) phenylhydrazine and 1.5 mol (189.3 g) cyclohexylmethylketone in 1.25 l glacial acetic acid are heated to reflux for 3 h. The glacial acetic acid is evaporated in vacuum, 1 l of water is added to the residue and the solution is extracted 4 x with ether. After the etheric layer is washed with sodium hydrogencarbonate solution and dried with sodium sulfate, the ether is evaporated and the residue is distilled in vacuum.

[0125] bp.(2 mbar): 133-136° C.

[0126] yield: 192.6 g (77%)

[0127] 1-(4-sulfobutyl)-2-methyl-3-spiro-1'-cyclohexan-3H-indolium hydroxide inner salt (FIG. 3, 2b)

[0128] 28.7 mmol (5.72 g) 2-methyl-3-spiro-1'-cyclohexan-3H-indole and 32 mmol (4.36 g) butanesultone are heated to 120° C. for 4 h. After cooling the mixture is triturated with 50 ml ether, washed 2 x with ethylacetate, again with ether and finally dried in vacuum.

[0129] Yield: 7.93 g (82%)

[0130] 2-(4-acetanilino-1,3-butadienyl)-3-spiro-1'-cyclohexan-1-(4-sulfobutyl)-3H-indolium hydroxide inner salt (FIG. 3, 2c)

[0131] 4 mmol (1.34 g) 1-(4-sulfobutyl)-2-methyl-3-spiro-1'-cyclohexan-3H-indolium hydroxide inner salt and 5 mmol malondianil (prepared from 5 mmol (1.62 g) malondianilhydroperchlorate and 5 mmol (490 mg) potassiumacetate in abs. ethanol, diluted with diethylether and evaporation of the filtrate) in 5 ml acetic anhydride and 5 ml glacial acetic acid are heated at 80° C. for 3 h. The solvent is evaporated in vacuum and the residue triturated with ether,

filtered off and washed with ether until the filtrate is colourless. The product is dried in vacuum and used for the next steps without further purification. Yield: 2.00 g (98% d.Th.)

[0132] 1-(4-acetoxybutyl)-2-methyl-3-spiro-1'-cyclohexan-3H-indolium iodide (FIG. 3, 2d)

[0133] 0.27 mol (49.8 g) 2-methyl-3-spiro-1'-cyclohexan-3H-indole and 0.27 mol (70.0 g) 4-iodobutylacetate are heated for 5 h at 110° C. After cooling the residue is dissolved in small amount of methylenchloride and stirred with 500 ml of ether. After stirring for a while the product crystallises, it is sucked off, washed with ether and dried in vacuum.

[0134] Yield: 104.9 g (87%)

[0135] Analogous to the preparation of 1-(4-acetoxybutyl)-2-methyl-3-spiro-1'-cyclohexan-3H-indolium iodide (2d) the according derivatives of 2-methylbenzoxazole, 2-methylbenzothiazole, 4-methylpyridine and 4-methylchinoline are accessible.

[0136] 1-(4-acetoxybutyl)-2-methyl benzoxazolium iodide (2e)

[0137] 10 mmol (1.33 g) 2-methylbenzoxazole and 10 mmol (2.42 g) 4-iodobutylacetate are heated for 6 h at 130° C. and after cooling mixed with 10 ml of ethylacetate. The crystals obtained are washed with ether and dried in vacuum.

[0138] Yield: 2.17 g (58%)

[0139] 1-(4-acetoxybutyl)-2-methyl benzothiazolium iodide (2f)

[0140] 10 mmol (1.49 g) 2-methyl-benzothiazole and 10 mmol (2.42 g) 4-iodobutylacetate are heated for 4 h at 120° C. and after cooling mixed with ether sucked off, again washed with ether and dried in vacuum.

[0141] Yield: 3.01 g (77%)

[0142] 1-(4-acetoxybutyl)-4-methyl-pyridinium iodide (2g)

[0143] 10 mmol (0.93 g) y-picoline and 10 mmol (2.42 g) 4-iodobutylacetate are heated 4 h at 120° C. and after cooling repeatedly stirred with ether, decanted and finally dried in vacuum.

[0144] Yield: 2.35 g (70% d.Th.) viscous mass

[0145] 1-(4-acetoxybutyl)-4-methyl-chinolinium iodide (2h)

[0146] 10 mmol (1.43 g) lepidine and 10 mmol (2.42 g) 4-iodobutylacetate are heated 4 h at 120° C. and after cooling repeatedly stirred with ether, decanted and finally dried in vacuum.

[0147] Yield: 3.13 g (81% d.Th.) viscous mass

[0148] Dye S4

[0149] 1-(4-hydroxybutyl)-2-[5-(1,3-dihydro-1-(4-sulfobutyl)-3-spiro-1'-cyclohexan-2H-indole-2-ylidene)-penta-1,3-dienyl]-benzoxazolium hydroxide inner salt (FIG. 4, 2i)

[0150] 0.20 mmol (101 mg) 2-(4-acetanilino-1,3-butadienyl)-3-spiro-1'-cyclohexan-1-(4-sulfobutyl)-3H-indolium hydroxide inner salt, 0.21 mmol (79 mg) 1-(4-acetoxybutyl)-benzoxazolium iodide and 0.21 mmol (29.3 µl) triethylamine are refluxed in 1 ml abs. ethanol for 30 min. After

cooling 5 ml ether is added and decanted. The residue is dissolved in a mixture of 1 ml conc. HCl and 9 ml methanol and left overnight at 4° C. The solvent is evaporated in vacuum and the residue is purified by chromatography on silicagel RP-18 and acetonitrile-water (1:1).

## EXAMPLE 5:

[0151] Synthesis of dye S5

[0152] 1-(4-hydroxybutyl)-2-[5-(1,3-dihydro-1-(4-sulfo-butyl)-3-spiro-1'-cyclohexan-2H-indole-2-yliden)-penta-1,3-dienyl]-benzothiazolium hydroxide inner salt (FIG. 4, 2j)

[0153] Analogously to the procedure cited above 0.21 mmol (82 mg) 1-(4-acetoxybutyl)-benzothiazolium iodide are reacted with the appropriate reagents.

## EXAMPLE 6:

[0154] Synthesis of dye S6

[0155] 1-(4-hydroxybutyl)-4-[5-(1,3-dihydro-1-(4-sulfo-butyl)-3-spiro-1'-cyclohexan-2H-indole-2-yliden)-penta-1,3-dienyl]-pyridinium hydroxide inner salt (FIG. 4, 2k)

[0156] Analogously to the procedure cited above 0.21 mol (71 mg) 1-(4-acetoxybutyl)-pyridinium iodide are reacted with the appropriate reagents.

## EXAMPLE 7:

[0157] Synthesis of dye S7

[0158] 1-(4-hydroxybutyl)-4-[5-(1,3-dihydro-1-(4-sulfo-butyl)-3-spiro-1'-cyclohexan-2H-indole-2-yliden)-penta-1,3-dienyl]-chinolinium hydroxide inner salt (FIG. 4, 2l)

[0159] Analogously to the procedure cited above 0.21 mmol (81 mg) 1-(4-acetoxybutyl)-chinolinium iodide are reacted with the appropriate reagents.

[0160] Due to the state of the art, the hydroxylsubstituted fluorophores, prepared according to 2i-2l may be converted to the corresponding phosphoramidites with 2-cyanoethyl-N,N-diisopropyl-chlorophosphoramidite and coupled with DNA.

[0161] Preparation of dyes of the asymmetric 3-spiro-1'-4-tetrahydropyrene-indole type

## EXAMPLE 8:

[0162] Synthesis of dye S8

[0163] 2-methyl-5-sulfo-3-spiro-4'-tetrahydropyrene-3H-indole (FIG. 5, 3a)

[0164] 40 mmol (7.53 g) phenylhydrazine-4-sulfonic acid and 44 mmol 4-acetyltetrahydropyrene in 40 ml glacial acetic acid were refluxed 48 h under nitrogen. Afterwards the precipitate is filtered and the acetic acid is evaporated. The residue is dissolved in a small amount of water and 4 ml of 10M NaOH is added. After evaporation to dryness the residue is extracted with 2-propanole. The extract is evaporated and the residue dried in vacuum.

[0165] Yield: 6.94 g (57%)

[0166] 1-ethyl-2-methyl-5-sulfo-3-spiro-4'-tetrahydropyrene-3H-indolium hydroxide inner salt (FIG. 5, 3b)

[0167] 5.2 mmol (1.58 g) 2-methyl-5-sulfo-3-spiro-4'-tetrahydropyrene-3H-indole and 5.72 mmol (882 mg) diethylsulfate are heated for 5 h at 110° C. in 3 ml 1,2-

dichlorobenzene. After cooling the mixture is diluted with acetone and filtered. The resulting cake is thoroughly washed with acetone and dried in vacuum.

[0168] Yield: 1.089 g (67%)

[0169] 1-(5-carboxypentyl)-2-methyl-5-sulfo-3-spiro-4'-tetrahydropyrene-3H-indolium inner salt (FIG. 5, 3c)

[0170] 10 mmol (3.03 g) 2-ethyl-5-sulfo-3-spiro-4'-tetrahydropyrene-3H-indole and 15 mmol (22.93 g) 6-bromohexanoic acid are heated in 10 ml of 1,2-dichlorobenzene for 24 h at 120° C. After cooling the solution is diluted with acetone and decanted. The precipitate is triturated with hot 2-propanole, cooled, filtered, washed with 2-propanole and acetone and finally dried in vacuum.

[0171] Yield 3.07 g (78%)

[0172] 2-(4-acetanilino-1,3-butadienyl)-1-ethyl-3-spiro-4'-tetrahydropyrene-5-sulfo-3H-indolium hydroxide inner salt (FIG. 5, 3d)

[0173] 3 mmol (0.93 g) 1-ethyl-2-methyl-3-spiro-4'-tetrahydropyrene-5-sulfo-3H-indolium toluenesulfonate and 3.6 mmol malondianile (prepared from 3.6 mmol (prepared from 3.6 mmol (1.165 g) malondianilehydroperchlorate and 3.6 mmol (353 mg) potassium acetate in abs. ethanol, diluted with diethylether and evaporation of the filtrate) in 15 ml of a mixture of glacial acetic acid and acetic anhydride (1:1 v/v) and heated for 3 h at 80° C. The solvent is evaporated in vacuum and the residue triturated with ether, filtered off and washed with ether until the filtrate is colourless. The product is dried in vacuum.

[0174] Yield: 1.33 g (92%)

[0175] dye S8

[0176] 1-(5-carboxypentyl)-2-[5-(1,3-dihydro-1-ethyl)-3-spiro-4'-tetrahydropyrene-5-sulfo-2H-indole-2-yliden)-penta-1,3-dienyl]-3-spiro-4'-tetrahydropyrene-5-sulfo-3H-indolium hydroxide inner salt potassium salt, (FIG. 5, 3e)

[0177] 2.5 mmol (1.20 g) 2-(4-acetanilino-1,3-butadienyl)-3-spiro-4'-tetrahydropyrene-1-ethyl-5-sulfo-3H-indolium hydroxide inner salt in a mixture of 5 ml pyridine and 5 ml acetic anhydride are treated with 2.7 mmol (1.07 g) 1-(5-carboxypentyl)-2-methyl-5-sulfo-3-spiro-4'-tetrahydropyran-3H-indolium inner salt and refluxed for 40 min. The solution is concentrated in vacuum to one third and the crude dye is precipitated with ether and decanted. The residue is dissolved in a small amount of 1M HCl and reprecipitated with a saturated solution of KCl. The precipitate is separated, dissolved in water and purified by reversed-phase chromatography on silicagel RP-18 with an acetonitrile-water gradient.

[0178] Preparation of dyes of the asymmetric 3-spiro-1'-4'-piperidyl-indole type

## EXAMPLE 9:

[0179] Syntheses of dye S9

[0180] Synthesis of 2-methyl-3-spiro-4'-(1'-carboxybenzyl)-piperidine-3H-indole (FIG. 6, 4a)

[0181] 4.31 g (16.5 mmol) of the piperidylketone and 1.62 g (1.48 ml, 15 mmol) phenylhydrazine are dissolved in 15 ml glacial acetic acid under inert gas atmosphere and refluxed for 3.5 h. The excess acetic acid is distilled in

vacuum on a rotary evaporator remaining a brown oil. The oil is taken up in 80 ml of water and extracted thrice with each 40 ml diethylether. The combined organic layers are washed once again with 40 ml of water and dried over magnesiumsulfate. The solvent is distilled off in vacuum on a rotary evaporator. The remaining brown oil is purified by column chromatography (dichloromethane/ethylacetate =2:1) The indole is obtained as reddish oil. Yield=3.23 g, 9.7 mmol,(64%).

[0182] Synthesis of 1-ethyl-2-methyl-3-spiro-4'-(1'-carboxybenzyl)-piperidine-3H-indolenium ethyl sulfate (**FIG. 6, 4b**)

[0183] Under inert gas atmosphere 2.06 g (6 mmol) of 2-methyl-3-spiro-4'-(1'-carboxybenzyl)-piperidine-3H-indole (**4a**) are dissolved in 3 ml toluene. 1.02 g (0.89 ml, 6.6 mmol) diethylsulfate are added through a syringe. The mixture is refluxed for 3.5 h, precipitating a dark purple oil, which turns solid on cooling. The supernatant liquid is decanted and the residue repeatedly washed with small portions of toluene. The solid is purified by column chromatography with the aid of a gradient system (1. ethylacetate, 2. acetone/glacial acetic acid=10:2). The quaternary indole is eluted twice with the solvent and the product is obtained as highly viscous oil (0.93 g, 1.9 mmol, 32%)

[0184] Synthesis of 2-(4-acetanilino-1,3-butadienyl)-1-ethyl-3-spiro-4'-(1'-carboxybenzyl)-piperidine-3H-indolenium ethyl sulfate (**FIG. 6, 4c**)

[0185] 0.93 g (1.9 mmol) of the quaternary indole (**4b**) and 2.4 mmol malondianil (prepared from 0.79 g (2.4 mmol) malondianil-hydroperchlorate and 0.23 g (2.4 mmol) potassium acetate in abs. ethanol, diluted with diethylether and evaporation of the filtrate) in 3 ml of acetic anhydride and 3 ml glacial acetic acid are heated for 3 h at 80° C. The solvent is removed under reduced pressure on a rotary evaporator. The solid residue is triturated with diethylether, filtered and washed with ether until the filtrate is colourless. The crude product (1.19 g, 1.8 mmol, 97%) is dried in vacuum and used in the next step without further purification.

[0186] Synthesis of 1-carboxypentyl-2-methyl-3-spiro-4'-(1'-carboxybenzyl)-piperidine-3H-indolium bromide (**FIG. 6, 4d**)

[0187] 2.06 g (6 mmol) 2-methyl-3-spiro-4'-(1'-carboxybenzyl)-piperidine-3H-indole (**4a**) are heated in 5 ml 6-bromohexanoic acid 3,5 h at 120° C. The supernatant is decanted from the reddish solid. Purification is carried out by column chromatography (water/acetic acid=10:2). The quaternary indole is obtained as reddish highly viscous oil (0.95 g, 1.8 mmol, 30%)

[0188] Dye S9

[0189] Synthesis of 1-(5-carboxypentyl)-2-[5-(1,3-dihydro-1-(1-ethyl)-3-spiro-4'-(1'-carboxybenzyl)-piperidine-2H-indole-2-yliden)-penta-1,3-dienyl]-3-spiro-4'-(1'-carboxybenzyl)-piperidin-3H-indolenium ethyl sulfate (**FIG. 6, 4e**)

[0190] 0.62 g (1.8 mmol) of the hemicyanine (**4c**) and 0.71 g (1.8 mmol) the indole (**4d**) in 3 ml acetic anhydride and 3 ml pyridine are refluxed for 30 min. The mixture is evaporated to dryness in vacuum on a rotary evaporator. The solid residue is taken up in a mixture of 10 ml conc. HCl and 100 ml methanol and left overnight at 4° C. The solvent is

distilled off at reduced pressure and the residue is purified by column chromatography (isopropanol:water, 2:1 (v/v)). The dye S9 is obtained as dark blue crystalline solid (0.36 g, 0.38 mmol, 21% ).

[0191] Synthesis of 1-(5-carboxypentyl)-2-[5-(1,3-dihydro-1-(1-ethyl)-3-spiro-4'-piperidine-2H-indole-2-yliden)-penta-1,3-dienyl]-3-spiro-4'-piperidine-3H-indolenium ethylsulfate (**FIG. 7, 4f**)

[0192] 0.36 g (0.38 mmol) of the dye (**4e**) is dissolved under an inert gas atmosphere in 5 ml of chloroform. With a syringe 0.09 g (0.07 ml, 0.46 mmol) of iodotrimethylsilane is added and the mixture is stirred at room temperature until the educt is not detectable again by thinlayer chromatography (isopropanol/water=10:2, v/v) . After that 2ml of methanol is added and the mixture is stirred for 5 min. at room temperature. The liquid components are distilled off in vacuum on a rotary evaporator and the solid residue is purified by column chromatography with the eluent isopropanol:water=2:1. Yield: 0.23 g, 0.3 mmol (80%)

[0193] The deprotected dye S9 (**4f**) can be converted into a polyethylen glycole-substituted dye (4 g) by melting together with glycidyl-PEG.

[0194] Absorption spectra of the dyes (S1-S3) prepared in examples 1-3 are shown in **FIG. 8**.

[0195] In order to covalently label target molecules the carbonyl moiety has to be converted into an activated ester (reactive group R<sub>x</sub>) . Due to state of the art, this can be carried out with N-hydroxysuccinimide in the presence of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) in aqueous solution or with dicyclohexylcarbodiimide (DCC) in organic solution (**FIG. 2, 1f**)

#### EXAMPLE 10

[0196] Coupling of the NHS activated pentamethine-spiro-cyanine (compound S2) to protein

[0197] 0,2 mg activated dye are added while shaking to 1 mg of bovine serum albumin in 1 ml 10 mM phosphate buffer, pH 8.0. After a reaction time of 60 min, the coupling is stopped by the addition of 40 µl glycine solution (10% m/v, in PBS-buffer, pH 7,4) and the excess of dye is removed through dialysis or gel filtration.

[0198] **FIG. 9** shows the emission spectrum of the dye-conjugate obtained with dye S2 (example 2) as described in example 10.

[0199] In comparison, **FIG. 10** shows the corresponding spectrum of a common conjugate created under identical conditions. A band in the absorption spectrum at about λ=600 nm due to dimer formation which leads to reduced fluorescence can clearly be seen.

[0200] The mobility of the fluorescent dye is reduced due to the influence of the bulky spiro-substituent, which leads to an increased photostability. **FIG. 11** displays the photostability of the compound S3 prepared according to example 3, determined in 10 mM phosphate buffer, pH 7.4, compared to a commercial state-of-the-art heptamethine dye. Irradiation was performed using a krypton-lamp and 20 mW/cm<sup>2</sup> at λ>400 nm.

[0201] By determination of the fluorescence of the S3-conjugate, according to example 10, in comparison to a

conjugate, created using a common cyanine-dye, the results displayed in FIG. 12 are obtained. The resulting fluorescence of the conjugates, created under identical conditions, was determined at different incubation times after separation of the excess of the dye. The longer the incubation time, the higher the label degree fluorophor:protein. The formation of a narrow maximum is observed with the conjugate according to the state of the art, the resulting fluorescence decreases more than 50% at higher labeling degrees due to self-quenching phenomenon. In contrast, conjugates with the fluorescent dye according to the invention (examples 2 and 10) show a substantially reduced tendency towards aggregation and thus exhibit higher fluorescence. The fluorescence of the conjugate with the invention based fluorophor (example 10) after an incubation time of 15 minutes is more than double as high as the one of the conjugate with a state-of-the-art fluorophor.

[0202] The spiro-substituent also causes a shielding of the central polymethine chain against the influence of the surroundings of the fluorophor. As known, many fluorophores exhibit a shift of their absorption-maximum towards longer wavelengths and a simultaneous increase of fluorescence with an increase of protein concentration. This is caused by the changed environment created by the protein and the reduced mobility of the fluorescent dye. This effect is minimized with the fluorescent dye according to the invention. As shown in FIG. 13 (normalized display), there only is a minimal increase of fluorescence in the presence of high concentrations of bovine serum albumin (BSA). The fluorescence of the commercial fluorophor, in contrast, is highly dependent on the environment. Latter effect is a very strong disadvantage when analyzing substances in blood serum.

[0203] In table 2 the characteristic data of selected fluorescent dyes according to the invention are shown.

TABLE 1

electrophilic group	nucleophilic group	type of chemical bondage formed
activated ester <sup>1)</sup>	amine/aniline	carboxamide
acylazide	amine/aniline	carboxamide
acylhalide	amine/aniline	carboxamide
acylhalogene	alkohole/phenole	ester
acylnitrile	alkohole/phenole	ester
acylnitrile	amine/aniline	carboxamide
aldehyde	amine/aniline	imine
aldehyde/ketone	hydrazine	hydrazone
aldehyde/ketone	hydroxylamine	oxime
alkylhalogene	amine/aniline	alkylamine
alkylhalogene	carbonic acids	ester
alkylhalogene	thiole	thio ether
alkylhalogene	alcohol/phenol	ether
alkylsulfonate	thiole	thio ether
alkylsulfonate	carbonic acids	ester
alkylsulfonate	alcohol/phenol	ester
anhydride	alcohol/phenol	ester
anhydride	amine/aniline	carboxamide
arylhalogene	thiole	thiophenole
arylhalogene	amine	arylamine
aziridine	thiole	thioester
boronate	glycole	boronatester
carbonic acids	amine/aniline	carboxamide
carbonic acids	alcohol	ester
carbonic acids	hydrazine	hydrazide
carbodiimide	carbonic acids	N-acyl-urea
diazoalkane	carbonic acids	ester
epoxide	thiole	thio ether
haloacetamide	thiole	thio ether

TABLE 1-continued

electrophilic group	nucleophilic group	type of chemical bondage formed
halotriazine	amine/aniline	aminotriazine
halotriazine	alcohol/phenol	triazinether
imidoester	amin/aniline	amidine
isocyanate	amine/aniline	urea
isocyanate	alcohol/phenol	urethane
isothiocyanate	amine/aniline	thiourea
maleimide	thiole	thio ether
phosphoramidite	alcohol	phosphite triester
silylhalogene	alcohol	silyl ether
sulfonic acid ester	amine/aniline	alkylamine
sulfonic acid ester	carbonic acids	ester
sulfonic acid ester	thiole	thio ether
sulfonic acid ester	alcohol	ether
sulfonic acid ester	amine/aniline	sulfonamide
sulfonylchloride	alcohol/phenol	sulfonic acid ester

<sup>1)</sup>activated ester with the general structure —CO-W, where in W is a suitable leaving group, e.g., nitro-, fluoro-, chloro-, cyano-, trifluoromethyl-, tosyl- etc.

[0204]

TABLE 2

characteristic data of selected fluorescent dyes according to the invention		
Compound	absorption maximum (nm)	emission-maximum (nm)
S1*	557	572
S2*	656	672
S3*	756	772
S4**	612	638
S5**	650	676
S6**	603	626
S7**	702	728
S8**	662	684
S9**	660	682


\*determined in phosphate buffer, pH 7.4

\*\*determined in ethanol

What is claimed is


1. A fluorescent dye having the general structure A—Z—A— in which

A is a headgroup having the general formula I




I

wherein X is






O, S, SO, SO<sub>2</sub> or




W<sup>-</sup>=counter ion, preferably selected from the group of halogenide, perchlorate or tosylate,

A' is selected from a headgroup having the general formulas II to V,




-continued


V



wherein Y is O, S, NR<sub>17</sub>,




and T is



II

O, S, SO, SO<sub>2</sub> or




and

III

Z denotes a group with the general formula VI or VII


VI



with n is an integer selected from the group consisting of 0, 1, 2, or 3 or

IV

VII



wherein at least one of the residues is selected from a group consisting of

- a) a chemical reactive group for covalent coupling to a target molecule or
- b) an ionic group for coupling to a target molecule by ionic interaction forces or
- c) a lipophilic group for coupling to a target molecule by adsorption.

and the other residues are independently selected from the group of H, alkyl (C<sub>1</sub>-C<sub>10</sub>), alkoxy (C<sub>1</sub>-C<sub>10</sub>), trifluoromethyl, halogen, sulfonic acid, sulfonate, phosphoric acid and phosphonate.

2. A fluorescent dye according to claim 1, wherein at least one of the residues R<sub>1</sub>, R<sub>4</sub>, R<sub>6</sub> to R<sub>10</sub>, R<sub>13</sub>, R<sub>15</sub> and R<sub>17</sub> to R<sub>21</sub> is a chemical reactive group.

3. A fluorescent dye according to claim 2, wherein at least one of the residues R<sub>6</sub> to R<sub>9</sub>, and R<sub>18</sub> to R<sub>21</sub> is a chemical reactive group.

4. A fluorescent dye according to at least one of the preceding claims,

wherein the reactive group is selected from the group of carbonic acids, activated esters, acylazides, acylhalogenides, acylnitriles, aldehydes, anhydrides, arylamides, alkylhalogenides, anilines, alkylsulfonates, arylhalogenides, thioles, azides, aziridines, borates, carbodiimides, diazoalkanes, epoxides, glyceroles, haloacetamines, halotriazines, hydrazines, hydroxylamines, imidoesters, isocyanates, isothiocyanates, maleimides, phosphoramidites, silylhalogene, sulfonic acid esters and sulfonylchlorides.

5. A fluorescent dye according to at least one of the preceding claims,

wherein the reactive group is selected from the group of succinimidylesters, maleimides, epoxides, hydrazines and phosphoramidites.

6. A fluorescent dye according to at least one of the preceding claims,

wherein the ionic residues are selected from the group of carbonic acids, sulfonic acids, sulfonates, phosphoric acids, phosphonates, phosphordiester, phosphortriesters, primary, secondary, tertiary and quarternary amines.

7. A fluorescent dye according to claim 6,

in which at least one of the residues R<sub>1</sub>, R<sub>4</sub>, R<sub>6</sub> to R<sub>10</sub>, R<sub>13</sub>, R<sub>15</sub> and R<sub>17</sub> to R<sub>21</sub> is an ionic group.

8. A fluorescent dye according to claim 7,

in which at least one of the residues R<sub>6</sub> to R<sub>9</sub>, and R<sub>18</sub> to R<sub>21</sub> is an ionic group.

9. A fluorescent dye according to at least one of the claims 6 to 8,

in which one residue of group A selected from residues R<sub>1</sub>, R<sub>6</sub> to R<sub>9</sub> and one residue of group A' selected from residues R<sub>10</sub>, R<sub>17</sub> to R<sub>21</sub>, are counter charged ionic residues.

10. A fluorescent dye according to claim 9,

in which both ionic residues are equally charged.

11. A fluorescent dye according to claim 10,

in which two anionic residues are complexed by a metal cation selected from the 1.-3. main and sub group of the elements.

12. A fluorescent dye according to at least one of the preceding claims,

in which at least one of the residues R<sub>1</sub> to R<sub>26</sub> is hydrophilic, selected from the group of carbonic acids, carbohydrates, peptides, nucleic acids, sulfonic acids, sulfonates, thioles, phosphoric acids, phosphonates, phosphordiester, phosphortriester, amines, halogenes, alcohols, polyols or polyether.

13. A fluorescent dye according to claim 12,

in which the carbonic acid is selected from the group of mono-, bi- and multifunctional aliphatic carbonic acids with a carbon chain of up to 10 atoms and

the carbohydrates selected from the group of monosaccharides, such as trioses, tetroses, pentoses, hexoses or heptoses, disaccharides like sucrose, lactose, trehalose or maltose, oligosaccharides like cyclodextrine or raffinose and polysaccharides like dextran or chitosane, including all O- and N-glycosidic modifications and combinations thereof.

14. A fluorescent dye according to at least one of the claims 12 or 13,

in which at least one of the residues R<sub>1</sub>, R<sub>4</sub>, R<sub>6</sub> to R<sub>10</sub>, R<sub>13</sub>, R<sub>15</sub> and R<sub>17</sub> to R<sub>21</sub> is a hydrophilic group.

15. A fluorescent dye according to claim 14,

in which at least one of the residues R<sub>6</sub> to R<sub>9</sub>, and R<sub>18</sub> to R<sub>21</sub> is a hydrophilic group.

16. A fluorescent dye according to at least one of the preceding claims,

in which at least one of the residues R<sub>1</sub> to R<sub>26</sub> is a lipophilic group selected from saturated or unsaturated alkyl chain (C<sub>8</sub>-C<sub>18</sub>) and/or fatty acid esters, each having 0, 1, 2 or 3 double bonds.

17. A fluorescent dye according to claim 16,

in which at least one of the residues R<sub>1</sub>, R<sub>4</sub>, R<sub>6</sub> to R<sub>10</sub>, R<sub>13</sub>, R<sub>15</sub> and R<sub>17</sub> to R<sub>21</sub> is a lipophilic group.

18. A fluorescent dye according to at least one of the claims 16 or 17,

in which at least one of the residues R<sub>6</sub> to R<sub>9</sub>, and R<sub>18</sub> to R<sub>21</sub> is a lipophilic group.

19. A fluorescent dye according to at least one of the preceding claims,


in which the counter ion W<sup>-</sup> is selected from the group of halogenide, tosylate and perchlorate.

20. A fluorescent dye, according to at least one of the preceding claims

in which each pair of aromatic ortho-substituted residues can be combined to form at least one adjacent carbo- or heterocyclic aromatic ring.


21. A fluorescent dye according to at least one of the preceding claims,

characterized thereby that the target molecule is a biomolecule selected from the group of antibodies, proteins, peptides, enzyme substrates, hormones lymphokines, metabolites, receptors, antigens, haptens, lecithines, toxins, carbohydrates, oligosaccharides, polysaccharides, nucleic acids, desoxy ribonucleic acids (DNA), derivatized DNA, ribonucleic acids (RNA), derivatized RNA(new!!), derivatized nucleic acids, DNA-fragments, RNA-fragments, drugs, virus particles, virus components, yeast, yeast components,



where  $M^+$  is a metal cation. X, T,  $W^-$  and n are defined according to claim 1.

25. A fluorescent dye according to at least one of the claims 1-23, characterized by a compound with the following formula IX



bacteria, bacterial components, blood cells, components of blood cells, biological cells, non cellular blood components and toxins

22. A fluorescent dye according to at least one of the preceding claims,

characterized thereby that the target molecule is a non biological molecule (compound) selected from the group of polymeres, polymer particles, glass particles, glass surfaces, plastic surfaces, plastic particles, polymer membranes, metals, electrical conductors and/or electrical semiconductors.


23. A fluorescent dye according to at least one of the preceding claims,

characterized thereby that the fluorescent dye is bound onto the surface or is incorporated into nano- or micro-particles of the basis of organic polymers and the use of said particles for analytical applications (methods).

24. A fluorescent dye according to at least one of the preceding claims 1-23, characterized by a compound with the following formula VIII

where  $M^+$  is a metal cation. X, T,  $W^-$  and n are defined according to claim 1.


26. A fluorescent dye according to at least one of the preceding claims 1-23, characterized by a compound with the following formula X



where  $M^+$  is a metal cation. X, T,  $W^-$  and n are defined according to claim 1.


$m=0, 1-17$

27. A fluorescent dye according to at least one of the preceding claims 1-23, characterized by a compound with the following formula XI




with  $M^+$  = metal cation, X, Y, T,  $W^-$ , m and n are defined according to claim 1.

**28.** A fluorescent dye according to at least one of the preceding claims **1-23**, characterized by a compound with the following formula XII




mit X, Y, T und n entsprechend Anspruch 1 definiert,  $p=1-8$ .

**29.** A fluorescent dye according to at least one of the preceding claims **1-23**, characterized by a compound with the following formula XIII




with X and n defined according to claim 1, q is equal to 1 to 8.

**30.** A fluorescent dye according to at least one of the preceding claims **1-23**, characterized by a compound with the following formula XIV




with  $\text{M}^+$ =metal cation and  $n$  defined according to claim 1,  
 NHS=N-hydroxy-succinimide,  $r$  and  $s$  independently  
 from each other equal to 1 to 8.

**31.** A fluorescent dye according to at least one of the  
 preceding claims **1-23**, characterized by a compound with  
 the following formula XV




with with  $\text{W}^-$  and  $n$  defined according to claim 1.

**32.** A fluorescent dye according to at least one of the  
 preceding claims **1-23**, characterized by a compound with  
 the following formula XVI




with  $\text{W}^-$  and  $n$  defined according to claim 1.

**33.** A fluorescent dye according to at least one of the  
 preceding claims **1-23**, characterized by a compound with  
 the following formula XVII




with  $W^-$  and  $n$  defined according to claim 1, NHS =N-hydroxy-succinimid, PEG=poly ethylen glycole and with  $t$  and  $u$  independently from each other equal to 1 to 8.

**34.** A fluorescent dye according to at least one of the preceding claims 1-23, characterized by a compound with the following formula XVIII




with  $W^-$  and  $n$  defined according to claim 1,  $R'$  equal trityl,  $R''$ =phosphoramidite,  $v$  and  $w$  independently equal to 0 to 6.

**35.** A fluorescent dye according to at least one of the preceding claims 1-23, characterized by a compound with the following formula XIX




with  $M^+$ =metal cation,  $X$ ,  $Y$ ,  $T$ ,  $W^-$  and  $n$  defined according to claim 1.

**36.** A fluorescent dye according to at least one of the preceding claims 1-23, characterized by a compound with the following formula XX



with  $M^+$ =metal cation,  $X$ ,  $Y$ ,  $T$ ,  $W^-$  and  $n$  defined according to claim 1,  $z$  equal to 0 to 6.

**37.** A fluorescent dye according to at least one of The preceding claims 1-23, characterized by a Compound with the following formula XXI



with  $X$ ,  $Y$ ,  $T$ ,  $W^-$  and  $n$  defined according to claim 1.

**38.** A method, where a compound according to claim 1 to 23 is used together with another fluorescent dye or another conjugate in a fluorescence resonance energy transfer system.

**39.** A method, where a compound according to claim 1 to 23 is used for labelling and/or detection of nucleic acids

**40.** A method according to claim 39, where the nucleic acid is detected as product of an amplification reaction.

**41.** A method, where a compound according to claims 1 to 23 is used as fluorescent dye for coupling to biological and non-biological target molecules, which are used in analytical methods.

\* \* \* \* \*

专利名称(译)	3-螺-cyanin荧光染料及其在生物测定中的应用		
公开(公告)号	<a href="#">US20030138791A1</a>	公开(公告)日	2003-07-24
申请号	US10/130547	申请日	2001-09-03
[标]申请(专利权)人(译)	HAALCK LUTZ GEDIG ERK HOELPERT HANS JUERGEN PODSADLOWSKI VIOLA		
申请(专利权)人(译)	HAALCK LUTZ GEDIG ERK PODSADLOWSKI VIOLA		
当前申请(专利权)人(译)	HAALCK LUTZ GEDIG ERK PODSADLOWSKI VIOLA		
[标]发明人	HAALCK LUTZ GEDIG ERK HOELPERT HANS JUERGEN PODSADLOWSKI VIOLA		
发明人	HAALCK, LUTZ GEDIG, ERK HOELPERT, HANS-JUERGEN PODSADLOWSKI, VIOLA		
IPC分类号	C07D491/107 C09B23/02 G01N33/533 C12Q1/68 C07D417/02 C07D43/02 C07D413/02		
CPC分类号	G01N33/533 C09B23/02		
优先权	10046215 2000-09-19 DE		
其他公开文献	US7855293		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

本发明涉及基于对称或不对称3-螺萘菁的空间屏蔽的稳定荧光染料及其在分析方法中作为标记分子的用途。根据本发明的荧光染料的特征在于由于插入庞大的螺-取代基而大大降低了聚集倾向，因此能够达到与靶分子的高度偶联，而不会由于自猝灭作用而显著损失荧光。因此，它们具有高量子产率，因此非常适合作为荧光染料，特别是作为NIR荧光染料，用于偶联或结合分析方法中使用的生物和非生物靶分子。分析方法包括使用荧光光学方法检测生物和非生物分子的所有方法。

