



US 20110014635A1

(19) **United States**

(12) **Patent Application Publication**
SUZUKI et al.

(10) **Pub. No.: US 2011/0014635 A1**

(43) **Pub. Date: Jan. 20, 2011**

(54) **MARKER PEPTIDE FOR ALZHEIMER'S DISEASE**

(76) Inventors: **Toshiharu SUZUKI**, Chiba-shi (JP); **Yoichi Araki**, Sapporo-Shi (JP); **Naomi Miyagi**, Sapporo-Shi (JP); **Haruyasu Yamaguchi**, Maebashi-Shi (JP); **Masaki Nishimura**, Kusatsu-Shi (JP); **Kazuo Yamamoto**, Tokyo (JP)

Correspondence Address:
BIRCH STEWART KOLASCH & BIRCH
PO BOX 747
FALLS CHURCH, VA 22040-0747 (US)

(21) Appl. No.: **12/859,950**

(22) Filed: **Aug. 20, 2010**

Related U.S. Application Data

(62) Division of application No. 10/577,008, filed on Aug. 7, 2006, now Pat. No. 7,807,777, filed as application No. PCT/JP2004/016209 on Nov. 1, 2004.

(30) **Foreign Application Priority Data**

Nov. 5, 2003 (JP) 2003-375363

Publication Classification

(51) **Int. Cl.**
G01N 33/53 (2006.01)
C07K 14/00 (2006.01)
C07K 16/18 (2006.01)

(52) **U.S. Cl. 435/7.21; 530/350; 436/86; 530/387.9**

(57) **ABSTRACT**

To provide a peptide obtainable by cleaving an N-terminal region and a C-terminal region of Alcadein α , Alcadein β , or Alcadein γ ; and capable of being a diagnostic marker for Alzheimer's disease. It is possible to detect Alzheimer's disease at an early stage without burdening subjects to be tested by using the peptide as a diagnostic marker.




Fig. 2




Fig. 3





Fig. 4

$A\beta$



$A\beta$



Merge




Fig. 5




Fig. 6




Fig. 7




Fig. 8






Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 19




Fig. 20




Fig. 21






Fig. 22



MARKER PEPTIDE FOR ALZHEIMER'S DISEASE

[0001] This application is a Divisional under 35 U.S.C. §120, of co-pending application Ser. No. 10/577,008 filed on Apr. 25, 2006, which is a national phase of PCT Application No. PCT/JP2004/016209 filed on Nov. 1, 2004, and this application claims priority of Application No. 2003-375363 filed in Japan on Nov. 5, 2003, respectively, under 35 U.S.C. §119; the entire contents of all of the aforementioned applications are hereby incorporated by reference.

TECHNICAL FIELD

[0002] The present invention relates to peptides which can be used as diagnostic markers for Alzheimer's disease, methods for diagnosing Alzheimer's disease using the peptides, methods for collecting data for diagnosing Alzheimer's disease by using the peptides, methods for screening for therapeutic agents for Alzheimer's disease by using the peptides, antibodies to the peptides, and diagnostic reagents including the antibodies.

BACKGROUND ART

[0003] Alzheimer's disease is currently diagnosed by carrying out an interview with a specialized physician and evaluating the degree of brain atrophy using MRI or the like. However, it is difficult to obtain an objective and correct diagnostic conclusion by interview only. Furthermore, it is impossible to identify a so-called pre-patient, before the onset of symptoms. Additionally, apparatuses such as MRI apparatuses are expensive and consequently can be used only in large special hospitals.

[0004] Under such circumstances, biochemical diagnosis using a marker is adopted as a simple and objective method. Among main markers for Alzheimer's disease, intracellular tau protein and β -amyloid (hereinafter referred to as "A β ") are known at present (Non-Patent Document 1 and Non-Patent Document 2).

[0005] Tau protein is a component constituting microtubules in nerve cells and is leaked out from the cells when the nerve cells are degenerated during an Alzheimer's disease process. As a result, tau protein is detected in cerebrospinal fluid, and is a useful marker. However, tau protein cannot be detected until the condition of the disease progresses. Furthermore, since the leakage amount is small, tau protein is hardly detected in body fluid (for example, blood) other than the cerebrospinal fluid.

[0006] A β is a causative substance of Alzheimer's disease. Therefore, A β can be a most effective marker provided that a quantitative change (an increase in the production) or a qualitative change (an increase in the ratio of highly aggregative A β) can be precisely measured. However, since A β shows aggregative nature, the amount of A β detected in a patient's cerebrospinal fluid is rather lower than that of healthy subjects.

[0007] Non-Patent Document 1: "Decreased beta-amyloid 1-42 and increased tau levels in cerebrospinal fluid of patients with Alzheimer disease"; Sunderland, T., Linker, G., Mirza, N., Putnam, K. T., Friedman, D. L., Kimmel, L. H., Bergeson, J., Manetti, G. J., Zimmermann, M., Tang, B., Bartko, J. J., and Cohen, R. M. JAMA 2003, 289, 2094-2103.

[0008] Non-Patent Document 2: "Cerebrospinal fluid biomarkers for disease stage and intensity in cognitively impaired patients"; Wahlund, L. O., and Blennow, K. Neurosci. Lett. 2003, 339, 99-102.

DISCLOSURE OF INVENTION

Problems to be Solved by the Invention

[0009] As mentioned above, it is difficult to detect Alzheimer's disease at an early stage by a diagnostic method using currently known markers. Furthermore, it is also difficult to diagnose Alzheimer's disease by a method such as blood examination, which is less of a burden on patients.

[0010] The present invention has been accomplished under the above-mentioned technical background, and it is an object of the present invention to provide a method for easily and accurately diagnosing Alzheimer's disease.

Means for Solving the Problem

[0011] The present inventors have performed intensive studies in order to solve the above-mentioned problems and, as a result, have found that Alcadein, which is a protein, is cleaved by an enzyme that cleaves a precursor protein of A β (hereinafter referred to as "APP") and then is extracellularly secreted in the same manner as in A β . It has been already reported that Alcadein forms a triple complex with X11L and APP and the formation of the complex suppresses the production of A β (Araki, Y. et al., J. Biol. Chem. 2003, 278, 49448-49458 and Japanese Unexamined Patent Application Publication No. 2003-164298). However, the fact that Alcadein is cleaved by the enzyme which cleaves APP and is extracellularly secreted in the same manner as in A β is a completely new finding. Additionally, the present inventors have found that the amount of a high-molecular-weight peptide generated from Alcadein by its consecutive cleavages is increased under the conditions that the amount of a high-molecular-weight A β , which is highly aggregative and highly neurotoxic, is increased.

[0012] The present invention has been accomplished on the basis of the foregoing findings.

[0013] Accordingly, the present invention provides the following aspects (1) to (15):

[0014] (1) a peptide obtainable by cleaving an N-terminal region and a C-terminal region of Alcadein α , Alcadein β , or Alcadein γ ; and capable of being a diagnostic marker for Alzheimer's disease (hereinafter the peptide is simply referred to as "peptide of the present invention");

[0015] (2) the peptide according to the aspect (1), wherein the N-terminal region to be cleaved is a portion of an extracellular domain at the N-terminal;

[0016] (3) the peptide according to the aspect (1) or (2), wherein the C-terminal region is cleaved by presenilin);

[0017] (4) the peptide according to the aspect (1), wherein the peptide is obtained by cleaving an N-terminal region and a C-terminal region of Alcadein α ; and the cleavage site of the N-terminal region is between amino acids 815 and 816, amino acids 820 and 821, or amino acids 838 and 839 of the amino acid sequence represented by SEQ ID NO: 1;

[0018] (5) the peptide according to the aspect (1) or (2), wherein the peptide is obtained by cleaving an N-terminal region and a C-terminal region of Alcadein α ; and the cleavage site of the C-terminal region is between amino acids 842

and 843, amino acids 843 and 844, or amino acids 851 and 852 of the amino acid sequence represented by SEQ ID NO: 1;

[0019] (6) the peptide according to the aspect (1), consisting of an amino acid sequence represented by any one of SEQ ID NOS: 4 to 12;

[0020] (7) a method for collecting data for diagnosing Alzheimer's disease, including a process of detecting or quantitatively determining the peptide according to any one of the aspects (1) to (6) in body fluid or tissues taken from an animal;

[0021] (8) the method for collecting data for diagnosing Alzheimer's disease according to the aspect (7), wherein the body fluid is blood or cerebrospinal fluid;

[0022] (9) the method for collecting data for diagnosing Alzheimer's disease according to aspect (7) or (8), wherein a ratio of a high-molecular-weight peptide in the detected or quantitatively determined peptide is used as an indicator for diagnosing Alzheimer's disease;

[0023] (10) a method for diagnosing Alzheimer's disease, including a process of detecting or quantitatively determining the peptide according to any one of the aspects (1) to (6) in body fluid or tissues taken from an animal;

[0024] (11) the method for diagnosing Alzheimer's disease according to the aspect (10), wherein the body fluid is blood or cerebrospinal fluid;

[0025] (12) the method for diagnosing Alzheimer's disease according to the aspect (10) or (11), wherein a ratio of a high-molecular-weight peptide in the detected or quantitatively determined peptide is used as an indicator for diagnosing Alzheimer's disease;

[0026] (13) a method for screening a therapeutic agent for Alzheimer's disease by contacting cells secreting the peptide according to any one of the aspects (1) to (6) with an agent to be screened and determining a change in the secreted amount of the peptide or a change in the molecular species of the secreted peptide;

[0027] (14) an antibody against the peptide according to any one of the aspects (1) to (6); and

[0028] (15) a diagnostic reagent for Alzheimer's disease, including the antibody according to the aspect (14).

ADVANTAGEOUS EFFECT OF THE INVENTION

[0029] By utilizing the peptide of the present invention, Alzheimer's disease can be detected before clinical symptoms or at an early stage by a simple method which does not put a burden on subjects to be tested.

BRIEF DESCRIPTION OF THE DRAWINGS

[0030] FIG. 1 is a photograph showing the results of Western blotting of proteins separated by density-gradient centrifugation.

[0031] FIG. 2 is a photograph showing the results of Western blotting of proteins recovered by immunoprecipitation.

[0032] FIG. 3 is photographs of immunostained brain sections from an Alzheimer's disease patient (Alc α and APP were detected).

[0033] FIG. 4 is photographs of immunostained brain sections from an Alzheimer's disease patient (Alc α and A β were detected).

[0034] FIG. 5 is a diagram schematically illustrating a process of obtaining A β from APP.

[0035] FIG. 6 is a diagram showing the results of Western blotting of cell lysates using an anti-APP antibody.

[0036] FIG. 7 is diagrams showing the results of Western blotting of cell lysates using an anti-Alc antibody and media using an anti-FLAG antibody.

[0037] FIG. 8 is diagrams showing the results of Western blotting of membrane fractions using an anti-Alc antibody.

[0038] FIG. 9 is a diagram illustrating a structure of Alc α Δ E.

[0039] FIG. 10 is a diagram showing the results of Western blotting of cell lysates and a medium using an anti-FLAG antibody.

[0040] FIG. 11 is diagrams showing the results of Western blotting of cell lysates expressing APP or Alc α and BACE1.

[0041] FIG. 12 is a diagram schematically illustrating a process of preparing β -Alc α (a peptide obtained by cleaving an N-terminal region and a C-terminal region of Alc α) including a FLAG sequence.

[0042] FIG. 13 is diagrams showing the results of Western blotting of an extract solution of cells expressing Alc α Δ and immunoprecipitate of a medium when an anti-FLAG antibody is used.

[0043] FIG. 14 is a diagram showing the result of mass spectrometry of (3-Alc α secreted from cells expressing Alc α A.

[0044] FIG. 15 (SEQ ID NO: 1) is a diagram showing a secondary cleavage site of Alc α determined by the MALDI-TOF/MS method.

[0045] FIG. 16 (SEQ ID NO: 13) is a diagram showing a cleavage site of human APP695.

[0046] FIG. 17 is a diagram showing the results of electrophoresis of a C-terminal region obtained by a primary cleavage of Alc α 1.

[0047] FIG. 18 (SEQ ID NO: 1) is a diagram showing cleavage sites of Alc α .

[0048] FIG. 19 is a diagram showing the results of Western blotting of cells expressing various PS1 variants and Alc α Δ E.

[0049] FIG. 20 is a diagram showing the results of Western blotting of cells expressing various PS1 variants (including L166P variant-type PS1) and Alc α Δ E.

[0050] FIG. 21 is diagrams showing the results of mass spectrometry of β -Alc obtained from wild-type PS1 or L166P variant-type PS1.

[0051] FIG. 22 (SEQ ID NO: 1) is a diagram showing cleavage sites of β -Alc.

[0052] FIG. 23 (SEQ ID NO: 1) is a diagram schematically showing molecular species of β -Alc.

BEST MODE FOR CARRYING OUT THE INVENTION

[0053] The present invention will now be described in detail.

[0054] The peptide of the present invention is obtained by cleaving at N-terminal and C-terminal regions of Alcadin α , Alcadin β , or Alcadin γ . The peptide generated by these cleavages can be a diagnostic marker for Alzheimer's disease.

[0055] There are three types of Alcadin: Alcadin α (hereinafter referred to as "Alc α "), Alcadin β (hereinafter referred to as "Alc β "), and Alcadin γ (hereinafter referred to as "Alc γ "). Alc α is a protein including an amino acid sequence which is the same or substantially the same as the amino acid sequence represented by SEQ ID NO: 1, Alc β is a protein including an amino acid sequence which is the same or substantially the same as the amino acid sequence represented by

SEQ ID NO: 2, and Alc γ is a protein including an amino acid sequence which is the same or substantially the same as the amino acid sequence represented by SEQ ID NO: 3.

[0056] Alc α , Alc β , and Alc γ may be proteins derived from any kind of cell (for example, hepatocytes, splenocytes, neurons, glia cells, pancreatic β -cells, myelocytes, mesangial cells, Langerhan's cells, epidermal cells, epithelial cells, goblet cells, endothelial cells, smooth muscle cells, fibroblasts, fibrous cells, muscle cells, fat cells, immune cells (e.g., macrophages, T-cells, B-cells, natural killer cells, mast cells, neutrophils, basophils, eosinophils, and monocytes), megakaryocytes, synovial cells, chondrocytes, osteocytes, osteoblasts, osteoclasts, mammary cells, hepatocytes, interstitial cells, progenitor cells of these cells, stem cells, and cancer cells) of human and warm-blooded animals (for example, guinea pig, rat, mouse, chicken, rabbit, swine, sheep, bovine, and monkey) or all tissues in which such cells are present, such as brain, various parts of brain (e.g., olfactory bulb, amygdaloid nucleus, cerebral basal nucleus, hippocampus, thalamus, hypothalamus, cerebral cortex, medulla oblongata, and cerebellum), spinal cord, pituitary gland, stomach, pancreas, kidney, liver, gonad, thyroid, gall-bladder, bone marrow, adrenal gland, skin, muscle, lung, gastrointestinal tract (e.g., large intestine and small intestine), blood vessels, heart, thymus, spleen, submandibular gland, peripheral blood, prostate, testis, ovary, placenta, uterus, bone, joint, and skeletal muscle, and furthermore, may be synthetic proteins as well.

[0057] Examples of the amino acid sequence which is substantially the same as the amino acid sequence represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3 include amino acid sequences which contain not less than about 50%, preferably not less than about 60%, more preferably not less than about 70%, still more preferably not less than about 80%, furthermore preferably not less than about 90%, most preferably not less than about 95% identity to the amino acid sequence represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3, respectively. Preferable examples of the protein which includes an amino acid sequence substantially the same as the amino acid sequence represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3 are proteins having the amino acid sequence substantially the same as the amino acid sequence represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3 and having an activity of substantially the same quality as that of the protein including the amino acid sequence represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3.

[0058] An example of the activity which has substantially the same quality as that of a protein is an activity that binds with the PI domain of X11L. The term substantially the same quality means the natures are equivalent (for example, physiologically or pharmacologically). Therefore, it is preferable that the above-mentioned activities are equivalent (e.g., about 0.01 to 100 times, preferably about 0.1 to 10 times, more preferably about 0.5 to 2 times), but it is allowable that the degrees of the activities and the quantitative elements such as molecular weight of the proteins are at different levels.

[0059] Alc α , Alc β , and Alc γ of the present invention each include, for example, so-called muteins such as proteins including (1) an amino acid sequence wherein one or more amino acids (preferably about 1 to 30, more preferably about 1 to 10, and further preferably a few (1 to 5) amino acids) are deleted from the amino acid sequence represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3, (2) an amino acid sequence wherein one or more amino acids (preferably about

1 to 30, more preferably about 1 to 10, and further preferably a few (1 to 5) amino acids) are added to the amino acid sequence represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3, (3) an amino acid sequence wherein one or more amino acids (preferably about 1 to 30, more preferably about 1 to 10, and further preferably a few (1 to 5) amino acids) are inserted into the amino acid sequence represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3, (4) an amino acid sequence wherein one or more amino acids (preferably about 1 to 30, more preferably about 1 to 10, and further preferably a few (1 to 5) amino acids) in the amino acid sequence represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3 are substituted with other amino acids, or (5) a combination thereof. When an amino acid sequence has the above-mentioned insertion, deletion, or substitution, the site of the insertion, deletion, or substitution is not limited as long as the activity is maintained. Specifically, as regards Alc α , Alc α 1 having the amino acid sequence represented by SEQ ID NO: 1 and Alc α 2 having an amino acid sequence wherein 10 amino acids are inserted between amino acids 71 and 72 of the amino acid sequence represented by SEQ ID NO: 1 are known.

[0060] The site where an N-terminal region is cleaved is not limited as long as the obtained peptide can be a diagnostic marker for Alzheimer's disease, but it is preferable that the site is within an extracellular domain at the N-terminal side. Generally, such a site is, in the case of Alc α , between amino acids 815 and 816, amino acids 820 and 821, or amino acids 838 and 839 of the amino acid sequence represented by SEQ ID NO: 1 or in the vicinities thereof; in the case of Alc β , between amino acid 825 and 826 of the amino acid sequence represented by SEQ ID NO: 2 or in the vicinity thereof (SEQ ID NO: 18); and in the case of Alc γ , between amino acids 804 and 805 of the amino acid sequence represented by SEQ ID NO: 3 or in the vicinity thereof.

[0061] Furthermore, Alc α is cleaved at another site at the N-terminal side too in the same manner as in APP: Alc α is also cleaved extracellularly by BACE which cleaves APP at the (3-site. Generally, such a site is between amino acid 708 and 709 of the amino acid sequence represented by SEQ ID NO: 1 or in the vicinity thereof.

[0062] The site where a C-terminal region is cleaved is not limited as long as the obtained peptide can be a diagnostic marker for Alzheimer's disease, but it is preferable that the site is cleaved by presenilin. Generally, such a site is, in the case of Alc α , between amino acids 842 and 843, amino acids 843 and 844, or amino acids 851 and 852 of the amino acid sequence represented by SEQ ID NO: 1 or in the vicinities thereof; in the case of Alc β , between amino acids 875 and 876 of the amino acid sequence represented by SEQ ID NO: 2 or in the vicinity thereof (SEQ ID NO: 18); and in the case of Alc γ , between amino acids 847 and 848 of the amino acid sequence represented by SEQ ID NO: 3 or in the vicinity thereof. Here, the term "vicinity" means generally a range within 10 amino acids, preferably a range within 5 amino acids from the cleavage site. Specific examples of the peptide of the present invention include a peptide consisting of an amino acid sequence represented by any one of SEQ ID NOS: 4 to 12.

[0063] It is thought that the peptide of the present invention can be used as a diagnostic marker for Alzheimer's disease because of the following reasons:

(1) the peptide of the present invention is obtained from Alc α that forms a triple complex with APP and X11L

(Araki, Y. et al., J. Biol. Chem. 2003, 278, 49448-49458 and Japanese Unexamined Patent Application Publication No. 2003-164298) and is distributed in the brain of an Alzheimer's disease patient in the same manner as in APP (Examples 2 and 3);

(2) Alcadin is cleaved by BACE as in APP (Example 8);

(3) the peptide of the present invention is obtained by the cleavage by presenilin as in A β (Examples 4, 6, and 7) and further secreted extracellularly as in A β (Example 7), and when the molecular species of A β is pathologically changed, the molecular species of the peptide of the present invention is also similarly changed (Example 11). On the basis of these facts, it is thought that the generation amount of A β can be predicted from the generation amount of the peptide of the present invention and that the qualitative change of A β can be predicted from the qualitative change of the peptide of the present invention; and

(4) A β cannot be used as a quantitative diagnostic marker for Alzheimer's disease because of its aggregative ability. A β has an α -helix structure at the N-terminal side and a 13-sheet structure at the C-terminal side, and a sequence composed of the 26th to the 29th amino acids forms a β -turn structure at the central portion. Consequently, an antiparallel β -sheet structure is formed by the N-terminal side and the C-terminal side. It is understood that this causes the aggregation of A β ("Oligomerization and fibril assembly of the amyloid β -protein" by Roher, A. E., et al., Biochem. Biophys. Acta 2000, 1502, 31-43). On the other hand, though the peptide of the present invention has an α -helix structure at the N-terminal side and a β -sheet structure at the C-terminal side as in the basic structure of A β , the peptide does not have a sequence to form a β -turn structure. Therefore, it is thought that since it is predicted that the α -helix structure is not converted to a β -sheet structure, the peptide does not have aggregative ability.

[0064] By using the peptide of the present invention as a diagnostic marker for Alzheimer's disease, Alzheimer's disease can be diagnosed and data for the diagnosis can be collected. Specifically, the diagnosis and the collection of data can be performed by detecting or quantitatively determining the peptide of the present invention in body fluid or tissues taken from animals.

[0065] The peptide of the present invention is available in various molecular weights. When a large amount of high-molecular-weight peptide is contained in the peptide of the present invention to be detected or quantitatively determined, it suggests a high possibility of Alzheimer's disease or its pre-stage. This is based on the fact that the amount of the high-molecular-weight peptide of the present invention is increased under the conditions that the amount of a high-molecular-weight A β (A β 42), which is highly aggregative and highly toxic, is increased (Examples 11 and 12). Therefore, Alzheimer's disease can be diagnosed by using the ratio of the amount of high-molecular-weight peptide to the total amount of the peptide of the present invention as an indicator, in addition to the determination of whether a certain amount of the peptide of the present invention is present in body fluid or the like. Here, the term "high-molecular-weight peptide" means a peptide which is obtained when the cleavage site of an N-terminal region is closer to the N-terminal end, or the cleavage site of a C-terminal region is closer to the C-terminal end, or a combination of both. For example, the high-molecular-weight peptide is defined as a molecular species which is obtained when the cleavage site is shifted to the N-terminal

end, the C-terminal end, or both ends from the site of each of the β -Alc molecular species shown in Table 1 described below. Thus, the high-molecular-weight peptide is not specifically defined by its molecular weight. When a primary cleavage site is ζ 1 and a secondary cleavage site is γ 3 as shown in Table 1, it is predicted that a peptide composed of 36 amino acids and having a molecular weight of about 4000 is obtained. If β -Alc has a molecular weight higher than that of this peptide by the shift of the cleavage site to the N-terminal end or the C-terminal end, the β -Alc is categorized as the above-mentioned "high-molecular-weight peptide". When a primary cleavage site is ζ 3 and a secondary cleavage site is γ 1 as shown in Table 1, it is predicted that a peptide composed of 4 amino acids and having a molecular weight of 500 to 600 is obtained. If β -Alc has a molecular weight higher than that of this peptide by the shift of the cleavage site to the N-terminal end or the C-terminal end, the β -Alc is categorized as the above-mentioned "high-molecular-weight peptide" even if the molecular weight is about 1000.

[0066] Examples of the animal from which body fluid or the like is taken include not only human but also warm-blooded animals other than human such as guinea pig, rat, mouse, chicken, rabbit, swine, sheep, bovine, and monkey.

[0067] Examples of the body fluid and tissues include blood, plasma, serum, cerebrospinal fluid, and brain tissues. Among them, blood and cerebrospinal fluid are preferable.

[0068] The method for detecting or quantitatively determining the peptide is not limited. Examples of the method include methods using an antibody, e.g., Western blotting, dot blotting, ELISA, sandwich ELISA, radioimmunoassay, and immunoprecipitation; mass spectrometry using a MALDI-TOF/MS; and combinations thereof. Among them, sandwich ELISA is most preferable. The sandwich ELISA may be conducted according to the description in the document of Tomita, et al. ("Cleavage of Alzheimer's amyloid precursor protein (APP) by secretases occurs after O-glycosylation of APP in the protein secretory pathway" Tomita, S., Kirino, Y., and Suzuki, T. J. Biol. Chem. 1998, 273, 6277-6284), for example. Specifically, the peptide of the present invention in a sample solution can be detected or quantitatively determined by (1) immobilizing an antibody specific to the peptide of the present invention on a solid phase, (2) adding the sample solution to the solid phase, (3) washing the solid phase, (4) adding another antibody specific to the peptide of the present invention, (5) adding an enzyme-labeled antibody (anti-IgG antibody) against the antibody, and (6) adding a substrate specific for the enzyme to detect the coloring or the like as an indicator. Here, the antibody specific to the peptide of the present invention can be prepared by a method described below. The anti-IgG antibody which is commercially available may be used. Examples of the solid phase include a micro-titer well and latex particles. Examples of the enzyme label include horseradish peroxidase, alkali phosphatase, and galactosidase.

[0069] The antibody of the present invention may be a monoclonal antibody or a polyclonal antibody.

[0070] The monoclonal antibody can be prepared, for example, by the method disclosed in the above-mentioned document of Tomita, et al. Specifically, a desired monoclonal antibody can be prepared by (1) administering the peptide of the present invention to an animal, (2) isolating antibody-producing cells from the animal, (3) fusing the antibody-producing cells with myeloma cells to prepare hybridomas, (4) selecting a hybridoma producing an antibody of the

present invention from the hybridomas, and (5) separating and purifying the antibody from the culture supernatant of the antibody-producing hybridoma.

[0071] The peptide of the present invention to be administered to an animal may be the whole peptide or a partial peptide thereof. The partial peptide to be administered is not limited, but it is preferable that, in the case of a peptide derived from Al α , the peptide has the amino acid at position 816, 821, or 839 of the amino acid sequence represented by SEQ ID NO: 1 as the N-terminal end and the amino acid at position 842, 843, or 851 of the amino acid sequence represented by SEQ ID NO: 1 as the C-terminal end; in the case of a peptide derived from Al β , the peptide (SEQ ID NO: 19) is composed of the amino acids at positions 826 to 845 of the amino acid sequence represented by SEQ ID NO: 2; and in the case of a peptide derived from Al γ , the peptide is composed of the amino acids at positions 805 to 824 of the amino acid sequence represented by SEQ ID NO: 3. In addition, the peptide may be administered with a complete or incomplete Freund's adjuvant in order to enhance the antibody productivity. The animal to be administered with the peptide is not limited. For example, monkey, rabbit, dog, guinea pig, mouse, rat, sheep, goat, and chicken may be used. The peptide administration intervals and number of times are not limited. In general, the peptide is administered about 2 to 10 times for every 2 to 6 weeks. The antibody-producing cells can be obtained by extracting spleen cells or lymph nodes from the animal 2 to 5 days after the last immunization. The myeloma cells to be used are not limited. For example, NS-1, P3U1, SP2/0, or AP-1 may be used. The hybridization can be performed by a general method using a polyethylene glycol or Sendai virus. The selection of a hybridoma which produces the antibody of the present invention can be performed, for example, by applying a culture supernatant of hybridomas to a micro-plate on which the peptide of the present invention is adsorbed, adding an anti-IgG antibody labeled with an enzyme, and detecting the anti-IgG antibody bound to the micro-plate. The isolation of the antibody of the present invention from the hybridoma culture supernatant can be performed by a general method for isolating and purifying immunoglobulin, e.g., salting-out, alcohol precipitation, isoelectric precipitation, electrophoresis, adsorption and desorption by an ion-exchanger, ultracentrifugation, or gel-filtration.

[0072] The polyclonal antibody can be also prepared, for example, according to the method disclosed in the documents of Araki, et al. (Araki, Y., et al., J. Biol. Chem. 2003, 278, 49448-49458 and Araki, Y., et al., J. Biol. Chem. 2004, 279, 24343-24354). Specifically, a desired polyclonal antibody can be prepared by (1) administering the peptide of the present invention to an animal, (2) extracting blood or ascites fluid from the animal, and (3) isolating and purifying the antibody from the blood or the like. The administration of the peptide and the isolation and purification of the antibody can be performed by the same manner as in the monoclonal antibody.

[0073] A diagnostic reagent of the present invention is generally prepared by adding the above-mentioned antibody of the present invention to an appropriate buffer solution. The concentration of the antibody and the kind of the buffer solution are not limited. They are properly determined according to the method for detecting or quantitatively determining the peptide of the present invention. Additionally, the diagnostic reagent may contain a component in addition to the antibody

of the present invention. Examples of such a component are an enzyme-labeled secondary antibody and a coloring agent.

[0074] The peptide of the present invention and Alcadein which is a precursor thereof are similar to A β and APP, respectively, in various respects. Therefore, it is highly suggested that a substance which suppresses the production of the peptide of the present invention also suppresses the production of A β . Furthermore, it is highly suggested that a substance which changes the molecular species of the peptide of the present invention from a high-molecular-weight peptide to a low-molecular-weight peptide (i.e., a peptide of the present invention other than the high-molecular-weight peptide) also changes the molecular species of A β from a high-molecular-weight peptide of highly toxic to other type. Therefore, it is thought that the screening for a therapeutic agent for Alzheimer's disease can be performed by contacting the cells secreting the peptide of the present invention with an agent to be screened and determining a change in the secretion of the peptide or a change in the molecular species of the secreted peptide.

[0075] The cells secreting the peptide of the present invention may be such cells that originally secrete the peptide of the present invention or may be such cells that have been transformed so as to secrete the peptide of the present invention by gene transfer. Examples of the former cells include fibroblasts (Araki, Y., et al., J. Biol. Chem. 2004, 279, 24343-24354) and HEK293 (in the Examples, an Alcadein gene is introduced into this cell, but endogenous Alcadein is expressed without the introduction). The latter cells can be prepared by introducing the full-length gene of Alcadein or a DNA encoding a first cleavage product (or a mimic construct thereof) into cells, for example. In addition, when a change in the molecular species of the peptide of the present invention is investigated, cells secreting a high-molecular-weight peptide of the present invention are preferably used. Such cells can be prepared by introducing a gene of a presenilin variant (1143F, 278T, 434C, L35F, etc.) into the cells so as to stably express the gene or inducing a mutation in the Alcadein gene, as shown in Examples 11 and 12 described below.

[0076] The method for contacting cells with an agent to be screened is not limited as long as the agent can act on the cells. Examples of the method include a method of directly inoculating an agent into cells and a method of adding an agent to a cell-culture medium.

[0077] The change in the amount of secreted peptide and the change in the molecular species of the secreted peptide can be investigated according to the method for detecting or quantitatively determining the peptide of the present invention described above. When a decrease in the secretion amount of the peptide caused by an agent to be screened is observed, the agent can be a candidate for a therapeutic agent for Alzheimer's disease. When a change in the molecular species of the peptide, i.e., a change from a high-molecular-weight peptide to a low-molecular-weight peptide, caused by an agent to be screened is observed, the agent also can be a candidate for a therapeutic agent for Alzheimer's disease.

[0078] Here, the term "therapeutic agent" includes not only an agent for treating Alzheimer's disease but also an agent having a preventive effect to suppress the onset of symptoms or delay the onset of symptoms of Alzheimer's disease.

EXAMPLES

[0079] The present invention will now be described further in detail with reference to Examples.

Example 1

[0080] The brain of five 8-week-old C57BL6 mice was homogenized with 10-strokes of a loose-fit Teflon homog-

enizer (clearance: 0.12 μm) in 30 ml of ice-cooled buffer A (10 mM HEPES of pH 7.4, 0.32 M sucrose, 5 $\mu\text{g}/\text{ml}$ chymostatin, 5 $\mu\text{g}/\text{ml}$ leupeptin, and 5 $\mu\text{g}/\text{ml}$ pepstatin). The homogenate was centrifuged (1000 \times g, 10 min) to remove unbroken cells and nuclei and to obtain a nucleus-removed cell homogenate. The nucleus-removed cell homogenate was further centrifuged (100000 \times g, 60 min) to obtain a pellet of a membrane fragment. The membrane fragment was resuspended in 2 ml of the buffer A and then gently overlaid on a solution (10 ml) of the buffer A with an iodixanol density-gradient of 0 to 28% in a Beckmann SW41 tube so as not to disturb the interface between the two solutions, and then centrifuged at 41000 rpm at 4° C. for 115 min. After the centrifugation, 13 fractions each of 900 ml were collected from the bottom of the tube. To 7.5 μl of each fraction, 5 μl of 5 \times SDS sample buffer (43% glycerol (Wako), 16% SDS (Wako), 64 ng/ml bromophenol blue (Wako), 5 mM EDTA, and 0.22 M Tris-HCl of pH 6.8) and 2.5 μl of 8 M urea solution were added. The resulting mixture was boiled for 5 min and then subjected to SDS-PAGE using an 8% gel according to the Laemmli method. Proteins on the gel were transferred on a nitrocellulose membrane for performing Western blotting. The detection was performed by using an ECL kit (Pharmacia). The used antibodies were an anti-APP cytoplasmic domain antibody (reactive to both full-length APP and a C-terminal fragment), anti-X11L antibody, anti-Alc α antibody, anti-protein disulfide-isomerase (PDI) antibody, anti-Golgi body 130 kDa matrix protein (GM-130) antibody, anti-synaptotagmin (SYT) antibody, anti-mouse kinesin heavy chain (KHC) antibody, and anti-presenilin 1 (PS1) C-terminal fragment antibody. Among them, commercially available anti-X11L antibody (mint2, BD Biosciences), anti-PDI antibody (1D3, Stressgen Biotechnologies), anti-GM130 antibody (#35, BD Biosciences), anti-SYT antibody (#41, BD Biosciences), anti-KHC antibody (H2, CHEMICON International), and anti-PS1 C-terminal fragment antibody (PS1-CTF, CHEMICON International) were used. The anti-APP cytoplasmic domain antibody was G369 and the anti-Alc α antibody was UT83. The G369 was prepared according to the method disclosed in Oishi, M., et al., Mol. Med. 1997, 3, 11-113. The UT83 was a polyclonal antibody derived from a rabbit immunized with an antigen peptide which was prepared by adding Cys to a C-terminal peptide (amino acids at positions 954 to 971) of human Alc α 1 (Araki, Y., et al., J. Biol. Chem. 2003, 278, 49448-49458). FIG. 1 shows the results of the Western blotting.

[0081] It was confirmed from the results of the Western blotting that large amounts of APP, X11L, and Alc were contained in the 8th fraction. Five hundred microliters of this fraction was added to an equal quantity of 2 \times CHAPS buffer (20 mM CHAPS, 20 mM sodium phosphate of pH 7.4, and 280 mM sodium chloride) to solubilize membrane components. Then, conjugate immunoprecipitation with G369 (anti-APP cytoplasmic domain antibody) was performed. Specifically, 4 W. of G369 was added to the solubilized membrane components. The resulting mixture was reacted at 4° C. for 1 hr, and then, to the mixture, 30 μl of 50% protein G-sepharose equilibrated with the 2 \times CHAPS buffer was further added. The resulting mixture was reacted at 4° C. for 1 hr. After the reaction, the beads were washed with 800 μl of the 2 \times CHAPS buffer, and then the components attached to the beads were solubilized by boiling the beads for 5 min in 45 μl of a sample-buffer mixture (a mixture of 30 μl of 5 \times SDS sample buffer and 15 μl of 8 M urea solution). The solubilized com-

ponents were subjected to SDS-PAGE using an 8% gel, followed by Western blotting as above. An antibody against IgG heavy chain (IgG (H)), in addition to the above-used anti-APP cytoplasmic domain antibody, anti-X11L antibody, anti-Alc α antibody, and anti-SYT antibody, was also used. In addition, as a control, components obtained by conjugate immunoprecipitation using an equal quantity of non-immunized rabbit serum instead of G369 were also subjected to Western blotting. Furthermore, similarly, the solubilized membrane components before the conjugate immunoprecipitation were also subjected to Western blotting. FIG. 2 shows the results.

[0082] As shown in FIG. 2, the components obtained by the conjugate immunoprecipitation with G369 contain not only APP but also X11L and Alc α . Therefore, it is thought that APP binds to X11L and Alc α to form a complex composed of the three.

Example 2

[0083] Frontal lobe tissues derived from 5 Alzheimer's disease patients were fixed in Kryofix (a mixture of ethanol, polyethylene glycol, and water:Merck) for 1 to 7 days and embedded in paraffin. The embedded tissues were cut into serial sections with a thickness of 4 μm . The sections were de-paraffined and then immunostained using ABC elite kit (Vector Laboratory).

[0084] The immunostain was performed by incubating the sections in a 0.8 $\mu\text{g}/\text{ml}$ anti-Alc α antibody (UT83) solution or a 0.5 $\mu\text{g}/\text{ml}$ anti-APP extracellular domain antibody (22C11: Roche Diagnostics) solution, reacting a secondary antibody with them, and visualizing the peroxidase activity by using a diaminobenzidine-hydrogen peroxide solution. As a control, the sections were incubated in a 0.8 $\mu\text{g}/\text{ml}$ non-immunized rabbit IgG solution and similarly immunostained. Additionally, the sections were incubated in a solution containing both the anti-Alc α antibody and its antigen peptide (40 nM) and similarly immunostained.

[0085] FIGS. 3-1, 3-2, and 3-3 show the results of the immunostain performed using the anti-Alc α antibody, anti-APP extracellular domain antibody, and non-immunized rabbit IgG, respectively.

[0086] As shown in these Figures, Alc α and APP are detected at similar brain regions of Alzheimer's disease patients. When the antigen peptide of the anti-Alc α antibody was presented in the solution (the results are not shown), nothing was detected as in the results shown in FIG. 3-3.

Example 3

[0087] The sections prepared in Example 2 were de-paraffined and incubated in a solution containing an anti-Alc α antibody (0.8 $\mu\text{g}/\text{ml}$) and an anti-APP extracellular domain antibody (0.5 $\mu\text{g}/\text{ml}$) or a solution containing an anti-Alc α antibody (0.8 $\mu\text{g}/\text{ml}$) and an anti-A β antibody (1/1000 dilution). As the anti-A β antibody, 4G8 (Sigma Lab) was used.

[0088] Then, the sections were incubated in solutions which each contain antibodies at the respective combinations, and then further incubated in a solution containing an FITC-labeled anti-rabbit IgG goat antibody (Jacson immunoresearch lab., 1/30 dilution) and a Cy3-labeled anti-mouse IgG goat antibody (Jacson immunoresearch lab., 1/50 dilution). The autofluorescence of lipofuscin granules was quenched by Sudan Black B staining before the immunoreaction.

[0089] The results when the anti-Alc α antibody and the anti-APP extracellular domain antibody were used as the primary antibodies are shown in FIG. 3-4 (only FITC was detected), FIG. 3-5 (only Cy3 was detected), and FIG. 3-6 (both FITC and Cy3 were detected). The results when the anti-Alc α antibody and the anti-A β antibody were used as the primary antibodies are shown in FIG. 4-1 (only FITC was detected), FIG. 4-2 (only Cy3 was detected), and FIG. 4-3 (both FITC and Cy3 were detected).

[0090] As shown in FIGS. 3-4, 3-5, and 3-6, Alc α and APP were detected at similar brain regions of Alzheimer's disease patients. This result agrees with that in Example 2. Additionally, as shown in FIGS. 4-1, 4-2, and 4-3, APP was detected in the vicinity of regions where senile plaques were formed by the accumulation of A β .

[0091] From the above-mentioned results, it is suggested that APP and Alc α similarly act in the pathogenesis process of Alzheimer's disease.

Example 4

[0092] A DNA encoding Alc α , Alc β , Alc γ , or APP695 (an isoform of human APP consisting of 695 amino acids) was inserted into a mammalian expression vector pcDNA3.1 (Invitrogen).

[0093] HEK293 cells were seeded in DMEM (D5796: Sigma) containing 10% fetal bovine serum in a 6-well culture plate (area of base: 10 cm²) and transfected with the expression vector prepared above by using a transfection reagent (LipofectAMINE 2000: Invitrogen). As a control, HEK293 cells were similarly transfected with an empty pcDNA3.1 vector.

[0094] One microliter of a DMSO solution containing 1 mM of a presenilin inhibitor L-685,458 (Calbiochem) was added to 1 ml of the culture medium. After the incubation for 24 hrs, a sample of the culture medium was taken. As a control, an equal quantity of DMSO instead of the L-685,458 solution was added to the culture medium and similarly incubated. Then, a sample of the culture medium was taken. The sample of the culture medium was added to 1 ml of HEST buffer (10 mM HEPES of pH 7.4, 150 mM sodium chloride, 0.5% Triton X-100, 5 μ g/ml chymostatin, 5 μ g/ml leupeptin, and 5 μ g/ml pepstatin) for extracting proteins of the cells. The solubilized cells were centrifuged (12000 \times g, 10 min) and supernatant was collected to recover the solubilized components. The solubilized components (7.5 μ l) were added to 7.5 μ l of a sample-buffer mixture (a mixture of 5 μ l of 5 \times SDS sample buffer and 2.5 μ l of 8 M urea solution) and boiled for 5 min. This sample was subjected to SDS-PAGE of 8/15% 2-stage gel and then the proteins were transferred on a nitrocellulose membrane for conducting Western blotting. The SDS-PAGE was performed according to the general method of Laemmli. As the primary antibody, an anti-APP cytoplasmic domain antibody (G369, 1/2000 dilution), anti-Alc α antibody (UT83, 0.3 μ g/ml), anti-Alc β antibody (UT99, 0.5 μ g/ml), and anti-Alc γ antibody (UT105, 1/500 dilution) were used. The detection was performed by using an ECL kit (Pharmacia). The UT99 and UT105 are antibodies recognizing the C-terminals of Alc β and Alc γ , respectively. FIG. 6 and FIGS. 7A to C show the results.

[0095] The C-terminal fragment obtained by a primary cleavage of APP is secondarily cleaved by presenilin, and the resulting cleavage fragment is extracellularly secreted (FIG. 5). At this point, the secondary cleavage is inhibited by adding a presenilin inhibitor and the C-terminal fragment of APP is

accumulated inside cells. As shown in FIG. 6, this fact is reflected to the result that a large amount of the C-terminal fragment (CTF α) was detected in the cell lysate only when the presenilin inhibitor (L-685,458) was added.

[0096] In the cases of Alc α and Alc γ , similarly to the case of APP, C-terminal fragments (CTF1) were detected only when the presenilin inhibitor was added (FIGS. 7A and C). From these results, it is thought that the C-terminal fragments of Alc α and Alc γ are cleaved by presenilin as in APP. However, the C-terminal fragment of Alc β was detected even when the presenilin inhibitor was not added (FIG. 7B). Therefore, it is not confirmed whether the C-terminal fragment of Alc β is cleaved by presenilin from this experiment only.

Example 5

[0097] DNAs encoding proteins (FLAG-Alc α , FLAG-Alc β , and FLAG-Alc γ), Alc α , Alc β , or Alc γ including a FLAG-tag sequence downstream of the N-terminal signal sequence thereof, were prepared. Each of the proteins was inserted into a mammalian expression vector pcDNA3.1 (Invitrogen).

[0098] HEK293 cells were seeded in DMEM (D5796: Sigma) containing 10% fetal bovine serum in a 6-well culture plate (area of base: 10 cm²) and the cells were transfected with the expression vector prepared above by using a transfection reagent (LipofectAMINE 2000: Invitrogen). As a control, HEK293 cells were transfected with an empty pcDNA3.1 vector.

[0099] One microliter of a DMSO solution containing 1 mM presenilin inhibitor L-685,458 (Calbiochem) was added to 1 ml of the culture medium. After the incubation for 24 hrs, a sample of the culture medium was taken. As a control, an equal quantity of DMSO instead of the L-685,458 solution was added to the culture medium and similarly incubated. Then, a sample of the culture medium was taken.

[0100] One hundred and fifty microliters of buffer B (7.7% SDS, 16.7 mM Tris-HCl of pH 7.4, 0.3 mg/ml chymostatin, 0.3 mg/ml leupeptin, and 0.3 mg/ml pepstatin) was added to 1 ml of the culture medium and boiled for 5 min to denature the protein components. Then, 3.75 ml of buffer C (6.7% NP-40, 0.4 M NaCl, 26 mM EDTA, and 200 mM Tris-HCl of pH 7.4) and 1.75 ml of an enzyme inhibition solution (distilled water containing 10 ng/ml leupeptin, 10 ng/ml pepstatin A, and 10 ng/ml chymostatin) were sequentially added. After the addition of 2 μ l of anti-FLAG antibody (Sigma), the resulting mixture was mixed by inverting the tube in a low-temperature chamber (4 $^{\circ}$ C.) for 8 hrs for an antigen-antibody reaction. Then, 50 μ l of rinse buffer (0.1% Triton X-100, 1 mM EDTA, 150 mM NaCl, and 10 mM Tris-HCl of pH 7.4) containing 25% protein G-sepharose/25% sepharose 4B (Pharmacia Biotech) was added, and the tube was rotated at 4 $^{\circ}$ C. for 3 hrs. The resin components were precipitated by centrifugation (3000 rpm, 5 min, 4 $^{\circ}$ C.) and recovered. The recovered resins were washed, in order to eliminate non-specific binding, with washing buffer I (0.1% Triton X-100, 1 M NaCl, and 20 mM Tris-HCl of pH 7.4), washing buffer II (0.05% SDS, 1% Triton X-100, 5 mM EDTA, 150 mM NaCl, and 50 mM Tris-HCl of pH 7.4), and the rinse buffer, sequentially. Then, 30 μ l of a sample-buffer mixture (a mixture of 20 μ l of 5 \times SDS sample buffer and 10 μ l of 8 M urea solution) was added to the resins and mixed. The mixture was boiled for 5 min for solubilizing components which were attached to the resins. After the centrifugation, the supernatant components were subjected to SDS-PAGE using a 6% gel and then the

proteins were transferred on a nitrocellulose membrane for conducting Western blotting. The SDS-PAGE was performed according to the general method of Lammler. As the primary antibody, an anti-FLAG antibody (M2, Sigma) was used. The detection was performed by using an ECL kit. FIGS. 7D to F show the results.

[0101] As shown in these Figures, fragments which can be recognized by the anti-FLAG antibody were detected in all culture media of Al α , Al β , and Al γ . Since the FLAG tag sequence was bound to the N-terminal of mature Al α , Al β , and Al γ , it is thought that the N-terminal fragments obtained by the primary cleavage of Al α , Al β , and Al γ are secreted extracellularly.

Example 6

[0102] DNAs encoding Al α , Al β , or Al γ were each inserted into a mammalian expression vector pcDNA3.1 (Invitrogen).

[0103] HEK293 cells were seeded in DMEM containing 10% fetal bovine serum in a 10-cm culture plate (area of base: 60 cm²) and transfected with the expression vector prepared above by using a transfection reagent (LipofectAMINE 2000: Invitrogen). As a control, HEK293 cells were transfected with an empty pcDNA3.1 vector.

[0104] After the incubation for 24 hrs, the culture medium was removed and the cells were washed with ice-cooled PBS. Then, 10 ml of PBS was added again. The cells were detached from the plate by pipetting and transferred into a 15 ml Falcon tube. The cells were collected by centrifugation (1500 rpm, 10 min, low-speed refrigerated centrifuge: Beckmann) as a pellet of the cells. One milliliter of buffer D (0.25 M sucrose, 10 mM triethanolamine-acetate of pH 7.8, 5 μ g/ml chymostatin, 5 μ g/ml leupeptin, and 5 μ g/ml pepstatin) was added to the pellet of the cells, and the cells were broken by 12 passages through a 27 G needle. Then, the broken cells were centrifuged with a TOMY TMA-6 rotor at 3000 rpm (1000 \times g) for 10 min at 4 $^{\circ}$ C. to remove unbroken cells and nuclei and to obtain a nucleus-removed cell homogenate. The nucleus-removed cell homogenate was further centrifuged with a Beckmann TLA-45 rotor at 45000 rpm (100000 \times g) for 60 min at 4 $^{\circ}$ C. to obtain a supernatant (cytoplasm fragment) and a precipitate (membrane fragment). The membrane fragment was resuspended in 100 μ l of the buffer D.

[0105] A sample of 20 μ l of the resuspended membrane-fragment was incubated at 37 $^{\circ}$ C. for 1 or 3 hrs. Separately, a sample of the resuspended membrane-fragment to which a presenilin inhibitor (L-685,458) was added at a final concentration of 1 μ M was also prepared. Twenty microliters of a sample-buffer mixture (a mixture of 13.4 μ l of 5 \times SDS sample buffer and 6.6 μ l of 8 M urea solution) was added to each sample for terminating the reaction. The sample was boiled for 5 min and was subjected to SDS-PAGE using 8/15% 2-stage gel. Then, the proteins were transferred on a nitrocellulose membrane for conducting Western blotting. The SDS-PAGE was performed according to the general method of Lammler. As the primary antibody, an anti-Al α antibody (UT83), anti-Al β antibody (UT99), and anti-Al γ antibody (UT105) were used. The detection was performed by using an ECL kit (Pharmacia). FIG. 8A to C shows the results.

[0106] As shown in these Figures, fragments (such as Al α -ICD) containing the C-terminal of each Alc were detected by incubating the membrane fragments of Al α , Al β , and Al γ . However, such fragments were not detected in the samples to which the presenilin inhibitor was added. On the basis of the

results above, it is thought that a fragment containing the C-terminal of Al α and so on was cleaved by presenilin.

Example 7

[0107] A DNA encoding Al α Δ E was prepared for efficiently detecting a cleavage product (A β -like fragment) obtained by a secondary cleavage of Al α . As shown in FIG. 9, in Al α Δ E, a fragment between the signal peptide and the primary cleavage site (indicated by δ in the Figure) was removed and the FLAG-tag sequence was introduced into the C-terminal end of the signal peptide.

[0108] A DNA encoding Al α Δ E was inserted into a mammalian expression vector pcDNA3.1 (Invitrogen). HEK293 cells were seeded in DMEM containing 10% fetal bovine serum in a 10-cm culture plate (area of base: 60 cm²) and transfected with the expression vector prepared above by using a transfection reagent (LipofectAMINE 2000: Invitrogen). As a control, HEK293 cells were similarly transfected with an empty pcDNA3.1 vector. A DMSO solution containing 10 μ M LLnL (Calbiochem), 1 μ M DAPT (Calbiochem), or 1 μ M L-685,458 (Calbiochem), which are presenilin inhibitors, was added to 1 ml of the culture medium at a final concentration of 1 μ M. After the incubation for 24 hrs, a sample of the culture medium was taken. As a control, an equal quantity of DMSO instead of the solution of the presenilin inhibitors such as L-685,458 was added to the culture medium and similarly incubated. Then, a sample of the culture medium was taken.

[0109] The recovered media were each subjected to immunoprecipitation using an anti-FLAG antibody (M2: Sigma) and Western blotting to detect a presenilin-induced N-terminal cleavage product. Specifically, 4 μ l of the anti-FLAG antibody was added to each recovered culture medium and reacted at 4 $^{\circ}$ C. for 1 hr. Then, 30 μ l of rinse buffer (0.1% Triton X-100, 1 mM EDTA, 150 mM NaCl, and 10 mM Tris-HCl of pH 7.4) containing 50% protein G-sepharose was added and further reacted at 4 $^{\circ}$ C. for 1 hr. Then, the beads were recovered and washed with 800 μ l of each of the washing buffer I (the composition is the same as in Example 5), the washing buffer II (the composition is the same as in Example 5), and the rinse buffer (the composition is the same as in above) sequentially. Then, the beads were boiled in 30 μ l of 2 \times tricine-sample buffer (900 mM Tris-HCl of pH 8.45, 24% glycerol, 8% SDS, and 0.005% Coomassie brilliant blue) for 5 min to solubilize components which were attached on the beads. The solubilized components were separated by using a 15% Tris-Tricine gel (as per Schagger & von Jagow method) and subjected to Western blotting using an anti-FLAG antibody (1/2000 dilution) and an anti-Al α antibody (UT83). The presenilin-induced cleavage product (A β -like fragment) and Al α Δ E were detected by an ECL kit (Pharmacia).

[0110] At the same time, cells were subjected to immunoprecipitation and Western blotting. Specifically, proteins were extracted from the cells in 4 ml of HBST buffer (the composition is the same as in Example 4). The solubilized cells were centrifuged (12000 \times g, 10 min) and the supernatant was collected to recover the solubilized components. Two microliters of the anti-FLAG antibody was added to 1 ml of the solubilized components and the mixture was reacted for 1 hr. Then, 30 μ l of HBST buffer containing 50% protein G-sepharose was added and further reacted at 4 $^{\circ}$ C. for 1 hr. Then, the beads were recovered and washed with 800 μ l of the HBST buffer 3 times. Then, the beads were boiled in 45 μ l of a sample-buffer mixture (a mixture of 30 μ l of a 5 \times SDS

sample buffer and 15 μ l of 8 M urea solution) for 5 min to solubilize components which were attached on the beads. The solubilized components were separated by using a 15% Tris-Tricine gel (as per Lamili method) and subjected to Western blotting using an anti-FLAG antibody (1/2000 dilution) and an anti-Alc α antibody (UT83). FIG. 10 shows the results mentioned above.

[0111] As shown in FIG. 10, Alc α Δ E which was transfected into the cells were detected in the cell lysates, but the presenilin-induced cleavage product (β -Alc α) was detected only in the culture media and was not detected in the cell lysates. From these results, it is thought that the presenilin-induced cleavage fragment has a property that the majority is secreted into the culture medium, as in A β .

Example 8

[0112] DNAs encoding Alc α or APP695 were each inserted into a mammalian expression vector pcDNA3.1 (Invitrogen), and a DNA (provided from Dr. Doms) encoding human BACE 1 was inserted into a mammalian expression vector pcDNA3.1Zeo(+) (Invitrogen).

[0113] HEK293 cells were seeded in DMEM (D5796: Sigma) containing 10% fetal bovine serum in a 6-well culture plate (area of base: 10 cm²) and transfected with the expression vector prepared above by using a transfection reagent (LipofectAMINE 2000: Invitrogen). Combinations of the introduced DNAs are shown in FIG. 11.

[0114] Each of the cells was incubated for 24 hrs. Proteins of the cells were solubilized in HBST buffer (the composition is the same as in Example 4) and subjected to SDS-PAGE using 8/15% gel. The proteins on the gel were transferred on a nitrocellulose membrane for conducting Western blotting using an anti-APP antibody (APP/c, Sigma) and an anti-Alc α antibody (UT83 antibody). FIGS. 11A and B show the results of Western blotting using the anti-APP antibody and anti-Alc α antibody, respectively.

[0115] As shown in FIG. 11A, in the cells not expressing BACE1 (3rd lane from the left), CTF α , which is a cleavage product at the α -site, was mainly detected, but in the cells expressing BACE1 (4th lane from the left), CTF β , which is a cleavage product at the β -site, was also detected. Additionally, only two bands indicated by arrows correspond to APP695 and another two bands detected in the vicinity thereof correspond to endogenous APPs (APP770 and APP751).

[0116] As shown in FIG. 11B, in the cells expressing Alc α but not expressing BACE1 (5th lane from the left), a fragment (CTF1) was mainly detected at a position of 30 kDa. Since the molecular weight of this fragment was approximately equal to that of Alc α Δ E expressed in Example 7, it is predicted that the cleavage site is between Met-815 and Ala-816 or in the vicinity thereof and that the number of amino acids is about 156. In the cells expressing both Alc α and BACE1 (6th lane from the left), CTF1 and a fragment (CTF β , the number of amino acid residues calculated from the molecular weight is about 280) with a molecular weight larger than that of CTF1 were detected. It is thought that this fragment is generated when Alc α is cleaved by BACE1. Namely, it is thought that Alc α is cleaved by BACE1 as in APP. Additionally, with reference to FIG. 11, Alc α is detected as two bands. This is due to sugar chain modification.

Example 9

[0117] The cleavage site of γ -secretase in Alc α was identified as follows: A cDNA construct (FIG. 9) expressing

Alc α Δ E protein was prepared. As shown in FIG. 12, cells transfected with this cDNA construct produce and secrete β -Alc α having a FLAG-tag sequence. The β -Alc α secreted by the cells was recovered by immunoprecipitation using an anti-FLAG antibody, and the molecular weight of the immunoprecipitate was analyzed by using a MALDI-TOF/MS. On the basis of this molecular weight, the cleavage site of γ -secretase was identified. The details will now be described.

(1) Analysis of β -Alc α having a FLAG sequence by Western blotting

[0118] HEK293 cells were seeded in a 10-cm dish (Corning). When the cells became confluent, the cells were transfected with an expression vector of Alc α Δ E (pcDNA3-FLAG-hAlc α Δ E) by using a transfection reagent (LipofectAMINE 2000: Invitrogen). The cells were incubated in a CO₂ incubator for 24 hrs. Hereat, 2 μ l of a DMSO solution containing 1 mM γ -secretase inhibitor L-685,458 (Calbiochem) was added to 1 ml of the culture medium. As a control, cells to which added only the DMSO solution were also prepared. Six milliliters of each culture medium was recovered and centrifuged (15000 rpm, 5 min, high-speed refrigerated microcentrifuge: TOMY). To the resulting supernatant, 1/1000 volume of an enzyme inhibition solution (a DMSO solution containing 5 mg/ml leupeptin, 5 mg/ml pepstatin A, and 5 mg/ml chymostatin) was added to prepare a sample for immunoprecipitation. This sample was mixed with 6 μ l of an anti-FLAG antibody solution (M2: Sigma) by inverting at 4 $^{\circ}$ C. for 1 hr. Then, to the mixture, 50 μ l of rinse buffer (10 mM Tris-HCl of pH 7.4, 1 mM EDTA, 0.1% Triton X-100, and 150 mM NaCl) containing 25% protein G-sepharose was added, and an antigen-antibody reaction was conducted by mixing by inverting the mixture at 4 $^{\circ}$ C. for 1 hr. After the reaction, the beads were precipitated and recovered by centrifugation (3000 rpm, 5 min, 4 $^{\circ}$ C., high-speed refrigerated microcentrifuge: TOMY SEIKO CO., LTD.), and washed with washing buffer 1 (1 M NaCl, 20 mM Tris-HCl of pH 7.4, and 0.1% Triton X-100), washing buffer 2 (150 mM NaCl, 5 mM EDTA, 50 mM Tris-HCl of pH 7.4, 1% Triton X-100, and 0.05% SDS), and rinse buffer, sequentially. Then, the beads were stirred in 20 μ l of a sample-buffer mixture (a mixture of 10 μ l of 2 \times SDS sample buffer and 10 μ l of 8 M urea solution) and then boiled for 5 min to elute components which were adsorbed on the beads. After the centrifugation, the supernatant components were separated by 20% acrylamide Tris-Tricine gel electrophoresis and then subjected to Western blotting using an anti-FLAG antibody (M2: Sigma). The reacting β -Alc having the FLAG-tag was detected by an ECL kit (Pharmacia).

[0119] Ice-cooled PBS (1.5 ml) was added to the cells isolated from the culture medium. Then, the cells were detached from the plate by pipetting and transferred into an eppendorf tube. The cells were collected by centrifugation (6000 rpm, 5 min, high-speed refrigerated microcentrifuge: TOMY SEIKO CO., LTD.). The resulting cell pellet was mixed with 0.9 ml of HBST buffer (the composition is the same as in Example 4) by inverting at 4 $^{\circ}$ C. for 1 hr to extract proteins of the cells. The solubilized cells were centrifuged (15000 rpm, 15 min, 4 $^{\circ}$ C., high-speed refrigerated microcentrifuge: TOMY SEIKO CO., LTD.) to recover the solubilized components as the supernatant. To 5 μ l of the solubilized components, 10 μ l of 2 \times SDS sample buffer and 5 μ l of 1% SDS were added. The resulting mixture was boiled for 5 min. The solubilized components were separated by 20% acrylamide Tris-Tricine gel electrophoresis and then transferred to a nitrocel-

lulose membrane (S&S). After the reaction with an anti-FLAG antibody (M2: Sigma), the reaction product was detected by an ECL kit (Pharmacia) on the membrane.

[0120] FIG. 13 shows the results of the Western blotting mentioned above. In the immunoprecipitate sample of the culture medium, 13-Alc α having the FLAG sequence was mainly detected as two bands at approximately 5 kDa (the lower diagram, the right lane). However, the β -Alc α was not detected in the immunoprecipitate when the γ -secretase inhibitor L-685,458 was added to the cell culture solution (the lower diagram, the right lane). From this experiment, it was revealed that β -Alc α having the FLAG sequence can be recovered by using the anti-FLAG antibody and it was confirmed that the β -Alc α is a cleavage product of γ -secretase. (2) Mass spectrometry of β -Alc α having a FLAG sequence by MALDI-TOF/MS

[0121] HEK293 cells were transfected with an expression vector of Alc α Δ E (pcDNA3-FLAG-hAlc α Δ E) by using a transfection reagent (LipofectAMINE 2000: Invitrogen) to establish a cell line stably expressing Alc α Δ E having the FLAG sequence. The established cell line cells were seeded in a 225-cm² flask. When the cells became confluent, the culture solution was changed to 40 ml of DMEM medium (D5796: Sigma) containing 10% fetal bovine serum and the cells were incubated in a CO₂ incubator for 24 hrs. Hereat, 1 μ l of a DMSO solution containing 1 mM γ -secretase inhibitor L-685,458 (Calbiochem) was added to 1 ml of the culture medium. As a control, cells to which only the DMSO solution was added were also prepared. The each culture medium was recovered and centrifuged (10000 \times g, 5 min, high-speed refrigerated centrifuge: Beckmann). An enzyme inhibition solution (a DMSO solution containing 5 mg/ml leupeptin, 5 mg/ml pepstatin A, and 5 mg/ml chymostatin) at a volume ratio of 1/1000 and 10% sodium azide solution at a volume ratio of 1/1000 were added to the resulting supernatant to prepare a sample for immunoprecipitation. Twenty milliliters of this sample was mixed with 20 μ l agarose beads (A2220: Sigma) conjugated to 50% (v/v) anti-FLAG antibody (M2) by inverting in a low-temperature chamber (4 $^{\circ}$ C.) overnight (about 12 hrs) for an antigen-antibody reaction. After the reaction, the beads were precipitated and recovered by centrifugation (3000 rpm, 5 min, 4 $^{\circ}$ C., low-speed refrigerated centrifuge: Beckmann).

[0122] The recovered beads were washed, in order to eliminate non-specific binding, with 800 μ l of washing buffer 1 (0.1% N-octylglucoside, 140 mM NaCl, 10 mM Tris-HCl of pH 8.0, and 0.025% sodium azide) twice and washing buffer 2 (10 mM Tris-HCl of pH 8.0 and 0.025% sodium azide) twice, sequentially. Then, the beads were stirred with 10 μ l of a matrix solution (trifluoroacetate (Wako Pure Chemical Industries, Ltd.)/acetonitrile (Sigma)/water (1:20:20) saturated with sinapinic acid (Applied Biosystem)) to elute components from the beads. In order to completely remove the beads, the eluted components were loaded onto a spin column (Amersham Bioscience) and centrifuged to separate the eluted components only. Two microliters of the eluted components were loaded onto a sample plate (Applied Biosystems) and dried, and then analyzed by using a MALDI-TOF/MS (PerSpective Biosystems).

[0123] Signals of the cell-culture solution of the cells not expressing Alc α Δ E were used as a background. Peptides having molecular weights of 3619.96, 5359.85, 5507.02, and 6233.83 were detected as β -Alc α having a FLAG sequence (FIG. 14).

[0124] Cleavage sites corresponding to these molecular weights were indicated by arrows in FIG. 15. An amino acid sequence predicted to be a transmembrane domain is shown by a solid line and a potential region to be included in the transmembrane domain is shown by a dotted line. On the basis of the fact that cleavage sites determined from the molecular weights of 5359.85, 5507.02, and 6233.83 exist in the transmembrane domain or the potential region to be the transmembrane domain, the cleavage sites can be determined to be in a cleavage domain of γ -secretase. On the other hand, since the cleavage site determined from the molecular weight of 3619.96 obviously exists in an extracellular domain, there is a high possibility that a cleavage product of β -Alc α is generated by the cleavage at the γ -site. Therefore, the secondary cleavage sites of Alc α by γ -secretase are at least three. This agrees with that fact that APP has a plurality of 7-sites and the variety of A β is produced and secreted (FIG. 16).

Example 10

[0125] FIG. 16 shows cleavage sites of APP (human APP695) and cleavage products obtained by the cleavage which have been already revealed (the numbers shown in the Figure are amino acid Nos. in the human APP695 isoform). Examples of the cleavage sites include the α -site and β -site which are primary cleavage sites, the γ -site which is a secondary cleavage site, and the ϵ -site which was identified recently (Gu, Y., Misonou, H., Sato, T., Dohmae, N., Takio, K., and Ihara, Y. J. Biol. Chem. 2002, 276, 35235-35238). Examples of the cleavage products include A β 40 and A β 42 which are main A β species and p3 peptide which is a cleavage product of the α -site.

[0126] Since Alc α has various similarities to APP, there is a high possibility that Alc α also has a plurality of primary cleavage sites. Hence, the primary cleavage site was identified by determining an amino acid sequence of the N-terminal of a C-terminal fragment (CTF) obtained by the primary cleavage of Alc α . However, since the CTF tends to subsequently receive a secondary cleavage, it is difficult to recover a sufficient amount of the protein for determining its N-terminal sequence from the cell extract. Therefore, a cell line of HEK293 cells stably expressing a dominant-negative protein so that a CTF obtained from Alc α does not receive a secondary cleavage was established by inducing a variant into an active site of presenilin (PS) which is a catalytic subunit of γ -secretase. The cell line was further transfected with an expression vector of Alc α 1-FLAG which is Alc α having a FLAG tag at the C-terminal (pcDNA3-hAlc α 1-FLAG) by using a transfection reagent (LipofectAMINE 2000: Invitrogen) to establish a cell line stably expressing Alc α 1-FLAG.

[0127] From the extract of this cell line, CTF α 1-FLAG obtained by a primary cleavage was immunoprecipitated. The precipitate was separated by discontinuous SDS electrophoresis with 8% (upper gel)-15% (lower gel) of acrylamide gels and transferred on a PVDF membrane (Immunobilon-PSQ: Millipore). The CTF α 1-FLAG was detected by Coomassie brilliant blue staining. Two kinds of protein of approximately 30 kDa, which were detected by antibody-specific detection, were analyzed by a gas-phase protein sequencer to determine three amino acid sequences derived from Alc α 1. The details will now be described.

[0128] HEK293 cells were transfected with an expression vector (pcDNA3-PS1D385A) of a PS1(D385A) variant protein obtained by substituting alanine for aspartic acid at position 385 in presenilin 1 (PS1) by using a transfection reagent

(LipofectAMINE 2000: Invitrogen); thus, a cell line stably expressing PS1 (D385A) was established. The cells of this cell line were further transfected with an Al α FLAG expression vector (pcDNA3-hAl α 1-FLAG) by using a transfection reagent (LipofectAMINE 2000: Invitrogen).

[0129] The cells of this cell line (denoted as hAl α 1FLAG/PS1D385A-293cell) were incubated in four 10-cm dishes until reaching a confluent state (about 1×10^8 cells/dish). The cells were washed with ice-cooled PBS. Then, 15 ml of an HBST solution (10 mM HEPES of pH 7.4, 150 mM NaCl, and 0.5% Triton X-100) containing an enzyme inhibition solution (a DMSO solution containing 5 μ g/ml leupeptin, 5 μ g/ml pepstatin A, and 5 μ g/ml chymostatin) was added to the cells and the resulting mixture was stirred by inverting at 4° C. for 0.5 hrs to solubilize the cells. The solubilized cells were centrifuged (12000 \times g, 10 min, 4° C., high-speed microcentrifuge: TOMY SEIKO CO., LTD.); thus, soluble components were recovered as the supernatant. Fifteen milliliters of the solubilized components was mixed with 30 μ l of agarose beads (A2220: Sigma) conjugated to 50% (v/v) anti-FLAG antibody (M2) by inverting in a low-temperature chamber (4° C.) overnight (about 12 hrs) for an antigen-antibody reaction. After the reaction, the beads were precipitated and recovered by centrifugation (3000 rpm, 5 min, 4° C., low-speed refrigerated centrifuge: Beckmann). Then, the beads were washed with 1 ml of HBST 3 times. After the addition of 2 mg/ml FLAG-peptide-containing HBST (30 μ l), the immunoprecipitate was competitively eluted; which is a method for performing antigen elution under moderate conditions in order to avoid contamination of immunoglobulin light chains (30 kDa or less) because the molecular weight of CTF1 of Al α is about 30 kDa. The eluate was centrifuged (12000 \times g, 10 min, 4° C., high-speed microcentrifuge: TOMY SEIKO CO., LTD.). Then, 5 \times SDS sample buffer was added to the supernatant at a volume ratio of 1/5 and acrylamide gel electrophoresis was performed according to the general method of Laemmli. The separated proteins were transferred on an Immobilon-PSQ membrane (Millipore) and detected by staining using Coomassie brilliant blue (CBB) stain (FIG. 17).

[0130] Two kinds of protein (proteins indicated as CTF1s) were detected at approximately 30 kDa. The amino acid sequences of these proteins were determined by using a gas-phase protein sequencer 492HT (Applied BioSystem). The results revealed that the protein with a higher molecular weight includes two types of amino acid sequence. The primary cleavage sites deduced based on these sequences are between Met-815 and Ala-816 (this cleavage site is referred to as “ ζ 1”) and between Gln-820 and Phe-821 (this cleavage site is referred to as “ ζ 2”) (FIG. 18). The ζ 1 agrees with the site determined in Example 8. The ζ 2 is a newly found primary cleavage site. Additionally, it was revealed that the protein with a lower molecular weight of CTF1s includes one type of amino acid sequence. The primary cleavage site deduced based on this sequence is between Ala-838 and Asn-839 (this cleavage site is referred to as “ ζ 3”) (FIG. 18).

[0131] As shown above, Al α has 3 primary cleavage sites (ζ 1, ζ 2, and ζ 3) and 3 secondary cleavage sites (γ 1, γ 2, and γ 3). As a result, it is understood that, in human beings, at least 9 types of β -Al α are produced (Table 1).

TABLE 1

PRIMARY CLEAVAGE SITE	SECONDARY CLEAVAGE SITE	THE NUMBER OF AMINO ACIDS	SEQUENCE LISTING
ζ 1	γ 1	27	SEQ ID NO: 4
ζ 1	γ 2	28	SEQ ID NO: 5
ζ 1	γ 3	36	SEQ ID NO: 6
ζ 2	γ 1	22	SEQ ID NO: 7
ζ 2	γ 2	23	SEQ ID NO: 8
ζ 2	γ 3	31	SEQ ID NO: 9
ζ 3	γ 1	4	SEQ ID NO: 10
ζ 3	γ 2	5	SEQ ID NO: 11
ζ 3	γ 3	13	SEQ ID NO: 12

[0132] The result that a plurality of types of β -Al α are generated well agrees with the fact that a plurality of types of A β generated from APP which is synchronously metabolized.

Example 11

[0133] In Alzheimer’s disease (AD), it is recognized not only an increase in the generation of A β but also a change in the molecular species of A β . Furthermore, it is reported that the ratio of the amount of A β 42, which is highly aggregative, to the total amount of generated A β is increased in AD patients. The increase of the ratio of A β 42 is thought to be highly involved in the onset of Alzheimer’s disease. For example, it is known that the ratio of A β 42 to A β is prominently increased in patients of familial Alzheimer’s disease (FAD) having a variant in the presenilin gene. Since Al α has various similarities to APP, there is a possibility in Al α that the molecular species of β -Al α generated by a presenilin variant is changed as in APP. Consequently, in order to confirm this possibility, the following experiment was conducted.

[0134] Expression vectors (pcDNA3-PS1I143F, pcDNA3-PS1R278T, pcDNA3-PS1A434C, and pcDNA3-PS1L435F) expressing four types of PS1 variant found in AD patients, i.e., I143F (substituting Phe for Ile at position 143), R278T (substituting Thr for Arg at position 278), A434C (substituting Cys for Ala at position 434), and L435F (substituting Phe for Leu at position 435) were prepared. These vectors and an expression vector of C99/CTF of APP (pcDNA3-APPC99) were introduced to HEK293 cells by using a transfection reagent (LipofectAMINE 2000: Invitrogen) to establish a cell line stably expressing both proteins. The cells of the cell line were seeded in a 10-cm dish (Corning). When the cells became confluent, pcDNA3-FLAG-hAl α Δ E was introduced into the cells by using a transfection reagent (LipofectAMINE 2000: Invitrogen) to transiently express CTF1 of Alcadin α .

[0135] Separately, expression vectors (pcDNA3-PS1 and pcDNA3-PS1D385A) expressing wild-type PS1 (wt) and inactive-type PS1 (D385A, Asp at the catalytic site of PS1 is substituted with Ala not to have γ -secretase activity) were prepared and introduced into HEK293 cells by using a transfection reagent (LipofectAMINE 2000: Invitrogen) to establish cell lines stably expressing both types of PS1. The cells of each cell line were seeded in a 10-cm dish (Corning). When the cells became confluent, pcDNA3-APPC99 and pcDNA3-FLAG-hAl α Δ E were introduced into the cells by using a transfection reagent (LipofectAMINE 2000: Invitrogen) to transiently express CTF of APP and CTF1 of Alcadin α .

[0136] The cells transfected with genes were incubated in a CO₂ incubator for 24 hrs. The culture solution was recovered and centrifuged (15000 rpm, 5 min, 4° C., high-speed refrigerated centrifuge: Beckmann). To 7.5 ml of the supernatant, 7.5 µl of an enzyme inhibition solution (a DMSO solution containing 5 mg/ml leupeptin, 5 mg/ml pepstatin A, and 5 mg/ml chymostatin) was added to prepare a sample. After the addition of 6 µl of an anti-FLAG antibody solution (M2: Sigma, lot No. 103k6043) to the sample, the resulting mixture was mixed by inverting at 4° C. for 1 hr. Then, 50 µl of rinse buffer containing 25% protein G-sepharose was added and the mixture was mixed by inverting at 4° C. overnight for an antigen-antibody reaction. After the reaction, the beads were washed with washing buffer 1 (1 M NaCl, 20 mM Tris-HCl of pH 7.4, and 0.1% Triton X-100), washing buffer 2 (150 mM NaCl, mM EDTA, 50 mM Tris-HCl of pH 7.4, 1% Triton X-100, and 0.05% SDS), and rinse buffer (10 mM Tris-HCl of pH 7.4, 1 mM EDTA, 0.1% Triton X-100, and 150 mM NaCl), sequentially. Then, 20 µl of a sample-buffer mixture (a mixture of 10 ml of 2×SDS sample buffer and 10 ml of 8 M urea solution) was added to the beads and stirred. The beads were boiled for 5 min to elute components which were adsorbed to the beads.

After the centrifugation, the supernatant components were separated by 20% acrylamide Tris-Tricine gel electrophoresis and then subjected to Western blotting using an anti-FLAG antibody solution (M2: Sigma). The reacted β-Alcα having a FLAG tag was detected by using an ECL kit (Pharmacia) and quantitatively determined by using a NIH image software. At the same time, Aβ40 and Aβ42 in the culture medium were quantitatively determined by sELISA according to the method of Tomita, et al. (J. Biol. Chem. 1988, 273, 6277-6284). FIG. 19 shows the results of the Western blotting and ratios of the amount of long β-Alcα to the total amount of generated β-Alcα (1.0) and ratios of the amount of Aβ42 to the total amount of generated Aβ (1.0). The N.D. in the Figure means the value was lower than the detection limit.

[0137] In the cells expressing the wild-type PS1, 2 types of β-Alc (indicated as short β-Alc and medium (3-Alc in the Figure) were mainly detected. On the other hand, in the cells expressing PS1 having a FAD variant, the amount of β-Alc (indicated as long (3-Alc) having a higher molecular weight was increased. The ratio of the long β-Alc to the total β-Alcα was increased in the cells expressing a PS1 variant; which is the same tendency as the increase in the ratio of Aβ42 to the total Aβ. Namely, it was revealed that a qualitative change in β-Alc reflects a quantitative change in Aβ. A qualitative change in β-Alc (such as the increase in the ratio of long β-Alc) in cerebrospinal fluid or blood of patients reflects a

qualitative change in Aβ. Therefore, the detection of β-Alc instead of the detection of Aβ42, which is highly aggregative, can find patients at an early stage or pre-patients of whom qualitative change is difficult to detect.

Example 12

[0138] In order to use in the determination of a cleavage site at the C-terminal of β-Alc shown in FIG. 19 by using a MALDI-TOF/MS, a PS1 variant expressing a larger amount of a high-molecular-weight β-Alc was intensively searched by conducting the same experiment as in Example 11. As a result, as shown in FIG. 20, it was found that a large amount of high-molecular-weight β-Alc was secreted in the culture medium of the cells expressing L166P (Leu at position 166 is substituted with Pro) PS1 variant, compared with other PS1 variants. Consequently, the molecular weights of β-Alc produced and secreted by the wild-type PS1 and the L166P variant-type PS1 were determined by mass spectrometry according to the method in Example 9. FIG. 21 shows the results. In the cells expressing the L166P variant-type PS1, the amount of short β-Alc (indicated by a bold downward arrow) is decreased and the amount of medium β-Alc (indicated by a bold upward arrow) is increased compared to those in the cells expressing the wild-type PS1. This is also obvious from the results of the Western blotting analysis shown in FIG. 20. Furthermore, long β-Alc, which was not detected in the cells expressing the wild-type PS1, was also detected.

[0139] FIG. 22 shows the cleavage sites of β-Alc which were revealed from the results shown in FIG. 21. The dotted line arrows indicate the cleavage sites in the cells expressing the wild-type PS1 and the solid line arrows indicate the cleavage sites in the cells expressing the L166P variant-type PS1. It was revealed that the cleavage sites of β-Alc shifted to the C-terminal side in the L166P variant-type PS1. This result well agrees with the fact that the cleavage site at the C-terminal side of Aβ shifts to the C-terminal side in a variant-type PS1 of familial Alzheimer's disease. FIG. 23 schematically shows 3 cleavage sites (ζ1, ζ2, and ζ3) at the N-terminal side and 3 cleavage sites (γ1, γ2, and γ3) at the C-terminal side of β-Alc and β-Alc molecular species generated by the cleavage at the γ-site shifted to the C-terminal side by the variant-type PS1 (the cleavage site γ1 was not detected in the experiment in Example 12).

[0140] The present invention contains subject matter disclosed in the specification and/or the drawings of Japanese Patent Application No. 2003-375363 on which the claim for a priority right is based, and the entire contents of the documents, patents, and patent applications cited as references are incorporated herein by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 19

<210> SEQ ID NO 1
<211> LENGTH: 971
<212> TYPE: PRT
<213> ORGANISM: human

<400> SEQUENCE: 1

Met Leu Arg Arg Pro Ala Pro Ala Leu Ala Pro Ala Ala Arg Leu Leu

-continued

1	5	10	15
Leu Ala Gly	Leu Leu Cys Gly	Gly Gly Val Trp Ala Ala Arg Val Asn	
	20	25	30
Lys His Lys	Pro Trp Leu Glu Pro	Thr Tyr His Gly Ile Val Thr Glu	
	35	40	45
Asn Asp Asn	Thr Val Leu Leu Asp	Pro Pro Leu Ile Ala Leu Asp Lys	
	50	55	60
Asp Ala Pro	Leu Arg Phe Ala Gly	Glu Ile Cys Gly Phe Lys Ile His	
	65	70	75
Gly Gln Asn	Val Pro Phe Asp Ala	Val Val Val Asp Lys Ser Thr Gly	
	85	90	95
Glu Gly Val	Ile Arg Ser Lys Glu	Lys Leu Asp Cys Glu Leu Gln Lys	
	100	105	110
Asp Tyr Ser	Phe Thr Ile Gln Ala	Tyr Asp Cys Gly Lys Gly Pro Asp	
	115	120	125
Gly Thr Asn	Val Lys Lys Ser His	Lys Ala Thr Val His Ile Gln Val	
	130	135	140
Asn Asp Val	Asn Glu Tyr Ala Pro	Val Phe Lys Glu Lys Ser Tyr Lys	
	145	150	155
Ala Thr Val	Ile Glu Gly Lys Gln	Tyr Asp Ser Ile Leu Arg Val Glu	
	165	170	175
Ala Val Asp	Ala Asp Cys Ser Pro	Gln Phe Ser Gln Ile Cys Ser Tyr	
	180	185	190
Glu Ile Ile	Thr Pro Asp Val Pro	Phe Thr Val Asp Lys Asp Gly Tyr	
	195	200	205
Ile Lys Asn	Thr Glu Lys Leu Asn	Tyr Gly Lys Glu His Gln Tyr Lys	
	210	215	220
Leu Thr Val	Thr Ala Tyr Asp Cys	Gly Lys Lys Arg Ala Thr Glu Asp	
	225	230	235
Val Leu Val	Lys Ile Ser Ile Lys	Pro Thr Cys Thr Pro Gly Trp Gln	
	245	250	255
Gly Trp Asn	Asn Arg Ile Glu Tyr	Glu Pro Gly Thr Gly Ala Leu Ala	
	260	265	270
Val Phe Pro	Asn Ile His Leu Glu	Thr Cys Asp Glu Pro Val Ala Ser	
	275	280	285
Val Gln Ala	Thr Val Glu Leu Glu	Thr Ser His Ile Gly Lys Gly Cys	
	290	295	300
Asp Arg Asp	Thr Tyr Ser Glu Lys	Ser Leu His Arg Leu Cys Gly Ala	
	305	310	315
Ala Ala Gly	Thr Ala Glu Leu Leu	Pro Ser Pro Ser Gly Ser Leu Asn	
	325	330	335
Trp Thr Met	Gly Leu Pro Thr Asp	Asn Gly His Asp Ser Asp Gln Val	
	340	345	350
Phe Glu Phe	Asn Gly Thr Gln Ala	Val Arg Ile Pro Asp Gly Val Val	
	355	360	365
Ser Val Ser	Pro Lys Glu Pro Phe	Thr Ile Ser Val Trp Met Arg His	
	370	375	380
Gly Pro Phe	Gly Arg Lys Lys Glu	Thr Ile Leu Cys Ser Ser Asp Lys	
	385	390	395
Thr Asp Met	Asn Arg His His Tyr	Ser Leu Tyr Val His Gly Cys Arg	
	405	410	415

-continued

Leu Ile Phe Leu Phe Arg Gln Asp Pro Ser Glu Glu Lys Lys Tyr Arg
 420 425 430

Pro Ala Glu Phe His Trp Lys Leu Asn Gln Val Cys Asp Glu Glu Trp
 435 440 445

His His Tyr Val Leu Asn Val Glu Phe Pro Ser Val Thr Leu Tyr Val
 450 455 460

Asp Gly Thr Ser His Glu Pro Phe Ser Val Thr Glu Asp Tyr Pro Leu
 465 470 475 480

His Pro Ser Lys Ile Glu Thr Gln Leu Val Val Gly Ala Cys Trp Gln
 485 490 495

Glu Phe Ser Gly Val Glu Asn Asp Asn Glu Thr Glu Pro Val Thr Val
 500 505 510

Ala Ser Ala Gly Gly Asp Leu His Met Thr Gln Phe Phe Arg Gly Asn
 515 520 525

Leu Ala Gly Leu Thr Leu Arg Ser Gly Lys Leu Ala Asp Lys Lys Val
 530 535 540

Ile Asp Cys Leu Tyr Thr Cys Lys Glu Gly Leu Asp Leu Gln Val Leu
 545 550 555 560

Glu Asp Ser Gly Arg Gly Val Gln Ile Gln Ala His Pro Ser Gln Leu
 565 570 575

Val Leu Thr Leu Glu Gly Glu Asp Leu Gly Glu Leu Asp Lys Ala Met
 580 585 590

Gln His Ile Ser Tyr Leu Asn Ser Arg Gln Phe Pro Thr Pro Gly Ile
 595 600 605

Arg Arg Leu Lys Ile Thr Ser Thr Ile Lys Cys Phe Asn Glu Ala Thr
 610 615 620

Cys Ile Ser Val Pro Pro Val Asp Gly Tyr Val Met Val Leu Gln Pro
 625 630 635 640

Glu Glu Pro Lys Ile Ser Leu Ser Gly Val His His Phe Ala Arg Ala
 645 650 655

Ala Ser Glu Phe Glu Ser Ser Glu Gly Val Phe Leu Phe Pro Glu Leu
 660 665 670

Arg Ile Ile Ser Thr Ile Thr Arg Glu Val Glu Pro Glu Gly Asp Gly
 675 680 685

Ala Glu Asp Pro Thr Val Gln Glu Ser Leu Val Ser Glu Glu Ile Val
 690 695 700

His Asp Leu Asp Thr Cys Glu Val Thr Val Glu Gly Glu Glu Leu Asn
 705 710 715

His Glu Gln Glu Ser Leu Glu Val Asp Met Ala Arg Leu Gln Gln Lys
 725 730 735

Gly Ile Glu Val Ser Ser Ser Glu Leu Gly Met Thr Phe Thr Gly Val
 740 745 750

Asp Thr Met Ala Ser Tyr Glu Glu Val Leu His Leu Leu Arg Tyr Arg
 755 760 765

Asn Trp His Ala Arg Ser Leu Leu Asp Arg Lys Phe Lys Leu Ile Cys
 770 775 780

Ser Glu Leu Asn Gly Arg Tyr Ile Ser Asn Glu Phe Lys Val Glu Val
 785 790 795 800

Asn Val Ile His Thr Ala Asn Pro Met Glu His Ala Asn His Met Ala
 805 810 815

-continued

Ala Gln Pro Gln Phe Val His Pro Glu His Arg Ser Phe Val Asp Leu
820 825 830

Ser Gly His Asn Leu Ala Asn Pro His Pro Phe Ala Val Val Pro Ser
835 840 845

Thr Ala Thr Val Val Ile Val Val Cys Val Ser Phe Leu Val Phe Met
850 855 860

Ile Ile Leu Gly Val Phe Arg Ile Arg Ala Ala His Arg Arg Thr Met
865 870 875 880

Arg Asp Gln Asp Thr Gly Lys Glu Asn Glu Met Asp Trp Asp Asp Ser
885 890 895

Ala Leu Thr Ile Thr Val Asn Pro Met Glu Thr Tyr Glu Asp Gln His
900 905 910

Ser Ser Glu Glu Glu Glu Glu Glu Glu Glu Glu Ser Glu Asp
915 920 925

Gly Glu Glu Glu Asp Asp Ile Thr Ser Ala Glu Ser Glu Ser Ser Glu
930 935 940

Glu Glu Glu Gly Glu Gln Gly Asp Pro Gln Asn Ala Thr Arg Gln Gln
945 950 955 960

Gln Leu Glu Trp Asp Asp Ser Thr Leu Ser Tyr
965 970

<210> SEQ ID NO 2
<211> LENGTH: 968
<212> TYPE: PRT
<213> ORGANISM: human

<400> SEQUENCE: 2

Met Val Leu Gly Cys Glu Leu Ser Gly Ser Thr Arg Val Val Val Gly
1 5 10 15

Val Glu Ala Leu Leu Thr Gly Ala Ser Ser Pro Leu Pro Gly Val Gly
20 25 30

Pro Ala Asn Lys His Lys Pro Trp Ile Glu Ala Glu Tyr Gln Gly Ile
35 40 45

Val Met Glu Asn Asp Asn Thr Val Leu Leu Asn Pro Pro Leu Phe Ala
50 55 60

Leu Asp Lys Asp Ala Pro Leu Arg Tyr Ala Gly Glu Ile Cys Gly Phe
65 70 75 80

Arg Leu His Gly Ser Gly Val Pro Phe Glu Ala Val Ile Leu Asp Lys
85 90 95

Ala Thr Gly Glu Gly Leu Ile Arg Ala Lys Glu Pro Val Asp Cys Glu
100 105 110

Ala Gln Lys Glu His Thr Phe Thr Ile Gln Ala Tyr Asp Cys Gly Glu
115 120 125

Gly Pro Asp Gly Ala Asn Thr Lys Lys Ser His Lys Ala Thr Val His
130 135 140

Val Arg Val Asn Asp Val Asn Glu Phe Ala Pro Val Phe Val Glu Arg
145 150 155 160

Leu Tyr Arg Ala Ala Val Thr Glu Gly Lys Leu Tyr Asp Arg Ile Leu
165 170 175

Arg Val Glu Ala Ile Asp Gly Asp Cys Ser Pro Gln Tyr Ser Gln Ile
180 185 190

Cys Tyr Tyr Glu Ile Leu Thr Pro Asn Thr Pro Phe Leu Ile Asp Asn
195 200 205

-continued

Asp Gly Asn Ile Glu Asn Thr Glu Lys Leu Gln Tyr Ser Gly Glu Arg
 210 215 220
 Leu Tyr Lys Phe Thr Val Thr Ala Tyr Asp Cys Gly Lys Lys Arg Ala
 225 230 235 240
 Ala Asp Asp Ala Glu Val Glu Ile Gln Val Lys Pro Thr Cys Lys Pro
 245 250 255
 Ser Trp Gln Gly Trp Asn Lys Arg Ile Glu Tyr Ala Pro Gly Ala Gly
 260 265 270
 Ser Leu Ala Leu Phe Pro Gly Ile Arg Leu Glu Thr Cys Asp Glu Pro
 275 280 285
 Leu Trp Asn Ile Gln Ala Thr Ile Glu Leu Gln Thr Ser His Val Ala
 290 295 300
 Lys Gly Cys Asp Arg Asp Asn Tyr Ser Glu Arg Ala Leu Arg Lys Leu
 305 310 315 320
 Cys Gly Ala Ala Thr Gly Glu Val Asp Leu Leu Pro Met Pro Gly Pro
 325 330 335
 Asn Ala Asn Trp Thr Ala Gly Leu Ser Val His Tyr Ser Gln Asp Ser
 340 345 350
 Ser Leu Ile Tyr Trp Phe Asn Gly Thr Gln Ala Val Gln Val Pro Leu
 355 360 365
 Gly Gly Pro Ser Gly Leu Gly Ser Gly Pro Gln Asp Ser Leu Ser Asp
 370 375 380
 His Phe Thr Leu Ser Phe Trp Met Lys His Gly Val Thr Pro Asn Lys
 385 390 395 400
 Gly Lys Lys Glu Glu Glu Thr Ile Val Cys Asn Thr Val Gln Asn Glu
 405 410 415
 Asp Gly Phe Ser His Tyr Ser Leu Thr Val His Gly Cys Arg Ile Ala
 420 425 430
 Phe Leu Tyr Trp Pro Leu Leu Glu Ser Ala Arg Pro Val Lys Phe Leu
 435 440 445
 Trp Lys Leu Glu Gln Val Cys Asp Asp Glu Trp His His Tyr Ala Leu
 450 455 460
 Asn Leu Glu Phe Pro Thr Val Thr Leu Tyr Thr Asp Gly Ile Ser Phe
 465 470 475 480
 Asp Pro Ala Leu Ile His Asp Asn Gly Leu Ile His Pro Pro Arg Arg
 485 490 495
 Glu Pro Ala Leu Met Ile Gly Ala Cys Trp Thr Glu Glu Lys Asn Lys
 500 505 510
 Glu Lys Glu Lys Gly Asp Asn Ser Thr Asp Thr Thr Gln Gly Asp Pro
 515 520 525
 Leu Ser Ile His His Tyr Phe His Gly Tyr Leu Ala Gly Phe Ser Val
 530 535 540
 Arg Ser Gly Arg Leu Glu Ser Arg Glu Val Ile Glu Cys Leu Tyr Ala
 545 550 555 560
 Cys Arg Glu Gly Leu Asp Tyr Arg Asp Phe Glu Ser Leu Gly Lys Gly
 565 570 575
 Met Lys Val His Val Asn Pro Ser Gln Ser Leu Leu Thr Leu Glu Gly
 580 585 590
 Asp Asp Val Glu Thr Phe Asn His Ala Leu Gln His Val Ala Tyr Met
 595 600 605

-continued

```

Asn Thr Leu Arg Phe Ala Thr Pro Gly Val Arg Pro Leu Arg Leu Thr
 610                               615                               620

Thr Ala Val Lys Cys Phe Ser Glu Glu Ser Cys Val Ser Ile Pro Glu
 625                               630                               635                               640

Val Glu Gly Tyr Val Val Val Leu Gln Pro Asp Ala Pro Gln Ile Leu
                               645                               650                               655

Leu Ser Gly Thr Ala His Phe Ala Arg Pro Ala Val Asp Phe Glu Gly
                               660                               665                               670

Thr Asn Gly Val Pro Leu Phe Pro Asp Leu Gln Ile Thr Cys Ser Ile
                               675                               680                               685

Ser His Gln Val Glu Ala Lys Lys Asp Glu Ser Trp Gln Gly Thr Val
 690                               695                               700

Thr Asp Thr Arg Met Ser Asp Glu Ile Val His Asn Leu Asp Gly Cys
 705                               710                               715                               720

Glu Ile Ser Leu Val Gly Asp Asp Leu Asp Pro Glu Arg Glu Ser Leu
                               725                               730                               735

Leu Leu Asp Thr Thr Ser Leu Gln Gln Arg Gly Leu Glu Leu Thr Asn
                               740                               745                               750

Thr Ser Ala Tyr Leu Thr Ile Ala Gly Val Glu Ser Ile Thr Val Tyr
 755                               760                               765

Glu Glu Ile Leu Arg Gln Ala Arg Tyr Arg Leu Arg His Gly Ala Ala
 770                               775                               780

Leu Tyr Thr Arg Lys Phe Arg Leu Ser Cys Ser Glu Met Asn Gly Arg
 785                               790                               795                               800

Tyr Ser Ser Asn Glu Phe Ile Val Glu Val Asn Val Leu His Ser Met
                               805                               810                               815

Asn Arg Val Ala His Pro Ser His Val Leu Ser Ser Gln Gln Phe Leu
                               820                               825                               830

His Arg Gly His Gln Pro Pro Pro Glu Met Ala Gly His Ser Leu Ala
 835                               840                               845

Ser Ser His Arg Asn Ser Met Ile Pro Ser Ala Ala Thr Leu Ile Ile
 850                               855                               860

Val Val Cys Val Gly Phe Leu Val Leu Met Val Val Leu Gly Leu Val
 865                               870                               875                               880

Arg Ile His Ser Leu His Arg Arg Val Ser Gly Ala Gly Gly Pro Pro
                               885                               890                               895

Gly Ala Ser Ser Asp Pro Lys Asp Pro Asp Leu Phe Trp Asp Asp Ser
                               900                               905                               910

Ala Leu Thr Ile Ile Val Asn Pro Met Glu Ser Tyr Gln Asn Arg Gln
 915                               920                               925

Ser Cys Val Thr Gly Ala Val Gly Gly Gln Gln Glu Asp Glu Asp Ser
 930                               935                               940

Ser Asp Ser Glu Val Ala Asp Ser Pro Ser Ser Asp Glu Arg Arg Ile
 945                               950                               955                               960

Ile Glu Thr Pro Pro His Arg Tyr
                               965

```

```

<210> SEQ ID NO 3
<211> LENGTH: 955
<212> TYPE: PRT
<213> ORGANISM: human

<400> SEQUENCE: 3

```

-continued

Met Leu Pro Gly Arg Leu Cys Trp Val Pro Leu Leu Leu Ala Leu Gly
 1 5 10 15
 Val Gly Ser Gly Ser Gly Gly Gly Gly Asp Ser Arg Gln Arg Arg Leu
 20 25 30
 Leu Ala Ala Lys Val Asn Lys His Lys Pro Trp Ile Glu Thr Ser Tyr
 35 40 45
 His Gly Val Ile Thr Glu Asn Asn Asp Thr Val Ile Leu Asp Pro Pro
 50 55 60
 Leu Val Ala Leu Asp Lys Asp Ala Pro Val Pro Phe Ala Gly Glu Ile
 65 70 75 80
 Cys Ala Phe Lys Ile His Gly Gln Glu Leu Pro Phe Glu Ala Val Val
 85 90 95
 Leu Asn Lys Thr Ser Gly Glu Gly Arg Leu Arg Ala Lys Ser Pro Ile
 100 105 110
 Asp Cys Glu Leu Gln Lys Glu Tyr Thr Phe Ile Ile Gln Ala Tyr Asp
 115 120 125
 Cys Gly Ala Gly Pro His Glu Thr Ala Trp Lys Lys Ser His Lys Ala
 130 135 140
 Val Val His Ile Gln Val Lys Asp Val Asn Glu Phe Ala Pro Thr Phe
 145 150 155 160
 Lys Glu Pro Ala Tyr Lys Ala Val Val Thr Glu Gly Lys Ile Tyr Asp
 165 170 175
 Ser Ile Leu Gln Val Glu Ala Ile Asp Glu Asp Cys Ser Pro Gln Tyr
 180 185 190
 Ser Gln Ile Cys Asn Tyr Glu Ile Val Thr Thr Asp Val Pro Phe Ala
 195 200 205
 Ile Asp Arg Asn Gly Asn Ile Arg Asn Thr Glu Lys Leu Ser Tyr Asp
 210 215 220
 Lys Gln His Gln Tyr Glu Ile Leu Val Thr Ala Tyr Asp Cys Gly Gln
 225 230 235 240
 Lys Pro Ala Ala Gln Asp Thr Leu Val Gln Val Asp Val Lys Pro Val
 245 250 255
 Cys Lys Pro Gly Trp Gln Asp Trp Thr Lys Arg Ile Glu Tyr Gln Pro
 260 265 270
 Gly Ser Gly Ser Met Pro Leu Phe Pro Ser Ile His Leu Glu Thr Cys
 275 280 285
 Asp Gly Ala Val Ser Ser Leu Gln Ile Val Thr Glu Leu Gln Thr Asn
 290 295 300
 Tyr Ile Gly Lys Gly Cys Asp Arg Glu Thr Tyr Ser Glu Lys Ser Leu
 305 310 315 320
 Gln Lys Leu Cys Gly Ala Ser Ser Gly Ile Ile Asp Leu Leu Pro Ser
 325 330 335
 Pro Ser Ala Ala Thr Asn Trp Thr Ala Gly Leu Leu Val Asp Ser Ser
 340 345 350
 Glu Met Ile Phe Lys Phe Asp Gly Arg Gln Gly Ala Lys Ile Pro Asp
 355 360 365
 Gly Ile Val Pro Lys Asn Leu Thr Asp Gln Phe Thr Ile Thr Met Trp
 370 375 380
 Met Lys His Gly Pro Ser Pro Gly Val Arg Ala Glu Lys Glu Thr Ile
 385 390 395 400

-continued

<211> LENGTH: 22
<212> TYPE: PRT
<213> ORGANISM: human

<400> SEQUENCE: 7

Phe Val His Pro Glu His Arg Ser Phe Val Asp Leu Ser Gly His Asn
1 5 10 15

Leu Ala Asn Pro His Pro
20

<210> SEQ ID NO 8
<211> LENGTH: 23
<212> TYPE: PRT
<213> ORGANISM: human

<400> SEQUENCE: 8

Phe Val His Pro Glu His Arg Ser Phe Val Asp Leu Ser Gly His Asn
1 5 10 15

Leu Ala Asn Pro His Pro Phe
20

<210> SEQ ID NO 9
<211> LENGTH: 31
<212> TYPE: PRT
<213> ORGANISM: human

<400> SEQUENCE: 9

Phe Val His Pro Glu His Arg Ser Phe Val Asp Leu Ser Gly His Asn
1 5 10 15

Leu Ala Asn Pro His Pro Phe Ala Val Val Pro Ser Thr Ala Thr
20 25 30

<210> SEQ ID NO 10
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: human

<400> SEQUENCE: 10

Asn Pro His Pro
1

<210> SEQ ID NO 11
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: human

<400> SEQUENCE: 11

Asn Pro His Pro Phe
1 5

<210> SEQ ID NO 12
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: human

<400> SEQUENCE: 12

Asn Pro His Pro Phe Ala Val Val Pro Ser Thr Ala Thr
1 5 10

<210> SEQ ID NO 13
<211> LENGTH: 73

-continued

<212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 13

```

Ile Ser Glu Val Lys Met Asp Ala Glu Phe Arg His Asp Ser Gly Tyr
1           5           10           15
Glu Val His His Gln Lys Leu Val Phe Phe Ala Glu Asp Val Gly Ser
                20           25           30
Asn Lys Gly Ala Ile Ile Gly Leu Met Val Gly Gly Val Val Ile Ala
            35           40           45
Thr Val Ile Val Ile Thr Leu Val Met Leu Lys Lys Lys Gln Tyr Thr
        50           55           60
Ser Ile His His Gly Val Val Gln Asn
65           70
  
```

<210> SEQ ID NO 14
 <211> LENGTH: 37
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 14

```

Ala Ala Gln Pro Gln Phe Val His Pro Glu His Arg Ser Phe Val Asp
1           5           10           15
Leu Ser Gly His Asn Leu Ala Asn Pro His Pro Phe Ala Val Val Pro
            20           25           30
Ser Thr Ala Thr Val
        35
  
```

<210> SEQ ID NO 15
 <211> LENGTH: 38
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 15

```

Ala Ala Gln Pro Gln Phe Val His Pro Glu His Arg Ser Phe Val Asp
1           5           10           15
Leu Ser Gly His Asn Leu Ala Asn Pro His Pro Phe Ala Val Val Pro
            20           25           30
Ser Thr Ala Thr Val Val
        35
  
```

<210> SEQ ID NO 16
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 16

```

Ala Ala Gln Pro Gln Phe Val His Pro Glu His Arg Ser Phe Val Asp
1           5           10           15
Leu Ser Gly His Asn Leu Ala Asn Pro His Pro Phe Ala Val Val Pro
            20           25           30
Ser Thr Ala Thr Val Val Ile
        35
  
```

<210> SEQ ID NO 17
 <211> LENGTH: 40

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Human

<400> SEQUENCE: 17

Ala Ala Gln Pro Gln Phe Val His Pro Glu His Arg Ser Phe Val Asp
1           5           10          15

Leu Ser Gly His Asn Leu Ala Asn Pro His Pro Phe Ala Val Val Pro
          20          25          30

Ser Thr Ala Thr Val Val Ile Val
          35          40

<210> SEQ ID NO 18
<211> LENGTH: 50
<212> TYPE: PRT
<213> ORGANISM: Human

<400> SEQUENCE: 18

Leu Ser Ser Gln Gln Phe Leu His Arg Gly His Gln Pro Pro Pro Glu
1           5           10          15

Met Ala Gly His Ser Leu Ala Ser Ser His Arg Asn Ser Met Ile Pro
          20          25          30

Ser Ala Ala Thr Leu Ile Ile Val Val Cys Val Gly Phe Leu Val Leu
          35          40          45

Met Val
          50

<210> SEQ ID NO 19
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human

<400> SEQUENCE: 19

Leu Ser Ser Gln Gln Phe Leu His Arg Gly His Gln Pro Pro Pro Glu
1           5           10          15

Met Ala Gly His
          20

```

1. A peptide capable of being a diagnostic marker for Alzheimer's disease, the peptide being obtained by cleaving an N-terminal region and a C-terminal region of Alcadein β .

2. The peptide according to claim 1, wherein the N-terminal region to be cleaved is a portion of an extracellular domain of the N-terminal end of Alcadein β .

3. The peptide according to claim 1, wherein the C-terminal region is cleaved by presenilin.

4. The peptide according to claim 1, wherein the peptide is obtained by cleaving an N-terminal and a C-terminal region of Alcadein β ; and the cleavage site of the N-terminal region is between amino acids 825 and 826 of the amino acid sequence represented by SEQ ID NO: 2.

5. The peptide according to claim 1, wherein the peptide is obtained by cleaving an N-terminal and a C-terminal region of Alcadein β ; and the cleavage site of the C-terminal region is between amino acids 875 and 876 of the amino acid sequence represented by SEQ ID NO: 2.

6. An isolated peptide consisting of the amino acid sequence selected from the group consisting of SEQ ID NO:2.

7. A method for diagnosing Alzheimer's disease, which comprises detecting or quantitatively determining the peptide according to any one of claims 1 to 6 in body fluid or tissues taken from an animal.

8. The method for diagnosing Alzheimer's disease according to claim 7, wherein the body fluid is blood or cerebrospinal fluid.

9. The method for diagnosing Alzheimer's disease according to claim 7, wherein a ratio of a high-molecular-weight peptide to the detected or quantitatively determined peptide is used as an indicator for diagnosing Alzheimer's disease.

10. A method for diagnosing Alzheimer's disease, which comprises:

obtaining a sample of brain tissue taken from the subject, determining co-localization of the peptide according to claim 6 with amyloid precursor protein (APP) present in said sample,

wherein Alzheimer's disease is indicated when the co-localization is detected as compared to the absence of said co-localization in a control non-Alzheimer's disease sample.

11. A method for screening a therapeutic agent for Alzheimer's disease, which comprises contacting cells secreting the peptide according to any one of claims 1 to 6 with an agent to be screened; and determining a change in the secreted amount of the peptide or a change in the molecular species of the secreted peptide.

12. An antibody against the peptide according to any one of claims 1 to 6.

13. A diagnostic reagent for Alzheimer's disease, the reagent comprising the antibody according to claim 12.

* * * * *

专利名称(译)	阿尔茨海默病的标记肽		
公开(公告)号	US20110014635A1	公开(公告)日	2011-01-20
申请号	US12/859950	申请日	2010-08-20
[标]申请(专利权)人(译)	铃木利治 荒木洋一 宫城NAOMI 山口晴康 西村MASAKI KAZUO YAMAMOTO		
申请(专利权)人(译)	铃木利治 荒木洋一 宫城NAOMI 山口晴康 西村MASAKI KAZUO YAMAMOTO		
当前申请(专利权)人(译)	免疫生物研究所CO. , LTD.		
[标]发明人	SUZUKI TOSHIHARU ARAKI YOICHI MIYAGI NAOMI YAMAGUCHI HARUYASU NISHIMURA MASAKI YAMAMOTO KAZUO		
发明人	SUZUKI, TOSHIHARU ARAKI, YOICHI MIYAGI, NAOMI YAMAGUCHI, HARUYASU NISHIMURA, MASAKI YAMAMOTO, KAZUO		
IPC分类号	G01N33/53 C07K14/00 C07K16/18 A61K38/00 C07K14/47 G01N33/68		
CPC分类号	A61K38/00 C07K14/4711 G01N2800/2821 G01N33/6896 C07K16/18		
优先权	2003375363 2003-11-05 JP PCT/JP2004/016209 2004-11-01 WO		
外部链接	Espacenet USPTO		

摘要(译)

提供可通过切割Alcadeina α , Alcadein β 或Alcadein γ 的N末端区域和C末端区域获得的肽;并且能够成为阿尔茨海默病的诊断标志物。通过使用肽作为诊断标记,可以在早期阶段检测阿尔茨海默氏病而不会使受试者负担。

