



US009005992B2

(12) **United States Patent**  
**Cha et al.**

(10) **Patent No.:** **US 9,005,992 B2**  
(45) **Date of Patent:** **Apr. 14, 2015**

(54) **IMMOBILIZING FUSION PROTEIN FOR EFFECTIVE AND ORIENTED IMMOBILIZATION OF ANTIBODY ON SURFACES**

2010/0266455 A1\* 10/2010 Sommer ..... 422/100  
2012/0202748 A1\* 8/2012 Cha et al. .... 514/19.1  
2013/0052712 A1\* 2/2013 Cha et al. .... 435/177

FOREIGN PATENT DOCUMENTS

(75) Inventors: **Hyung Joon Cha**, Pohang-si (KR);  
**Chang Sup Kim**, Pohang-si (KR)

KR 10-2011-0051778 5/2011

OTHER PUBLICATIONS

(73) Assignee: **Postech Academy-Industry Foundation**, Pohang (KR)

Lin et al. "Adhesion mechanisms of the mussel foot proteins mfp-1 and mfp-3", vol. 104 No. 10 (2007).\*

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1062 days.

C. S. Kim, W. Ko, and H. J. Cha, "Novel linker material based on mussel adhesive protein fused with BC domain of protein A for efficient surface immobilization of antibody", Proceedings of the 239th ACS National Meeting, San Francisco, Mar. 21-25, 2010.

(21) Appl. No.: **13/051,219**

Chang Sup Kim, Wooree Ko and Hyung Joon Cha, "Effective Immobilization of Antibodies with Fc-specific BC Domains-fused Mussel Adhesive Protein", Proceedings of the 2010 KSBB Spring Meeting and International Symposium, Kyungwon University, Korea, Apr. 15-16, 2010.

(22) Filed: **Mar. 18, 2011**

Chang Sup Kim, Wooree Ko and Hyung Joon Cha, "A novel material of BC-mussel adhesive protein fusion protein for efficient immobilization of antibody", Proceeding of the 2010 KICHe Spring Meeting, Theories and Applications of Chemical Engineering, vol. 16, No. 1, 2010, p. 402.

(65) **Prior Publication Data**

US 2012/0238039 A1 Sep. 20, 2012

\* cited by examiner

(51) **Int. Cl.**

**G01N 33/543** (2006.01)

**G01N 33/531** (2006.01)

**G01N 21/65** (2006.01)

**C07K 14/435** (2006.01)

*Primary Examiner* — Anh Lam

(52) **U.S. Cl.**

CPC ..... **G01N 33/54353** (2013.01); **G01N 33/531** (2013.01); **G01N 21/658** (2013.01); **C07K 14/43504** (2013.01); **C07K 2319/705** (2013.01); **C07K 2319/21** (2013.01)

(74) *Attorney, Agent, or Firm* — Lexyoume IP Meister, PLLC

(58) **Field of Classification Search**

None  
See application file for complete search history.

(57) **ABSTRACT**

The present invention relates to a novel fusion protein comprising Staphylococcal protein A and mussel adhesive protein, a biochip comprising a solid substrate to which the fusion protein is attached, and a method for detecting a target antigen in a biological sample using the biochip. Furthermore, the present invention relates to a polynucleotide encoding the fusion protein, a recombinant vector comprising the polynucleotide, a transformed cell comprising the recombinant vector, and a method of preparing the fusion protein by transformed cell comprising the recombinant vector.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,817,470 A \* 10/1998 Burzio et al. .... 435/7.9  
8,673,986 B2 \* 3/2014 Cha et al. .... 514/772.1  
8,765,682 B2 \* 7/2014 Cha et al. .... 514/19.1  
2007/0042427 A1 \* 2/2007 Gerdes et al. .... 435/7.1  
2010/0261286 A1 \* 10/2010 Kim et al. .... 436/149

**19 Claims, 8 Drawing Sheets**

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7





Fig. 8



# IMMOBILIZING FUSION PROTEIN FOR EFFECTIVE AND ORIENTED IMMOBILIZATION OF ANTIBODY ON SURFACES

## BACKGROUND OF THE INVENTION

### (a) Field of the Invention

The present invention relates to a novel fusion protein comprising Staphylococcal protein A and mussel adhesive protein, a biochip comprising a solid substrate to which the fusion protein is attached, and a method for detecting a target antigen in a biological sample using the biochip. Furthermore, the present invention relates to a polynucleotide encoding the fusion protein, a recombinant vector comprising the polynucleotide, a transformed cell comprising the recombinant vector, and a method of preparing the fusion protein by transformed cell comprising the recombinant vector.

### (b) Description of the Related Art

Early detection of diseases is a key for their successful treatments despite of advancement of therapeutic tools. Immunoassay is one of the vital technologies for disease diagnosis due to high specificity and affinity of antibodies to target antigens. In most of immunoassays, antibody-antigen interactions are measured on solid supports. Because effective immobilization of antibody on solid support controls specificity and sensitivity of assay devices, immobilization of antibody with efficient manner is crucial and one of key issues for various fields including immunoassays and immunosensors.

Antibodies have been immobilized on solid supports by diverse methods including physical adsorption and covalent binding (R. Pei, X. Yang, E. Wang, *Analyst* 2001, 126, 4.; A. Subramanina, J. Irudayaraj, T. Ryan, *Biosens. Bioelectron.* 2006, 21, 998.). However, these methods have been suffered from reduced antibody activity (that is, antigen binding ability) due to random orientation, denaturation, and chemical modification of antibodies (G. T. Hermanson, *Bioconjugate Techniques*, Academic press, California 1996.; H. Zhu, M. Synder, *Curr. Opin. Chem. Biol.* 2003, 7, 55.). Therefore, antibody-binding proteins such as protein A and G have been widely employed to overcome limitations of the previous physical and chemical methods. Those proteins specifically recognize and bind to Fc region of antibodies with high affinity, and realized oriented-immobilization of antibodies on surfaces without reduced antibody activity (M. D. P. Boyle, K. J. Reis, *Biotechnology* 1987, 5, 697.; W. L. Hoffman, D. J. O' Shannessy, *J. Immunol. Methods* 1998, 112, 113.). Thus, the antibody-binding protein-based antibody immobilization methods have significantly improved antigen-binding ability, sensitivity, and stability compared to the previous methods.

However, efficient attachment of antibody-binding proteins on solid surfaces still remains as a major challenge. For efficient binding with functional orientation, antibody-binding proteins have been fused with several biomolecules such as cysteine residue(s), oligonucleotide, hexahistidine (His<sub>6</sub>) peptide, gold-binding peptide, and glutathione-S-transferase (J. M. Lee, et al., *Anal. Chem.* 2007, 79, 2680.; Y. Jung, et al., *Anal. Chem.* 2007, 79, 6534.; S. M. Patrie, et al., *Anal. Chem.* 2007, 79, 5878.; S. Ko, et al., *Biosens. Bioelectron.* 2009, 24, 2592.; T. H. Ha, et al., *Anal. Chem.* 2007, 79, 546.). However, the above fusion proteins have still limitation because immobilization is possible onto only corresponding reactive group-derived surfaces. For protein immobilization, preparation of modified surface is time-consuming and sometimes difficult for non-experts.

Mussel adhesive proteins (MAPs) from marine mussels are water-insoluble bioadhesives that adhere tightly to substrata. MAPs are able to form strong bond to diverse substrates including glass, metal, and plastics without any surface pretreatments (J. H. Waite, *Int. J. Adhesion Adhesive* 1987, 7, 9.; L. O. Burzio, et al., *The Japanese Soc. Mar. Biotechnology*, Tokyo, 1989.). However, the amounts of MAPs that can be extracted directly from mussels are extremely low and attempts to produce functional recombinant MAPs have been failed (D. Morgan, *Scientist* 1990, 4, 1; M. Kitamura, et al., *J. Polym. Sci. Ser. A* 1999, 37, 729; A. J. Salerno, et al., *Appl. Microbiol. Biotechnol.* 1993, 58, 209.).

In the present work, we constructed a novel molecule, BC-MAP, as a functional immobilizing linker by genetic fusion of MAP with two domains (B and C) of protein A for facile and effective immobilization of antibodies on diverse solid supports with oriented manner without any additional steps and/or pretreatments. A representative scheme of effective immobilization of antibody onto diverse surfaces using functional BC-MAP linker material is shown in FIG. 8.

## SUMMARY OF THE INVENTION

The present invention provides a fusion protein comprising protein A or a peptide derived from protein A and mussel adhesive protein or a peptide derived from a mussel adhesive protein.

The present invention also provides a polynucleotide encoding the fusion protein.

The present invention also provides a recombinant vector comprising the polynucleotide.

The present invention also provides a transformed cell comprising the recombinant vector.

The present invention also provides a method of preparing the fusion protein by transformed cell comprising the recombinant vector.

The present invention also provides a biochip comprising a solid substrate to which the fusion protein is attached.

The present invention also provides a method for detecting a target antigen in a biological sample using the biochip.

## BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1(a) shows schematic representation of pET-BC-MAP and pET-BC plasmids for the expression of BC-MAP fusion protein and BC domain protein used in the present invention. FIG. 1(b) shows Coomassie-blue-stained SDS-PAGE analysis of purified BC-MAP and BC. Lane: M, molecular weight.

FIG. 2 shows direct coatings of BC-MAP and sole BC with color image analyses on diverse surfaces. Alkaline phosphatase-conjugated anti-rabbit IgG was used as a model antibody.

FIG. 3(a) shows color image analysis depending on immobilized alkaline phosphatase-conjugated anti-rabbit IgG concentration on glass slide coated by BC-MAP and BC.

FIG. 3(b) shows color signal intensities are plotted in the bottom graph. Each value and error bar represents the mean of six independent spots and its standard deviation.

FIG. 4(a) shows color images of anti-alkaline phosphatase functionalized glass slide coated by BC-MAP response to different concentrations of alkaline phosphatase. 100  $\mu\text{g mL}^{-1}$  anti-alkaline phosphatase was used.

FIG. 4(b) shows color signal intensities are plotted in bottom graph. Each value and error bar represents the mean of six independent spots and its standard deviation.

FIG. 5(a) shows schematic procedure to investigate the specificity of antibodies functionalized glass slide coated by BC-MAP.

FIG. 5(b) Specificity of antibodies functionalized glass slide coated by BC-MAP. Antibodies for alkaline phosphatase (1) and transferrin (2) were added onto each defined region of BC-MAP-coated glass slide and the slide was reacted with sample containing alkaline phosphatase.

FIG. 6(a) shows QCM response for the binding of BC and BC-MAP on gold at flow rate  $10 \mu\text{L min}^{-1}$ .

FIG. 6(b) shows QCM response for the binding of antibody on BC- and BC-MAP-coated gold at a flow rate of  $10 \mu\text{L min}^{-1}$ .

FIG. 6(c) shows data analysis of frequency shift upon the binding of BC and BC-MAP on gold at a flow rate of  $10 \mu\text{L min}^{-1}$ . Values are the means from three independent experiments. Standard derivations are indicated in bar graphs.

FIG. 6(d) shows data analysis of frequency shift upon the binding of antibody on BC- and BC-MAP-coated gold at a flow rate of  $10 \mu\text{L min}^{-1}$ . Values are the means from three independent experiments. Standard derivations are indicated in bar graphs.

FIG. 7(a) shows QCM response for the binding of BC and BC-MAP on gold at flow rate  $50 \mu\text{L min}^{-1}$ .

FIG. 7(b) shows QCM response for the binding of antibody on BC- and BC-MAP-coated gold at a flow rate of  $50 \mu\text{L min}^{-1}$  and.

FIG. 7(c) shows data analysis of frequency shift upon the binding of BC and BC-MAP on gold at a flow rate of  $50 \mu\text{L min}^{-1}$ . Values are the means from three independent experiments. Standard derivations are indicated in bar graphs.

FIG. 7(d) shows data analysis of frequency shift upon the binding of antibody on BC- and BC-MAP-coated gold at a flow rate of  $50 \mu\text{L min}^{-1}$ . Values are the means from three independent experiments. Standard derivations are indicated in bar graphs.

FIG. 8 shows schematic representation of antibody immobilization on diverse surfaces coated by BC-MAP.

#### DETAILED DESCRIPTION OF THE EMBODIMENTS

In the following detailed description, only certain exemplary embodiments of the present invention have been shown and described, simply by way of illustration.

As those skilled in the art would realize, the described embodiments may be modified in various ways, all without departing from the spirit or scope of the present invention.

Accordingly, the drawings and description are to be regarded as illustrative in nature and not restrictive.

Like reference numerals designate like elements throughout the specification.

In addition, unless explicitly described to the contrary, the word "comprise", and variations such as "comprises" or "comprising", will be understood to imply the inclusion of stated elements, but not the exclusion of any other elements.

In one aspect, the present invention relates to a fusion protein comprising protein A or a peptide derived from protein A and a mussel adhesive protein or a peptide derived from a mussel adhesive protein.

In a strategy to immobilize antibodies efficiently on surfaces in order to assess sensitivity and specificity of various immunosensors and immunoassays, the present invention intends to provide a fusion protein in which a mussel adhesive protein is fused to an antibody-binding protein, especially protein A derived from *Staphylococcus Aureus* or a peptide derived from protein A.

The proposed fusion protein of the present invention could be an invaluable linker material to facilitate an efficient immobilization of antibodies onto diverse solid supports.

The term "antibody-binding protein" as used herein refers to a protein that specifically recognizes and binds to Fc region of antibodies with high affinity, and realizes oriented-immobilization of antibodies on surfaces without reduced antibody activity. Preferably, the antibody-binding protein includes a protein A derived from *Staphylococcus aureus* or a peptide derived from protein A.

In one embodiment, the protein A has an amino acid sequence of SEQ ID No: 7.

In another embodiment, the peptide derived from protein A comprises a peptide or polypeptide comprising continuous subsequence of protein A that is capable of binding to Fc region of antibodies.

Preferably, the peptide derived from protein A comprises an amino acid sequence of 50 to 500 continuous amino acid residues, more preferably, 58 to 500 continuous amino acid residues of amino acid sequence of SEQ ID No:7. In a preferred embodiment, the peptide derived from protein A is a protein, a fragment, and a fusion peptide of at least one selected from the group consisting of domain A, domain B, domain C, domain D, domain E, or BC domain of the protein A.

In another embodiment, the peptide derived from protein A is a protein, a fragment, and a fusion peptide of at least one selected from group consisting of domain A having an amino acid sequence of SEQ ID No: 8, domain B having an amino acid sequence of SEQ ID No:9, domain C having an amino acid sequence of SEQ ID No: 10, domain D having an amino acid sequence of SEQ ID No: 11 and domain E having an amino acid sequence of SEQ ID No: 12.

In another embodiment, the peptide derived from protein A is BC domain of the protein A having an amino acid sequence of SEQ ID No: 13.

The term "mussel adhesive protein (MAP)" as used herein refers to a protein from marine mussels and is water-insoluble bioadhesives that adhere tightly to substrata. MAP is able to form a strong bond to diverse substrates including glass, metal, and plastics without any surface pretreatment. In the present invention, MAP acts as an immobilization agent for antibodies.

The peptide derived from the mussel adhesive protein in the present invention comprises a protein, a fragment, and a fusion peptide of at least one selected from mussel foot protein (FP)-1, FP-2, FP-3, FP-4, FP-5, and FP-6.

In one embodiment, the FP-5 comprises an amino acid sequence of SEQ ID No: 6.

The term "fusion protein" as used herein, generally indicates a polypeptide in which heterogenous polypeptides having different origins are linked, and in the present invention, refers to a polypeptide in which an antibody-binding protein and a MAP are linked.

In one embodiment, the protein A or a peptide derived from protein A is fused to the N-terminus or C-terminus of MAP or a peptide derived from MAP.

In another embodiment, the fusion protein comprises MAP having an amino acid sequence of SEQ ID No: 6, and protein A is fused to the N-terminus or C-terminus of MAP. Preferably, the fusion protein comprises BC domain of the protein A fused to the N-terminus or C-terminus of MAP having an amino acid sequence of SEQ ID No: 6. More preferably, the fusion protein comprises BC domain of the protein A having an amino acid sequence of SEQ ID No: 13 fused to the N-terminus or C-terminus of MAP having an amino acid

sequence of SEQ ID No: 6, which in turn forms a fusion protein having an amino acid sequence of SEQ ID No: 14.

This fusion protein may be obtained by chemical synthesis, or expression and purification processes of genetic recombination. Preferably, a hybrid gene, in which a gene sequence encoding an antibody-binding protein is linked to another gene sequence encoding a MAP, is expressed in a cell expression system. Then, the present fusion protein may be produced by transforming a host cell with a recombinant vector, and isolating and purifying a fusion protein expressed by the host cell.

In such a fusion protein, the antibody-binding protein and the MAP may be linked directly or by means of a connector, such as a linker. When a linker is used, it should not negatively affect the induction of antibody immobilization by the fusion protein.

Thus, in another aspect, the present invention relates to a polynucleotide encoding the fusion protein, a recombinant vector comprising the polynucleotide, and a transformed cell comprising the recombinant vector.

In a further aspect, the present invention relates to a method of preparing the fusion protein by transformed cell comprising the recombinant vector.

A process of producing the fusion protein of the present invention by genetic recombination comprises the following four steps.

The first step is to insert a gene encoding the fusion protein into a vector to construct a recombinant vector. Preferably, a gene encoding the fusion protein has a nucleotide acid sequence of SEQ ID No: 15, but not limited thereto.

A vector into which a gene encoding the fusion protein is introduced may be a plasmid, a virus, a cosmid, or the like. The recombinant vector includes a cloning vector and an expression vector. A cloning vector contains a replication origin, for example, a replication origin of a plasmid, phage or cosmid. The origin may be characterized as a "replicon" at which the replication of an exogenous DNA fragment attached thereto is initiated. An expression vector is developed for use in protein synthesis. A recombinant vector serves as a carrier for a foreign DNA fragment inserted thereto, which typically means a double-stranded DNA fragment. The term "foreign DNA" as used herein refers to DNA derived from a heterogeneous species, or a substantially modified form of native DNA from a homogenous species. In this case, a foreign gene is a specific target nucleic acid to be transcribed to encode a polypeptide.

The recombinant vector contains a target gene that is operably linked to transcription and translation expression regulatory sequences, which exert their functions to a selected host cell, in order to increase expression levels of the transfected gene in the host cell. The recombinant vector is a genetic construct that contains essential regulatory elements to which a gene insert is operably linked to be expressed in cells of an individual. Such a genetic construct is prepared using a standard recombinant DNA technique. The type of the recombinant vector is not specifically limited as long as the vector expresses a target gene in a variety of host cells including prokaryotes and eukaryotes, and functions to produce a target protein. However, a vector capable of mass-producing a foreign protein in a form similar to a native form while possessing a strong promoter to achieve strong expression of the target protein is preferred. The recombinant vector preferably contains at least a promoter, a start codon, a gene encoding a target protein, a stop codon and a terminator. The recombinant vector may further suitably contain DNA coding

a signal peptide, an enhancer sequence, 5'- and 3'-untranslational regions of a target gene, a selection marker region, a replication unit, or the like.

The second step is to transform a host cell with the recombinant vector and culture the host cell. The recombinant vector is introduced into a host cell to generate a transformant by a method described by Sambrook, J. et al., *Molecular Cloning, A Laboratory Manual* (2nd Ed.), Cold Spring Harbor Laboratory, 1. 74, 1989, the method including a calcium phosphate or calcium chloride/rubidium chloride method, electroporation, electroinjection, chemical treatments such as PEG treatment, and gene gun. A useful protein can be produced and isolated on large scale by culturing a transformant expressing the recombinant vector in a nutrient medium. Common media and culture conditions may be suitably selected according to host cells. Culture conditions, including temperature, pH of a medium and culture time, should be maintained suitably for a cell growth and mass production of a protein of interest. Host cells capable of being transformed with the recombinant vector according to the present invention include both prokaryotes and eukaryotes. Host cells having high introduction efficiency of DNA and having high expression levels of an introduced DNA may be typically used. Examples of host cells include known prokaryotic and eukaryotic cells such as *Escherichia* sp., *Pseudomonas* sp., *Bacillus* sp., *Streptomyces* sp., fungi and yeast, insect cells such as *Spodoptera frugiperda* 9 (Sf9), and animal cells such as CHO, COS 1, COS 7, BSC 1, BSC 40 and BMT 10. *E. coli* may be preferably used.

The third step is to induce expression as well as accumulation by the fusion protein. In the present invention, the inducer IPTG was used for the induction of peptide expression, and induction time was adjusted to obtain maximal protein yield.

The final step is to isolate and purify the fusion protein. Typically, a recombinantly produced peptide can be recovered from a medium or a cell lysate. When the peptide is in a membrane-bound form, it may be liberated from the membrane using a suitable surfactant solution (e.g., Triton-X 100) or by enzymatic cleavage. Cells used in the expression of the fusion protein may be destroyed by a variety of physical or chemical means, such as repeated freezing and thawing, sonication, mechanical disruption or a cell disrupting agent, and the fusion protein may be isolated and purified by commonly used biochemical isolation techniques (Sambrook et al., *Molecular Cloning: A laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, 1989; Deutscher, M., *Guide to Protein Purification Methods Enzymology*, Vol. 182. Academic Press. Inc., San Diego, Calif., 1990). Non-limiting examples of the biochemical isolation techniques include electrophoresis, centrifugation, gel filtration, precipitation, dialysis, chromatography (ion-exchange chromatography, affinity chromatography, immunosorbent affinity chromatography, reverse phased HPLC, gel permeation HPLC), isoelectric focusing, and variations and combinations thereof.

Protein A consists of five domains (D, E, A, B, and C) which bind to Fc region of immunoglobulin G. The present inventor constructed a novel immobilizing linker protein, in *Escherichia coli* by genetically fusing mussel adhesive protein (MAP) with two domains (B and C) of protein A for efficient immobilization of antibodies on diverse surfaces. BC domain without MAP was also biosynthesized as a comparative control (FIG. 1).

Based on direct surface coating analyses with color imaging, it was found that BC-MAP successfully was coated on diverse surfaces including glass slide, polypropylene, and titanium while BC domain alone exhibited partial coating.

Importantly, antibody was efficiently immobilized on BC-MAP-coated surfaces, and immobilized antibody interacted selectively with the corresponding antigen (FIG. 3 to FIG. 5).

Quartz crystal microbalance (QCM) analyses showed that BC-MAP has an excellent antibody binding ability compared to BC protein on the gold surface. These results demonstrated that unique strong and underwater adhesive properties of MAP, which can promote a direct and efficient immobilization of BC-MAP molecules on diverse surfaces without any additional surface treatments (FIG. 6 and FIG. 7). Thus, the proposed BC-MAP fusion protein could be a valuable linker material to facilitate an efficient immobilization of antibodies onto diverse solid supports.

In yet another aspect, the present invention relates to a biochip comprising a solid substrate to which the fusion protein is attached.

The biochip comprises a suitable solid substrate to which the fusion protein is attached. Protein A or a peptide derived from protein A can afford oriented antibody immobilization on surfaces of the solid substrate by binding Fc region of the antibody. MAP or a peptide derived from MAP can attach to a variety of surfaces and therefore, the fusion protein can attach to various surfaces of the solid substrate as well without loss of activity.

“Substrate” or “solid support” or other grammatical equivalents used herein refer to any materials that can be modified to contain discrete individual sites appropriate for the attachment or association of the antibody and is amenable to at least one detection method. As will be appreciated by those in the art, the number of possible substrates are extensive, and include, but are not limited to, glass and modified or functionalized glass, plastics (including acrylic, polystyrene and copolymers of styrene and other materials, polypropylene, polyethylene, polybutylene, polyurethanes, Teflon, tetrafluoroethylene, etc.), polysaccharides, nylon or nitrocellulose, resins, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses, etc. In general, the substrates allow optical detection and do not appreciably fluoresce.

The substrate provides a place for antigen-antibody interactions as well as functions as a solid support. The size or shape of the substrate and sites of immobilizing antibody on the substrate would be diverse according to the purpose of the analysis or apparatus such as spotting machine or scanner.

In another aspect, the present invention relates to a method for detecting antigen in a biological sample using the biochip.

Preferably, the detecting method of the present invention comprises the step of immobilizing an antibody which specifically binds to the target antigen on the solid substrate of the biochip, adding a biological sample comprising the target antigen to the biochip, and detecting the target antigen by measuring antigen-antibody interactions.

The first step is to immobilize an antibody which specifically binds to the target antigen on the solid substrate of the biochip. In the present invention, an antibody is fixed to a solid substrate by the fusion protein to facilitate washing and isolation of the antigen-antibody complex. In the fusion protein, protein A or a peptide derived from protein A can afford oriented antibody immobilization on the solid substrate by binding Fc region of the antibody, and MAP can attach to a variety of surfaces. Therefore, the fusion protein can attach to various solid substrates as well without loss of activity.

The term “antibody” refers to an immunoglobulin molecule having a specific structure that binds specifically to a molecule comprising the antigen used for synthesizing the antibody or with an antigen closely related to it. As used herein, the term “antibody” broadly includes full length anti-

bodies and may also include certain antibody fragments thereof. Also included are monoclonal and polyclonal antibodies, multivalent and monovalent antibodies, multispecific antibodies (for example bi-specific antibodies), chimeric antibodies, human antibodies, and humanized antibodies.

As used herein, an “antigen-binding fragment” or “antibody fragment” means a portion of the intact antibody that preferably retains most or all, or minimally at least one of, the normal functions of that antibody fragment. An antibody fragment, for example, may comprise an Fc region that retains all or most or some of the functions of the corresponding Fc region in the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub> and Fv fragments, linear antibodies, diabodies, single chain antibodies (ScFV) and multispecific antibodies.

As used herein, a “monoclonal antibody” means an antibody that is a highly specific antibody directed against a single target antigen. A monoclonal antibody may be obtained from a population of homogenous or substantially homogenous antibodies wherein each monoclonal antibody is identical and/or bind the same epitope, except for natural mutations which may occur in minor amounts.

The second step is to add a biological sample comprising the target antigen to the biochip.

Biological sample as used herein means any sample derived from a subject to be detected. The sample may be any sample known in the art in which the target antigen can be detected by antigen-antibody interactions. Included are any body fluids such as plasma, blood, saliva, interstitial fluid, serum, urine, synovial, cerebrospinal, lymph, seminal, amniotic, pericardial fluid and ascites, but not limited thereto.

The term “target antigen” as used herein refers to any solid or non-solid material capable of binding to an antibody immobilized on the biochip or a fragment or variant thereof. In one embodiment the term refers to any natural or non-natural molecule that binds to an antibody immobilized on the biochip or a fragment or variant thereof. Examples of target antigen include proteins, peptides, carbohydrates, lipids, and small molecule compounds, but not limited thereto.

The third step is to detect the target antigen by measuring antigen-antibody interactions.

To facilitate detection, antibodies and fragments herein may be labelled with detectable markers such as fluorescent, bioluminescent, and chemiluminescent compounds, as well as radioisotopes, magnetic beads and affinity labels (e.g. biotin and avidin). Examples of labels which permit indirect measurement of binding include enzymes where the substrate may provide for a coloured fluorescent product. Suitable enzymes include horseradish peroxidase, alkaline phosphatase, malate dehydrogenase and the like. Fluorochromes (e.g. Texas Red, fluorescein, phycobiliproteins, and phycoerythrin) can be used with a fluorescence activated cell sorter. Labelling techniques well known in the art can be utilized.

An antigen-antibody interaction can be detected by any means known in the art. Most commonly, antigen-antibody interactions herein are detected using an assay such as ELISA or RIA, competitive binding assays, sandwich assays, non-competitive assays, fluoroimmunoassay, immunofluorometric assay, or immunoradiometric assays, luminescence assays, or chemiluminescence assays.

The present invention is further explained in more detail with reference to the following examples. These examples, however, should not be interpreted as limiting the scope of the present invention in any manner.

## Construction of Expression Vectors

## 1.1: Construction of Expression Vector for BC-MAP Fusion Protein

The gene coding of MAP fp-5 was amplified by a polymerase chain reaction (PCR) from the previously constructed vector pMDG05. The forward primer (F1: 5'-GGGC-CATGGGGTGGCGGAGGGAGCTCTGAA-GAATATAAAG-3', SEQ ID No:1) was designed to contain a NcoI restriction enzyme site and the sequence encoding a GGGG amino acid linker. The backward primer (B1: 5'-CGCGCTCGAGGCTGCTGCCGCCAT-AATATTTTTATAG-3', SEQ ID No:2) was designed to contain a XhoI restriction enzyme site. The PCR product was digested with two restriction enzymes of NcoI and XhoI (Fermentas, Burlington, Ontario, Canada). The digested DNA fragment was ligated with a pET22b(+) vector (Novagen, San Diego, Calif., USA), which predigested with NcoI and XhoI, by using a ligation kit (TaKaRa, Shiga, Japan) to make the plasmid pET-MAP. The gene, encoding two domains (B and C domain) of protein A, was amplified by PCR from the genomic DNA of *Staphylococcus aureus* ATCC 6538 (B. H. Hwang, H. J. Cha, *Biotechnol. Bioeng.* 2010, 106, 183). The forward primer (F2: 5'-GCGC-CATATGGATAACAAATTCAACAAAGAACAAC-3', SEQ ID No:3) was designed to contain a NdeI restriction enzyme site and the backward primer (B2: 5'-GCGCCCATG-GTTTTGGTGTCTTGAGCATCGTTTAGC-3', SEQ ID No:4) was designed to contain a NcoI restriction enzyme site. The PCR product was digested with two restriction enzymes of NdeI (Fermentas, Burlington, Ontario, Canada) and NcoI. The digested PCR was ligated with pET-MAP, which predigested with NdeI and NcoI, resulting in the plasmid pET-BC-MAP (FIG. 2(a)).

## 1.2: Construction of Expression Vector for Sole BC Protein

For the construction of BC protein as control protein, the corresponding gene was amplified by PCR with the forward primer (F2) and backward primer (B3: 5'-GCGCCTC-GAGTTTTGGTGTCTTGAGCATCGTTTAGC-3', SEQ ID No:5). The PCR product was ligated into the NdeI and XhoI sites of pET22b(+) to make pET-BC (FIG. 2a).

## Example 2

## Expression and Purification of BC-MAP and Sole BC Proteins

Each plasmid containing pET-BC-MAP or pET-BC was transformed into *Escherichia coli* BL21 (DE3). The transformants were grown to 0.8-0.9 OD<sub>600</sub> at 37° C. in shake flasks containing of 400 mL of Luria-Bertani (LB) medium (0.5% yeast extract, 1% tryptophan, and 1% NaCl) with 50 µg mL<sup>-1</sup> ampicillin. The expression was induced by adding isopropyl β-D-thiogalactopyranoside (IPTG) to a final concentration of 1 mM. The transformed cells were grown for an additional 10 h at 37° C. The cells were harvested by centrifugation at 4000 g for 10 min. The cell pellets were resuspended in lysis buffer (50 mM NaH<sub>2</sub>PO<sub>4</sub>, 300 mM NaCl, 10 mM imidazole, pH 8.0) and disrupted with a sonic dismembrator (Fisher scientific, USA) for 10 min at 50% power (3 sec pulse on and 2 sec pulse off). The soluble and insoluble fractions were then separated by centrifugation at 9000 g for 30 min at 4° C. The soluble fractions were loaded into a Ni-NTA column (Qiagen, Valencia, Calif., USA). BC and BC-MAP were eluted with elution buffer (50 mM NaH<sub>2</sub>PO<sub>4</sub>, 300 mM NaCl, 250 mM

imidazole, pH 8.0) and desalted by a PD-10 column (GE Healthcare, Sweden). The protein concentration was determined by Bradford method using bovine serum albumin (BSA) (Sigma-Aldrich, St. Louis, Mo., USA) as a standard.

## Example 3

## MALDI-TOF MS Analysis

Matrix-assisted laser desorption ionization mass spectrometry with time-of-flight (MALDI-TOF MS) analysis was performed on a 4700 Proteomics Analyzer (AB Sciex) in the positive ion linear mode. A-cyano-4-hydroxycinnamic acid in 50% acetonitrile and 0.1% trifluoroacetic acid was used as the matrix solution. Sample were diluted 1:20 with matrix solution, and 1 µL of the mixture was spotted onto the MALDI sample target plates and evaporated using a vacuum pump. Spectra were obtained in the mass range between 5,000 and 25,000 Da with ~2000 laser shots. Internal calibration was performed using mix 3 of sequezyme peptide mass standard kit (AB Sciex).

## Example 4

## Direct Surface Coating Analysis

Three types of surfaces (glass (Marienfeld GmbH & Co., Lauda-Königshofen, Germany), titanium (Alfa Aesar, Ward Hill, Mass., USA), and polypropylene) were used for direct surface coating analyses. Glass slide was cleaned with hot piranha solution (10% (v/v) H<sub>2</sub>O<sub>2</sub>, 30% (v/v) H<sub>2</sub>SO<sub>4</sub>) and was thoroughly rinsed with ethanol and dH<sub>2</sub>O. Titanium was treated with 40% sulfuric acid at room temperature for 15 min and rinsed extensively with dH<sub>2</sub>O. And then, the titanium was boiled in dH<sub>2</sub>O and washed with dH<sub>2</sub>O and acetone. The titanium was dried under N<sub>2</sub> gas. BC-MAP (2 mg mL<sup>-1</sup>) and BC (2 mg mL<sup>-1</sup>) were coated on each surface and the surfaces were dried for complete immobilization. After washing with PBS (pH 7.5) at 200 rpm shaker for 5 min, the surfaces was treated with 5% non-fat milk solution for 1 h at room temperature, followed by washing with TTBS (20 mM Tris-HCl, 500 mM NaCl, 0.0005% Tween 20 (pH 7.5)) (5 min×3 times). Anti-rabbit antibody conjugated with alkaline phosphatase solution (as a target antibody) (Sigma-Aldrich) was added onto BC-MAP-coated surfaces for 1 h at room temperature. To analyze antibody-binding capability of BC-MAP on the surfaces, antibody-immobilized surfaces were treated with NBT/BCIP solution (Bio Basic Inc., Canada) for 20 min after washing with sequential TTBS (5 min×2 times) and TBS (20 mM Tris-HCl, 500 mM NaCl (pH 7.5)) (5 min×1 time).

For analyzing immobilizing degree on BC-MAP-coated glass slide according to antibody concentrations, various antibody concentrations (20 µg mL<sup>-1</sup>-200 ng mL<sup>-1</sup>) were used.

## Example 5

## Antigen-Antibody Interaction on BC-MAP-Coated Glass Slide

BC-MAP- and BC-coated glass slides were constructed as described above. The glass slide was immersed onto anti-alkaline phosphatase (Sigma-Aldrich) solution (100 µg mL<sup>-1</sup>), followed by washing with TTBS (5 min×3). Alkaline phosphatase (antigen) solutions with different concentrations (100, 80, 60, 40, 20, and 10 µg mL<sup>-1</sup>) were added onto the antibody immobilized surface for 1 h at room temperature.

After washing as described above, the surface was reacted with NBT/BCIP solution for 1 h

#### Example 6

##### Selectivity of Antibodies Functionalized BC-MAP-Coated Glass Slide

BC-MAP-coated glass slide was constructed (FIG. 5c). After washing and blocking, antibody solutions (anti-alkaline phosphatase and anti-human transferrin (Sigma-Aldrich), 2 mg mL<sup>-1</sup>) were added onto each defined block by Gene Frame® (Thermo Scientific, San Jose, Calif., USA) and then the glass slide was immersed with alkaline phosphatase solution (100 µg mL<sup>-1</sup>) after washing with sequential TTBS/TBS. Color development processes were performed by same procedure described above except reaction time (90 min) with NBT/BCIP solution.

#### Example 7

##### Quartz Crystal Microbalance (QCM) Analysis

Gold-coated quartz crystals (5 MHz, Stanford Research System, Sunnyvale, Calif., USA) were cleaned by UV irradiation for 20 min before a syringe pump (WPI, Sarasota, Fla., USA) equipped with a low-pressure liquid chromatography injector valve and a 250 µL injection loop (Upchurch Scientific, Oak Harbor, Wash., USA) was used to inject either sample solution or carrier solution (1×PBS, pH 7.4) into the flow cell at a flow rate of 10 or 50 µL min<sup>-1</sup>. After rinsing with DW and drying with ultra-high purity N<sub>2</sub>, the crystals were mounted inside QCM flow modules. PBS (1×) was then injected into the flow module until stable baseline *f* shifts were achieved. A total of 250 µL of BC-MAP (0.2 mg mL<sup>-1</sup>), BC (0.12 mg mL<sup>-1</sup>), or MAP (0.2 mg mL<sup>-1</sup>) was injected into the flow module for attachment to the gold surface. After rinsing with PBS until the frequency shift was stable, the surface was treated with 250 µL BSA solution (1 mg mL<sup>-1</sup>) to block non-specific interactions. This step was followed by a PBS rinse to remove unattached BSA. Then 250 µL of polyclonal anti-DsRed antibody solution (20 ng mL<sup>-1</sup>; Santa Cruz Biotechnology, Santa Cruz, Calif., USA) was injected into the flow module. The adsorbed mass was calculated using the Sauerbrey equation (1):

$$\Delta f = -\frac{2f_0^2}{A\sqrt{\rho_q\mu_q}}\Delta m \quad (1)$$

where  $\Delta f$  is the change in fundamental frequency of the coated crystal,  $f_0$  is the resonant frequency of the unloaded crystal (5 MHz),  $A$  is the active area of the crystal between electrodes (1.267 cm<sup>2</sup>),  $\rho_q$  is the density of quartz (2.648 g cm<sup>-3</sup>),  $\mu_q$  is the shear modulus of quartz (2.947×10<sup>11</sup> g cm<sup>-1</sup> s<sup>-2</sup>), and  $\Delta m$  is the change in mass per unit surface area.

#### Experimental Example 1

##### Expression and Purification of BC-MAP and BC

Protein A consists of five domains (D, E, A, B, and C) which bind to Fc region of immunoglobulin G (IgG). We constructed novel immobilizing linker by fusion of BC domain of protein A with MAP. BC domain without MAP was also constructed as a comparative control. To produce

recombinant BC-MAP and sole BC in *E. coli*, each BC-MAP and BC gene was cloned into expression vector containing His<sub>6</sub> affinity purification tag sequence and an inducible T7 promoter (FIG. 1a). Both BC-MAP and BC were majorly expressed as soluble form in *E. coli* (data not shown).

To investigate potential activity of BC-MAP as immobilizing agent for efficient immobilization of antibodies on diverse surfaces, recombinant BC-MAP and BC proteins were purified by immobilized metal affinity chromatography (IMAC). Each actual molecular weight of BC-MAP and BC from sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE; FIG. 1b) and matrix-assisted laser desorption/ionization-time of flight mass spectrometry (MALDI/TOF-MS) was similar to the calculated molecular weight of ~23.5 kDa and ~14.2 kDa, respectively. Finally, we obtained high purity of BC-MAP (~90.7%) and BC (~97.0%) by one-step affinity purification (FIG. 1b).

#### Experimental Example 2

##### Antibody Immobilization on BC-MAP-Coated Surfaces

In order to investigate antibody-immobilizing activity of BC-MAP, we performed direct coatings on diverse surfaces with color image analysis. We used three different surfaces (glass slide, titanium, and polypropylene) and anti-rabbit IgG conjugated with alkaline phosphatase as a model antibody. Glass slide, polymer, and titanium have been utilized as solid supports for protein arrays as well as immunosensors. Sole BC protein was used a comparative control. First, BC-MAP- and BC-coated surfaces were severely washed with PBS at 200 rpm shaker by changing washing time (5, 10, and 15 min) to analyze coated pattern along with washing time. After washing, the surfaces were stained using Coomassie blue. When the surfaces were washed for 5 min, sole BC was completely disappeared on all surfaces. However, most BC-MAP was not removed on all surfaces under 15 min washing time (data not shown). Note that sole BC was removed on the surfaces even under gentle washing condition (data not shown). This implies that surface binding of BC-MAP is much more stable and stronger than interaction between BC and the surfaces owing to the unique property of MAP. This result was consistent with our previous result that MAP have strong binding ability to diverse surfaces such as glass slide, poly(methyl methacrylate) plate, polystyrene plate, and aluminum. Using this adhesive characteristic of MAP, BC-MAP could be used as immobilizing agent for efficient immobilization of antibody on diverse surfaces without any additional steps. In the subsequent experiments, we set washing time as 5 min for protein coating.

After washing and blocking by 5% non-fat milk, BC-MAP-coated the surfaces was immersed in antibody solution, followed by washing with TTBS/TBS buffer solution and reacting with NBT/BCIP solution. As expected, the model antibody, anti-rabbit IgG, was successfully immobilized on all three BC-MAP-coated surfaces (FIG. 2). This result indicates that BC-MAP-based immobilization of antibody onto diverse surfaces does not require any additional steps including chemical modification and BC domain of BC-MAP interacts specifically with Fc region of IgG through proper orientation of BC-MAP through binding of MAP domain onto surfaces and exposure of BC domain toward IgG-containing solution. If BC domain of BC-MAP was immobilized on surfaces but not exposed outward, surface-immobilized BC-MAP would be easily removed during washing step. Thus, MAP domain of BC-MAP played a criti-

cal role for facile antibody immobilization on diverse surfaces and exposure of functional BC domain.

### Experimental Example 3

#### Antigen-Antibody Interaction on BC-MAP-Coated Glass Slide

In diagnosis, quantitative analysis is one of important issues. Thus, before analyzing antigen-antibody interaction on BC-MAP-coated surface, we investigated immobilization degree according to antibody concentrations. Among three surfaces, glass slide was used in further subsequent experiments because it is commonly utilized as solid surface in antibody array. While the spot shapes on hydrophobic polypropylene were somewhat clean, the shapes on glass slide were not relatively uniform because hydrophilic surface leads to larger printed spots. FIG. 3 shows antibody-BC-MAP interaction in a range of antibody concentrations from 200 ng mL<sup>-1</sup> to 20 μg mL<sup>-1</sup>. We found that the relative signal intensities of color spots increased according to applied antibody concentrations (FIG. 3a). We determined that detection limit and saturation point were corresponded to 200 ng mL<sup>-1</sup> and 10 μg mL<sup>-1</sup>, respectively, at 20 min fixed color development time (FIG. 3b).

Next, we used anti-alkaline phosphatase IgG as a model antibody and alkaline phosphatase as a model antigen to investigate interaction of the immobilized antibody on BC-MAP-coated glass slide with the corresponding antigen and potency of quantitative analysis. Alkaline phosphatase solutions with different concentrations were applied on the anti-alkaline phosphatase IgG-immobilized BC-MAP-coated glass slide. As shown in FIG. 4, the signal intensities increased along with increasing antigen concentrations. With considering background intensity value (11.18±1.70) on bare glass slide and fixed color development time, we found that at least 20 μg mL<sup>-1</sup> antigen could be detected on the glass slide. Importantly, linear correlation between signal intensity and

antigen concentration was shown with high coefficient of determination (R<sup>2</sup>) of over 0.97 at the range of 20-100 μg mL<sup>-1</sup> antigen concentration (FIG. 4b). The result indicates that antibody array using BC-MAP as immobilizing agent can be used for quantitative analysis of antigen-antibody interaction. Furthermore, the selectivity of this BC-MAP-based immunosensor system was investigated by loading two antibody solutions (anti-alkaline phosphatase IgG as a positive antibody and human transferrin IgG as a negative control) onto each defined blocks and then reacting with alkaline phosphatase solution (FIG. 5a). FIG. 5b shows that the antibody array using BC-MAP linker was very specific to the corresponding antigen without detection of cross-reaction, demonstrating the good selectivity of the system.

### Experimental Example 4

#### QCM Analysis of BC-MAP Material

We also investigated the function of BC-MAP as a linker molecule for efficient immobilization of antibody using quartz crystal microbalance (QCM). We examined the ability of BC-MAP at two different conditions. First, BC-MAP was tested at relatively low flow rate on bare gold surface. Second, relatively high flow rate was used to identify markedly the capability of BC-MAP although it was reported that conventional QCM design in which a circular resonator is sandwiched between two O-rings shows unstable signal at high flow rate. BC-MAP and sole BC (control) were immobilized on the gold surfaces, and anti-DsRed antibodies were applied on BC-MAP- and BC-coated gold surfaces, respectively. QCM plots for the binding events between protein and gold and between antibody and protein are shown in FIGS. 6a-b and FIG. 7a-b, respectively, under two different conditions. FIG. 6a and FIG. 7a shows QCM response for the binding of BC-MAP and sole BC onto gold surfaces at 10 μL min<sup>-1</sup> and 50 μL min<sup>-1</sup>, respectively. FIG. 6b and FIG. 7b represents QCM response for the binding of anti-DsRed antibody onto BC-MAP- and BC-coated gold surfaces at 10 μL min<sup>-1</sup> and 50 μL min<sup>-1</sup>, respectively. We compared QCM responses upon the bindings of BC-MAP and BC on gold surfaces (FIG. 6c & FIG. 7c). At the 10 μL min<sup>-1</sup> flow rate condition, the changed frequency values of BC and BC-MAP on gold were 24.0±0.9 and 31.1±7.6 Hz, which corresponded to the similar bindings of 1.8×10<sup>13</sup> and 1.4×10<sup>13</sup> molecules cm<sup>-2</sup> on the surface, respectively (Table 1). Different with the result of direct surface coating analysis, sole BC also showed a similar binding level with BC-MAP on gold surface in QCM system. This was resulted from smooth washing condition (relatively low flow rate) used in QCM system. At the higher flow rate condition (50 μL min<sup>-1</sup>), the frequency shift values of BC and BC-MAP on gold were 6.5±2.0 and 16.2±0.8 Hz, indicating about 73% and 47% decreases of these on gold at low flow rate, respectively (Table 1). We confirmed that binding of BC-MAP on gold surface was more stable and stronger than that of sole BC.

TABLE 1

Flow rate [μL min <sup>-1</sup> ]	Sample	Protein immobilization [Hz]	Surface density [molecule cm <sup>-2</sup> ]	Antibody binding [Hz]	Surface density [molecule cm <sup>-2</sup> ]	Binding efficiency [%]
10	BC	24.0 ± 0.9	1.8 × 10 <sup>13</sup>	6.1 ± 3.0	4.3 × 10 <sup>11</sup>	2.4
	BC-MAP	31.1 ± 7.6	1.4 × 10 <sup>13</sup>	28.2 ± 14.1	2.0 × 10 <sup>12</sup>	14.2
50	BC	6.5 ± 2.0	4.9 × 10 <sup>12</sup>	1.4 ± 0.6	9.9 × 10 <sup>10</sup>	2.0
	BC-MAP	16.2 ± 0.8	7.3 × 10 <sup>12</sup>	21.7 ± 0.5	1.5 × 10 <sup>12</sup>	20.5

Next, we compared QCM responses upon the bindings of antibody on BC-MAP and sole BC-coated gold surfaces (FIG. 6d & FIG. 7d). Nonspecific interactions were blocked by treatment of BSA before addition of anti-DsRed antibody. At the lower flow rate condition 10 μL min<sup>-1</sup>, the antibody binding signal to BC-MAP-coated gold surface was 28.2±14.1 Hz, whereas the value to BC-coated gold surface was 6.1±3.0 Hz (Table 1). By considering the molecular weights and the mass changes due to adsorption (Δm) of the antibody (150 kDa), BC-MAP (23.5 kDa), and BC (14.2 kDa), we determined that BC-coated gold surface had only 2.4% of immobilized BC which was involved in interaction with the Fc region of antibody (Table 1). We suspect that active binding sites of BC were not exposed toward the anti-

body solution due to non-oriented physical adsorption on gold surface. Importantly, over 14.2% of immobilized BC-MAP interacted with the antibody on gold surface and this was about 6-fold more antibody binding compared to the sole BC molecule. At the higher flow rate (50  $\mu\text{L min}^{-1}$ ), the antibody binding signals to sole BC- and BC-MAP-coated gold surfaces were  $1.4\pm 0.6$  and  $21.7\pm 0.5$  Hz, respectively (Table 1). As a result, the BC-MAP showed increase of binding efficiency of 20.5% while the binding efficiency of sole BC maintained at similar low level of  $\sim 2.0\%$  (Table 1). We speculated that this is because BC domain of BC-MAP on gold at high flow rate well exposed the antibody solution compared this on gold at low flow rate by immobilizing MAP domain of BC-MAP onto the surface but preventing BC

domain of BC-MAP from interacting with the surface. Two explanations might be possible for higher antibody binding efficiency of BC-MAP. First, active binding sites of BC domain in BC-MAP molecule were effectively exposed outward. Second is that MAP domain in BC-MAP was also involved in antibody binding. To investigate possibility for direct involvement of MAP in antibody binding, we performed antibody binding on sole MAP-coated gold surface. But, we found that MAP was not involved in antibody binding after BSA blocking although much larger quantity of MAP was used. Thus, we can exclude the possibility about direct involvement of MAP in antibody binding and conclude that BC-MAP binds to surface with oriented manner for BC domain.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 15

<210> SEQ ID NO 1

<211> LENGTH: 41

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: forward primer for MAP protein

<400> SEQUENCE: 1

gggccatgg ggtggcggag ggagctctga agaataaaa g 41

<210> SEQ ID NO 2

<211> LENGTH: 38

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: reverse primer for MAP protein

<400> SEQUENCE: 2

cgcgctcgag gctgctgcc ccataatatt ttttag 38

<210> SEQ ID NO 3

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: forward primer for BC protein

<400> SEQUENCE: 3

gcgccatag gataacaaat tcaacaaga acaac 35

<210> SEQ ID NO 4

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: reverse primer for BC protein

<400> SEQUENCE: 4

gcgccatgg ttttggctgct tgagcatcgt ttagc 35

<210> SEQ ID NO 5

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: reverse primer for BC protein

<400> SEQUENCE: 5

gcgcctcgag ttttggctgct tgagcatcgt ttagc 35

-continued

---

<210> SEQ ID NO 6  
 <211> LENGTH: 75  
 <212> TYPE: PRT  
 <213> ORGANISM: Mytilus galloprovincialis

<400> SEQUENCE: 6

Ser Ser Glu Glu Tyr Lys Gly Gly Tyr Tyr Pro Gly Asn Thr Tyr His  
 1 5 10 15  
 Tyr His Ser Gly Gly Ser Tyr His Gly Ser Gly Tyr His Gly Gly Tyr  
 20 25 30  
 Lys Gly Lys Tyr Tyr Gly Lys Ala Lys Lys Tyr Tyr Tyr Lys Tyr Lys  
 35 40 45  
 Asn Ser Gly Lys Tyr Lys Tyr Leu Lys Lys Ala Arg Lys Tyr His Arg  
 50 55 60  
 Lys Gly Tyr Lys Lys Tyr Tyr Gly Gly Ser Ser  
 65 70 75

<210> SEQ ID NO 7  
 <211> LENGTH: 500  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 7

Met Lys Lys Lys Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile  
 1 5 10 15  
 Ala Ser Val Thr Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro  
 20 25 30  
 Ala Ala Asn Ala Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr  
 35 40 45  
 Gln Val Leu Asn Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe  
 50 55 60  
 Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly  
 65 70 75 80  
 Glu Ala Gln Lys Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln  
 85 90 95  
 Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu  
 100 105 110  
 Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser  
 115 120 125  
 Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys  
 130 135 140  
 Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys  
 145 150 155 160  
 Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn  
 165 170 175  
 Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser  
 180 185 190  
 Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln  
 195 200 205  
 Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe  
 210 215 220  
 Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly  
 225 230 235 240  
 Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu  
 245 250 255

-continued

---

Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Ala Asp Asn  
 260 265 270

Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu  
 275 280 285

Pro Asn Leu Thr Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys  
 290 295 300

Asp Asp Pro Ser Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys Leu  
 305 310 315 320

Asn Asp Ala Gln Ala Pro Lys Glu Glu Asp Asn Asn Lys Pro Gly Lys  
 325 330 335

Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys  
 340 345 350

Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp Asn Lys Lys Pro Gly Lys  
 355 360 365

Glu Asp Asn Lys Lys Pro Gly Lys Glu Asp Asn Lys Lys Pro Gly Lys  
 370 375 380

Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp Asn Lys Lys Pro Gly Lys  
 385 390 395 400

Glu Asp Gly Asn Gly Val His Val Val Lys Pro Gly Asp Thr Val Asn  
 405 410 415

Asp Ile Ala Lys Ala Asn Gly Thr Thr Ala Asp Lys Ile Ala Ala Asp  
 420 425 430

Asn Lys Leu Ala Asp Lys Asn Met Ile Lys Pro Gly Gln Glu Leu Val  
 435 440 445

Val Asp Lys Lys Gln Pro Ala Asn His Ala Asp Ala Asn Lys Ala Gln  
 450 455 460

Ala Leu Pro Glu Thr Gly Glu Glu Asn Pro Phe Ile Gly Thr Thr Val  
 465 470 475 480

Phe Gly Gly Leu Ser Leu Ala Leu Gly Ala Ala Leu Leu Ala Gly Arg  
 485 490 495

Arg Arg Glu Leu  
 500

<210> SEQ ID NO 8  
 <211> LENGTH: 58  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 8

Ala Asp Asn Asn Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile  
 1 5 10 15

Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln  
 20 25 30

Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ser Glu Ala  
 35 40 45

Lys Lys Leu Asn Glu Ser Gln Ala Pro Lys  
 50 55

<210> SEQ ID NO 9  
 <211> LENGTH: 58  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 9

Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile  
 1 5 10 15

-continued

---

Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln  
                   20                                  25                                  30

Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala  
                   35                                  40                                  45

Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys  
           50                                  55

<210> SEQ ID NO 10  
 <211> LENGTH: 58  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 10

Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile  
   1                  5                                  10                                  15

Leu His Leu Pro Asn Leu Thr Glu Glu Gln Arg Asn Gly Phe Ile Gln  
                   20                                  25                                  30

Ser Leu Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu Ala Glu Ala  
                   35                                  40                                  45

Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys  
           50                                  55

<210> SEQ ID NO 11  
 <211> LENGTH: 59  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 11

Ala Asp Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile  
   1                  5                                  10                                  15

Ile Asn Met Pro Asn Leu Asn Glu Ala Gln Asn Arg Asn Gly Phe Ile  
                   20                                  25                                  30

Gln Ser Leu Lys Ile Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu  
                   35                                  40                                  45

Ala Lys Lys Leu Asn Glu Ser Gln Ala Pro Lys  
           50                                  55

<210> SEQ ID NO 12  
 <211> LENGTH: 59  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 12

Ala Asp Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr Glu Ile  
   1                  5                                  10                                  15

Ile Asn Met Pro Asn Leu Asn Glu Glu Gln Asn Arg Asn Gly Phe Ile  
                   20                                  25                                  30

Gln Ser Leu Lys Ile Asp Pro Ser Gln Ser Ala Asn Leu Leu Ser Glu  
                   35                                  40                                  45

Ala Lys Lys Leu Asn Glu Ser Gln Ala Pro Lys  
           50                                  55

<210> SEQ ID NO 13  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 13

Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile

-continued

---

```

1           5           10           15
Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln
      20           25           30
Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala
      35           40           45
Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Ala Asp Asn Lys Phe Asn
      50           55           60
Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu
      65           70           75           80
Thr Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro
      85           90           95
Ser Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala
      100          105          110
Gln Ala Pro Lys
      115

```

```

<210> SEQ ID NO 14
<211> LENGTH: 206
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Fusion protein

```

```

<400> SEQUENCE: 14

```

```

Met Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu
  1           5           10           15
Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile
      20           25           30
Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu
      35           40           45
Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Ala Asp Asn Lys Phe
      50           55           60
Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn
      65           70           75           80
Leu Thr Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp
      85           90           95
Pro Ser Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys Leu Asn Asp
      100          105          110
Ala Gln Ala Pro Lys Pro Trp Gly Gly Gly Gly Ser Ser Glu Glu Tyr
      115          120          125
Lys Gly Gly Tyr Tyr Pro Gly Asn Thr Tyr His Tyr His Ser Gly Gly
      130          135          140
Ser Tyr His Gly Ser Gly Tyr His Gly Gly Tyr Lys Gly Lys Tyr Tyr
      145          150          155          160
Gly Lys Ala Lys Lys Tyr Tyr Tyr Lys Tyr Lys Asn Ser Gly Lys Tyr
      165          170          175
Lys Tyr Leu Lys Lys Ala Arg Lys Tyr His Arg Lys Gly Tyr Lys Lys
      180          185          190
Tyr Tyr Gly Gly Ser Ser Leu Glu His His His His His His
      195          200          205

```

```

<210> SEQ ID NO 15
<211> LENGTH: 618
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: polynucleotide encoding fusion protein

```

-continued

<400> SEQUENCE: 15

```

atggcggata acaaatcaa caaagaacaa caaaatgctt tctatgaaat cttacattta    60
cctaaacttaa acgaagaaca acgcaatggt ttcatccaaa gcttaaaaga tgaccaaacg    120
caaagcgcta acottttagc agaagctaaa aagctaaatg atgcacaagc accaaaagct    180
gacaacaaat tcaacaaaga acaacaaaat gctttctatg aaattttaca tttacctaac    240
ttaactgaag aacaacgtaa cggcttcctc caaagcctta aagacgatcc ttcagtgage    300
aaagaaattt tagcagaagc taaaagccta aacgatgctc aagcaccaaa accatggggg    360
ggcggaggga gctctgaaga atataaaggc ggctattatc cgggcaacac ctaccattat    420
cattctggcg gcagctatca tggtctggcg tatcatggcg gctataaagg caaatattat    480
ggcaaagcga aaaaatatta ttataaatat aaaaacagcg gcaaatataa atatctgaaa    540
aaagcgagaa aatatcatag aaaaggctat aaaaaatatt atggcggcag cagcctcgag    600
caccaccacc accaccac                                     618
    
```

What is claimed is:

1. A fusion protein comprising protein A or a peptide derived from protein A and a mussel adhesive protein or a peptide derived from a mussel adhesive protein.
2. The fusion protein according to claim 1, wherein the peptide derived from a mussel adhesive protein is a protein, a fragment or a fusion peptide of at least one selected from group consisting of mussel foot protein (FP)-1, FP-2, FP-3, FP-4, FP-5 and FP-6.
3. The fusion protein according to claim 2, wherein the peptide derived from a mussel adhesive protein is FP-5 having an amino acid sequence of SEQ ID No: 6.
4. The fusion protein according to claim 1, wherein the protein A has an amino acid sequence of SEQ ID No: 7.
5. The fusion protein according to claim 1, wherein the peptide derived from protein A comprises 50 to 500 continuous amino acid residues of amino acid sequence of SEQ ID No:7.
6. The fusion protein according to claim 1, wherein the peptide derived from protein A is a protein, a fragment, and a fusion peptide of at least one selected from the group consisting of domain A, domain B, domain C, domain D, domain E, and BC domain of the protein A.
7. The fusion protein according to claim 6, wherein the peptide derived from protein A is a protein, a fragment, and a fusion peptide of at least one selected group from the domain A having an amino acid sequence of SEQ ID No: 8, domain B having an amino acid sequence of SEQ ID No:9, domain C having an amino acid sequence of SEQ ID No: 10, domain D having an amino acid sequence of SEQ ID No: 11 and domain E having an amino acid sequence of SEQ ID No: 12.
8. The fusion protein according to claim 6, wherein the peptide derived from protein A is BC domain having an amino acid sequence of SEQ ID No: 13.
9. The fusion protein according to claim 1, wherein the protein A or the peptide derived from protein A is fused to the N-terminus or C-terminus of the mussel adhesive protein or the peptide derived from a mussel adhesive protein.

10. The fusion protein according to claim 1, which has an amino acid sequence of SEQ ID No: 14.
11. A polynucleotide encoding the fusion protein of claim 1.
12. The polynucleotide according to claim 11, which has a nucleotide acid sequence of SEQ ID No: 15.
13. A recombinant vector comprising the polynucleotide of claim 12.
14. A transformed cell comprising the recombinant vector of claim 13.
15. A method of preparing the fusion protein of claim 1 by culturing the transformed cell of claim 14.
16. A biochip comprising a solid substrate to which the fusion protein of claim 1 is attached.
17. The biochip according to claim 16, wherein the solid substrate is one or more selected group consisting of glass, modified or functionalized glass, plastics, polysaccharides, nylon or nitrocellulose, resins, silica or silica-based materials including silicon and modified silicon, carbon, metals, and inorganic glasses, and wherein the plastics include acrylic, polystyrene, copolymers of styrene and other materials, polypropylene, polyethylene, polybutylene, polyurethanes, Teflon and tetrafluoroethylene.
18. A method for detecting antigen, which comprises the steps of:
  - immobilizing an antibody which specifically binds to the target antigen on the solid substrate of the biochip of claim 16;
  - adding a biological sample comprising the target antigen to the biochip; and
  - detecting the target antigen by measuring antigen-antibody interactions.
19. The method according to claim 18, wherein the biological sample is one or more selected from the group consisting of plasma, blood, saliva, interstitial fluid, serum, urine, synovial, cerebrospinal, lymph, seminal, amniotic, pericardial fluid and ascites.

\* \* \* \* \*

专利名称(译)	固定融合蛋白，有效和定向固定表面抗体		
公开(公告)号	<a href="#">US9005992</a>	公开(公告)日	2015-04-14
申请号	US13/051219	申请日	2011-03-18
[标]申请(专利权)人(译)	浦项工科大学校产学协力团		
申请(专利权)人(译)	POSTECH奥斯卡产业基础		
当前申请(专利权)人(译)	POSTECH奥斯卡产业基础		
[标]发明人	CHA HYUNG JOON KIM CHANG SUP		
发明人	CHA, HYUNG JOON KIM, CHANG SUP		
IPC分类号	G01N33/543 C07K14/435 G01N21/65 G01N33/531		
CPC分类号	G01N33/54353 G01N33/531 G01N21/658 C07K14/43504 C07K2319/705 C07K2319/21		
其他公开文献	US20120238039A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

本发明涉及包含葡萄球菌蛋白A和贻贝粘附蛋白的新型融合蛋白，包含附着有融合蛋白的固体基质的生物芯片，以及使用该生物芯片检测生物样品中的靶抗原的方法。此外，本发明涉及编码融合蛋白的多核苷酸，包含该多核苷酸的重组载体，包含该重组载体的转化细胞，以及通过包含该重组载体的转化细胞制备该融合蛋白的方法。

Fig. 1

