



US009464325B2

(12) United States Patent
Petrovics et al.**(10) Patent No.: US 9,464,325 B2**
(45) Date of Patent: Oct. 11, 2016**(54) METHODS OF DIAGNOSING OR TREATING PROSTATE CANCER USING THE ERG GENE, ALONE OR IN COMBINATION WITH OTHER OVER OR UNDER EXPRESSED GENES IN PROSTATE CANCER**WO 2005007830 1/2005
WO WO 2006/028655 A2 * 3/2006 C12Q 1/68
WO 2010/126972 11/2010**OTHER PUBLICATIONS****(75) Inventors: Gyorgy Petrovics**, Bethesda, MD (US);
Shiv Srivastava, Potomac, MD (US)**(73) Assignee: The Henry M. Jackson Foundation**
for the Advancement of Military
Medicine, Inc., Bethesda, MD (US)**(*) Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.**(21) Appl. No.: 13/534,529****(22) Filed: Jun. 27, 2012****(65) Prior Publication Data**

US 2012/0295282 A1 Nov. 22, 2012

Related U.S. Application Data**(62)** Division of application No. 11/579,695, filed as application No. PCT/US2005/015926 on May 6, 2005.**(60)** Provisional application No. 60/568,822, filed on May 7, 2004, provisional application No. 60/622,021, filed on Oct. 27, 2004.**(51) Int. Cl.**
G01N 33/53 (2006.01)
C12Q 1/68 (2006.01)
C07K 16/30 (2006.01)**(52) U.S. Cl.**
CPC **C12Q 1/6886** (2013.01); **C07K 16/30**
(2013.01); **C12Q 2600/112** (2013.01); **C12Q**
2600/118 (2013.01); **C12Q 2600/158** (2013.01)**(58) Field of Classification Search**
None
See application file for complete search history.**(56) References Cited****U.S. PATENT DOCUMENTS**6,514,751 B2 2/2003 Johann et al.
2001/0010934 A1 8/2001 Libermann et al.**FOREIGN PATENT DOCUMENTS**FR 2 817 043 5/2002
JP 2002/360255 12/2002
WO 98/45420 10/1998
WO 9845420 10/1998
WO 01/88122 11/2001
WO 0188124 11/2001
WO 03/016484 2/2003
WO 03009814 2/2003
WO 03053223 7/2003
WO 2004/031231 4/2004
WO 2004/076814 9/2004Tockman et al (Cancer Res., 1992, 52:2711s-2718s).
Alberts et al. (Molecular Biology of the Cell, 3rd edition, 1994, p. 465).*

Greenbaum et al. (Genome Biology, 2003, vol. 4, Issue 9, pp. 117.1-117.8).*

Reddy et al (PNAS, 1987, 84: 6131-6135).*

Gavrilov et al (European Journal of Cancer, 2001, 37: 1033-1040).*

Ernst et al (American Journal of Pathology, 2002, 160(6): 2169-2180).*

Tsujimoto et al (International Journal of Urology, 2002, 9: 167-172).*

Luo et al (Cancer Res, 2002, 62: 2220-2226).*

Rao et al (Science, 1987, 237: 635-639).*

Luo et al. α -Methylacetyl-CoA Racemase: A New Molecular Marker for Prostate Cancer. Cancer Research, 2002, vol. 62, pp. 2220-2226.

Shaheduzzaman et al. Evaluation of Prostate Cancer Associated Gene Expression Patterns in High Risk Prostate Cancer Patients. Journal of Urology, 2003, vol. 169, No. 4, (Supp.), pp. 51-52.

Xu et al. Quantitative Expression Profile of Androgen-Regulated Genes in Prostate Cancer Cells and Identification of Prostate-Specific Genes. International Journal of Cancer, 2001, vol. 92, pp. 332-328.

Sun et al. A Human Novel Gene DERPC Located on 16q22.1 Inhibits Prostate Tumor Cell Growth and Its Expression is Decreased in Prostate and Renal Tumors. Molecular medicine. 2002, vol. 8, pp. 655-663.

Brooks, James D. Microarray Analysis in Prostate Cancer Research. Current Opinion in Urology, 2002, vol. 12, pp. 395-399.

Hinzman et al. Human prostatic carcinoma derived DNA SEQ ID 84 #4, Dec. 2, 2004, retrieved from EBI accession No. GSN:ADR66792.

Baldus et al. Acute myeloid leukemia with complex karyotypes and abnormal chromosome 21: Amplification discloses overexpression of APP, ETS2, and ERG Genes. PNAS, 2004 vol. 101, pp. 3915-3920.

Carrere et al. Erg proteins, transcription factors of the Ets family, form homo, heterodimers, and ternary complexes via two distinct domains. Institute de Biologie de Lille, France, Jan. 25, 1998, pp. 3261-3268.

(Continued)

Primary Examiner — Sean Aeder**(74) Attorney, Agent, or Firm** — MH2 Technology Law Group, LLP**(57) ABSTRACT**

The present invention relates to oncogenes or tumor suppressor genes, as well as other genes, involved in prostate cancer and their expression products, as well as derivatives and analogs thereof. Provided are therapeutic compositions and methods of detecting and treating cancer, including prostate and other related cancers. Also provided are methods of diagnosing and/or prognosing prostate cancer by determining the expression level of at least one prostate cancer-cell-specific gene, including, for example, the ERG gene or the LTF gene alone, or in combination with at least one of the AMACR gene and the DD3 gene.

28 Claims, 17 Drawing Sheets

(56)

References Cited

OTHER PUBLICATIONS

de Alava et al. Ewing Tumor: Tumor Biology and Clinical Applications. *International Journal of Surgical Pathology*, Jan. 1, 2001, vol. 9, pp. 7-17.

Gavrilov et al. Expression of urokinase plasminogen activator and receptor in conjunction with the ets family and AP-1 complex transcription factors in high grade prostate cancers. *European Journal of Cancer*, Jan. 18, 2001, vol. 37, pp. 1033-1040.

Liu et al. Identification of Differentially Expressed Prostate Genes: Increased Expression of Transcription Factor ETS-02 in Prostate Cancer. *The Prostate*, 1997. 1997. vol. 30, pp. 145-153.

Nelson et al. Preneoplastic Prostate Lesions, and Opportunity for Prostate Cancer Prevention. *Annals New York Academy of Sciences*, pp. 135-144.

Oikawa et al. Molecular biology of the Ets family transcription factors. *Gene*, 2003, vol. 303, pp. 11-34.

Owczarek et al. Detailed mapping of the ERG-ETS2 interval of human chromosome 21 and comparison with the region of conserved synteny on mouse chromosome 16. *Gene*, 2004, vol. 324, pp. 65-77.

Park et al. Organization of the human transferrin gene: Direct evidence that it originated by gene duplication. *Proc. Natl. Acad. Sci. USA*, May 1985, vol. 82, pp. 3149-3153.

Rao et al. erg, a Human ets-Related Gene on Chromosome 21: Alternative Splicing, Polyadenylation, and Translation, *Science*, Aug. 7, 1987, vol. 237, pp. 635-639.

Rubin et al. α -Methylacyl Coenzyme A Racemase as a Tissue Biomarker for Prostate Cancer, *Journal of the American Medical Association* Jan. 9, 2002, vol. 287, No. 13, pp. 1662-1670.

Sementchenko et al. ETS2 function is required to maintain the transformed state of human prostate cancer cells. *Oncogene*, Jun. 16, 1998, vol. 17, pp. 2883-2888.

Seth et al. c-ets-2 protooncogene has mitogenic and oncogenic activity. *Proc. Natl. Acad. Sci.*, Oct. 1989, vol. 86, pp. 7833-7837.

Sharrocks, Andrew D. The ETS-Domain Transcription Factor Family. *Nature Reviews—Molecular Cell Biology*, Nov. 2001, vol. 2, pp. 827-837.

Shimizu et al., An ets-related gene, ERG, is rearranged in human myeloid leukemia with t(16:21) chromosomal translocation. *Proc. Natl. Acad. Sci. USA*, Nov. 1993, vol. 90, pp. 10280-10284.

Simpson et al. Altered expression of Erg and Ets-2 transcription factors associated with genetic changes at 21q22.2-22.3 in immortal and cervical carcinoma cell lines. *Oncogene*, 1997, vol. 154, pp. 2149-2157.

Teng, Christina T. Lactoferrin gene expression and regulation: an overview. *Biochem. Cell Biol.*, 2002, vol. 80, pp. 7-16.

Teng et al. Lactoferrin gene expression is estrogen responsive in human and rhesus monkey endometrium. *Molecular Human Reproduction*, 2002, vol. 8, No. 1, pp. 58-67.

Touda et al. Cancer prevention by bovine lactoferrin and underlying mechanisms—a review of experimental and clinical studies. *Biochem. Cell Biol.*, 2002, vol. 80, pp. 131-136.

Tsuda et al., Prevention of colon carcinogenesis and carcinoma metastasis by orally administered bovine lactoferrin in animals. *Biofactors*, 2002, vol. 12, pp. 83-88.

Tsuda et al. Milk and dairy products in cancer prevention: focus on bovine lactoferrin. *Mutation Research*, 2000, vol. 162, pp. 227-233.

van Sande et al. Lactoferrin in Human Prostate Tissue. *Urological Research*, 1981, vol. 9, pp. 241-244.

Vlaerninck-Guillern et al. Mutual repression of transcriptional activation between the ETS-related factor ERG and estrogen receptor. *Oncogene*, 2003, vol. 22, pp. 8072-8084.

Bowman et al. Transferrin: evolution and genetic regulation of expression. *Adv. Genet.*, 1988, vol. 25, pp. 1-38.

Hart et al. Human ERG is a proto-oncogene with mitogenic and transforming activity, *Oncogene*, 1995, 10(7), pp. 1423-1430.

Papes et al. ETS family of genes in leukemia and Down syndrome. *Am. J. Med. Genet. Suppl.*, 1990, vol. 7, pp. 251-281.

International Search Report and the Written Opinion of the International Searching Authority from PCT/US2005/015926.

Oikawa, Tsuneyuki. ETS transcription factors: Possible targets for cancer therapy. *Cancer Sci.*, 95(8): 626-633, Aug. 2004.

Gavrilov, D. et al. Expression of urokinase plasminogen activator and receptor in conjunction with the ets family and AP-1 complex transcription factors in high grade prostate cancers. *European Journal of Cancer*, 37 (2001) 1033-1040.

Vanaja, D. K. et al. Transcriptional Silencing of Zinc Finger Protein 185 Identified by Expression Profiling Is Associated with Prostate Cancer Progression, *Cancer Research*, 63, Jul. 15, 2003, 3877-3882.

Bussemakers, Martion J. G. et al., "DD3: A New Prostate-specific Gene, Highly Overexpressed in Prostate Cancer", *Cancer Research* 59, Dec. 1, 1999, pp. 5975-5979.

Petrovics, Gyorgy et al., "Frequent overexpression of ETS-related gene-1 (ERG1) in prostate cancer transcriptome", *Oncogene*, 2005, 24, pp. 3847-3852.

Ernst, Thomas et al., "Decrease and Gain of Gene Expression Are Equally Discriminatory Markers for Prostate Carcinoma, A Gene Expression Analysis on Total and Microdissected Prostate Tissue," *American Journal of Pathology*, vol. 160, No. 6, Jun. 2002, pp. 2169-2180.

Amanda M. de Mestre et al., "Regulation of Inducible Heparanase Gene Transcription in Activated T Cells by Early Growth Response 1," *The Journal of Biological Chemistry*, vol. 278, 50, Dec. 23, 2003, pp. 50377-50385.

Ogle B.A. et al. *Nucleic Acids Research* (2003) vol. 31, No. 22, e139, pp. 1-6.

Reddy E.S.P. et al, *Proc. Natl. Acad. Sci. USA* (Sep. 1987) vol. 84, pp. 6131-6135.

Traynor J. et al. *BMC Medical Genetics* (2002) vol. 3, No. 12, Printed pp. 1-15.

Ahern H. *The Scientist*, Jul. 24, 1995, pp. 20 and 22.

Liu G. et al. *Nucleic Acids Research*, 2003, vol. 31, No. 1, pp. 82-86.

Details for HG-U133B 232572_AT (2009), from www.affymetrix.com, Printed pp. 1-2.

Details for HG-U95AV2:36383_AT (2009), from www.affymetrix.com, Printed pp. 1-4.

Details for HG-U133B 230549_AT (2009), from www.affymetrix.com, Printed pp. 1-3.

Details for HG-U95AV2:41706_AT (2009), from www.affymetrix.com, Printed pp. 1-4.

Details for HG-U133A:217113_AT (2009), from www.affymetrix.com, Printed pp. 1-3.

Patent Examination Report No. 2 from Australian Patent Office for Australian Patent Application No. 2011213720 issued Sep. 19, 2012, 4 pages.

Erg-1/2/3 antibody (C-20), Santa Cruz Biotechnology, Inc., accessed online Sep. 19, 2012 <[URL:http://www.scbt.com/datasheet-353-erg-1-2-3-c-20-antibody.html](http://www.scbt.com/datasheet-353-erg-1-2-3-c-20-antibody.html)>.

European Patent Office Communication Pursuant to Article 94(3) EPC from European Patent Application No. 05 779 973.6 dated Feb. 11, 2011, 7 Pages.

European Patent Office Communication Pursuant to Rule 114(2) EPC dated Oct. 24, 2012, reporting Third Party Observations in European Patent Application No. 05779973.6, 9 Pages.

European Patent Office Communication Pursuant to Article 94(3) EPC from European Patent Application No. 05 779 973.6 dated Jul. 2, 2012, 7 Pages.

European Patent Office Communication, European Search Report from European Patent Application No. 11172817.6 dated Sep. 30, 2011, 12 Pages.

European Patent Office Communication Pursuant to Rule 114(2) EPC dated Feb. 28, 2013, reporting Third Party Observations in European Patent Application No. 05779973.6, 9 Pages.

Communication dated Mar. 12, 2014 from European Patent Application No. 11172817.6, pp. 1-11.

Mohamed et al., Ets Family Protein, Erg Expression in Developing and Adult Mouse Tissues by a Highly Specific Monoclonal Antibody, *Journal of Cancer*, 1:197-208 (2010).

Rosen, Philip et al., Clinical Potential of the ERG Oncoprotein in Prostate Cancer, 9:131-137 (2012).

(56)

References Cited

OTHER PUBLICATIONS

Santa Cruz product description of sc-353 printed from Santa Cruz Biotechnology website in Jan. 2014.

Non-Final Office Action dated Jan. 30, 2015 from U.S. Appl. No. 13/756,028, filed Jan. 31, 2013, 7 Pages.

Non-Final Office Action dated Feb. 9, 2015 from U.S. Appl. No. 11/579,695, filed Oct. 9, 2008, 17 Pages.

European Patent Office Communication dated Mar. 12, 2014 from European Patent Application No. 11172817.6, 11 Pages.

De Kok, Jacques B. et al. DD3(PCA3), a Very Sensitive and Specific Marker to Detect Prostate Tumors. *Cancer Research*, May 1, 2002, vol. 62, pp. 2695-2698.

Non-Final Office Action dated Aug. 30, 2013 from U.S. Appl. No. 13/445,706, filed Apr. 12, 2012, 13 Pages.

Dhanasekaran, Saravana M. et al. Delineation of prognostic biomarkers in prostate cancer. *Nature*, Aug. 23, 2001, vol. 412, pp. 822-826 with supplementary information available at www.nature.com.

Luo, Jun et al. α -Methylcayl-CoA Racemase: A New Molecular Marker for Prostate Cancer. *Cancer Research*, Apr. 15, 2002, vol. 62, pp. 2220-2226.

Wang, Shunyou et al. Prostate-specific deletion of the murine Pten tumor suppressor gene leads to metastatic prostate cancer. *Cancer Cell*, Sep. 2003, vol. 4, pp. 209-221.

Reddy, E. S. et al. Human erg protein (ets-related gene) mRNA, complete cds. GenBank Locus HUMERG11, Apr. 27, 1993, pp. 1-2, www.ncbi.nih.gov.

European Patent Office Communication dated Aug. 6, 2013 from European Patent Application No. 05 779 9736, pp. 1-7.

Third Party Observations dated Oct. 24, 2012 from European Patent Application No. 05 779 9736, pp. 1-9.

European Patent Office Communication dated Jul. 2, 2012 from European Patent Application No. 05 779 9736, pp. 1-7.

Third Party Observations dated Feb. 28, 2013 from European Patent Application No. 05 779 9736, pp. 1-6.

Hollenhorst, Peter C. et al. Genomic and Biochemical Insights into the Specificity of ETS Transcription Factors. *Annual Review of Biochemistry*, May 3, 2011, pp. 437-471.

Details for HG-U95E:91858_AT (2009), from www.affymetrix.com, Printed pp. 1-3.

Details for HG-U133A:211626_X_AT (2009), from www.affymetrix.com, Printed pp. 1-3.

GenBank Locus HUMERG11 (Apr. 11, 1993) "Human erg protein (ets-related gene) mRNA, complete cds" from www.ncbi.nlm.nih.gov, pp. 1-3.



Communication dated Jan. 9, 2014 from Canadian Patent Application No. 2,565,450, pp. 1-5.



Schalken, Jack A. et al. New targets for therapy in prostate cancer: different display code 3 (DD3(PCA3)), a highly prostate cancer-specific gene. *Urology*, Nov. 2003, vol. 62 (Supplement 5A), pp. 34-43.


DePrimo, Samuel E. et al. Expression profiling of blood samples from an SU5416 Phase III metastatic colorectal cancer clinical trial: a novel strategy for biomarker identification. *BMC Cancer*, Feb. 7, 2003, 3:3, pp. 1-12.


Shing, Danielle C. et al. FUS/ERG Gene Fusions in Ewing's Tumors. *Cancer Research*, Aug. 1, 2003, vol. 63, pp. 4568-4576.

* cited by examiner

**FIG. 1A****FIG. 1B**

**FIG. 1C****FIG. 1D**

**FIG. 2A**



2025



**FIG. 3A**

FIG. 3B







FIG. 3B CONT.

**FIG. 3C**






FIG. 3D

**FIG. 3E**

**FIG. 4A**


**FIG. 4B**

FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E





FIG. 5F








FIG. 6

**FIG. 7**

**FIG. 8**




FIG. 9

METHODS OF DIAGNOSING OR TREATING PROSTATE CANCER USING THE ERG GENE, ALONE OR IN COMBINATION WITH OTHER OVER OR UNDER EXPRESSED GENES IN PROSTATE CANCER

CROSS REFERENCE TO RELATED APPLICATIONS

The present application is a divisional of U.S. patent application Ser. No. 11/579,695, which is a national phase application of PCT/US2005/015926, filed May 6, 2005, and claims the benefit of U.S. provisional applications No. 60/568,822, filed May 7, 2004, and No. 60/622,021, filed Oct. 27, 2004, the entire disclosures of which are relied upon and incorporated by reference.

GOVERNMENT INTEREST

This invention was made with government support under grant numbers DK065977 and CA162383 awarded by the National Institutes of Health. The government has certain rights in the invention.

FIELD OF THE INVENTION

The present invention relates to oncogenes, tumor suppressor genes, as well as other genes, and their expression products, involved in prostate cancer, as well as derivatives and analogs thereof. The invention further relates to therapeutic compositions and methods of detecting, diagnosing, and treating cancer, including prostate and other related cancers.

BACKGROUND OF THE INVENTION

Prostate cancer (CaP) is the most common malignancy in American men and second leading cause of cancer mortality (Landis et al. (1999) *Cancer J. Clin.*, 49:8-31; Jemal et al. (2004) *Cancer J Clin* 54:8-29). The molecular determinants in the development and progression of this disease are poorly understood. In recent years, there have been intensive investigations of molecular genetics of the CaP. To date, however, oncogene, tumor suppressor gene, or other gene alterations common to most CaPs have not been found. Alterations of tumor suppressors such as p53, PTEN and p27, or oncogenes such as BCL2, HER2 and C-MYC associate with only small subsets of primary CaP, with more frequent association observed in advanced CaP.

Current clinical parameters, including serum Prostate Specific Antigen (PSA), tumor stage, and Gleason score are routinely used as risk factors at the time of diagnosis, but have limited application to identify patients at a greater risk for developing aggressive CaP. Approximately 30-40% of patients treated with radical prostatectomy for localized CaP have been found to have microscopic disease that is not organ-confined and a significant portion of these patients relapse. (Singh et al., *Cancer Cell* (2000) 1:203-209; Henshell et al., *Can. Res.* (2003) 63: 4196-4203). Therefore, discovery of novel biomarkers or gene expression patterns defining CaP onset and progression is crucial in predicting patients with greater risk to develop aggressive CaP.

CaP-specific genetic alterations have been the subject of intensive research by several investigations in the past five years (Srikatan et al., *In Prostate Cancer, Diagnosis and Surgical Treatment* (2002) Springer-Verlag, 25-40; Karan et al., *Int. J. Can.* (2003) 103:285-293; Augustus et al., *In*

Molecular Pathology of Early Cancer (1999) IOS press: 321-340; Moul et al., *Clin Prostate Cancer* (2002) 1:42-50; Lalani et al., *Cancer and Mets Rev* (1997) 16: 29-66; Issacs et al., *Epidemiol Rev* (2001) 23:36-41; Ozen et al., *Anti-cancer Res* (2000) 20:1905-1912; Morton et al., *J Natl Med Assoc* (1998) 90:S728-731). Promising leads both in biology and translational research areas are beginning to emerge from recent genomics and proteomics technology, as well as traditional approaches. However, the inherent heterogeneity of CaP has hampered the molecular characterization of CaP.

One of the challenges in studying molecular alterations in human cancers, including prostate tumors, is to define the relative contributions of genetic alterations in epithelial and non-epithelial components of the target organ in the process of tumorigenesis. Despite advances in technology, changes in human CaP-specific epithelial and stromal cell-associated gene expression are still not well understood.

Despite recent advances in the identification of molecular alterations associated with certain prostate cancers, the heterogeneous nature of prostate tissue has hindered the identification of genetic targets common to all, or at least the vast majority of, prostate cancers. The complexity and heterogeneity of prostate cancer has also hindered the identification of targets that allow differentiation between clinically aggressive and non-aggressive cancers at the time of diagnosis. Therefore, there remains a need to identify molecular alterations specific for a pathologically defined cell population that can provide important clues for optimal diagnosis and prognosis, and help to establish individualized treatments tailored to the molecular profile of the tumor.

Citation of references herein shall not be construed as an admission that such references are prior art to the present invention.

SUMMARY OF THE INVENTION

It is one of the objects of the present invention to provide methods and kits for detecting cancer, in particular prostate cancer. These methods and kits can be used to detect (either qualitatively or quantitatively) nucleic acids or proteins that serve as cancer markers. For example, the expression of the prostate cancer-cell-specific gene ERG, when detected in a biological sample from a subject, either alone or in combination with other cancer markers, including the expression of other prostate cancer-cell-specific genes, can be used to indicate the presence of prostate cancer in the subject or a higher predisposition of the subject to develop prostate cancer. Detecting ERG expression, alone or in combination with the expression of any gene identified in Tables 1-6, can thus be used to diagnose or prognose cancer, particularly prostate cancer.

According to one aspect of the invention, the method for detecting the expression of one or more prostate cancer cell-specific genes, such as ERG, AMACR, and LTF or the DD3 gene, in a biological sample, comprises:

- (a) combining a biological sample with at least a first and a second oligonucleotide primer under hybridizing conditions, wherein the first oligonucleotide primer contains a sequence that hybridizes to a first sequence in a target sequence from a prostate cancer cell-specific gene, such as ERG (SEQ ID NO:1), AMACR (SEQ ID NO:3), and/or LTF (SEQ ID NO:5) and/or DD3 (SEQ ID NO:4), and the second oligonucleotide primer contains a sequence that hybridizes to a second sequence in a nucleic acid strand complementary to the target sequence, wherein the first sequence does not overlap with the second sequence,

- (b) adding at least one polymerase activity to produce a plurality of amplification products when the target sequence is present in the biological sample,
- (c) adding an oligonucleotide probe that hybridizes to at least one amplification product of the target sequence, and
- (d) detecting whether a signal results from hybridization between the oligonucleotide probe and the at least one amplification product, wherein detection of the signal indicates the expression of a prostate cancer cell-specific gene in the biological sample.

The method preferably comprises detecting the expression of the following combinations of genes: 1) ERG and AMACR; 2) ERG and DD3; and 3) ERG, AMACR and DD3. In another embodiment, the method comprises detecting LTF and one or more of ERG, AMACR and DD3. Expression of these genes can also be detected by measuring ERG, AMACR or LTF polypeptides in the biological sample.

The biological sample is preferably a prostate tissue, blood, or urine sample. Detecting a signal resulting from hybridization between the oligonucleotide probe and the at least one amplification product can be used to diagnose or prognose cancer, particularly prostate cancer.

The oligonucleotide probe may be optionally fixed to a solid support. When detecting ERG expression in a biological sample, the oligonucleotide probe, first oligonucleotide primer, and second oligonucleotide primer, each comprise a nucleic acid sequence that is capable of hybridizing under defined conditions (preferably under high stringency hybridization conditions, e.g., hybridization for 48 hours at 65° C. in 6×SSC followed by a wash in 0.1×SSC at 50° C. for 45 minutes) to SEQ ID NO:1. Thus, the oligonucleotide probe, first oligonucleotide primer, and second oligonucleotide primer can include, for example, SEQ ID NO:1 itself, or a fragment thereof or a sequence complementary thereto. Preferably the oligonucleotide probe, first oligonucleotide primer, or second oligonucleotide primer is a fragment of SEQ ID NO:1 having at least about 15, at least about 20, or at least about 50 contiguous nucleotides of SEQ ID NO:1 or a sequence complementary thereto. When detecting ERG expression, the target sequence is preferably a fragment of SEQ ID NO:1. Probes, primers, and target sequences can be similarly derived from other genes of interest, such as DD3 (SEQ ID NO:4), and other prostate cancer-cell-specific genes, including, for example, AMACR (SEQ ID NO:3) and LTF (SEQ ID NO:5).

In another aspect of the invention, the method of diagnosing or prognosing prostate cancer comprises:

- measuring the expression level (e.g. mRNA or polypeptide) of an over expressed prostate cancer cell-specific gene, such as ERG and/or AMACR, and/or the DD3 gene in a biological sample, and
- correlating the expression level of the ERG, AMACR, and/or DD3 gene with the presence of prostate cancer or a higher predisposition to develop prostate cancer in the subject.

In a related aspect of the invention, the method of diagnosing or prognosing prostate cancer comprises:

- measuring the expression level (e.g. mRNA or polypeptide) of an under expressed prostate cancer cell-specific gene, such as LTF in a biological sample, and
- correlating the expression level of the LTF gene with the presence of prostate cancer or a higher predisposition to develop prostate cancer in the subject.

The skilled artisan will understand how to correlate expression levels or patterns of the desired genes with the

presence of prostate cancer or a higher predisposition to develop prostate cancer. For example, the expression levels can be quantified such that increased or decreased expression levels relative to a control sample or other standardized value or numerical range indicate the presence of prostate cancer or a higher predisposition to develop prostate cancer.

The increased or decreased expression levels in the methods of the invention may be measured relative to the expression level of the prostate cancer cell-specific gene or polypeptide in normal, matched tissue, such as benign prostate epithelial cells from the same subject. Alternatively, the expression level of a gene or polypeptide may be measured relative to the expression of the gene or polypeptide in other noncancerous samples from the subject or in samples obtained from a different subject without cancer. Expression of a gene may also be normalized by comparing it to the expression of other cancer-specific markers. For example, in prostate cancer, a prostate-cell specific marker, such as PSA, can be used as a control to compare and/or normalize expression levels of other genes, such as ERG, LTF, DD3, and/or AMACR. By way of example, the method of diagnosing or prognosing prostate cancer comprises measuring the expression levels of the ERG, DD3 and/or AMACR gene and diagnosing or prognosing prostate cancer, where an increased expression level of the ERG, DD3, and/or AMACR gene of at least two times as compared to the control sample indicates the presence of prostate cancer or a higher predisposition in the subject to develop prostate cancer. Conversely, by way of example, in such a method of diagnosing or prognosing prostate cancer, a decreased expression of the LTF gene of at least two times as compared to the control sample indicates the presence of prostate cancer or a higher predisposition in the subject to develop prostate cancer.

The expression levels of prostate cancer cell-specific genes (e.g., mRNA or polypeptide expression) can be detected according to the methods described herein or using any other known detection methods, including, without limitation, immunohistochemistry, Southern blotting, Northern blotting, Western blotting, ELISA, and nucleic acid amplification procedures, including but not limited to PCR, transcription-mediated amplification (TMA), nucleic acid sequence-based amplification (NASBA), self-sustained sequence replication (3SR), ligase chain reaction (LCR), strand displacement amplification (SDA), and Loop-Mediated Isothermal Amplification (LAMP).

It is yet another object of the present invention to provide a method of determining a gene expression pattern in a biological sample, where the pattern can be correlated with the presence or absence of tumor cells, particularly prostate tumor cells. For example, ERG is detected in combination with other prostate cancer cell-specific genes (identified in Tables 1-6), including AMACR and/or LTF, to obtain expression profiles from biological samples. The expression profiles of these prostate cancer-cell-specific genes are useful for detecting cancer, particularly prostate cancer. ERG can also be detected in combination with DD3, with or without other prostate cancer cell-specific genes, such as AMACR and/or LTF, to obtain expression profiles from biological samples. These expression profiles are also useful for detecting cancer, particularly prostate cancer. Increased levels of ERG, AMACR, and/or DD3 in a biological sample indicate the presence of prostate cancer or a higher predisposition in the subject to develop prostate cancer. Decreased levels of LTF in a biological sample indicate the presence of prostate cancer or a higher predisposition in the subject to develop prostate cancer.

It is yet another object of the present invention to provide a method of determining a gene expression pattern in a biological sample, where the pattern can be used to indicate or predict the pathologic stage of cancer, particularly prostate cancer. For example, the gene expression pattern can be used to indicate or predict a moderate risk prostate cancer or a high risk prostate cancer or to predict whether the prostate cancer is progressing or regressing or in remission. The gene expression pattern can also be used as a prognostic indicator of disease-free survival following radical prostatectomy. In a particular embodiment, gene expression patterns are derived from the expression level of the ERG gene, alone or in combination with other prostate cancer-cell-specific genes (identified in Tables 1-6), including AMACR and LTF, or DD3.

Kits for detecting cancer, particularly prostate cancer, are also provided. These kits comprise a nucleic acid probe, such as the ones described herein, that hybridizes to a prostate cancer-cell-specific gene. In one embodiment the nucleic acid probe hybridizes to SEQ ID NO:1 (ERG) or the complement thereof under defined hybridization conditions (preferably under high stringency hybridization conditions, e.g., hybridization for 48 hours at 65° C. in 6×SSC followed by a wash in 0.1×SSX at 50° C. for 45 minutes) and includes SEQ ID NO:1, itself, or a fragment of SEQ ID NO:1 having at least about 15, at least about 20, or at least about 50 contiguous nucleotides of SEQ ID NO:1 or a sequence complementary thereto. In a particular embodiment, the probe selectively hybridizes to the ERG1 and ERG2 isoforms but not to ERG isoforms 3-9. In another embodiment, the probe selectively hybridizes to the ERG1 isoform but not to ERG isoforms 2-9. The nucleic acid probe may be optionally fixed to a solid support.

The kit may also contain at least one additional nucleic acid probe that hybridizes (preferably high stringency hybridization conditions, e.g., hybridization for 48 hours at 65° C. in 6×SSC followed by a wash in 0.1×SSX at 50° C. for 45 minutes) to DD3 (SEQ ID NO:4) or a gene identified in Tables 1-6, including for example, AMACR (SEQ ID NO:3) or LTF (SEQ ID NO:5). In one embodiment, the kit comprises a first oligonucleotide probe capable of hybridizing to SEQ ID NO:1 (ERG) or a sequence complementary thereto under conditions of high stringency and at least one other oligonucleotide probe capable of hybridizing to SEQ ID NO:3 (AMACR) or a sequence complementary thereto, or to SEQ ID NO:4 (DD3) or a sequence complementary thereto, or to a gene identified in Tables 1-6 under conditions of high stringency. In a related embodiment, the kit having an ERG and AMACR probe further comprises a third oligonucleotide probe capable of hybridizing to SEQ ID NO:4 (DD3) or a sequence complementary thereto. The kits described herein may optionally contain an oligonucleotide probe capable of hybridizing to SEQ ID NO:5 (LTF) or a sequence complementary thereto under conditions of high stringency.

The kits may further comprise a first oligonucleotide primer and a second oligonucleotide primer, where the first oligonucleotide primer contains a sequence that hybridizes to a first sequence in SEQ ID NO:1, and the second oligonucleotide primer contains a sequence that hybridizes to a second sequence in a nucleic acid strand complementary to SEQ ID NO:1, wherein the first sequence does not overlap with the second sequence. The first and second oligonucleotide primers are capable of amplifying a target sequence of interest in SEQ ID NO:1. Similarly, the kits can further comprise first and second oligonucleotide primers derived

from DD3 (SEQ ID NO:4) or a prostate cancer-cell-specific gene, including, for example AMACR (SEQ ID NO:3) or LTF (SEQ ID NO:5).

It is another object of the invention to provide therapeutic methods of treating cancer, in particular prostate cancer.

It is yet another object of the present invention to provide screening methods for identifying compounds that modulate expression of a CaP-cell-specific gene, such as ERG, in prostate cancer cells.

The present invention is based in part on the identification of gene expression signatures that correlate with a high risk of CaP progression. Over expression or under expression of specific genes are predictive of tumor progression. The invention provides genes, such as the ERG gene, and analogs of specific genes that can be used alone or in combination with DD3 or other CaP-cell-specific genes, such as AMACR or LTF, to function as diagnostic and prognostic targets for cancer, particularly prostate tumors. The invention further provides genes, such as the ERG gene, and analogs of specific genes that can be used alone or in combination as therapeutic targets for cancer, in particular prostate tumors.

The invention further discloses diagnostic kits comprised of an anti-CaP-cell-specific gene antibody, for example, an anti-ERG gene antibody, which is optionally, detectably labeled. A kit is also provided that comprises nucleic acid primer sequences and/or a nucleic acid probe capable of hybridizing under defined conditions (preferably high stringency hybridization conditions, e.g., hybridization for 48 hours at 65° C. in 6×SSC followed by a wash in 0.1×SSX at 50° C. for 45 minutes) to an ERG nucleic acid. The kits may also contain an anti-DD3 gene antibody or a second anti-CaP-cell-specific gene antibody, such as an anti-AMACR or anti-LTF gene antibody, or a second set of nucleic acid primer sequences and/or a nucleic acid probe capable of hybridizing under defined conditions to the DD3 gene or another CaP-cell-specific gene, such as the AMACR or LTF gene.

The disclosed CaP-cell-specific genes, such as ERG, can be used alone or in combination as biomarkers of cancer, and in particular, prostate cancers and other related diseases, as targets for therapeutic intervention, or as gene therapy agents.

The invention provides for treatment of disorders of hyperproliferation (e.g., cancer, benign tumors) by administering compounds that modulate expression of the specific genes.

Methods of screening cancer cells, and in particular, prostate cancer cells, for specific gene expression signatures, including ERG gene expression signatures, alone or in combination with DD3 gene expression signatures or other CaP-cell-specific gene expression signatures, such as AMACR or LTF, are provided.

Additional objects of the invention will be set forth in part in the description following, and in part will be understood from the description, or may be learned by practice of the invention.

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention as claimed.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1: Relative expression level of ERG (A), AMACR (B), GSTP1 (C), and LTF (D) in matched tumor and benign prostate epithelial cells analyzed by QRT-PCR (TaqMan)

X-axis: CaP patients analyzed (1-20); Y-axis: Expression ratio between tumor versus benign laser capture microdissection (LCM) sample pairs.

FIG. 2: Identification of genes by a distance based MDS and weighted analysis that discriminates between cancerous and benign tissue. (A) Two-dimensional MDS plot elucidating discrimination of 18 tumor samples and 18 benign samples. (B) Hierarchical clustering dendrogram with two major clusters of 18 tumor samples in the right cluster and 18 benign samples in the left cluster.

FIG. 3: A distance based MDS and weighted gene analysis using the tumor over benign ratio (or fold change) data for the identification of genes that can discriminate between high risk CaP and moderate risk CaP. (A) A supervised MDS analysis of 18 samples (9 samples from high risk group and 9 samples from moderate risk group) that ranks the genes according to their impact on minimizing cluster volume and maximizing center-to-center inter cluster distance. (B) Hierarchical clustering of the first 55 genes of the top 200 obtained by the MDS analysis. Genes and samples are arranged as ordered by cluster and treeview. Expression of each gene in each sample is obtained by the tumor over benign ratio or fold change (T/N). Dendrogram at the top of the cluster shows two major clusters, 9 samples of the MR groups in the right cluster and 9 samples of the HR groups in the left cluster. (C) Two-dimensional MDS plot of 18 CaP tumor epithelia that shows the differentiation between the high risk group (9 tumor epithelia) and moderate risk group (9 tumor epithelia) on the basis of the impact of the rank of the genes that discriminate between the HR and MR groups. (D) Hierarchical clustering dendrogram with two major clusters of 9 samples of the MR groups in the left cluster and 8+1 samples of the HR groups in the right cluster. (E) Two-dimensional MDS plot of 18 CaP benign epithelia that shows the discrimination between the high risk group (9 benign epithelia) and moderate risk group (9 benign epithelia) samples.

FIG. 4: In silico validation: the discriminatory potential of the genes that we obtained from our supervised MDS analysis on two independent data sets (Welsh et al. 2001, Singh et al. 2002). Two-dimensional MDS plot that shows the discrimination between 7 tumor epithelia of the high risk group and 7 tumor epithelia of the moderate risk group using data from Welsh et al. (A), as well as discrimination between 4 tumor epithelia of the high risk group and 5 tumor epithelia of the moderate risk group using data from Singh et al. (B).

FIG. 5: Combined gene expression analysis of ERG, AMACR and DD3 in tumor and benign prostate epithelial cells of 55 CaP patients. The graphs represent patient distribution by tumor versus benign gene expression ratios according to five gene expression categories: 1) "Up:" greater than 2 fold over expression in tumor compared to benign; 2) "Down:" less than 0.5 fold under expression in tumor compared to benign; 3) "Same:" no significant difference (0.5 to 2 fold); 4) "No expr.:" no detectable gene expression; and 5) "Other:" collectively defines patients with expression category 2, 3 and 4 for the indicated genes (i.e., other than category 1). (A) ERG Expression. (B) AMACR Expression. (C) DD3 Expression. (D) ERG or AMACR Expression. (E) ERG or DD3 Expression. (F) ERG, AMACR, or DD3 Expression.

FIG. 6: Map of ERG1 and ERG2 isoforms with probe and primer locations. The numbered boxes represent exons, the darker boxes after exon 16 are the 3' non-coding exon regions. Translational start and stop codons are indicated by star and pound signs, respectively. The locations of the

Affymetrix probe set (213541_s_at), the TaqMan probes, the traditional RT-PCR primers, and the in situ hybridization probe are indicated.

FIG. 7: Correlation of ERG1 expression and PSA recurrence-free survival. Kaplan-Meier analysis of correlation with post-prostatectomy PSA recurrence-free survival was performed on 95 CaP patients having detectable levels of ERG1 mRNA by real time QRT-PCR (TaqMan). Kaplan-Meier survival curves were stratified by the following ERG1 expression categories: 1) greater than 100 fold over expression; 2) 2-100 fold over expression; and 3) less than 2 fold over expression or under expression of ERG1 in the prostate tumor cells. The p value was 0.0006.

FIG. 8: In situ hybridization images in 7 CaP patients were analysed by the Open-Lab image analysis software (Improvisation, Lexington, Mass.) coupled to a microscope via a cooled digital camera (Leica Microsystems, Heidelberg, Germany). Density (OD) values for tumor (dark columns) and benign (light columns) epithelium are shown on the Y axis, and patients 1-7 are shown in the X axis. Patient No. 7 was added as a control with no significant ERG1 expression difference between tumor and benign cells by QRT-PCR (TaqMan). Statistical analysis was performed with the SPSS software package.

FIG. 9: ERG1 is represented as a modular structure. The two conserved regions namely SAM-PNT Domain (Protein/RNA interaction domain) and ETS Domain (Interaction with DNA) are shaded.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

The term "CaP-cell-specific gene," or "prostate cancer-cell-specific gene," refers to a gene identified in Tables 1-6. The definition further encompasses CaP-cell-specific gene analogs, e.g., orthologues and homologues, and functionally equivalent fragments of CaP-cell-specific genes or their analogs, the expression of which is either upregulated or downregulated in prostate cancer cells.

The term "CaP-cell-specific gene expression signature" refers to the pattern of upregulation or downregulation of product expression as measured by the Affymetrix GeneChip assay described in Example 1, the QRT-PCR assay described in Example 2, or any other quantitative expression assay known in the art.

The term "ERG" refers to the ERG gene or ERG cDNA or mRNA described herein, and includes ERG isoforms, such as ERG1 and ERG2. The cDNA sequence of the ERG1 gene is published in GenBank under the accession number M21535. The cDNA sequence of the ERG2 gene is published in GenBank under the accession number M17254.

The term "AMACR" refers to the AMACR gene or AMACR cDNA or mRNA described herein, and includes AMACR isoforms. The cDNA sequence of the AMACR gene is published in GenBank under the accession number NM_014324.

The term "DD3" refers to the DD3 gene or DD3 cDNA or mRNA described herein, and includes DD3 isoforms. The cDNA sequence of the DD3 gene is published in GenBank under the accession number AF 103907 and is also disclosed in WO 98/45420 (1998). Although DD3 was originally used to describe a fragment of exon 4 of the prostate cancer antigen 3 (PCA3) gene, the term, as used in herein, is not so

limited. DD3 is intended to refer to the entire DD3 gene or cDNA or mRNA, which in the art is also commonly referred to as PCA3.

The term "LTF" refers to the LTF gene or LTF cDNA or mRNA described herein and includes LTF isoforms. The cDNA sequence of the LTF gene is published in GenBank under the accession number NM_002343.

The term "polypeptide" is used interchangeably with the terms "peptide" and "protein" and refers to any chain of amino acids, regardless of length or posttranslational modification (e.g., glycosylation or phosphorylation), or source (e.g., species).

The phrase "substantially identical," or "substantially as set out," means that a relevant sequence is at least 70%, 75%, 80%, 85%, 90%, 95%, 97, 98, or 99% identical to a given sequence. By way of example, such sequences may be allelic variants, sequences derived from various species, or they may be derived from the given sequence by truncation, deletion, amino acid substitution or addition. For polypeptides, the length of comparison sequences will generally be at least 20, 30, 50, 100 or more amino acids. For nucleic acids, the length of comparison sequences will generally be at least 50, 100, 150, 300, or more nucleotides. Percent identity between two sequences is determined by standard alignment algorithms such as, for example, Basic Local Alignment Tool (BLAST) described in Altschul et al. (1990) *J. Mol. Biol.*, 215:403-410, the algorithm of Needleman et al. (1970) *J. Mol. Biol.*, 48:444-453, or the algorithm of Meyers et al. (1988) *Comput. Appl. Biosci.*, 4:11-17.

The terms "specific interaction," "specific binding," or the like, mean that two molecules form a complex that is relatively stable under physiologic conditions. The term is also applicable where, e.g., an antigen-binding domain is specific for a particular epitope, which is carried by a number of antigens, in which case the specific binding member carrying the antigen-binding domain will be able to bind to the various antigens carrying the epitope. Specific binding is characterized by a high affinity and a low to moderate capacity. Nonspecific binding usually has a low affinity with a moderate to high capacity. Typically, the binding is considered specific when the affinity constant K_a is higher than $10^6 M^{-1}$, more preferably higher than $10^7 M^{-1}$, and most preferably $10^8 M^{-1}$. If necessary, non-specific binding can be reduced without substantially affecting specific binding by varying the binding conditions. Such conditions are known in the art, and a skilled artisan using routine techniques can select appropriate conditions. The conditions are usually defined in terms of concentration of antibodies, ionic strength of the solution, temperature, time allowed for binding, concentration of non-related molecules (e.g., serum albumin, milk casein), etc. The term "detectably labeled" refers to any means for marking and identifying the presence of a molecule, e.g., an oligonucleotide probe or primer, a gene or fragment thereof, or a cDNA molecule. Methods for labeling a molecule are well known in the art and include, without limitation, radioactive labeling (e.g., with an isotope such as ^{32}P , ^{35}S , or ^{125}I) and nonradioactive labeling (e.g., fluorescent and chemiluminescent labeling).

The term "modulatory compound" is used interchangeably with the term "therapeutic" as used herein means any compound capable of "modulating" either CaP-cell-specific gene expression at the transcriptional, translational, or post-translational levels or modulating the biological activity of a CaP-cell-specific polypeptide. The term "modulate" and its cognates refer to the capability of a compound acting as either an agonist or an antagonist of a certain reaction or activity. The term modulate, therefore, encompasses the

terms "activate" and "inhibit." The term "activate," for example, refers to an increase in the expression of the CaP-cell-specific gene or activity of a CaP-cell-specific polypeptide in the presence of a modulatory compound, relative to the activity of the gene or the polypeptide in the absence of the same compound. The increase in the expression level or the activity is preferably at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or higher. Analogously, the term "inhibit" refers to a decrease in the expression of the CaP-cell-specific gene or activity of a CaP-cell-specific polypeptide in the presence of a modulatory compound, relative to the activity of the gene or the polypeptide in the absence of the same compound. The decrease in the expression level or the activity is preferably at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or higher. The expression level of the CaP-cell-specific gene or activity of a CaP-cell-specific polypeptide can be measured as described herein or by techniques generally known in the art.

The term "treatment" is used interchangeably herein with the term "therapeutic method" and refers to both therapeutic treatment and prophylactic/preventative measures. Those in need of treatment may include individuals already having a particular medical disorder as well as those who may ultimately acquire the disorder.

The term "isolated" refers to a molecule that is substantially free of its natural environment. Any amount of that molecule elevated over the naturally occurring levels due to any manipulation, e.g., over expression, partial purification, etc., is encompassed with the definition. With regard to partially purified compositions only, the term refers to an isolated compound that is at least 50-70%, 70-90%, 90-95% (w/w), or more pure.

The term "effective dose," or "effective amount," refers to that amount of the compound that results in amelioration of symptoms in a patient or a desired biological outcome, e.g., inhibition of cell proliferation. The effective amount can be determined as described in the subsequent sections.

The terms "polynucleotide," "oligonucleotide," "nucleic acid," and "DNA" are used interchangeably herein and refer to deoxyribonucleic acid (DNA), and, where appropriate, ribonucleic acid (RNA). The term should also be understood to include nucleotide analogs, and single or double stranded polynucleotides. Examples of polynucleotides include, but are not limited to, plasmid DNA or fragments thereof, viral DNA or RNA, anti-sense RNA, etc. The term "plasmid DNA" refers to double stranded DNA that is circular.

As used herein the term "hybridization under defined conditions," or "hybridizing under defined conditions," is intended to describe conditions for hybridization and washes under which nucleotide sequences that are significantly identical or homologous to each other remain bound to each other. The conditions are such that sequences, which are at least about 6 and more preferably at least about 20, 50, 100, 150, 300, or more nucleotides long and at least about 70%, more preferably at least about 80%, even more preferably at least about 85-90% identical, remain bound to each other. The percent identity can be determined as described in Altschul et al. (1997) *Nucleic Acids Res.*, 25: 3389-3402.

Appropriate hybridization conditions can be selected by those skilled in the art with minimal experimentation as exemplified in Ausubel et al. (2004), *Current Protocols in Molecular Biology*, John Wiley & Sons. Additionally, stringent conditions are described in Sambrook et al. (2001) *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press. A nonlimiting example of defined conditions of low stringency is as follows. Filters

containing DNA are pretreated for 6 hours at 40° C. in a solution containing 35% formamide, 5×SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA. Hybridizations are carried out in the same solution with the following modifications: 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, 10% (wt/vol) dextran sulfate, and 5-20×10⁶ cpm ³²P-labeled probe is used. Filters are incubated in hybridization mixture for 18-20 hours at 40° C., and then washed for 1.5 hours at 55° C. in a solution containing 2×SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS. The wash solution is replaced with fresh solution and incubated an additional 1.5 hours at 60° C. Filters are blotted dry and exposed for autoradiography. Other conditions of low stringency well known in the art may be used (e.g., as employed for cross-species hybridizations).

A non-limiting example of defined conditions of high stringency is as follows. Prehybridization of filters containing DNA is carried out for 8 hours to overnight at 65° C. in buffer composed of 6×SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are hybridized for 48 hours at 65° C. in the prehybridization mixture containing 100 µg/ml denatured salmon sperm DNA and 5-20×10⁶ cpm of ³²P-labeled probe. Washing of filters is done at 37° C. for 1 hour in a solution containing 2×SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA. This is followed by a wash in 0.1×SSC at 50° C. for 45 minutes. Other conditions of high stringency well known in the art may be used. An oligonucleotide hybridizes specifically to a target sequence under high stringency conditions.

The term "solid support" means a material that is essentially insoluble under the solvent and temperature conditions of the assay method, comprising free chemical groups available for joining an oligonucleotide or nucleic acid. Preferably, the solid support is covalently coupled to an oligonucleotide designed to directly or indirectly bind a target nucleic acid. When the target nucleic acid is an mRNA, the oligonucleotide attached to the solid support is preferably a poly-T sequence. A preferred solid support is a particle, such as a micron- or submicron-sized bead or sphere. A variety of solid support materials are contemplated, such as, for example, silica, polyacrylate, polyacrylamide, a metal, polystyrene, latex, nitrocellulose, polypropylene, nylon or combinations thereof. More preferably, the solid support is capable of being attracted to a location by means of a magnetic field, such as a solid support having a magnetite core. Particularly preferred supports are monodisperse magnetic spheres (i.e., uniform size.+-about 5%).

The term "detecting" or "detection" means any of a variety of methods for determining the presence of a nucleic acid, such as, for example, hybridizing a labeled probe to a portion of the nucleic acid. A labeled probe is an oligonucleotide that specifically binds to another sequence and contains a detectable group which may be, for example, a fluorescent moiety, a chemiluminescent moiety (such as an acridinium ester (AE) moiety that can be detected chemilu-

minescently under appropriate conditions (as described in U.S. Pat. No. 5,283,174)), a radioisotope, biotin, avidin, enzyme, enzyme substrate, or other reactive group. Other well known detection techniques include, for example, gel filtration, gel electrophoresis and visualization of the amplicons, and High Performance Liquid Chromatography (HPLC). As used throughout the specification, the term "detecting" or "detection" includes either qualitative or quantitative detection.

The term "primer" or "oligonucleotide primer" means an oligonucleotide capable of binding to a region of a target nucleic acid or its complement and promoting nucleic acid amplification of the target nucleic acid. Generally, a primer will have a free 3' end that can be extended by a nucleic acid polymerase. Primers also generally include a base sequence capable of hybridizing via complementary base interactions either directly with at least one strand of the target nucleic acid or with a strand that is complementary to the target sequence. A primer may comprise target-specific sequences and optionally other sequences that are non-complementary to the target sequence. These non-complementary sequences may comprise a promoter sequence or a restriction endonuclease recognition site.

CaP-Cell-Specific Gene Expression Signature Identification

The present invention is based in part on the identification and validation of consistent CaP epithelial cell specific gene expression signatures. These gene expression signatures define patients with CaP who are at risk to develop advanced disease by identifying genes and pathways in prostate epithelial cells that differentiate between aggressive and non-aggressive courses of cancer development. Two patient groups were selected, a high risk (HR) group having, for example, PSA recurrence, Gleason score 8-9, T3c stage, seminal vesicle invasion, poor tumor differentiation, and a moderate risk (MR) group having, for example, no PSA recurrence, Gleason score 6-7, T2a-T3b stage, no seminal vesicle invasion, well or moderate tumor differentiation. The two patient groups were matched for known risk factors: age, race, and family history of CaP. LCM derived epithelial cells from tumor and normal prostate of the two patient groups were compared by GeneChip analyses, as described in the following Example 1. Results were validated using quantitative reverse transcriptase PCR (QRT-PCR), as described in the following Example 2. The group of genes identified and validated as having the highest association with aggressive or non-aggressive CaP can be used to reliably determine the likely course of CaP progression.

Strikingly, one of the most consistently over expressed genes in CaP cells identified in this study was the ERG (ETS related gene) oncogene, a member of the ETS transcription factor family. (Oikawa et al., Gene (2003) 303:11-34; Sharrocks, A D, Nat Rev Mol Cell Biol (2001) 2(11):827-37; Hart et al., Oncogene (1995) 10(7):1423-30; Rao et al., Science (1987) 237(4815): 635-639). Two isoforms of ERG, ERG1 and ERG2, are over expressed with the highest frequency. The ERG1 coding sequence (with start and stop codons underlined) is publicly available through GenBank under the accession number M21535, as follows:

(SEQ ID NO: 1)

```

1  gaattccctc caaagcaaga caaatgactc acagagaaaa aagatggcag aaccaagggc
61  aactaaagcc gtcaggttct gaacagctgg tagatgggct ggcttactga aggacatgat
121  tcgactgtc cgggaccag cagtcatat caaggaactc tcctgatgaa tgcagtgtgg
181  ccaaaggcgg gaagatgggtg ggcagcccag acaccgttgg gatgaactac ggcagctaca

```

-continued

241 tggaggagaa gcacatgcca ccccaaaaca tgaccacgaa cgagcgcaga gttatcgtgc
301 cagcagatcc tacgctatgg agtacagacc atgtgcggca gtggctggag tgggcggtga
361 aagaatatgg ccttcagac gtcaacatct tgttattcca gaacatcgat gggaaggaa
421 tgtgaagat gaccaaggac gacttcaga ggctcacc cagctacaac gccgacatcc
481 ttctctcaca tctccactac ctcagagaga ctctcttcc acatttgact tcagatgatg
541 ttgataaagc cttacaaaac tctccacggg taatgcatgc tagaaacaca gatttaccat
601 atgagccccc caggagatca gcctggaccg gtcacggcca cccacgccc cagtcgaaag
661 ctgctcaacc atctccttcc acagtgcaca aaactgaaga ccagcgtcct cagttagatc
721 cttatcagat tcttgacca acaagtagcc gccttgcaaa tccaggcagt ggcagatcc
781 agctttggca gttcctcctg gagtcctgt cggacagctc caactccagc tgcacacct
841 gggaaggcac caacggggag ttcaagatga cggatccga cgaggtggcc cggcgctggg
901 gagagcggaa gagcaaaccc aacatgaact acgataagct cagccgcgcc ctccgttact
961 actatgacaa gaacatcatg accaagggtc atgggaagcg ctacgcctac aagttcgact
1021 tccacgggat cgcacaggcc ctccagcccc acccccggga gtcacatctg tacaagtacc
1081 cctcagacct cccgtacatg ggctcctatc acgcccacc acagaagatg aactttgtgg
1141 cgcaccacc tccagccctc cccgtgacat cttccagttt ttttgcgtcc ccaaacccat
1201 actggaattc accaactggg ggtatatacc ccaacactag gctccccacc agccatatgc
1261 cttctcatct gggcacttac tactaaagac ctggcggagg cttttcccat cagcgtgcat
1321 tcaccagccc atcgccacaa actctatcgg agaacatgaa tcaaaagtgc ctcaagagga
1381 atgaaaaaag ctttactggg gctggggaag gaagccgggg aagagatcca aagactcttg
1441 ggagggagtt actgaagtct tactgaagtc ttactacaga aatgaggagg atgctaaaaa
1501 tgtcacgaat atggacatat catctgtgga ctgacctgtg aaaagacagt gtatgtagaa
1561 gcatgaagtc ttaaggacaa agtgccaaag aaagtgtct taagaaatgt ataaacttta
1621 gagtagagtt tgaatccac taatgcaaac tgggatgaaa ctaaagcaat agaaacaaca
1681 cagtttttag ctaacatacc gtttataatg ccattttaag gaaaactacc tgtatttaaa
1741 aatagtttca tatcaaaaac aagagaaaag acacgagaga gactgtggcc catcaacaga
1801 cgttgatagc caactgcatg gcatgtgctg ttttggtga aatcaaatc attcogtttg
1861 atggacagct gtcagcttcc tcaaactgtg aagatgaccc aaagtttcca actcctttac
1921 agtattaccg ggactatgaa ctaaaagggtg ggactgagga tgtgtataga gtgagcgtgt
1981 gattgtagac agaggggtga agaaggagga ggaagaggca gagaaggagg agaccaggct
2041 gggaaagaaa cttctcaagc aatgaagact ggactcagga catttgggga ctgtgtacaa
2101 tgagttatgg agactcgagg gttcatgcag tcagtgttat accaaaccca gtgttaggag
2161 aaaggacaca gcgtaatgga gaaagggaag tagtagaatt cagaaacaaa aatgcgcatc
2221 tctttctttg tttgtcaaat gaaaatttta actggaattg tctgatattt aagagaaaca
2281 ttcaggacct catcattatg tgggggcttt gttctccaca gggtcaggta agagatggcc
2341 ttcttggtcg ccacaatcag aaatcacgca ggcatttttg gtaggcggcc tccagtttcc
2401 ctttgagtcg cgaacgctgt gcgtttgtca gaatgaagta tacaagtcaa tgttttccc
2461 cctttttata taataattat ataacttatg catttataca ctacgagttg atctcgccca
2521 gccaaagaca cagcaaaaa gagacaatcg atataatgtg gccttgaatt ttaactctgt
2581 atgcttaatg tttacaatat gaagttatta gttcttagaa tgcagaatgt atgtaaaaa
2641 ataagcttgg cctagcatgg caaatcagat ttatacagga gtctgcattt gcactttttt

-continued

2701 tagtgactaa agttgcttaa tgaaaacatg tgctgaatgt tgtggatttt gtgttataat
 2761 ttactttgtc caggaacttg tgcaaggag agccaaggaa ataggatgtt tggcacccaa
 2821 atggcgctcag cctctccagg tcctctctgc cteccctcct gtcttttatt tctagccctc
 2881 tttggaacag gaaggacccc ggggtttcaa ttggagcctc catatttatg cctggaagga
 2941 aagaggccta tgaagctggg gttgtcattg agaaattcta gtccagcacc tggtcacaaa
 3001 tcacccttaa ttctgctatg attaaaatac atttgttgaa cagtgaacaa gctaccactc
 3061 gtaaggcaaa ctgtattatt actggcaaat aaagcgcat ggatagctgc aatttctcac
 3121 tttaaca

Nucleotides 195-1286 of SEQ ID NO:1 represent the¹⁵
 coding sequence of SEQ ID NO:1.

The ERG2 coding sequence is publicly available through
 GenBank under the accession number M17254, as follows
 (with start and stop codons underlined):

(SEQ ID NO: 2)

1 gtccgcgcgt gtccgcgccc gcgtgtgcca gcgcgcgtgc cttggccgtg cgccgccgagc
 61 cgggtgcac taactccctc ggccgcgacg gcgcgcgctaa cctctcggtt attccaggat
 121 ctttggagac ccgaggaaag ccgtgttgac caaaagcaag acaaatgact cacagagaaa
 181 aaagatggca gaaccaaggg caactaaagc cgtcaggttc tgaacagctg gtagatgggc
 241 tggcttactg aaggacatga ttcagactgt cccggaccca gcagctcata tcaaggaagc
 301 cttatcagtt gtgagttagg accagtcgtt gtttgagtgt gctacggaa cgccacacct
 361 ggctaagaca gagatgaccg cgtcctctc cagcgactat ggacagactt ccaagatgag
 421 cccacgcgtc cctcagcagg attggctgtc tcaaccccca gccagggtca ccatcaaaat
 481 ggaatgtaac cctagccagg tgaatggctc aaggaactct cctgatgaat gcagtgtggc
 541 caaaggcggg aagatggtag gcagcccaga caccgttggg atgaactacg gcagctacat
 601 ggaggagaag cacatgccac ccccaaacat gaccacgaac gagcgagag ttatcgtgcc
 661 agcagatcct acgctatgga gtacagacca tgtcggcag tggctggagt gggcggtgaa
 721 agaatatggc cttccagacg tcaacatctt gttattccag aacatcgatg ggaaggaaat
 781 gtgcaagatg accaaggacg acttccagag gctcaccacc agctacaacg ccgacatcct
 841 tctctcacat ctccactacc tcagagagac tcctcttcca catttgactt cagatgatgt
 901 tgataaagcc ttacaaaact ctccacggtt aatgcagctc agaaacacag atttaccata
 961 tgagccccc aggagatcag cctggaccgg tcacggccac cccacgccc agtcgaaagc
 1021 tgctcaacca tctccttcca cagtgcccaa aactgaagac cagcgtcctc agttagatcc
 1081 ttatcagatt cttggaccaa caagtagccg ccttgcaaat ccaggcagtg gccagatcca
 1141 gctttggcag ttctcctg agctcctgtc ggacagctcc aactccagct gcatcacctg
 1201 ggaaggcacc aacggggagt tcaagatgac ggatcccac gaggtggccc ggcgtgggg
 1261 agagcggaag agcaaaccca acatgaacta cgataagctc agccgcgccc tccgttacta
 1321 ctatgacaag aacatcatga ccaagggtcca tgggaagcgc tacgcctaca agttcgactt
 1381 ccacgggatc gccagggccc tccagcccca ccccccggag tcatctctgt acaagtaccc
 1441 ctcagacctc ccgtacatgg gctcctatca cgcacccca cagaagatga actttgtggc
 1501 gccccaccct ccagccctcc ccgtgacatc ttccagtttt ttgctgccc caaaccata
 1561 ctggaattca ccaactgggg gtatataccc caacactagg ctcaccca gccatagcc
 1621 ttctcatctg ggcacttact actaaagacc tggcggaggc ttttccatc agcgtgcatt

-continued

1681 caccagccca tcgccacaaa ctctatcggg gaacatgaat caaaagtggc tcaagaggaa
 1741 tgaaaaaagc tttactgggg ctgggggaagg aagccgggga agagatccaa agactcttgg
 1801 gagggagtta ctgaagtctt actacagaaa tgaggaggat gctaaaaatg tcacgaatat
 1861 ggacatatca tctgtggact gaccttgtaa aagacagtgt atgtagaagc atgaagtctt
 1921 aaggacaaaag tgccaaaagaa agtgggtctta agaaatgtat aaactttaga gtagagtttg
 1981 aatcccacta atgcaaactg ggatgaaact aaagcaatag aaacaacaca gttttgacct
 2041 aacataccgt ttataatgcc attttaagga aaactacctg tatttaaaaa tagtttcata
 2101 tcaaaaaacaa gagaaaagac acgagagaga ctgtggccca tcaacagacg ttgatatgca
 2161 actgcatggc atgtgctggt ttggttgaaa tcaaatatcat tccgtttgat ggacagctgt
 2221 cagctttctc aaactgtgaa gatgacccaa agtttccaac tcctttacag tattaccggg
 2281 actatgaact aaaaggtggg actgaggatg tgtatagagt gagcgtgtga ttgtagacag
 2341 aggggtgaag aaggaggagg aagaggcaga gaaggaggag accaggctgg gaaagaaact
 2401 tctcaagcaa tgaagactgg actcaggaca tttggggact gtgtacaatg agttatggag
 2461 actcgagggt tcatgcagtc agtgtttatc caaacccagt gttaggagaa aggacacagc
 2521 gtaattggaga aagggaagta gtagaattca gaaacaaaaa tgcgcactctc tttctttggt
 2581 tgtcaaatga aaattttaac tggaattgtc tgatatttaa gagaaacatt caggacctca
 2641 tcattatgtg ggggctttgt tctccacagg gtcaggtaag agatggcctt cttggctgcc
 2701 acaatcagaa atcacgcagg cattttgggt aggcggcctc cagttttcct ttgagtcgag
 2761 aacgctgtgc gtttgtcaga atgaagtata caagtcaatg tttttccccc tttttatata
 2821 ataattatat aacttatgca ttatacact acgagttgat ctcggccagc caaagacaca
 2881 cgacaaaaga gacaatcgat ataattgtgc cttgaatttt aactctgtat gcttaattgt
 2941 tacaatatga agttattagt tcttagaatg cagaatgtat gtaataaaaat aagcttggcc
 3001 tagcatggca aatcagattt atacaggagt ctgcatttgc acttttttta gtgactaaag
 3061 ttgcttaatg aaaacatgtg ctgaatgttg tggattttgt gttataattt actttgtcca
 3121 ggaacttgtg caaggagag ccaaggaaat aggatgtttg gcaccc

Nucleotides 257-1645 of SEQ ID NO:2 represent the coding sequence of SEQ ID NO:2.

Validation by QRT-PCR (TaqMan) in microdissected tumor and benign prostate epithelial cells of 20 CaP patients confirmed a consistent, tumor associated over expression of ERG isoforms ERG1 and/or ERG2 in 95% of patients (19 of 20) (FIG. 1A). As a quality test and comparison, the expression of AMACR, a recently identified CaP tissue marker (Rubin et al, JAMA (2002) 287:1662-1670; Luo et al., Cancer Res (2002) 62: 2220-2226), and of GSTP1, a gene known to have decreased expression in CaP (Nelson et al., Ann N Y Acad Sci (2001) 952: 135-144), was also determined (FIGS. 1B and 1C). Robust over expression in CaP cells of 95% of the patients, similarly to ERG, was observed for AMACR, while the GSTP1 expression was significantly decreased in the tumor cells of each CaP patient, confirming the high quality of the tumor and benign LCM specimens and the reliability of the QRT-PCR.

Recently a detailed mapping of the chromosomal region (21q22.2-q22.3) containing the ERG gene, as well as a complete exon-intron structure with 9 alternative transcripts (or isoforms) has been described. (Owczarek et al., Gene (2004) 324: 65-77). The probes on the Affymetrix GeneChip used in our initial discovery of consistent ERG over expres-

sion in CaP, as well as the TaqMan probe designed for the validation experiment, recognize a region specific to the ERG 1 and 2 isoforms only.

Both ERG and ETS are proto-oncogenes with mitogenic and transforming activity. (Sharrocks, A D, Nat Rev Mol Cell Biol (2001) 2(11):827-37; Seth et al., Proc Natl Acad Sci USA (1989) 86:7833-7837). Deregulation or chromosomal reorganization of ERG is linked to Ewing sarcoma, myeloid leukemia and cervical carcinoma. (DeAlva et al., Int J Surg Pathol (2001) 9: 7-17; Simpson et al., Oncogene (1997) 14: 2149-2157; Shimizu et al., Proc Natl Acad Sci USA (1993) 90:10280-284; Papas, et al., Am J Med Genet Suppl. (1990) 7:251-261). ETS2 has been implicated in CaP, but it is over expressed only in a small proportion of CaP specimens. (Liu et al., Prostate (1997) 30:145-53; Semchenko, et al., Oncogene (1998) 17:2883-88). ERG over expression without amplification of DNA copy number was recently reported in acute myeloid leukemia. (Baldue et al., Proc. Natl. Acad. Sci. USA (2004) 101:3915-20). Gavrilov et al., Eur J Cancer (2001) 37:1033-40 examined the expression of various transcription factors, including several proteins from the ETS family, in a very limited number of high-grade prostate cancer samples. Antibodies against the ETS family proteins, Elf-1 and Fli-1, caused intense staining of most of the high-grade prostate cancer

samples. In contrast, ERG protein, while being detected in the noncancerous endothelial cells (microvessels in the stroma) of most samples tested, was detected in only a minority of the high-grade prostate cancers. ETS family proteins have a variety of expression patterns in human tissues. (Oikawa et al., Gene (2003) 303:11-34). ERG is expressed in endothelial tissues, hematopoietic cells, kidney, and in the urogenital tract. ERG proteins are nuclear transcription factors that form homodimers, as well as heterodimers with several other members of the ETS family of transcription factors. (Carrere et al., Oncogene (1998) 16(25): 3261-68). A negative crosstalk observed between ERG and estrogen receptor (ER-alpha) may be relevant in urogenital tissues, where both transcription factors are

expressed. (Vlaeminck-Guillem et al., Oncogene (2003) 22(50):8072-84). The present invention is based in part upon the surprising discovery that ERG is over expressed in the majority of CaP specimens analyzed, indicating that this oncogene plays a role in prostate tumorigenesis, most likely by modulating transcription of target genes favoring tumorigenesis in prostate epithelium.

The present invention is further based in part upon the over expression of the AMACR gene in prostate cancer epithelium. The cDNA sequence of the AMACR is publicly available through GenBank under the accession numbers NM_014324 and AF047020. The sequence (with start and stop codons underlined) corresponding to accession number NM_014324 is as follows:

(SEQ ID NO: 3)

```

1  gggattggga gggcttcttg caggctgctg ggctggggct aagggtctgt cagtttcctt
61  cagcggggca ctgggaagcg ccatggcact gcagggcac tcggctcgtg agctgtccgg
121 cctggccccc ggcccggtct gtgctatggt cctggctgac ttcggggcgc gtgtggtacg
181 cgtggaccgg cccggctccc gctacgacgt gagccgcttg ggcgggggca agcgtctcgt
241 agtgctggac ctgaagcagc cgcggggagc cgccgtgctg cgcgctctgt gcaagcggtc
301 ggatgtgctg ctggagccct tccgccgcgg tgcctatggag aaactccagc tgggccccaga
361 gattctgcag cgggaaaatc caagccttat ttatgccagg ctgagtggat ttggccagtc
421 aggaagcttc tgccgggttag ctggccacga tatcaactat ttggctttgt caggtgttct
481 ctcaaaaatt ggcagaagtg gtgagaatcc gtatgccccg ctgaatctcc tggtgacttc
541 tgctgggtgt ggcccttatgt gtgactggg cattataatg gctctttttg accgcacacg
601 cactggcaag ggtcagggtc ttgatgcaaa tatggtggaa ggaacagcat atttaagttc
661 ttttctgttg aaaactcaga aattgagtct gtgggaagca cctcgaggac agaacatggt
721 ggatggtgga gcacctttct atacgactta caggacagca gatggggaat tcatggctgt
781 tggagcaata gaacccagct tctacgagct gctgatcaaa ggacttggac taaagtctga
841 tgaacttccc aatcagatga gcatggatga ttggccagaa atgaagaaga agtttcgaga
901 tgtatttgca gagaagacga aggcagagtg gtgtcaaatc tttgacggca cagatgcctg
961 tgtgactccg gttctgactt ttgaggaggt tgttcacat gatcacaaca aggaacgggg
1021 ctggtttatc accagtgagg agcaggacgt gagccccgc cctgcacctc tgctgttaaa
1081 caccacagcc atcccttctt tcaaaaggga tcctttcata ggagaacaca ctgaggagat
1141 acttgaagaa tttggattca gccgcgaaga gatttatcag cttaactcag ataaaatcat
1201 tgaagtaat aaggtaaaag ctagtctcta acttcaggc ccacggctca agtgaatttg
1261 aatactgcat ttacagtgtg gagtaacaca taacattgta tgcattgaaa catggaggaa
1321 cagtattaca gtgtcctacc actctaatac agaaaagaat tacagactct gattctacag
1381 tgatgattga attctaaaaa tgggttatcat tagggctttt gatttataaa accttgggta
1441 cttatactaa attatggtag ttattctgcc ttccagtttg ctgatataat ttgttgatat
1501 taagattctt gacttatatt ttgaatgggt tctagtgaac aaggaatgat atattcttga
1561 agacatcgat atacatttat ttacactctt gattctacaa tgtagaaaaa gaggaaatgc
1621 cacaaattgt atggtgataa aagtcacgtg aaacagagtg attggttgca tccaggcctt
1681 ttgtcttggt gttcatgac tcctctaaag cacattccaa actttagcaa cagttatcac
1741 accttgtaat ttgcaaagaa aagtttcacc tgtattgaat cagaatgcct tcaactgaaa
1801 aaaacatata caaaataatg aggaatgtg ttggctcact acgtagagtc cagagggaca

```

-continued

1861 gtcagtttta ggggtgcctg tatccagtaa ctcggggcct gtttccccgt ggggtctctgg
 1921 gctgtcagct ttcttttctc catgtgtttg atttctctc aggetggtag caagtctctg
 1981 atctttatac caacacacag caacatccag aaataaagat ctccaggacc cccagcaagt
 2041 cgttttgtgt ctcttggac tgagttaagt tacaagcctt tctttacct gtctttgaca
 2101 aagaagacgg gattgtcttt acataaaacc agcctgctcc tggagcttcc ctggactcaa
 2161 ctctctaaag gcatgtgagg aaggggtaga ttccacaatc taatccgggt gccatcagag
 2221 tagagggagt agagaatgga tgttgggtag gccatcaata aggtccattc tgcgcagtat
 2281 ctcaactgcc gttcaacaat cgcaagagga aggtggagca ggtttcttca tcttacagtt
 2341 gagaaaacag agactcagaa gggtctctta gttcatgttt cccttagcgc ctcagtgatt
 2401 ttttcatggt ggcttaggcc aaaagaaata tctaaccatt caatttataa ataattaggt
 2461 ccccaacgaa ttaaatatta tgtcctacca acttattagc tgcctgaaaa atataatata
 2521 cataaataaa aaaa

Nucleotides 83-1231 of SEQ ID NO:3 represent the coding sequence of AMACR.

The present invention is further based in part upon the over expression of the DD3 gene in prostate cancer epithe-

lium. The cDNA sequence of the DD3 gene is publicly available through GenBank under the accession number AF103907. The sequence corresponding to accession number AF103907 is as follows:

(SEQ ID NO: 4)

1 acagaagaaa tagcaagtgc cgagaagctg gcatcagaaa aacagagggg agatttgtgt
 61 ggctgcagcc gagggagacc aggaagatct gcatgggtgg aaggacctga tgatacagag
 121 gaattacaac acatatactt agtgtttcaa tgaacaccaa gataaataag tgaagagcta
 181 gtccgctgtg agtctctca gtgacacagg gctggatcac catcgacggc actttctgag
 241 tactcagtc agcaaagaaa gactacagac atctcaatgg caggggtgag aaataagaaa
 301 ggctgctgac ttaccatct gaggccacac atctgctgaa atggagataa ttaacatcac
 361 tagaaacagc aagatgacaa tataatgtct aagtagtgac atgtttttgc acatttccag
 421 cccctttaaa tatccacaca cacaggaagc acaaaaggaa gcacagagat cccctgggaga
 481 aatgcccgcc cgccatcttg ggtcatcgat gagcctcgcc ctgtgcctgg tcccgttgt
 541 gaggggaagga cattagaaaa tgaattgatg tgttcttaa aggatgggca ggaaaacaga
 601 tcctgttgtg gatatttatt tgaacgggat tacagatttg aaatgaagtc acaaagttag
 661 cattaccaat gagaggaaaa cagacgagaa aatcttgatg gcttcacaag acatgcaaca
 721 aacaaaatgg aatactgtga tgacatgagg cagccaagct ggggaggaga taaccacggg
 781 gcagagggtc aggattctgg ccctgctgcc taaactgtgc gtccataacc aaatcatttc
 841 atatttctaa ccctcaaaac aaagctgttg taatatctga tctctacggg tcttctggg
 901 cccaacattc tccatatatc cagccacact catttttaat atttagttcc cagatctgta
 961 ctgtgacctt tctacactgt agaataacat tactcatttt gttcaagac ccttcgtgtt
 1021 gctgcctaata atgtagctga ctgtttttcc taaggagtgt tctggcccag gggatctgtg
 1081 aacaggctgg gaagcatctc aagatcttcc caggggtata ctactagca cacagcatga
 1141 tcattacgga gtgaattatc taatcaacat catcctcagt gtctttgccc atactgaaat
 1201 tcatttccca cttttgtgcc cattctcaag acctcaaaat gtcattccat taatatcaca
 1261 ggattaactt ttttttttaa cctggaagaa ttcaatgtta catgcagcta tgggaattta
 1321 attacatatt ttgttttcca gtgcaaagat gactaagtcc tttatccctc ccctttgttt
 1381 gatttttttt ccagtataaa gttaaaatgc ttagccttgt actgaggtg tatacagcac

-continued

1441 agcctctccc catccctcca gccttatctg tcatcaccat caaccctcc cataccacct
1501 aaaaaaate taacttgtaa ttccttgaa atgtcaggac atacattatt ccttctgcct
1561 gagaagctct tccctgtctc ttaaactag aatgatgtaa agttttgaat aagttgacta
1621 tcttacttca tgcaagaag ggacacatat gagattcatc atcacatgag acagcaaata
1681 ctaaaagtgt aatttgatta taagagtta gataaatata tgaaatgcaa gagccacaga
1741 gggaatgttt atggggcacg tttgtaagcc tgggatgtga agcaaaggca gggaacctca
1801 tagtatctta tataatatat ttcatttctc tatctctatc acaatatcca acaagctttt
1861 cacagaattc atgcagtga aatcccaaaa ggtaaccttt atccatttca tggtagtgcc
1921 gctttagaat tttggcaaat catactggtc acttatctca actttgagat gtgtttgtcc
1981 ttgtagttaa ttgaagaaa tagggcactc ttgtgagcca ctttaggggt cactcctggc
2041 aataaagaat ttacaaagag ctactcagga ccagttgtta agagctctgt gtgtgtgtgt
2101 gtgtgtgtgt gagtgatcat gccaaagtgt gcctctctct cttgacccat tatttcagac
2161 ttaaaaaaag catgttttca aatggcacta tgagctgcca atgatgtatc accaccatat
2221 ctcattatct tccagtaaat gtgataataa tgtcatctgt taacataaaa aaagtttgac
2281 ttcacaaaag cagctggaaa tggacaacca caatatgcac aaatctaact cctaccatca
2341 gctacacact gcttgacata tattgttaga agcacctcgc atttgtgggt tctcttaagc
2401 aaaatacttg cattaggtct cagctggggc tgtgcatcag gcggtttgag aaatattcaa
2461 ttctcagcag aagccagaat ttgaattccc tcactcttta ggaatcattt accaggtttg
2521 gagaggattc agacagctca ggtgctttca ctaatgtctc tgaacttctg tccctctttg
2581 tgttcatgga tagtccaata aataatgtta tctttgaact gatgctcata ggagagaata
2641 taagaactct gagtgatctc aacattaggg attcaaagaa atattagatt taagctcaca
2701 ctgggtcaaaa ggaaccaaga tacaaagaac tctgagctgt catcgtcccc atctctgtga
2761 gccacaacca acagcaggac ccaacgcatg tctgagatcc ttaaatcaag gaaaccagtg
2821 tcatgagttg aattctctca ttatggatgc tagcttctgg ccatctctgg ctctcctctt
2881 gacacatat agcttctagc ctttgcttcc acgactttta tctttctctc aacacatcgc
2941 ttaccaatcc tctctctgct ctggtgcttt ggacttcccc acaagaattt caacgactct
3001 caagtctttt cttccatccc caccactaac ctgaatgcct agacccttat ttttattaat
3061 ttccaataga tgctgcctat gggctatat gctttagatg aacattagat atttaaagct
3121 caagaggttc aaaaaccaac tcattatctt ctctttcttt cactccctg ctctctccc
3181 tatattactg attgcactga acagcatggt cccaatgta gccatgcaaa tgagaaaccc
3241 agtggctcct tgtggtacat gcatgcaaga ctgctgaagc cagaaggatg actgattacg
3301 cctcatgggt ggaggggacc actcctgggc cttcgtgatt gtcaggagca agacctgaga
3361 tgctccctgc cttcagtgtc ctctgcatct cccctttcta atgaagatcc atagaatttg
3421 ctacatttga gaattccaat taggaactca catgttttat ctgccctatc aattttttaa
3481 acttgctgaa aattaagttt tttcaaaatc tgccttgta aattactttt tcttacagtg
3541 tcttggcata ctatatcaac tttgattctt tgttacaact tttcttactc ttttatcacc
3601 aaagtggctt ttattctctt tattattatt atttctttt actactatat tacgttggtta
3661 ttattttggt ctctatagta tcaatttatt tgatttagtt tcaatttatt tttattgctg
3721 acttttaaaa taagtattc ggggggtggg agaacagggg agggagagca ttaggacaaa
3781 tacctaagtc atgtgggact taaaacctag atgatgggtt gatagtgca gcaaacctac

-continued

3841 atggcacacg tatacctgtg taacaaacct acacattctg cacatgtatc ccagaacgta

3901 aagtaaaatt taaaaaaaaag tga

5

The DD3 gene appears to represent a non-coding nucleic acid. Therefore, no start and stop codons have been indicated.

The present invention is further based in part upon the under expression of the LTF gene in prostate cancer epithe-

lium. The cDNA sequence of the lactotransferrin (LTF) gene is publicly available through GenBank under the accession number NM_002343. The sequence (with start and stop codons underlined) corresponding to accession number NM_002343 is as follows:

(SEQ ID NO: 5)

```

1 agagccttcg tttgccaagt cgctccaga cgcagacat_gaaacttgtc ttccctcgtec
61 tgctgttctt cggggccctc ggactgtgtc tggctggccg taggaggagt gtccagtggg
121 ggcgcgtatc ccaaccgcag gccacaaaat gcttccaatg gcaaaggaat atgagaaaaa
181 tgcgtggccc tctgtgcagc tgcataaaga gagactcccc catccagtgt atccaggcca
241 ttgcggaaaa cagggccgat gctgtgacct ttgatggtgg ttccatatac gaggcaggcc
301 tggcccccta caaactgcga cctgtagcgg cggaagtcta cgggaccgaa agacagccac
361 gaactcacta ttatgccgtg gctgtggtga agaagggcgg cagctttcag ctgaacgaac
421 tgcaagggtt gaagtctgc caccacaggcc ttgcaggac cgtcggatgg aatgtcccta
481 tagggacact tcgtccattc ttgaattgga cgggtccacc tgagccatt gaggcagctg
541 tggccagggt cttctcagcc agctgtgttc ccggtgcaga taaggacag ttccccaacc
601 tgtgtgcct gtgtgcgggg acaggggaaa acaaatgtgc cttctcctcc caggaaccgt
661 acttcagcta ctctggtgcc ttcaagtgtc tgagagacgg ggtcggagac ttggctttta
721 tcagagagag cacagtgttt gaggacctgt cagacaggc tgaaggagac gagtatgagt
781 tactctgccc agacaacact cggaagccag tggacaagtt caaagactgc catctggccc
841 gggctccctc tcatgccgtt gtggcacgaa gtgtgaatgg caaggaggat gccatctgga
901 atcttctccg ccaggcacag gaaaagtgtt gaaaggacaa gtcaccgaaa ttccagctct
961 ttggctcccc tagtgggcag aaagatctgc tgttcaagga ctctgccatt gggttttcga
1021 ggggtccccc gaggatagat tctgggctgt acctggctc cggctacttc actgccatcc
1081 agaacttgag gaaaagttag gaggaagtgg ctgcccggcg tgcgcgggtc gtgtggtgtg
1141 cgggtggcga gcaggagctg cgcaagtgt accagtggag tggcttgagc gaaggcagcg
1201 tgacctgtct ctgcgcctcc accacagagg actgcacgc cctggtgctg aaaggagaag
1261 ctgatgccat gagtttgat ggaggatatg tgtacactgc aggcacaaatgt ggtttggtgc
1321 ctgtcctggc agagaactac aaatcccaac aaagcagtga cctgatcct aactgtgtgg
1381 atagacctgt ggaaggatat ctgtgtgtg cggtggttag gagatcagac actagcetta
1441 cctggaactc tgtgaaaggc aagaagtcct gccacaccgc cgtggacagg actgcaggct
1501 ggaatatccc catgggcctg ctcttcaacc agacgggtc ctgcaaatgt gatgaatatt
1561 tcagtcaaaag ctgtgccctt gggctcgacc cgagatctaa tctctgtgct ctgtgtattg
1621 gcgacgagca ggggtgagaat aagtgcgtgc ccaacagcaa cgagagatac tacggctaca
1681 ctggggcttt ccgtgcctg gctgagaatg ctggagacgt tgcatttgtg aaagatgtca
1741 ctgtcttgca gaacctgat ggaaataaca atgaggcatg ggctaaggat ttgaagctgg
1801 cagactttgc gctgctgtgc ctcgatggca aacggaagcc tgtgactgag gctagaagct
1861 gccatcttgc catggccccg aatcatgccg tgggtgtctc gatggataag ttggaacgcc
1921 tgaaacaggt gttgctccac caacaggcta aatttgggag aaatggatct gactgccggg
1981 acaagttttg cttattccag tctgaaacca aaaaccttct gttcaatgac aacactgagt

```

-continued

2041 gctctggccag actccatggc aaaacaacat atgaaaaata ttggggacca cagtatgtcg
 2101 caggcattac taatctgaaa aagtgtctcaa cctccccct cctggaagcc tgtgaattcc
 2161 tcaggaagta aaaccgaaga agatggccca gctccccaag aaagctcag ccattcactg
 2221 cccccagctc ttctccccag gtgtgttggg gccttggcct cccctgtctga aggtggggat
 2281 tgccccacca tctgcttaca attccctgct gtcgtcttag caagaagtaa aatgagaaat
 2341 ttgtgtgata ttctctcctt aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa

Nucleotides 39-2171 of SEQ ID NO:5 represent the coding sequence of LTF.

LTF is a non-heme iron binding glycoprotein and a member of the transferring gene family. Bowman et al., *Adv. Genet.* 25:1-38 (1988); Park et al., *Proc. Natl. Acad. Sci. U.S.A.*, 82:3149-53 (1985). The concentration of LTF in human prostate is hormone dependent and its expression is regulated by estrogen. van Sande et al., *Urol. Res.*, 9(5): 241-244 (1981); Teng et al., *Biochem. Cell Biol.*, 80:7-16 (2002); Teng et al., *Mol. Human Reproduction.*, 8, (1):58-67 (2002). LTF has also been implicated in certain cancers. For example, bovine LTF inhibits colon, esophagus, lung, and bladder carcinomas in rats. Tsuda et al., *Biochem. Cell Biol.*, 80:131-136 (2002); Tsuda et al., *Biofactors.*, 12(1-4):83-8 (2000); Tsuda et al., *Biofactors.*, 12(1-4):83-8 (2000); Tsuda et al., *Mutat Res.*, 462(2-3):227-33 (2000). In a study published over 20 years ago, van Sande et al., *Urol. Res.* 9:241-244 (1981), examined lactoferrin protein levels in human benign prostatic hypertrophy samples. They also detected low levels of lactoferrin protein in 3 carcinoma samples. However, we are the first to report the consistent and significant under expression of LTF mRNA in prostate cancer epithelial cells from a large number of patient samples. The observed under expression of LTF mRNA in such a statistically significant sample size indicates that under expression of LTF is a useful diagnostic marker for prostate cancer.

In one experiment, when screened using the Affymetrix GeneChip, CaP tumor cells exhibited upregulated AMACR expression in comparison to matched benign cells. In this studied patient cohort (n=73), AMACR was upregulated in tumor compared to matched benign prostate epithelium in 89.04% of the patients (65 of 73), while ERG was upregulated in 78.08% (57 of 73). When these two markers were combined, we observed a 100% CaP detection rate (under the criteria that at least one marker was upregulated) in the studied patient cohort (73 of 73). These data indicate that the combination of ERG and AMACR screening provides a highly accurate tool for CaP detection.

In another experiment, 96.4% of patients showed upregulation of either the ERG or AMACR gene in laser microdissected matched tumor and benign prostate epithelial cells from 55 CaP patients (FIG. 5). Similarly, 96.4% of patients showed upregulation of either the ERG or DD3 gene (FIG. 5). When the expression data for the ERG, AMACR, and DD3 genes was combined, 98.2% of the CaP patients showed upregulation of at least one of the three genes in tumor cells (FIG. 5). Thus, the combination of ERG, AMACR, and DD3 screening also provides a highly accurate tool for CaP detection.

In yet another experiment, validation by QRT-PCR (TaqMan) in microdissected tumor and benign prostate epithelial cells of 20 CaP patients confirmed a consistent, tumor associated under expression of LTF in 100% of patients (20 of 20) (FIG. 1D). Further validation studies by QRT-PCR in

microdissected tumor and benign prostate epithelial cells of 103 CaP patients were consistent with the initial results, showing tumor associated under expression in 76% of patients (78 of 103).

Diagnostic Uses

In one embodiment, the present invention comprises a method of CaP diagnosis comprising screening biological samples for CaP-cell-specific gene expression signatures. In particular, the invention comprises screening for at least one of the CaP-cell-specific genes listed in Tables 1-6, particularly the ERG gene, the AMACR gene, the LTF gene or a combination of the ERG gene and the AMACR genes. The invention also comprises methods of diagnosing CaP comprising screening biological samples for expression of the ERG and DD3 genes, or a combination of the ERG, DD3, and AMACR genes.

In a further embodiment, the present invention comprises a method of CaP diagnosis comprising screening biological samples for CaP-cell-specific gene expression signatures using methods known in the art, including, for example, immunohistochemistry, ELISA, in situ RNA hybridization, and any oligonucleotide amplification procedure known or later developed, including PCR (including QRT-PCR), transcription-mediated amplification (TMA), nucleic acid sequence-based amplification (NASBA), self-sustained sequence replication (3SR), ligase chain reaction (LCR), strand displacement amplification (SDA), and Loop-Mediated Isothermal Amplification (LAMP). See, e.g., Mullis, U.S. Pat. No. 4,683,202; Erlich et al., U.S. Pat. No. 6,197,563; Walker et al., *Nucleic Acids Res.*, 20:1691-1696 (1992); Fahy et al., *PCR Methods and Applications*, 1:25-33 (1991); Kacian et al., U.S. Pat. No. 5,399,491; Kacian et al., U.S. Pat. No. 5,480,784; Davey et al., U.S. Pat. No. 5,554,517; Birkenmeyer et al., U.S. Pat. No. 5,427,930; Marshall et al., U.S. Pat. No. 5,686,272; Walker, U.S. Pat. No. 5,712,124; Notomi et al., European Patent Application No. 1 020 534 A1; Dattagupta et al., U.S. Pat. No. 6,214,587; and HELEN H. LEE ET AL., *NUCLEIC ACID AMPLIFICATION TECHNOLOGIES: APPLICATION TO DISEASE DIAGNOSIS* (1997). Each of the foregoing amplification references is hereby incorporated by reference herein. In particular, the invention comprises generating antibodies to CaP-cell-specific genes, including ERG, AMACR, LTF, and DD3 for use in an immunohistochemistry assay. Other known diagnostic assays may be used to detect gene expression.

In a specific embodiment, the present invention comprises a method of diagnosing CaP comprising screening biological samples for expression of the ERG and AMACR genes, the ERG and DD3 genes, or the ERG, AMACR, and DD3 genes, or the LTF gene using methods known in the art, including, for example, immunohistochemistry, ELISA, in situ hybridization, PCR (including QRT-PCR), transcription-mediated amplification (TMA), nucleic acid sequence-based amplification (NASBA), self-sustained sequence replication (3SR), ligase chain reaction (LCR), strand

displacement amplification (SDA), and Loop-Mediated Isothermal Amplification (LAMP).

ERG, LTF, or AMACR polypeptides, their fragments or other derivatives, or analogs thereof, may be used as immunogens in order to generate antibodies that specifically bind such immunogens. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain and Fab fragments. In a specific embodiment, antibodies to a human ERG, LTF or AMACR protein are produced. Antibodies can then be used in standard diagnostic assays to detect the protein produced by the desired gene.

Various procedures known in the art may be used for the production of polyclonal antibodies to an ERG, LTF, or AMACR protein or derivative or analog. In a particular embodiment, rabbit polyclonal antibodies to an epitope of a ERG, LTF, or AMACR protein can be obtained. For the production of antibody, various host animals can be immunized by injection with the native ERG, LTF, or AMACR protein, or a synthetic version, or derivative (e.g., fragment) thereof, including but not limited to rabbits, mice, rats, etc. Various adjuvants may be used to increase the immunological response, depending on the host species, and including, but not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum.

For preparation of monoclonal antibodies directed toward a ERG, LTF, or AMACR protein sequence or analog thereof, any technique, which provides for the production of antibody molecules by continuous cell lines in culture may be used. For example, the hybridoma technique originally developed by Kohler et al (1975) *Nature*, 256:495-497, as well as the trioma technique, the human B-cell hybridoma technique (Kozbor et al. (1983) *Immunology Today*, 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al. (1985) *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). According to the invention, human antibodies may be used and can be obtained by using human hybridomas (Cote et al. (1983) *Proc. Natl. Acad. Sci. U.S.A.*, 80:2026-2030) or by transforming human B cells with EBV virus in vitro (Cole et al. (1985) *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, pp. 77-96). According to the invention, techniques developed for the production of chimeric antibodies (Morrison et al. (1984) *Proc. Natl. Acad. Sci. U.S.A.*, 81:6851-6855; Neuberger et al. (1984) *Nature*, 312:604-608; Takeda et al. (1985) *Nature*, 314:452-454) by splicing the genes from a mouse antibody molecule specific for ERG, LTF, or AMACR together with genes from a human antibody molecule of appropriate biological activity can be used; such antibodies are within the scope of this invention.

Techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778) can be used to produce ERG-, LTF-, or AMACR-specific single chain antibodies. An additional embodiment of the invention utilizes the techniques described for the construction of Fab expression libraries (Huse et al. (1989) *Science*, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for ERG, LTF or AMACR proteins, derivatives, or analogs.

Antibody fragments which contain the idiotype of the molecule can be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragment which can be produced by pepsin digestion

of the antibody molecule; the Fab' fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragment, the Fab fragments which can be generated by treating the antibody molecule with papain and a reducing agent, and Fv fragments, including single chain Fv (scFv) fragments.

In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, e.g., ELISA. For example, to select antibodies that recognize a specific domain of a ERG, LTF, or AMACR protein, one may assay generated hybridomas for a product which binds to a ERG, LTF, or AMACR fragment containing such domain.

A second aspect of the invention provides for use of the expression profiles resulting from these methods in diagnostic methods, including, but not limited to, characterizing the treatment response to any therapy, correlating expression profiles with clinico-pathologic features, distinguishing indolent prostate cancers from those with a more aggressive phenotype (e.g. moderate risk versus high risk), analyzing tumor specimens of patients treated by radical prostate surgery to help define prognosis, screening candidate genes for the development of a polynucleotide array for use as a blood test for improved prostate cancer detection, and identifying further genes that may serve as biomarkers for response to treatment to screen drugs for the treatment of advanced prostate cancer.

As will be readily appreciated by persons having skill in the art, the ERG, LTF, DD3, and/or the AMACR nucleic acid sequences described herein can easily be synthesized directly on a support, or pre-synthesized polynucleotide probes may be affixed to a support as described, for example, in U.S. Pat. Nos. 5,744,305, 5,837,832, and 5,861,242, each of which is incorporated herein by reference.

Such arrays may be used to detect specific nucleic acid sequences contained in a target cell or sample, as described in U.S. Pat. Nos. 5,744,305, 5,837,832, and 5,861,242, each of which is incorporated herein by reference. More specifically, in the present invention, these arrays may be used in methods for the diagnosis or prognosis of prostate cancer, such as by assessing the expression profiles of genes, in biological samples. In a preferred embodiment, computer models may be developed for the analysis of expression profiles. Moreover, such polynucleotide arrays are useful in methods to screen drugs for the treatment of advanced prostate cancer. In these screening methods, the polynucleotide arrays are used to analyze how drugs affect the expression of the ERG, LTF, AMACR, and/or DD3 genes.

Therapeutic Uses

The invention provides for treatment or prevention of various diseases and disorders by administration of a therapeutic compound (termed herein "therapeutic"). "Therapeutics" include but are not limited to: ERG or LTF proteins and analogs and derivatives (including fragments) thereof (e.g., as described herein above); nucleic acids encoding the ERG or LTF proteins, analogs, or derivatives; ERG or LTF antisense nucleic acids, ERG or LTF dominant negative mutants, siRNA against ERG or LTF, ERG or LTF antibodies and ERG or LTF agonists and antagonists. ERG or LTF agonists and antagonists, including small molecules, can be identified using the methods disclosed in this application or any standard screening assay to identify agents that modulate ERG or LTF expression or function, particularly in prostate cancer cells. For example, ERG or LTF expression or function can be readily detected, e.g., by obtaining a biological sample from a patient, e.g., a tissue sample (e.g., from biopsy tissue), a blood sample, or a urine sample, and

assaying it in vitro for mRNA or protein levels, structure and/or activity of the expressed ERG or LTF mRNA or protein. Many methods standard in the art can be employed, including but not limited to, kinase assays, immunoassays to detect and/or visualize ERG or LTF protein (e.g., Western blot, immunoprecipitation followed by SDS-PAGE, immunocytochemistry, etc.) and/or hybridization assays to detect ERG or LTF expression by detecting and/or visualizing ERG or LTF mRNA (e.g., Northern assays, dot blots, in situ hybridization, PCR (including RT-PCR), TMA, NASAB, 3SR, LCR, SDA, LAMP, etc.).

Hyperproliferative Disorders

Disorders involving hyperproliferation of cells are treated or prevented by administration of a therapeutic that antagonizes (reduces or inhibits) ERG function or expression or enhances LTF function or expression. In certain embodiments, ERG function is inhibited by use of ERG antisense nucleic acids. The present invention provides the therapeutic or prophylactic use of nucleic acids of at least 10, 15, 100, 200, 500, 1000, 1500, 2000, or 2500 contiguous nucleotides in antisense to any of the ERG nucleotides described herein. In a particular embodiment, the ERG antisense nucleic acid comprises at least 10, 15, 100, 200, 500, 1000, 1500, 2000, or 2500 contiguous nucleotides in antisense orientation to the ERG nucleotide sequence. An ERG "antisense" nucleic acid as used herein refers to a nucleic acid capable of hybridizing under defined conditions to a portion of an ERG nucleic acid by virtue of some sequence complementarity. The antisense nucleic acid may be complementary to a coding and/or noncoding region of an ERG nucleic acid. Such antisense nucleic acids have utility as therapeutics that inhibit ERG function, and can be used in the treatment or prevention of disorders as described herein.

The antisense nucleic acids of the invention can be oligonucleotides that are double-stranded or single-stranded, RNA or DNA or a modification or derivative thereof, which can be directly administered to a cell, or which can be produced intracellularly by transcription of exogenously introduced coding sequences.

The dominant negative mutants of the invention can be produced by expression plasmids containing a nucleic acid encoding a non-functional domain of ERG, such as the DNA binding domain of ERG. These expression plasmids can be introduced into a target cell or tissue and can induce tumor growth inhibition and apoptosis by acting as a dominant negative form against the wild-type ERG transcription factors influencing cell hyperproliferation (Oikawa, *Cancer Sci* (2004), 95:626-33).

RNA interference can be achieved using siRNA against the ERG gene. The siRNA is a short double stranded RNA molecule of about 18-25 nucleotides that comprises a nucleotide sequence complementary to a region of the target gene. The siRNA can be introduced into a target cell or tissue, for example using an expression plasmid, where it interferes with the translation of the ERG gene. RNA interference techniques can be carried out using known methods as described, for example, in published U.S. Patent Applications 20040192626, 20040181821, and 20030148519, each of which is incorporated by reference.

Therapeutics which are useful according to this embodiment of the invention for treatment of a disorder may be selected by testing for biological activity in promoting the survival or differentiation of cells. For example, in a specific embodiment relating to cancer therapy, including therapy of prostate cancer, a therapeutic decreases proliferation of tumor cells. These effects can be measured as described in the Examples or using any other method standard in the art.

In specific embodiments, malignancy or dysproliferative changes (such as metaplasias and dysplasias), or hyperproliferative disorders, are treated or prevented in the prostate.

The therapeutics of the invention that antagonize ERG activity can also be administered to treat premalignant conditions and to prevent progression to a neoplastic or malignant state, including but not limited to those disorders described herein, such as prostate cancer.

Gene Therapy

In a specific embodiment, nucleic acids comprising a sequence encoding an ERG or LTF protein or functional derivative thereof, are administered to promote ERG or LTF function, by way of gene therapy. Alternatively, nucleic acids comprising an antisense ERG sequence are administered to antagonize ERG expression or function. Gene therapy refers to therapy performed by the administration of a nucleic acid to a subject.

Any of the methods for gene therapy available in the art can be used according to the present invention. For specific protocols, see Morgan (2001) *Gene Therapy Protocols*, 2nd ed., Humana Press. For general reviews of the methods of gene therapy, see Goldspiel et al. (1993) *Clinical Pharmacy*, 12:488-505; Wu et al. (1991) *Biotherapy*, 3:87-95; Tolstoshev (1993) *Ann. Rev. Pharmacol. Toxicol.*, 32:573-596; Mulligan (1993) *Science*, 260:926-932; and Morgan et al. (1993) *Ann. Rev. Biochem.*, 62:191-217; May (1993) *TIBTECH*, 11(5):155-215. Methods commonly known in the art of recombinant DNA technology which can be used are described in *Current Protocols in Molecular Biology* (2004), Ausubel et al., eds., John Wiley & Sons, NY; and Kriegler (1990) *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY.

In one embodiment, the therapeutic comprises an ERG or LTF nucleic acid or antisense ERG nucleic acid that is part of a vector. In particular, such a nucleic acid has a regulatory sequence, such as a promoter, operably linked to the ERG or LTF coding region or antisense molecule, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, a nucleic acid molecule is used in which the ERG or LTF coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the ERG or LTF nucleic acid (Koller et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.*, 86:8932-8935; Zijlstra et al. (1989) *Nature*, 342:435-438).

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the desired nucleic acids, such that expression of the nucleic acid is controllable by the appropriate inducer of transcription.

Delivery of the nucleic acid into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vector, or indirect, in which case, cells are first transformed with the nucleic acid in vitro, then transplanted into the patient. These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

In a specific embodiment, the nucleic acid is directly administered in vivo, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by infection using a defective or attenuated retroviral or other viral vector (see U.S. Pat. No. 4,980,286, which is incorporated herein by reference), or by direct injection of naked

DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, DuPont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering it in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu et al. (1987) *J. Biol. Chem.*, 262:4429-4432). In another embodiment, a nucleic acid-ligand complex can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell-specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Pubs. WO 92/06180; WO 92/22635; WO92/20316; WO93/14188; WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.*, 86:8932-8935; Zijlstra et al. (1989) *Nature*, 342:435-438).

In a specific embodiment, a viral vector that contains an ERG or LTF nucleic acid is used. For example, a retroviral vector can be used (see, Miller et al. (1993) *Meth. Enzymol.*, 217:581-599). These retroviral vectors have been modified to delete retroviral sequences that are not necessary for packaging of the viral genome and integration into host cell DNA. The ERG or LTF nucleic acid to be used in gene therapy is cloned into the vector, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al. (1994) *Biotherapy*, 6:291-302, which describes the use of a retroviral vector to deliver the MDRL gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al. (1994) *J. Clin. Invest.*, 93:644-651; Kiem et al. (1994) *Blood*, 83:1467-1473; Salmons et al. (1993) *Hum. Gene Ther.*, 4:129-141; and Grossman et al. (1993) *Curr. Opin. Gen. Devel.*, 3:110-114.

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky et al. (1993, *Curr. Opin. Gen. Devel.*, 3:499-503) present a review of adenovirus-based gene therapy. Bout et al. (1994, *Hum. Gene Ther.*, 5:3-10) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al. (1991) *Science*, 252:431-434; Rosenfeld et al. (1992) *Cell*, 68:143-155; and Mstrangeli et al. (1993) *J. Clin. Invest.*, 91:225-234.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al. (1993) *Proc. Soc. Exp. Biol. Med.*, 204:289-300).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recom-

binant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, e.g., Loeffler et al. (1993) *Meth. Enzymol.*, 217:599-618; Cohen et al. (1993) *Meth. Enzymol.*, 217:618-644; Cline (1985) *Pharmac. Ther.*, 29:69-92) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. In one preferred embodiment, epithelial cells are injected, e.g., subcutaneously. In another embodiment, recombinant skin cells may be applied as a skin graft onto the patient. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) may be administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include, but are not limited to, epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes, lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc. In certain embodiments, the cells used for gene therapy are autologous to the patient.

In one embodiment, an ERG or LTF nucleic acid or antisense molecule is introduced into the cells such that it is expressible by the cells or their progeny, and the recombinant cells are then administered in vivo for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained in vitro can potentially be used in accordance with this embodiment of the present invention. Such stem cells include, but are not limited to, hematopoietic stem cells (HSC), stem cells of epithelial tissues such as the skin and the lining of the gut, embryonic heart muscle cells, liver stem cells (PCT Pub. WO 94/08598), and neural stem cells (Stemple et al. (1992) *Cell*, 71:973-985).

Epithelial stem cells (ESCs) or keratinocytes can be obtained from tissues such as the skin and the lining of the gut by known procedures (Rheinwald (1980) *Meth. Cell Bio.*, 21A:229). In stratified epithelial tissue such as the skin, renewal occurs by mitosis of stem cells within the germinal layer, the layer closest to the basal lamina. Stem cells within the lining of the gut provide for a rapid renewal rate of this tissue. ESCs or keratinocytes obtained from the skin or lining of the gut of a patient or donor can be grown in tissue culture (Rheinwald (1980) *Meth. Cell Bio.*, 21A:229; Pitelkow et al. (1986) *Mayo Clinic. Proc.*, 61:771). If the ESCs are provided by a donor, a method for suppression of host versus graft reactivity (e.g., irradiation, drug or antibody administration to promote moderate immunosuppression) can also be used.

With respect to hematopoietic stem cells (HSC), any technique which provides for the isolation, propagation, and

maintenance in vitro of HSC can be used in this embodiment of the invention. Techniques by which this may be accomplished include (a) the isolation and establishment of HSC cultures from bone marrow cells isolated from the future host, or a donor, or (b) the use of previously established long-term HSC cultures, which may be allogeneic or xenogeneic. Non-autologous HSC may be used in conjunction with a method of suppressing transplantation immune reactions of the future host/patient. In a particular embodiment, human bone marrow cells can be obtained from the posterior iliac crest by needle aspiration (see, e.g., Kodo et al. (1984) *J. Clin. Invest.*, 73:1377-1384). In one embodiment, the HSCs can be made highly enriched or in substantially pure form. This enrichment can be accomplished before, during, or after long-term culturing, and can be done by any techniques known in the art. Long-term cultures of bone marrow cells can be established and maintained by using, for example, modified Dexter cell culture techniques (Dexter et al. (1977) *J. Cell Physiol.*, 91:335) or Witlock-Witte culture techniques (Witlock et al. (1982) *Proc. Natl. Acad. Sci. U.S.A.*, 79:3608-3612).

Pharmaceutical Compositions and Administration

The invention further provides pharmaceutical compositions comprising an effective amount of an ERG or LTF therapeutic, including ERG or LTF nucleic acids (sense or antisense) or ERG or LTF polypeptides of the invention, in a pharmaceutically acceptable carrier, as described below.

Compositions comprising an effective amount of a polypeptide of the present invention, in combination with other components such as a physiologically acceptable diluent, carrier, or excipient, are provided herein. The polypeptides can be formulated according to known methods used to prepare pharmaceutically useful compositions. They can be combined in admixture, either as the sole active material or with other known active materials suitable for a given indication, with pharmaceutically acceptable diluents (e.g., saline, Tris-HCl, acetate, and phosphate buffered solutions), preservatives (e.g., thimerosal, benzyl alcohol, parabens), emulsifiers, solubilizers, adjuvants and/or carriers. Suitable formulations for pharmaceutical compositions include those described in Remington's *Pharmaceutical Sciences*, 16th ed., Mack Publishing Company, Easton, Pa., 1980.

In addition, such compositions can be complexed with polyethylene glycol (PEG), metal ions, or incorporated into polymeric compounds such as polyacetic acid, polyglycolic acid, hydrogels, dextran, etc., or incorporated into liposomes, microemulsions, micelles, unilamellar or multilamellar vesicles, erythrocyte ghosts or spheroblasts. Such compositions will influence the physical state, solubility, stability, rate of in vivo release, and rate of in vivo clearance, and are thus chosen according to the intended application.

The compositions of the invention can be administered in any suitable manner, e.g., topically, parenterally, or by inhalation. The term "parenteral" includes injection, e.g., by subcutaneous, intravenous, or intramuscular routes, also including localized administration, e.g., at a site of disease or injury. Sustained release from implants is also contemplated. One skilled in the art will recognize that suitable dosages will vary, depending upon such factors as the nature of the disorder to be treated, the patient's body weight, age, and general condition, and the route of administration. Preliminary doses can be determined according to animal tests, and the scaling of dosages for human administration is performed according to art-accepted practices.

Compositions comprising nucleic acids of the invention in physiologically acceptable formulations, e.g., to be used for gene therapy are also contemplated. In one embodiment,

the nucleic acid can be administered in vivo to promote expression of the encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular as described in other sections herein.

Various delivery systems are known in the art and can be used to administer a therapeutic of the invention. Examples include, but are not limited to encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the therapeutic, receptor-mediated endocytosis (see, e.g., Wu et al. (1987) *J. Biol. Chem.*, 262:4429-4432), construction of a therapeutic nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to administer the pharmaceutical compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, a suppository, an implant, wherein the said implant is of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. In one embodiment, administration can be by direct injection at the site (or former site) of a malignant tumor or neoplastic or pre-neoplastic tissue.

In another embodiment, the therapeutic can be delivered in a vesicle, in particular a liposome (see Langer (1990) *Science*, 249:1527-1533; Treat et al. (1989) in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein et al., eds., Liss, New York, pp. 353-365; Lopez-Berestein, *ibid.*, pp. 317-327. In yet another embodiment, the therapeutic can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton (1987) *CRC Crit. Rev. Biomed. Eng.*, 14:201; Buchwald et al. (1980) *Surgery*, 88:507; Saudek et al. (1989) *New Engl. J. Med.*, 321:574). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer et al., eds., CRC Press, Boca Raton, Fla., 1974; *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen et al., eds., Wiley, N.Y., 1984; Ranger et al. (1983) *J. Macromol. Sci. Rev. Macromol. Chem.*, 23:61; see also Levy et al. (1985) *Science*, 228:190; During et al. (1989) *Ann. Neurol.*, 25:351; Howard et al. (1989) *J. Neurosurg.*, 71:105. In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson (1984) in *Medical Applications of Controlled*

Release, *supra*, vol. 2, pp. 115-138). Other controlled release systems are discussed in the review by Langer (1990, *Science*, 249:1527-1533).

Diagnosis and Screening

ERG, LTF, and/or AMACR proteins, analogues, derivatives, and fragments thereof, and antibodies thereto; ERG, LTF, DD3, and/or AMACR nucleic acids (and their complementary and homologous sequences) and antibodies thereto, including anti-ERG, anti-DD3, anti-LTF and/or anti-AMACR antibodies, have uses in diagnostics. Such molecules can be used in assays, such as immunoassays, to detect, prognose, diagnose, or monitor various conditions, diseases, and disorders affecting ERG, LTF, DD3, and/or AMACR expression, or monitor the treatment thereof, particularly cancer, and more particularly prostate cancer. In particular, such an immunoassay is carried out by a method comprising contacting a sample derived from an individual with an anti-ERG, anti-LTF, anti-DD3, and/or anti-AMACR antibody (directed against either a protein product or a nucleic acid) under conditions such that specific binding can occur, and detecting or measuring the amount of any specific binding by the antibody. In one embodiment, such binding of antibody, in tissue sections, can be used to detect aberrant ERG, LTF, DD3, and/or AMACR localization or aberrant (e.g., high, low or absent) levels of ERG, LTF, DD3, and/or AMACR. In a specific embodiment, antibody to ERG, LTF, DD3, and/or AMACR can be used to assay in a biological sample (e.g., tissue, blood, or urine sample) for the presence of ERG, LTF, DD3, and/or AMACR where an aberrant level of ERG, LTF, DD3, and/or AMACR is an indication of a diseased condition, such as cancer, including, for example, prostate cancer.

Any biological sample in which it is desired to detect an oligonucleotide or polypeptide of interest can be used, including tissue, cells, blood, lymph, semen, and urine. The biological sample is preferably derived from prostate tissue, blood, or urine. The tissue sample comprises cells obtained from a patient. The cells may be found in a prostate tissue sample collected, for example, by a prostate tissue biopsy or histology section, or a bone marrow biopsy. The blood sample can include whole blood, plasma, serum, or any derivative thereof, including, for example, circulating cells, such as prostate cells, isolated from the blood sample, or nucleic acid or protein obtained from the isolated cells. Blood may contain prostate cells, particularly when the prostate cells are cancerous, and, more particularly, when the prostate cancer metastasizes and is shed into the blood. Similarly, the urine sample can be whole urine or any derivative thereof, including, for example, cells, such as prostate cells, obtained from the urine.

The immunoassays which can be used include, but are not limited to, competitive and non-competitive assay systems using techniques such as Western blots, radioimmunoassays, ELISA, immunoprecipitation assays, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few.

ERG, LTF, DD3, and/or AMACR genes and related nucleic acid sequences and subsequences, including complementary sequences, can also be used in hybridization assays. ERG, LTF, DD3, and/or AMACR nucleic acid sequences, or subsequences thereof comprising about at least 8, 15, 20, 50, 100, 250, or 500 nucleotides can be used as hybridization probes. Hybridization assays can be used to detect, prognose, diagnose, or monitor conditions, disorders, or disease states associated with aberrant changes in ERG, LTF, DD3, and/or AMACR expression and/or activity as described

above. In particular, such a hybridization assay is carried out by a method comprising contacting a sample containing nucleic acid with a nucleic acid probe capable of hybridizing under defined conditions (preferably under high stringency hybridization conditions, e.g., hybridization for 48 hours at 65° C. in 6×SSC followed by a wash in 0.1×SSC at 50° C. for 45 minutes) to an ERG, LTF, DD3, and/or AMACR nucleic acid, and detecting (i.e., measuring either qualitatively or quantitatively) the degree of the resulting hybridization. As described herein, any nucleic acid amplification procedure, including, PCR/RT-PCR, TMA, NASBA, 3SR, LCR, SDA, and LAMP can be used to detect the presence of the ERG, LTF, DD3 and/or AMACR gene and/or the level of its mRNA expression.

In some applications, probes exhibiting at least some degree of self-complementarity are desirable to facilitate detection of probe:target duplexes in a test sample without first requiring the removal of unhybridized probe prior to detection. Molecular torch probes are a type of self-complementary probes that are disclosed by Becker et al., U.S. Pat. No. 6,361,945. The molecular torch probes disclosed by Becker et al. have distinct regions of self-complementarity, referred to as "the target binding domain" and "the target closing domain," which are connected by a joining region and which hybridize to one another under predetermined hybridization assay conditions. When exposed to denaturing conditions, the complementary regions (which may be fully or partially complementary) of the molecular torch probe melt, leaving the target binding domain available for hybridization to a target sequence when the predetermined hybridization assay conditions are restored. And when exposed to strand displacement conditions, a portion of the target sequence binds to the target binding domain and displaces the target closing domain from the target binding domain. Molecular torch probes are designed so that the target binding domain favors hybridization to the target sequence over the target closing domain. The target binding domain and the target closing domain of a molecular torch probe include interacting labels (e.g., luminescent/quencher) positioned so that a different signal is produced when the molecular torch probe is self-hybridized as opposed to when the molecular torch probe is hybridized to a target nucleic acid, thereby permitting detection of probe:target duplexes in a test sample in the presence of unhybridized probe having a viable label or labels associated therewith.

Another example of detection probes having self-complementarity are the molecular beacon probes disclosed by Tyagi et al. in U.S. Pat. No. 5,925,517. Molecular beacon probes include nucleic acid molecules having a target complement sequence, an affinity pair (or nucleic acid arms) holding the probe in a closed conformation in the absence of a target nucleic acid sequence, and a label pair that interacts when the probe is in a closed conformation. Hybridization of the target nucleic acid and the target complement sequence separates the members of the affinity pair, thereby shifting the probe to an open conformation. The shift to the open conformation is detectable due to reduced interaction of the label pair, which may be, for example, a fluorophore and quencher, such as DABCYL and EDANS.

By way of example, ERG, LTF, AMACR, or DD3 hybridization probes can comprise a nucleic acid having a contiguous stretch of at least about 8, 15, 20, 50, 100, 250, 500, 750, 1000, 1250, or 1500 contiguous nucleotides of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, or SEQ ID NO:5 or a sequence complementary thereto. Such contiguous fragments of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, or SEQ ID NO:5 may also contain at

least one mutation so long as the mutant sequence retains the capacity to hybridize to SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO: 3, SEQ ID NO:4, or SEQ ID NO:5 under low or high stringency conditions (preferably under high stringency hybridization conditions, e.g., hybridization for 48 hours at 65° C. in 6×SSC followed by a wash in 0.1×SSX at 50° C. for 45 minutes).

In specific embodiments, diseases and disorders involving hyperproliferation of cells, such as cancers, including, for example, prostate cancer, can be diagnosed, or their suspected presence can be screened for, or a predisposition to develop such disorders can be predicted, by detecting levels of the ERG, LTF, and/or AMACR protein, ERG, DD3, and/or AMACR RNA, or ERG, DD3, and/or AMACR functional activity, or by detecting mutations in ERG, DD3, LTF and/or AMACR RNA, DNA, or protein (e.g., translocations in ERG, LTF, DD3, or AMACR nucleic acids, truncations in the ERG, LTF, DD3, or AMACR gene or protein, changes in nucleotide or amino acid sequence relative to wild-type ERG, LTF, DD3, or AMACR) that cause increased or decreased expression or activity of ERG, LTF, DD3, and/or AMACR. By way of example, levels of ERG, LTF, and/or AMACR protein can be detected by immunoassay, levels of ERG, LTF, DD3, and/or AMACR mRNA can be detected by hybridization assays (e.g., Northern blots, dot blots, or any nucleic acid amplification procedure, including, PCR/RT-PCR, TMA, NASBA, 3SR, LCR, SDA, and LAMP), translocations and point mutations in ERG, LTF, DD3, and/or AMACR nucleic acids can be detected by Southern blotting, RFLP analysis, any nucleic acid amplification procedure, including, PCR/RT-PCR, TMA, NASBA, 3SR, LCR, SDA, LAMP, sequencing of the ERG, LTF, DD3, and/or AMACR genomic DNA or cDNA obtained from the patient, etc.

In one embodiment, levels of the ERG, DD3, LTF and/or AMACR mRNA or protein in a subject sample are detected or measured and compared to the mRNA or protein expression levels of the corresponding gene in a control sample or to a standard numerical value or range. For example, increased expression levels of ERG, DD3, and/or AMACR or decreased levels of LTF, relative to a matched, normal tissue sample, indicate that the subject has a malignancy or hyperproliferative disorder, including, for example, prostate cancer, or a predisposition to develop the same. Other appropriate controls include other noncancerous samples from the subject, samples obtained from a different subject without cancer, or other cancer-specific markers. For example, in prostate cancer, a prostate-cell specific marker, such as PSA, can be used as a control to compare and/or normalize expression levels of other genes, such as ERG, LTF, DD3, and/or AMACR. In one embodiment, a method of diagnosing cancer, such as prostate cancer, comprises obtaining a biological sample from a subject (e.g., a tissue sample (e.g., from biopsy tissue), a blood sample, or a urine sample), determining the expression level of a ERG, LTF, DD3, and/or AMACR gene and/or ERG, LTF, DD3, and/or AMACR activity in the samples, and diagnosing or prognosing cancer in said subject. In further embodiments, the expression level of the ERG, LTF, DD3, and/or AMACR gene and/or ERG, LTF, DD3, and/or AMACR activity is determined by Southern blotting, Northern blotting, Western blotting, ELISA, any nucleic acid amplification procedure, including, PCR/RT-PCR, TMA, NASBA, 3SR, LCR, SDA, and LAMP, or other techniques as described herein or known in the art. Without limiting the instant invention, increased or decreased expression of at least two times, as

compared to the control sample indicates the presence of prostate cancer or a higher predisposition to developing prostate cancer.

Another aspect of the invention provides a means for monitoring a response to “hormonal therapy” by evaluating the expression profiles of the ERG gene, alone or in combination with the AMACR and/or DD3 genes and/or LTF genes, and correlating these profiles with the clinical signs of the disease.

Kits for diagnostic use are also provided. A kit comprises an anti-ERG gene antibody or an antibody directed against the ERG protein and/or an anti-AMACR gene antibody or an antibody directed against the AMACR protein and/or an anti-DD3 gene antibody and/or an anti-LTF gene antibody or an antibody directed against the LTF protein, which can be optionally detectably labeled. A kit is also provided that comprises a nucleic acid probe capable of hybridizing under defined conditions (preferably under high stringency hybridization conditions, e.g., hybridization for 48 hours at 65° C. in 6×SSC followed by a wash in 0.1×SSX at 50° C. for 45 minutes) to ERG, LTF, DD3, and/or AMACR nucleic acid. In a specific embodiment, a kit comprises at least a pair of primers (e.g., each in the size range of at least about 6, 17, 30, or 60 nucleotides) that are capable of priming amplification, by any nucleic acid amplification procedure (including e.g., PCR/RT-PCR, TMA, NASBA, 3SR, LCR, SDA, LAMP), of the ERG, LTF, DD3, and/or AMACR gene or a fragment thereof. A kit can comprise a predetermined amount of a purified ERG, LTF, DD3, and/or AMACR protein or nucleic acid for use, e.g., as a standard or control. The kit can also comprise one or more components for detecting the nucleic acid probe, including components described herein or known in the art.

In one embodiment, the kit comprises a nucleic acid that hybridizes under defined conditions (and preferably under conditions of high stringency, e.g., hybridization for 48 hours at 65° C. in 6×SSC followed by a wash in 0.1×SSX at 50° C. for 45 minutes) with at least one gene chosen from those genes identified in Tables 1-6 or the DD3 gene, and is affixed to a support, alone, or in combination with other nucleic acids. For example, an ERG and/or LTF nucleic acid can be affixed to the support, with or without other nucleic acids. In a specific embodiment, the support comprises at least an ERG nucleic acid and an AMACR nucleic acid or at least an ERG nucleic acid and a DD3 nucleic acid. In another embodiment, the support comprises at least an ERG nucleic acid, an AMACR nucleic acid, and a DD3 nucleic acid. This support can be used as part of a kit for detecting cancer, such as prostate cancer. These kits can further comprise at least a pair of primers (e.g., each in the size range of at least about 6, 17, 30, or 60 nucleotides) that are capable of priming amplification, by any nucleic acid amplification procedure (including e.g., PCR/RT-PCR, TMA, NASBA, 3SR, LCR, SDA, LAMP), of the ERG, LTF, DD3, and/or AMACR gene or a fragment thereof.

EXAMPLES

Example 1

Screening of CaP Cell-Specific Gene Expression Signatures Using Affymetrix GeneChip

Patient Selection

Specimens were obtained under an IRB-approved protocol from patients treated by radical prostatectomy (RP) at Walter Reed Army Medical Center (WRAMC). From over

300 patients two groups were selected which had prostate tumors with either moderate (MR) or high risk (HR) of disease progression after RP. The HR group had PSA recurrence, Gleason score 8-9, T3c stage, seminal vesicle invasion, and poorly differentiated tumor cells; the MR group had no PSA recurrence, Gleason score 6-7, T2a-T3b stage, no seminal vesicle invasion, and well to moderately differentiated tumor cells. LCM compatible specimens were selected from age and race matched HR or MR patients with no family history of CaP.

Tissue Samples and Laser-Capture Microdissection

Normal and cancer cells were laser capture microdissected (LCM) from OCT embedded and Hematoxylin-eosin (H&E) stained frozen prostate sections of radical prostatectomy specimens (2000 laser shots for one sample). Laser capture microdissection (LCM) facilitates the isolation of morphologically defined, homogenous cell populations from complex tissues by selectively adhering the cells of interest to a transparent film with focused pulses of low energy infrared laser under a microscope. Emmert-Buck et al., Science (1996); 274(5289): 921-922; Schutz et al., Nat Biotechnol (1998) 16(8): 737-742.

RNA Extraction and T7-Based Linear RNA Amplification

Total RNA was isolated from the LCM samples with the MicroRNA kit (Stratagene, La Jolla, Calif.), quantified using RiboGreen dye (Molecular Probes, Eugene, Oreg.) and VersaFluor fluorimeter (BioRad, Hercules, Calif.), and quality tested by RT-PCR using NKX3.1 and GAPDH primers. Linear RNA amplification was performed using RiboAmp RNA amplification kit (Arcturus, Mountain View, Calif.). Precisely, 2 nanograms of total RNA from LCM derived epithelial cells of normal as well as tumor tissue from each patient was used for the first round of amplification. During the second round of amplification after cDNA synthesis and purification the samples were biotinylated during in vitro transcription which was used for the GeneChip analysis.

Gene Chip Analysis

Linearly amplified aRNA was hybridized to high-density oligonucleotide human genome array (HG U133A array) (Affymetrix, Santa Clara, Calif., USA). The array contains 22,283 probe sets, about 18,000 of which represent well annotated genes, while the remainder represent various expressed sequence tags (EST) and hypothetical genes. Biotinylation was carried out using aRNA by in vitro transcription using MEGA script T7 in vitro Transcription Kit (Ambion, Austin, Tex., USA) cDNA and biotinylated UTP and biotinylated CTP (ENZO, Farmingdale, N.Y., USA) (34). The biotin labeled cRNA was purified using the QIAGEN RNeasy spin columns (QIAGEN, Valencia, Calif.) following the manufacturer's protocol. The biotin labeled cRNA was fragmented in a 40 μ l reaction mixture containing 40 mM Tris-acetate, pH 8.1, 100 mM potassium acetate, and 30 mM magnesium acetate incubated at 94° C. for 35 minutes and then put on ice.

Hybridization, Staining and Scanning of the GeneChip

The biotin labeled and fragmented aRNA was hybridized to the HG U133A array. Briefly, a 220 μ l hybridization solution consisting of: 1M NaCl, 10 mM Tris pH 7.6, 0.005% Triton X-100, 50 pM control Oligo B2 (5' bioGT-CAAGATGCTACCGTTCAG 3') (SEQ ID NO:6) (Affymetrix); the control cRNA cocktail of: Bio B (150 pM), Bio C (500 pM), Bio D (2.5 nM) and Cre X (10 nM) (American Type Tissue Collection, Manassas, Va. and Lofstrand Labs, Gaithersburg, Md.), 0.1 mg/ml herring sperm DNA and 0.05 μ g/ μ l of the fragmented labeled sample cRNA was heated to 95° C. for 35 min., cooled to 40° C. and clarified by centrifugation. Hybridization was at 42° C. in a rotisserie

oven (Model 320, Affymetrix) at 60 rpm for 16 hours. Following hybridization, the GeneChip arrays were washed 10 times at 25° C. with 6 \times SSPE-T buffer (1 M NaCl, 0.006 M EDTA, and 0.06 M Na₃PO₄, 0.005% Triton X-100, pH 7.6) using the automated fluidics station protocol. GeneChip arrays were incubated at 50° C. in 0.5 \times SSPE-T, 0.005% Triton X-100 for 20 minutes at 60 rpm in the rotisserie oven. GeneChip arrays were stained for 15 minutes at room temperature and at 60 rpm, with streptavidin phycoerythrin (Molecular Probes, Inc., Eugene, Oreg.) stain solution at a final concentration of 10 μ g/ml in 6 \times SSPE-T buffer and 1.0 mg/ml acetylated bovine serum albumin (Sigma). GeneChip arrays were washed twice at room temperature with 6 \times SSPE-T buffer, and then were scanned with the HP GeneArray Scanner (Hewlett-Packard, Santa Clara, Calif.) controlled by GeneChip 3.1 Software (Affymetrix).

Example 2

Analysis of GeneChip Results by Supervised Multi-Dimensional Scaling (MDS)

Image Analysis and Data Collection

Affymetrix GeneChip Microarray Analysis Software, version 3.1 and Affymetrix Micro DB and Data Mining Tool version 2.0 (Affymetrix), Microsoft Excel 2000 (Microsoft, Seattle, Wash.) and Statistica version 4.1 (Stat Soft, Inc., Tulsa, Okla.) were used. In the Affymetrix system, the average difference fluorescence is the average of the difference between every perfect match probe cell and its control mismatch probe cell and is directly related to the level of expression of a transcript. A comparative file indicates the relative change in abundance (fold change) for each transcript between a baseline and an experimental sample. For further detail and advanced bioinformatic analysis we used the Microarray Data Analysis software from NHGRI and the GeneSpring software (Silicon Genetics, Calif.).

Data Analysis

For clustering analysis, National Human Genome Research Institute (NHGRI) Microarray Data Analysis software was used, which partitioned the samples of the high risk and moderate risk groups into well-separated and homogeneous groups based on the statistical behavior of their genes expression. To achieve the objective of clustering each of the groups, all pair-wise similarities between samples were evaluated, and then grouped via the average linkage algorithm. Pearson correlation coefficient or Euclidean distance were typically used to quantify the similarity. Unsupervised hierarchical and or non hierarchical clustering was also performed using the same distance matrix.

Using a matrix of Euclidean distance measurements from complete pair wise comparison of all the prostate specimens, a multidimensional scaling (MDS) method was performed using an implementation of MDS in the MATLAB package to determine the overall similarities and dissimilarities in gene expression profiles. A weighted gene analysis was performed to generate a subset of genes statistically significant in separating the high risk group from the moderate risk group.

Briefly, for two different groups e.g., epithelium of high risk tumor and epithelium of moderate risk tumor with a given number of samples 25 and 25, the discriminative weight for each gene is determined by the formula: $w = d_B / (k_1 d_{w1} + k_2 d_{w2} + \alpha)$; where d_B is the Euclidean distance between two groups (center-to-center or between cluster Euclidean distance), d_{w1} is the average Euclidean distance

among all the epithelial samples of high risk group, d_{w2} is average Euclidean distance among all the epithelial samples of moderate risk group, $k_1=25/(25+25)$, $k_2=25/(25+25)$, and α is a small constant to ensure the denominator is never equal to zero. Genes were ranked according to their w values. Genes with high w values created greater separation between groups and denser compaction within the group. In other words, the subset of genes with high w values have the most discriminative power to differentiate a high risk group from a moderate risk group and vice versa. Sample labels were randomly permuted and the w value was computed again for each gene to test the statistical significance of the discriminative weights. Genes with the most significant expression differences were selected by p -values. A hierarchical clustering algorithm was used to verify the predictor model obtained from the supervised MDS analysis.

From this analysis, specific genes were identified whose expression signature in tumor tissue varied from their expression signature in benign matched tissue. Genes with a p -value of not more than 0.05 were selected and ranked by p -value, as shown in Tables 1-6.

In Silico Validation:

We have tested the discriminatory potential of the genes that we obtained from our analysis on some independent data sets. Affymetrix oligonucleotide GeneChip Hum95Av2 data were obtained from Welsh et al. 2001, Singh et al. Genes from these data bases that correspond with the genes of our discriminatory list were selected and their tumor specific expression intensities and/or tumor over normal ratio were used for an MDS analysis as described above in the data analysis section. MDS plots were obtained depicting the discriminatory capability of the genes on the independent data sets.

TABLE 1

The first 50 genes obtained from the supervised MDS analysis of tumor versus benign tissues of all the high risk and moderate risk CaP patients, ranked by p -value. (T vs B in All 18 Samples)							
GenBank					Expression Regulation		
No.	Accession	Common Name of Genes	Description of Genes	Map	p-Value	Tumor	Benign
1.	AF047020	AMACR	Alpha-methylacyl-CoA racemase	5p13.2-q11.1	0	Up	Down
2.	NM_002343	LTF	Lactotransferrin	3q21-q23	0	Down	Up
3.	NM_002275	KRT15	Keratin 15	17q21	0.000001	Down	Up
4.	BC000915	CLIM1, CLP36, CLP-36	PDZ and LIM domain 1 (elfin)	10q22-q26.3	0.000001	Down	Up
5.	X90579	CYP3A5	Cytochrome P450, subfamily 3A, polypeptide 5	7	0.000001	Down	Up
6.	NM_003671	CDC14B1, CDC14B2	<i>H. sapiens</i> CDC14 cell division cycle 14 homolog B	9q22.2-q22.31	0.000005	Down	Up
7.	AI424243	CEGP1	<i>H. sapiens</i> cDNA clone IMAGE: 2094442	11	0.000005	Down	Up
8.	NM_022370	Rbig1	Hypothetical protein FLJ21044 similar to Rbig1	11q24.2	0.000009	Down	Up
9.	AI356398	ZFP36L2	TISD_HUMAN P47974 TIS11D PROTEIN	2	0.000018	Down	Up
10.	NM_005213	STF1, STFA	Cystatin A (stefin A)	3q21	0.000018	Down	Up
11.	NM_006394	RIG	Regulated in glioma	11p15.1	0.000018	Down	Up
12.	AF275945	EVA1	Epithelial V-like antigen 1	11q23.3	0.000018	Down	Up
13.	NM_020186	DC11	DC11 protein	7q21.3	0.000018	Up	Down
14.	AI922538	TMEM1	Transmembrane protein 1	21	0.000018	Down	Up
15.	NM_014863	BRAG, KIAA0598	B cell RAG associated protein	10q26	0.000018	Down	Up
16.	AI669229	RARRES1	<i>Homo sapiens</i> cDNA clone IMAGE: 2315074	3q25.33	0.000036	Down	Up
17.	NM_006017	AC133, CD133	Prominin (mouse)-like 1	4p15.33	0.000036	Down	Up
18.	NM_004503	HOXC6	Homeo box C6	12q12-q13	0.000036	Up	Down
19.	NM_005084	PAFAH, LDL-PLA2	Phospholipase A2, group VII	6p21.2-p12	0.000036	Up	Down
20.	NM_001511	MGSA, CXCL1, SCYB1	GRO1 oncogene	4q21	0.000071	Down	Up
21.	BG054844	ARHE	<i>H. sapiens</i> cDNA clone IMAGE: 3441573	2q23.3	0.000071	Down	Up
22.	NM_007191	WIF-1	Wnt inhibitory factor-1	12q14.2	0.000071	Down	Up
23.	X99268	TWIST	Twist (<i>Drosophila</i>) homolog	7p21.2	0.000071	Up	Down
24.	AI826799	EFEMP1	EXTRACELLULAR PROTEIN S1-5 PRECURSOR	2p16	0.000071	Down	Up
25.	NM_001018	RPS15	Ribosomal protein S15	19p13.3	0.000071	Up	Down
26.	AV711904	LYZ	Lysozyme (renal amyloidosis)		0.000071	Down	Up
27.	AI433463	MME	NEPRILYSIN (HUMAN)	3q25.1-q25.2	0.000071	Down	Up
28.	BE908217	ANXA2	<i>H. sapiens</i> cDNA clone IMAGE: 3902323	15q21-q22	0.000071	Down	Up
29.	NM_000441	PDS, DFNB4	Solute carrier family 26, member 4	7q31	0.000071	Down	Up
30.	BC003068	SLC19A1	Solute carrier family 19, member 1	21q22.3	0.000071	Up	Down
31.	NM_005950	MT1	Metallothionein 1G	16q13	0.000071	Down	Up
32.	NM_013281	FLRT3	Fibronectin leucine rich transmembrane protein 3	20p11	0.000071	Down	Up
33.	AI351043	ESTs	<i>H. sapiens</i> cDNA clone IMAGE: 1948310	21	0.000145	Up	Down
34.	NM_001099	PAP	Acid phosphatase, prostate	3q21-q23	0.000145	Down	Up
35.	NM_006113	VAV3	Vav 3 oncogene	1p13.1	0.000145	Down	Up
36.	NM_005980	S100P	S100 calcium-binding protein P	4p16	0.000145	Down	Up
37.	NM_000165	GJA1	Gap junction protein, alpha 1, 43 kD (connexin 43)	6q21-q23.2	0.000145	Down	Up
38.	NM_003897	DIF2, IEX1, PRG1	Immediate early response 3	6p21.3	0.000145	Down	Up

TABLE 1-continued

The first 50 genes obtained from the supervised MDS analysis of tumor versus benign tissues of all the high risk and moderate risk CaP patients, ranked by p-value. (T vs B in All 18 Samples)						
GenBank					Expression Regulation	
No.	Accession	Common Name of Genes	Description of Genes	Map	p-Value	Tumor Benign
39.	BC001388	ANX2, LIP2, CALIH	Annexin A2	15q21-q22	0.000145	Down Up
40.	BC003070	HDR, MGC5445	GATA-binding protein 3	10p15	0.000145	Down Up
41.	NM_020139	LOC56898	Oxidoreductase UCPA	4	0.000145	Down Up
42.	AK002207	KIAA0610	KIAA0610 protein	13	0.000145	Down Up
43.	NM_000574	CR, TC, CD55	Decay accelerating factor for complement	1q32	0.000145	Down Up
44.	NM_006926	SP-A2, COLEC5	Surfactant, pulmonary-associated protein A2	10q22-q23	0.000145	Up Down
45.	U37546	API2, MIHC, CIAP2	Baculoviral IAP repeat-containing 3	11q22	0.000145	Down Up
46.	AU148057	DKK3	<i>H. sapiens</i> cDNA clone MAMMA1002489	11pter-p15.5	0.000145	Down Up
47.	NM_002600	DPDE4, PDEIVB	Phosphodiesterase 4B, cAMP-specific	1p31	0.000145	Down Up
48.	S59049	BL34, IER1, IR20	Regulator of G-protein signalling 1	1q31	0.0003	Down Up
49.	NM_001275	CGA, CgA	Chromogranin A (parathyroid secretory protein 1)	14q32	0.0003	Down Up
50.	AL575509	ETS2	<i>H. sapiens</i> cDNA clone CS0DI059YP21 = 21q22.2		0.0003	Down Up

TABLE 2

The first 50 genes from the supervised MDS analysis of tumor over benign (T/B) tissues ratio (Fold Change) of the high risk versus moderate risk CaP patients, ranked by p-value.: (T/B Fold Change in HR vs MR)					
No	Genbank accession	Common Name of Genes	Description of Genes	Map	p-Value
1.	NM_004522	KINN, NKHC, NKHC2, NKHC-2	Kinesin family member 5C	2q23.3	0.00011
2.	J03198	GNAI3	Guanine nucleotide binding protein G (K), alpha subunit	1p13	0.000981
3.	NM_018010	HIPPI, FLJ10147	Hypothetical protein FLJ10147	3q13.13	0.003257
4.	NM_005479	FRAT1	Frequently rearranged in advanced T-cell lymphomas	10q23.33	0.004964
5.	NM_021795	SAP1	ELK4, ETS-domain protein (SRF accessory protein 1)	1q32	0.004964
6.	NM_003113	LEU5, RFP2	Nuclear antigen Sp100	2q37.1	0.004964
7.	NM_002053	GBP1	Guanylate binding protein 1, interferon-inducible, 67 kD	1p22.1	0.004964
8.	AF064092	GSA, GSP, GPSA, GNAS1,	Guanine nucleotide regulatory protein	20q13.2-q13.3	0.007579
9.	BC003070	HDR, MGC5445	GATA-binding protein 3	10p15	0.007579
10.	NM_012245	SKIP, NCOA-62	SKI-interacting protein	14q24.3	0.007579
11.	NM_015895	LOC51053	Geminin	6p22.2	0.007579
12.	AA083478	TRIM22	Stimulated trans-acting factor (50 kDa)	11	0.007579
13.	NM_000100	PME, CST6, EPM1, STFB	Cystatin B (stefin B)	21q22.3	0.007579
14.	NM_003031	SIAH1	Seven in absentia (<i>Drosophila</i>) homolog 1	16q12	0.007579
15.	NM_003407	TFP, GOS24, TIS11, NUP475	Zinc finger protein 36, C3H type, homolog (mouse)	19q13.1	0.007579
16.	BF979419	ESTs	ESTs, Highly similar to 60S ribosomal protein 13A [<i>H. sapiens</i>]	19q13.33	0.007579
17.	NM_021038	MBNL	Muscleblind (<i>Drosophila</i>)-like	3q25	0.007579
18.	NM_014454	PA26	P53 regulated PA26 nuclear protein	6q21	0.007579
19.	BC004399	DEME-6	DEME-6 protein	1p32.3	0.007579
20.	NM_018490	LGR4	G protein-coupled receptor 48	11p14-p13	0.007579
21.	NM_004328	BCS, BCS1, h-BCS, Hs.6719	BCS1 (yeast homolog)-like	2q33	0.007579
22.	D87445	KIAA0256	KIAA0256 gene product	15	0.007579
23.	NM_006326	NIFIE14	<i>Homo sapiens</i> seven transmembrane domain protein, mRNA	19q13.12	0.007579
24.	D83077	TTC3	Tetratricopeptide repeat domain 3	Xq13.1	0.007579
25.	NM_006732	GOS3	FBJ murine osteosarcoma viral oncogene homolog B	19q13.32	0.007579
26.	NM_003760	EIF4G3	Eukaryotic translation initiation factor 4 gamma, 3	1pter-p36.13	0.007579
27.	NM_004905	AOP2	Anti-oxidant protein 2	1q24.1	0.01159
28.	NM_018439	IMPACT	Hypothetical protein IMPACT	18	0.01159
29.	BC000629	DARS	Aspartyl-tRNA synthetase	2q21.2	0.01159
30.	AK002064	DKFZP564A2416	DKFZP564A2416 protein	2	0.01159
31.	NM_013387	HSPC051	Ubiquinol-cytochrome c reductase complex (7.2 kD)	22	0.01159
32.	AA135522	KIAA0089	<i>Homo sapiens</i> KIAA0089 mRNA sequence.	3	0.01159
33.	NM_015545	KIAA0632	KIAA0632 protein	7q22.1	0.01159
34.	NM_005767	P2Y5	Purinergic receptor (family A group 5)	13q14	0.01159
35.	BC003682	G25K, CDC42Hs	Cell division cycle 42 (GTP-binding protein, 25 kD)	1p36.1	0.01159
36.	NM_005053	RAD23A	RAD23 (<i>S. cerevisiae</i>) homolog A	19p13.2	0.017805
37.	A1672541	IPW	Human non-translated mRNA sequence.	15q11-q12	0.017805
38.	AK023938	<i>H. sapiens</i> cDNA FLJ13876 clone	SELECTED MODEL ORGANISM PROTEIN SIMILARITIES	2q37.3	0.017805
39.	NM_000062	C1IN, C1NH, C1-INH	Serine (or cysteine) proteinase inhibitor, clade G (C1 inhibitor)	11q12-q13.1	0.017805
40.	AA576961	PHLDA1	Pleckstrin homology-like domain, family A, member 1	12q15	0.017805

TABLE 2-continued

The first 50 genes from the supervised MDS analysis of tumor over benign (T/B) tissues ratio (Fold Change) of the high risk versus moderate risk CaP patients, ranked by p-value.: (T/B Fold Change in HR vs MR)					
No	Genbank accession	Common Name of Genes	Description of Genes	Map	p-Value
41.	AI796269	NBS1, ATF, NIBRIN	<i>H. sapiens</i> cDNA similar to Cell Cycle Regulatory Protein P95.	8q21-q24	0.017805
42.	NM_000016	ACADM	Acyl-Coenzyme A dehydrogenase, C-4 to C-12 straight chain	1p31	0.017805
43.	AI867102	KIAA0906, NUP210, gp210	Nuclear pore membrane glycoprotein 210	3p25.2-p25.1	0.017805
44.	AI263909	ARHB, RHOB, RHOH6	Oncogene RHO6; Aplysia RAS-related homolog 6	2pter-p12	0.017805
45.	NM_016021	NCUBE1	Non-canonical ubiquitin conjugating enzyme 1	6	0.017805
46.	NM_012192	TIM9B, TIM10B	Fracture callus 1 (rat) homolog	11p15.5-p15.3	0.017805
47.	NM_025087	FLJ21511	Hypothetical protein FLJ21511	4	0.017805
48.	NM_014959	CARD8, CARDINAL, KIAA0955	Tumor up-regulated CARD-containing antagonist of caspase 9	19q13.33	0.017805
49.	AA923354	MAOA	Monoamine oxidase A.	Xp11.4-p11.3	0.017805
50.	NM_021964	ZNF148	Zinc finger protein 148 (pHZ-52)	3q21	0.017805
51.	NM_001674	ATF3	Activating transcription factor 3	1q32.3	0.017805

TABLE 3

The first 50 genes obtained from the supervised MDS analysis of tumor versus benign tissues of all the high risk CaP patients, ranked by p-value. (T vs N Intensities of 9 HR)							
No.	GenBank Accession	Common Name of Genes	Description of Genes	Map	p-Value	Expression Regulation	
1.	U65585	HLA-DR1B	Major histocompatibility complex, class II, DR beta 1	6p21.3	0.00002	Down	Up
2.	NM_002053	GBP1	Guanylate binding protein 1, interferon-inducible,	1p22.1	0.000076	Down	Up
3.	NM_021983	HLA-DRB4	Major histocompatibility complex, class II, DR beta 4	6	0.000076	Down	Up
4.	AI424243	CEGP1	<i>Homo sapiens</i> cDNA clone IMAGE: 2094442	11	0.000102	Down	Up
5.	NM_002343	LTF	Lactotransferrin	3q21-q23	0.000138	Down	Up
6.	NM_014575	SCHIP-1	Schwannomin-interacting protein 1	3q26.1	0.000257	Down	Up
7.	BC001169	ESD	Esterase D/fornylglutathione hydrolase	13q14.1-q14.2	0.000357	Up	Down
8.	BF970427	UGCG	UDP-glucose ceramide glucosyltransferase	9	0.000357	Down	Up
9.	NM_002275	KRT15	Keratin 15	17q21	0.000495	Down	Up
10.	AU148057	DKK3	<i>H. sapiens</i> cDNA clone MAMMA1002489	11pter-p15.5	0.000495	Down	Up
11.	AI922538	TMEM1	Transmembrane protein 1	21	0.000689	Down	Up
12.	NM_004481	GALNAC-T2	UDP-GalNAc transferase 2	1q41-q42	0.000689	Down	Up
13.	BC003070	HDR, MGC5445	GATA-binding protein 3	10p15	0.00097	Down	Up
14.	BF979419	ESTs, similar to RPL13A	<i>H. sapiens</i> 60S Ribosomal protein L13A		0.00097	Up	Down
15.	BG054844	ARHE	<i>H. sapiens</i> cDNA clone IMAGE: 3441573	2q23.3	0.00097	Down	Up
16.	L42024	HLA-B	Major histocompatibility complex, class I, B	6p21.3	0.00138	Down	Up
17.	AL545982	CCT2	<i>H. sapiens</i> cDNA clone CS0DI023YD15	12q15	0.001992	Up	Down
18.	NM_001993	TF, TFA, CD142	Coagulation factor III (thromboplastin, tissue factor)	1p22-p21	0.001992	Up	Down
19.	NM_004198	CHRNA6	Cholinergic receptor, nicotinic, alpha polypeptide 6	8p11.1	0.001992	Down	Up
20.	AV711904	LYZ	Lysozyme (renal amyloidosis)	12q15	0.001992	Down	Up
21.	NM_013387	HSPC051	Ubiquinol-cytochrome c reductase complex (7.2 kD)	22	0.001992	Up	Down
22.	AW514210	HLA-F	HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, F A	6p21.3	0.001992	Down	Up
23.	NM_005032	PLS3	Plastin 3 (T isoform)	Xq24	0.002894	Down	Up
24.	NM_003407	TTP, GOS24, NUP475	Zinc finger protein 36, C3H type, homolog (mouse)	19q13.1	0.002894	Down	Up
25.	NM_000165	GJA1	Gap junction protein, alpha 1, 43 kD (connexin 43)	6q21-q23.2	0.002894	Down	Up
26.	AF275945	EVA1	Epithelial V-like antigen 1	11q23.3	0.002894	Down	Up
27.	NM_002450	MT1	Metallothionein 1L	16q13	0.002894	Down	Up
28.	NM_005950	MT1	Metallothionein 1G	16q13	0.002894	Down	Up
29.	NM_006994	BTN3A3	Butyrophilin, subfamily 3, member A3	6p21.33	0.002894	Down	Up
30.	AI049962	KIAA0191	<i>H. sapiens</i> cDNA clone IMAGE: 1700970	1	0.002894	Down	Up
31.	X99268	TWIST	Twist (<i>Drosophila</i>) homolog	7p21.2	0.002894	Up	Down
32.	NM_016021	NCUBE1	Non-canonical ubiquitin conjugating enzyme 1	6	0.002894	Up	Down
33.	NM_016205	SCDF	Platelet derived growth factor C	4q32	0.002894	Up	Down
34.	AI681120	RANBP2	<i>H. sapiens</i> cDNA clone IMAGE: 2272403	2q11-q13	0.004205	Up	Down
35.	NM_000574	CR, TC, CD55	Decay accelerating factor for complement	1q32	0.004205	Down	Up
36.	NM_014937	KIAA0966	Sac domain-containing inositol phosphatase 2	10q26.13	0.004205	Down	Up
37.	NM_005213	STF1, STFA	Cystatin A (stefin A)	3q21	0.004205	Down	Up
38.	NM_005952	MT1	Metallothionein 1X	16q13	0.004205	Down	Up
39.	AF130095	FN1	Fibronectin 1	2q34	0.004205	Down	Up

TABLE 3-continued

The first 50 genes obtained from the supervised MDS analysis of tumor versus benign tissues of all the high risk CaP patients, ranked by p-value. (T vs N Intensities of 9 HR)							
GenBank		Common Name of	Expression Regulation				
No.	Accession	Genes	Description of Genes	Map	p-Value	Tumor	Benign
40.	BE568219	PDE8A	<i>H. sapiens</i> cDNA clone IMAGE: 3683966	15q25.1	0.004205	Up	Down
41.	D50925	STK37, PASKIN,	PAS-serine/threonine kinase	2q37.3	0.004205	Down	Up
42.	NM_006113	VAV3	Vav 3 oncogene	1p13.1	0.004205	Down	Up
43.	NM_001018	RPS15	Ribosomal protein S15	19p13.3	0.006189	Up	Down
44.	NM_021038	MBNL	Muscleblind (<i>Drosophila</i>)-like	3q25	0.006189	Down	Up
45.	NM_012323	U-MAF	V-maf musculoaponeurotic fibrosarcoma, protein F	22q13.1	0.006189	Down	Up
46.	NM_005138	SCO1L	SCO (cytochrome oxidase deficient, yeast) homolog 2	22q13.33	0.006189	Down	Up
47.	AF186779	KIAA0959	RalGDS-like gene	1q25.2	0.006189	Down	Up
48.	D26054	FBP	Fructose-1,6-bisphosphatase 1	9q22.3	0.006189	Up	Down
49.	U37546	API2, MIHC, HIAP1	Baculoviral IAP repeat-containing 3	11q22	0.006189	Down	Up
50.	AB046845	SMURF1	E3 ubiquitin ligase SMURF1	7q21.1-q31.1	0.006189	Down	Up

TABLE 4

The first 50 genes obtained from the supervised MDS analysis of tumor versus benign tissues of all the moderate risk CaP patients, ranked by p-value: (T vs N Intensities of 9 MR)							
Genbank		Common Name	Expression Regulation				
No.	Accession	of Genes	Description of Genes	Map	p-Value	Tumor	Benign
1.	NM_014324	AMACR	Alpha-methylacyl-CoA racemase	5p13.2-q11.1	0	Up	Down
2.	NM_006457	ENH	LIM protein (similar to rat protein kinase C-binding enigma)	4q22	0.000009	Up	Down
3.	AI351043	ESTs	<i>H. sapiens</i> cDNA clone IMAGE: 1948310	21	0.000011	Up	Down
4.	AI433463	MME	<i>H. sapiens</i> cDNA clone similar to NEPRILYSIN (HUMAN)	3q25.1-q25.2	0.000028	Down	Up
5.	BE256479	HSPD1	<i>H. sapiens</i> cDNA clone IMAGE: 3352031	12p13.31	0.000037	Up	Down
6.	NM_015900	PS-PLA1	Phosphatidylserine-specific phospholipase A1alpha	3q13.13-q13.2	0.000083	Up	Down
7.	NM_002343	LTF	Lactotransferrin	3q21-q23	0.000083	Down	Up
8.	NM_001099	PAP	Acid phosphatase, prostate	3q21-q23	0.000083	Down	Up
9.	T15991	CHRM3	IB2413 Infant brain, Bento Soares <i>Homo sapiens</i> cDNA	1q41-q44	0.00011	Up	Down
10.	NM_005084	PAFAH	Phospholipase A2, group VII	6p21.2-p12	0.00011	Up	Down
11.	NM_004503	HOXC6	Homeo box C6	12q12-q13	0.00011	Up	Down
12.	N74607	AQP3	<i>H. sapiens</i> cDNA clone IMAGE: 296424	9p13	0.000149	Down	Up
13.	BC003068	SLC19A1	Solute carrier family 19 (folate transporter), member 1	21q22.3	0.000149	Up	Down
14.	M21535	ERG (ets-related gene)	ERG v-ets erythroblastosis virus E26 oncogene like (avian)	21q22.3	0.000149	Up	Down
15.	NM_013451	MYOF,	Fer-1 (<i>C. elegans</i>)-like 3 (myoferlin)	10q24	0.0002	Down	Up
16.	NM_006017	AC133, CD133	Prominin (mouse)-like 1	4p15.33	0.0002	Down	Up
17.	BE550599	CACNA1D	<i>H. sapiens</i> cDNA clone IMAGE: 3220210	3p14.3	0.0002	Up	Down
18.	U22178	PSP57, PSP94	Microseminoprotein, beta-	10q11.2	0.0002	Down	Up
19.	NM_015865	JK, UT1, UTE	Solute carrier family 14 (urea transporter), member 1	18q11-q12	0.000275	Down	Up
20.	NM_000441	PDS, DFNB4	Solute carrier family 26, member 4	7q31	0.000275	Down	Up
21.	AA877789	MYO6	<i>H. sapiens</i> cDNA clone IMAGE: 1161091	6q13	0.000275	Up	Down
22.	AI356398	ZFP36L2	<i>H. sapiens</i> cDNA clone IMAGE: 2028039	2	0.000275	Down	Up
23.	BC000915	CLIM1, CLP36	PDZ and LIM domain 1 (elfin)	10q22-q26.3	0.000275	Down	Up
24.	NM_000286	PEX12	Peroxisomal biogenesis factor 12	17q11.2	0.000275	Up	Down
25.	NM_003671	CDC14B1, CDC14B2,	<i>Homo sapiens</i> CDC14 cell division cycle 14 homolog B (<i>S. cerevisiae</i>) (CDC14B), transcript variant 1, mRNA	9q22.2-q22.31	0.000386	Down	Up
26.	NM_016545	SBB148	Immediate early response 5	1q24.3	0.000386	Down	Up
27.	NM_002443	PSP57, PSP94	Microseminoprotein, beta-	10q11.2	0.000386	Down	Up
28.	NM_004999	DFNA22	Myosin VI	6q13	0.000386	Up	Down
29.	X99268	TWIST	Twist (<i>Drosophila</i>) homolog	7p21.2	0.000386	Up	Down
30.	NM_023009	MACMARCKS	Macrophage myristoylated alanine-rich C kinase substrate	1p34.3	0.000386	Up	Down
31.	AI721219	TRAF3	as68b11.x1 Barstead colon HPLRB7 <i>Homo sapiens</i> cDNA clone IMAGE: 2333853 3', mRNA sequence.	14q32.33	0.000547	Down	Up
32.	NM_001584	D11S302E	Chromosome 11 open reading frame 8	11p13	0.000547	Down	Up
33.	NM_018846	SBB126	SBB126 protein	7p15.3	0.000547	Up	Down
34.	M87771	BEK, KGFR,	Fibroblast growth factor receptor 2	10q26	0.000547	Down	Up
35.	AF275945	EVA1	Epithelial V-like antigen 1	11q23.3	0.000547	Down	Up
36.	AI791860	ESTs	<i>H. sapiens</i> cDNA clone IMAGE: 1011110		0.000547	Up	Down
37.	BC001282	NHC	High-mobility group (nonhistone chromosomal) protein 17-like 3	6p21.3	0.000547	Down	Up

TABLE 4-continued

The first 50 genes obtained from the supervised MDS analysis of tumor versus benign tissues of all the moderate risk CaP patients, ranked by p-value: (T vs N Intensities of 9 MR)							
Genbank		Common Name		Map	p-Value	Expression Regulation	
No.	Accession	of Genes	Description of Genes			Tumor	Benign
38.	NM_002015	FKH1, FKHR	Forkhead box O1A (rhabdomyosarcoma)	13q14.1	0.000547	Down	Up
39.	X15306	NF-H	<i>H. sapiens</i> NF-H gene, exon 1 (and joined CDS).	22q12.2	0.000547	Down	Up
40.	BE965029	EST	<i>H. sapiens</i> cDNA clone IMAGE: 3886131	11	0.000775	Up	Down
41.	NM_002275	KRT15	Keratin 15	17q21	0.000775	Down	Up
42.	NM_001511	MGSA, CXCL1	GRO1 oncogene	4q21	0.000775	Down	Up
43.	NM_005213	STF1, STFA	Cystatin A (stefin A)	3q21	0.000775	Down	Up
44.	NM_007191	WIF-1	Wnt inhibitory factor-1	12q14.2	0.000775	Down	Up
45.	H15129	MEIS3	EPIDERMAL GROWTH FACTOR-LIKE CRIPTO PROTEIN	17	0.000775	Down	Up
46.	AW452623	EST	<i>H. sapiens</i> cDNA clone IMAGE: 3068608	13	0.000775	Up	Down
47.	X90579	EST	<i>H. sapiens</i> DNA for cyp related pseudogene	7	0.000775	Down	Up
48.	BC001388	ANX2, ANX2L4	Annexin A2	15q21-q22	0.001116	Down	Up
49.	NM_014863	BRAG,	B cell RAG associated protein	10q26	0.001116	Down	Up
50.	NM_021076	NEFH	Neurofilament, heavy polypeptide (200 kD)	22q12.2	0.001116	Down	Up

TABLE 5

Top 50 Unregulated Genes in All the 18 Samples (HR and MR) obtained from Tumor over Benign (T/B) ratio.					
Genbank No ID	T/N ratio	Common Name of Genes	Description	Map	
1. AF047020	39.86910	AMACR	Alpha-methylacyl-CoA racemase	5p13.2-q11.1	
2. M54886	20.86411	LOC51334	Mesenchymal stem cell protein DSC54	5p13.1	
3. AF070581	19.07263	ESTs	<i>Homo sapiens</i> cDNA clone IMAGE: 1948310	21	
4. NM_014324	18.04841	TRG@	T cell receptor gamma locus	7p15-p14	
5. NM_001669	15.98177	NPY	Neuropeptide Y	7p15.1	
6. NM_018360	13.34037	HOXC6	Homeo box C6	12q12-q13	
7. AF092132	9.588665	IMP2	IMP (inosine monophosphate) dehydrogenase 2	3p21.2	
8. NM_023067	7.712272	HSPC028	HSPC028 protein	7p21.2	
9. NM_014439	7.031155	LTBP1	Latent transforming growth factor beta binding protein 1	2p22-p21	
10. AI613045	6.739595	GMF	Glia maturation factor, beta	14q22.1	
11. AB051446	6.563991	DSC2	HUMAN Q02487 DESMOCOLLIN 2A/2B PRECURSOR	18q12.1	
12. NM_005342	6.442383	TRG, TCRG	T cell receptor gamma locus	7p15-p14	
13. D87012	6.327042	PAWR	<i>H. sapiens</i> cDNA clone IMAGE: 1950862	12q21	
14. NM_018221	6.098105	SNX2	Sorting nexin 2	5q23	
15. NM_005114	5.769173	HS3ST1	Heparan sulfate (glucosamine)-3-O-sulfotransferase 1	11	
16. NM_022831	5.624385	RA70, SAPS, SKAP55R	Src family associated phosphoprotein 2	7p21-p15	
17. NM_014324	5.621786	TRG, TCRG	T cell receptor gamma locus	7p15-p14	
18. NM_006820	5.550019	BICD1	Bicaudal D (<i>Drosophila</i>) homolog 1	12p11.2-p11.1	
19. NM_005574	5.454622	FOLH1	Folate hydrolase (prostate-specific membrane antigen) 1	11p11.2	
20. AL365343	5.451875	KIAA0615	<i>Homo sapiens</i> mRNA for KIAA0615 protein, complete cds.	16q11.2	
21. NM_022580	5.318270	TBCE	Tubulin-specific chaperone e	1q42.3	
22. AK022765	5.315669	CLDN8	Claudin 8	21	
23. AF067173	5.272626	P21, NSG1, D4S234	Neuron-specific protein	4p16.3	
24. NM_006220	5.180025	SHMT2	<i>Homo sapiens</i> cDNA clone IMAGE: 2676158	12q12-q14	
25. AL133600	5.146792	ANK2	<i>Homo sapiens</i> cDNA clone by03a08	4q25-q27	
26. AY009108	5.097967	PSM	PROSTATE-SPECIFIC MEMBRANE ANTIGEN (HUMAN)	2	
27. AL035603	5.076761	FLJ10907	Ribonuclease 6 precursor	6q27	
28. NM_014017	5.058610	MAPBPIP	Mitogen-activated protein-binding protein-interacting protein	13	
29. BF247098	5.030722	PHLP, DKFZp564M1863	Phosducin-like	9q12-q13	
30. U62296	4.992345	GOLPH2	Golgi phosphoprotein 2	9	
31. AF130082	4.988912	EST	<i>Homo sapiens</i> clone FLC1492 PRO3121 mRNA, complete cds	8	
32. NM_020373	4.969535	C8orf4	Chromosome 8 open reading frame 4	6	
33. U90030	4.873056	BICD1	Bicaudal D homolog 1 (<i>Drosophila</i>)	6p22.2-p21.3	
34. NM_021071	4.821960	KIAA0426	KIAA0426 gene product	12q13.3-q14.1	
35. NM_030817	4.753895	KIAA1157	KIAA1157 protein	Xq26.1	
36. NM_019844	4.700642	HPRT, HGPRT	Hypoxanthine phosphoribosyltransferase 1	3p21.3-p21.2	
37. NM_004721	4.689246	RPL29	Ribosomal protein L29	19pter-q12	
38. NM_004866	4.669274	EF2, EEF-2	Eukaryotic translation elongation factor 2	Xq28	
39. NM_014501	4.610132	BGN	Biglycan	8q22-q23	
40. NM_020655	4.575193	SDC2	Syndecan 2 (heparan sulfate proteoglycan 1, fibroglycan)	19p13.11	
41. NM_006716	4.557526	ASK	Activator of S phase kinase	11q14.3	
42. NM_002968	4.541752	FOLH1	Folate hydrolase (prostate-specific membrane antigen) 1		

TABLE 5-continued

Top 50 Unregulated Genes in All the 18 Samples (HR and MR) obtained from Tumor over Benign (T/B) ratio.				
Genbank No ID	T/N ratio	Common Name of Genes	Description	Map
43. X06268	4.539479	NCUBE1	Non-canonical ubiquitin conjugating enzyme 1	6
44. AK021609	4.520464	PTH2, PTEN2, PSPTEN	Phosphatase and tensin homolog (mutated in multiple advanced cancers 1), pseudogene 1	9p21
45. NM_001133	4.479513	TCTEX1L	T-complex-associated-testis-expressed 1-like	Xp21
46. D38491	4.477160	KIAA0461, POGZ,	Pogo transposable element with ZNF domain, KIAA0461 protein	1q21.2
47. NM_006426	4.385531	DDX26	Deleted in cancer 1; RNA helicase HDB/DICE1	13q14.12-q14.2
48. AW058148	4.347362	SPHAR	S-phase response (cyclin-related)	1q42.11-q42.3
49. U55209	4.293919	MYO7A	myosin VIIA (Usher syndrome 1B)	4
50. NM_004610	4.275521	KIAA0634, ASTN2	Astrotactin 2	9q33.1

TABLE 6

Top 35 Downregulated Genes in All the 18 Samples (HR and MR) obtained from Tumor over Benign (T/B) ratio.					
No.	Genbank ID	T/N Ratio	Common Name of the Genes	Description	Map
1.	X90579	0.181138	CYP3A5	Cytochrome P450, family 3, subfamily A, polypeptide 5	7
2.	NM_005213	0.198502	STF1, STFA	Cystatin A (stefin A)	3q21
3.	NM_005864	0.254524	EF51, HEFS	Signal transduction protein (SH3 containing)	14q11.2-q12
4.	X15306	0.291665	NF-H	<i>H. sapiens</i> NF-H gene, exon 1 (and joined CDS).	22q12.2
5.	BE908217	0.319347	ANXA2	Annexin A2	15q21-q22
6.	BC001388	0.320110	ANX2, LIP2, ANX2L4	Annexin A2	15q21-q22
7.	U22178	0.326560	PSP57, PSP94, PSP-94	Microseminoprotein, beta-	10q11.2
8.	NM_002443	0.338948	PSP57, PSP94, PSP-94	Microseminoprotein, beta-	10q11.2
9.	NM_021076	0.359039	NEFH	Neurofilament, heavy polypeptide (200 kD)	22q12.2
10.	AI433463	0.360636	MME, CD10, NEP, CALLA	Nephrilysin	3q25.1-q25.2
11.	AF275945	0.366939	EVA1	Epithelial V-like antigen 1	11q23.3
12.	NM_002343	0.370305	LTF	Lactotransferrin	3q21-q23
13.	NM_013451	0.378555	MYOF, KIAA1207	Fer-1 (<i>C. elegans</i>)-like 3 (myoferlin)	10q24
14.	NM_001584	0.385272	239FB, D11S302E	Chromosome 11 open reading frame 8	11p13
15.	AL390736	0.391520	BA209J19.1, GW112	GW112(differentially expressed in hematopoietic lineages)	
16.	NM_000441	0.392117	PDS, DFNB4	Solute carrier family 26, member 4	7q31
17.	AL031602	0.399115	ESTs	ESTs	1p34.1-35.3
18.	NM_004039	0.399796	ANXA2	Annexin A2	15q21-q22
19.	NM_001546	0.402261	ID4	DNA binding inhibitor protein of ID-4	6p22-p21
20.	NM_001099	0.406234	PAP	Acid phosphatase, prostate	3q21-q23
21.	X57348	0.422692	9112	<i>H. sapiens</i> mRNA (clone 9112).	1p35.2
22.	NM_020139	0.440648	LOC56898	Oxidoreductase UCPA	4
23.	AU148057	0.444528	DKK3, REIC	Dickkopf related protein-3 precursor (Dkk-3) (Dickkopf-3) (hDkk-3)	11pter-p15.5
24.	BF059159	0.446108	ROBO1, DUTT1, SAX3	Roundabout, axon guidance receptor, homolog 1 (<i>Drosophila</i>)	3p12
25.	BC001120	0.448109	MAC2, GALBP, MAC-2,	Lectin, galactoside-binding, soluble, 3 (galectin 3)	14q21-q22
26.	N74607	0.451123	AQP3	Aquaporin 3	9p13
27.	NM_013281	0.454835	FLRT3	Fibronectin leucine rich transmembrane protein 3	20p11
28.	NM_000700	0.456566	ANX1, LPC1	Annexin A1	9q12-q21.2
29.	X57348	0.458169	9112	<i>H. sapiens</i> mRNA (clone 9112).	1p35.2
30.	AI356398	0.467028	ZFP36L2, ERF-2, TIS11D	EGF-respons factor 2	2
31.	AF016266	0.467787	DR5, TRAILR2, TRICK2A,	Tumor necrosis factor receptor superfamily, member 10b	8p22-p21
32.	S59049	0.467913	BL34, IER1, IR20	Regulator of G-protein signalling 1	1q31
33.	NM_000165	0.470393	GJA1	Gap junction protein, alpha 1, 43 kD (connexin 43)	6q21-q23.2
34.	AI826799	0.471081	EFEMP1, DRAD, FBNL	EGF-CONTAINING FIBULIN-LIKE EXTRACELLULAR MATRIX PROTEIN 1	2p16
35.	AL575509	0.476538	ETS2	V-ets erythroblastosis virus E26 oncogene homolog 2 (avian)	21q22.2

Classification between Tumor and Benign Prostate Epithelium:

A class prediction analysis using distance based Multi Dimensional Scaling (MDS) was used to determine expression differences between tumor and benign epithelial cells in 18 patients with radical prostatectomy. All the genes that meet a minimum level of expression were included in the

analysis. We used the normalized intensities of all the 18 tumor and 18 normal samples for a class prediction analysis by distance based MDS to determine differentiation between tumor and benign tissue specific gene expression profile among all the 18 patients. Using a matrix of Pearson correlation coefficients from the complete pair-wise comparison of all the experiments we observed a significant

overall difference in gene expression pattern between the tumor and benign tissue as displayed as a two-dimensional MDS plot in FIG. 2A. The position of the each tumor and benign samples is displayed in the MDS plot in two-dimensional Euclidean space with the distance among the samples reflecting correlation among the samples in each individual group (distance within the cluster) and as well as reflecting distinct separation between the two groups (center-to center distance) (FIG. 2A). The MDS plot was obtained from the top 200 genes obtained by 10,000 permutations of the tumor and benign intensities of 4566 genes. Out of these 200 genes that define the tumor specific alteration of gene expression, 53 genes had higher expression in the tumor samples and the remaining 147 genes had higher expression in the benign samples. A partial list of genes that distinctly discriminate the tumor and benign samples from all the 18 patients is shown in Table 1. We also performed a hierarchical clustering analysis using the 200 discriminatory genes. The hierarchical clustering algorithm resulted in a hierarchical dendrogram that identified two major distinct clusters of 16 tumor samples and 17 benign samples (FIG. 2B).

Classification of CaP into HR and MR Groups Using the Ratio of Tumor over Benign Gene Expression Intensities

We used the tumor over benign gene expression intensity ratio (T/B ratio) (FIG. 3A) from the HR (9 patients) and MR (9 patients) groups for a class prediction analysis using distance based MDS method to determine if the 18 patients can be differentiated into the two patient groups. Pathological and clinical features of the 18 tumors used in our study were clearly distinguishable between the HR and MR groups. We observed a significant overall difference in expression pattern between the HR and MR groups. The distance between the samples reflects both the extent of correlation within each individual selected group (distance within the cluster) as well as distinct separation between the two selected groups (center-to-center distance) (FIG. 3A). The MDS plot obtained from top 200 genes by 10,000

permutations of the 4868 genes based on the T/B ratio is shown in FIG. 3A. Out of the top 200 genes of the MDS analysis 135 were over expressed in the HR group and 65 genes were over expressed in the MR group. The top 50 genes with best p-values identified by the T/B ratio based MDS analysis discriminating the HR and MR groups are listed in FIG. 3B. The approach we used for the interpretation of discrimination between the HR and MR groups was empirical. The 'weighted list' (FIG. 3B) of individual genes whose variance of change across all the tumor samples defines the boundary of a given cluster to predict a class that correlates with the pathological and clinical features of CaP. We also performed a hierarchical clustering to verify the results of the MDS analysis and also to test the potential of those 200 genes to predict class/group (HR and or MR) using another approach of analysis. The resulting hierarchical dendrogram of TB ratio demonstrates that 9 samples of the HR group formed a very distinct and tight cluster, as did the 9 samples of MR group (FIG. 3B).

Classification of CaP into HR and MR groups based on Gene Expression Intensities in Tumor Cells

MDS analysis was used to determine differentiation among 18 patients into HR and MR groups. An overall difference in tumor specific expression between the HR and MR groups is displayed as a two-dimensional MDS plot (FIG. 3C). The MDS plot obtained from 10,000 permutations of the gene expression intensities of 4115 genes from the tumor samples of 18 patients differentiated them into HR and MR groups based on the selected top 200 genes (FIG. 3C). Out of this 200 genes, 94 had higher expression in the HR groups and the remaining 106 genes had higher expression in the MR groups. We performed a hierarchical clustering analysis using the 200 discriminatory genes obtained from the supervised MDS analysis. The resulting hierarchical dendrogram of 18 tumor samples demonstrates that 9 tumor samples of the HR group and 9 tumor samples of the MR group were separated into two tight clusters. (FIG. 3D). The approach we utilized on the basis of the linear correlation of global gene expression in FIG. 3 to obtain 'gene cluster' interpretation to discriminate the HR and MR groups was empirical. Genes that discriminate the HR and MR groups are shown in Table 7.

TABLE 7

Top 17 genes analysis based on T/B fold change of HR vs MR groups								
Gene Bank ID	Common Name	Description	Map	p-Value	HR	MR	Absent	Positive
1 NM_004522	KINN, NKIIC	Kinesin family member 5C	2q23.3	0.0001	Up	Down	4	60%
3 NM_018010	HIPPI, FLJ10	Hypothetical protein FLJ10147	3q13.13	0.0033	Up	Down	0	56%
10 NM_012245	SKIP, NCOA-	SKI-interacting protein	14q24.3	0.0076	Up	Down	2	42.80%
11 NM_015895	LOC51053	Geminin	6p22.2	0.0076	Up	Down	2	71.40%
14 NM_003031	SLAH1	Seven in absentia (<i>Drosophila</i>) homolog 1	16q12	0.0076	Up	Down	3	66%
42 NM_000016	ACADM	Acyl-Coenzyme A dehydrogenase, C-4 to C-12 straight	1p31	0.0178	Up	Down	1	75%
47 NM_025087	FLJ21511	Hypothetical protein FLJ21511	4	0.0178	Up	Down	1	50%
17 NM_021038	MBNL	Muscleblind (<i>Drosophila</i>)-like	3q25	0.0076	Down	Up	2	71
25 NM_006732	GOS3	FBJ murine osteosarcoma viral oncogene homolog B	19q13.32	0.0076	Down	Up	3	83%
51 NM_001674	ATF3	Activating transcription factor 3	1q32.3	0.0178	Down	Up	0	100%
7 NM_002053	GBP1	Guanylate binding protein 1, interferon-inducible, 67 KD	1p22.1	0.005	Down	Up	4	83%
15 NM_003407	TTP, GOS24	Zinc finger protein 36, C3H type, homolog (mouse)	19q13.1	0.0076	Down	Up	1	62%
26 NM_003760	EIF4G3	Eukaryotic translation initiation factor 4 gamma, 3	1pter-p3	0.0076	Down	Up	4	40%
38 AK023938	<i>Homo sapien</i>	SELECTED MODEL ORGANISM PROTEIN SIMILARITIES	2q37.3	0.0178	Down	Up	4	80%
45 NM_016021	NCUBE1	Non-canonical ubiquitin conjugating enzyme 1	6	0.0178	Down	Up ?	3	66%
5 NM_021795	SAP1	ELK4, ETS-domain protein (SRF accessory protein 1)	1q32	0.005	Up	Down	4	80%
18 NM_014454	PA26	P53 regulated PA26 nuclear protein	6q21	0.0076	Up	Down	3	83%

Classification of CaP into High Risk and Medium Risk Groups Based on Gene Expression Intensities in Benign Prostate Epithelial

We used a similar MDS and Cluster analysis as in the tumor versus tumor sample gene expression intensities for the normalized intensities of 9 benign samples of HR group and 9 benign samples of MR group for a class prediction. Strikingly the MDS plot of the benign samples depicted distinct separation between the HR and MR groups (FIG. 3E). We observed a significant overall difference in expression pattern between the HR and MR groups. The MDS plot obtained from the top 200 genes by 10,000 permutations of the 3358 genes from the benign versus benign intensities (FIG. 3E). Out of this 200 genes 61 were over expressed in benign samples of the HR groups and the remaining 139 genes were over expressed in the MR groups. The 'weighted list' of individual genes whose variance of expression alteration across all the normal samples depicts the capability of a given cluster to predict classification. The hierarchical clustering algorithm identified a similar major cluster of the 9 benign samples of the HR group and a cluster of 9 benign samples of the MR group.

The weighted gene analysis by distance based supervised multidimensional scaling method we used, (depicted in FIGS. 3A, 3C, and 3E) utilizing the gene expression ratio of tumor and benign intensities, gene expression intensities of tumor samples and as well as normal for obtaining a 'weighted list' of individual genes, whose variance of change across all the tumor and benign samples distinctly delineate the boundary of a given cluster, to predict a class that correlates with the pathological and clinical features of CaP.

Independent In silico Cross Validation

In silico analysis for the predicted classifier was carried out using two independent data sets. The HR and MR groups were selected on the basis of Gleason score as that was the only criterion available for these data. At least 200 genes were extracted from all the MDS analysis (see methods for detail description). This subset of 200 classifier genes were found in the data of Welsh et al. 2001 and Sing et al. 2002. Exactly similar MDS analysis ($p < 0.001$ as measured by 10,000 permutation testing) as described above was performed using the expression intensities of these 200 genes from Welsh and Singh data. MDS analysis using tumor over benign ratio of as low as 50 genes from the subset of 200 genes from Welsh data (FIG. 4A) as well as Singh data (FIG. 4B) clearly separated samples from HR group and samples from MR group. Thus, this observation elucidates that the differential expression profile of this small set of genes can be used to predict the identity or class or group of unknown prostate cancer samples on the basis of their clinico-pathological features. The outcome of this analysis depicts that the expression profile of this small number of genes is conserved across the independent data sets.

Validation of GeneChip Results by Real-Time PCR

To further validate the expression alterations of genes identified by GeneChip analysis with an indicated biological relevance to prostate cancer, primers and probes were obtained for real-time PCR analysis using AMACR and GSTP1. These genes were chosen for validation purposes because it has been reported previously by several investigators that AMACR is elevated and GSTP1 decreased in CaP. Each sample demonstrated a unique pattern of down-regulation of GSTP1 gene in 18 of 20 samples as well as up-regulation of AMACR (FIG. 1) the other two samples did show significant change (fold change less than 1.5).

One ng of total RNA samples from paired tumor and normal specimens was reverse-transcribed using Omnisensecript RT-kit (Qiagen, Valencia, Calif.) according to the manufacturer's protocol.

Quantitative gene expression analysis was performed using TaqMan Master Mix Reagent and an ABI prism 7700 Sequence Detection System (PE Applied Biosystems Foster, Calif.). All sets of primer and probe for tested genes were Assays-on-Demand Gene expression products obtained from PE Applied Biosystems. The expression of house keeping gene, GAPDH was simultaneously analyzed as the endogenous control of same batch of cDNA, and the target gene expression of each sample was normalized to GAPDH. For each PCR run, a master-mix was prepared on ice with 1x TaqMan Master Mix, 1x target gene primer/probe and 1xGAPDH primer/probe. Two microliters of each diluted cDNA sample was added to 28 μ l of PCR master-mix. The thermal cycling conditions comprised an initial denaturation step at 95° C. for 10 minutes and 50 cycles at 95° C. for 15 seconds and 60° C. for 1 minute. RNA samples without reverse transcription were included as the negative control in each assay. All assays were performed in duplicate. Results were plotted as average C_T (threshold cycle) of duplicated samples. The relative gene expression level was presented as "Fold Change" of tumor versus matched normal cells, which is calculated as: $\text{Fold change} = 2^{(\Delta C_{T\text{normal}} - \Delta C_{T\text{tumor}})}$, where ΔC_T means normalized C_T value of target genes to GAPDH.

Example 3

Distinguishing Between ERG1 and ERG2 Isoforms

The Affymetrix GeneChip probe set (213541_s_at) and TaqMan probes used in the experiments described above recognize a region specific for both ERG1 and ERG2 isoforms (FIG. 6), but exclude isoforms 3 to 9. Although other primers and probes could be used, by way of example, TaqMan primers and probe recognizing both ERG1 and ERG2, but not other ERG isoforms were as follows:

```
Fwd primer: (SEQ ID NO: 7)
5'-AGAGAAACATTCAGGACCTCATCATTATG-3'

Reverse primer: (SEQ ID NO: 8)
5'-GCAGCCAAGAAGGCCATCT-3'

Probe: (SEQ ID NO: 9)
5'-TTGTTCTCCACAGGGT-3'
```

The probe has the reporter dye, 6-FAM, attached to the 5' end and TAMRA attached to the 3' end. The 3'-TAMRA effectively blocks extension during PCR.

To further distinguish between these two ERG isoforms, the expression of the ERG1 and ERG2 isoforms were tested in PC3 cells and in normal prostate tissue (pooled prostate RNA from 20 men, Clontech), as well as in microdissected tumor and normal prostate epithelial cells from 5 CaP patients (data not shown). Only ERG1 was expressed in the prostate cells and in PC3 cells. ERG2 expression was not detectable. A TaqMan QRT-PCR probe and primers were designed that specifically recognize only the ERG1 isoform (FIG. 6). Although other primers and probes could be used, by way of example, we designed TaqMan primers and probes recognizing only the ERG1 isoform as follows:

Forward primer:
 5' - CAGGTCCTTCTTGCCTCCC - 3' (SEQ ID NO: 10)

Reverse primer:
 5' - TATGGAGGCTCCAATTGAAACC - 3' (SEQ ID NO: 11)

Probe:
 5' - TGTCTTTTATTCTAGCCCTTTTGGACAGGA - 3' (SEQ ID NO: 12)

The probe has the reporter dye, 6-FAM, attached to the 5' end and TAMRA attached to the 3' end. The 3'-TAMRA effectively blocks extension during PCR.

ERG1 expression was determined in 228 RNA specimens from microdissected matched tumor and benign prostate epithelial cells of 114 CaP patients. Overall, 62.4% of the 114 CaP patients analyzed had significant over expression of ERG1 isoform in their tumor cells (i.e., greater than 2 fold ERG1 expression in tumor versus benign cells), while 16.6% of CaP patients had no detectable ERG1 expression, 15.0% had under expression of ERG1 (less than 0.5 fold difference in ERG1 expression in tumor versus benign cells), and 6.0% had no significant difference (0.5 to 2 fold difference in ERG1 expression between tumor versus benign cells).

In a further study, ERG expression was analyzed in 82 CaP patients. Using the TaqMan primers and probes discussed above, we observed tumor-associated over expression of ERG1 (isoform 1 only) and ERG (isoforms 1 and 2) in 63.4% and 72.0% of the patients, respectively. Therefore, ERG1 isoform specific expression may actually reflect an underestimate of the overall ERG expression in CaP.

Example 4

Correlation of ERG1 Expression With Various Clinico-Pathologic Features

Since the ERG1 tumor versus benign expression ratio data did not have normal distribution, the Wilcoxon Rank Sum Test was used to analyze its relationship with various clinico-pathologic features, as shown in Table 8.

TABLE 8

Relationship of ERG1 expression ratios in tumor versus benign prostate epithelial cells with patient clinical factors				
Clinical factors	N	Median of ERG1 fold changes	Mean scores of ERG1 fold changes	P
PSA recurrence				
No	75	142.2	52.19	0.0042
Yes	20	1.2	32.30	
Tumor Differentiation				
Well & Moderate	40	362.3	57.62	0.0020
Poor	54	13.9	40.00	
Pathologic T stage				
pT2	38	502.0	53.45	0.0136
pT3-4	52	33.5	39.69	
Margin status				
Negative	64	197.0	52.55	0.0209
Positive	31	20.4	38.61	
Seminal vesicle				
Negative	82	106.7	49.28	0.2555
Positive	13	6.9	39.92	
Race				
Caucasian	73	172.1	52.08	0.0086

TABLE 8-continued

Relationship of ERG1 expression ratios in tumor versus benign prostate epithelial cells with patient clinical factors				
Clinical factors	N	Median of ERG1 fold changes	Mean scores of ERG1 fold changes	P
African American	22	3.8	34.45	
Family history				
No	70	106.7	49.46	0.3887
Yes	25	4.8	43.92	
Diagnostic PSA (ng/ml)				
<=4	13	101.3	57.15	0.1801
>4-10	62	112.1	48.03	
>10	19	20.5	39.16	
Gleason sum				
<7	33	112.0	52.06	0.2923
=7	45	118.1	47.16	
>7	16	21.0	39.06	

As shown in Table 8, 95 CaP patients with detectable ERG1 expression were analyzed by Wilcoxon rank sum test. N represents the number of CaP patients falling into the indicated clinical factor category. Significant p values (<0.05) are in bold face.

We also found a significant correlation of high ERG1 over expression with Caucasian over African American ethnicity (p=0.0086) (Table 8). To further explore the correlation with PSA recurrence, Kaplan-Meier survival analysis was performed based on three patient groups: 1) CaP patients with tumor versus benign ERG1 expression ratio of less than 2 fold; 2) CaP patients with tumor versus benign ERG1 expression ratio of 2-100 fold; and 3) CaP patients with tumor versus benign ERG1 expression ratio of greater than 100 fold (FIG. 7). The results show that patients with higher ERG1 over expression in their prostate tumor tissue had significantly longer PSA recurrence-free survival (log rank test, P=0.0006) (FIG. 7). The 36-months PSA recurrence-free survival for patients with less than 2 fold ERG1 expression ratio (n=24) was 54.4%, while for patients with greater than 100 fold ERG1 expression ratio (n=47) it was 87.7%. From a univariate COX proportional hazard ratio regression analysis for PSA recurrence-free time using ERG1 tumor/benign cells expression ratio, race, diagnostic PSA, Gleason sum, pathologic T stage, margin status, and seminal vesicle invasion status, we found that five of these variables (ERG1 tumor/benign cells expression ratio, Gleason sum, pathologic T stage, margin status, seminal vesicle invasion) had a significant p value (Table 9).

TABLE 9

Correlation of clinical parameters and ERG1 expression ratios in tumor versus benign prostate epithelial cells with PSA recurrence-free time after radical prostatectomy		
Factors	Crude Hazard Ratio (95% CI)	P
ERG1 fold changes		
2-100 fold vs. <2 fold	0.291 (0.093-0.915)	0.0347
>100 fold vs. <2 fold	0.173 (0.060-0.498)	0.0011
Race		
Caucasian vs. African American	1.092 (0.395-3.016)	0.8657
Diagnostic PSA		0.8723
>4-10 vs. <=4	0.976 (0.275-3.468)	0.9705
>10 vs. <=4	1.285 (0.307-5.378)	0.7313
Gleason Sum		
7 vs. 2-6	1.574 (0.393-6.296)	0.0001
8-10 vs. 2-6	9.899 (2.752-35.610)	0.0004

TABLE 9-continued

Correlation of clinical parameters and ERG1 expression ratios in tumor versus benign prostate epithelial cells with PSA recurrence-free time after radical prostatectomy		
Factors	Crude Hazard Ratio (95% CI)	P
Pathologic T stage		
pT3/4 vs. pT2 Margin status	6.572 (1.517-28.461)	0.0118
Positive vs. Negative Seminal Vesicle	2.825 (1.169-6.826)	0.0210
Positive vs. Negative	3.792 (1.487-9.672)	0.0053

In Table 9, crude hazard ratios with 95% confidence interval are shown for ERG1 fold change (tumor versus benign) and six clinical parameter categories in a univariate COX proportional hazard ratio analysis. Significant p values are in bold face. The multivariate COX proportional hazard ratio regression analysis of the significant variables from the univariate analysis shows that ERG1 overexpression (greater than 100 fold vs. less than 2 fold: p=0.0239, RR=0.274, overall p value 0.0369), and Gleason sum (Gleason 8-10 vs. Gleason 2-6: p=0.0478, RR=4.078, overall p value 0.0148) are independent predictors of PSA recurrence after radical prostatectomy (Table 10). These results demonstrate that the status of ERG1 expression ratios (tumor vs. benign) in radical prostatectomy specimens carries a predictive value for patient prognosis.

TABLE 10

Factors	Crude Hazard Ratio (95% CI)	P
ERG1 fold changes		
2-100 fold versus <2 fold	0.320 (0.097-1.059)	0.0369
>100 fold versus <2 fold	0.274 (0.089-0.843)	0.0620
Gleason Sum		0.0148
7 versus 2-6	0.948 (0.223-4.033)	0.9424
8-10 versus 2-6	4.078 (1.014-16.401)	0.0478
Pathologic T stage		
PT3/4 versus pT2 Margin status	3.306 (0.636-17.177)	0.1550
Positive versus Negative Seminal Vesicle	1.116 (0.421-2.959)	0.8254
Positive versus Negative	1.308 (0.466-3.670)	0.6098

ERG1 expression in prostate tumor tissue showed highly significant association with longer PSA recurrence free survival (p=0.0042), well and moderately differentiated grade (p=0.0020), lower pathologic T stage (p=0.0136), and negative surgical margin status (p=0.0209), suggesting that ERG1 over expression in tumor cells is generally higher in less aggressive CaP than in more aggressive CaP (Table 8).

The ERG1 over expression in tumor cells identified by GeneChip analysis and verified by real time QRT-PCR assays was further validated by in situ hybridization. Based on the real time QRT-PCR data, 6 patients with high ERG1 over expression in their tumor cells (and as a control one patient with no ERG1 over expression) were selected for in situ hybridization and quantitative image analysis in a blinded fashion. As expected, in each case the in situ expression data confirmed the over expression of ERG1 in the tumor epithelial cells (FIG. 8).

Example 5

Generation and Characterization of ERG Antibody

Cloning of ERG1 into Tetracycline Regulated Mammalian Expression Vectors:

ERG1 cDNA was subcloned into tetracycline-regulated mammalian expression vectors (pTet-off, EC1214A). The constructs generated include, pTet-off-ERG1 (sense), pTet-off-ERG1 (antisense), pTet-off-FlagERG1 (sense) and pTet-off-FlagERG1 (antisense). Originally, ERG1 construct in a riboprobe vector pGEM was obtained from Dr. Dennis K. Watson, Medical University of South Carolina. The constructs were verified by dideoxy sequencing and agarose gel analysis.

Generation of Polyclonal ERG Antibody:

Antibodies against ERG were generated using peptide antigens derived from the full length ERG1 coding sequence. The epitope for the antigen were carefully selected such that the antibody recognizes specifically ERG1/2/3 and not other members of the ETS family (FIG. 9). The following peptides, having the highest hydrophilicity (-1.26 and -0.55) and antigenicity in the desired region, were used to generate antibodies:

Peptide M-50-mer:

(SEQ ID NO: 13)

CKALQNSPRLMHARNTDLPYEPPRRSAWTGHGHPTQSKAAQSPSTVP

K-[NH₂]

Peptide C-49-mer:

(SEQ ID NO: 14)

CDFHGIAQALQPHPPESLYKYPYSDLPYMGSYHAHPQKMNFFVAPHPAL

Cysteine was added to each peptide for conjugation. Peptide M is amidated at the C-terminal residue because it is an internal peptide.

The synthesis of the peptide epitopes and the immunization of rabbits were carried out in collaboration with Bio-Synthesis Inc. Two rabbits were immunized for each of the two epitopes. Bleeds obtained post immunization were collected and tested. Subsequently, bleeds from one of the rabbits from each epitope were affinity purified using SulfoLink kit (Pierce) and were verified by immunoblot analysis.

Characterization of Polyclonal ERG Antibody by Immunoblot Analysis:

To characterize the affinity purified antibody, we transiently transfected HEK-293 (Human embryonic kidney cell line, ATCC, Manassas, Va.) with ERG1 constructs pTet-off-ERG1 (sense) and pTet-off-FlagERG1 (sense) using Lipofectamine reagent (Invitrogen, Carlsbad, Calif.) as per manufacturers instructions. HEK-293 that were not transfected with the plasmid served as a transfection control. The cells were harvested 48 hours post-transfection and processed for immunoblot analysis. Expression of ERG1 following transfection was determined by immunoblotting using the affinity purified polyclonal antisera generated against the unique M- and C-ERG epitopes described above. Endogenous ERG1 expression was not detected in non-transfected HEK-293 cells. However, the ERG antibodies detected ERG1 expression in HEK-293 cells transfected with the various ERG1 constructs. Tetracycline (2 ug/ml) abolished ERG1 expression in both tetracycline-regulated constructs, pTet-off-ERG1 (sense) and pTet-off-FlagERG1

(sense). The M2-Flag antibody specifically recognized only the Flag-tagged ERG1 protein.

Example 6

Combined Expression of ERG, AMACR, and DD3 Genes in Prostate Tumors

The strikingly high frequency of ERG over expression in CaP cells led to a comparison of ERG expression with two other genes, AMACR and DD3, that are also over expressed in CaP cells. We have evaluated quantitative gene expression features of AMACR and DD3, along with the ERG gene, in laser microdissected matched tumor and benign prostate epithelial cells from 55 CaP patients.

Although other primers and probes can be used, by way of example, we designed the following TaqMan primers and probe recognizing the DD3 gene:

Forward primer: (SEQ ID NO: 15)
5'-CACATTTCAGCCCCCTTAAATA-3'
Reverse primer: (SEQ ID NO: 16)
5'-GGGCGAGGCTCATCGAT-3'
Probe: (SEQ ID NO: 17)
5'-GGAAGCACAGAGATCCCTGGGAGAAATG-3'.

The probe has the reporter dye, 6-FAM, attached to the 5' end and TAMRA attached to the 3' end. The 3'-TAMRA effectively blocks extension during PCR.

AMACR TaqMan primers and probe were purchased from Applied Biosystems.

AMACR and DD3 showed upregulation in tumor cells of 78.2% and 87.3% of CaP patients, respectively (FIG. 5). ERG over expression in tumor cells was detected in 78.2% of the same group of CaP patients (FIG. 5). Comparative expression analysis revealed that when the AMACR and ERG expression data are combined, 96.4% of the CaP patients showed upregulation of either of the two genes in tumor cells (FIG. 5). Similarly, the combination of the ERG and DD3 expression data improved the cancer detection power of either of the genes to 96.4% (FIG. 5). When combining the expression data from all the three genes, 98.2% of the CaP patients showed upregulation of at least one of the three genes in tumor cells (FIG. 5). Thus,

screening for ERG gene expression, alone, or in combination with other genes that are over expressed in CaP, such as AMACR and DD3, provides a new, powerful diagnostic and prognostic tool for CaP.

Example 7

Under Expression of LTF in Malignant Prostate Epithelium

One of the most consistently under expressed genes in CaP cells was LTF (Table 1). Validation by QRT-PCR (TaqMan) in LCM-derived tumor and benign prostate epithelial cells confirmed a consistent, tumor associated LTF under expression in 100% of CaP cells tested (FIG. 1D). As a quality control, the expression of AMACR, a recently identified CaP tissue marker, and of GSTP1, a gene showing commonly reduced or absent expression in CaP (Nelson et al., Ann. N.Y. Acad. Sci., 952:135-44 (2001)), was also determined (FIGS. 1B and 1C, respectively). Robust under expression similar to LTF, was observed for GSTP1, while the increased expression of AMACR was noted in 95% of the tumor cells tested, confirming the high quality of the tumor and benign LCM specimens and the reliability of the QRT-PCR. In a further study, LTF expression was analyzed by QRT-PCR in in microdissected tumor and benign prostate epithelial cells of 103 CaP patients. The results were consistent with the initial results, showing tumor associated under expression in 76% of patients (78 of 103).

LTF under expression was also validated at the protein level with anti-LTF goat polyclonal antibody (Santa Cruz, Calif., sc-14434) using Western blot analysis on protein lysates and immunohistochemistry techniques. Hematoxylin-eosin (H&E) and LTF staining was performed on tissue samples from 30 CaP patients by immunocytochemical analysis. In 30 of 30 (100%) cases, benign epithelial cells adjacent to tumor cells were highly positive for LTF, whereas, on average, less than 10% of prostate tumor cells revealed LTF positive cytoplasmic staining.

The specification is most thoroughly understood in light of the teachings of the references cited within the specification which are hereby incorporated by reference. The embodiments within the specification provide an illustration of embodiments of the invention and should not be construed to limit the scope of the invention. The skilled artisan readily recognizes that many other embodiments are encompassed by the invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1

<211> LENGTH: 3126

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```
gaattccctc caaagcaaga caaatgactc acagagaaaa aagatggcag aaccaagggc      60
aactaaagcc gtcaggttct gaacagctgg tagatgggct ggcttactga aggacatgat      120
tcagactgtc ccggaccagc cagctcatat caaggaactc tcttgatgaa tgcagtgtgg      180
ccaaaggcgg gaagatgggtg ggcagcccag acaccgttgg gatgaactac ggcagctaca      240
tggaggagaa gcacatgcc accccaaaca tgaccacgaa cgagcgaga gttatcgtgc      300
```

-continued

cagcagatcc tacgctatgg agtacagacc atgtgcggca gtggctggag tgggcgggtga	360
aagaatatgg ccttcacagac gtcaacatct tgttattcca gaacatcgat gggaagggaac	420
tgtgcaagat gaccaaggac gacttcacaga ggctcacccc cagctacaac gccgacatcc	480
ttctctcaca tctccactac ctccagagaga ctctctctcc acatttgact tcagatgatg	540
ttgataaagc cttacaaaac tctccacggg taatgcatgc tagaaacaca gattttaccat	600
atgagccccc caggagatga gcctggaccg gtcacggcca cccacgccc cagtcgaaag	660
ctgctcaacc atctccttcc acagtgccca aaactgaaga ccagcgtcct cagttagatc	720
cttatcagat tcttggaaca acaagtagcc gccttgcaaa tccaggcagt ggccagatcc	780
agcttttgca gttcctcttg gagctcctgt cggacagctc caactccagc tgcacacct	840
gggaaggcac caacggggg ttcaagatga cggatcccga cgaggtgcc cgcgctggg	900
gagagcggaa gagcaaacc aacatgaact acgataagct cagccgcgcc ctccgttact	960
actatgacaa gaacatcatg accaaggtcc atgggaagcg ctacgcctac aagttcgact	1020
tccacgggat cgccaggcc ctccagcccc acccccggga gtcactcttg tacaagtacc	1080
cctcagacct ccgctacatg ggctcctatc acgcccaccc acagaagatg aactttgttg	1140
cgccccaccc tccagccctc ccctgacat ctccagttt ttttgcgtgc ccaaaccat	1200
actggaattc accaactggg ggtatatacc ccaacactag gctccccacc agccatatgc	1260
cttctcatct gggcacttac tactaaagac ctggcggagg cttttcccat cagcgtgcat	1320
tcaccagccc atcgccacaa actctatcgg agaacatgaa tcaaaagtgc ctcaagagga	1380
atgaaaaaag ctttactggg gctggggaag gaagccgggg aagagatcca aagactcttg	1440
ggagggagtt actgaagtct tactgaagtc ttactacaga aatgaggagg atgctaaaaa	1500
tgtcacgaat atggacatat catctgtgga ctgaccttgt aaaagacagt gtatgtagaa	1560
gcatgaagtc ttaaggacaa agtgccaaag aaagtggctc taagaaatgt ataaacttta	1620
gagtagagtt tgaatccca taatgcaaac tgggatgaaa ctaagcaat agaaacaaca	1680
cagttttgac ctaacatacc gtttataatg ccattttaag gaaaactacc tgtattttaa	1740
aatagtttca tatcaaaaa aagagaaaag acacgagaga gactgtggcc catcaacaga	1800
cgttgatatg caactgcatg gcatgtgctg ttttggttga aatcaaatc attccgtttg	1860
atggacagct gtcagcttcc tcaaaactgt aagatgaccc aaagtttcca actcctttac	1920
agtattaccg ggactatgaa ctaaaagggt ggactgagga tgtgtataga gtgagcgtgt	1980
gattgtagac agaggggtga agaaggagga ggaagggca gagaaggagg agaccaggct	2040
gggaaagaaa cttctcaagc aatgaagact ggactcagga catttgggga ctgtgtacaa	2100
tgagttatgg agactcgagg gttcatgcag tcagtgttat accaaaccca gtgttaggag	2160
aaaggacaca cggtaatgga gaaaggggag tagtagaatt cagaaacaaa aatgcgcac	2220
tctttctttg tttgtcaaat gaaaatttta actggaattg tctgatattt aagagaaaca	2280
ttcaggacct catcattatg tgggggcttt gttctccaca gggtcaggta agagatggcc	2340
ttcttggtg ccacaatcag aaatcacgca ggcattttgg tagggcgccc tccagttttc	2400
ctttgagtcg cgaacgctgt gcgtttgtca gaatgaagta tacaagtcaa tgtttttccc	2460
cctttttata taataattat ataacttatg catttataca ctacgagttg atctcggcc	2520
gccaaagaca cagacaaaa gagacaatcg atataatgtg gccttgaatt ttaactctgt	2580
atgcttaatg tttaacaatat gaagttatta gttcttagaa tgcagaatgt atgtaataaa	2640
ataagcttgg cctagcatgg caaatcagat ttatacagga gtctgcattt gcactttttt	2700

-continued

tagtgactaa agttgcttaa tgaaaacatg tgctgaatgt tgtggatttt gtgttataat	2760
ttactttgtc caggaacttg tgcaaggag agccaaggaa ataggatgtt tggcacccaa	2820
atggcgctcag cctctccagg tccttcttgc ctccccctct gtcttttatt tctagccct	2880
tttggaacag gaaggacccc ggggtttcaa ttggagcctc catatttatg cctggaagga	2940
aagaggccta tgaagctggg gttgtcattg agaaattcta gttcagcacc tggtcacaaa	3000
tcacccttaa ttctgctatg attaaaatac atttgttgaa cagtgaacaa gctaccactc	3060
gtaaggcaaa ctgtattatt actggcaaat aaagcgcat ggatagctgc aatttctcac	3120
tttaca	3126

<210> SEQ ID NO 2

<211> LENGTH: 3166

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

gtccgcgcgt gtccgcgcc gcgtgtgcc gcgcgcgtgc cttggcgtg cgcgcgcgagc	60
cgggtcgcac taactccctc ggcgcgcagc gcggcgctaa cctctcggtt attccaggat	120
ctttggagac ccgaggaaa cgtgttgac caaaagcaag acaaatgact cacagagaaa	180
aaagatggca gaaccaaggg caactaaagc cgtcagggtc tgaacagctg gtagatgggc	240
tggcttactg aaggacatga ttcagactgt ccgggaccca gcagctcata tcaaggaagc	300
cttatcagtt gtgagtggg accagtcgtt gtttgagtgt gcctacggaa cgcacacct	360
ggctaagaca gagatgaccg cgtcctcctc cagcgactat ggacagactt ccaagatgag	420
cccacgcgtc cctcagcagg attggtgtc tcaaccccca gccagggtca ccatcaaat	480
ggaatgtaac cctagccagg tgaatggctc aaggaactct cctgatgaat gcagtgtggc	540
caaaggcggg aagatgggtg gcagcccaga caccgttggg atgaactacg gcagctacat	600
ggaggagaag cacatgccac ccccaaact gaccacgaac gagcgagag ttatcgtgcc	660
agcagatcct acgctatgga gtacagacca tgtgcggcag tggctggagt ggcgggtgaa	720
agaatatggc cttccagacg tcaacatctt gttattccag aacatcgatg ggaaggaact	780
gtgcaagatg accaaggacg acttcagag gctcaccccc agctacaacg ccgacatcct	840
tctctcatc ctccactacc tcagagagac tcctcttcca catttgactt cagatgatgt	900
tgataaagcc ttacaaaact tcaccagggt aatgcatgct agaaacacag atttaccata	960
tgagcccccc aggagatcag cctggaccgg tcacggccac cccacgcccc agtcgaaagc	1020
tgctcaacca tctccttcca cagtgcctaa aactgaagac cagcgtcctc agttagatcc	1080
ttatcagatt cttggaccaa caagtagccg ccttgcaaat ccaggcagtg gccagatcca	1140
gctttggcag ttctcctggt agctcctgtc ggacagctcc aactecagct gcataccctg	1200
ggaaggcacc aacggggagt tcaagatgac ggatcccgac gaggtggccc ggcgctgggg	1260
agagcggaag agcaaaccca acatgaacta cgataagctc agccgcgccc tccgttacta	1320
ctatgacaag aacatcatga ccaagggtcc tgggaagcgc tacgcctaca agttcgactt	1380
ccacgggatc gccaggccc tcacgcccc cccccggag tcatctctgt acaagtaccc	1440
ctcagacctc ccgtacatgg gctcctatca cgcacccca cagaagatga actttgtggc	1500
gccccaccct ccagccctcc ccgtgacatc ttccagtttt ttgctgccc caaacccata	1560
ctggaattca ccaactgggg gtatatatccc caacactagg ctccccacca gccatatgcc	1620
ttctcatctg ggcacttact actaaagacc tggcgagggc ttttcccatc agcgtgcatt	1680

-continued

caccagccca tcgccacaaa ctctatcgga gaacatgaat caaaagtgcc tcaagaggaa	1740
tgaaaaaagc ttactgagg ctggggaagg aagccgggga agagatccaa agactcttgg	1800
gagggagtta ctgaagtctt actacagaaa tgaggaggat gctaaaaatg tcacgaatat	1860
ggacatatca tctgtggact gacctgttaa aagacagtgt atgtagaagc atgaagtctt	1920
aaggacaaag tgccaaagaa agtgggtctta agaaatgtat aaactttaga gtagagtttg	1980
aatcccacta atgcaaactg ggatgaaact aaagcaatag aaacaacaca gttttgacct	2040
aacataccgt ttataatgcc attttaagga aaactacctg tatttaaaaa tagtttcata	2100
tcaaaaacaa gagaaaagac acgagagaga ctgtggccca tcaacagacg ttgatatgca	2160
actgcatggc atgtgctgtt ttggttgaaa tcaatacat tccgtttgat ggacagctgt	2220
cagctttctc aaactgtgaa gatgaccaa agtttccaac tcctttacag tattaccggg	2280
actatgaact aaaaggtggg actgaggatg tgtatagagt gagcgtgtga ttgtagacag	2340
aggggtgaag aaggaggagg aagaggcaga gaaggaggag accaggctgg gaaagaaact	2400
tctcaagcaa tgaagactgg actcaggaca ttgggggact gtgtacaatg agttatggag	2460
actcgagggt tcatgcagtc agtggtatac caaacccagt gttaggagaa aggacacagc	2520
gtaatggaga aagggaaagta gtagaattca gaaacaaaaa tgcgcattctc tttctttgtt	2580
tgtcaaatga aaattttaac tggaattgtc tgatatattaa gagaaacatt caggacctca	2640
tcattatgtg ggggctttgt tctccacagg gtcaggtaag agatggcctt cttggctgcc	2700
acaatcagaa atcacgcagg cattttgggt aggcggcctc cagttttcct ttgagtgcg	2760
aacgctgtgc gtttgcaga atgaagtata caagtcaatg tttttcccc tttttatata	2820
ataattatat aacttatgca ttatatact acgagttgat ctggccagc caaagacaca	2880
cgacaaaaga gacaatcgat ataattgtggc cttgaatttt aactctgtat gcttaattgt	2940
tacaatatga agttattagt tcttagaatg cagaatgtat gtaataaaat aagcttggcc	3000
tagcatggca aatcagattt atacaggagt ctgcatttgc acttttttta gtgactaaag	3060
ttgcttaatg aaaacatgtg ctgaatgttg tggattttgt gttataattt actttgtcca	3120
ggaacttggt caagggagag ccaaggaaat aggatgtttg gcaccc	3166

<210> SEQ ID NO 3

<211> LENGTH: 2534

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

gggattggga gggcttcttg caggtctgtg ggctggggct aagggtctgt cagtttcctt	60
cagcggggca ctgggaagcg ccatggcact gcagggcac tcggtcgtgg agctgtccgg	120
cctggccccg ggcccgttct tgctataggt cctggctgac ttcggggcgc gtgtggtacg	180
cgtggaccgg cccggctccc gctacgacgt gagccgcttg ggccggggca agcgtctgct	240
agtcttggtg ctgaagcagc cgcggggagc cgcctgtctg cggcgtctgt gcaagcggtc	300
ggatgtgctg ctggagccct tccgcccggt tgctcatggg aaactccagc tgggccccaga	360
gattctgcag cgggaaaatc caaggcttat ttatgccagg ctgagtggat ttggccagtc	420
aggaagcttc tgccgggttag ctggccacga tatcaactat ttggctttgt cagggtgtct	480
ctcaaaaatt ggcagaagtg gtgagaatcc gtatgccccg ctgaatctcc tggctgactt	540
tgctgggtgt ggcccttatgt gtgcactggg cattataatg gctctttttg accgcacag	600
cactggcaag ggtcagggtc ttgatgcaaa tatggtggaa ggaacagcat atttaagttc	660

-continued

ttttctgtgg aaaactcaga aattgagtct gtgggaagca cctcgaggac agaacatgtt	720
ggatggtgga gcacctttct atacgactta caggacagca gatggggaat tcatggctgt	780
tggagcaata gaacccagct tctacgagct gctgatcaaa ggacttggac taaagtctga	840
tgaacttccc aatcagatga gcatggatga ttggccagaa atgaagaaga agtttgcaga	900
tgtatttgca gagaagacga aggagagtg gtgtcaaato tttgacggca cagatgcctg	960
tgtgactccg gttctgactt ttgaggaggt tgttcacat gatcacaaca aggaacgggg	1020
ctcgtttata accagtgagg agcaggagct gagccccgc cctgcacctc tgetgtttaa	1080
cacccagcc atcccttctt tcaaaagga tcctttcata ggagaacaca ctgaggagat	1140
acttgaagaa ttggattca gccgcgaaga gatttatcag cttactcag ataaaatcat	1200
tgaagtaat aaggtaaaag ctagtctcta acttcaggc ccacggctca agtgaatttg	1260
aatactgcat ttacagtga gagtaacaca taacattgta tgcattgaaa catggaggaa	1320
cagtattaca gtgtcctacc actctaatac agaaaagaat tacagactct gattctacag	1380
tgatgattga attctaaaaa tggttatcat tagggctttt gatttataaa actttgggta	1440
cttatactaa attatggtag ttattctgcc ttccagtttg cttgatata ttgttgatat	1500
taagattctt gacttatatt ttgaatgggt tctagtgaag aaggaatgat atattcttga	1560
agacatcgat atacatttat ttacactctt gattctacaa ttagagaaaat gaggaatgc	1620
cacaaattgt atgggtataa aagtcacgtg aaacagagtg attggttgca tccaggcctt	1680
ttgtcttggt gttcatgac tcctctaaag cacattccaa actttagcaa cagttatcac	1740
actttgtaat ttgcaaagaa aagtttcacc tgtattgaat cagaatgcct tcaactgaaa	1800
aaaacatata caaaataatg aggaatgtg ttggctcact acgtagagtc cagagggaca	1860
gtcagtttta gggttgcctg tatccagtaa ctggggcct gtttcccgt ggtctctctg	1920
gctgtcagct ttcccttctc catgtgtttg atttctctc aggtctgtag caagttcttg	1980
atcttatacc caacacacag caacatccag aaataaagat ctcaggaccc cccagcaagt	2040
cgttttgtgt ctccctggac tgagttaagt tacaagcctt tcttatacct gtctttgaca	2100
aagaagacgg gattgtcttt acataaaaac agcctgctcc tggagcttcc ctggactcaa	2160
cttcctaaag gcatgtgagg aaggggtaga ttccacaato taatccgggt gccatcagag	2220
tagaggagat agagaatgga tgttgggtag gccatcaata aggtccattc tgcgcagtat	2280
ctcaactgcc gttcaacaat cgcaagagga aggtggagca ggtttcttca tcttacagtt	2340
gagaaaacag agactcagaa gggcttctta gttcatgttt cccttagcgc ctcaagtatt	2400
ttttcatggt ggcttaggcc aaaagaaata tctaaccatt caatttataa ataattaggt	2460
ccccaacgaa ttaaatatta tgtcctacca acttattagc tgcttgaaaa atataatata	2520
cataaataaa aaaa	2534

<210> SEQ ID NO 4

<211> LENGTH: 3923

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

acagaagaaa tagcaagtgc cgagaagctg gcatcagaaa aacagagggg agatttgtgt	60
ggctgcagcc gagggagacc aggaagatct gcatggtggg aaggacctga tgatacagag	120
gaattacaac acatatactt agtgtttcaa tgaacaccaa gataaataag tgaagagcta	180
gtccgctgtg agtctctca gtgacacagg gctggatcac catcgacggc actttctgag	240

-continued

tactcagtgc agcaaaagaaa gactacagac atctcaatgg caggggtgag aaataagaaa	300
ggctgctgac ttaccatct gaggccacac atctgctgaa atggagataa ttaacatcac	360
tagaaacagc aagatgacaa tataatgtct aagtagtgac atgtttttgc acatttcag	420
cccctttaa tatccacaca cacaggaagc acaaaaggaa gcacagagat ccctgggaga	480
aatgccggc cgcacatctg ggtcatcgat gagcctcgcc ctgtgcctgg tcccgcttgt	540
gaggaagga cattagaaaa tgaattgatg tgttccttaa aggatgggca ggaaaacaga	600
tcctgttgat gatatttatt tgaacgggat tacagatttg aaatgaagtc acaaatgag	660
cattaccaat gagaggaaaa cagacgagaa aatcttgatg gcttcacaag acatgcaaca	720
aacaaatgg aatactgtga tgacatgagg cagccaagct ggggaggaga taaccacggg	780
gcagagggc aggattctgg ccctgctgcc taaactgtgc gttcataacc aaatcatttc	840
atatttctaa cctcaaaac aaagctgttg taatatctga tctctacggt tccttctggg	900
cccaacattc tccatatatc cagccacact catttttaat atttagttcc cagatctgta	960
ctgtgacctt tctacactgt agaataacat tactcatttt gttcaaagac ccttcgtgtt	1020
gctgccta atgtagctga ctgtttttcc taaggagtgt tctggccag gggatctgtg	1080
aacaggctgg gaagcatctc aagatcttcc cagggttata cttactagca cacagcatga	1140
tcattacgga gtgaattatc taatcaacat catcctcagt gtctttgccc atactgaaat	1200
tcatttccca cttttgtgcc cattctcaag acctcaaat gtcattccat taatatcaca	1260
ggattaactt ttttttttaa cctggaagaa ttcaatgtta catgcageta tgggaattta	1320
attacatatt ttgttttcca gtgcaaagat gactaagtcc tttatccctc ccctttgttt	1380
gatttttttt ccagtataaa gttaaaatgc ttagccttgt actgaggctg tatacagcac	1440
agcctctccc catccctcca gccttatctg tcatcccat caaccctcc cataccacct	1500
aaacaaaatc taacttgtaa ttccttgaac atgtcaggac atacattatt ccttctgcct	1560
gagaagctct tcctgtctc ttaaatctag aatgatgtaa agttttgaat aagttgacta	1620
tcttacttca tgcaagaag ggacacatat gagattcatc atcacatgag acagcaaata	1680
ctaaaagtgt aatttgatta taagagtta gataaatata tgaaatgcaa gagccacaga	1740
gggaatgttt atggggcacg tttgtaagcc tgggatgtga agcaaaggca gggaaacctca	1800
tagtatctta tataatatat ttcatttctc tatctctatc acaatatcca acaagctttt	1860
cacagaattc atgcagtga aatcccaaaa ggtaaccttt atccatttca tggtgagtgc	1920
gctttagaat tttggcaaat catactggc acttatctca actttgagat gtgtttgtcc	1980
ttgtagttaa ttgaaagaa tagggcactc ttgtgagcca ctttaggggt cactcctggc	2040
aataaagaat ttacaaagag ctactcagga ccagtgttta agagctctgt gtgtgtgtgt	2100
gtgtgtgtgt gagtgtacat gccaaagtgt gcctctctct cttgacctat tatttcagac	2160
ttaaaacaag catgttttca aatggcacta tgagctgcca atgatgtatc accaccatat	2220
ctcattattc tccagtaaat gtgataataa tgtcatctgt taacataaaa aaagtttgac	2280
ttcacaaaag cagctggaaa tggacaacca caatatgcat aaatctaact cctaccatca	2340
gctacacact gcttgacata tattgttaga agcacctcgc atttgtgggt tctcttaagc	2400
aaaatacttg cattaggtct cagctggggc tgtgcatcag gcggtttgag aaatattcaa	2460
ttctcagcag aagccagaat ttgaattccc tcatctttta ggaatcattt accaggtttg	2520
gagaggattc agacagctca ggtgctttca ctaatgtctc tgaacttctg tccctctttg	2580
tgttcattga tagtccaata aataatgtta tctttgaact gatgctcata ggagagaata	2640

-continued

taagaactct gagtgatata aacattaggg attcaaagaa atattagatt taagctcaca	2700
ctggtcaaaa ggaaccaaga tacaagaac tctgagctgt catcgcccc atctctgtga	2760
gccacaacca acagcaggac ccaacgcatg tctgagatcc ttaaatcaag gaaaccagtg	2820
tcatgagttg aattctccta ttatggatgc tagcttctgg ccatctctgg ctctcctctt	2880
gacacatatt agcttctaga ctttgcctcc acgactttta tcttttctcc aacacatcgc	2940
ttaccaatcc tctctctgct ctgttgcttt ggacttcccc acaagaattt caacgactct	3000
caagtctttt cttccatccc caccactaac ctgaatgcct agacccttat ttttattaat	3060
ttccaataga tgctgcttat gggctatatt gcttttagatg aacattagat atttaaagct	3120
caagaggttc aaaatccaac tcattatctt ctctttcttt cactccctg ctctctctcc	3180
tatattactg attgcaactg acagcatggg ccccaatgta gccatgcaa tgagaaaccc	3240
agtggctcct tgtggtacat gcatgcaaga ctgctgaagc cagaaggatg actgattacg	3300
cctcatgggt ggaggggacc actcctgggc cttcgtgatt gtcaggagca agacctgaga	3360
tgctccctgc cttcagtgtc ctctgcatct cccctttcta atgaagatcc atagaatttg	3420
ctacatttga gaattccaat taggaactca catgttttat ctgccctatc aatttttta	3480
acttgctgaa aattaagttt tttcaaaatc tgccttgta aattactttt tcttacagtg	3540
tcttggcata ctatatcaac tttgattctt tgttacaact tttcttactc ttttatcacc	3600
aaagtggctt ttattctctt tattattatt attttcttt actactatat tacgttggtta	3660
ttattttggt ctctatagta tcaatttatt tgatttagtt tcaatttatt tttattgctg	3720
acttttaaaa taagtgatcc ggggggtggg agaacagggg agggagagca ttaggacaaa	3780
tacctaatgc atgtgggact taaaacctag atgatgggtt gatagggtgca gcaaacact	3840
atggcacacg tatacctgtg taacaaacct acacattctg cacatgtatc ccagaacgta	3900
aagtaaaatt taataaaaag tga	3923

<210> SEQ ID NO 5

<211> LENGTH: 2390

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

agagccttcg tttgccaagt cgctccaga ccgcagacat gaaacttgtc ttctctgccc	60
tgctgttccg cggggccctc ggactgtgtc tggctggcgg taggaggagt gttcagtgg	120
gcgcctgata ccaaccgag gccacaaaat gcttccaatg gcaaaggaat atgagaaaag	180
tgctgggccc tctgtcagc tgcataaaga gagactcccc catccagtgt atccaggcca	240
ttgcggaaaa cagggccgat gctgtgaccc ttgatgggtg tttcatatac gaggcaggcc	300
tggcccccta caaactgca cctgtagcgg cggaagtcta cgggaccgaa agacagccac	360
gaactcacta ttatgccgtg gctgtggtga agaagggcgg cagctttcag ctgaacgaac	420
tgcaaggtct gaagtccgc cacacaggcc ttgcaggac cgctggatgg aatgtcccta	480
tagggacact tcgtccattc ttgaattgga cgggtccacc tgagccatt gaggcagctg	540
tggccagggt cttctcagcc agctgtgttc ccggtgcaga taaaggacag tcccccaacc	600
tgtgtgcctt gtgtgcgggg acaggggaaa acaaatgtgc cttctctccc caggaaccgt	660
acttcagcta ctctggtgcc ttcaagtgtc tgagagacgg ggctggagac gtggctttta	720
tcagagagag cacagtgttt gaggacctgt cagacaggcc tgaaagggac gagtatgagt	780

-continued

tactctgccc agacaacact cggaagccag tggacaagtt caaagactgc catctggccc	840
gggtcccttc tcatgccgtt gtggcacgaa gtgtgaatgg caaggaggat gccatctgga	900
atcttctccg ccaggcacag gaaaagtttg gaaaggacaa gtcaccgaaa ttccagctct	960
ttggctcccc tagtgggcag aaagatctgc tgttcaagga ctctgccatt gggttttcga	1020
gggtgcccc gaggatagat tctgggctgt accttggtctc cggtacttct actgccatcc	1080
agaacttgag gaaaagttag gaggaagtgg ctgcccggcg tgcgcgggtc gtgtggtgtg	1140
cgggtggcga gcaggagctg cgcaagtgtg accagtggag tggcttgagc gaaggcagcg	1200
tgacctgtct ctgcgcctcc accacagagg actgcatcgc cctgggtgtg aaaggagaag	1260
ctgatgccat gagtttgat ggaggatatg tgtacctgc aggc aaatgt ggtttggtgc	1320
ctgtcctggc agagaactac aaatcccaac aaagcagtga cctgatcct aactgtgtgg	1380
atagacctgt ggaaggatat cttgctgtgg cgggtggttag gagatcagac actagcctta	1440
cctggaactc tgtgaaagcg aagaagtctc gccacaccgc cgtggacagg actgcaggct	1500
ggaatatccc catgggctct ctcttcaacc agacgggctc ctgcaaattt gatgaatatt	1560
tcagtcaaag ctgtgcccc gggctctgacc cgagatctaa tctctgtgct ctgtgtattg	1620
gcgacgagca gggtgagaat aagtgcgtgc ccaacagcaa cgagagatac tacggctaca	1680
ctggggcttt ccgggtcctg gctgagaatg ctggagacgt tgcatttgtg aaagatgtca	1740
ctgtcttgca gaacctgat ggaaataaca atgaggcacg ggctaaggat ttgaagctgg	1800
cagactttgc gctgctgtgc ctcgatggca aacggaagcc tgtgactgag gctagaagct	1860
gccatcttgc catggccccg aatcatgccg tgggtgtctc gatggataag gtggaacgcc	1920
tgaaacaggt gttgtccac caacaggcta aatttgggag aaatggatct gactgcccgg	1980
acaagttttg cttattccag tctgaaacca aaaaccttct gttcaatgac aacctgagt	2040
gtctggccag actccatggc aaaacaacat atgaaaaata tttgggacca cagtatgtcg	2100
caggcattac taatctgaaa aagtgtctaa cctccccct cctggaagcc tgtgaattcc	2160
tcaggaagta aaaccgaaga agatggccca gctccccaag aaagcctcag ccattcactg	2220
ccccagctc ttctccccag gtgtgttggg gccttggcct cccctgctga aggtggggat	2280
tgcccacca tctgcttaca attcctgct gtcgtcttag caagaagtaa aatgagaaat	2340
ttgttgata ttctctcctt aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	2390

<210> SEQ ID NO 6

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 6

gtcaagatgc taccgttcag

20

<210> SEQ ID NO 7

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

agagaaacat tcaggacctc atcattatg

29

-continued

<210> SEQ ID NO 8
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

gcagccaaga aggccatct 19

<210> SEQ ID NO 9
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

ttgttctcca cagggt 16

<210> SEQ ID NO 10
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

caggtccttc ttgcctccc 19

<210> SEQ ID NO 11
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

tatggaggct ccaattgaaa cc 22

<210> SEQ ID NO 12
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

tgtcttttat ttctagcccc ttttgaaca gga 33

<210> SEQ ID NO 13
<211> LENGTH: 50
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

Cys Lys Ala Leu Gln Asn Ser Pro Arg Leu Met His Ala Arg Asn Thr
1 5 10 15

Asp Leu Pro Tyr Glu Pro Pro Arg Arg Ser Ala Trp Thr Gly His Gly
20 25 30

His Pro Thr Pro Gln Ser Lys Ala Ala Gln Pro Ser Pro Ser Thr Val
35 40 45

Pro Lys
50

<210> SEQ ID NO 14
<211> LENGTH: 49
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 14

Cys Asp Phe His Gly Ile Ala Gln Ala Leu Gln Pro His Pro Pro Glu
 1 5 10 15

Ser Ser Leu Tyr Lys Tyr Pro Ser Asp Leu Pro Tyr Met Gly Ser Tyr
 20 25 30

His Ala His Pro Gln Lys Met Asn Phe Val Ala Pro His Pro Pro Ala
 35 40 45

Leu

<210> SEQ ID NO 15

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

cacatttcca gccctttaa ata 23

<210> SEQ ID NO 16

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

gggcgaggct catcgat 17

<210> SEQ ID NO 17

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

ggaagcacag agatccctgg gagaaatg 28

What is claimed is:

1. A method of diagnosing prostate cancer in a subject, 40 comprising:

a) measuring the expression level of a human ERG polypeptide in a biological sample from the subject, wherein the biological sample comprises prostate epithelial cells and wherein the expression level of the human ERG polypeptide is measured by an immuno- 45 assay using a monoclonal antibody that binds to the human ERG1 polypeptide and does not bind to the human ETS domain; and

b) correlating the expression level of the human ERG polypeptide with the presence of prostate cancer in the subject, wherein increased expression of the human ERG polypeptide in the biological sample as compared to the expression of the human ERG polypeptide in a control sample indicates the presence of prostate cancer 55 in the subject.

2. The method of claim 1, wherein the monoclonal antibody is generated using a fragment of the human ERG1 polypeptide as an immunogen.

3. The method of claim 1, wherein the monoclonal antibody binds to the human ERG1, ERG2, and ERG3 polypeptides. 60

4. A method of diagnosing prostate cancer in a subject, comprising:

a) measuring the expression level of a human ERG polypeptide in a biological sample from the subject, wherein the biological sample comprises prostate epi- 65

thelial cells and wherein the expression level of the human ERG polypeptide is measured by an immuno-assay using a monoclonal antibody that binds to the human ERG1 polypeptide; and

b) correlating the expression level of the human ERG polypeptide with the presence of prostate cancer in the subject, wherein increased expression of the human ERG polypeptide in the biological sample as compared to the expression of the human ERG polypeptide in a control sample indicates the presence of prostate cancer in the subject, wherein the monoclonal antibody is generated using a fragment of the human ERG1 polypeptide as an immunogen, and wherein the monoclonal antibody binds to an epitope formed by amino acids 275-322 of the polypeptide encoded by a nucleic acid comprising the sequence of SEQ ID NO: 1.

5. The method of claim 1, wherein the monoclonal antibody binds to an epitope formed by amino acids 118-166 of the polypeptide encoded by a nucleic acid comprising the sequence of SEQ ID NO:1.

6. The method of claim 1, wherein the biological sample is chosen from a tissue sample, a blood sample, or a urine sample.

7. The method of claim 6, wherein the biological sample is a prostate tissue sample.

8. The method of claim 1, further comprising measuring the expression levels of a human AMACR polypeptide and correlating the expression levels of the human ERG poly-

peptide and the human AMACR polypeptide with the presence of prostate cancer in the subject.

9. The method of claim 8, wherein the human AMACR polypeptide is encoded by a nucleic acid comprising the sequence of SEQ ID NO. 3.

10. The method of claim 1, wherein the control sample is a noncancerous biological sample from the subject.

11. The method of claim 8, wherein the expression levels of the human ERG polypeptide and the human AMACR polypeptide in the biological sample are compared to the expression levels of the human ERG polypeptide and the human AMACR polypeptide in a control sample and wherein increased expression of the human ERG polypeptide and the human AMACR polypeptide in the biological sample as compared to the expression of the ERG polypeptide and the human AMACR polypeptide in the control sample indicates the presence of prostate cancer in the subject.

12. The method of claim 1, wherein the immunoassay is an immunohistochemistry assay.

13. A method of identifying prostate cancer in a subject, comprising:

- a) providing a prostate tissue sample from the subject;
- b) contacting the prostate tissue sample with a monoclonal antibody that binds to a human ERG polypeptide and does not bind to the human ETS domain; and
- c) measuring the level of expression of the human ERG polypeptide in the prostate tissue sample by detecting the amount of binding between the monoclonal antibody and the human ERG polypeptide in the prostate tissue sample;
- d) comparing the level of expression of the ERG polypeptide in the prostate tissue sample to the level of expression of the ERG polypeptide in a control sample; and
- e) identifying prostate cancer in the prostate tissue sample from the subject if the level of expression of the ERG polypeptide in the prostate tissue sample is increased as compared to the expression level of the ERG polypeptide in the control sample.

14. The method of claim 13, wherein the monoclonal antibody binds to the ERG1, ERG2, and ERG3 polypeptides.

15. A method of identifying prostate cancer in a subject, comprising:

- a) providing a prostate tissue sample from the subject;
- b) contacting the prostate tissue sample with a monoclonal antibody that binds to a human ERG polypeptide; and
- c) measuring the level of expression of the human ERG polypeptide in the prostate tissue sample by detecting the amount of binding between the monoclonal antibody and the human ERG polypeptide in the prostate tissue sample;
- d) comparing the level of expression of the ERG polypeptide in the prostate tissue sample to the level of expression of the ERG polypeptide in a control sample; and
- e) identifying prostate cancer in the prostate tissue sample from the subject if the level of expression of the ERG polypeptide in the prostate tissue sample is increased as

compared to the expression level of the ERG polypeptide in the control sample, wherein the monoclonal antibody binds to an epitope formed by amino acids 275-322 of the polypeptide encoded by a nucleic acid comprising the sequence of SEQ ID NO:1.

16. The antibody of claim 13, wherein the monoclonal antibody binds to an epitope formed by the amino acid sequence of SEQ ID NO:14.

17. The method of claim 13, wherein the control sample is a noncancerous biological sample from the subject.

18. The method of claim 17, wherein the noncancerous biological sample from the subject is normal prostate tissue comprising benign prostate epithelial cells.

19. A method of detecting the expression of a human ERG polypeptide in a biological sample, wherein the biological sample comprises human prostate epithelial cells, the method comprising:

- a) contacting the human prostate epithelial cells with a monoclonal antibody that binds to a human ERG1, ERG2, and ERG3 polypeptide, wherein the monoclonal antibody binds to an epitope formed by amino acids 275-322 of the polypeptide encoded by a nucleic acid comprising the sequence of SEQ ID NO:1 or to an epitope formed by the amino acid sequence of SEQ ID NO:14; and

- b) measuring the level of expression of the human ERG polypeptide in the human prostate epithelial cells sample by detecting the amount of binding between the monoclonal antibody and the human ERG polypeptide in the human prostate epithelial cells.

20. The method of claim 19, further comprising a step of comparing the level of expression of the human ERG polypeptide in the human prostate epithelial cells to the level of expression of the human ERG polypeptide in a control sample.

21. The method of claim 19, wherein the biological sample is obtained from a patient that has prostate cancer or is suspected of having prostate cancer.

22. The method of claim 21, wherein the patient has been treated by radical prostate surgery.

23. The method of claim 19, further comprising detecting the expression of a human AMACR polypeptide in the biological sample.

24. The method of claim 1, wherein the expression level of the ERG polypeptide is correlated with longer PSA recurrence free survival, well and moderate tumor differentiation, a pathologic T stage of pT2 or lower, or a negative surgical status.

25. The method of claim 1, wherein the monoclonal antibody does not bind to a SMAP-PNT domain.

26. The method of claim 13, wherein the monoclonal antibody does not bind to a SMAP-PNT domain.

27. The method of claim 1, wherein the monoclonal antibody specifically binds to ERG1, ERG2, and ERG3, but does not specifically bind to another protein of the ETS family.

28. The method of claim 13, wherein the monoclonal antibody specifically binds to ERG1, ERG2, and ERG3, but does not specifically bind to another protein of the ETS family.

专利名称(译)	使用erg基因单独或与前列腺癌中其他过表达或低表达基因组合诊断或治疗前列腺癌的方法		
公开(公告)号	US9464325	公开(公告)日	2016-10-11
申请号	US13/534529	申请日	2012-06-27
[标]申请(专利权)人(译)	HENRY中号JACKSON发现的军事医学的进步		
申请(专利权)人(译)	亨利M.杰克逊基金会为军事医学，INC进步.		
当前申请(专利权)人(译)	亨利M.杰克逊基金会为军事医学，INC进步.		
[标]发明人	PETROVICS GYORGY SRIVASTAVA SHIV		
发明人	PETROVICS, GYORGY SRIVASTAVA, SHIV		
IPC分类号	G01N33/53 C12Q1/68 C07K16/30		
CPC分类号	C12Q1/6886 C07K16/30 C12Q2600/112 C12Q2600/118 C12Q2600/158 C07K16/32 C07K2317/33 H05K999/99		
优先权	60/568822 2004-05-07 US PCT/US2005/015926 2005-05-06 WO 60/622021 2004-10-27 US		
其他公开文献	US20120295282A1		
外部链接	Espacenet USPTO		

摘要(译)

本发明涉及癌基因或肿瘤抑制基因，以及涉及前列腺癌及其表达产物的其他基因，以及其衍生物和类似物。提供了治疗组合物和检测和治疗癌症的方法，包括前列腺癌和其他相关癌症。还提供了通过测定至少一种前列腺癌细胞特异性基因的表达水平来诊断和/或预测前列腺癌的方法，所述基因包括例如单独的ERG基因或LTF基因，或与至少一种组合。AMACR基因和DD3基因。




FIG. 1A




FIG. 1B