



US009720000B2

(12) **United States Patent**  
**Ponniah et al.**

(10) **Patent No.:** **US 9,720,000 B2**  
(45) **Date of Patent:** **Aug. 1, 2017**

(54) **TARGETED IDENTIFICATION OF IMMUNOGENIC PEPTIDES**

(56) **References Cited**

- (71) Applicant: **The Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc.**, Bethesda, MD (US)
- (72) Inventors: **Sathibalan Ponniah**, Columbia, MD (US); **George E Peoples**, San Antonio, TX (US); **Catherine E Storrer**, Columbia, MD (US); **Michael Flora**, Mt. Airy, MD (US)
- (73) Assignee: **The Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc.**, Bethesda, MD (US)
- (\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

U.S. PATENT DOCUMENTS

4,554,101 A	11/1985	Hopp	
5,550,214 A	8/1996	Eberlein et al.	
5,726,023 A	3/1998	Cheever et al.	
6,780,598 B1	8/2004	Kokulus	
6,861,234 B1 *	3/2005	Simard	A61K 39/0011 435/23
7,179,645 B2	2/2007	Humphreys et al.	
7,211,432 B2	5/2007	Schlom et al.	
7,446,185 B2	11/2008	Nelson	
8,883,164 B2	11/2014	Ponniah et al.	
8,945,573 B2	2/2015	Ponniah et al.	
9,050,322 B2	6/2015	Ponniah et al.	
2004/0018971 A1 *	1/2004	Fikes	C07K 14/71 424/185.1
2005/0119288 A1	6/2005	Bhattacharya et al.	
2010/0209443 A1	8/2010	Peoples et al.	
2011/0256164 A1	10/2011	Peoples	

FOREIGN PATENT DOCUMENTS

WO	94/20127	9/1994
WO	2004053139 A1	6/2004
WO	2005007694	1/2005
WO	2005/076975	8/2005
WO	2008150577	12/2008

(21) Appl. No.: **14/713,721**

(22) Filed: **May 15, 2015**

(65) **Prior Publication Data**

US 2015/0260727 A1 Sep. 17, 2015

**Related U.S. Application Data**

(63) Continuation of application No. 13/774,829, filed on Feb. 22, 2013, now Pat. No. 9,050,322, which is a continuation of application No. 12/045,402, filed on Mar. 10, 2008, now Pat. No. 8,945,573, which is a continuation of application No. PCT/US2006/035171, filed on Sep. 8, 2006.

(60) Provisional application No. 60/714,865, filed on Sep. 8, 2005.

(51) **Int. Cl.**

**G01N 33/53** (2006.01)  
**G01N 33/68** (2006.01)  
**A61K 39/00** (2006.01)  
**G01N 33/50** (2006.01)  
**G01N 33/564** (2006.01)  
**G01N 33/574** (2006.01)  
**A61K 39/395** (2006.01)  
**C07K 16/32** (2006.01)  
**C40B 30/02** (2006.01)  
**G01N 33/577** (2006.01)

(52) **U.S. Cl.**

CPC ..... **G01N 33/6803** (2013.01); **A61K 39/0011** (2013.01); **A61K 39/39558** (2013.01); **C07K 16/32** (2013.01); **C40B 30/02** (2013.01); **G01N 33/5011** (2013.01); **G01N 33/564** (2013.01); **G01N 33/574** (2013.01); **G01N 33/577** (2013.01); **G01N 33/6845** (2013.01); **G01N 2500/00** (2013.01); **G01N 2500/04** (2013.01)

(58) **Field of Classification Search**

None  
See application file for complete search history.

OTHER PUBLICATIONS

Jasinska et al (Int. J. Cancer, 2003, 107:976-983).\*

Ayyoub et al (Journal of Immunology, 2002, 168:1717-1722).\*

Ishihara et al (International Journal of Oncology, Apr. 2004, 24:967-975).\*

Parker et al (Molecular Biotechnology, 2002, 49-62).\*

Papac et al (Protein Science, 1994, 3:1485-1492).\*

Suckau et al (PNAS, 1990, 87:9848-9852).\*

English Translation of Office Action dated Jun. 20, 2012 from Japanese Patent Application No. 2008-530244, 2 pages.

Kono, Koji et al. Trastuzumab (Herceptin) Enhances Class 1-Restricted Antigen Presentation Recognized by HER-2. neu-Specific T Cytotoxic Lymphocytes. Clinical Cancer Research, Apr. 1, 2004, vol. 10, 2538-2544.

zum Buschenfelde et al. (Cancer Research, 2002, 62:2244-2247).

Mittendorf et al. (Proc. Amer. Assoc. Cancer Research, Apr. 2005, 46; abstract#3449).

Fisk, Bryan et al. Identification of an immunodominant Peptide of HER-2/neu Protooncogene Recognized by Ovarian Tumor-specific Cytotoxic T Lymphocytes Lines. J. Exp. Med., vol. 181, Jun. 1995, pp. 2109-2117.

(Continued)

Primary Examiner — Laura B Goddard

(74) Attorney, Agent, or Firm — MH2 Technology Law Group, LLP

(57) **ABSTRACT**

This invention relates generally to identifying peptide sequences involved in antibody binding to any protein for synthesis of vaccine treatments. This novel method allows for a more manageable vaccine peptide discovery and specific generation of unique immunogenic peptides from self-tumor associated proteins and/or foreign proteins from infectious organisms for specific and/or enhanced expression only in the presence of the antibody.

**7 Claims, 8 Drawing Sheets**

(56)

## References Cited

## OTHER PUBLICATIONS

Yao et al. Identification of Parathyroid Hormone-related Protein-derived Peptides Immunogenic in HLA-A2 Positive Prostate Gland Cancer Patients. Proceedings of 63th Annual Meeting of the Japanese Cancer Association, 2004, p. 295, P-0735 (with English translation).

Hird et al. Journal of Surgical Research, vol. 144, No. 2 (Feb. 2008), pp. 180-181.

Webster et al. Breast Cancer Research and Treatment, vol. 100, Supp. 1 (2006), pp. S79-S80.

Mittendorf et al. Journal of the American College of Surgeons, vol. 203, No. 3, (Sep. 2006), p. S79.

Mittendorf et al. Annals of Surgical Oncology, vol. 13, No. 8 (Aug. 2006), pp. 1085-1098.

Pero et al. British Journal of Cancer, vol. 96, No. 10 (May 2007), pp. 1520-1525.

Knutson et al. The Journal of Clinical Investigation, vol. 107, No. 4 (Feb. 2001), pp. 477-484.

Tanaka et al. International Journal of Cancer, vol. 94, No. 4 (Nov. 2001), pp. 540-544.

Mittendorf et al. Breast Diseases: A Year Book Quarterly, vol. 17, No. 4 (Jan. 2007), pp. 318-320.

International Search Report mailed Oct. 17, 2008 in PCT/GB2008/050227.

Office Action dated Jul. 16, 2014 from Canadian Patent Application No. 2,812,110, pp. 1-2.

Communication dated Apr. 11, 2014 from European Patent Application No. 06824918.4, pp. 1-14.

Knutson, Keith L. et al. Immunization of Cancer Patients with a HER-2/neu, HLA-A2 Peptide, p369-377, Results in Short-lived Peptide-specific Immunity. Clinical Cancer Research, 2002, 8:1014-1018.

Knutson, K. L. et al. Tumor antigen-specific T helper cells in cancer immunity and immunotherapy. Cancer Immunology Immunotherapy, Aug. 1, 2005, vol. 54, No. 8, pp. 721-728.

Mittelman, Abraham et al. Monoclonal and Polyclonal Humoral Immune Response to EC HER-2/neu Peptides with Low Similarity to the Host's Proteome. International Journal of Cancer, Apr. 10, 2002, vol. 98, No. 5, pp. 741-747.

Dakappagari, Naveen K. et al. Prevention of Mammary Tumors with a Chimeric HER-2 B-cell Epitope Peptide Vaccine. Cancer Research, Jul. 15, 2000, vol. 60, No. 14, pp. 3782-3789.

Ramasamy, R. et al. Characterisation of an inhibitory monoclonal antibody-defined epitope on a malaria vaccine candidate antigen. Immunology Letters, Feb. 1, 1990, vol. 23, No. 4, pp. 305-309.

Wagner, C. et al. Identification of an HLA-A\*02 restricted immunogenic peptide derived from the cancer testis antigen HOM-MEL-40/SSX2. Cancer Immunity, Academy of Cancer Immunology, Dec. 1, 2003, vol. 3, No. 18, pp. 1-15.

Non-Final Office Action dated Jan. 31, 2014 from U.S. Appl. No. 13/336,068, pp. 1-14.

Murray, James L. et al. Toxicity, Immunogenicity, and Induction of E75-specific Tumor-lytic CTLs by HER-2 Peptide E75 (369-377) Combines with Granulocyte Macrophage Colony-stimulating Factor in HLA-A2+ Patients with Metastatic Breast and Ovarian Cancer. Clinical Cancer Research, Nov. 2002, vol. 8, pp. 3407-3418.

Non-Final Office Action dated Apr. 21, 2014 from U.S. Appl. No. 12/045,402, pp. 1-18.

Emens, Leisha A. et al. Augmenting the Potency of Breast Cancer Vaccines: Combined Modality Immunotherapy. Breast Disease, 2004, vol. 20, pp. 13-24.

Baxevasis, Constatin N. et al. Immunobiology of HER-2/neu oncoprotein and its potential application in cancer immunotherapy. Cancer Immunol Immunother, 2004, vol. 53, pp. 166-175.

Communication dated May 21, 2013 from Canadian Application No. 2,622,036, pp. 1-4.

Zimmerman et al. (Drug Discovery Today, Jan. 2007, 12: 34-42).

Emens et al. (Endocrine Related Cancer, Mar. 2005, 12:1-7).

Cruz et al. (Vaccine, 2003, 21:1317-1326).

Communication dated Jan. 19, 2015 from European Patent Application No. 06824918.4, pp. 1-8.

Dakappagari, Naveen K. et al., "Conformational HER-2/neu B-cell Epitope Peptide Vaccine Designed to Incorporate Two Native Disulfide Bonds Enhances Tumor Cell Binding and Antitumor Activities", The Journal of Biological Chemistry, Jan. 7, 2005, vol. 280 No. 1, pp. 54-63.

Disis et al., "Peptide-Based, but Not Whole Protein, Vaccines Elicit Immunity to HER-2/neu, an Oncogenic Self Protein", The Journal of Immunology, May 1, 1996, vol. 156, No. 9, 9 pages.

English Translation of Office Action Dated Sep. 29, 2015 from Japanese Patent Application No. 2014-136512, 3 pages.

European Search Report and Written Opinion dated May 17, 2017 for European Application No. 17159896.4, 9 pages.

\* cited by examiner

Figure 1




FIGURE 2

SEQ ID NO: 1 – Her2/neu SEQUENCE

1255 AA; 137909 MW; 398DFDA04DCF962 CRC64;

MELAALCRWG LLLALLPEGA ASTQVCTGTD MCLRLLFASPE THLDMLRHLX QGQOVVQGNL  
ELTYLPTNAS LSEFLQDIEV QGYVLLAHNQ VROVFLQRLR IVRGTLQLED NYALAVLDNG  
DPLNNTTPVT GASPGELREL QRLSLTEILK GSVLIQRNEQ LCYQDTILWK DIFKKNQOLA  
LTLIDTNRSR ACHFCSMCK GSKCWGSESE DCQSLTRIVC AGGCARCKGP LPTDCCHEQC  
AAGCTGPKHS DCLACLHENH SGICELHCPA LVYNTIDTFE SMRFESRYT FGASCVTACP  
YNYLSTDVGS CELYVCPLHNQ EVTAEDGTQR CEKCSKPCAR VCYGLGMEHL REVRAVTSAN  
IOEFAGCKKI FGSLLAFERS FDGDFASNTA PLOPEQLQVF ETHEETGYL YISAWPDSLF  
DLSVFQNLQV IGRILHNGA YSLTLQGLGI SWLGLRSLRE LQSGLALIHK NTHLCFVHTV  
PWEQLFRNEH QALLTANR EDECVGEGIA CHOLCARGHC WGEPTCCVN CSQFLRGQEC  
VEECRVLQGL PREYVN  
VKPDLSY MPIWKFEDDE GACQPCFINC THSCVDLDDK GCPAEQRASP LTSIIISAVVG  
LLLVVVLGVV FGLIKRQQ KIRKYTMRL LQETELVPEL TFSGAMENQA QMRILKETEL  
RKVKVLGSGA FGTWVKGIWI PDGENVKIPV AIKVLRENTS PRANKEILDE AYVMAGVGGSP  
YVSRLLGICL TSTVQLVTQL MFIGCLLDHV RENRGRLSGQ DDLNWCQIA KMSYLEDVK  
LVHRDLAARN VLKSENHVK ITDFGLARLL DIDEITYHAD CCKVPIKWA LESILRRFT  
HQSDVWSYGV TVWELMFGA KPYDGIKARE IEDLLEKGER LEQPPYCTID YVMIMVECRM  
IDSECRFRF ELVSESRMA RDFQRFVVIQ NEDLGPASPL DSTFYRSLLE DDDMGDLVDA  
EYLVLPQQGF FCPDFAPGAG GMVHHRHRS STREGGGLT LGLBFSSEEA PRSPLAPSEG  
AGSDVFDGDL GMGAANGLQS LPTHDPSPLO RYSEDPTVPL PSETDGYVAP LTCSPQPEYV  
NCPDVRFPQP SREGELPAA RFAGATLERP KILSPGKNGV VKDVFAPGGA VENPEYLLTPQ  
GGAAPQPHFP PAFSPADNL YYWDQDPPER GAPPSTFKGT PTAENPEYLG LDVVFV

SEQ ID NO: 2 – E75 Vaccine: KIFGSLAFL

SEQ ID NO: 3 – HER2 Herceptin binding region  
PREYVNAHCLPCHPECCPQNGSVTCFGPEADQCACAHYKDPPEFCVARCPG

FIGURE 3

**HLA peptide motif search results**

User Parameters and Scoring Information	
method selected to limit number of results	explicit number
number of results requested	20
HLA molecule type selected	A_0201
length selected for subsequences to be scored	9
echoing mode selected for input sequence	Y
echoing format	numbered lines
length of user's input peptide sequence	53
number of subsequence scores calculated	45
number of top-scoring subsequences reported back in scoring output table	20

Scoring Results				
Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)	SEQ ID NO:
1	27	FGPERDQCV	8.563	4
2	23	SVTCFPEEA	1.652	5
3	17	EOPONGSVT	0.310	6
4	2	REYVRAHHC	0.062	7
5	38	AHYKDFEFC	0.061	8
6	5	VNARKCLPC	0.055	9
7	18	QPONGSVTC	0.032	10
8	40	YKDFEFCVA	0.010	11

FIGURE 4

**HLA peptide motif search results**

User Parameters and Scoring Information	
method selected to limit number of results	explicit number
number of results requested	50
HLA molecule type selected	A_0201
length selected for subsequences to be scored	9

Scoring Results			SEQ ID NO:	
Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)	
1	64	FLPEADQCV	1311.751	12
2	46	GILGFVFTL	550.927	13
3	10	KIPESLAFI	481.186	2
4	100	IMDQVFFSV	198.115	14
5	28	RILCFTEIV	126.098	15
6	73	FVPEADQCV	114.778	16
14	82	FGPEADQCV	8.563	4
31	19	IISAVVGIL	0.612	17
32	37	IVSAVVGIL	0.390	18

FIGURE 5

T2 Peptide-binding Assay	HLA-A2 levels are indicated as Mean Fluorescence Intensity units	
	Expt 1	Expt 2
NP	683	367
Flu M	1357	857
E75	1151	474
GP2	682	579
<b>2G-577</b>	<b>1275</b>	<b>849</b>
2L-577		831
2V-577		796

25ug of peptide incubated with T2 cells

Levels of HLA-A2 were measured by flow cytometry

FIGURE 6

2G-577-stimulated cells can lyse tumor cells			
Results expressed as % lysis			
	Prostate Ca LNCaP	Ovarian Ca SKOV3-A2	Breast Ca MCF-7
PBMC 3			
CM	18	29	36
E75	22	35	38
2G-577	36	38	50
PBMC 7			
CM	22	37	38
E75	32	38	30
2G-577	42	47	39
PBMC 14			
CM	47	41	46
E75	47	43	52
2G-577	52	50	52

FIGURE 7

**Enhanced lysis of Herceptin-treated tumor targets by 2G, 2V, 2L-577-stimulated cells**

**Results expressed as % lysis**

	MCF-7	MCF7+10 Hercp	MCF7+ 50 Hercp	LNCaP	LNCaP+ 50 Hercp
PBMC 8					
CM	46	49	53	15	46
E75	47	51	53	30	47
GP2	47	45	51	18	45
2G-577	35	48	52	27	48
2V-577	48	56	58	21	51
2L-577	50	54	50	26	52
PBMC 10					
CM	45	52	58	20	33
E75	54	60	62	30	52
GP2	51	58	66	35	41
2G-577	54	61	64	32	61
2V-577	57	64	64	27	85
2L-577	56	64	67	31	56

FIGURE 8

Enhanced lysis of Herceptin-treated tumor targets by 2G-577-stimulated cells from Breast Cancer patients. (Results expressed as % lysis)

	Sample	MCF-7	MCF7+50ug Hercep
BCa	NNV24	33	68
BCa	NNV24	36	77
BCa	B53V20	45	93
BCa	B53V20	47	100
Normal Donor	PBMC 3	32	80
Normal Donor	PBMC 3	29	88

## TARGETED IDENTIFICATION OF IMMUNOGENIC PEPTIDES

### REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 13/774,829, filed Feb. 22, 2013 (allowed), which is a continuation of U.S. patent application Ser. No. 12/045,402, filed Mar. 10, 2008 (now U.S. Pat. No. 8,945,573), which is a continuation of International Application No. PCT/US06/35171 filed Sep. 8, 2006, which claims priority to U.S. Provisional Application No. 60/714,865 filed Sep. 8, 2005, the entirety of which is hereby incorporated by reference.

### RIGHTS IN THE INVENTION

This invention was made with government support under grant number W81XWH-05-2-0053 awarded by the U.S. Army Medical Research and Materiel Command. The government has certain rights in this invention.

### BACKGROUND OF THE INVENTION

#### 1. Field of the Invention

This invention, in the field of immunology/immunotherapy, vaccine discovery and development, relates generally to the identification of immunogenic peptides from regions of proteins and molecules that are involved in the binding interactions with polyclonal and monoclonal antibodies and other specific binding peptides/molecules. The present invention is directed to methods for identification and use of the peptides for preventing, suppressing and treating immune-related diseases. Specifically, the invention provides therapy that result in clinical improvement in cancer patients.

#### 2. Description of the Background

Autoimmune diseases are characterized by an unwanted and unwarranted attack by the immune system on the tissues of the host. While the mechanism for progress of these diseases is not well understood, at least some of the details with respect to antigen presentation are known. It is thought that antigens, including autoantigens, are processed by antigen-presenting cells (APC), and the resulting fragments are then associated with one of the cell surface proteins encoded by the major histocompatibility complex (MHC). As a result, recognition of a peptide antigen is said to be MHC "restricted." When the MHC/antigen fragment complex binds to a complementary T cell receptor (TCR) on the surface of a T lymphocyte, activation and proliferation of the clone or subpopulation of T cells result bearing that particular TCR. Once activated, T cells have the capacity to regulate other cells of the immune system which display the processed antigen and to destroy the cells or tissues which carry epitopes of the recognized antigen.

Antibody therapies in which antibodies are directed to MHC molecules and CD4 molecules have been generally successful in several animal models of autoimmunity. However, these approaches may be too nonspecific and potentially overly suppressive. This may be because 70% of T cells bear the CD4 marker and because all T cell-mediated responses and most antibody responses require MHC-associated antigen presentation.

A major difficulty with present approaches is that they require the use of complex biological preparations which do not comprise well-defined therapeutic agents. Such preparations suffer from complex production and maintenance

requirements (e.g., the need for sterility and large quantities of medium for producing large number of "vaccine" T cells), and lack reproducibility from batch to batch. To be useful in humans, T cell "vaccine" preparations must be both autologous and individually specific. This means they must be uniquely tailored for each patient. Furthermore, the presence of additional antigens on the surface of such T cells may result in a broader, possibly detrimental, immune response not limited to the desired T cell clones (Offner et. al., *J. Neuroimmunol.* 21:13-22 (1989).

There is a need, therefore, for agents and pharmaceutical compositions which have the properties of specificity for the targeted immune response. These agents and compositions should also have predictability in their selection, convenience and reproducibility of preparation, and sufficient definition in order to permit precise control of dosage.

As effective vaccine is capable of generating a long-lasting immunity while being relatively harmless to the recipient. Attenuated organisms and purified antigens from organisms have traditionally been used as vaccines. However, such agents often produce deleterious side effects or fail to protect against subsequent challenges. Because of the inherent difficulties in growing pathogenic organisms and producing effective vaccines from them, many viral, bacterial and parasitic diseases have no effective vaccine.

A further difficulty with the use of peptides as vaccines is that, in most instances, peptides alone are not good immunogens. It is a well known phenomenon that most immune responses to peptide antigens are T cell-dependent. Accordingly, "carrier" molecules have been attached to peptide antigens that bind, for example, to B cell surface immunoglobulin in order to generate a high affinity, IgG response. In other words, nonresponsiveness to peptide antigens may sometimes be overcome by attaching another peptide that induces helper T cell activity.

In general, peptides that induce helper T cell activity are generated by B cells from enzymatic digestion of native proteins internalized by way of an antibody receptor. These T cell stimulating peptides are then presented on the surface of the B cell in association with class II major histocompatibility complex (MHC) molecules. In a similar fashion, peptides that induce cytotoxic T cell activity may be generated by accessory cells, including B cells. These peptides are presented on the cell surface of accessory cells in association with class I MHC molecules. As used herein, the term "T cell stimulatory peptide" means any peptide which activates or stimulates T cells, including (but not limited to) helper T cells and/or cytotoxic T cells.

Peptides represent a promising approach to the production and design of vaccines. However the difficulties in making peptides that induce the desired immune response have hampered their success. This includes the difficulties inherent in making peptides that closely mimic the native structure of antigenic determinants.

These antigenic determinants, or epitopes, of a protein antigen represent the sites that are recognized as binding sites by certain immune components such as antibodies or immunocompetent cells. While epitopes are defined only in a functional sense, i.e. by their ability to bind to antibodies or immunocompetent cells, there is a structural basis for their immunological activity.

Epitopes are classified as either being continuous and discontinuous. Discontinuous epitopes are composed of sequences of amino acids throughout an antigen and rely on the tertiary structure or folding of the protein to bring the sequences together and form the epitope. In contrast, con-

tinuous epitopes are linear peptide fragments of the antigen that are able to bind to antibodies raised against the intact antigen.

Many antigens have been studied as possible serum markers for different types of cancer because the serum concentration of the specific antigen may be an indication of the cancer stage in an untreated person. As such, it would be advantageous to develop immunological reagents that react with the antigen. More specifically, it would be advantageous to develop immunological reagents that react with the epitopes of the protein antigen.

Conventional methods using biochemical and biophysical properties have attempted to determine the location of probable peptide epitopes. These methods include a careful screening of a protein's primary structure, searching for critical turns, helices, and even the folding of the protein in the tertiary structure. Continuous epitopes are structurally less complicated and therefore may be easier to locate. However, the ability to predict the location, length and potency of the site is limited.

Various other methods have been used to identify and predict the location of continuous epitopes in proteins by analyzing certain features of their primary structure. For example, parameters such as hydrophilicity, accessibility and mobility of short segments of polypeptide chains have been correlated with the location of epitopes.

Hydrophilicity has been used as the basis for determining protein epitopes by analyzing an amino acid sequence in order to find the point of greatest local hydrophilicity. As discussed in U.S. Pat. No. 4,554,101, each amino acid is assigned a relative hydrophilicity numerical value which is then averaged according to local hydrophilicity so that the locations of the highest local average hydrophilicity values represent the locations of the continuous epitopes. However, this method does not provide any information as to the optimal length of the continuous epitope. Similarly, U.S. Pat. No. 6,780,598 B1 determines the immunopotency of the epitope by providing a ranking system delineating between dominant and subdominant epitopes.

Computer-driven algorithms have been devised to take advantage of the biochemical properties of amino acids in a protein sequence by sorting information to search for T cell epitopes. These algorithms have been used to search the amino acid sequence of a given protein for characteristics known to be common to immunogenic peptides. They can often locate regions that are likely to induce cellular immune response in vitro. Computer-driven algorithms can identify regions of proteins that contain epitopes which are less variable among geographic isolates, or regions of each geographic isolate's more variable proteins, or perform as a preliminary tool to evaluate the evolution of immune response to an individual's own quasi species.

Peptides presented in conjunction with class I MHC molecules are derived from foreign or self protein antigens that have been synthesized in the cytoplasm. Peptides presented with class II MHC molecules are usually derived from exogenous protein antigens. Peptides binding to class I molecules are usually shorter (about 8-10 amino acid residues) than those that bind to class II molecules (8 to greater than 20 residues).

Identification of T cell epitopes within protein antigens has traditionally been accomplished using a variety of methods. These include the use of whole and fragmented native or recombinant antigenic protein, as well as the more commonly employed "overlapping peptide" method for the identification of T cell epitopes within protein antigens which involves the synthesis of overlapping peptides span-

ning the entire sequence of a given protein. Peptides are then tested for their capacity to stimulate T cell cytotoxic or proliferation responses in vitro.

The overlapping peptide method is both cost and labor intensive. For example, to perform an assay using 15 amino acid long peptides overlapping by 5 amino acids spanning a given antigen of length n (a small subset of the possible 15-mers spanning the protein), one would need to construct and assay  $(n/5)-1$  peptides. For most types of analyses, this number would be prohibitive.

Accordingly, a simple method to identify immunogenic peptides from regions of self-proteins and other proteins and molecules involved in binding interactions with polyclonal and monoclonal antibodies is needed.

#### DESCRIPTION OF THE FIGURES

FIG. 1 depicts one embodiment of the invention in which a unique immunogenic region of the HER-2/neu is identified.

FIG. 2 shows the Her2/neu sequence (SEQ ID NO: 1) and highlights the antibody-binding site of Herceptin to the Her2/new protein. Also shown is the E75 vaccine sequence (SEQ ID NO: 2) and the Her2 loops binding to Herceptin (SEQ ID NO: 3),

FIG. 3 shows HLA peptide motif search results and scoring results for SEQ ID NOs: 4-11.

FIG. 4 shows amino acid substitutions for increased binding affinity of native peptide vaccine development. A table indicating HLA peptide motif search results, as well as the scoring results for SEQ ID NOs: 12-20 are included.

FIG. 5 depicts a T2 Peptide-binding assay, in which 25  $\mu$ g of peptide were incubated with T2 cells, levels of HLA-A2 levels were measured by flow cytometry and are indicated as Mean Fluorescence Intensity units.

FIG. 6 is a table depicting an experiment looking at whether 2G-577- and E75-stimulated cells can lyse tumor cells, in prostate cancer, ovarian cancer, and breast cancer, with results expressed as % lysis.

FIG. 7 is a table showing enhanced lysis of Herceptin-treated tumor targets by B75, GP2, 2G-577, 2V-577, and 2L-577-stimulated cells, with results expressed as % lysis. These data suggest the benefit of combination therapy with Herceptin treatment and vaccine-induced T cell targeting for synergistic tumor cell killing.

FIG. 8 is a table showing enhanced lysis of Herceptin-treated tumor targets by 2G-577-stimulated cells from breast cancer patients, with results expressed as % lysis.

#### SUMMARY OF THE INVENTION

The invention overcomes the problems and disadvantages associated with current methods and provides tools and methods of generating an immune response in a patient in need thereof.

One embodiment of the invention is directed to a method for synthesizing an immunogenic peptide from a self-protein comprising the steps of identifying one or more peptide sequences of a self protein that are directly or indirectly involved with antibody-binding, subjecting the one or more peptide sequences to an algorithm that identifies sequences suspected of being immunogenic, screening all peptide fragments from the one or more peptide sequences, and identifying an immunogenic peptide of the protein fragment wherein the antibody-binding interaction is polyclonal or monoclonal. Further, a patient is treated with the immunogenic peptide to generate an immune response.

Another embodiment of the invention is directed to immunogenic peptides identified by the method described above.

Another embodiment of the invention is directed to an immunogenic peptide that produces an immune response to a self-protein.

Another embodiment is directed to a method of presenting epitopes for recognition by the immune system to generate an immune response comprising the steps of identifying a protein fragment that is recognized by a pool of unused and immunoreactive T cells, subjecting the protein fragment to an algorithm, identifying one or more specific sequences of the protein fragment that is immunogenic, synthesizing at least one immunogenic peptide corresponding to the sequence and treating a patient with the immunogenic peptide to generate an immune response. Further, an antibody that binds to the protein is generated.

Another embodiment of the invention is directed to immunogenic peptides identified by the method described above.

Another embodiment of the invention is directed to a vaccine comprising the immunogenic peptides described above.

Another embodiment of the invention is directed to a vaccine comprising antibodies that react with the immunogenic peptides described above.

Another embodiment of the invention is directed to a method of identifying a vaccine treatment comprising the step of binding an antibody to a protein molecule forming a complex, subjecting the complex to proteasome digestion, obtaining digestion products comprising peptides, and identifying an immunogenic peptide sequence from the digestion products.

Another embodiment of the invention is directed to a method of identifying a patient-specific treatment comprising the steps of obtaining a pre-existing immuno-reactive precursor to said patient's AGIE/ABIE, culturing tumor cells obtained from said patient, incubating the cultured tumor cells that are reactive against generated antibodies, examining dataset responses in presence and absence of the generated antibodies and identifying the patient-specific immunogenic epitopes. This method may include the generation of antibodies that are reactive against self antigens, the generation of antibodies that are reactive against foreign antigens, and/or the generation of antibodies, once administered to said patient, that are therapeutic or prophylactic.

Another embodiment of the invention is directed to a method of identifying a vaccine treatment comprising the steps of binding an antibody to a protein molecule with a specific binding activity forming a complex, subjecting the complex to proteasome digestion, obtaining digestion producing comprising peptides, identifying an immunogenic peptide sequence from the digestion products.

Other embodiments and technical advantages of the invention are set forth below and may be apparent from the drawings and the description of the invention which follow, or may be learned from the practice of the invention.

#### DESCRIPTION OF THE INVENTION AND EXAMPLES

Treatments for complex diseases involving the immune system and immune responses caused by endogenous self antigens and/or foreign antigens that are involved in producing autoimmune antibody are extremely difficult to discover. Antigens involved with such diseases are either foreign or self antigens (or both). Administration of foreign

antigens for passive immunization can result in serum sickness-like immune complex diseases. Also, reactive T cells capable of recognizing self peptides are typically deleted or processed and destroyed. These peptides, generated and displayed under normal and constitutive conditions are degraded by cell protein degradation machinery resulting in the absence of an immune response.

A simple method for identification of immunogenic regions of self-proteins and other proteins and molecules involved in the binding interactions with polyclonal and monoclonal antibodies has been surprisingly discovered. From this method, new and unique epitopes are generated that are presented in the presence of bound ligands, such as, preferably antibodies. Once these immunogenic regions are identified, vaccines comprising the antigen, modifications of the antigen, or antibodies specifically reactive to the antigen or the modified antigen can be prepared. Thus, the present invention makes vaccine peptide discovery manageable, and allows for the specific generation of unique immunogenic peptides from self-tumor associated proteins that can be induced or generated for specific expression in the presence of the antibody, allowing for vaccine end/or novel combination therapy.

The invention is described more fully herein and refers to many preferred embodiments. This invention, however, should not be construed as limited to those embodiments.

The proteasome is a multi-subunit complex with proteolytic cleavage activities that result in the generation of a wide variety of peptides from proteins. The susceptibility of a given protein to the proteolytic activities of the proteasome is dependent upon the various primary, secondary and tertiary structural and post-translational modifications that take place within the proteasome. These activities expose certain sequences or regions of the protein and not others, to the system.

In one embodiment of the invention, cancer cells are induced to express immunogenic peptides from unique or self tumor-specific antigens to stimulate anti-tumor immune responses for immunotherapy. Normally these peptides are not generated from self-proteins. Binding of antibodies (or other molecules) to the sites normally accessible and processed by proteasomes alters the pattern of accessibility and the resulting proteolytic cleavage pattern by the proteasome. Such alteration of the site results in generation of novel peptides that may be intrinsically immunogenic because they have not been previously expressed and displayed for the deletion of immune-reactive T cells.

In one preferred embodiment, a unique immunogenic region of the HER-2/neu was identified. HER-2/neu is an over-expressed oncogenic protein. Conventional vaccine strategies, normally effective in immunizing patients, did not work with a "self-protein" such as HER-2/neu. Tolerance to self-proteins may only be directed to dominant epitopes of the protein and not the entire protein. Therefore, immunization to just a specific protein fragment, and not the entire protein, alleviates this problem. This specific protein fragment is located within the sequence directly involved with antibody-binding interaction or in the proximity of that region.

After identifying this restricted, shorter peptide sequence, the sequence is subjected to algorithms to identify likely functionally active or target sequences or regions. Running algorithms on this segment, as opposed to the whole segment, provides a manageable set of peptides to test as candidates for vaccine development. In the past, computer-driven tests were run on the entire sequence consuming much time, money and effort. The algorithms searched the

amino acid sequence provided for characteristic immune response *in vitro*. Regions of the proteins identified as containing epitopes may be useful as a vaccine.

Next, treatment of the tumor cells with the antibody against the identified peptide sequence was performed. Inducibility of the altered turnover and subsequent generation of the newly identified peptides only in the tumor cells and only in the presence of the antibody gave it a specific targeting and triggering feature that was controllable. See FIG. 1. An antibody booster can be given to increase the peptide-specific cytotoxic T cell response. In cases where the antibody already existed, discovery of the peptide only was sufficient for triggering the specific immune response.

The method to generate new and unique epitopes included the identification of immunogenic peptides from regions of proteins and molecules involved in the binding interactions with most any ligand including, but not limited to polyclonal and monoclonal antibodies, receptors, ligands and any molecule with high affinity to the antigen. These new epitopes were presented to antigen-processing cells in the presence of bound ligand. In certain cases, binding protected the epitope or a portion of the epitope revealing the immunogenic portion of the antigen. These immunogenic regions become new and uniquely enhanced targets for recognition by the immune system and for use in modulation of the immune responses for the treatment of disease states and immunotherapy either by themselves, an antibody vaccines, or when used in combination with other molecules or antibodies. These antibody-generated immunogenic epitopes (AGIE)/antibody-bound immunogenic epitopes (ABIE) were useful for stimulating immune response in cancer and infectious diseases. They can also suppress immune responses in autoimmune diseases and transplantation based upon the activity of the antibody in modulating the immune response in the direction of upregulation or down regulation.

Peptides varying in length including single chain antigen-binding polypeptides that can specifically interact, protect and/or modulate any given epitope such that it results in specifically protecting and/or enhancing the preservation and presentation of the epitope when subjected to the various proteolytic activities of the antigen processing machinery of the proteasome and immunoproteasome subunits and complexes.

Another embodiment of this invention comprises a combination of specific antibodies and/or treatment with other molecules that involve the same regions of the peptide sequences and targets unique populations of cells. This technology allows for the development of novel vaccines made of antibodies and/or antigen-binding fragments Fab or F(ab)<sub>2</sub>-fragments bound to specific length peptides that contain immunogenic epitopes(s) of interest that are specifically processed and presented.

Specific epitopes of proteins may be protected, both intracellular and extracellular, that are normally destroyed by antigen processing machinery of proteasomes and immunoproteasomes by single chain antibodies and/or peptides that can specifically bind to selected site sequences so that these epitopes become novel targets that are developed as unique vaccines for the specific recognition of cancer cells. HSPs that are released during inflammatory processes perform similar activities and enhance immune responses because they were made to bind to many different peptide sequences non-covalently.

Another preferred embodiment of the invention is directed to a method for the identification of highly immunogenic peptides. These peptides are used to enhance or suppress immune responses from normal cell proteins that

are not processed and presented naturally because of the constitutive destruction of these epitopes by the normal activities of the proteasome and immunoproteasome complexes. These peptides are used for generation for specific antibodies that bind and generate the novel presentation of these epitopes for recognition by the immune system.

In a preferred example, a large pool of immunoreactive T cells that are present or may be generated that are capable of reacting with these inert/naturally non-existent epitopes, but are not utilized and are considered "wasteful" because these epitopes are not being generated normally. This allows access to the pools of unused T cells for specifically directed therapeutic benefits. Again, computer-driven algorithms are run. Upon identification of the peptide sequences involved with antibody-binding interaction, these peptides are used to immunize animals and/or patients to generate specific antibodies that bind the protein and generate these novel peptides for subsequent immune recognition and response.

Another embodiment of the invention is directed to a direct method of binding antibody or antibodies to the protein molecules and/or polypeptide regions and subjecting these bound and unbound/native complexes to *ex vivo* or *in vitro* to all forms of the proteasome machinery. Proteolytic digestion or cleavage products are obtained to observe differential yield of peptides for use in vaccine development. However, there is a drawback to this method in that it is not fully representative of all "mechanisms and proteolytic activities" that may occur *in vivo* within the cell itself, thereby limiting application. Such an approach however, is useful for the proof of concept in systems where it provides clean reproducible and predictable results, e.g. using various individual components of proteasomes and combinations thereof and then observing peptide patterns by mass spectroscopy analysis. It is also useful for the definition of peptide patterns generated in the presence and/or absence of proteasome inhibitors with and without bound antibody or antibodies.

Another embodiment of the present invention involves application of this approach for identification of patient-specific treatment. Optimal antibody choices are made for the patient based upon patient's personal pre-existing immuno-reactive precursors to the AGIE/ABIE. First, PBMC (peripheral blood mononuclear cells) are obtained from patient and culture in presence or absence of cytokines (eg IL-2, IL-7, and IL-15) in the presence of FCS (fetal calf serum) or autologous serum. After optimal culture and expansion during a set number of days, the cells are tested in cytotoxicity assays against tumor cell lines that have been pre-incubated in the presence and absence of specific antibodies or antibody combination(s). Datasets from responses seen specifically to presence of antibodies and those where individual antibodies do not yield a response, but the combinations that do are noted. The epitopes/vaccine peptides predicted or defined by the binding-sites of the antibodies are tested for fine reactivities in determination of which vaccines to use on patients.

Another embodiment of the invention is directed to identifying antigens that are not expressed on the cell surface or for parts of protein that are always cytoplasmic or for proteins that are completely intracellular/intranuclear or cytoplasmic (e.g. p53, telomerase, etc.). Methods for expressing the "modulating" antibody or antibodies as endogenous proteins, so that they are able to bind these proteins intracellularly and modulate their processing and presentation from within. The endogenous expression methods included recombinant DNA expression systems as well as viral/bacterial vector deliver and expression systems.

Another embodiment of the invention is directed to “shuttling” in the antibodies or single-chain antigen-binding fragment encapsulated within various delivery systems to include liposomes, micelles, nanoparticles and others.

Generation of “protective” or “suppressive” antibody preparations or combination capable of being administered passively for treatment of disease similar to the gamma-globulin shots were generated as polyclonal preps or as specific combinations of selected antibodies against one or more tumor or tissue specific antigens for that particular disease or condition.

The present invention is not restricted to HLA-A2 or class I peptides because one or more longer-length peptides from a known region or sequence can be used for the generation of AGIE/ABIE by the antibody or antibodies in question. The benefit is not having to be restricted by only specific or limited HLA types (Class I and II) in terms of patients that can be treated. Furthermore, although the majority of Class I and Class II peptides are derived from the processing of endogenous and exogenous proteins respectively, the well described and accepted mechanisms of “cross-presentation” allows for the presentation of all sources or peptides on both classes of MHC molecules.

In another embodiment of the invention, the invention is used for identifying inducible vaccine responses by “discovering” the MAb/s that will generate AGIE/ABIE. This is achieved by first screening 10-mer or 20-mer consecutive or overlapping peptides from antigen for the highest precursor T-cell responses present in cancer and/or normal individuals. These “ultra-immunogenic” peptides are then used for generation of MAb in mice. These MAbs are then tested for ability to generate AGIE/ABIE specific responses in MAb-treated targets. MAbs that are promising are then humanized for therapeutic purposes. To ensure involvement of T cell specific immune responses the promising Fab' fragments are first used to eliminate purely antibody-dependent cell-mediated cytotoxicity (ADCC)-mediated activity.

Another way of specifically identifying peptides capable of generating antibodies that induce AGIE/ABIE that are most protective is by the method of screening serum sample from high-risk cancer individuals (those with family history of cancer, e.g. smokers who do and do not get lung cancer, patients who are ‘cured’) who do and do not develop cancer (or from progressor vs non-progressors in the case of AIDS) to look for Ab responses against the overlapping peptides from specific well characterized or defined tumor antigens (HER2/neu, prostate specific antigen or PSA, prostate specific membrane antigen or PSMA, Tyrosinase, melanoma Ags, etc.) that are present in the protected individuals or present in fully recovered/cured cancer patients. Based on the specific Ab responses that are found to be uniquely present or predominantly present in the ‘protected’ individuals, peptides are targeted for generation or ‘discovering’ of MAb that generated AGIE/ABIE. All descriptions herein were done for both Class I and Class II epitopes and responses.

In a preferred embodiment, data of appropriate combinations of antibodies from published sources with well documented “pre-existing” corresponding CD4-helper and CB8 CTL-specific responses is used for developing a vaccine for the AIDS virus and other immunotherapy treatment using the approach from the present invention. The HIV Molecular Immunology Database provided by the National Institute of Allergy and Infectious Diseases at [www.hiv.lanl.gov](http://www.hiv.lanl.gov) offers a current comprehensive listing of defined HIV epitopes. The website also includes the exact epitope and binding site

of all known monoclonal and polyclonal antibodies to the protein sequences as well as neutralization activity.

The present invention provides enhanced lytic activity of antibody-treated tumor cells by vaccines identified from the binding sites of these antibodies. The present invention also provides a novel indication for antibodies already in clinical use for cancer treatment as combination therapy furthering development of vaccine treatment discovery.

## EXAMPLE

The binding of trastuzumab (Herceptin) to HER2/neu for the development of novel vaccines against HER2/neu-expressing tumors was studied. The Ab-binding site of trastuzumab (Tz) was analyzed for peptides that bind HLA-A2 and A3. A peptide Her577 (aa; 577-585) within the Ab-binding site on HER2/neu was identified, synthesized and tested. T2 HLA-stabilization assays were performed to confirm HLA-A2 binding activity of Her577 by flow cytometry. PBMC from 9 healthy donors were stimulated with Her577 and tested in <sup>51</sup>Cr-release cytotoxicity (CTX) assays against 3 HER2/neu-expressing tumor cell lines. Her577-stimulated PBMC from 5 HLA-A2<sup>+</sup> healthy donors were also tested in CTX with HER2/neu<sup>+</sup> targets pre-treated with Tz. Simultaneous experiments were done with E75 and GP2, two other immunogenic peptides from HER2/neu.

The Her577 peptide bound HLA-A2 comparable to E75 and GP2 with mean fluorescence intensity values of 1275, 1151 and 682, respectively. The average specific CTX by Her577-stimulated cytotoxic T lymphocytes (CTL) against LNCaP, SKOV-3 and MCF-7 tumor cells was found to be similar to CTX achieved with E75- and GP2-stimulated CTL (% lysis; Table 1).

TABLE 1

	LNCaP (HLA-A2*) (n = 2)	SKOV-3 (HLA-A3*) (n = 2)	MCF-7 (HLA-A2*) (n = 5)
Her577	30 ± 3	49 ± 28	45 ± 3
E75	30 ± 1	51 ± 20	51 ± 4
GP2	26 ± 9	46 ± 5	48 ± 1

Pre-treatment with Tz at 50 µg/ml for 24 h increased specific CTX by Her577-stimulated CTL versus untreated MCF-7, SKOV-3 and LNCaP by 16%, 47% and 83%, respectively (P=0.19). A dose-dependent incremental increase in CTX against each tumor cell line was seen with 10 and 50 µg/ml doses of Tz.

It can be concluded that Her577 is a newly described T cell-epitope from HER2/neu identified by using a novel method for the discovery of immunogenic peptide(s) for vaccine development by assessing the binding site of Tz. There is a potential for combination immunotherapy with therapeutic Ab and unique peptide vaccines derived from the Ab-binding site.

Other embodiments and uses of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. All references cited herein, including all patents and publications that are cited for any reason, including U.S. Provisional Application No. 60/714,865, on which priority is based, are specifically and entirely incorporated by reference. The specification and examples should be considered exemplary only with the true scope and spirit of the invention embodied within the following claims.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1

<211> LENGTH: 1255

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic construct

<400> SEQUENCE: 1

```

Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu
  1           5           10           15

Pro Pro Gly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys
  20           25           30

Leu Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His
  35           40           45

Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu Glu Leu Thr Tyr
  50           55           60

Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val
  65           70           75           80

Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu
  85           90           95

Gln Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr
 100           105           110

Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro
 115           120           125

Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser
 130           135           140

Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln
 145           150           155           160

Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn
 165           170           175

Asn Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys
 180           185           190

His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser
 195           200           205

Ser Glu Asp Cys Gln Ser Leu Thr Arg Thr Val Cys Ala Gly Gly Cys
 210           215           220

Ala Arg Cys Lys Gly Pro Leu Pro Thr Asp Cys Cys His Glu Gln Cys
 225           230           235           240

Ala Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys Leu
 245           250           255

His Phe Asn His Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu Val
 260           265           270

Thr Tyr Asn Thr Asp Thr Phe Glu Ser Met Pro Asn Pro Glu Gly Arg
 275           280           285

Tyr Thr Phe Gly Ala Ser Cys Val Thr Ala Cys Pro Tyr Asn Tyr Leu
 290           295           300

Ser Thr Asp Val Gly Ser Cys Thr Leu Val Cys Pro Leu His Asn Gln
 305           310           315           320

Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser Lys
 325           330           335

Pro Cys Ala Arg Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg Glu
 340           345           350

```

-continued

---

Val Arg Ala Val Thr Ser Ala Asn Ile Gln Glu Phe Ala Gly Cys Lys  
 355 360 365

Lys Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly Asp  
 370 375 380

Pro Ala Ser Asn Thr Ala Pro Leu Gln Pro Glu Gln Leu Gln Val Phe  
 385 390 395 400

Glu Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp Pro  
 405 410 415

Asp Ser Leu Pro Asp Leu Ser Val Phe Gln Asn Leu Gln Val Ile Arg  
 420 425 430

Gly Arg Ile Leu His Asn Gly Ala Tyr Ser Leu Thr Leu Gln Gly Leu  
 435 440 445

Gly Ile Ser Trp Leu Gly Leu Arg Ser Leu Arg Glu Leu Gly Ser Gly  
 450 455 460

Leu Ala Leu Ile His His Asn Thr His Leu Cys Phe Val His Thr Val  
 465 470 475 480

Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His Thr  
 485 490 495

Ala Asn Arg Pro Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His  
 500 505 510

Gln Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Gln Cys  
 515 520 525

Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu Cys  
 530 535 540

Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val Asn Ala Arg His Cys  
 545 550 555 560

Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys  
 565 570 575

Phe Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp  
 580 585 590

Pro Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp Leu  
 595 600 605

Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu Gly Ala Cys Gln  
 610 615 620

Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Asp Lys  
 625 630 635 640

Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Ile Ile Ser  
 645 650 655

Ala Val Val Gly Ile Leu Leu Val Val Val Leu Gly Val Val Phe Gly  
 660 665 670

Ile Leu Ile Lys Arg Arg Gln Gln Lys Ile Arg Lys Tyr Thr Met Arg  
 675 680 685

Arg Leu Leu Gln Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly  
 690 695 700

Ala Met Pro Asn Gln Ala Gln Met Arg Ile Leu Lys Glu Thr Glu Leu  
 705 710 715 720

Arg Lys Val Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys  
 725 730 735

Gly Ile Trp Ile Pro Asp Gly Glu Asn Val Lys Ile Pro Val Ala Ile  
 740 745 750

Lys Val Leu Arg Glu Asn Thr Ser Pro Lys Ala Asn Lys Glu Ile Leu  
 755 760 765

-continued

---

Asp Glu Ala Tyr Val Met Ala Gly Val Gly Ser Pro Tyr Val Ser Arg  
 770 775 780

Leu Leu Gly Ile Cys Leu Thr Ser Thr Val Gln Leu Val Thr Gln Leu  
 785 790 795 800

Met Pro Tyr Gly Cys Leu Leu Asp His Val Arg Glu Asn Arg Gly Arg  
 805 810 815

Leu Gly Ser Gln Asp Leu Leu Asn Trp Cys Met Gln Ile Ala Lys Gly  
 820 825 830

Met Ser Tyr Leu Glu Asp Val Arg Leu Val His Arg Asp Leu Ala Ala  
 835 840 845

Arg Asn Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe  
 850 855 860

Gly Leu Ala Arg Leu Leu Asp Ile Asp Glu Thr Glu Tyr His Ala Asp  
 865 870 875 880

Gly Gly Lys Val Pro Ile Lys Trp Met Ala Leu Glu Ser Ile Leu Arg  
 885 890 895

Arg Arg Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Val  
 900 905 910

Trp Glu Leu Met Thr Phe Gly Ala Lys Pro Tyr Asp Gly Ile Pro Ala  
 915 920 925

Arg Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro  
 930 935 940

Pro Ile Cys Thr Ile Asp Val Tyr Met Ile Met Val Lys Cys Trp Met  
 945 950 955 960

Ile Asp Ser Glu Cys Arg Pro Arg Phe Arg Glu Leu Val Ser Glu Phe  
 965 970 975

Ser Arg Met Ala Arg Asp Pro Gln Arg Phe Val Val Ile Gln Asn Glu  
 980 985 990

Asp Leu Gly Pro Ala Ser Pro Leu Asp Ser Thr Phe Tyr Arg Ser Leu  
 995 1000 1005

Leu Glu Asp Asp Asp Met Gly Asp Leu Val Asp Ala Glu Glu Tyr Leu  
 1010 1015 1020

Val Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala Pro Gly Ala Gly  
 1025 1030 1035 1040

Gly Met Val His His Arg His Arg Ser Ser Ser Thr Arg Ser Gly Gly  
 1045 1050 1055

Gly Asp Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu Glu Ala Pro Arg  
 1060 1065 1070

Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp Gly  
 1075 1080 1085

Asp Leu Gly Met Gly Ala Ala Lys Gly Leu Gln Ser Leu Pro Thr His  
 1090 1095 1100

Asp Pro Ser Pro Leu Gln Arg Tyr Ser Glu Asp Pro Thr Val Pro Leu  
 1105 1110 1115 1120

Pro Ser Glu Thr Asp Gly Tyr Val Ala Pro Leu Thr Cys Ser Pro Gln  
 1125 1130 1135

Pro Glu Tyr Val Asn Gln Pro Asp Val Arg Pro Gln Pro Pro Ser Pro  
 1140 1145 1150

Arg Glu Gly Pro Leu Pro Ala Ala Arg Pro Ala Gly Ala Thr Leu Glu  
 1155 1160 1165

Arg Pro Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp Val  
 1170 1175 1180

-continued

---

Phe Ala Phe Gly Gly Ala Val Glu Asn Pro Glu Tyr Leu Thr Pro Gln  
1185 1190 1195 1200

Gly Gly Ala Ala Pro Gln Pro His Pro Pro Pro Ala Phe Ser Pro Ala  
1205 1210 1215

Phe Asp Asn Leu Tyr Tyr Trp Asp Gln Asp Pro Pro Glu Arg Gly Ala  
1220 1225 1230

Pro Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr  
1235 1240 1245

Leu Gly Leu Asp Val Pro Val  
1250 1255

<210> SEQ ID NO 2

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 2

Lys Ile Phe Gly Ser Leu Ala Phe Leu  
1 5

<210> SEQ ID NO 3

<211> LENGTH: 53

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 3

Pro Arg Glu Tyr Val Asn Ala Arg His Cys Leu Pro Cys His Pro Glu  
1 5 10 15

Cys Gln Pro Gln Asn Gly Ser Val Thr Cys Phe Gly Pro Glu Ala Asp  
20 25 30

Gln Cys Val Ala Cys Ala His Tyr Lys Asp Pro Pro Phe Cys Val Ala  
35 40 45

Arg Cys Pro Ser Gly  
50

<210> SEQ ID NO 4

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 4

Phe Gly Pro Glu Ala Asp Gln Cys Val  
1 5

<210> SEQ ID NO 5

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 5

Ser Val Thr Cys Phe Gly Pro Glu Ala  
1 5

-continued

---

<210> SEQ ID NO 6  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 6

Cys Gln Pro Gln Asn Gly Ser Val Thr  
1 5

<210> SEQ ID NO 7  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 7

Arg Glu Tyr Val Asn Ala Arg His Cys  
1 5

<210> SEQ ID NO 8  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 8

Ala His Tyr Lys Asp Pro Pro Phe Cys  
1 5

<210> SEQ ID NO 9  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 9

Val Asn Ala Arg His Cys Leu Pro Cys  
1 5

<210> SEQ ID NO 10  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 10

Gln Pro Gln Asn Gly Ser Val Thr Cys  
1 5

<210> SEQ ID NO 11  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

-continued

---

<400> SEQUENCE: 11

Tyr Lys Asp Pro Pro Phe Cys Val Ala  
1 5

<210> SEQ ID NO 12

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 12

Phe Leu Pro Glu Ala Asp Gln Cys Val  
1 5

<210> SEQ ID NO 13

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 13

Gly Ile Leu Gly Phe Val Phe Thr Leu  
1 5

<210> SEQ ID NO 14

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 14

Ile Met Asp Gln Val Pro Phe Ser Val  
1 5

<210> SEQ ID NO 15

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 15

Arg Leu Leu Gln Glu Thr Glu Leu Val  
1 5

<210> SEQ ID NO 16

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 16

Phe Val Pro Glu Ala Asp Gln Cys Val  
1 5

-continued

---

```

<210> SEQ ID NO 17
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      peptide

```

```

<400> SEQUENCE: 17

```

```

Ile Ile Ser Ala Val Val Gly Ile Leu
  1             5

```

```

<210> SEQ ID NO 18
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      peptide

```

```

<400> SEQUENCE: 18

```

```

Ile Val Ser Ala Val Val Gly Ile Leu
  1             5

```

---

25

The invention claimed is:

1. A method of identifying an immunogenic region of a self-protein comprising the steps of:

identifying one or more peptide sequences of a self protein that are directly or indirectly involved with antibody binding, wherein the identifying of the one or more peptide sequences of the self protein comprises: binding an antibody to the self-protein forming a complex; subjecting the complex to proteasome digestion; and obtaining digestion products comprising one or more peptide sequences;

subjecting the one or more peptide sequences to an algorithm that identifies peptide fragments suspected of being immunogenic;

screening all peptide fragments from the one or more peptide sequences for their ability to induce an immune response; and

identifying an immunogenic peptide of the peptide fragment.

2. The method of claim 1, wherein the antibody binding comprises polyclonal antibody binding.

3. The method of claim 1, wherein the antibody binding comprises monoclonal antibody binding.

4. The method of claim 1, further comprising treating a patient with the immunogenic peptide to generate an immune response.

5. The method of claim 1, wherein the self-protein is HER2/neu.

6. The method of claim 1, wherein the immune response is a cytotoxic T cell response.

7. The method of claim 1, wherein screening all peptide fragments from the one or more peptide sequences for their ability to induce an immune response comprises:

stimulating Peripheral Blood Mononuclear Cells with the peptide fragments from the one or more peptide sequences; and

determining the peptide fragments with the highest T cell responses; or

contacting the peptide fragments from the one or more peptide sequences with serum from high risk cancer individuals who do not develop cancer; and

selecting the one or more peptide fragments that bind to the antibodies in the serum; and

determining if the one or more peptide fragments selected in the previous step can induce an antibody response.

\* \* \* \* \*

专利名称(译)	有针对性地鉴定免疫原性肽		
公开(公告)号	<a href="#">US9720000</a>	公开(公告)日	2017-08-01
申请号	US14/713721	申请日	2015-05-15
[标]申请(专利权)人(译)	HENRY中号JACKSON发现的军事医学的进步		
申请(专利权)人(译)	亨利M.杰克逊基金会为军事医学, INC进步.		
当前申请(专利权)人(译)	亨利M.杰克逊基金会为军事医学, INC进步.		
[标]发明人	PONNIAH SATHIBALAN PEOPLES GEORGE E STORRER CATHERINE E FLORA MICHAEL		
发明人	PONNIAH, SATHIBALAN PEOPLES, GEORGE E STORRER, CATHERINE E FLORA, MICHAEL		
IPC分类号	G01N33/53 A61K39/00 G01N33/68 G01N33/577 C07K16/32 A61K39/395 G01N33/50 G01N33/564 C40B30/02 G01N33/574		
CPC分类号	G01N33/6803 A61K39/0011 A61K39/39558 C07K16/32 C40B30/02 G01N33/5011 G01N33/564 G01N33/574 G01N33/577 G01N33/6845 G01N2500/00 G01N2500/04 A61P35/00 G16B35/00 G16C20 /60 A61K2300/00 A61K39/001106		
优先权	12/045402 2015-02-03 US 60/714865 2005-09-08 US		
其他公开文献	US20150260727A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

本发明一般涉及鉴定参与抗体结合任何蛋白质的肽序列, 用于合成疫苗处理。这种新方法允许更易控制的疫苗肽发现和特异性产生来自自身肿瘤相关蛋白的独特免疫原性肽和/或来自感染性生物的外来蛋白, 仅在抗体存在下进行特异性和/或增强表达。

