



US007279562B2

(12) **United States Patent**  
**Molnar-Kimber et al.**(10) **Patent No.:** **US 7,279,562 B2**  
(45) **Date of Patent:** **Oct. 9, 2007**(54) **RAPAMYCIN CONJUGATES**(75) Inventors: **Katherine L. Molnar-Kimber**,  
Worcester, PA (US); **Craig E. Caufield**,  
New York, NY (US); **Timothy D.**  
**Ocain**, Framingham, MA (US)(73) Assignee: **Wyeth**, Madison, NJ (US)(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 812 days.5,194,447 A \* 3/1993 Kao ..... 514/542  
5,233,036 A 8/1993 Hughes  
5,252,579 A 10/1993 Skotnicki  
5,504,091 A 4/1996 Molnar-Kimber  
5,532,137 A 7/1996 Niwa  
5,665,772 A 9/1997 Cottens  
5,912,253 A 6/1999 Cottens  
5,985,890 A 11/1999 Cottens  
6,187,547 B1 2/2001 Legay  
6,328,970 B1 12/2001 Molnar-Kimber et al.  
6,635,745 B2 10/2003 Sedrani et al.  
2002/0002273 A1 1/2002 Sedrani et al.  
2002/0022717 A1 2/2002 Sedrani et al.(21) Appl. No.: **10/124,386**(22) Filed: **Apr. 18, 2002**(65) **Prior Publication Data**

US 2002/0151088 A1 Oct. 17, 2002

**Related U.S. Application Data**(60) Division of application No. 09/576,951, filed on May  
24, 2000, which is a continuation-in-part of applica-  
tion No. 08/424,983, filed on Apr. 19, 1995, now  
abandoned, which is a continuation of application No.  
08/224,205, filed on Apr. 14, 1994, now abandoned,  
which is a continuation-in-part of application No.  
08/053,030, filed on Apr. 23, 1993, now abandoned.(51) **Int. Cl.****G01N 1/13** (2006.01)**G01N 1/04** (2006.01)(52) **U.S. Cl.** ..... **530/403**; 530/404; 530/405;  
530/406(58) **Field of Classification Search** ..... 530/403,  
530/404, 405, 406

See application file for complete search history.

(56) **References Cited****U.S. PATENT DOCUMENTS**3,929,992 A 12/1975 Sehgal  
3,993,749 A 11/1976 Sehgal  
4,316,885 A 2/1982 Rakhit  
4,401,653 A 8/1983 Eng  
4,650,803 A 3/1987 Stella  
4,885,171 A 12/1989 Surendra  
5,078,999 A 1/1992 Warner  
5,080,899 A 1/1992 Sturm  
5,089,390 A 2/1992 Davalian  
5,091,389 A 2/1992 Ondeyka  
5,100,883 A 3/1992 Schiehser  
5,100,899 A 3/1992 Calne  
5,102,876 A 4/1992 Caufield  
5,118,677 A \* 6/1992 Caufield ..... 514/183  
5,118,678 A 6/1992 Kao  
5,120,842 A 6/1992 Failli  
5,122,511 A 6/1992 Patchett  
5,130,307 A 7/1992 Failli  
5,138,051 A 8/1992 Hughes  
5,151,413 A 9/1992 Caufield  
5,164,495 A 11/1992 Lunetta  
5,169,773 A 12/1992 Rosenthaler  
5,169,851 A 12/1992 Hughes  
5,177,203 A \* 1/1993 Failli et al. .... 540/456**FOREIGN PATENT DOCUMENTS**EP 0 201751 11/1986  
EP 283801 3/1988  
EP 473961 8/1991  
EP 487289 11/1991  
EP 506032 3/1992  
EP 507555 3/1992  
EP 525960 6/1992  
EP 0 693 132 1/1996  
WO WO 90/08957 8/1990  
WO WO 90/06763 9/1990  
WO WO 92/05179 4/1992  
WO WO 92/22332 12/1992  
WO WO 93 19752 10/1993  
WO WO 93 25533 12/1993  
WO WO 94/24304 10/1994**OTHER PUBLICATIONS**Hultsch et al, *Proceedings of the National Academy of Sciences of*  
*USA*, 88:6229-6233 (1991).  
Chen et al, *Journal of the American Chemical Society*, 116:2683-  
2684 (1994).  
Morris, *Med. Sci. Res.*, 17:877 (1989).  
Baeder, *Fifth Int. Conf. Inflamm. Res. Assoc.*, Abstract, p. 121  
(1990).  
Gregory, *J. Heart Lung Transplant.*, 11(pt 2):197 (1992).  
Meiser, *J. Heart Lung Transplant.*, 9:55 (1992).  
Stepkowski, *Transplantation Proceedings*, 21(1):507-508 (1991).  
Tamura, *Transplantation Proceedings*, 19(6):23-29 (1987).  
Cadoff, *Transplantation Proceedings*, 22(1):50-51 (1990).  
Sevier et al, *Clin. Chem.*, 27(11):1797-1806 (1981).  
Yelton et al, *American Scientist*, 68:510-516 (1980).

(Continued)

*Primary Examiner*—Long V. Le  
*Assistant Examiner*—Shafiquel Haq  
(74) *Attorney, Agent, or Firm*—Sughrue Mion, PLLC(57) **ABSTRACT**Provided are rapamycin conjugates which are useful as  
immunogenic molecules for the generation of antibodies  
specific for rapamycin, for measuring levels of rapamycin or  
derivatives thereof; for isolating rapamycin binding pro-  
teins; and detecting antibodies specific for rapamycin or  
derivatives thereof. This invention also provides a rapamy-  
cin specific monoclonal antibody.**11 Claims, No Drawings**

OTHER PUBLICATIONS

Campbell et al, *Monoclonal Antibody and Immunosensor Technology*, Elsevier, Chapter 1 (1991).  
Linskins et al, *Immunology in Plant Sciences*, pp. 86-141 (1986).  
Holt et al, *Chemical Abstracts*, vol. 120, No. 25 (1994).  
Venzina et al, *J. Antibiot.*, 28:721 (1975).  
Sehgal, *J. Antibiot.*, 28:727 (1975).

Baker, *Antibiot.*, 31:539 (1978).  
Martel, *Can. J. Physiol. Pharmacol.*, 55:48 (1997).  
Staruch, *FASEB*, 3:3411 (1989).  
Dumont, *FASEB*, 3:5256 (1989).  
Calne, *Lancet*, p. 1183 (1978).

\* cited by examiner

## RAPAMYCIN CONJUGATES

This application is a Divisional of U.S. patent application Ser. No. 09/576,951, filed May 24, 2000; which in turn is a Continuation-in-part of U.S. patent application Ser. No. 08/424,983, filed Apr. 19, 1995 (now abandoned); which in turn is a Continuation of U.S. patent application Ser. No. 08/224,205, filed Apr. 14, 1994 (now abandoned); which in turn is a Continuation-In-Part of U.S. patent application Ser. No. 08/053,030, filed Apr. 23, 1993 (now abandoned). The disclosure of each of which is incorporated herein by reference.

## BACKGROUND OF THE INVENTION

This invention relates to derivatives of rapamycin which are useful as immunogenic molecules for the generation of antibodies specific for rapamycin, for measuring levels of rapamycin or derivatives thereof; for isolating rapamycin binding proteins; and detecting antibodies specific for rapamycin or derivatives thereof.

Rapamycin is a macrocyclic triene antibiotic produced by *Streptomyces hygroscopicus*, which was found to have antifungal activity, particularly against *Candida albicans*, both in vitro and in vivo [C. Vezina et al., J. Antibiot. 28, 721 (1975); S. N. Sehgal et al., J. Antibiot. 28, 727 (1975); H. A. Baker et al., J. Antibiot. 31, 539 (1978); U.S. Pat. No. 3,929,992; and U.S. Pat. No. 3,993,749].

Rapamycin alone (U.S. Pat. No. 4,885,171) or in combination with picibanil (U.S. Pat. No. 4,401,653) has been shown to have antitumor activity. R. Martel et al. [Can. J. Physiol. Pharmacol. 55, 48 (1977)] disclosed that rapamycin is effective in the experimental allergic encephalomyelitis model, a model for multiple sclerosis; in the adjuvant arthritis model, a model for rheumatoid arthritis; and effectively inhibited the formation of IgE-like antibodies.

The immunosuppressive effects of rapamycin have been disclosed in FASEB 3, 3411 (1989). Cyclosporin A and FK-506, other macrocyclic molecules, also have been shown to be effective as immunosuppressive agents, therefore useful in preventing transplant rejection [FASEB 3, 3411 (1989); FASEB 3, 5256 (1989); R. Y. Calne et al., Lancet 1183 (1978); and U.S. Pat. No. 5,100,899].

Rapamycin has also been shown to be useful in preventing or treating systemic lupus erythematosus [U.S. Pat. No. 5,078,999], pulmonary inflammation [U.S. Pat. No. 5,080,899], insulin dependent diabetes mellitus [Fifth Int Conf. Inflamm. Res. Assoc. 121 (Abstract), (1990)], adult T-cell leukemia/lymphoma [European Patent Application 525,960 A1], and smooth muscle cell proliferation and intimal thickening following vascular injury [Morris, R. J. Heart Lung Transplant 11 (pt 2): 197 (1992)].

Mono- and diacylated derivatives of rapamycin (esterified at the 28 and 43 positions) have been shown to be useful as antifungal agents (U.S. Pat. No. 4,316,885) and used to make water soluble prodrugs of rapamycin (U.S. Pat. No. 4,650,803). Recently, the numbering convention for rapamycin has been changed; therefore according to Chemical Abstracts nomenclature, the esters described above would be at the 31- and 42-positions. U.S. Pat. No. 5,100,883 discloses fluorinated esters of rapamycin. U.S. Pat. No. 5,118,677 discloses amide esters of rapamycin. U.S. Pat. No. 5,118,678 discloses carbamates of rapamycin. U.S. Pat. No. 5,130,307 discloses aminoesters of rapamycin. U.S. Pat. No. 5,177,203 discloses sulfonates and sulfamates of rapamycin. U.S. Pat. No. 5,194,447 discloses sulfonylcarbamates of


rapamycin. PCT Publication WO 92/05179 discloses carboxylic acid esters of rapamycin.

Yatscoff has reported that rapamycin levels can be quantitated using HPLC method with a sensitivity of 1 ng/ml [Ther. Drug Monitoring 14: 138 (1992)] This method is time consuming and each sample must be assayed individually.

Immunoassays have been developed for numerous proteins as well as various drugs including cyclosporin A [Morris, R. G., Ther. Drug Monitoring 14: 226-(1992)], and FK506 [Tamura, Transplant Proc. 19: 23 (1987); Cadoff, Transplant Proc. 22: 50 (1990)]. Numerous types of immunoassays, that have been developed to measure proteins or compounds, have been based on competitive inhibition, dual antibodies, receptor-antibody interactions, antigen capture, dipstick, antibody or receptor trapping, or on affinity chromatography. Affinity columns with rapamycin have been reported in which a rapamycin analog was covalently attached to a matrix [Fretz J. Am. Chem. Soc. 113: 1409 (1991)]. These columns have been used to isolate rapamycin binding proteins.

## DESCRIPTION OF THE INVENTION

This invention provides a rapamycin conjugate of formula I, having the structure



wherein  $R^1$  and  $R^2$  are each, independently, hydrogen or  $-(R^3-L-R^4)_a-$ ;

L is a linking group;

$R^3$  is selected from the group consisting of carbonyl,  $-S(O)-$ ,  $-S(O)_2-$ ,  $-P(O)_2-$ ,  $-P(O)(CH_3)-$ ,  $-C(S)-$ , and  $-CH_2C(O)-$ ;

$R^4$  is selected from the group consisting of carbonyl,  $-NH-$ ,  $-S-$ ,  $-CH_2-$ , and  $-O-$ ;

$a=1-5$ ;

$x=0-1$ ;

$y=0-1$ ;


$z$  is from about 1 to about 120;

and Carrier is immunogenic carrier material, detector carrier material, or a solid matrix, or a salt thereof with the proviso that  $R^1$  and  $R^2$  are both not hydrogen; and further provided that when  $a$  is greater than 1, each L group can be the same or different; and still further provided that  $x$  is 0 if  $R^1$  is hydrogen and  $y$  is 0 if  $R^2$  is hydrogen, and if  $x$  and  $y$  are both 1, the Carrier moiety is the same in both cases.

## 3

The linking group, L, is any moiety that contains the group R<sup>3</sup> on one end and R<sup>4</sup> on other end therefore enabling the linking group to be connected to the 42- and/or 31-hydroxyl groups of rapamycin on one end and connected to another linking group or the immunogenic carrier material, detector material, or matrix on the other end. When a is greater than 1, each L group can be the same or different. In such cases, the first L group is designated as L<sup>1</sup>, the second L group designated as L<sup>2</sup> and so on. The rapamycin conjugates of the present invention may be prepared in such ways as to encompass a wide range of linking groups (L) and terminal functional groups R<sup>4</sup>. For example, L may be linear or branched alkyls comprising from 1 to as many as 15, more usually 10 or less, and normally less than 6 carbon atoms (i.e., methylene, ethylene, n-propylene, iso-propylene, n-butylene, and so forth). In addition, such alkyls can contain other substituent groups such as cyano, amino (including substituted amino), acylamino, halogen, thiol, hydroxyl, carbonyl groups, carboxyl (including substituted carboxyls such as esters, amides, and substituted amides). The linking group L can also contain or consist of substituted or unsubstituted aryl, aralkyl, or heteroaryl groups (e.g., phenylene, phenethylene, and so forth). Additionally, such linkages can contain one or more heteroatoms selected from nitrogen, sulfur and oxygen in the form of ether, ester, amido, amino, thio ether, amidino, sulfone, or sulfoxide. Also, such linkages can include unsaturated groupings such as olefinic or acetylenic bonds, disulfide, imino, or oximino groups. Preferably L will be a chain, usually aliphatic comprising between 1 and about 20 atoms, more usually between 1 and 10, excluding hydrogen, of which between 0 and 5 are heteroatoms preferably selected from nitrogen, oxygen, and sulfur. Therefore, the choice of linking group L is not critical to the present invention and may be selected by one of ordinary skill taking normal precautions to assure that stable compounds are produced.

A preferred embodiment of this invention provides a conjugate of formula II, having the structure



R<sup>1</sup> and R<sup>2</sup> are each, independently, hydrogen or —R<sup>3</sup>-L-R<sup>4</sup>—;

L is —A-(CR<sup>5</sup>R<sup>6</sup>)<sub>b</sub>[B-(CR<sup>7</sup>R<sup>8</sup>)<sub>d</sub>]-E—

A is —CH<sub>2</sub>— or —NR<sup>9</sup>—;

## 4

B is —O—, —NR<sup>9</sup>—, —S—, —S(O)—, or —S(O)<sub>2</sub>—;

R<sup>3</sup> is selected from the group consisting of carbonyl, —S(O)—, —S(O)<sub>2</sub>—, —P(O)<sub>2</sub>—, —P(O)(CH<sub>3</sub>)—, —C(S)—, and —CH<sub>2</sub>C(O)—;

R<sup>4</sup> is selected from the group consisting of carbonyl, —NH—, —S—, —CH<sub>2</sub>—, and —O—;

R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, and R<sup>8</sup> are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, halo, hydroxy, trifluoromethyl, arylalkyl of 7-10 carbon atoms, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-4 carbon atoms, alkoxy of 1-6 carbon atoms, carbalkoxy of 2-7 carbon atoms, cyano, amino, —CO<sub>2</sub>H, or phenyl which is optionally mono-, di-, or tri-substituted with a substituent selected from alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, hydroxy, cyano, halo, nitro, carbalkoxy of 2-7 carbon atoms, trifluoromethyl, amino, or —CO<sub>2</sub>H;

R<sup>9</sup> is hydrogen, alkyl of 1-6 carbon atoms, or aralkyl of 7-10 carbon atoms;

b=0-10;

d=0-10;

e=0-2;


x=0-1;

y=0-1;

z is from about 1 to about 120;

and Carrier is immunogenic carrier material, detector carrier material, or a solid matrix, or a salt thereof with the proviso that R<sup>1</sup> and R<sup>2</sup> are both not hydrogen; and further provided that when b is greater than 1, each of the CR<sup>5</sup>R<sup>6</sup> groups can be the same or different, and when d is greater than 1, each of the CR<sup>7</sup>R<sup>8</sup> groups can be the same or different; and still further provided that x is 0 if R<sup>1</sup> is hydrogen and y is 0 if R<sup>2</sup> is hydrogen, and if x and y are both 1, the Carrier moiety is the same in both cases.

A second preferred embodiment of this invention provides a conjugate of formula III, having the structure





R<sup>1</sup> and R<sup>2</sup> are each, independently, hydrogen or —(R<sup>3</sup>-L<sup>1</sup>-R<sup>4</sup>)<sub>f</sub>-(R<sup>10</sup>-L<sup>2</sup>-R<sup>11</sup>)<sub>g</sub>-carrier—;

L<sup>1</sup> is —(CH<sub>2</sub>)<sub>h</sub>-CHR<sup>12</sup>—(CH<sub>2</sub>)<sub>j</sub>—;

L<sup>2</sup> is —(CH<sub>2</sub>)<sub>k</sub>-D-(CH<sub>2</sub>)<sub>m</sub>-E—;

5

D is  $-\text{CH}_2-$ ,  $-\text{S}-$ , orE is  $-\text{CH}_2-$  or

$\text{R}^3$  and  $\text{R}^{10}$  are each, independently, selected from the group consisting of carbonyl,  $-\text{S}(\text{O})-$ ,  $-\text{S}(\text{O})_2-$ ,  $-\text{P}(\text{O})_2-$ ,  $-\text{P}(\text{O})(\text{CH}_3)-$ ,  $-\text{C}(\text{S})-$ , and  $-\text{CH}_2\text{C}(\text{O})-$ ;

$\text{R}^4$  and  $\text{R}^{11}$  are each, independently, selected from the group consisting of carbonyl,  $-\text{NH}-$ ,  $-\text{S}-$ ,  $-\text{CH}_2-$ , and  $-\text{O}-$ ;

$\text{R}^{12}$  is hydrogen, alkyl of 1-6 carbon atoms, arylalkyl of 7-10 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms,  $-(\text{CH}_2)_n\text{CO}_2\text{R}^{13}$ ,  $-(\text{CH}_2)_p\text{NR}^{14}\text{R}^{15}$ , carbamylalkyl of 2-3 carbon atoms, aminoalkyl of 1-4 carbon atoms, hydroxyalkyl of 1-4 carbon atoms, guanylaalkyl of 2-4 carbon atoms, mercaptoalkyl of 1-4 carbon atoms, alkylthioalkyl of 2-6 carbon atoms, indolylmethyl, hydroxyphenylmethyl, imidazolylmethyl, halo, trifluoromethyl, or phenyl which is optionally mono-, di-, or tri-substituted with a substituent selected from alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, hydroxy, cyano, halo, nitro, carbalkoxy of 2-7 carbon atoms, trifluoromethyl, amino, or  $-\text{CO}_2\text{H}$ ;

$\text{R}^{14}$ , and  $\text{R}^{15}$  are each, independently, hydrogen, alkyl of 1-6 carbon atoms, or arylalkyl of 7-10 carbon atoms;

$\text{R}^{13}$  is hydrogen, alkyl of 1-6 carbon atoms, arylalkyl of 7-10 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenyl which is optionally mono-, di-, or tri-substituted with a substituent selected from alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, hydroxy, cyano, halo, nitro, carbalkoxy of 2-7 carbon atoms, trifluoromethyl, amino, or  $-\text{CO}_2\text{H}$ ;

f=0-3;

g=0-1;

j=0-10;

k=0-10;

m=0-10;

n=0-6;

p=0-6;

x=0-1;

y=0-1;


z is from about 1 to about 120;

and Carrier is immunogenic carrier material, detector carrier material, or a solid matrix, or a salt thereof with the proviso that  $\text{R}^1$  and  $\text{R}^2$  are both not hydrogen; and further provided that f and g are both not 0 and when f is greater than 1, each of the  $-(\text{R}^3-\text{L}^1-\text{R}^4)-$  moieties can be the same or different; and still further provided that x is 0 if  $\text{R}^1$  is hydrogen and y is 0 if  $\text{R}^2$  is hydrogen, and if x and y are both 1, the Carrier moiety is the same in both cases.

6

This invention also provides a conjugate of formula IV, having the structure

IV



wherein  $\text{R}^1$  is  $-\text{OCH}_2(\text{CH}_2)_q\text{R}^4-$ ;

$\text{R}^4$  is selected from the group consisting of carbonyl,  $-\text{NH}-$ ,  $-\text{S}-$ ,  $-\text{CH}_2-$ , and  $-\text{O}-$ ;

q=0-6;

z is from about 1 to about 120;

and Carrier is immunogenic carrier material, detector carrier material, or a solid matrix, or a salt thereof.

The immunogenic carrier material can be selected from any of those conventionally known. In most cases, the carrier will be a protein or polypeptide, although other materials such as carbohydrates, polysaccharides, lipopolysaccharides, nucleic acids and the like of sufficient size and immunogenicity can likewise be used. For the most part, immunogenic proteins and polypeptides will have molecular weights between 5,000 and 10,000,000, preferably greater than 15,000 and more usually greater than 40,000. Generally, proteins taken from one animal species will be immunogenic when introduced into the blood stream of another species. Particularly useful proteins are those such as albumins, globulins, enzymes, hemocyanins, glutelins or proteins having significant non-proteinaceous constituents, e.g., glycoproteins, and the like. Further reference for the state-of-the-art concerning conventional immunogenic carrier materials and techniques for coupling haptens thereto may be had to the following: Parker, Radioimmunoassay of Biologically Active Compounds, Prentice-Hall (Englewood Cliffs, N.J., USA, 1976), Butler, J. Immunol. Meth. 7:1-24 (1975) and Pharmacol. Rev. 29(2):103-163 (1978); Weinryb and Shroff, Drug Metab. Rev. 10:P271-283 (1975); Broughton and Strong, Clin. Chem. 22:726-732 (1976); and Playfair et al., Br. Med. Bull. 30:24-31 (1974). Preferred immunogenic carrier materials for use in the present invention are ovalbumin and keyhole limpet hemocyanin. Particularly preferred for use in the present invention is ovalbumin. The detector carrier material can be a rapamycin-linking moiety conjugated to an enzyme such as horseradish peroxidase, alkaline phosphatase, luciferase, a fluorescent moiety such as fluorescein, Texas Red, or rhodamine, a chemiluminescent moiety, and the like. The solid matrix carrier material can be resin beads, an ELISA plate, glass beads as commonly used in a radioimmunoassay, plastic beads, solid matrix material typically used in a dipstick-type assay. When rapamycin is conjugated to a solid matrix, the resulting conjugate can be used in a dipstick

assay, as described in this disclosure, for the affinity purification of antibodies, or for isolating rapamycin binding proteins.

It should be noted that as used in the formulae above describing the specific rapamycin conjugates, *z* represents the number of rapamycin conjugated to the carrier material. The value *z* is sometimes referred to as the epitopic density of the immunogen, detector, or solid matrix and in the usual situation will be on the average from about 1 to about 120 and more typically from 1 to 50. The densities, however, may vary greatly depending on the particular carrier material used.

When any of the compounds of this invention contain an aryl or arylalkyl moiety, it is preferred that the aryl portion is a phenyl, naphthyl, pyridyl, quinolyl, isoquinolyl, quinoxalyl, thienyl, thionaphthyl, furyl, benzofuryl, benzodioxyl, benzoxazolyl, benzoisoxazolyl, or benzodioxolyl group that may be optionally mono-, di-, or tri-substituted with a group selected from alkyl of 1-6 carbon atoms, arylalkyl of 7-10 carbon atoms, alkoxy of 1-6 carbon atoms, cyano, halo, nitro, carbalkoxy of 2-7 carbon atoms, trifluoromethyl, amino, dialkylamino of 1-6 carbon atoms per alkyl group, alkylthio of 1-6 carbon atoms, —SO<sub>3</sub>H, —PO<sub>3</sub>H, and —CO<sub>2</sub>H. It is more preferred that the aryl moiety is a phenyl group that is optionally mono-, di-, or tri-substituted with a group selected from alkyl of 1-6 carbon atoms, arylalkyl of 7-10 carbon atoms, alkoxy of 1-6 carbon atoms, cyano, halo, nitro, carbalkoxy of 2-7 carbon atoms, trifluoromethyl, amino, dialkylamino of 1-6 carbon atoms per alkyl group, alkylthio of 1-6 carbon atoms, —SO<sub>3</sub>H, —PO<sub>3</sub>H, and —CO<sub>2</sub>H.

The salts are those derived from such inorganic cations such as sodium, potassium, and the like; organic bases such as: mono-, di-, and trialkyl amines of 1-6 carbon atoms, per alkyl group and mono-, di-, and trihydroxyalkyl amines of 1-6 carbon atoms per alkyl group, and the like; and organic and inorganic acids as: acetic, lactic, citric, tartaric, succinic, maleic, malonic, gluconic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, and similarly known acceptable acids.

The compounds of this invention can be prepared by reacting the 42- and/or 31-hydroxyl groups of rapamycin with a suitable electrophilic reagent that will serve as the linker moiety. The following patents exemplify the preparation of the 42- and/or 31-derivatives of rapamycin that can be used as linking groups for the preparation of the compounds of this invention. The preparation of fluorinated esters of rapamycin is described in U.S. Pat. No. 5,100,883. The preparation of amide esters is disclosed in U.S. Pat. No. 5,118,677. The preparation of carbamates of rapamycin is disclosed in U.S. Pat. No. 5,118,678. The preparation of aminoesters of rapamycin is described in U.S. Pat. No. 5,130,307. The preparation of sulfonates and sulfamates of rapamycin are described in U.S. Pat. No. 5,177,203. The preparation of sulfonylcarbamates of rapamycin are described in U.S. Pat. No. 5,194,447. The disclosures of the above cited U.S. patents are hereby incorporated by reference. From these patents, it can be seen that reactive electrophiles such as isocyanates, used in the preparation of carbamates, or sulfonyl chlorides, used in the preparation of sulfonates, can be reacted with the hydroxyl groups of rapamycin without the need for an activating agent. For the esterification of the rapamycin hydroxyl groups with a carboxylic acid, activation is usually required through the use of a coupling reagent such as DCC, or a water soluble analog thereof, such as dimethylaminopropyl-3-ethyl carbodiimide (DAEC). Representative examples of the preparation of rapamycin-linking group moieties are provided as

examples below. The preparation of ether derivatives of rapamycin can be accomplished using the methodology disclosed in Example 18.

For the compounds of this invention in which the linker group is attached to the 42- or the 31,42-hydroxyls, the electrophile (or activated electrophile) is reacted with rapamycin to typically provide a mixture of the 42- and 31,42-derivatized rapamycin that can be separated by chromatography. For the compounds of this invention in which the linker group is attached to the 31-hydroxyl of rapamycin, the 42-hydroxyl group must be protected with a suitable protecting group, such as with a tert-butyldimethyl silyl group. The 31-hydroxyl can then be reacted with a suitable electrophile to provide the derivatized rapamycin, followed by deprotection of the 42-hydroxyl group. The preparation of 42-O-silyl ethers of rapamycin and subsequent deprotection is described in U.S. Pat. No. 5,120,842, which is hereby incorporated by reference. Preparation of compounds containing different linkers at the 31- and 42-positions can be accomplished by first preparing the 42-derivatized compound and then using a different linker to derivatize the 31-position. The preparation of the 27-oxime linking groups can be accomplished using the methodology disclosed in U.S. Pat. No. 5,023,264, which is hereby incorporated by reference; and as described in Example 21.

The linker group attached to rapamycin can be coupled to a second linker group using standard methodology described in the peptide literature; typically by activating the electrophilic moiety, with DCC type coupling reagent, or with N-hydroxysuccinimide, or as an activated ester or anhydride. The activated electrophilic end of one linking moiety can then be reacted with the nucleophilic end of the other linker moiety.

The coupling of the rapamycin linking group moiety to the immunogenic carrier can be accomplished under standard literature conditions. In general, for reaction with a nucleophilic group on the immunogenic carrier material, an electrophilic moiety, such as a carboxylic acid, on the linking group is activated with a suitable activating agent such as N-hydroxysuccinimide, and then reacted with the nucleophilic moiety on the immunogenic carrier material. Examples 2 and 3 specifically exemplify this technique. Similar methodology is employed for the coupling of a nucleophilic moiety on the linking group to an electrophilic moiety on the immunogenic carrier material. In such cases, the electrophilic moiety on the immunogenic carrier material is activated as described above, and then reacted with the nucleophilic end of the linking group.

The reagents used to prepare the compounds of the invention are commercially available or can be prepared by methods that are disclosed in the literature.

This invention also covers analogous conjugates of other rapamycins such as, but not limited to, 29-demethoxyrapamycin, [U.S. Pat. No. 4,375,464, 32-demethoxyrapamycin under C.A. nomenclature]; rapamycin derivatives in which the double bonds in the 1-, 3-, and/or 5-positions have been reduced [U.S. Pat. No. 5,023,262]; 42-oxorapamycin [U.S. Pat. No. 5,023,262]; 27-oximes of rapamycin [U.S. Pat. No. 5,023,264]; 27-hydrazones of rapamycin [U.S. Pat. No. 5,120,726]; 29-desmethylrapamycin [U.S. Pat. No. 5,093,339, 32-desmethylrapamycin under C.A. nomenclature]; 7,29-bisdesmethylrapamycin [U.S. Pat. No. 5,093,338, 7,32-desmethylrapamycin under C.A. nomenclature]; and 15-hydroxy- and 15,27-bishydroxy-rapamycin [U.S. Pat. No. 5,102,876]. The disclosures in the above cited U.S. patents are hereby incorporated by reference. Also covered are conjugates of the rapamycin 1,3-Diels Alder adduct with diethyl azidodicarboxylate and rapamycin 1,3-Diels Alder adduct with phenyltriazoline dione. The preparation of these compounds is described in Examples 14 and 15.

The compounds of this invention are rapamycin immunogen, detector, and matrix bound conjugates that are useful for the generation and detection of antibodies specific for rapamycin and derivatives thereof, for measuring levels of rapamycin or a derivative thereof in biological or laboratory fluids, and for isolating rapamycin binding proteins. Rapamycin derivatives as defined here are compounds containing a rapamycin nucleus in which one or more of the hydroxyl groups has been esterified into a carboxylic ester, a carbamate, a sulfonate ester, an amide, or the like, or one or more of the ketones has been reduced to a hydroxyl group, or one or more of the double bonds has been reduced, or one ketones has been converted to an oxime or a hydrazone. Other rapamycin derivatives for which the compounds of this invention can be used for measuring levels of or generating antibodies to will be apparent to one skilled in the art based on this disclosure.

Antibodies specific for rapamycin using the rapamycin immunogen conjugates of this invention may be generated by standard techniques that are known in the art. Typically, a host animal is inoculated at one or more sites with the immunogen conjugate, either alone or in combination with an adjuvant. The typical host mammals include, but are not limited to, mice, goats, rabbits, guinea pigs, sheep, or horses. Subsequent injections can be made until a sufficient titer of antibodies are produced. The antibodies generated from the rapamycin immunogen conjugates of this invention can be used in numerous immunoassays, for determining rapamycin levels, in ELISAs, radioimmunoassays, in chemiluminescence immunoassays, and in fluorescent immunoassays. Although many variations of the immunoassay can be used (antigen capture, antibody capture, competitive inhibition, or two antibody immunoassay), a basic competitive inhibition immunoassay can be performed as follows: Antibody specific for the ligand is usually bound to a matrix. A solution is applied to decrease nonspecific binding of the ligand to the matrix. After rinsing the excess away, the antibody coupled matrix may be treated in some cases so it can be stored. In a competitive inhibition assay, the ligand standard curve is made and added with the rapamycin detector conjugate to compete for binding to the rapamycin-specific antibody. If necessary, the excess is removed. The detector molecule is detected by the standard methods used by one skilled in the art. Different formats can be used, which include but are not limited to, dipstick assays, FPIA, EMIT, ELISA, VISTA, RIA, and MEIA. Detector conjugates of the present invention can be prepared to use in the above assays. For example, the detector conjugates can be Carrier material with labeled fluorescent, chemiluminescent, or enzymatic moieties.

This invention also provides for the use of the rapamycin immunogen conjugates or rapamycin specific antibodies in a test kit that can be commercially marketed. The test kit may be used for measuring levels of rapamycin in biological or laboratory fluids. Test kit components may include rapamycin antibodies, antisera, or rapamycin carrier conjugates. The conjugates or antibodies may be bound to a solid matrix, and rapamycin derivatives or antibodies may be radiolabeled if the assay so requires. Standard concentrations of rapamycin can be included so that a standard concentration curve can be generated. Suitable containers, microtiter plates, solid supports, test tubes, trays, can also be included in any such kit. Many variations of reagents can be included in the kit depending on the type of assay used.

The following is illustrative of the use of a rapamycin immunogen conjugate of this invention to generate rapamycin specific antibodies and detect them using an ELISA

format immunoassay. Five mice were immunized with 50 µg rapamycin 31,42-diester with glutaric acid conjugate with keyhole limpet hemocyanin in Complete Freund's Adjuvant intrasplenically and after about one month were boosted with 50 µg of rapamycin 31,42-diester with glutaric acid conjugate with keyhole limpet hemocyanin in incomplete Freund's Adjuvant into the footpads. Microtiter plates (IMMUNOLON I) were coated overnight with 100 µl of goat anti-mouse antibody (10 µg/ml in 10 mM potassium phosphate buffer, pH 7.2) at 4° C. The plates were flicked and blocked with 100 µl of 1% bovine sera albumin in phosphate buffered saline overnight at 4° C. After flicking and washing the plates thrice with 10 mM phosphate buffer, pH 7.05, 30 mM NaCl, 0.02% TRITON X-100 (polyethylene glycol tert-octylphenyl ether), and 0.004% thimerosal wash buffer, 100 µl of each mouse sera diluted with phosphate buffer solution was added to a well and incubated at room temperature for overnight. After flicking and washing the plates thrice with wash buffer, rapamycin 31,42-diester with glutaric acid conjugate with horseradish peroxidase (compound of Example 10 (100 µl, 0.5 ng/ml) was added and incubated for 1 hour at room temperature in the dark. After flicking and washing the plates thrice with wash buffer, tetramethyl benzidine (TMB) substrate with H2O2 was added and the plates were incubated covered for 30 mm. at room temperature in the dark. The optical density was read on a spectrophotometer at 450 nm. As shown in Table I, five of the five mice had antibodies reactive for rapamycin 31,42-diester with glutaric acid conjugate with horseradish peroxidase (compound of Example 10).

TABLE I

MOUSE #	DILUTION <sup>a</sup>	O.D.
6902	1/300	0.199
6903	1/100	0.231
6904	1/500	0.412
6905	1/100	0.121
6906	1/300	0.321
background	—	0.076

<sup>a</sup>Dilution of mouse sera in PBS

The results in Table 1 show that mouse 6904 produced the most antibodies to the compound of Example 10. Hybridomas were generated using standard methodology. Following a splenectomy of a mouse immunized and boosted 3 times with the compound of Example 4, spleen cells were fused to SP20 cells to produce hybridomas. The hybridomas were evaluated for the production of rapamycin specific antibodies using an ELISA assay as briefly described below.

Microtiter plates (IMMUNOLON I) were coated overnight with 100 µl of goat anti-mouse antibody (10 µg/ml in 10 mM potassium phosphate buffer, pH 7.2) at 4° C. The plates were flicked and blocked with 100 µl of 1% bovine sera albumin in phosphate buffered saline (PBS) overnight at 4° C. After flicking and washing the plates thrice with 0.2xPBS containing 0.02% TRITON X-100 and 0.004% thimerosal, 100 µl of each hybridoma supernatant was added to a well and incubated at room temperature for overnight. After flicking and washing the plates thrice with 0.2xPBS containing 0.02% TRITON X-100 and 0.004% thimerosal, the compound of Example 22 (100 µl, 0.17 µM) was added and incubated for 1 hour at 4° C. After flicking and washing the plates thrice with 0.2xPBS containing 0.02% TRITON X-100 and 0.004% thimerosal, streptavidin or avidin conjugated to horseradish peroxidase (100 µl, 0.2 µg/ml) was added and incubated at room temperature for 1 hour in the

dark. After flicking and washing the plates thrice with 0.2xPBS containing 0.02% TRITON X-100 and 0.004% thimerosal, TMB substrate and H<sub>2</sub>O<sub>2</sub> was added and the plates were incubated covered for 30 mm. at room temperature in the dark. The optical density was read on a spectrophotometer at 450 nm. An optical density reading of 0.25-3 indicates specific antibody binding. The results in Table 2 show that the hybridoma from well P4G1 is positive for binding to the compound of Example 22, and is therefore specific for rapamycin.

TABLE 2

Screening for Rapamycin Specific Monoclonal Antibodies	
WELL	OPTICAL DENSITY
P3H4	0.120
P3H5	0.105
P4G1	1.940

The hybridoma cell line in P4G1 was cloned by limiting dilution and is designated as hybridoma cell line, RAP-42-OVAF<sub>2</sub>#1hc. The rapamycin-specific antibody, designated as RAP-42-OVAF<sub>2</sub>#1MoAb, was isolated and purified using conventional methodology. Hybridoma RAP-42-OVAF<sub>2</sub>#1hc was deposited under the terms of the Budapest Treaty at the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209, USA, on Mar. 10, 1994, and was granted accession number NB 11568.

The compounds of Examples 12 and 13 can be used in an assay for the detection of polyclonal antibodies and monoclonal antibodies specific for rapamycin as described below.

Microtiter plates (IMMUNOLON I) were coated overnight with 100 µl of goat anti-mouse antibody (10 µg/ml in 10 mM potassium phosphate buffer, pH 7.2) at 4° C. The plates were flicked and blocked with 100 µl of 1% bovine sera albumin in phosphate buffered saline overnight at 4° C. After flicking and washing the plates thrice with wash buffer, 100 µl of rabbit sera diluted 1:5 in phosphate buffered saline was added to a well and incubated at room temperature for overnight. After flicking and washing the plates thrice with wash buffer, rapamycin 42-ester with 3-[3-(4-imino-butylthio)succinimidy]phenylglycine conjugate with horseradish peroxidase (compound of Example 12) (100 µl, 0.5 ng/ml) or rapamycin 42 ester with (N-(3-carboxyphenyl)-3-thiosuccinimidy]glycine conjugate with horseradish peroxidase (compound of Example 13) (100 µl, 0.5 ng/ml) was added and incubated for 1 hour at room temperature in the dark. After flicking and washing the plates thrice with wash buffer, TMB substrate with H<sub>2</sub>O<sub>2</sub> was added and the plates were incubated covered for 30 mm. at room temperature in the dark. The optical density was read on a spectrophotometer at 450 nm. The results are shown in Table III.

TABLE 3

Comparison of Anti-rapamycin Antibody Levels in Rabbits Immunized with the Compound of Example 3 vs. Naive Rabbits Using a Capture ELISA Assay				
Rabbit No.	Prebleed	AA450 (3rd Bleed-Prebleed)		
	Example 10	Example 10	Example 12	Example 13
81	0.119	0.713	0.217	0.114
89	0.136	0.037	0.026	0.020

The data in Table 3 show that the compounds of Examples 12 and 13 can be used to detect antibodies specific for rapamycin in an a mammal, as seen in rabbit number 81.

The following is an example of the measurement of rapamycin concentrations using a competitive inhibition assay for rapamycin with an ELISA format using an antibody specific for rapamycin. Microtiter plates (IMMUNOLON I) were coated overnight with 100 µl of goat anti-mouse antibody (10 µg/ml in 10 mM potassium phosphate buffer, pH 7.2) at 4° C. The plates were flicked and blocked with 100 µl of 1% bovine sera albumin in phosphate buffered saline overnight at 4° C. After flicking and washing the plates thrice with wash buffer, the rapamycin specific antibody described above (100 µl of 1 µg/ml) was added to each well and incubated at room temperature for 1-4 hour. After flicking and washing the plates thrice with wash buffer, rapamycin 31,42-bis(hemiglutarate) conjugate with horseradish peroxidase (100 µl, 0.5 ng/ml) was added and incubated for 1 hour at room temperature in the dark. After flicking and washing the plates thrice with wash buffer, TMB substrate was added and the plates were incubated covered for 5 mm at room temperature in the dark. The optical density was read on a spectrophotometer at 450 nm. Results of the competition between rapamycin and rapamycin 31,42-diester with glutaric acid conjugate with horseradish peroxidase binding to mouse sera are shown in Table 4. From these results, a standard curve can be constructed and the concentration of rapamycin in a sample can be determined.

TABLE 4

Free	OPTICAL DENSITY x1000			
RAPAMYCIN	1	2	Avg	% Inhibition
10 µM	158	158	158	74.1
5	182	194	188	69.2
0.5	304	322	313	48.6
0.05	494	501	498	18.4
0.005	528	546	537	11.9
0.0005	601	611	606	0.6
0	583	636	610	—

The compound of Example 11 (rapamycin 42-ester with N-[9H-fluoren-9-ylmethoxy]carbonyl]glycine) can be deprotected by the procedure used in Example 12 (to give rapamycin 42-ester with glycine) and conjugated to a solid matrix. It can bind rapamycin specific antibodies as used in some dipstick immunoassay methods or to isolate rapamycin binding proteins. The following example illustrates that 803 resonance units (RU) of the compound of Example 11 can be immobilized on a solid matrix using the BIAcore's standard protocol based on EDC and NHS used in a BIAcore. This matrix bound 1401 RU units of rapamycin specific antibody. The kinetics of association and dissociation were determined for each concentration of antibody tested (0.625, 1.25, 2.5, 5.0, 10.0 ug/ml). These data show that the compound of Example 11, even when bound to a matrix was accessible to binding by a rapamycin-specific antibody and the interaction could be characterized. Similar procedures can be used to bind a rapamycin-binding protein to deprotected rapamycin 42-ester with N-[9H-fluoren-9-ylmethoxy]carbonyl]glycine conjugated matrix. This matrix can also be used for the isolation of novel binding proteins, as practiced by one skilled in the art. Deprotected rapamycin 42-ester with N-[9H-fluoren-9-ylmethoxy]carbonyl]glycine can be used to isolate binding proteins of rapamycin-FKBP complex by one of the following methods. In one approach,



## 13

tissue or cell lysates containing the appropriate protease inhibitors are incubated with FKBP which has been incubated with a deprotected-rapamycin 42-ester with N-[9H-fluoren-9-ylmethoxy)carbonyl]glycine conjugated matrix for a sufficient time to allow binding. Various buffers are used to rinse the proteins which are nonspecifically bound. Proteins are released by the addition of additional buffers which disrupt the bond between the rapamycin nucleus-FKBP and the binding proteins.

The following examples represent the preparation of representative compounds of this invention.

## EXAMPLE 1

## Rapamycin 42-ester with Succinic Acid

1.1 g (11 mmol) of succinic anhydride and 400 mg of dimethylaminopyridine (DMAP) were added to a stirring solution of 5 g (5.5 mmol) of rapamycin and 880  $\mu$ l of pyridine in 15 ml of methylene chloride. The reaction mixture was stirred for 2 days at room temperature, diluted with methylene chloride and washed with three 50 ml portions of 1N HCl. The organic layer was then dried over  $\text{Na}_2\text{SO}_4$  and concentrated in vacuo affording crude product. Pure material was obtained by reverse phase HPLC with 55% acetonitrile/water as eluant affording 1 g (18%) of the title compound. Spectral data follows:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz) 4.650 (m, 1H,  $\text{H}_2\text{COC}=\text{O}$ ), 4.168 (d, 1H,  $\text{H}_2\text{COH}$ ), 2.795 (s, 4H,  $\text{OC}=\text{OCH}_2\text{CH}_2\text{C}=\text{O}$ ).

## EXAMPLE 2

Rapamycin 42-ester with  
(N-hydroxysuccinimide(hemisuccinate))

21 mg (0.098 mmol) of DCC and 12 mg (0.098 mmol) of N-hydroxysuccinimide were added to a stirring solution of 100 mg of rapamycin 42-ester with succinic acid in 3 ml ethyl acetate. The reaction mixture was stirred overnight at room temperature, filtered, and concentrated in vacuo affording crude product. Pure material was obtained by reverse phase HPLC with 80% acetonitrile/water as eluant affording 75 mg (69%) of the title compound. Spectral data follows:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz) 4.650 (m, 1H,  $\text{H}_2\text{COC}=\text{O}$ ), 4.168 (d, 1H,  $\text{H}_2\text{COH}$ ), 2.951 (m, 2H,  $\text{OC}=\text{OCH}_2$ ), 2.795 (m, 4H,  $\text{OC}=\text{OCH}_2\text{CH}_2\text{C}=\text{O}$ ), 2.705 (m, 2H,  $\text{OC}=\text{OCH}_2$ ); MS (neg.ion FAB) 1110 ( $\text{M}^-$ ), 1056, 1012, 913, 148 (100).

## EXAMPLE 3

Rapamycin 42-ester with Succinic Acid Conjugate  
with Keyhole Limpet Hemocyanin

197 mg of keyhole limpet hemocyanin in 6 ml of 0.05 M phosphate buffer was added to a stirring solution of 37 mg of rapamycin 42-ester with (N-hydroxysuccinimide(hemisuccinate)) in 3 ml of 1,4 dioxane and the reaction was left stirring for 3 days at  $4^\circ\text{C}$ . The reaction mixture was then dialyzed for 24 hr at  $4^\circ\text{C}$ . in 1500 ml of 0.05 M phosphate buffer to give the title compound which could be used without further purification. The number of rapamycin 42-ester with succinic acid moieties per keyhole limpet hemocyanin was approximately 42:1.

## 14

## EXAMPLE 4

Rapamycin 42-ester with Succinic Acid Conjugate  
with Ovalbumin

197 mg of ovalbumin in 6 ml of 0.05 M phosphate buffer was added to a stirring solution of 37 mg of rapamycin 42-ester with (N-hydroxysuccinimide(hemisuccinate)) in 3 ml of 1,4 dioxane and the reaction was left stirring for 3 days at  $4^\circ\text{C}$ . The reaction mixture was then dialyzed for 24 hr at  $4^\circ\text{C}$ . in 1500 ml of 0.05 M phosphate buffer to give the title compound which could be used without further purification.

## EXAMPLE 5

Rapamycin 42-ester with Succinic Acid Conjugate  
with Horseradish Peroxidase

16 mg of horseradish peroxidase in a solution of 0.4 ml of 1,4 dioxane and 0.4 ml of 0.5% sodium bicarbonate was added to 1 mg of rapamycin 42-ester with (N-hydroxysuccinimide(hemisuccinate)) in 40  $\mu$ l of 1,4 dioxane and the reaction left stir for 2.5 hr at  $4^\circ\text{C}$ . The reaction mixture was then dialyzed for 24 hr at  $4^\circ\text{C}$ . in 1500 ml of 0.05 M phosphate buffer to give the title compound which could be used without further purification.

## EXAMPLE 6

## Rapamycin 31,42 Diester with Glutaric Acid

The title compound was prepared according to the method used in Example 1.

## EXAMPLE 7

Rapamycin 31,42-diester with  
(N-hydroxysuccinimide(hemiglutarate))

To a solution of 15.9 mg of rapamycin 31,42-diester with glutaric acid in 160  $\mu$ L of dimethyl formamide was added 3.65 mg of N,N-dimethylaminopropyl-ethylcarbodiimide and 1.8 mg of N-hydroxysuccinimide. The reaction mixture was allowed to stir until reaction was complete, poured into water, and extracted with ethyl acetate. The organic layers were combined, dried over sodium sulfate, filtered, and concentrated in vacuo to give the title compound, which was stored at  $4^\circ\text{C}$ . at 0.1 N sodium phosphate buffer and used without further purification.

## EXAMPLE 8

Rapamycin 31,42-diester with Glutaric Acid  
Conjugate with Keyhole Limpet Hemocyanin

To 20 mg of keyhole limpet hemocyanin in 2 mL of 0.1 M  $\text{NaHCO}_3$  was added 55  $\mu$ L of rapamycin 31,42-diester with (N-hydroxysuccinimide(hemiglutarate)) at  $0^\circ\text{C}$ . in 10  $\mu$ L increments over a 30 min period. The solution was gently shaken until reaction was complete, centrifuged at 6000 rpm for 20 min, and unconjugated starting material was separated from the title compound on a G-25 column with phosphate buffer solution. The conjugate was mixed with glycerol at 50% and stored at  $-70^\circ\text{C}$ . The number of rapamycin 31,42-diester with glutaric acid moieties per keyhole limpet hemocyanin ranged from 17-45:1.

## 15

## EXAMPLE 9

Rapamycin 31,42-diester with Glutaric Acid  
Conjugate with Ovalbumin

To 20 mg of ovalbumin in 2 mL of 0.1 M NaHCO<sub>3</sub> was added 55 µL of rapamycin 31,42-diester with (N-hydroxysuccinimide(hemiglutarate)) at 0° C. in 10 µL increments over a 30 min period. The solution was gently shaken until reaction was complete, centrifuged at 6000 rpm for 20 min, and unconjugated starting material was separated from the title compound on a G-25 column with phosphate buffer solution. The conjugate was mixed with glycerol at 50% and stored at -70° C.

## EXAMPLE 10

Rapamycin 31,42-diester with Glutaric Acid  
Conjugate with Horseradish Peroxidase

To 10 mg of horseradish peroxidase in 1 mL of 0.1 M NaHCO<sub>3</sub> was added 105 µL of rapamycin 31,42-diester with (N-hydroxysuccinimide(hemiglutarate)) in 10 µL increments over a 30 min period. The solution was gently shaken until complete, centrifuged at 6000 rpm for 20 min, and eluted from a G-25 column with phosphate buffer solution. The conjugate was mixed with glycerol at 50% and stored at -20° C.

## EXAMPLE 11

Rapamycin 42-ester with  
N-[9H-fluoren-9-ylmethoxy]carbonyl]glycine

To a chilled (0° C.) solution of rapamycin (0.73 g, 0.08 mmol) in methylene chloride (5 mL) was added 0.6 g (1.19 mmol) of N-[9H-fluoren-9-ylmethoxy]carbonyl]glycine pentafluorophenyl ester, followed by pyridine (0.85 mL, 10.5 mmol) and dimethylaminopyridine (18 mg, 0.14 mmol) to form a heterogeneous solution, which became homogeneous upon warming to room temperature. The reaction mixture was stirred at room temperature overnight. A large excess of EtOAc was added. The organic layer was washed with 0.5 N HCl (2x) and brine, dried (MgSO<sub>4</sub>), and concentrated to yield an off-white foam. Flash chromatography (30-50% hexane/EtOAc) yielded the title compound in 71% yield (0.679 g, 0.57 mmol). Mass spec (negative ion FAB) M<sup>-</sup> at m/z 1192.

## EXAMPLE 12

Rapamycin 42-ester with 3-[3-(4-iminobutylthio)succinimidy]phenacyl]glycine Conjugate with Horseradish Peroxidase

To a solution of rapamycin 42-ester with N-[9H-fluoren-9-ylmethoxy]carbonyl]glycine (10 mg, 8.4 µmol) in acetonitrile (84 µL) was added 10 µL (in acetonitrile at 0.84 M) of diethylamine. The reaction mixture was stirred at room temperature for 60 minutes and the solvent was removed with a stream of nitrogen. The residue was dissolved in acetonitrile (100 µL) and washed with hexane (5 times, 200 µL), followed by concentration of the solvent with a nitrogen stream. The resulting rapamycin 42-ester with glycine was taken up in a solution of m-maleimidobenzoyl-N-hydroxysuccinimide (MBS) (2 mg) in DMF (200 µL) and allowed to incubate for two hours at 4° C., followed by the addition

## 16

of 50 nM ethanolamine (20 µL) in 50 mM Tris HCl, pH 8.0. Horseradish peroxidase (5 mg) and Rabbit IgG (10 mg) were treated with 2-iminothiolane and purified with Sephadex G-25, followed by the addition of the MBS-rapamycin glycine ester adduct. The mixture was incubated overnight at 4° C. and purified by gel filtration on Sephadex G-25 to provide the title compound.

## EXAMPLE 13

Rapamycin 42 Ester with  
(N-(3-carboxyphenyl)-3-thiosuccinimidy]-glycine  
Conjugate with Horseradish Peroxidase

To a solution of rapamycin 42-ester with N-[9H-fluoren-9-ylmethoxy]carbonyl]glycine (10 mg, 8.4 µmol) in acetonitrile (84 µL) was added 10 µL (in acetonitrile at 0.84 M) of diethylamine. The reaction mixture was stirred at room temperature for 60 minutes and the solvent was removed with a stream of nitrogen. The residue was dissolved in acetonitrile (100 µL) and washed with hexane (5 times, 200 µL), followed by concentration of the solvent with a nitrogen stream. The resulting rapamycin 42-ester with glycine was taken up in a solution of N-succinimidy] S-acetylthioacetate (2 mg) in DMF (200 µL). The reaction mixture was stirred at room temperature for 15 minutes and then at 4° C. overnight. A solution of hydroxylamine HCl (7 mg in 50 µL DMF) was added to the solution of rapamycin reaction mixture, incubated for one hour, followed by the addition of MBS-horseradish peroxidase adduct and MBS-Rabbit IgG to give the title compound which was purified by Sephadex G-25 gel filtration.

## EXAMPLE 14

Rapamycin 1,3, Diels Alder Adduct with Diethyl  
Azidodicarboxylate

Rapamycin (1 g, 1.093 mmol) and diethyl azodicarboxylate (0.381 g, 2.187 mmol) were dissolved in dichloromethane (10 ml) and heated at 65° C. overnight, TLC showed that the reaction was complete. The mixture was purified on a silica gel column using ethyl acetate as eluant to provide a white solid (0.750 g) which was triturated with hexane and air dried to give the title compound (0.666 g) as a powder.

Anal Calc for C<sub>57</sub>H<sub>89</sub>N<sub>3</sub>O<sub>17</sub>: C, 62.91; H, 8.24; N, 3.86. Found: C, 62.81; H, 8.12; N, 3.91. IR (KBr, cm<sup>-1</sup>) 3450, 1720 NMR (CDCl<sub>3</sub>) δ 6.15 (m, 1H), 5.20 (d, 1H), 3.40 (s, 3H), 3.30 (s, 3H), 3.15 (s, 3H), 0.9 (t, 3H), 0.72 (q, 1H) MS (-FAB) 1087 (M<sup>-</sup>)

## EXAMPLE 15

Rapamycin 1,3, Diels Alder Adduct with  
Phenyltriazolinedione

Rapamycin (0.66 g, 721 µmol) was dissolved in dichloromethane (10 ml) and cooled to 0° C. To this was added, dropwise, a solution of phenyltriazolinedione (0.133 g, 758 µmol) in dichloromethane (10 ml). The solution was stirred overnight, TLC showed the reaction was not complete. Additional phenyltriazolinedione (0.025 g, 27 µmol) was added. The reaction was purified using HPLC (4.1x31 cm, SiO<sub>2</sub>) with ethyl acetate as eluant to provide the title compound as a solid. The solid was triturated with 30 ml of hexane and 1 ml of ethyl acetate filtered and air dried to give the title compound as a powder (0.383 g).

## 17

Anal Calc for  $C_{59}H_{84}N_4O_{15}$ : C, 65.05; H, 7.77; N, 5.14. Found: C, 65.39; H, 7.98; N, 4.92. IR (KBr,  $cm^{-1}$ ) 3450, 1715 NMR (DMSO)  $\delta$  7.50 (m, 3H), 7.40 (m, 2H), 3.11 (s, 3H), 3.00 (s, 3H), 2.95 (s, 3H), 0.8 (q, 1H) MS (-FAB) 1088 ( $M^-$ )

The following are representative examples of fluorescent rapamycin derivatives that can be conjugated via a linker at the 31-position of rapamycin.

## EXAMPLE 16

## 42-Dansylrapamycin

Rapamycin (200 mg, 0.22 mmol) in dry pyridine (2 ml) was cooled to 0° C. and was treated with dansyl chloride (840 mg, 3.1 mmol). The reaction was warmed to room temperature and stirred for 24 hours. The reaction mixture was poured into cold 2N HCl (30 ml) and was extracted with ethyl acetate (4x25 ml). The ethyl acetate was pooled and washed with brine, dried over  $MgSO_4$ , filtered and concentrated in vacuo. The residue was chromatographed on silica with 25% ethyl acetate in benzene. This afforded 150 mg of the title compound as a yellow powder, mp 101-104° C.

## EXAMPLE 17

## Rapamycin 42-ester with Pyrene Butyric Acid

Rapamycin (459 mg, 0.5 mmol) and pyrenebutyric acid (216 mg, 0.75 mmol) were dissolved in THF/ $CH_2Cl_2$  (10 ml, 1:1). 1-(3-Dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (146 mg, 0.67 mmol) and 4-dimethylaminopyridine (15 mg) were added to the solution. The reaction was allowed to warm to room temperature over 15 hours. The reaction was diluted with  $CH_2Cl_2$  and washed with 5% HCl, then brine. The solution was dried over  $MgSO_4$ , filtered and evaporated to a solid. The solid was applied to a 3 mm silica gel Chromatron plate which was eluted with 50% ethyl acetate in hexane to provide 180 mg of the title compound as a foam. The reaction also afforded 100 mg of 31,42-diesterified rapamycin.

IR (KBr,  $cm^{-1}$ ) 3420, 1740 NMR ( $CDCl_3$ )  $\delta$  8.3 (d, 1H), 8.14 (dd, 2H), 8.10 (d, 2H), 7.85 (d, 1H), 3.34 (s, 3H), 3.30 (s, 3H), 3.11 (s, 3H) MS (-FAB) 1183 ( $M^-$ )

The following are representative examples of rapamycin derivatives that can be conjugated to immunogenic carriers by the procedures described above or can be connected to another linker and then conjugated.

## EXAMPLE 18

## Rapamycin 42-carbomethoxymethyl Ether and Rapamycin 42-bis(carbomethoxymethyl ether)

Rapamycin (2.0 g, 2.187 mmol) and rhodium (II) acetate (0.37 g, 0.08 mmol) were heated to reflux in benzene and treated with a solution ethyl diazoacetate (500 ml) in benzene (10 ml) over 10 minutes. The solution was cooled to room temperature and was stirred overnight. TLC showed that the reaction was incomplete. Two additional portions of ethyldiazoacetate (3 ml) were added at 24 hour intervals. The mixture was concentrated and purified by flash chromatography over silica using ethyl acetate. This provided the 42-monoether (1 g) and the 31,42 diether (0.850 g) as oils. The 42-monoether was triturated in a mixture of hexane, ethyl acetate and dichloromethane over the weekend to give the product as a powder. The diether was purified on HPLC

## 18

on a silica gel column with ethyl acetate as eluant. This provided the product as a solid.

Analytical data for the monoether:

Analysis Calc for  $C_{55}H_{85}NO_{15}$ : C, 66.04; H, 8.57; N, 1.40. Found: C, 65.29; H, 8.64; N, 1.60. IR (KBr,  $cm^{-1}$ ) 3420, 1715 NMR ( $CDCl_3$ )  $\delta$  4.82 (s, 1H), 3.41 (s, 3H), 3.33 (s, 3H), 3.13 (s, 3H), 1.28 (t, 3H), 0.70 (q, 1H) MS (-FAB) 999 ( $M^-$ )

Analytical data for the diether:

Analysis Calc for  $C_{59}H_{91}NO_{17}$ : C, 65.23; H, 8.44; N, 1.29. Found: C, 63.29; H, 8.40; N, 1.44. IR (KBr,  $cm^{-1}$ ) 1740 NMR ( $CDCl_3$ )  $\delta$  6.36 (q, 2H), 5.24 (s, 1H), 3.39 (s, 3H), 3.32 (s, 3H), 3.12 (s, 3H), 0.65 (q, 1H) MS (-FAB) 1085 ( $M^-$ )

## EXAMPLE 19

## Rapamycin 42-(4-nitrophenyl)carbonate and Rapamycin 31,42-bis(4-nitrophenyl)carbonate

Rapamycin (0.450 g, 0.49 mmol) was dissolved in dry dichloromethane (10 ml) and cooled to 0° C. To this solution was added pyridine (0.4 ml, 5.7 mmol) and a crystal of 4-dimethyl aminopyridine. A solution of 4-nitrophenyl chloroformate (0.3 g 1.49 mmol) in dichloromethane (3 ml) was added. The solution was allowed to warm to room temperature overnight and was stirred at room temperature for 24 hours. The reaction was quenched into 0.1N HCl (5 ml) and the aqueous layer was washed with dichloromethane. The organic layer was dried over  $MgSO_4$ , filtered, and evaporated in vacuo to afford a yellow solid. Chromatography over silica gel with 75% Ethyl acetate in hexane afforded 180 mg of the 42-monocarbonate and 47 mg of the 31,42-dicarbonate as yellow solids.

## EXAMPLE 20

## 42-O-(Phenoxythiocarbonyl)-rapamycin

Rapamycin (1.030 g, 1.12 mmol) was dissolved in dry dichloromethane (100 ml) and was cooled to 0° C. To this solution was added pyridine (0.27 ml, 3.33 mmol) and a crystal of 4-dimethyl aminopyridine. A solution of thiophenyl chloroformate (0.47 ml 1.49 mmol) in dichloromethane (5 ml) was added to the reaction mixture. The solution was allowed to warm to room temperature overnight and was stirred at room temperature for 24 hours. The reaction was quenched into 0.1N HCl (5 ml) and the aqueous layer was washed with dichloromethane. The organic layer was dried over  $MgSO_4$ , filtered and evaporated in vacuo to afford a yellow solid. Chromatography on a 4 mm silica gel Chromatron plate with a gradient of 40% to 70% ethyl acetate in hexane afforded 520 mg of the title compound as a yellow foam.

Analysis Calc for  $C_{58}H_{83}NOS_{14}$ : C, 66.32; H, 7.97; N, 1.33. Found: C, 66.48; H, 8.05; N, 1.12. IR (KBr,  $cm^{-1}$ ) 3420, 1715 NMR ( $CDCl_3$ )  $\delta$  7.41 (t, 1H), 7.25 (t, 2H), 7.12 (d, 1H), 3.45 (s, 3H), 3.33 (s, 3H), 3.13 (s, 3H) MS (-FAB) 1049 ( $M^-$ )

## EXAMPLE 21

## Rapamycin-O-carboxymethyl-27-oxime

To a solution of 600 mg (650  $\mu$ M) of rapamycin in 6 mL of methanol was added at room temperature, 100 mg (1.2 mmol) of anhydrous sodium acetate and 140 mg (660  $\mu$ M)

19

of carboxymethoxylamine hemihydrochloride. After stirring overnight at room temperature, the reaction was complete. The reaction mixture was concentrated in vacuo and the residue was triturated with water. The solids were filtered and washed thoroughly with water. The product was dried under high vacuum to give 575 mg (89.7%) of a white solid. <sup>13</sup>C and <sup>1</sup>H NMR indicated a mixture of E and Z isomers for the oxime derivative at position 27.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz): 3.43 and 3.41 (2s, 3H, CH<sub>3</sub>O), 3.30 (s, 3H, CH<sub>3</sub>O), 3.18 and 3.12 (2s, 3H, CH<sub>3</sub>O), 1.82 (s, 3H, CH<sub>3</sub>C=C), 1.695 and 1.633 (2s, 3H, CH<sub>3</sub>C=C); <sup>13</sup>C NMR (CDCl<sub>3</sub>, MHz): 215.8 (C=O), 211.5 (C=O), 194.5 (C=O), 191.0 (C=O), 172.5 (C=O), 169.0 (C=O), 168.5 (C=O), 167.0 (C=O), 161.5 (C=NOC), 160.0 (C=NOC), 140.0; MS (neg. ion FAB: 985 (M-H)<sup>-</sup>, 590, 167, 128, 97, 75 (100%)

Analysis Calcd for C<sub>53</sub>H<sub>82</sub>N<sub>2</sub>O<sub>15</sub>·0.15 H<sub>2</sub>O: C, 63.90; H, 8.40; N, 2.81. Found: C, 63.81; H, 8.41; N, 2.85.

The following compound was used in the generation of rapamycin specific antibodies.

## EXAMPLE 22

## Rapamycin 42-ester with Glycylbiotin

To a solution of biotin (0.83 g, 3.4 mmol) in 60 mL of DMF was added glycine t-butyl ester hydrochloride (0.57 g, 3.4 mmol), N-methylmorpholine (0.92 mL, 8.36 mmol), 1-hydroxybenzotriazole (0.61 g, 3.99 mmol) and 1-(3-Dimethylaminopropyl)-3-ethylcarbo-diimide hydrochloride (0.65 g, 3.4 mmol). The reaction mixture was stirred at room temperature for 7 days. The DMF was concentrated, ethyl acetate was added, and the organic layer was washed with water, 0.5 N HCl, saturated sodium bicarbonate, and brine. The ethyl acetate layer was dried (MgSO<sub>4</sub>) and concentrated to yield tert-butylglycylbiotin as a white solid which was primarily one spot on TLC (0.611 g, 1.71 mmol, 50%). Mass spec [M+H]<sup>+</sup> at m/z 358.


To a solution of tert-butylglycylbiotin (0.271 g, 0.758 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL) was added 0.5 mL trifluoroacetic acid. The reaction mixture was stirred at room temperature for 2 h, concentrated, and triturated with anhydrous diethyl ether. The off-white precipitate was collected to yield 0.209 g (0.694 mmol, 92%) of glycylbiotin. Mass spec [M+H]<sup>+</sup> at m/z 302.

To a solution of glycylbiotin (0.65 g, 2.16 mmol) in 1-methylpyrrolidinone (5 mL) was added 6 mL of CH<sub>2</sub>Cl<sub>2</sub>, causing a precipitate to form which persisted even after the addition of 0.33 mL (2.36 mmol) of triethylamine. To this heterogeneous solution was added 2 g (2.19 mmol) of rapamycin, 0.43 g (2.24 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbo-diimide hydrochloride, and 30 mg (2.46 mmol) of DMAP. After several hours, the reaction mixture became homogenous, and was stirred an additional four days. A large excess of ethyl acetate was added and the organic layer was washed with water, 0.5 N HCl, saturated sodium bicarbonate, and brine. The organic layer was dried (MgSO<sub>4</sub>) and concentrated. The light yellow foam was triturated with hot anhydrous diethyl ether to yield 1.2 g of impure title compound as a light yellow solid. A portion (0.5 g) of this material was flash chromatographed in 5% MeOH/CHCl<sub>3</sub>, and triturated again in hot ether to yield 87 mg of the title compound contaminated with a small amount of rapamycin. This material was rechromatographed (gradient 0-5% MeOH/CHCl<sub>3</sub>), and triturated a final time with ether to yield 34 mg (0.028 mmol) of pure title compound as a white solid. Mass spec, negative FAB M<sup>-</sup> at m/z 1196.

20

What is claimed is:

1. A rapamycin conjugate represented by formula I, or a salt thereof:



wherein R<sup>1</sup> and R<sup>2</sup> are each, independently, hydrogen or —(R<sup>3</sup>-L-R<sup>4</sup>)<sub>a</sub>—;

L is a linking group;

R<sup>3</sup> is selected from the group consisting of carbonyl, —S(O)—, —S(O)<sub>2</sub>—, —P(O)<sub>2</sub>—, —P(O)(CH<sub>3</sub>)—, —C(S)—, and —CH<sub>2</sub>C(O)—;

R<sup>4</sup> is selected from the group consisting of carbonyl, —NH—, —S—, —CH<sub>2</sub>—, and —O—;

a=1-5;

x=0-1;


y=0-1;

z is from about 1 to about 120;

and Carrier is immunogenic carrier material, detector carrier material, or a solid matrix,

with the proviso that R<sup>1</sup> and R<sup>2</sup> are both not hydrogen; and further provided that when a is greater than 1, each L group can be the same or different; and still further provided that x is 0 if R<sup>1</sup> is hydrogen and y is 0 if R<sup>2</sup> is hydrogen, and if x and y are both 1, the Carrier moiety is the same in both cases.

2. A rapamycin conjugate represented by formula II, or a salt thereof:



## 21

wherein R<sup>1</sup> and R<sup>2</sup> are each, independently, hydrogen or —R<sup>3</sup>-L-R<sup>4</sup>—;

$$L \text{ is } -A-(CR^5R^6)_b[B-(CR^7R^8)_d]_e-;$$

A is  $-\text{CH}_2-$  or  $-\text{NR}^9-$ ;

B is  $-\text{O}-$ ,  $-\text{NR}^9-$ ,  $-\text{S}-$ ,  $-\text{S}(\text{O})-$ , or  $-\text{S}(\text{O})_2-$ ; 5

R<sup>3</sup> is selected from the group consisting of carbonyl, —S(O)—, —S(O)<sub>2</sub>—, —P(O)<sub>2</sub>—, —P(O)<sub>2</sub>(CH<sub>3</sub>)—, —C(S)—, and —CH<sub>2</sub>C(O)—;

R<sup>4</sup> is selected from the group consisting of carbonyl, —NH—, —S—, —CH<sub>2</sub>—, and —O—;

R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, and R<sup>8</sup> are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, halo, hydroxy, trifluoromethyl, arylalkyl of 7-10 carbon atoms, aminoalkyl of

1-6 carbon atoms, hydroxyalkyl of 1-4 carbon atoms, alkoxy of 1-6 carbon atoms, carbalkoxy of 2-7 carbon atoms, cyano, amino, -CO<sub>2</sub>H, or phenyl which is optionally mono-, di-, or tri-substituted with a substituent selected from the group consisting of alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, hydroxy, cyano, halo, nitro, carbalkoxy of 2-7 carbon atoms, trifluoromethyl, amino, and —CO<sub>2</sub>H;

R<sup>9</sup> is hydrogen, alkyl of 1-6 carbon atoms, or aralkyl of 7-10 carbon atoms;

```
b=0-10;
```

d=0-10;

e=0-2;

```
x=0-1;
```


```
y=0-1;
```

z is from about 1 to about 120;

and Carrier is immunogenic carrier material, detector carrier material, or a solid matrix,

with the proviso that R<sup>1</sup> and R<sup>2</sup> are both not hydrogen; and further provided that when b is greater than 1, each of the CR<sup>5</sup>R<sup>6</sup> groups can be the same or different, and when d is greater than 1, each of the CR<sup>7</sup>R<sup>8</sup> groups can be the same or different; and still further provided that x is 0 if R<sup>1</sup> is hydrogen and y is 0 if R<sup>2</sup> is hydrogen, and if x and y are both 1, the Carrier moiety is the same in both cases.

3. A rapamycin conjugate represented by formula III, or a salt thereof:




wherein R<sup>1</sup> and R<sup>2</sup> are each, independently, hydrogen or —(R<sup>3</sup>-L<sup>1</sup>-R<sup>4</sup>)<sub>n</sub>—(R<sup>10</sup>-L<sup>2</sup>-R<sup>11</sup>)<sub>m</sub>—;

$$L^1 \text{ is } \text{---}(\text{CH}_2)_h\text{---CHR}^{12}\text{---}(\text{CH}_2)_i\text{---};$$


$L^2$  is  $-(CH_2)_k-D-(CH_2)_m-E$ ;

## 22

D is  $\text{—CH}_2\text{—}$ ,  $\text{—S—S—}$ , or



E is  $\text{—CH}_2\text{—}$  or  $\text{—C—}$ ;



R<sup>3</sup> and R<sup>10</sup> are each, independently, selected from the group consisting of carbonyl, —S(O)—, —S(O)<sub>2</sub>—, —P(O)<sub>2</sub>—, —P(O)(CH<sub>3</sub>)—, —C(S)—, and —CH<sub>2</sub>C(O)—;

$R^4$  and  $R^{11}$  are each, independently, selected from the group consisting of carbonyl,  $-NH-$ ,  $-S-$ ,  $-CH_2-$ , and  $-O-$ ;

R<sup>12</sup> is hydrogen, alkyl of 1-6 carbon atoms, arylalkyl of 7-10 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms,  $-(CH_2)_nCO_2R^{13}$ ,  $-(CH_2)_pNR^{14}R^{15}$ , carbamylalkyl of 2-3 carbon atoms, aminoalkyl of 1-4 carbon atoms, hydroxyalkyl of 1-4 carbon atoms, guanlylalkyl of 2-4 carbon atoms, mercaptoalkyl of 1-4 carbon atoms, alkylthioalkyl of 2-6 carbon atoms, indolylmethyl, hydroxyphenylmethyl, imidazolylmethyl, halo, trifluoromethyl, or phenyl which is optionally mono-, di-, or tri-substituted with a substituent selected from the group consisting of alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, hydroxy, cyano, halo, nitro, carbalkoxy of 2-7 carbon atoms, trifluoromethyl, amino, and  $-CO_2H$ ;

R<sup>14</sup> and R<sup>15</sup> are each, independently, hydrogen, alkyl of 1-6 carbon atoms, or arylalkyl of 7-10 carbon atoms;

R<sup>13</sup> is hydrogen, alkyl of 1-6 carbon atoms, arylalkyl of 7-10 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenyl which is optionally mono-, di-, or tri-substituted with a substituent selected from the group consisting of alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, hydroxy, cyano, halo, nitro, carbalkoxy of 2-7 carbon atoms, trifluoromethyl, amino, and —CO<sub>2</sub>H;

f=0-3;

g=0-1;

```
j=0-10;
```

```
k=0-10;
```

```
m=0-10;
```

n=0-6;

p=0-6;

```
x=0-1;
```

```
y=0-1;
```

z is from about 1 to about 120;

and Carrier is immunogenic carrier material, detector carrier material, or a solid matrix, with the proviso that  $R^1$  and  $R^2$  are both not hydrogen; and further provided that  $f$  and  $g$  are both not 0 and when  $f$  is greater than 1, each of the  $-(R^3-L^1-R^4)-$  moieties can be the same or different; and still further provided that  $x$  is 0 if  $R^1$  is hydrogen and  $y$  is 0 if  $R^2$  is hydrogen, and if  $x$  and  $y$  are both 1, the Carrier moiety is the same in both cases.

23

4. The rapamycin conjugate of claim 1, wherein said immunogenic carrier material is selected from the group consisting of keyhole limpet hemocyanin and ovalbumin.

5. The rapamycin conjugate of claim 2, wherein said immunogenic carrier material is selected from the group consisting of keyhole limpet hemocyanin and ovalbumin.


6. The rapamycin conjugate of claim 2, wherein said immunogenic carrier material is selected from the group consisting of keyhole limpet hemocyanin and ovalbumin.

7. An immunoassay kit for measuring the blood level of a rapamycin comprising a conjugate of any one of claims 1, 2 or 3, wherein the carrier is a detector carrier material.

8. The immunoassay kit of claim 1, wherein said detector carrier material is horseradish peroxidase.

9. The rapamycin conjugate of claim 1, wherein in formula I,  $R^1$  and  $R^2$  are each  $-(R^3-L-R^4)_a-$ ;  $R^3$  is carbonyl; L is a  $CH_2CH_2$  linking group;  $R^4$  is a carbonyl, a is 1, x is 1, and y is 0.

10. The conjugate of claim 3, wherein in formula III,  $R^1$  is  $(R^3-L^1-R^4)_f-(R^{10}-L^2-R^{11})_g-$ ,  $R^2$  is hydrogen,  $R^3$  is carbonyl;  $R^4$  is NH, and h is 0,  $R^{12}$  is hydrogen, j is 0, and f is 1;  $R^{10}$  is carbonyl is 0, D is



m is 3, E is




and  $R^{11}$  is NH, g is 1, x is 1 and y is 0, and Carrier is horseradish peroxidase.


24

11. A conjugate represented by formula III

III



wherein  $R^1$  is  $(R^3-L^1-R^4)_f-(R^{10}-L^2-R^{11})_g-$ ,  $R^2$  is hydrogen,  $R^3$  is carbonyl, f is 1 and  $L^1$  is  $(CH_2)_h-CHR^{12}-$   $(CH_2)_j$ , h is 0,  $R^{12}$  is hydrogen, j is 0;  $R^4$  is NH,  $R^{10}$  is carbonyl;  $L^2$  is  $(CH_2)_k-D-(CH_2)_m-E$ , k is 1, D is



m is 0, E is carbonyl,  $R^{11}$  is NH, g is 1, x is 1 and y is 0, and Carrier is horseradish peroxidase.

\* \* \* \* \*

专利名称(译)	雷帕霉素缀合物		
公开(公告)号	<a href="#">US7279562</a>	公开(公告)日	2007-10-09
申请号	US10/124386	申请日	2002-04-18
[标]申请(专利权)人(译)	MOLNAR KIMBER凯瑟琳大号 CAUFIELD CRAIGê OCAIN TIMOTHYĐ		
申请(专利权)人(译)	MOLNAR-KIMBER凯瑟琳L. CAUFIELD CRAIG E. OCAIN TIMOTHY D.		
当前申请(专利权)人(译)	惠氏		
[标]发明人	MOLNAR KIMBER KATHERINE L CAUFIELD CRAIG E OCAIN TIMOTHY D		
发明人	MOLNAR-KIMBER, KATHERINE L. CAUFIELD, CRAIG E. OCAIN, TIMOTHY D.		
IPC分类号	G01N1/04 A61K31/435 A61K47/48 C07K16/12 G01N33/533		
CPC分类号	A61K31/435 A61K47/48284 A61K47/48507 C07K16/1292 C12N9/0065 G01N33/533 G01N33/9446 C07K16/14 Y10S530/809 A61K47/643 A61K47/6835		
其他公开文献	US20020151088A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

## 摘要(译)

本发明提供雷帕霉素缀合物，其可用作产生雷帕霉素特异性抗体的免疫原性分子，用于测定雷帕霉素或其衍生物的水平;用于分离雷帕霉素结合蛋白;检测对雷帕霉素或其衍生物特异的抗体。本发明还提供雷帕霉素特异性单克隆抗体。

