



US009169517B2

(12) **United States Patent**
Petit et al.

(10) **Patent No.:** **US 9,169,517 B2**
(45) **Date of Patent:** ***Oct. 27, 2015**

- (54) **MUTATION WITHIN THE CONNEXIN 26 GENE RESPONSIBLE FOR PRELINGUAL NON-SYNDROMIC DEAFNESS AND METHOD OF DETECTION**
- (71) Applicant: **INSTITUT PASTEUR**, Paris Cedex (FR)
- (72) Inventors: **Christine Petit**, Le Plessis-Robinson (FR); **Francoise Denoyelle-Gryson**, Arcueil (FR); **Dominique Weil**, Paris (FR); **Sandrine Marlin**, La Gareene-Colombes (FR); **Jean-Luc Guesdon**, Sevres (FR)

5,661,122	A	8/1997	Clark et al.
5,769,796	A	6/1998	Palermo et al.
5,855,577	A	1/1999	Murphy-Churtorian et al.
5,876,373	A	3/1999	Giba et al.
5,967,991	A	10/1999	Gardineer et al.
5,998,147	A	12/1999	Petit et al.
6,053,870	A	4/2000	Fulton, III
6,053,899	A	4/2000	Slanda et al.
6,106,473	A	8/2000	Violante et al.
6,146,339	A	11/2000	Biagtan et al.
6,165,139	A	12/2000	Damadian
6,165,178	A	12/2000	Bashiri et al.
6,179,809	B1	1/2001	Khairkhahan et al.
6,238,415	B1	5/2001	Sepetka et al.
6,248,112	B1	6/2001	Gambale et al.
6,277,082	B1	8/2001	Gambale

- (73) Assignee: **INSTITUT PASTEUR**, Paris (FR)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- This patent is subject to a terminal disclaimer.

- (21) Appl. No.: **13/899,963**
- (22) Filed: **May 22, 2013**

- (65) **Prior Publication Data**
US 2014/0113285 A1 Apr. 24, 2014

- Related U.S. Application Data**
- (63) Continuation of application No. 13/430,062, filed on Mar. 26, 2012, now Pat. No. 8,455,195, which is a continuation of application No. 11/826,141, filed on Jul. 12, 2007, now Pat. No. 8,143,000, which is a continuation of application No. 10/278,089, filed on Oct. 23, 2002, now Pat. No. 7,258,975, which is a continuation of application No. 09/485,415, filed as application No. PCT/EP98/05175 on Aug. 14, 1998, now Pat. No. 6,485,908.
- (60) Provisional application No. 60/055,863, filed on Aug. 15, 1997.

- (51) **Int. Cl.**
C12Q 1/68 (2006.01)
C12P 19/34 (2006.01)
- (52) **U.S. Cl.**
CPC *C12Q 1/6883* (2013.01); *C12Q 1/6886* (2013.01); *C12Q 2600/156* (2013.01)

- (58) **Field of Classification Search**
None
See application file for complete search history.

- (56) **References Cited**
U.S. PATENT DOCUMENTS

4,228,805	A	10/1980	Rosen et al.
4,709,703	A	12/1987	Lazarow et al.
5,190,050	A	3/1993	Nitzsche
5,312,415	A	5/1994	Palermo
5,358,479	A	10/1994	Wilson
5,480,382	A	1/1996	Hammerslag et al.

OTHER PUBLICATIONS

Beudet et al., A Suggested Nomenclature for Designating Mutations, *Human Mutation*, 245-248 (1993).

Borresen-Dale et al., Temporal Temperature Gradient Gel Electrophoresis on the DCode™ System—A Comparison to DGGE and CDGE in Mutation Screening, *Bio-Rad, Us/EG Bulletin* 2133.

Campbell et al., A new locus for non-syndromal, autosomal recessive, sensorineural hearing loss (DFNB16) maps to human chromosome 15q21-q22, *J. Med. Genet.*, 1015-1017 (1997).

Chevrier et al., PCR product quantification by non-radioactive hybridization procedures using an oligonucleotide covalently bound to microwells, *Molecular and Cellular Probes*, vol. 7, No. 3, 187-197 (1993).

Chevrier et al., Rapid detection of *Salmonella* subspecies I by PCR combined with non-radioactive hybridization using covalently immobilized oligonucleotide on a microplate, *FEMS—Immunology and Medical Microbiology*, vol. 10, 245-252 (1995).

Gasparini et al., Linkage of DFNB1 to Nonsyndromic Neurosensory Autosomal-Recessive Deafness in Mediterranean Families, *Eur. J. Hum. Genet.* 83-88 (1997).

Guilford et al., A non-syndromic form of neurosensory, recessive deafness maps to the pericentromeric region of chromosome 13q, *Nature Genetics*, vol. 6, 24-28 (1994).

Kelsell et al., Connexin 26 mutations in hereditary non-syndromic sensorineural deafness, *Nature*, vol. 387, 80-83 (1997).

Kiang et al., Upstream genomic sequence of the human connexin26 gene, *Gene*, 165-171, (1997).

Lee et al., Transcriptional Downregulation of Gap-Junction Proteins Blocks Junctional Communication in Human Mammary Tumor Cell Lines, *The Journal of cell Biology*, vol. 118, No. 5, 1213-1221 (1992).

(Continued)

Primary Examiner — Jehanne Sitton
(74) *Attorney, Agent, or Firm* — Foley & Lardner LLP

(57) **ABSTRACT**


A purified polynucleotide having a chain of nucleotides corresponding to a mutated sequence, which in a wild form encodes a polypeptide implicated in hereditary sensory defect, wherein said mutated purified polynucleotide presents a mutation responsible for prelingual non-syndromic deafness selected from the group consisting of a specific deletion of at least one nucleotide.

(56)

References Cited

OTHER PUBLICATIONS

- Lessa et al., Screening techniques for detecting allelic variation in DNA sequences, *Molecular Ecology*, vol. 2, 119-129 (1993).
- Liu et al., Mutations in the myosin VIIA gene cause non-syndromic recessive deafness, *Nature Genetics*, vol. 16, 188-190 (1997).
- Lynch, et al., Nonsyndromic Deafness DFNA 1 Associated with Mutation of α Human Homolog of the *Drosophilla* Gene *Diaphanous*, *Science*, vol. 278, pp. 1315-1318.
- Marazita et al., Genetic Epidemiological Studies of Early-Onset Deafness in the U.S. School-Age Population, *American Journal of Medical Genetics* 486-491 (1993).
- Maw et al., The Contribution of the DFNBI Locus to Neurosensory Deafness in a Caucasian Population, *Am. J. Hum. Genet.*, 629-635 (1995).
- Morton, N.E., Genetic Epidemiological of Hearing Impairment, *Annals of the New York Academy of Sciences*, vol. 630, 16-31 (1991).
- Pennisi, "The Architecture of Hearing," *Science*, vol. 278, pp. 1223-1224 (1997).
- Petit, C., Autosomal recessive non-syndromal hearing loss, *Genetics and Hearing Impairment*, Chap. 21, 197-211 (1996).
- Petit, C., Genes responsible for human hereditary deafness: symphony of a thousand, *Nature Genetics*, vol. 14, 385-391 (1996).
- Riesner et al., Temperature-gradient gel electrophoresis of nucleic acids: Analysis of conformational transitions, sequence variations, and protein-nucleic acid interactions, *Electrophoresis*, vol. 10, 377-389 (1989).
- Sambrook et al., Hybridization of Radiolabeled Oligonucleotides to Genomic DNA, *Molecular Cloning—A Laboratory Manual*, 9.56-9.57 (1989).
- Smith et al., Fluorescence detection in automated DNA sequence analysis, *Nature*, vol. 321, 674-679 (1986).
- Verhoeven et al., A Gene for Autosomal Dominant Nonsyndromic Hearing Loss (DFNA12) Maps to Chromosome 11q22-24, *Am. J. Hum. Genet.*, 1168-1173 (1997).
- Weil et al., The autosomal recessive isolated deafness, DFNB2, and the Usher 1 B syndrome are allelic defects of the myosin-VIIA gene, *Nature Genetics*, vol. 16, 191-193 (1997).
- Zelante et al., "Human Molecular Genetics," vol. 6, No. 9, Sep. 1, 1997, pp. 1605-1609.



**MUTATION WITHIN THE CONNEXIN 26
GENE RESPONSIBLE FOR PRELINGUAL
NON-SYNDROMIC DEAFNESS AND
METHOD OF DETECTION**

This is a continuation of application Ser. No. 13/430,062, filed Mar. 26, 2012, now U.S. Pat. No. 8,455,195, which is a continuation of Ser. No. 11/826,141, filed Jul. 12, 2007, now U.S. Pat. No. 8,143,000, which is a continuation of Ser. No. 10/278,089, filed Oct. 23, 2002, now U.S. Pat. No. 7,258,975, which is a continuation of Ser. No. 09/485,415, filed Feb. 10, 2000, now U.S. Pat. No. 6,485,908, which is a 371 of PCT/EP98/05175, filed Aug. 14, 1998, which claims the benefit of priority to U.S. Provisional Application No. 60/055,863, filed Aug. 15, 1997, all of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

The present invention concerns a mutation responsible for autosomal prelingual non-syndromic deafness and a method for the detection of this hereditary sensory defect for homozygous and heterozygous individuals. The invention concerns more particularly a specific deletion of at least one nucleotide in the connexin 26 (Cx26) gene and especially in a guanosine rich region, notably between the nucleotides 27 and 32. The invention is also directed to the use of polynucleotide, or fragments thereof, for example as tools useful for the in vitro detection of a mutation of a gene belonging to the Cx26 gene family.

Profound or severe prelingual deafness affects one child in a thousand in developed countries (Morton N E. Genetic epidemiology of hearing impairment. In Genetics of hearing impairment. (The New York Acad Sci, New York 1991; 630: 16-31). It is a major handicap as it impedes language acquisition.

According to studies performed in a U.S. population of children with non-syndromic (isolated) prelingual deafness and in whom an obvious environmental cause has been excluded, it is estimated that up to two-thirds of the cases have a genetic basis (Marazita M L, Ploughman L M, Rawlings B, Remington E, Amos K S, Nance W E. Genetic epidemiological studies of early-onset deafness in the U.S. school-age population. *Am J Med Genet* 1993; 46:486-91). These forms are mainly sensorineural and are almost exclusively monogenic. The major mode of inheritance is autosomal recessive (DFNB), involving 72% to 85% of cases, this fraction increasing to 90% when only profound deafness is taken into account.

Autosomal recessive prelingual deafness is known to be genetically highly heterogeneous. Estimates of the number of DFNB loci vary from thirty to one hundred (Petit C. Autosomal recessive non-syndromal hearing loss. In Genetics and Hearing Impairment. Martini A, Read A P, Stephens D, eds (Whurr, London) 1996; 197-212), for a review), of which fourteen have so far been mapped to the human chromosomes (Petit C. Genes responsible for human hereditary deafness: symphony of a thousand. *Nature Genet* 1996; 14:385-91) for review, (Verhoeven K, Van Camp G, Govaerts P J, et al. A gene for autosomal dominant non-syndromic hearing loss (DFNA12) maps to chromosome 11q22-24. *Am J Hum Genet* 1997; 60:1168-74 and Campbell D A, McHale D P, Brown K A, et al. A new locus for non-syndromal autosomal recessive sensorineural hearing loss (DFNB16) maps to human chromosome 15q21-q22. *J Med Genet* 1997; in press).

A majority of the families attending genetic counseling clinics consist of normal hearing parents with a single deaf

child who wish to know the risk of recurrence of the defect. In most cases, given the major role of environmental causes of prelingual deafness, it is not usually possible even to recognize whether the hearing loss is of genetic origin. Genetic counseling in such families would be greatly improved by an ability to detect DFNB mutations. In this respect, the high genetic heterogeneity of the condition represents a major obstacle.

After the initial identification of the DFNB1 locus on 13q11 in a large consanguineous Tunisian family (Guilford P, Ben Arab S, Blanchard S, et al. A non-syndromic form of neurosensory, recessive deafness maps to the pericentromeric region of chromosome 13q. *Nature Genet* 1994; 6:24-8), two studies performed on New Zealand/Australian families (Maw M A, Allen-Powell D R, Goodey R J, et al. The contribution of the DFNB1 locus to neurosensory deafness in a Caucasian population. *Am J Hum Genet* 1995; 57:629-35), and on Italian/Spanish families (Gasparini P, Estivill X, Volpini V, et al. Linkage of DFNB1 to non-syndromic neurosensory autosomal-recessive deafness in Mediterranean families. *Eur J Hum Genet* 1997; 5:83-8) suggested that this locus might be a major contributor to prelingual deafness in these populations, although individual lod scores obtained in these families were not significant owing to the small size of these families.

Recently, the Cx26 gene, which encodes a gap junction protein, connexin 26, has been shown to underlie DFNB1 deafness. Two different G→A substitutions resulting in premature stop codons in three DFNB1 linked consanguineous Pakistani families have been reported (Kelsell D P, Dunlop J, Stevens H P, et al. Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature* 1997; 387:80-3). These two substitutions were identified, respectively, at codon 77 and at codon 24. This result has offered the opportunity directly to assess this hypothesis.

The difficulties encountered in genetic counseling for prelingual non-syndromic deafness due to the inability to distinguish genetic and non-genetic deafness in the families presenting a single deaf child was one of the reasons that led the inventors to undertake a characterization of the spectrum and prevalence of mutations present in the Cx26 gene in 35 families from several parts of the world with autosomal recessive prelingual deafness.

SUMMARY OF THE INVENTION

The determination of a mutation in the Cx26 gene has notably rendered possible the use of a detection probe as a tool for the identification of a specific form of autosomal prelingual non-syndromic deafness, and more particularly the useful role of a newly identified 30delG (a G deletion at position 30; position 1 being the first base of the initiator codon) mutation in such families. This invention establishes that the contribution of the DFNB1 locus predominantly results essentially from the 30delG mutation. It is now believed that the 30delG accounts for about three-quarters of all recessive DFNB1 mutations.

The invention is thus intended to provide a purified polynucleotide having a chain of nucleotides corresponding to a mutated sequence, which in a wild form encodes a polypeptide implicated in hereditary sensory defect. The mutated purified polynucleotide presents a mutation responsible for prelingual non-syndromic deafness.

The invention also provides oligonucleotides comprising of 15 to 50 consecutive nucleotides of the mutated purified polynucleotide that are useful as primers or as probes.

In addition, the invention aims to supply a method and a kit for the detection of the hereditary sensory defect for homozygous as heterozygous individuals.

According to the invention, the purified polynucleotide having a chain of nucleotides corresponding to a mutated sequence, which encodes in a wild form a polypeptide implicated in hereditary sensory defect, presents a mutation responsible for prelingual non-syndromic deafness selected from the group consisting of a specific deletion of at least one nucleotide.

By mutation, according to the invention it means a specific deletion of at least one nucleotide. Thus, a mutated sequence means a polynucleotide sequence comprising at least a mutation.

A chain of nucleotides, according to the invention, means a polynucleotide, which encodes not necessarily a polypeptide, but which presents between 27 and 2311 nucleotides linked together.

The invention particularly concerns a purified polynucleotide wherein, the specific mutation is a deletion located in a region encoding connexin 26 of chromosome 13q11-12, preferably located in a guanosine rich region starting at nucleotide 27 preferably at nucleotide 30, and extending to nucleotide 32 or nucleotide 35, all the recited nucleotides being inclusive. More particularly according to the invention, the specific deleted purified polynucleotide encodes for a truncated polypeptide.

By truncated polypeptide, according to the invention it means a fragment of the polypeptide, which does not present the properties of the wild form of the polypeptide either in length, in amino acid composition, or in functional properties.

A preferred embodiment of a specific deletion is a guanosine deletion at position 30, also called "30delG mutation". Another preferred embodiment of the specific deletion is a 38 bp deletion beginning at position 30.

The invention also includes a purified polynucleotide, which hybridizes specifically with any one of the polynucleotides as defined above under the following stringent conditions: at low temperatures between 23° C. and 37° C., in the presence of 4×SSC buffer, 5×Denhardt's solution, 0.05% SDS, and 100 µg/ml of salmon sperm DNA. (1×SSC corresponds to 0.15 M NaCl and 0.05M sodium citrate; 1×Denhardt's solution corresponds to 0.02% Ficoll, 0.02% polyvinylpyrrolidone and 0.02% bovine serum albumin).

The invention also concerns an oligonucleotide useful as a primer or as a probe comprising 15 to 50 consecutive nucleotides of the polynucleotide according to any one of the polynucleotides as defined above. The oligonucleotide sequence is selected from the following group:

A first couple:

5' -TCTTTTCCAGAGCAAACCGCC-3' (SEQ ID NO. 1)

5' -TGAGCACGGGTGCCTCATC-3' (SEQ ID NO. 2)

The length of the PCR product has been obtained from 285 bp in length;

A second couple allowing to explore the other part of the reading frame:

5' -GACACGAAGATCAGCTGCAG-3' (SEQ ID NO. 3)

5' -CCAGGCTGCAAGAAGCTGTG-3' (SEQ ID NO. 4)

A third couple:

5' -CTAGTGATTCCTGTGTTGTGTGC-3'; (SEQ ID NO. 9)
and

5' -ATAATGCGAAAAATGAAGAGGA-3' (SEQ ID NO. 10)
and

A fourth couple:

5' -CGCCCGCCGCGCCCGCCCGCCCGCCCGCC (SEQ ID NO. 14)

GCCCCGCCCCCTAGTGATTCCTGTGTTGTGT

GC-3';
and

5' -ATAATGCGAAAAATGAAGAGGA-3'. (SEQ ID NO. 10)

Another oligonucleotide useful as a probe is selected from the following group:

5' -AGACGATCCTGGGGGTGTGAACAAA-3' (SEQ ID NO. 5)

5' -ATCCTGGGGGTGTGA-3' (SEQ ID NO. 6)

5' -AGACGATCCTGGGGCTCACCGTCTC-3'. (SEQ ID NO. 7)

In addition, the invention concerns a method for the detection of an hereditary sensory defect, namely autosomal prelingual non-syndromic deafness, for homozygous as heterozygous individuals in a biological sample containing DNA, comprising the steps of:

a) bringing the biological sample into contact with a oligonucleotide primers as defined above, the DNA contained in the sample having been optionally made available to hybridization and under conditions permitting a hybridization of the primers with the DNA contained in the biological sample;

b) amplifying the DNA;

c) revealing the amplification products;

d) detecting the mutation.

Step d) of the above-described method may comprise a Single-Strand Conformation Polymorphism (SSCP), a Denaturing Gradient Gel Electrophoresis (DGGE) sequencing (Smith, L. M., Sanders, J. Z., Kaiser, R. J., Fluorescence detection in automated DNA sequence analysis. *Nature* 1986; 321:674-9); a molecular hybridization capture probe or a temperature gradient gel electrophoresis (TGGE).

Step c) of the above-described method may comprise the detection of the amplified products with an oligonucleotide probe as defined above.

According to the invention, a biological sample can be a blood sample extracted from people suffering from any kind of deafness with any criteria as follows: neurosensorial or mixed isolated deafness, advanced or not, at any degree of severity, concerning familial or sporadic case, or individuals exposed to noise, or individuals suffering from a low acoustic, or individuals susceptible to carry an anomaly in the gene, or from an embryo for antenatal diagnostic.

Another aim of the invention comprises a method for the detection of an hereditary sensory defect, the autosomal prelingual non-syndromic deafness, for homozygous and heterozygous individuals in a biological sample containing DNA, comprising the steps of:

a) bringing the biological sample into contact with an oligonucleotide probe according to the invention, the DNA contained in the sample having been optionally made available to

hybridization and under conditions permitting a hybridization of the primers with the DNA contained in the biological sample; and

b) detecting the hybrid formed between the oligonucleotide probe and the DNA contained in the biological sample.

Step b) of the above-described method may consist in a single-strand conformation. Polymorphism (SSCP), a denaturing gradient gel electrophoresis (DGGE) or amplification and sequencing.

The invention also includes a kit for the detection of an hereditary sensory defect, the autosomal prelingual non-syndromic deafness, for homozygous as heterozygous individuals, said kit comprising:

- a) oligonucleotides according to the invention;
- b) the reagents necessary for carrying out DNA amplification; and
- c) a component that makes it possible to determine the length of the amplified fragments or to detect a mutation.

BRIEF DESCRIPTION OF THE DRAWINGS

This invention will be more described in greater detail by reference to the drawings in which:

FIG. 1 depicts the results of temperature gradient gel electrophoresis for detection of mutants in which:

Lanes 1 and 2: DNA from normal patients.

Lanes 3 and 4: DNA from homozygous patients with 30delG mutation.

Lanes 5 and 6: DNA from heterozygous patients.

Lane 7: PCR control without DNA.

Lane 8: PCR fragment amplified from a normal DNA and hybridized with a standard DNA fragment harboring the 30delG mutation.

Lane 9: PCR fragment amplified from a mutant homozygous DNA and hybridized with a normal standard DNA fragment harboring the guanine 30.

DETAILED DESCRIPTION OF THE INVENTION

Prelingual non-syndromic (isolated) deafness is the most frequent hereditary sensory defect in children. The inheritance in most is autosomal recessive. Several dozens of genes might be involved, only two of which, DFNB1 and DFNB2, have so far been identified (Kelsell, D. P., et al., Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature* 1997; 387:80-3; Liu, X-Z, et al., Mutations in the myosin VIIA gene cause non-syndromic recessive deafness, *Nature Genet* 1997; 16:188-90; and Weil, D., et al., The autosomal recessive isolated deafness, DFNB2, and the Usher 1B syndrome are allelic defects of the myosin-VIIA. *Nature Genet* 1997; 16:191-3). A search was made searched for mutations in the gene encoding connexin 26, Cx26, which has recently been shown to be responsible for DFNB1. Mutation analysis of Cx26 was performed by PCR amplification on genomic DNA and sequencing of the single coding exon.

Example 1

Patients

Thirty-five affected families from various geographical regions, mainly France, New Zealand and Australia, Tunisia and Lebanon, were studied. They could be classified into three categories: (1) consanguineous families each having a significant linkage to the DFNB1 locus; (2) small non-consanguineous families in which linkage analysis was compat-

ible with the involvement of DFNB1; and (3) small families in which no linkage analysis had been undertaken.

The first category consists of six large families living in geographically isolated regions. Five were from Tunisia, two from the north and three from the south. Linkage to the DFNB1 locus in the two families from northern Tunisia (families 20 and 60) had previously been reported (Guilford P, Ben Arab S, Blanchard S, et al., A non-syndromic form of neurosensory, recessive deafness maps to the pericentromeric region of chromosome 13q. *Nature Genet* 1994; 6:24-8); the three families from southern Tunisia (S15, S19 and ST) and the family from Lebanon (LH) comprise total of three, five, two, and five deaf children, respectively, the deafness being of severe or profound degree. The marriages were between first cousins (S15, ST and LH) and between first and second cousins (S19). Linkage analysis of these six families resulted in individual lod scores ranging from 2.5 to 10 with polymorphic markers from the DFNB1 region (D13S175, D13S141, D13S143 and D13S115).

The second category of patients comprises seven New Zealand families with at least two deaf siblings (families 51, 1160, 1548, 1608, 1773, 1873, 1877) and one Australian (9670) family. Family 1608 was atypical in that four siblings sharing the same DFNB1 marker haplotypes had a mild to moderate deafness (severe at high frequency), with the child of one of them being profoundly deaf. In family 1873, the unrelated parents (individuals II.2 and II.3) were deaf as well as their two children, and we have therefore considered this as two families, bringing to nine the total of independent families. Apart from families 1608 and 1873, no parent acknowledged any hearing impairment. These nine families showed cosegregation between deafness and polymorphic markers of the DFNB1 region with maximum individual lod scores ranging from 0.6 to 1.2. Ten other families in the original study of Maw et al. (Maw M A, Allen-Powell D R, Goodey R J, et al. The contribution of the DFNB1 locus to neurosensory deafness in a Caucasian population. *Am J Hum Genet* 1995; 57:629-35) had shown no cosegregation and one other cosegregating family was not tested for Cx26 mutations. The New Zealand families were all of Caucasian origin with no known Polynesian admixture. According to the antecedent family names, the ancestral proportion among the families reflected that of the general Caucasian New Zealand population with the great predominance being of Anglo-Celtic patrimony and a small fraction due to migration from continental Europe. Neither parental consanguinity, nor links between any of the families were recognized. In the Australian case, the father was from Northern Ireland and the mother from Yorkshire, England.

The third category is composed of nineteen families living in France and two in New Zealand, each with at least two children having a severe to profound deafness. No parent acknowledged any hearing impairment, except for the mother in family P16 and the father in family P17 who had moderate and progressive high-frequency hearing loss. Five of these families had foreign ancestors from Lebanon (family P3), Turkey (family P4), Portugal (family P9), Algeria (family P14) and Poland (father in family P16). In two of the families (P7 and P14), the parents were distantly related.

Example 2

Amplification of the coding exon of Cx26 PCRs were carried out on genomic DNA using a set of primers that allowed the amplification of the entire coding sequence of the Cx26 gene, which consists of a single coding exon (Kelsell D P, Dunlop J, Stevens H P, et al. Connexin 26 mutations in

7

hereditary non-syndromic sensorineural deafness. *Nature* 1997; 387: 80-4 Primer sequences were as follows:

5' -TCTTTTCCAGAGCAAACCGCC-3' (SEQ ID NO. 1)
and

5' -TGAGCACGGGTGCCTCATC-3'. (SEQ ID NO. 2)

PCR conditions were: 35 cycles of 95° C., 1 min; 58° C., 1 min; 72° C., 2 min. The PCR product obtained was 777 bp in length.

Example 3

DNA Sequencing

Sequencing of the PCR products was performed as previously described (Smith L. M. Sanders J Z, Kaiser R J, et al., Fluorescence detection in automated DNA sequence analysis, *Nature* 1986; 321:674-9) using the dideoxy chain terminator method on an Applied Biosystems DNA sequencer ABI373 with fluorescent dideoxynucleotides. The primers used were the same as those for the PCR amplification plus two internal primers:

5' -GACACGAAGATCAGCTGCAG-3' (SEQ ID NO. 3)
and

5' -CCAGGCTGCAAGAAGCTGTG-3'. (SEQ ID NO. 4)

Example 4

Mutations in Consanguineous Tunisian and Lebanese DFNB1 Families

In these families the involvement of the DFNB1 locus could be demonstrated by linkage analysis. In four of the five families from Tunisia (S15, S19, 20, and 60) and in the Lebanese family (LH), the same mutation was detected in all affected children on both Cx26 alleles, namely, a deletion of a guanosine (G) in a sequence of six G extending from position 30 to 35 (position 1 being the first base of the initiator codon) (Table 1). This mutation is hereafter referred to as 30delG mutation according to the nomenclature proposed by Beaudet and Tsui ((Beaudet A L, Tsui L-C. A suggested nomenclature for designating mutations, *Hum Mutation* 1993; 2: 245-8)). It creates a frameshift, which results in a premature stop codon at nucleotide position 38. The mutation segregating in the fifth family from Tunisia (ST) was identified as a G to T transversion at nucleotide position G39 creating a premature stop codon (GAG→TAG) at codon 47, and was designated E47X. In each family, normal hearing parents were found to be heterozygous for the corresponding mutation.

Example 5

Mutations in Small Nonconsanguineous New Zealand and Australian Families Consistent with DFNB1 Linkage

In these families, segregation analysis has previously been reported as compatible with the involvement of the DFNB1 locus (Maw M A, Allen-Powell D R, Goodey R J, et al. The contribution of the DFNB1 locus to neurosensory deafness in a Caucasian population. *Am J Hum Genet* 1995; 57: 629-35). The deaf individuals from five of the nine families (51, 1160, 1608 (III.20), 1873 (II.3) and 1877) were homozygous for the

8

30delG mutation. The deaf children from family 1773 were heterozygous for 30delG. Deaf individual II.2 from family 1873 (see "subjects" and Table 1) was heterozygous for a deletion of 38 bp beginning at nucleotide position G30, designated 30del38. No other mutation was detected in the deaf children of family 1773 and the deaf individual (II.2) in family 1873. Nevertheless, in this last individual, a deletion of the polymorphic marker immediately proximal to the Cx26 gene (locus D13S175) had previously been observed (Maw M A, Allen-Powell D R, Goodey R J, et al. The contribution of the DFNB1 locus to neurosensory deafness in a Caucasian population. *Am J Hum Genet* 1995; 57: 629-35), which may indicate that a DNA rearrangement has impaired the functioning of the other Cx26 allele of the gene in cis. In family 9670, compound heterozygosity for a missense mutation (R184P) and an in frame single-codon deletion (delE138) was observed in affected siblings. In only one family (1548) was no Cx26 mutation detected. Results are summarized in Table 1.

Example 6

Mutations in Small Families Uncharacterized for DFNB1 Linkage Living in France and New Zealand

Nineteen families (P1 to 17, L14190 and L13131) living in France and two in New Zealand (families 1885 and 2254) were studied. In these families, cosegregation of the deafness with polymorphic markers had not been analysed. Deaf children from six of the twenty-one families (P1, P3, P5, P9, P10, and P16) were found to be homozygous for the mutation 30delG. In five additional families (P6, P11, P14, P17, and 1885), deaf children were heterozygous for this mutation; no other mutation was detected in these families. In the ten remaining families, no mutation in the Cx26 gene was found.

Example 7

Molecular Hybridization Using Allele-Specific Capture Probes

Molecular hybridization capture probe (see, e.g., D. Chevrier et al. PCR product quantification by non-radioactive hybridization procedures using an oligonucleotide covalently bound to microwells. *Molecular and Cellular Probes* 1993; 7: 187-197 and D Chevrier et al. Rapid detection of *Salmonella* subspecies I by PCR combined with non-radioactive hybridization using covalently immobilized oligonucleotide on a microplate. *FEMS Immunology and Medical Microbiology* 1995; 10: 245-252 each of which is incorporated by reference herein) permit specific detection of the 30delG mutation. The technique has been adapted to permit rapid diagnosis of prelingual non-syndromic deafness caused by the 30delG mutation. The technique provides certain advantages in a clinical setting because it uses stable, nonradioactive molecules, it can be easily automated, and it is well adapted to large scale analysis.

Using primers designed for PCR amplification, the region of interest in the Cx26 gene is amplified from genomic DNA samples. The primer sequences are as follows:

CONN3:
5' -CTAGTGATTCTGTGTTGTGTGC-3' (SEQ ID NO. 9)

CONN4:
5' -ATAATGCCGAAAAATGAAGAGGA-3' (SEQ ID NO. 10)

PCR is performed with the CONN3 (SEQ ID NO:9) and CONN4 (SEQ ID NO:10) primers (1 μM each), an aliquot of the DNA to be analyzed (2 μl, 100-300 ng), 1.5 mM MgCl₂,

200 μ M dNTP, and Taq polymerase. The amplification program consists of the following steps: 1) 95° C., 5 min; 2) addition of enzyme, 95° C., 1 min; 3) 60° C., 1 min (ramp rate=0.25° C./s); 4) 72° C., 1 min; 5) repeat steps 2 to 4 for 40 cycles; and 6) 72° C., 10 min. PCR products are verified by a rapid gel electrophoresis.

The amplified PCR product contains either the normal or the mutant Cx26 sequence. To distinguish between the normal and mutant sequence, two capture probes are designed. The sequences of these two capture probes are as follows:

For detection of normal sequence:

CONN6:
5'-AAAAAAATCCTGGGGGTGTG-3' (SEQ ID NO. 11)

For detection of mutant sequence:

CONN7:
5'-AAAAAAATCCTGGGGGTGTGA-3' (SEQ ID NO. 12)

Each capture probe must be 22 nucleotides long. Furthermore, to be efficient, the capture probe must include an A₇ spacer at its 5' end and a hybridization region of 15 bases. Such a capture probe is able to specifically differentiate the mutant sequence from the normal sequence. Thus, CONN6 (SEQ ID NO:11) is designed to specifically hybridize with the normal sequence, whereas CONN7 (SEQ ID NO:12) is designed to specifically hybridize with the mutant sequence.

Before attaching the capture probes to a microtiter plate, they are phosphorylated at their 5' ends. The phosphorylation is carried out for 1 hour at 37° C. in presence of 20 nmoles of CONN6 (SEQ ID NO:11) or CONN7 (SEQ ID NO:12) oligonucleotides, 100 μ M ATP, 10 units T4 polynucleotide kinase in 200 μ l of buffer (50 mM Tris-HCl pH 7.4; 10 mM MgCl₂, 5 mM dithiothreitol; and 1 mM spermidine). The mixture is heated for 10 min. at 68° C. to inactivate the T4 polynucleotide kinase, then the oligonucleotide is precipitated by adding 145 μ l of 10 M CH₃COONH₄, 15 μ l H₂O, and 800 μ l iced ethanol. After a 30 min. incubation in ice, the mixture is centrifuged for 20 min. at 12,000 \times g at 4° C. The resulting pellet is washed with 500 μ l iced ethanol (70%) and dissolved in 800 μ l of TE buffer. The phosphorylated oligonucleotide concentration is determined by optical density at 260 nm.

Before attaching the phosphorylated oligonucleotides to microplates, they are denatured by heating at 95° C. for 10 min. and rapidly cooled in ice to avoid the formation of secondary structure. 500 ng of phosphorylated CONN6 (SEQ ID NO:11) or CONN7 (SEQ ID NO:12) and 1 μ l of 1 M 1-methylimidazole, pH 7, is added to each well of a microplate, which is kept on ice. The total volume of each well is adjusted to 70 μ l with distilled water, before adding 30 μ l of a cold, 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide solution (167 mM). The microplate is covered and incubated for 5 hours at 50° C. in an incubator (Thermomix® from Lab-systems). After the 5-hour incubation, the microplate is washed three times with a warm solution (50° C.) of 0.4 N NaOH containing 0.25% SDS. The microplate is incubated for 5 min. with the same warm solution and washed again with warm NaOH/SDS (50° C.). Finally, the microplate is washed five times with TE buffer. The coated microplate can be kept several months at 4° C., if the wells are filled with TE buffer.

The amplified sequences from the genomic DNA samples are incubated with a biotinylated detection probe in the coated microplates. Unlike the capture probes; which are allele specific, the detection probe can hybridize with both the normal and mutant sequences. The sequence of the detection probe is:

CONN12:
5'-CAGCATTGGAAAGATCTGGCTCA-3'. (SEQ ID NO. 13)

The amplified sequences and the detection probe, which is biotinylated at its 5' end, are denatured directly in the microplates by successively adding to each well: 95 μ l of water, 5 μ l of PCR reaction, 40 μ l of biotinylated probe (SEQ ID NO:13) at 22 nM diluted in water, and 14 μ l 1 N NaOH. After 10 min., 21 μ l of 1 M NaH₂PO₄ and 1% Sarkosyl is added to each well to bring the total volume to 175 μ l per well. The final concentration of the detection probe is 5 nM. The microplate is covered and incubated overnight at 40° C. in an incubator (Thermomix® from Lab-systems) and then extensively washed (5 times) with TBS-Tween to remove the excess biotinylated probe (SEQ ID NO:13).

An immunoenzymatic method is used to detect the hybridized probe. Each well receives 100 μ l of the conjugate (Ex-travidine—alkaline phosphatase, Sigma E-2636) diluted 1/4000 in TBS-BSA-Tween. The microplate is covered and incubated for 1 hour at 25° C. Following the incubation, the microplate is washed 5 times with TBS-Tween. Then 200 μ l of preheated (37° C.) substrate (7.5 mg para-nitro-phenyl-phosphate in 20 ml of the following buffer: 1 M diethanolamine pH 9.8 containing 1 mM MgCl₂) are added to each well. The microplate is covered and incubated for 3 hours at 37° C. The absorbance is measured at 405 nm to determine the specific signal and at 630 nm to determine the background noise.

The hybridization ratio (R) between the signal obtained with CONN6 (SEQ ID NO:11) probe (normal sequence) and that obtained with CONN7 (SEQ ID NO:12) probe mutant sequence) is calculated. The calculated R values are used to determine the genotypes of the sample DNA as follows: homozygous for the normal Cx26 sequence (R \geq 2), heterozygous for the 30delG mutation (0.5<R<2), and homozygous for the 30delG mutation (R \leq 0.5). The range of the hybridization ratio (R) can be slightly modified when the number of samples increases. The following table represents an example of results obtained with 39 samples.

Genotype:	Hybridization ratio (R)		
	Normal	Homozygous 30delG	Heterozygous
	5.96	0.48	1.33
	5.43	0.17	1.13
	3.39	0.21	0.73
	4.14	0.16	0.63
	4.09	0.28	1.4
	2.76	0.13	0.73
	2.2	0.21	0.76
	3.97	0.4	0.73
	4.07		1.06
	3		
	2.76		
	3.66		
	3.87		
	3.92		
	3.26		
	5.17		
	2.74		
	4.51		
	6.3		
	3.49		
	4.05		
	3.17		
Number	22	8	9
Mean value	3.91	0.26	0.94
Standard deviation	1.06	0.12	0.29
Range	(6.3-2.2)	(0.48-0.13)	(1.4-0.63)

Temperature Gradient Gel Electrophoresis

Temperature gradient gel electrophoresis (TGGE) permits the detection of any type of mutation, including deletions, insertions, and substitutions, which is within a desired region of a gene. (See, e.g. D. Reiner et al. Temperature-gradient gel electrophoresis of nucleic acids: Analysis of conformational transitions, sequence variations and protein-nucleic acid interactions. *Electrophoresis* 1989; 10: 377-389; E. P. Lessa and G. Applebaum Screening techniques for detecting allelic variation in DNA sequences. *Molecular Ecology* 1993; 2: 119-129 and A. L. Börresen-Dale et al. Temporal Temperature Gradient Gel Electrophoresis on the D Code™ System. *Bio-Rad US/EG Bulletin* 2133; the entire disclosure of each publication is incorporated by reference herein.) However, TGGE does not permit one to determine precisely the type of mutation and its location.

As in the previously described molecular hybridization technique, the region of interest in the Cx26 gene is first amplified from genomic DNA samples by PCR. The primer sequences are as follows:

CONN2:
5'-CGCCCGCGCGCCCGCGCCCGCCCGCC (SEQ ID NO. 14)

GCCCCGCCCCCTAGTGATTCTCTGTGTTGTGT

GC-3'

CONN4:
5' ATAATGCGAAAAATGAAGAGGA-3' (SEQ ID NO. 10)

PCR is performed with 1 μM of the CONN2 (SEQ ID NO:14) primer, which has a GC clamp at its 5' end, and 1 μM of the CONN4 (SEQ ID NO:10) primer, an aliquot of the DNA to be analyzed (2 μl, 100-300 ng), 1.5 mM MgCl₂, 200 μM dNTP, and Taq polymerase. The amplification program consists of the following steps: 1) 95° C., 5 min; 2) addition of enzyme, 95° C., 1 min; 3) 60° C., 1 min (ramp rate=0.25° C./s); 4) 72° C., 1 min; 5) repeat steps 2 to 4 for 40 cycles; and 6) 72° C., 10 min.

Analyzing these PCR amplification fragments by TGGE can differentiate between homozygous (normal or mutant) samples, which produce a single band on a gel, and heterozygous samples, which produce three bands. However, differentiating between genomic samples that are homozygous for the normal sequence and genomic samples that are homozygous for the 30delG mutants requires an additional step.

To differentiate normal homozygous versus mutant homozygous samples, an aliquot of the amplified PCR product is mixed with either a known, normal homozygous sample or a known, 30delG mutant homozygous sample and analyzed for heteroduplex formation. If the amplified PCR product derives from a normal, homozygous sample, it will form a heteroduplex with the known, 30delG mutant homozygous sample. On the other hand, if the amplified PCR product derives from a mutant, homozygous sample, it will form a heteroduplex with the known, normal homozygous sample. To promote heteroduplex formation in these mixtures, they are denatured at 95° C. for 5 min, followed by a renaturation step at 60° C. for 45 min.

The PCR fragments from the initial amplification and those that are subjected to the additional heating steps to permit heteroduplex formation are analyzed on a 10% polyacrylamide gel containing 7 M urea. By way of example, a 30 ml gel is prepared by combining the following ingredients:

12.6 g urea
0.75 ml 50×TAE
7.5 ml acrylamide:bisacrylamide (37.5:1) at 40%
water to bring volume to 30 ml
30 μl Temed (added extemporaneously)
300 μl 10% ammonium persulfate (added extemporaneously).

After adding the Temed and ammonium persulfate, the gel is poured between two glass plates (Dcode Universal Mutation Detection System® from BIORAD) and allowed to polymerize for 1 hour.

An aliquot (7.5 μl) of the PCR mixture is mixed with 7.5 μl of 2× sample solution (2 mM EDTA pH 8; 70% glycerol; 0.05% xylene cyanol; 0.05% bromophenol blue), and introduced into a gel well. Electrophoresis is performed for 4-5 hours at 150V in 1.25×TAE buffer with a temperature gradient ranging from 61° C. to 62° C. at a rate of 0.2° C. per hour. Following electrophoresis, the gel is incubated for 6 min. in 1.25×TAE containing 25 μg/ml ethidium bromide. Excess ethidium bromide is removed by a 20 min. wash in 1.25×TAE, and the DNA fragments are visualized with a UV transilluminator.

A typical TGGE result is represented in FIG. 1. The amplified DNA from homozygous patients (normal or mutant) produces only one band. The amplified DNA from heterozygous patients results in three different fragments in the polyacrylamide gel. The more intense band, which migrates more rapidly, corresponds to both homoduplexes, which cannot be separated in this gel. The other two bands, which migrate more slowly, correspond to both kinds of heteroduplexes.

The DNA of normal homozygous patients can be differentiated from the DNA of mutant homozygous patients by analyzing the PCR fragments that were subjected to the conditions that permitted heteroduplex formation. Heteroduplexes form when the PCR amplified fragment from a normal homozygous genome is mixed with sequences from a known, mutant homozygous genome, or when the PCR amplified fragment from a mutant homozygous genome is mixed with sequences from a known, normal homozygous genome. These heteroduplexes are visible by TGGE analysis. Consequently, the DNA of normal and mutant homozygous patients can be easily differentiated by this technique using the primers described in the present study.

In all the known DFNB1 families (6/6), in all but one (8/9) of the putatively DFNB1-linked families, and in about half (11/21) of the families not tested for DFNB1 linkage, a mutation in Cx26 was detected. Furthermore, of the 44 chromosomes reckoned to be independent upon which a Cx26 mutant allele was identified or inferred, 33 (75%) were found to carry the same deletion of a guanosine, G, at position 30 (30delG).

Cx26 mutations represent a major cause of recessively inherited prelingual deafness and would be implicated in about half of cases in the examined populations. In addition, one specific mutation, 30delG, accounts for the majority (about three-quarters in our series) of the Cx26 mutant alleles.

The wild type connexin 26 gene published in LEE S. W. et al. (1992) *J. Cell Biol.* 118: 1213-1221 has the following sequence:

(SEQ ID NO: 15)

1 GATTTAATCC TATGACAAAC TAAGTTGGTT CTGTCTTACC CTGTTTTGGT

51 GAGGTTGTGT AAGAGTTGGT GTTTGCTCAG GAAGAGATTT AAGCATGCTT

-continued

101 GCTTACCCAG ACTCAGAGAA GTCTCCCTGT TCTGTCCCTAG CTATGTTCTGT
 151 GTGTTGTGTG CATTTCGTCTT TTCCAGAGCA AACCGCCAG AGTAGAAGAT
 201 GGATTGGGGC ACGCTGCAGA CGATCCTGGG GGGTGTGAAC AAACACTCCA
 251 CCAGCATTGG AAAGATCTGG CTCACCGTCC TCTTCATTTT TCGCATTATG
 301 ATCCTCGTTG TGGCTGCAAA GGAGGTGTGG GGAGATGAGC AGGCCGACTT
 351 TGTCTGCAAC ACGCTGCAGC CAGGCTGCAA GAACGTGTGC TACGATCACT
 401 ACTTCCCCAT CTCCCACATC CGGCTATGGG CCCTGCAGCT GATCTTCTGT
 451 TCCAGCCAG CGTCCTAGT GGCCATGCAC GTGGCCTACC GGAGACATGA
 501 GAAGAAGAGG AAGTTCATCA AGGGGAGAT AAAGAGTGAA TTTAAGGACA
 551 TCGAGGAGAT CAAAACCCAC AAGGTCCGCA TCGAAGGCTC CCTGTGGTGG
 601 ACGTACACAA GCAGaATCTT CTCCCGGTC ATCTTCGAAG CCGCCTTCAT
 651 GTACGCTTC TATGTCATGT ACGACGGCTT CTCCATGCAG CGGCTGGTGA
 701 AGTGCAACGC CTGGCCTTGT CCCAACACTG TGGACTGCTT TGTGTCCOAG
 751 CCCACGGAGA AGACTGTCTT TCACAGTGTT CATGATTGCA GTGTCTGGAA
 801 TTTGCATCCT GCTGAATGTC ACTGAATTGT GTTATTTGCT AATTAGATAT
 851 TGTTCCTGGG AGTCAAAAAA GCCAGTTTAA CGCATTGCC AGTTGTTAGA
 901 TTAAGAAATA GACAGCATGA GAGGGATGAG GCAACCCGTG CTCAGCTGTC
 951 AAGGCTCAGT CGCCAGCATT TCCCAACACA AAGATTCTGA CCTTAAATGC
 1001 AACCATTGTA AACCCCTGTA GGCCTCAGGT GAAACTCCAG ATGCCACAAT
 1051 GAGCTCTGCT CCOCTAAACC CTCAAAACAA AGGCCTAATT CTATGCCTGT
 1101 CTTAATTTTC TTTCACTTAA GTTAGTTCCA CTGAGACCCC AGGCTGTTAG
 1151 GGGTTATTGG TGTAAGGTAC TTTTCATATTT TAAACAGAGG ATATCGGCAT
 1201 TTGTTCTTTT CTCTGAGGAC AAGAGAAAAA AGCCAGGTTC CACAGAGGAC
 1251 ACAGAGAAGG TTTGGGTGTC CTCCTGGGGT TCTTTTTGCC AACTTTCCCC
 1301 ACGTTAAAGG TGAACATTGG TTCTTTCATT TGCTTTGGAA GTTTTAATCT
 1351 CTAACAGTGG ACAAAGTTAC CAGTGCCTTA AACTCTGTTA CACTTTTGG
 1401 AAGTGAAAAC TTTGTAGTAT GATAGGTTAT TTTGATGTAA AGATGTTCTG
 1451 GATACCATTA TATGTTCCCC CTGTTTCAGA GGCTCAGATT GTAATATGTA
 1501 AATGGTATGT CATTTCGTAC TATGATTTAA TTTGAAATAT GGTCTTTTGG
 1551 TTATGAATAC TTTGCAGCAC AGCTGAGAGA GGCTGTCTGT TGTATTCAAT
 1601 GTGGTCATAG CACCTAACAA CATTGTAGCC TCAATCGAGT GAGACAGACT
 1651 AGAAGTTCCT AGTTGCCTTA TGATAGCAAA TGGCCTCATG TCAAATATTA
 1701 GATGTAATTT TGTGTAAGAA ATACAGACTG GATGTACCAC CAACTACTAC
 1751 CTGTAATGAC AGGCCTGTCC AACACATCTC CTTTTCCAT GCTGTGGTAG
 1801 CCAGCATCGG AAAGAACGCT GATTTAAAGA GGTGAGCTTG GGAATTTTAT
 1851 TGACACAGTA CCATTTAATG GGGAGACAAA AATGGGGGCC AGGGGAGGGA
 1901 GAAGTTTCTG TCGTTAAAAA CGAGTTTGGG AAGACTGGAC TCTAAATCT
 1951 GTTGATTAAG GATGAGCTTT GTOTACCTTC AAAAGTTTGT TTGGCTTACC
 2001 CCOTTACAGC TCCAATTTT TAAGTAAAAA TATAACTAAT AACATGTGAA
 2051 AAGAATAGTA GCTAAGGTTT AGATAAATAT TGAGCAGATC TATAGGAAGA
 2101 TTGAACCTGA ATATTGCCAT TATGCTTGAC ATGGTTTCCA AAAAAATGTA

-continued

2151 CTCACATAG TTCAGTGAGG GTAAGTATTT TCCTGTTGTC AAGAATAGCA
 2201 TTGTAAAAGC ATTTTGTAAAT AATAAAGAAT AGCTTTAATG ATATGCTTGT
 2251 AACTAAAATA ATTTTGTAAAT GTATCAAATA CATTTAAAC ATTAAAATAT
 2301 AATCTCTATA AT

The wild type connexin 26 gene is published in D. T. Kiang ¹⁰
 et al. (1997) Gene 199 (1-2): 165-171; has the following
 sequence:

1 GATTTAATCC TATGACAAAC TAAGTTGGTT CTGTCTTAC CTGTTTTGGT
 51 GAGGTTGTGT AAGAGTTGGT GTTTGCTCAG GAAGAGATTT AAGCATGCTT
 101 GCTTACCAG ACTCAGAGAA GTCTCCCTGT TCTGTCTTAG CTAGTGATTC
 151 CTGTGTTGTG TGCATTGTC TTTCCAGAG CAAACCGCCC AGAGTAGAAG
 201 ATGGATTGGG GCACGCTGCA GACGATCCTG GGGGGTGTGA ACAAACTC
 251 CACCAGCATT GGAAAGATCT GGCTCACCGT CCTCTTCATT TTTGCATTA
 301 TGATCCTCGT TGTGGCTGCA AAGGAGGTGT GGGGAGATGA GCAGGCCGAC
 351 TTTGTCTGCA ACACCCTGCA GCCAGGCTGC AAGAACGTGT GCTACGATCA
 401 CTACTTCCCC ATCTCCACA TCCGGCTATG GGCCCTGCAG CTGATCTTCG
 451 TGTCACGCC AGCGCTCCTA GTGGCCATGC ACGTGGCCTA CCGGAGACAT
 501 GAGAAGAAGA GGAAGTTCAT CAAGGGGAG ATAAGAGTG AATTTAAGGA
 551 CATCGAGGAG ATCAAAACCC AGAAGGTCCG CATCGAAGGC TCCCTGTGGT
 601 GGACCTACAC AAGCAGCATC TTCTCCGGG TCATCTTCGA AGCCGCCTTC
 651 ATGTACGTCT TCTATGTCAT GTACGACGGC TTCTCCATGC AGCGGCTGGT
 701 GAAGTGCAAC GCCTGGCCTT GTCCCAACAC TGTGGACTGC TTTGTGTCCC
 751 GGCCACGGA GAAGACTGTC TTTCACAGTG TTCATGATTC CAGTGTCTGG
 801 AATTTGCATC CTGCTGAATG TCACTGAATT GTGTTATTTG CTAATTAGAT
 851 ATTGTCTGG GAAGTCAAAA AAGCCAGTTT AACGCATTGC CCAGTTGTTA
 901 GATTAAGAAA TAGACAGCAT GAGAGGGATG AGGCAACCCG TGCTCAGCTG
 951 TCAAGGCTCA GTCGCCAGCA TTTCCCAACA CAAAGATTCT GACCTTAAAT
 1001 GCAACCATTT GAAACCCCTG TAGGCCTCAG GTGAACTCC AGATGCCACA
 1051 ATGAGCTCTG CTCCCCTAAA GCCTCAAAAC AAAGCCCTAA TTCTATGCCT
 1101 GTCTTAATTT TCTTCACTT AAGTTAGTTC CACTGAGACC CCAGGCTGTT
 1151 AGGGTTATT GGTGTAAGGT ACTTTCATAT TTTAAACAGA GGATATCGGC
 1201 ATTTGTTTCT TTCTCTGAGG ACAAGAGAAA AAAGCCAGGT TCCACAGAGG
 1251 ACACAGAGAA GGTGTTGGTG TCCTCCTGGG GTTCTTTTTG CCAACTTCC
 1301 CCACGTTAAA GGTGAACATT GGTCTTTCA TTTGCTTGG AAGTTTAAAT
 1351 CTCTAACAGT GGACAAAGTT ACCAGTGCCT TAAACTCTGT TACACTTTTT
 1401 GGAAGTGAAA ACTTTGTAGT ATGATAGGTT ATTTTGATGT AAAGATGTTT
 1451 TGATAACCAT TATATGTTCC CCCTGTTTCA GAGGCTCAGA TTGTAATATG
 1501 TAAATGGTAT GTCATTGCT ACTATGATTT AATTTGAAAT ATGGTCTTTT
 1551 GGTATGAAT ACTTTGCAGC ACAGCTGAGA GAGGCTGTCT GTTGATTTCA
 1601 TTGTGTCAT AGCACCTAAC AACATTGTAG CCTCAATCGA GTGAGACAGA

-continued

1651 CTAGAAGTTC CTAGTTGGCT TATGATAGCA AATGGCCTCA TGTCAAATAT
 1701 TAGATGTAAT TTTGTGTAAG AAATACAGAC TGGATGTACC ACCAACTACT
 1751 ACCTGTAATG ACAGGCCTGT CCAACACATC TCCCTTTTCC ATGCTGTGGT
 1801 AGCCAGCATC GGAAAGAACG CTGATTTAAA GAGGTGAGCT TGGGAATTTT
 1851 ATTGACACAG TACCATTTAA TGGGGAGACA AAAATGGGGG CCAGGGGAGG
 1901 GAGAAGTTTC TGTCGTTAAA AACGAGTTTG GAAAGACTGG ACTCTAAATT
 1951 CTGTTGATTA AAGATGAGCT TTGTCTACCT TCAAAAGTTT GTTTGGCTTA
 2001 CCCCCCTCAG CCTCCAATT TTTAAGTGAA AATATAACTA ATAACATGTG
 2051 AAAAGAATAG AAGCTAAGGT TTAGATAAAT ATTGAGCAGA TCTATAGGAA
 2101 GATTGAACCT GAATATTGCC ATTATGCTTG ACATGGTTTC CAAAAAATGG
 2151 TACTCCACAT ACTTCAGTGA GGGTAAGTAT TTTCTGTTG TCAAGAATAG
 2201 CATTGTAATA GCATTTTGTA ATAATAAAGA ATAGCTTTAA TGATATGCTT
 2251 GTAACATAAA TAATTTTGTA ATGTATCAAA TACATTTAAA ACATTAATAA
 2301 ATAATCTCTA TAAT

(SEQ ID No. 8). The ATG underlined in the sequences corresponds to the start codon. The guanine residue "G", which is in bold print, marks the end of the guanosine rich region between nucleotides 27 and 32, inclusive.

TABLE I

Mutations in the Cx26 coding exon in individuals affected with familial forms of prelingual deafness			
Family (geographical origin)	30delG mutation	Other mutation	Deafness
DFNB1-linked families			
S15 (sTu)	homozygous	—	profound
S19 (sTu)	homozygous	—	profound
ST (sTu)	—	homozygous E47X	profound
20 (nTu)	homozygous	—	profound
60 (nTu)	homozygous	—	profound
LH (Leb)	homozygous	—	severe-profound
Families consistent with DFNB1 linkage			
51 (NZ)	homozygous	—	severe-profound
1160 (NZ)	homozygous	—	moderate-severe*
1548 (NZ)	—	—	profound
1608 (NZ)	homozygous	—	profound**
1773 (NZ)	heterozygous	—	profound
1873 individual II.3 (NZ)	homozygous	—	moderate
1873 individual II.2 (NZ)	—	heterozygous 30del38	profound
1877 (NZ)	homozygous	—	profound
9670 (Aust)	—	delE118/R14 8P	moderate-severe

25

TABLE I-continued

Mutations in the Cx26 coding exon in individuals affected with familial forms of prelingual deafness			
Family (geographical origin)	30delG mutation	Other mutation	Deafness
Families uncharacterized for DFNB1 linkage			
35 P1 (Fr)	homozygous	—	severe-profound
P2 (Fr)	—	—	profound
P3 (Leb)	homozygous	—	severe-profound
P4 (Tur)	—	—	severe-profound
P5 (Fr)	homozygous	—	profound
P6 (Fr)	heterozygous	—	severe-profound
40 P7 (Fr)	—	—	moderate
P8 (Fr)	—	—	moderate
L13131 (Fr)	—	—	profound
L14190 (Fr)	—	—	mild-moderate
P9 (Por)	homozygous	—	severe-profound
P10 (Fr)	homozygous	—	severe-profound
P11 (Fr)	heterozygous	—	moderate-severe
45 P12 (Fr)	—	—	severe-profound
P13 (Fr)	—	—	profound
P14 (Alg)	heterozygous	—	moderate-severe
P15	—	—	severe-profound
P16 (Fr)	homozygous	—	severe**
50 (mother/Fr, father/Pol)	—	—	—
P17 (Fr)	heterozygous	—	severe***
1885 (NZ)	heterozygous	—	profound
2254 (NZ)	—	—	moderate-severe

55

The analysis reported here concerns deaf children of the various families except for family 1873 (see patients and methods).

* moderate in one ear, severe in the other ear.

60

** moderate hearing loss in mother (severe at high frequencies), *** mild hearing loss in father, who are heterozygous carriers for the 30delG mutation.

65

Geographical origins: (Alg) Algeria, (Aust) Australia, (Fr) France, (Leb) Lebanon, (NZ) New Zealand, (Pol) Poland, (Por) Portugal, (nTu) North Tunisia, (sTu) South Tunisia, (Tur) Turkey

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 15

<210> SEQ ID NO 1
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide

<400> SEQUENCE: 1

tcttttccag agcaaaccgc c 21

<210> SEQ ID NO 2
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide

<400> SEQUENCE: 2

tgagcacggg ttgcctcatc 20

<210> SEQ ID NO 3
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide

<400> SEQUENCE: 3

gacacgaaga tcagctgcag 20

<210> SEQ ID NO 4
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide

<400> SEQUENCE: 4

ccaggctgca agaactgtg 20

<210> SEQ ID NO 5
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide

<400> SEQUENCE: 5

agacgatcct gggggtgtga acaaa 25

<210> SEQ ID NO 6
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide

<400> SEQUENCE: 6

atcctggggg tgtga 15

-continued

<210> SEQ ID NO 7
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Oligonucleotide

<400> SEQUENCE: 7

agacgatcct gggggetcac cgtcctc 27

<210> SEQ ID NO 8
 <211> LENGTH: 2314
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

gatttaatcc tatgacaaac taagttgggt ctgtcttcac ctgttttggg gaggttgtgt 60
 aagagttggg gtttgcctcag gaagagattt aagcatgctt gcttaccag actcagagaa 120
 gtctccctgt tctgtcctag ctagtgatc ctgtgttggg tgcattcgtc tttccagag 180
 caaacccccc agagtagaag atggattggg gcacgctgca gacgatcctg ggggggtgtga 240
 acaaacactc caccagcatt ggaagatct ggctcaccgt cctcttcatt tttcgatta 300
 tgatcctcgt tgtggctgca aaggaggtgt ggggagatga gcaggccgac tttgtctgca 360
 acaccctgca gccaggctgc aagaactgtt gctacgatca ctacttcccc atctcccaca 420
 tccggctatg gccctcag ctgatcttcg tgtccacgcc agcgtccta gtggccatgc 480
 acgtggccta ccggagacat gagaagaaga ggaagtcat caagggggag ataaagagtg 540
 aatttaagga catcgaggag atcaaaaccc agaaggtccg catcgaaggc tcctctgtgt 600
 ggacctacac aagcagcctc ttcttccggg tcatcttcga agccgccttc atgtactgtc 660
 tctatgtcat gtacgacggc ttctccatgc agcggctggg gaagtgcaac gcttggcctt 720
 gtcccacac tgtggactgc tttgtgtccc ggcccacgga gaagactgct tttcacagtg 780
 ttcattgatg cagtgtctgg aatttgcctc ctgctgaatg tcaactgaatt gtgttatgtg 840
 ctaattagat attgttctgg gaagtcaaaa aagccagttt aacgcattgc ccagttgtta 900
 gattaagaaa tagacagcat gagagggatg aggcaacccg tgctcagctg tcaaggctca 960
 gtcgccagca tttcccaca caaagattct gaccttaaat gcaaccattt gaaaccctg 1020
 taggcctcag gtgaaactcc agatgccaca atgagctctg ctcccctaaa gctcaaaac 1080
 aaaggcctaa ttctatgcct gtcttaattt tctttcactt aagttagttc cactgagacc 1140
 ccaggctgtt aggggttatt ggtgtaaggt actttcatat tttaaacaga ggatctggc 1200
 atttgtttct ttctctgagg acaagagaaa aaagccaggt tccacagagg acacagagaa 1260
 ggtttgggtg tctctctggg gttctttttg ccaactttcc ccacgttaaa ggtgaacatt 1320
 ggttctttca tttgctttgg aagttttaat ctctaacagt ggacaaagtt accagtgctc 1380
 taaactctgt tacacttttt ggaagtgaaa actttgtagt atgatagggt attttgatgt 1440
 aaagatgttc tggataccat tatatgttcc ccctgtttca gaggtcaga ttgtaatatg 1500
 taaatggtat gtcattcgtc actatgatgt aatttgaaat atggctcttt ggttatgaat 1560
 actttgcagc acagctgaga gaggtctgtc gttgtattca ttgtggcat agcacctaac 1620
 aacattgtag cctcaatcga gtgagacaga ctagaagttc ctagttggct tatgatagca 1680
 aatggcctca tgtcaaatat tagatgtaat tttgtgtaag aaatacagac tggatgtacc 1740

-continued

```

accaactact acctgtaatg acaggcctgt ccaacacatc tcccttttcc atgctgtggt 1800
agccagcatc gaaaagaacg ctgatttaaa gaggtgagct tgggaatttt attgacacag 1860
taccatttaa tggggagaca aaaatggggg ccaggggagg gagaagtttc tgtcgttaaa 1920
aacgagtttg gaaagactgg actctaaatt ctgttgatta aagatgagct ttgtctacct 1980
tcaaaagtgt gtttggctta ccccttcag cctccaattt ttttaagttaa aatataacta 2040
ataacatgtg aaaagaatag aagctaaggt ttagataaat attgagcaga tctataggaa 2100
gattgaacct gaatattgcc attatgcttg acatggttcc caaaaaatgg tactccacat 2160
acttcagtga gggtaagtat tttcctgttg tcaagaatag cattgtaaaa gcattttgta 2220
ataataaaga atagctttaa tgatatgctt gtaactaaaa taattttgta atgtatcaaa 2280
tacatttaaa acattaaaat ataattctta taat 2314

```

```

<210> SEQ ID NO 9
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
      Oligonucleotide

```

```
<400> SEQUENCE: 9
```

```
ctagtgattc ctgtgttgtg tgc 23
```

```

<210> SEQ ID NO 10
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
      Oligonucleotide

```

```
<400> SEQUENCE: 10
```

```
ataatgcaa aatgaagag ga 22
```

```

<210> SEQ ID NO 11
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
      Oligonucleotide

```

```
<400> SEQUENCE: 11
```

```
aaaaaaaaatc ctggggggtg tg 22
```

```

<210> SEQ ID NO 12
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
      Oligonucleotide

```

```
<400> SEQUENCE: 12
```

```
aaaaaaaaatc ctggggggtg ga 22
```

```

<210> SEQ ID NO 13
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide

<400> SEQUENCE: 13

cagcattgga aagatctggc tca 23

<210> SEQ ID NO 14
<211> LENGTH: 63
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide

<400> SEQUENCE: 14

cgccccgcgc gcccccgcgc cggccccgcg ccccccccc ctagtgattc ctgtgttg 60
tgc 63

<210> SEQ ID NO 15
<211> LENGTH: 2312
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

gatttaatcc tatgacaaac taagttggtt ctgtcttcac ctgttttggg gaggttgtgt 60
aagagttggt gttgtctcag gaagagatatt aagcatgctt gcttaccag actcagagaa 120
gtctccctgt tetgtctcag ctatgttctt gtgtttgtgt cattcgtctt ttccagagca 180
aaccccccag agtagaagat ggattggggc acgctgcaga cgatcctggg ggggtgtgaa 240
aaacactcca ccagcattgg aaagatctgg ctaccgtcc tottcaatct tcgcattatg 300
atcctcgttg tggctgcaaa ggaggtgtgg ggagatgagc aggcogactt tgtctgcaac 360
accctgcagc caggctgcaa gaacgtgtgc tacgatcact acttccccat ctcccacatc 420
cggtatggg cctgcagct gatctctgtg tccagcccag cgctcctagt ggccatgcac 480
gtggcctacc ggagacatga gaagaagagg aagttcatca agggggagat aaagagtgaa 540
ttaaaggaca tcgaggagat caaaaccag aaggtccgca tcgaaggctc cctgtggtgg 600
acctacaaa gcagcatctt ctccggggtc atcttcgaag ccgccttcat gtacgtcttc 660
tatgtcatgt acgacggctt ctccatgcag cggctggtga agtgcaacgc ctggccttgt 720
cccaactctg tggactgctt tgtgtcccgg cccaecggaga agactgtctt tcacagtgtt 780
catgattgca gtgtctggaa tttgcatcct gctgaatgtc actgaattgt gttatttgc 840
aattagatat tgttctggga agtcaaaaa gccagtttaa cgcattgccc agttgtttaga 900
ttaagaaata gacagcatga gagggatgag gcaaccctgt ctacagctgtc aaggctcagt 960
cgccagcatt tccaacaca aagattctga ccttaaatgc aaccattga aaccctgta 1020
ggcctcaggt gaaactccag atgccacaat gagctctgct cccctaaagc ctcaaaaaca 1080
aggcctaatt ctatgcctgt cttaattttc tttcacttaa gttagttcca ctgagacccc 1140
aggctgttag gggttattgg tgtaaggtac tttcatatct taaacagagg atatcgccat 1200
ttgtttcttt ctctgaggac aagagaaaa agccaggctc cacagaggac acagagaagg 1260
tttgggtgtc ctccctgggt tctttttgcc aactttcccc acgttaaagg tgaacattgg 1320
ttctttcatt tgctttggaa gttttaatct ctaacagttg acaaagtac cagtgcctta 1380
aactctgtta cactttttgg aagtgaaaa tttgtagtat gataggttat tttgatgtaa 1440
agatgttctg gataaccatta tatgttcccc ctgtttcaga ggctcagatt gtaatatgta 1500

-continued

aatggtatgt cattcgctac tatgatttaa tttgaaatat ggtcttttgg ttatgaatac	1560
tttgcagcac agctgagaga ggctgtctgt tgtattcatt gtggcatag cacctaacaa	1620
cattgtagcc tcaatcgagt gagacagact agaagttcct agttggctta tgatagcaaa	1680
tggcctcatg tcaaatatta gatgtaattt tgtgtaagaa atacagactg gatgtaccac	1740
caactactac ctgtaatgac aggcctgtcc aacacatctc ccttttccat gctgtggtag	1800
ccagcatcgg aaagaacgct gatttaaaga ggtgagcttg ggaattttat tgacacagta	1860
ccatttaatg gggagacaaa aatgggggccc aggggagggga gaagtctctg tcgttaaaaa	1920
cgagtttggg aagactggac tctaatttct gttgattaaa gatgagcttt gtctaccttc	1980
aaaagtttgt ttggcttacc cccttcagcc tccaattttt taagtgaaaa tataactaat	2040
aacatgtgaa aagaatagaa gctaaggttt agataaatat tgagcagatc tataggaaga	2100
ttgaacctga atattgccat tatgcttgac atggtttcca aaaaatggta ctccacatag	2160
ttcagtgagg gtaagtattt tcctgttgtc aagaatagca ttgtaaaagc attttgtaat	2220
aataaagaat agctttaatg atatgcttgt aactaaaata attttgtaat gtatcaata	2280
catttaaaac attaaaatat aatctctata at	2312

What is claimed is:

1. A method for diagnosing the autosomal prelingual non-syndromic deafness in a human subject, comprising assaying a biological sample comprising a connexin 26 gene obtained from the human subject to determine the presence of a mutation in the connexin 26 gene,
 wherein the assaying comprises contacting the connexin 26 gene with an allele-specific probe, wherein the allele-specific probe specifically hybridizes with a connexin 26 gene comprising the mutation, but not to the wild-type connexin 26 gene, and detecting a hybrid formed between the allele-specific probe and the connexin 26 gene,
 wherein the mutation results in a deletion of a guanosine at a position corresponding to position 30 of the coding region of the connexin 26 gene,

and wherein the presence of the mutation indicates that the human subject has autosomal prelingual non-syndromic deafness.
 2. The method according to claim 1, wherein assaying comprises amplifying the connexin 26 gene using a pair of primers.
 3. The method according to claim 1, wherein the biological sample is a blood sample.
 4. The method according to claim 1, wherein assaying a sample comprising a connexin 26 gene comprises sequencing all of the connexin 26 gene.
 5. The method according to claim 1, wherein assaying a sample comprising a connexin 26 gene further comprises sequencing a portion of the connexin 26 gene.
 6. The method according to claim 2, wherein amplifying the connexin 26 gene using a pair of primers comprises use of nucleotide sequences as set forth in SEQ ID NOs: 1 and 2.

* * * * *

专利名称(译)	连接蛋白26基因内的突变负责语前非综合征性耳聋和检测方法		
公开(公告)号	US9169517	公开(公告)日	2015-10-27
申请号	US13/899963	申请日	2013-05-22
[标]申请(专利权)人(译)	PETIT CHRISTINE denoyelle格蕾森方丝 WEIL DOMINIQUE 马林duvernois桑德林 Guesdon Jean-Luc		
申请(专利权)人(译)	PETIT , CHRISTINE DENOYELLE-GRYSON弗朗索瓦· WEIL , DOMINIQUE MARLIN-DUVERNOIS , SANDRINE GUESDON , JEAN-LUC		
当前申请(专利权)人(译)	巴斯德研究所		
[标]发明人	PETIT CHRISTINE DENOYELLE GRYSON FRANCOISE WEIL DOMINIQUE MARLIN SANDRINE GUESDON JEAN LUC		
发明人	PETIT, CHRISTINE DENOYELLE-GRYSON, FRANCOISE WEIL, DOMINIQUE MARLIN, SANDRINE GUESDON, JEAN-LUC		
IPC分类号	C12Q1/68 C12P19/34 G01N33/53 C12N15/09 C12Q1/6883 C12Q1/6886 G01N33/566		
CPC分类号	C12Q1/6883 C12Q1/6886 C12Q2600/156		
代理机构(译)	FOLEY & Lardner的律师事务所		
优先权	11/826141 2012-03-27 US 10/278089 2007-08-21 US 09/485415 2002-11-26 US PCT/EP1998/005175 1998-08-14 WO 60/055863 1997-08-15 US		
其他公开文献	US20140113285A1		
外部链接	Espacenet USPTO		

摘要(译)

一种纯化的多核苷酸，其具有对应于突变序列的核苷酸链，其以野生形式编码与遗传性感觉缺陷有关的多肽，其中所述突变的纯化的多核苷酸具有引起前语非综合征性耳聋的突变，所述突变性耳聋选自自由以下组成的组：特异性缺失至少一个核苷酸。

